Sequence-based prediction of olfactory receptor responses

Shashank Chepurwar[1], Abhishek Gupta[1,2], Rafi Haddad[3], Nitin Gupta[1,*]

[1] Department of Biological Sciences and Bioengineering, Indian Institute of Technology Kanpur, Kanpur, Uttar Pradesh 208016, India

[2] Department of Chemistry, Indian Institute of Technology Kanpur, Kanpur, Uttar Pradesh 208016, India

[3] The Gonda Multidisciplinary Brain Research Center, Bar-Ilan University, Ramat-Gan, 5290002, Israel

* Corresponding author, email address: guptan@iitk.ac.in

Keywords: olfaction, olfactory, anopheles, drosophila, receptors

Summary

Computational prediction of how strongly an olfactory receptor responds to various odors can help in

bridging the widening gap between the large number of receptors that have been sequenced and the

small number of experiments measuring their responses. Previous efforts in this area have predicted the

responses of a receptor to some odors, using the known responses of the same receptor to other odors.

Here we present a method to predict the responses of a receptor without any known responses, by

using available data about the responses of other conspecific receptors and their sequences. We applied

this method to olfactory receptors in insects *Drosophila melanogaster* (both adult and larva) and

Anopheles gambiae, and to mouse and human olfactory receptors. We found the predictions to be in

significant agreement with the experimental measurements. The method also provides clues about the

response-determining positions within the receptor sequences.

Introduction

Odor sensing begins with the binding of odor molecules to olfactory receptors (ORs) expressed on the

membranes of olfactory sensory neurons. An organism often expresses a repertoire of tens or hundreds

of types of receptors, which combinatorially can detect a very large number of odor stimuli (Malnic et

al., 1999). The amino acid sequences of these receptors – and consequently their tuning profiles to

odors – vary from species to species, in accordance with their environmental niches.

Identification of the odors that a given OR responds to, also known as deorphanization, is fundamental

to understanding olfactory processing. In cases where the sensory neurons expressing a particular OR

are known and easy to target with electrodes, the cognate odors can be identified using extracellular

recordings from those neurons (Lu et al., 2007; Olsson and Hansson, 2013; Li et al., 2018). An alternative

approach is to express the OR in an easy-to-target cell using heterologous expression systems —

including cell lines, *Xenopus* oocytes, or the empty neuron system; these have been used in different

species including humans (Touhara et al., 1999; Mainland et al., 2015), mice (Oka et al., 2006; Saito et

al., 2009), fruit flies (de Bruyne et al., 2001; Hallem and Carlson, 2006; Lin and Potter, 2015), mosquitoes

(Carey et al., 2010a; Wang et al., 2010), moths (de Fouchier et al., 2017), and tsetse flies (Chahda et al.,

2019). However, such experiments need to be performed separately for each receptor and are time-

consuming; moreover, some receptors do not express well in the heterologous systems (Ronderos et al.,

2014). High-throughput methods for identifying OR-odor interactions have also been developed

recently, but these require elaborate experimental pipelines for each species (Jiang et al., 2015; Hu and

Matsunami, 2018; Jones et al., 2019). The high-throughput methods also generate many false positives,

so it is recommended that the identified responses be further verified with targeted experiments

(Nishizumi and Sakano, 2015).

With lowering costs of sequencing, the sequences of ORs are rapidly becoming available for various

organisms, including multiple species of fruit flies (Drosophila 12 Genomes Consortium, 2007; Gardiner

et al., 2008; Ramasamy et al., 2016), mosquitoes (Leal et al., 2013; Neafsey and Waterhouse, 2015;

Lombardo et al., 2017), bees (Elsik et al., 2015; Karpe et al., 2017) and tsetse flies (Attardo et al., 2019).

However experimental data about odor responses are available for relatively few ORs. Computational

predictions can potentially bridge this gap. Indeed, computational methods have been developed to

predict the response of an OR to new odors based on its known responses to other odors, by analyzing

the chemical structure (Schmuker et al., 2007), molecular volume (Saberi and Seyed-Allaei, 2016), or

other physicochemical parameters of odor molecules ((Haddad et al., 2008; Boyle et al., 2013; Gabler et

al., 2013; Bushdid et al., 2018; Kepchia et al., 2019)). However, these methods can only be used for ORs

whose responses to some odors are known.

Here, we present a computational approach for predicting the responses of an OR even when no odor

responses are available for that OR; our method instead uses the known responses of other conspecific

ORs and the sequence similarities among them. We demonstrate the effectiveness of this approach by

comparing our predictions with the experimentally measured responses recordings in insects. Our

method also provides insights into the response-determining positions in the receptor sequences.

Results

Lack of correlation between OR sequences and responses

As odor responses of an OR depend on its three-dimensional structure, which in turn depends on the

amino-acid sequence, we expected that the sequence similarity between a pair of ORs will relate to the

similarity in their responses. To test this idea, we used large scale datasets of odor responses of 24 ORs

in *D. melanogaster* (Hallem and Carlson, 2006) and 50 ORs in *A. gambiae* (Carey et al., 2010) to sets of

110 odors; these odor sets were partially different between the two species but identical for all ORs

within a species. For each pair of ORs within a species, we calculated the distance between their 110-

length response vectors (**see Methods**). We quantified amino-acid sequence similarity for each pair of

ORs within a species using global sequence alignment. While we expected to see a negative correlation

between sequence similarity and response distance, we surprisingly found no correlation at all (**Fig. 1**):

the Pearson's correlation coefficient was 0.0018 (P = 0.97, N = 276 pairs of receptors) in *D.*

melanogaster, and 0.026 (P = 0.35, N = 1225 pairs) in *A. gambiae*.

Identification of response-determining positions

The task of sequence-based predictions is made challenging by the very low level of conservation among

the ORs: the percentage amino-acid identity between pairs of ORs was only 20.23 ± 4.48 % in *D.*

melanogaster and 20.60 ± 8.17 % in *A. gambiae* (mean ± s.d. calculated over all pairs within a species)*.*

The lack of correlation between the sequences and the responses is partly because only a small fraction

of residues in the sequence are involved in determining odor-specificity, and the similarity of these

residues is not represented well by the overall sequence similarity (Man et al., 2007). However, as there

is very little structural information available for ORs, the positions of these response-determining

residues are not known. We took an empirical approach to rank each position in the multiple sequence

alignment of all receptor sequences according to its importance in determining the odor responses (see

Methods). Within a species, the ranking was done jointly over all ORs to avoid overfitting. Using this

ranked list of positions for each species, we selected the top 20 positions that should help in predicting

the odor responses of new ORs.

We first confirmed that the subsequences of amino acids formed by these 20 positions in each OR were

indeed related to the responses. We found the correlation between the response distance and sequence

similarity (calculated using only the subsequences formed by the 20 positions) to be -0.56 (P= 1.7×10^{-24},

N = 276 pairs; **Fig. 2A1**) in *D. melanogaster*, and -0.39 (P = 1.7×10^{-45}, N = 1225 pairs; **Fig. 2A2**) in *A.*

gambiae. Compared to the result seen with the full sequence (**Fig. 1**), these correlations were

significantly negative. Similar negative correlations were observed even if we used only half of the odors

for calculating the top 20 positions and the other half for calculating the response similarity

(**Supplementary Fig. S1**). If we varied the number of top ranked positions used in the analysis, the

correlation coefficients did not change for numbers between 10 and 40 but gradually reduced for larger

numbers of positions (**Figs. 2B1,2**), suggesting that lower-ranked residues are not important for the

prediction of responses. Subsequently, we used the top 20 positions as the response-determining

positions in our analysis (**Figs. 2B1,2** shows that our results would not change if we chose some other

number between 50 % and 200 % of this number).

We then checked the robustness of the top 20 positions by reducing the number of receptors used for

ranking the positions. When we removed one receptor (this was done by removing each receptor, one

at a time, to get the averages and error bars), we found that on average ~16 of the 20 positions in *D.*

melanogaster and ~19 of the 20 positions in *A. gambiae* were retained in the top 20. The number of

retained positions dropped gradually as more receptors were left out (again, all possible combinations

were tested for each number). But even with half of the receptors removed, at least 8 of the original 20

positions remained in top 20 in both the species (**Figs. 2C1,2**). This analysis indicated that the selected

positions were robust to minor perturbations in the datasets.

Although the top 20 alignment positions were chosen independently in *D. melanogaster* and *A.*

gambiae, we found that 3 of the top 20 positions were conserved between the species. This number

was three times the number of conserved positions expected by chance (20 x 20/369 = 1.08, where 369

is the number of alignment positions that were ranked in both the species), further increasing our

confidence in the reliability of the identified positions.

Predicting responses using the identified positions

Having identified reasonable candidates for response-determining positions, common to all ORs within a

species, we asked if it is possible to reliably predict the responses of a query OR to a panel of odors using

only the sequence information about the amino acids present at the identified positions. We tested this

idea using a simple and intuitive approach, in which the response of a query OR was estimated as the

mean of the responses of known ORs, weighted by their similarity to the query OR at the response-

determining positions using the BLOSUM62 amino-acid substitution matrix (see **Methods**).

We first applied this approach to predict the responses of the 24 and the 50 ORs in the *D. melanogaster*

and *A. gambiae* datasets, respectively. By turn, each OR was treated as a query OR and its response was

predicted using the responses of the remaining ORs in the same species (excluding the query OR). We

found that the predicted response profiles (i.e., the 110-length vectors of odor responses) of the ORs

were correlated with their actual response profiles (**Figs. 3A,B**); the correlations were positive and

statistically significant in 21 of the 24 ORs in *D. melanogaster*, and 29 of the 50 ORs in *A. gambiae*. Mean

of these correlation values was significantly greater than zero: 0.46 (P = 2.53 x 10^{-7}, t-test; P = 1.63 x 10^{-5},

sign-rank test; N = 24 ORs) in *D. melanogaster,* and 0.25 (P = 1.18 x 10^{-4}, t-test; P = 1.19 x 10^{-4}, sign-rank

test; N = 50 ORs) in *A. gambiae*. In contrast, control predictions obtained by shuffling the response

matrices (see **Methods**) were not correlated with the actual responses (**Fig. 3C**): mean correlation values

were 0.0002 (P = 0.95, t-test; N = 24 ORs) in *D. melanogaster,* and 0.0008 (P = 0.65, t-test; N = 50 ORs) in

A. gambiae.

The correlation coefficient between the actual and the predicted odor response vectors tells whether

these responses have the same relative magnitude across different odors, but does not tell how close

the two responses are in absolute terms (a predicted response that is several times larger than the

actual response for every odor will have a perfect correlation). We therefore also used a distance-based

metric and checked whether our predictions were closer to the actual responses than the *control*

predictions were. We define $D_{Pred,Actual}$ as the absolute difference between the predicted response

and the actual response of an OR to an odor, and $D_{Ctrl,Actual}$ as the absolute difference between the

control prediction and the actual response. Thus, these two terms indicate the error in the actual and

the control predictions. If our predicted response is better (closer to the actual response) than the

control prediction, the value of $D_{Pred,Actual} - D_{Ctrl,Actual}$ should be negative. The heat maps in **Fig. 3D**

show the comparisons for all the predictions in *D. melanogaster* and *A. gambiae* and reveal a very high

abundance of negative values. In *D. melanogaster*, the average $D_{Pred,Actual}$ of an OR (28.60) was

smaller than average $D_{Ctrl,Actual}$ (47.84) by 19.24 spikes (P = 7.30 x 10^{-15}, paired t-test; N = 24 ORs; **Fig.**

3E1), an improvement of more than 40 %. In *A. gambiae*, the average $D_{Pred,Actual}$ (20.11) was smaller

than average $D_{Ctrl,Actual}$ (33.29) by 13.18 spikes (P = 8.46 x 10^{-26}, paired t-test; N = 50 ORs; **Fig. 3E2**),

again an improvement of about 40 %. Thus, our predicted responses using the sequence similarity at the

response-determining positions were significantly closer to the actual responses than random

predictions were. These improvements were substantial considering the absolute values of odor

responses in *D. melanogaster* (mean ± s.d. = 32.43 ± 49.29 spikes, N = 24 x 110 OR-odor combinations)

and *A. gambiae* (mean ± s.d. = 20.31 ± 37.35 spikes, N = 50 x 110).

Test on an independent dataset

Next, we collected the responses of 26 other ORs in *D. melanogaster* from 10 different studies (de

Bruyne et al., 1999, 2001, 2010; Dobritsa et al., 2003; Goldman et al., 2005; Kreher et al., 2005, 2008;

Marshall et al., 2010; Ronderos et al., 2014; Dweck et al., 2015) deposited in the DoOR database (Münch

and Galizia, 2016). Each of these ORs had responses available for some of the 110 odors used previously;

overall, 68 of the previously used odors were represented in the new dataset. These 26 ORs were

entirely different from the 24 ORs used previously in selecting the response-determining positions, and

therefore provided an independent test for our approach. Using the set of top 20 response-determining

positions as selected earlier from the original *D. melanogaster* dataset, we found that the responses

predicted for the 26 novel ORs were significantly better with our approach compared to the control

predictions (**Fig. 4A,B**): the average $D_{Pred,Actual}$ of an OR (37.01) was smaller than average $D_{Ctrl,Actual}$

(51.36) by 14.35 spikes (P = 1.13 x 10^{-7}, paired t-test; P = 5.96 x 10^{-8}, sign-rank test; N = 26 ORs), an

improvement of about 28 %. Overall, among the 506 OR-odor responses predicted (non-grey squares in

Fig. 4A), in 380 (75.1%) cases the predictions were better than the control predictions. These results

show that once the response-determining positions have been identified using known responses of

some ORs in a species, the responses of other conspecific ORs can be predicted reasonably using only

their sequences.

While predicting the responses of the new ORs above, we had relied on the known responses of 24 ORs.

Can similar predictions be made if responses of fewer ORs are available? To check this, we sampled

subsets of ORs from the original dataset of 24 ORs, and used these samples to select response-

determining positions and make the response predictions. The sizes of these samples were varied from

3 to 21 in intervals of 3, and for each size the sampling was performed 10 times. We found that the

difference between the predicted and the actual response ($D_{Pred,Actual}$) decreased gradually as the

number of sampled ORs increased but saturated for 15 or more ORs (**Fig. 4C**). A similar trend was

observed when we checked how the number of sampled ORs affected the fraction of predictions that

were better than control (**Fig. 4D**). These results suggest that a relatively small set of ORs with known

responses may be enough to make reasonable predictions for new ORs.

Predicting responses in larval *Drosophila*

Although the OR responses in larval *Drosophila melanogaster* differ from the responses in adult

Drosophila (Kreher et al., 2008), the ORs themselves are the same at the molecular level. Therefore, we

checked if the top positions identified in adult *Drosophila* can be used to predict the larval responses.

We tested this idea on a dataset of 21 ORs and 26 odors (each available at two different concentrations,

10^{-2} and 10^{-4}) from larval *Drosophila* (Kreher et al., 2008). For each concentration, we found that the

predictions were significantly better than control predictions (**Supplementary Fig. S2**): at 10^{-2}

concentration, the average $D_{Pred,Actual}$ (38.01) was smaller than average $D_{Ctrl,Actual}$ (54.72) by 16.71

spikes (P = 6.32 x 10^{-12}, paired t-test; N = 21 ORs); at 10^{-4} concentration, the average $D_{Pred,Actual}$ (12.38)

was smaller than average $D_{Ctrl,Actual}$ (16.7) by 4.32 spikes (P = 1.1 x 10^{-8}, paired t-test; N = 21 ORs).

Properties of the response-determining positions

Our results show that the identified subset of sequence positions can allow reliable prediction of the

neural responses for new ORs. To check where these top 20 positions are located within the protein

structure, we visualized these positions in the secondary structure of two representative ORs in *D.*

melanogaster and *A. gambiae* using the Protter tool (Omasits et al., 2014). These residues were not

limited to any one region, but were instead found in various parts of the protein (**Fig. 5**). It is important

to note that our analysis does not imply a causal or a mechanistic role for each of these residues in

interacting with the odor molecules, but only suggests that these residues may be involved in some

aspect of determination of odor responses, perhaps indirectly. Notably, the region with the highest

density of the top 20 positions was the seventh transmembrane helix, including 4 positions in *D.*

melanogaster and 5 in *A. gambiae* (**Fig. 5**; similar numbers were seen in other receptors). Moreover, of

the three response-determining positions that were conserved between the two species, two were

located in the seventh transmembrane helix. These observations are in agreement with previous studies

suggesting that the seventh transmembrane helix plays an important role in determining odor specificity

(Hughes et al., 2014; Ray et al., 2014).

Next, we collected data from various experimental studies that have mutated individual residues in

various ORs and have checked their effect on odor responses. We collected a total of 47 such mutated

positions from 5 studies (Nichols and Luetje, 2010; Nakagawa et al., 2012; Xu and Leal, 2013; Hughes et

al., 2014; Ray et al., 2014) and compared them with our identified set of response-determining

positions. In addition to *D. melanogaster* and *A. gambiae*, these studies also included data from *Bombyx*

mori and *Culex quinquefasciatus.* Sequences from these other species were added to our existing

multiple sequence alignment (see **Methods**), so that the mutated positions in all these species can be

compared with our alignment positions.

Given 20 response-determining positions in *D. melanogaster* and *A. gambiae* each, and with 3 being

common between the two species, 37 unique positions in the multiple sequence alignment were

marked as the top positions out of a total of 817. Thus, each of the 47 mutated positions in the curated

experimental set had a 37/817 probability of matching a top position, and the expected number of

matches by chance = 47 x 37/817 = 2.13. So, while only about 2 of the 47 mutated positions were

expected to be among the top positions by chance, we found as many as 9 of them to be among the top

positions (P = 2.28 x 10^{-4}; binomial test), providing experimental support to our response-determining

positions. These 9 experimentally studied mutations, which map to the top positions in our alignment,

are listed in **Table 1** (further, **Table S1** shows all positions). Among these, residue 146 in OR85b of *D.*

melanogaster is one of the residues expected to play a crucial role in the activation of the odorant

(Nichols and Luetje, 2010). Residue 109 mutant in OR1 of *B. mori* was found to change the reversal

potential and rectification index of the BmOR1-Orco complex (Nakagawa et al., 2012). Point mutation of

residues 165 and 194 in BmorOR1 of *B. mori* led to significant reduction in response to bombykol

pheromone (Xu and Leal, 2013). Similarly, odor responses in *A. gambiae* were found to change with

mutation of residue 369 in OR15 (Hughes et al., 2014), and residues 405 and 406 in OR1 (Ray et al.,

2014). Among the remaining 38 out of the 47 mutations in the curated set, 10 and 8 were found to be

within a distance of 1 and 2 residues, respectively, from a top column. Thus, in total, 27 of the 47

experimentally confirmed important positions matched or were adjacent to our computationally

predicted response-determining positions.

Prediction of vertebrate olfactory responses

As our method does not make any assumption about the composition of the receptor sequences, it

should be applicable to any family of chemosensory receptors. We next sought to check its utility in

predicting the responses of the vertebrate ORs. Saito et al. (Saito et al., 2009) have provided a dataset of

EC50 values of 52 mouse and 10 human ORs that responded to at least one of the 63 odors tested. We

extracted a binary response matrix (1 for response, 0 for no response) from this dataset: among the

mouse responses (52 x 63 = 3276 values), this matrix included 262 (approximately 8%) 1s and rest were

0s. Using the multiple sequence alignment provided in the same study (Saito et al., 2009) and mouse

data, we calculated the top 20 response determining positions. We then calculated the response of each

mouse receptor using the weighted average of the responses of the remaining receptors, as done for

insect receptors (see **Methods**); for comparison with actual responses, this value was also converted to

a binary response using a threshold, set to keep the total fraction of 1s in the predicted responses also

at 8%. The predictions thus obtained had a false alarm rate of about 7% and a hit rate of about 21%,

resulting in sensitivity index d' = 0.68 (P < 0.001, estimated by comparing the d' value with those

243 obtained for 1000 shuffled control predictions). We next asked if human OR responses could also be

244 predicted using the mouse training data. Using the top positions, the responses, and the threshold

245 identified from the mouse data, we predicted the binary responses of the 10 human ORs. Among the 10

246 x 63 values in the original human dataset, 78 (~12%) were 1s. Our prediction yielded 581 0's and 49 1s,

247 with a false alarm rate of about 7% and a hit rate of about 13% (d' = 0.34, P = 0.047 based on 1000

248 shuffled predictions). In summary, these results show that our method can also be used to predict

249 vertebrate OR responses with better than chance accuracy.

251 **Discussion**

252 Sequence-based approaches to response prediction require a reasonably high homology among the

253 receptors. In vertebrates, receptors with less than 40 % sequence identity were found to have little

254 overlap in their responses (Li et al., 2015). The insects ORs presented a particularly difficult test case, as

255 they have only about 20 % homology among conspecific ORs; indeed, we found that there was no

256 correlation between the overall sequence similarity and the response similarity among the ORs. We

257 were able to overcome this difficulty by focusing on subsets of positions that may be important for

258 determining odor responses. Our approach was successful in predicting the responses for not only the

259 set of fly and mosquito receptors whose data was used while developing the method, but also for a

260 completely novel set of receptors whose responses were taken from 10 independent experimental

261 datasets. Further, we found that the predictions require only about 15 ORs with known responses (**Fig.**

262 **4**). This computational approach can be particularly useful for receptors expressed in neurons that are

263 difficult to access in electrophysiology experiments, or for receptors that do not express well in

264 heterologous expression systems (Ronderos et al., 2014). Finally, we also showed that our method can

265 also be used for vertebrate receptors.

266 Our method allows prediction of responses of novel ORs for already studied odors. Previously developed

267 methods allow prediction of responses of already studied ORs to novel odors (Schmuker et al., 2007;

268 Haddad et al., 2008; Boyle et al., 2013; Gabler et al., 2013; Saberi and Seyed-Allaei, 2016; Bushdid et al.,

269 2018; Kepchia et al., 2019). These two approaches are complementary and can be combined in future

270 work to enable the prediction of the responses of novel ORs to novel odors, which would help with

271 large-scale deorphanization. While applying these approaches, care must be taken regarding the

272 concentrations of the odors in the training data, as changes in odor concentration can affect neural

responses (Hallem and Carlson, 2006; Olsen et al., 2010) and behavior (Wright et al., 2005; Yarali et al.,

2009). Another caveat is that these approaches are currently limited to predicting the response

magnitudes, and ignore the temporal patterns of spikes observed in the olfactory receptor neurons

(Spors et al., 2006; Raman et al., 2010; Montague et al., 2011; Su et al., 2011; Grillet et al., 2016; Egea-

Weiss et al., 2018).

Unlike vertebrate ORs which are G-protein coupled receptors, insect ORs are heteromeric ligand-gated

ion channels (Wicher et al., 2008), composed of a specific OR and a conserved co-receptor Orco (Larsson

et al., 2004). The structure of any insect receptor was not known until very recently when the structure

of Orco from fig wasp was studied using Cryo-EM (Butterwick et al., 2018). The approach of predicting

responses using the three-dimensional modeling of ORs, that has seen some success in vertebrates

(Bavan et al., 2014), is unlikely to be very productive in insects in the near future given the very low

sequence similarity among the ORs and with other proteins, and the paucity of available structures.

Therefore sequence-based computational methods will be important for receptors whose structures

and responses have not been determined experimentally. The sequence-based predictions are expected

to fare better when the responses being predicted are in the same range as the responses in the training

dataset, but may not fare as well in predicting uniquely strong ligands for a novel receptor (for the latter

goal, three-dimensional modeling may be necessary). Our computational predictions can be used to

identify the candidate ligands to be tested experimentally. The method can also be extended to other

types of chemosensory receptors, such as the ionotropic receptors and the gustatory receptors.

We have used a simple algorithm in our approach, where we first identify the important positions in the

sequences and then use the similarity at those positions to make the predictions. It may be possible to

achieve good predictions without explicitly identifying the important positions and instead using a

machine learning-based approach that uses all the residues and learns a non-linear mapping between

the sequences and the responses. However, we preferred the simpler deterministic approach over the

blackbox approaches typically used in machine learning because the deterministic approach provides a

transparent rationale for and information about the sequence positions that are used for making the

predictions. Expectedly, a majority of the identified positions were located in the extracellular loops or

transmembrane helices (particularly the seventh transmembrane, which has been experimentally shown

to be important for determining responses), but some were also found in intracellular loops. A previous

study has shown that mutation of intracellular residues can affect the electrophysiological responses of

neurons (Nakagawa et al., 2012). We emphasize that our approach only looks for correlations between

the residues and responses, and it is likely that some of the identified positions, particularly those found

in intracellular regions, affect the odor responses indirectly rather than through direct interactions with

the odor molecules. By comparing the 47 positions previously found to be important in mutagenesis

studies in various organisms (Nichols and Luetje, 2010; Nakagawa et al., 2012; Xu and Leal, 2013;

Hughes et al., 2014; Ray et al., 2014), we found that 9 of them (a statistically significant fraction, more

than 4 times larger than that expected by chance) mapped to the top positions identified in our analysis

(**Table 1**). This shows that our method is using reliable positions for making the predictions. The other

top positions that have not been experimentally studied so far are good candidates for future

mutagenesis studies, and could help in understanding the mechanisms of odor-OR interactions. As more

experimental evidence accumulates, the response predictions can be further improved by giving more

weightage to the experimentally verified positions among the top positions.

Methods

Response datasets

Responses of 24 *D. melanogaster* ORs were obtained from a previous study (Hallem and Carlson, 2006)

and responses of 50 *A. gambiae* ORs were obtained from another study (Carey et al., 2010), for a set of

110 odors. The response values indicate the odor-elicited spiking response of a neuron expressing the

receptor, minus the response for the diluent or the spontaneous firing rate. To serve as an independent

dataset, responses of 26 other *D. melanogaster* ORs were obtained from 10 studies (de Bruyne et al.,

1999, 2001, 2010; Dobritsa et al., 2003; Goldman et al., 2005; Kreher et al., 2005, 2008; Marshall et al.,

2010; Ronderos et al., 2014; Dweck et al., 2015) deposited in the Database of Odorant Responses

(DoOR) (Münch and Galizia, 2016), for 68 odors that were common with the earlier set of 110 odors. If

the response of a particular OR to an odor was reported in multiple studies, we used the average of

those values as the actual response.

Sequence analysis

The olfactory receptor sequences for *D. melanogaster* and *A. gambiae* were obtained from the Database

of Olfactory Receptors (Nagarathnam et al., 2014) and the (Hill et al., 2002) study, respectively. The

sequence similarity between any two receptor sequences was determined using the Needleman

Wunsch Algorithm using the BLOSUM62 scoring matrix (Needleman and Wunsch, 1970; Henikoff and

Henikoff, 1992). Multiple sequence alignment was performed for a total of 140 (62 in *D. melanogaster*

and 78 in *A. gambiae*) olfactory receptor sequences using MAFFT multiple sequence alignment program,

and additional sequences from other species (for validation) were added individually using the '-add'

option (Katoh and Frith, 2012).

The OCTOPUS tool (Viklund and Elofsson, 2008) was used to predict the membrane topology of the

receptors and the Protter tool was used to visualize the 2-dimensional topology of the receptors

(Omasits et al., 2014).

Ranking the response-determining positions

We identified the response-determining sequence positions for *D. melanogaster* and *A. gambiae* using

the following approach, which quantifies at each position how well the differences in amino acids across

the receptors correlate with the differences in their responses. These positions corresponded to

columns in the multiple sequence alignment, and were same for all the receptors within a species

(allowing different positions for each receptor can provide better correlations for individual receptors

due to overfitting but will have poor generalizability for other receptors).

From the total of 817 alignment columns in the multiple sequence alignment, we first eliminated the

columns which were either fully conserved or had gaps in more than half of the sequences, leaving 378

columns to be ranked in *D. melanogaster* and 392 columns to be ranked in *A. gambiae*, of which 369

were common to both species (see **Table S1**). We denote an element of the alignment by $A_{r,c}$ where c

indicates the alignment column and $r \in [1, 24] \; or \; [1, 50]$ in *D. melanogaster* and *A. gambiae*,

respectively, indicates the row corresponding to a receptor with known responses. The following steps

were carried out separately for each species. To calculate the rank for each of the columns, we first

calculated a sequence similarity vector (SS_c) for each column, each element ($SS_{(r_1, r_2),c}$) of which

contained the pairwise amino acid similarity scores, calculated using BLOSUM62 matrix, between all

pairs of receptors (r_1, r_2) that did not have a gap in the alignment column c:

$$SS_{(r_1, r_2),c} = BLOSUM62\left(A_{r_1,c}, A_{r_2,c}\right) \forall \; (r_1, r_2) \; where \; A_{r_1,c} \neq \; gap \; \& \; A_{r_2,c} \neq gap$$

Similarly, we computed a response distance vector (DR_c^o) for the same pairs of ORs in the column, each

element ($DR_{(r_1, r_2),c}^o$) of which contained the absolute difference between the responses of receptors

$r_1 \; and \; r_2$ to an odor o:

$$DR_{(r1, r2),c}^o = \left|N_{r_1}^o - N_{r_2}^o\right| \forall \; (r_1, r_2) \; where \; A_{r_1,c} \neq \; gap \; \& \; A_{r_2,c} \neq \; gap,$$

where N_r^o denotes the neural response of receptor r to odor o. We used the L1-norm instead of the

Euclidean distance to avoid giving extra weightage to larger values. Note that SS_c and DR_c^o are both

vectors of the same length; this length is equal to the number of all possible pairs of receptors that did

not have a gap in column c. We then calculated the Pearson's correlation coefficient (R_c^o) between the

sequence similarity (SS_c) and response difference (DR_c^o) vectors for each column c and each odor o. The

P-value of this correlation, denoted by P_c^o, was also noted. A negative value of R_c^o indicates that the

receptors that have high sequence similarity at the alignment position c also have high response

similarity (small distance) for odor o. Finally, we calculated the total score of a column as

371
$$S_c = - \sum_{\substack{o=1, \\ R_c^o < 0}}^{110} \log P_c^o$$

Use of P-values allowed us to give more weightage to highly significant correlations. Columns with high

S_c scores were used subsequently as the response-determining positions.

Response prediction

We first generated the subsequences using the amino acids at the response-determining positions; this

was done for the receptors with known responses and also for the receptor whose response is being

predicted (*query*). Subsequence similarity ($SSS_{r,query}$) between the query receptor and each of the

known receptors was calculated by taking the average of BLOSUM62 score for all positions in the two

subsequences (excluding positions with gaps). The similarity values were then linearly scaled (denoted

by $\overline{SSS}_{r,query}$) to a range of 0 to 100 to avoid negative weights in the subsequent steps:

379
$$\overline{SSS}_{r,query} = \frac{SSS_{r,query} - \min_i SSS_{r_i,query}}{\max_i SSS_{r_i,query} - \min_i SSS_{r_i,query}} \times 100$$

380   The predicted response ($\widetilde{N}_{query}^o$) of the query receptor to an odor o was calculated as a weighted

381   average of the known responses:

382
$$\widetilde{N}_{query}^o = \frac{\sum_{j=1}^n (\overline{SSS}_{r_j,query} \times N_{r_j}^o)}{\sum_{j=1}^n \overline{SSS}_{r_j,query}}$$

where n is the number of receptors with known responses and $N_{r_j}^o$ denotes the neural response of

receptor r_j to odor o.

Control Predictions

Control predictions were obtained by shuffling the OR-odor response matrix and taking the shuffled

response as the control prediction for each OR-odor combination (this operation allowed generation of

random predictions while maintaining the overall statistical properties of the response values). The

shuffling was not limited to shifting of rows or columns, but included the shuffling of all elements in the

matrix. The shuffling was performed 50 times independently, and the results (correlations or $D_{Ctrl,Actual}$

values) were averaged over the 50 shufflings.

Statistical analysis and code availability

All analyses were performed in MATLAB. A modified version of Gramm plotting toolbox (Morel, 2018)

was used to draw the plots. We used Pearson correlation to measure the correlations, and t-tests and

Wilcoxon signed-rank tests to compare the means. All tests were two-tailed. The number of sample

points is shown for each test in the results. The code developed in this study can be accessed from

https://github.com/neuralsystems/OR_response_prediction.

Acknowledgements

We thank members of the N.G. laboratory for helpful comments.

Funding

This work was supported by the Wellcome Trust/DBT India Alliance Fellowship (grant number

IA/I/15/2/502091) awarded to N.G., and ISF-UGC joint research program in which R.H. was supported by

the ISF (grant No. 2307/15) and N.G. was supported by the UGC (F.No. 6-11/2016[IC]).

Declaration of interests

The authors declare no competing financial interests.

Author Contributions

N.G. conceived the study. S.C., A.G., R.H., and N.G. developed the methodology; S.C. and A.G. developed

the software and performed analysis; N.G. and R.H. secured funding; S.C., A.G., R.H., and N.G. wrote the

manuscript.

References

Attardo GM et al. (2019) The Glossina Genome Cluster: Comparative Genomic Analysis of the Vectors of African

Trypanosomes. bioRxiv.

Bavan S, Sherman B, Luetje CW, Abaffy T (2014) Discovery of Novel Ligands for Mouse Olfactory Receptor MOR42-

3 Using an In Silico Screening Approach and In Vitro Validation. PLoS One 9.

Boyle SM, McInally S, Ray A (2013) Expanding the olfactory code by in silico decoding of odor-receptor chemical

space. Elife:1–17.

Bushdid C, De March CA, Fiorucci S, Matsunami H, Golebiowski J (2018) Agonists of G-Protein-Coupled Odorant

Receptors Are Predicted from Chemical Features. J Phys Chem Lett 9:2235–2240.

Butterwick JA, del Mármol J, Kim KH, Kahlson MA, Rogow JA, Walz T, Ruta V, Mármol J, Kim KH, Kahlson MA,

Rogow JA, Walz T, Ruta V (2018) Cryo-EM structure of the insect olfactory receptor Orco. Nature 560:447–

452.

Carey AF, Wang G, Su C-Y, Zwiebel LJ, Carlson JR (2010) Odorant reception in the malaria mosquito Anopheles

gambiae. Nature 464:66–71.

Chahda JS, Soni N, Sun JS, Ebrahim SAM, Weiss L, Carlson JR (2019) The molecular and cellular basis of olfactory

response to tsetse fly attractants. PLOS Genet:1–22.

de Bruyne M, Clyne PJ, Carlson JR (1999) Odor coding in a model olfactory organ: the Drosophila maxillary palp. J

Neurosci 19:4520–4532.

de Bruyne M, Foster K, Carlson JR (2001) Odor coding in the Drosophila antenna. Neuron 30:537–552.

de Bruyne M, Smart R, Zammit E, Warr CG (2010) Functional and molecular evolution of olfactory neurons and

receptors for aliphatic esters across the Drosophila genus. J Comp Physiol A 196:97–109.

de Fouchier A, Walker WB, Montagne N, Steiner C, Binyameen M, Schlyter F, Chertemps T, Maria A, Francois M-C,

Monsempes C, Anderson P, Hansson BS, Larsson MC, Jacquin-Joly E (2017) Functional evolution of

Lepidoptera olfactory receptors revealed by deorphanization of a moth repertoire. Nat Commun 8:1–11.

Dobritsa AA, Naters WVDG Van, Warr CG, Steinbrecht RA, Carlson JR, Haven N, Vic C (2003) Integrating the

Molecular and Cellular Basis of Odor Coding in the Drosophila Antenna. Neuron 37:827–841.

Drosophila 12 Genomes Consortium (2007) Evolution of genes and genomes on the Drosophila phylogeny. Nature

450:203–218.

Dweck HKM, Ebrahim SAM, Farhan A, Hansson BS, Stensmyr MC (2015) Olfactory Proxy Detection of Dietary

Antioxidants in Drosophila. Curr Biol 25:455–466.

Egea-Weiss A, Renner A, Kleineidam CJ, Szyszka P (2018) High Precision of Spike Timing across Olfactory Receptor
Neurons Allows Rapid Odor Coding in Drosophila. ISCIENCE 4:76–83.

Elsik CG, Tayal A, Diesh CM, Unni DR, Emery ML, Nguyen HN, Hagen DE (2015) Hymenoptera Genome Database :
integrating genome annotations in HymenopteraMine. Nucleic Acids Res 44:793–800.

Gabler S, Soelter J, Hussain T, Sachse S, Schmuker M (2013) Physicochemical vs. vibrational descriptors for
prediction of odor receptor responses. Mol Inform 32:855–865.

Gardiner A, Barker D, Butlin RK, Jordan WC, Ritchie MG (2008) Drosophila chemoreceptor gene evolution :
selection , specialization and genome size. Mol Ecol 17:1648–1657.

Goldman AL, Van Der Goes Van Naters W, Lessing D, Warr CG, Carlson JR (2005) Coexpression of two functional
odor receptors in one neuron. Neuron 45:661–666.

Grillet M, Campagner D, Petersen R, Mccrohan C, Cobb M (2016) The peripheral olfactory code in Drosophila
larvae contains temporal information and is robust over multiple timescales. Proc R Soc B 283:1–8.

Haddad R, Carmel L, Sobel N, Harel D (2008) Predicting the Receptive Range of Olfactory Receptors. PLoS Comput
Biol 4:0001–0008.

Hallem EA, Carlson JR (2006) Coding of Odors by a Receptor Repertoire. Cell 125:143–160.

Henikoff S, Henikoff JG (1992) Amino acid substitution matrices from protein blocks. Proc Natl Acad Sci 89:10915–
10919.

Hill CA, Fox AN, Pitts RJ, Kent LB, Tan PL, Chrystal MA, Cravchik A, Collins FH, Robertson HM, Zwiebel LJ (2002) G
protein-coupled receptors in Anopheles gambiae. Science (80-) 298:176–178.

Hu XS, Matsunami H (2018) High-Throughput Odorant Receptor Deorphanization Via Phospho-S6 Ribosomal
Protein Immunoprecipitation and mRNA Profiling BT - Olfactory Receptors: Methods and Protocols. In
(Simoes de Souza FM, Antunes G, eds), pp 95–112. New York, NY: Springer New York.

Hughes DT, Wang G, Zwiebel LJ, Luetje CW (2014) A determinant of odorant specificity is located at the
extracellular loop 2-transmembrane domain 4 interface of an anopheles gambiae odorant receptor subunit.
Chem Senses 39:761–769.

Jiang Y, Gong NN, Hu XS, Ni MJ, Pasi R, Matsunami H (2015) Molecular profiling of activated olfactory neurons
identifies odorant receptors for odors in vivo. Nat Neurosci 18:1446–1454.

Jones EM, Jajoo R, Cancilla D, Lubock NB, Wang J, Satyadi M, Cheung R, de March C, Bloom JS, Matsunami H, Kosuri
S (2019) A Scalable, Multiplexed Assay for Decoding GPCR-Ligand Interactions with RNA Sequencing. Cell Syst
8:254–260.e6.

Karpe SD, Dhingra S, Brockmann A, Sowdhamini R (2017) Computational genome-wide survey of odorant receptors
from two solitary bees Dufourea novaeangliae (Hymenoptera : Halictidae) and Habropoda laboriosa (
Hymenoptera : Apidae). Sci Rep 7:1–13.

Katoh K, Frith MC (2012) Adding unaligned sequences into an existing alignment using MAFFT and LAST.

Bioinformatics 28:3144–3146.

Kepchia D, Xu P, Terryn R, Castro A, Schürer SC, Leal WS, Luetje CW (2019) Use of machine learning to identify

novel , behaviorally active antagonists of the insect odorant receptor co-receptor (Orco) subunit. Sci Rep

9:1–18.

Kreher SA, Kwon JY, Carlson JR (2005) The Molecular Basis of Odor Coding in the Drosophila Larva. Neuron 46:445–

456.

Kreher SA, Mathew D, Kim J, Carlson JR (2008) Translation of Sensory Input into Behavioral Output via an Olfactory

System. Neuron 59:110–124.

Larsson MC, Domingos AI, Jones WD, Chiappe ME, Amrein H, Vosshall LB (2004) Or83b encodes a broadly

expressed odorant receptor essential for Drosophila olfaction. Neuron 43:703–714.

Leal WS, Choo Y, Xu P, Silva CSB, Ueira-vieira C (2013) Differential expression of olfactory genes in the southern

house mosquito and insights into unique odorant receptor gene isoforms. Proc Natl Acad Sci U S A

110:18704–18709.

Li H, You Y, Zhang L (2018) Single Sensillum Recordings for Locust Palp Sensilla Basiconica. JoVE:e57863.

Li J, Haddad R, Santos V, Bavan S, Luetje CW (2015) Receptive range analysis of a mouse odorant receptor

subfamily. J Neurochem 134:47–55.

Lin C, Potter CJ (2015) Re-Classification of Drosophila melanogaster Trichoid and Intermediate Sensilla Using

Fluorescence-Guided Single Sensillum Recording. PLoS One:1–14.

Lombardo F, Salvemini M, Fiorillo C, Nolan T, Zwiebel LJ, Ribeiro JM, Arcà B (2017) Deciphering the olfactory

repertoire of the tiger mosquito Aedes albopictus. BMC Genomics:1–23.

Lu T, Qiu YT, Wang G, Kwon JY, Rutzler M, Kwon H-W, Pitts RJ, van Loon JJA, Takken W, Carlson JR, Zwiebel LJ

(2007) Odor Coding in the Maxillary Palp of the Malaria Vector Mosquito Anopheles gambiae. Curr Biol

17:1533–1544.

Mainland JD, Li YR, Zhou T, Liu WL in. L, Matsunami H (2015) Human olfactory receptor responses to odorants. Sci

data 2:1–9.

Malnic B, Hirono J, Sato T, Buck LB, Hughes H (1999) Combinatorial Receptor Codes for Odors. Cell 96:713–723.

Man O, Willhite DC, Crasto CJ, Shepherd GM, Gilad Y (2007) A Framework for Exploring Functional Variability in

Olfactory Receptor Genes. PLoS One 2:e682.

Marshall B, Warr CG, Bruyne M De (2010) Detection of Volatile Indicators of Illicit Substances by the Olfactory

Receptors of Drosophila melanogaster. Chem Senses 35:613–625.

Montague SA, Mathew D, Carlson JR (2011) Similar Odorants Elicit Different Behavioral and Physiological

Responses, Some Supersustained. J Neurosci 31:7891–7899.

Morel P (2018) Gramm : grammar of graphics plotting in Matlab. J open source Softw 3:1–4.

Münch D, Galizia CG (2016) DoOR 2 . 0 - Comprehensive mapping of Drosophila melanogaster odorant responses.

Sci Rep 6:1–9.

Nagarathnam B, Karpe SD, Harini K, Sankar K, Iftekhar M, Rajesh D, Giji S, Archunan G, Balakrishnan V, Gromiha
MM, Nemoto W, Fukui K, Sowdhamini R (2014) DOR - a Database of Olfactory Receptors - Integrated
Repository for Sequence and Secondary Structural Information of Olfactory Receptors in Selected Eukaryotic
Genomes. Bioinform Biol Insights 8:147–158.

Nakagawa T, Pellegrino M, Sato K, Vosshall LB, Touhara K (2012) Amino acid residues contributing to function of
the heteromeric insect olfactory receptor complex. PLoS One 7:1–9.

Neafsey DE, Waterhouse RM et al. (2015) Highly evolvable malaria vectors: The genomes of 16 Anopheles
mosquitoes. Science (80-) 347:1–8.

Needleman SB, Wunsch CD (1970) A general method applicable to the search for similarities in the amino acid
sequence of two proteins. J Mol Biol 48:443–453.

Nichols AS, Luetje CW (2010) Transmembrane segment 3 of Drosophila melanogaster odorant receptor subunit
85b contributes to ligand-receptor interactions. J Biol Chem 285:11854–11862.

Nishizumi H, Sakano H (2015) Decoding and deorphanizing an olfactory map. Nat Neurosci 18:1432–1433.

Oka Y, Katada S, Omura M, Suwa M, Yoshihara Y, Touhara K (2006) Odorant Receptor Map in the Mouse Olfactory
Bulb: In Vivo Sensitivity and Specificity of Receptor-Defined Glomeruli. Neuron 52:857–869.

Olsen SR, Bhandawat V, Wilson RI (2010) Divisive Normalization in Olfactory Population Codes. Neuron 66:287–
299.

Olsson SB, Hansson BS (2013) Electroantennogram and Single Sensillum Recording in Insect Antennae. In:
Pheromone Signaling: Methods and Protocols, pp 157–177. Totowa, NJ: Humana Press.

Omasits U, Ahrens CH, Müller S, Wollscheid B (2014) Protter: Interactive protein feature visualization and
integration with experimental proteomic data. Bioinformatics 30:884–886.

Raman B, Joseph J, Tang J, Stopfer M (2010) Temporally Diverse Firing Patterns in Olfactory Receptor Neurons
Underlie Spatiotemporal Neural Codes for Odors. J Neurosci 30:1994–2006.

Ramasamy S, Ometto L, Crava CM, Revadi S, Kaur R, Horner DS, Pisani D, Dekker T, Anfora G, Rota-stabelli O (2016)
The Evolution of Olfactory Gene Families in Drosophila and the Genomic Basis of chemical-Ecological
Adaptation in Drosophila suzukii. Genome Biol Evol 8:2297–2311.

Ray A, van Naters WG oe., Carlson JR (2014) Molecular determinants of odorant receptor function in insects. J
Biosci 39:555–563.

Ronderos DS, Lin C-C, Potter CJ, Smith DP (2014) Farnesol-Detecting Olfactory Neurons in Drosophila. J Neurosci
34:3959–3968.

Saberi M, Seyed-Allaei H (2016) Odorant receptors of Drosophila are sensitive to the molecular volume of
odorants. Sci Rep 6:1–11.

Saito H, Chi Q, Zhuang H, Matsunami H, Mainland JD (2009) Odor coding by a Mammalian receptor repertoire. Sci
Signal 2:1–14.

Schmuker M, De Bruyne M, Hähnel M, Schneider G (2007) Predicting olfactory receptor neuron responses from

odorant structure. Chem Cent J 1:1–10.

Spors H, Wachowiak M, Cohen LB, Friedrich RW (2006) Temporal Dynamics and Latency Patterns of Receptor
Neuron Input to the Olfactory Bulb. J Neurosci 26:1247–1259.

Su C, Martelli C, Emonet T, Carlson JR (2011) Temporal coding of odor mixtures in an olfactory receptor neuron.
Proc Natl Acad Sci 108:1–6.

Touhara K, Sengoku S, Inaki K, Tsuboi A, Hirono J, Sato T, Salano H, Haga T (1999) Functional identification and
reconstitution of an odorant. Proc Natl Acad Sci USA 96:4040–4045.

Viklund H, Elofsson A (2008) OCTOPUS: Improving topology prediction by two-track ANN-based preference scores
and an extended topological grammar. Bioinformatics 24:1662–1668.

Wang G, Carey AF, Carlson JR, Zwiebel LJ (2010) Molecular basis of odor coding in the malaria vector mosquito
Anopheles gambiae. Proc Natl Acad Sci 107:4418–4423.

Wicher D, Schäfer R, Bauernfeind R, Stensmyr MC, Heller R, Heinemann SH, Hansson BS (2008) Drosophila odorant
receptors are both ligand-gated and cyclic-nucleotide- activated cation channels. Nature 452:1007–1011.

Wright GA, Thomson MGA, Smith BH (2005) Odour concentration affects odour identity in honeybees. Proc R Soc B
272:2417–2422.

Xu P, Leal WS (2013) Probing insect odorant receptors with their cognate ligands: Insights into structural features.
Biochem Biophys Res Commun 435:477–482.

Yarali A, Ehser S, Hapil FZ, Huang J, Gerber B (2009) Odour intensity learning in fruit flies. Proc R Soc B 276:3413–
3420.

Legends

Fig. 1: Lack of correlation between response distance and sequence similarity among pairs of ORs.

Scatter plots of distances in response vectors versus the amino-acid sequence similarity (a larger value

indicates more similarity), among pairs of ORs, reveal the lack of correlation, in both *D. melanogaster*

(**A1**) and *A. gambiae* (**A2**). Distance is calculated as the L1-distance between the two 110-length

response vectors containing number of spikes for each of the 110 odors. Sequence similarity is

measured as the Needleman-Wunsch alignment score. Each point corresponds to a pair of conspecific

ORs (n = 276 pairs in *D. melanogaster* and n = 1225 pairs in *A. gambiae*).

Fig. 2: Analysis with response determining positions.

A1, **A2**, Scatter plots of distances in 110-odor response vectors versus the sequence similarity measured

at the top 20 positions, among pairs of ORs, show significant negative correlation, in both *D.*

melanogaster (**A1**) and *A. gambiae* (**A2**). Each point corresponds to a pair of conspecific ORs (n = 276

pairs in *D. melanogaster* and n = 1225 pairs in *A. gambiae*).

B1, **B2**, Bars indicate the negative correlation between the distance in response vectors and the

sequence similarity measured using the top N positions, plotted as a function of N in the range of 10 to

100. Note that the correlation values are relatively stable in both the species in the range of 10-40 and

decrease gradually in magnitude for larger values of N.

C1, **C2**, Bars indicate the number of top 20 positions in *D. melanogaster* (**C1**) or *A. gambiae* (**C2**) retained

among the top 20 positions when the positions were recalculated after removing different numbers of

receptors from the dataset. For each number, all possible combinations of that many receptors were

removed and the average number of positions preserved among the top 20 is reported. Error bars

represent s.e.m. over all combinations for each number.

Fig. 3: Reliability of responses predicted using the top 20 positions.

A, Scatter plots of the actual and the predicted odor responses for three representative ORs each from

D. melanogaster and *A. gambiae*. The correlation coefficients between the two responses are indicated.

The responses were spread around the y=x line (dashed line, corresponding to ideal predictions).

B1, B2, Violin plots showing the correlation coefficient between the predicted and the actual response

vectors of ORs in *D. melanogaster* **(B1)** or *A. gambiae* (n = 50) **(B2)**. Each point corresponds to one OR (n

= 24 ORs in *D. melanogaster* and n = 50 ORs in *A. gambiae*). Note that the correlations are mostly

positive. Statistically significant positive correlations (with P less than 0.05) are shown with black

outlines.

C1, C2, Violin plots showing the correlation coefficient between the control predictions and the actual

response vectors of ORs in *D. melanogaster* **(C1)** or *A. gambiae* (n = 50) **(C2)**. Note that the correlations

are close to zero in most cases.

D1, D2, Color maps showing the relative prediction error for each OR-odor pair, for the set of 110 odors,

and all 24 ORs in *D. melanogaster* **(D1)** or all 50 ORs in *A. gambiae* **(D2)**. The relative prediction error is

measured as $D_{Pred,Actual}$ - $D_{Ctrl,Actual}$, where $D_{Pred,Actual}$ is the absolute distance between the

predicted and actual response, and $D_{Ctrl,Actual}$ is the absolute distance between the control prediction

and the actual response. The abundance of negative values (red shades) indicates that the predictions

are usually better than the control. Column averages and row averages of the matrix values are shown

on top and right, respectively.

E1, E2, Violin plots shows the average (over odors) of $D_{Pred,Actual}$ - $D_{Ctrl,Actual}$ for each of the 24 ORs in

D. melanogaster **(E1)** or 50 ORs in *A. gambiae* **(E2)**.

In violin plots, horizontal line and the error bar indicate the mean and the s.e.m., respectively, within a

species.

Fig. 4: Response prediction on an independent set of ORs.

A, Color map showing the relative prediction error (measured as $D_{Pred,Actual}$ - $D_{Ctrl,Actual}$) for each OR-

odor pair, for the set of 68 odors and 26 novel ORs. Pairs whose values were not available are shown in

grey. Note the high frequency of negative values (red) and very low frequency of positive values (blue),

indicating that the predictions were closer to the actual value than the control were in most cases.

B, Violin plots shows the average (over odors) of $D_{Pred,Actual}$ - $D_{Ctrl,Actual}$ for each of the 26 ORs.

Horizontal line and the error bar indicate the mean and the s.e.m., respectively.

C, $D_{Pred,Actual}$ is plotted as a function of the number of ORs with known responses used in making the

predictions. These different numbers of ORs were sampled from the original set of 24 ORs with known

responses. For each number, the sampling was performed 10 times; for each sampling, selection of

response-determining positions and prediction of responses was performed.

D, For each number of sampled ORs, we calculated the fraction of the 506 OR-odor pairs (non-grey

values in panel **A**) for which the predicted responses were closer to the actual values than the control

predictions were. Both methods of generating control predictions (see **Methods**) were compared.

In panels **C** and **D**, errors bars indicate s.e.m. over the 10 samplings for each number from 3 to 21; for

24, all available ORs were used.

Fig. 5: Localization of the top 20 response-determining positions.

Protter plots showing the localization of the top 20 response-determining residues in representative

ORs: OR85b of *D. melanogaster* **(A1)** and OR1 of *A. gambiae* **(A2)**. Residues in the top 20 positions in the

corresponding species are shown with shaded backgrounds, while the residues that are among the top

20 positions in both the species are shown with a black outline. One of the top 20 positions in *A.*

gambiae mapped to a gap in OR1 in the multiple sequence alignment, so only 19 positions are shaded.

Table 1: Positions identified in previous mutagenesis studies that match the top positions identified in

our analysis.

Of the 47 mutations that we collected from the mutagenesis studies, the 9 that matched the top

positions in *D. melanogaster* or *A. gambiae* are listed. The complete set of the 47 mutations is shown in

Table S1. TM: transmembrane helix; EL: extracellular loop.

Supplementary Table S1: Complete multiple sequence alignment of all the ORs.

The table shows the multiple sequence alignment of all the ORs that have been used in this study. Each

row corresponds to one OR. The first two rows indicate with colored background the positions that were

ranked in the two species; among these the ranks of the top 20 positions are mentioned. The 47

residues that have been identified as important in previous mutagenesis studies are marked with yellow

color in the matrix.

Table 1

Study	Mutated receptor	Mutated residue	Position in multiple sequence alignment	Rank among top positions	Topological position
Hughes et al, 2014	*A. gambiae* OR15	F369L	783	2 in *D. melanogaster* & 17 in *A. gambiae*	TM 7
Nakagawa et al 2012	*B. mori* OR1	Y109A	234	14 in *D. melanogaster*	TM 3
Nichols and Luetje, 2010	*D. melanogaster* OR85b	C146S	335	20 in *D. melanogaster*	TM 3
Ray et al, 2014	*A. gambiae* OR1	Y405H	791	6 in *A. gambiae*	TM 7
Ray et al, 2014	*A. gambiae* OR1	S406A	792	20 in *A. gambiae*	TM 7
Xu and Leal, 2013	*B. mori* OR1	P165A	335	20 in *D. melanogaster*	TM 3
Xu and Leal, 2013	*B. mori* OR1	P194A	398	9 in *A. gambiae*	EL 2
Xu and Leal, 2013	*A. gambiae* OR10	P154A	398	9 in *A. gambiae*	EL 2
Xu and Leal, 2013	*C. quinquefasciatus* OR10	P157A	398	9 in *A. gambiae*	EL 2

Figure 1

A1

A2

Distance between response vectors of two ORs (spikes)

D . melanogaster

A. gambiae

Sequence similarity between two ORs (alignment score)

Figure 2

Figure 3

Figure 4

Figure 5

A1

OR85b

● Top 20 positions in *D. melanogaster*
○ Among top 20 positions in both species

A2

OR1

● Top 20 positions in *A. gambiae*
○ Among top 20 positions in both species

Supplementary Figure S1

Supplementary Fig. S1: Cross-validation of correlation between response similarity and sequence similarity at the top 20 positions.

Scatter plots of distances in response vectors (using half of the odors) versus the sequence similarity measured at the top 20 positions, among pairs of ORs. This figure is similar to Fig. 2A1,2, except that here only half of the odors (in odd-numbered rows of the original dataset) were used to calculate the top 20 positions, and the remaining half were used to calculate the response similarity, thus providing a cross-validation. The observed Pearson correlations are -0.56 in both *D. melanogaster* (**A1**), and -0.31 in *A. gambiae* (**A2**). Each point in the plots corresponds to a pair of conspecific ORs (n = 276 pairs in *D. melanogaster* and n = 1225 pairs in *A. gambiae*).

Supplementary Figure S2

Supplementary Fig. S2: Predictions in larval *Drosophila*.

A1, A2 Color maps showing the relative prediction error for each OR-odor pair, for 21 ORs in *D. melanogaster* larvae for the set of 26 odors, at the dilution of 10^{-2} **(A1)** or 10^{-4} **(A2)**. The relative prediction error is measured as $D_{Pred,Actual}$ - $D_{Ctrl,Actual}$, as described in **Fig. 3**.

B1, B2 Violin plots shows the average (over odors) of $D_{Pred,Actual}$ - $D_{Ctrl,Actual}$ for each of the 21 ORs in panels **A1** and **A2,** respectively. Horizontal line and the error bar indicate the mean and the s.e.m., respectively.