Phosphatidylcholine biosynthesis in Mitis group streptococci via host metabolite scavenging

Luke R. Joyce[a], Ziqiang Guan[b]#, and Kelli L. Palmer[a]#

[a]Department of Biological Sciences, The University of Texas at Dallas, Richardson, Texas, USA

[b]Department of Biochemistry, Duke University Medical Center, Durham, NC[2]

Running title: Phosphatidylcholine biosynthesis in streptococci

#Corresponding authors:

Ziqiang Guan: Ziqiang.Guan@duke.edu

Kelli Palmer: Kelli.Palmer@utdallas.edu

Abstract

The Mitis group streptococci include the major human pathogen *Streptococcus pneumoniae* and the opportunistic pathogens *S. mitis* and *S. oralis* which are human oral cavity colonizers and agents of bacteremia and infective endocarditis in immunocompromised patients. Bacterial membrane lipids play crucial roles in microbe-host interactions, yet for many pathogens, the composition of the membrane is poorly understood. In this study, we characterized the lipidomes of selected species of Mitis group streptococci and investigated the mechanistic basis for biosynthesis of the phospholipid phosphatidylcholine (PC). PC is a major lipid in eukaryotic cellular membranes, but it is considered to be comparatively rare in bacterial taxa. Using liquid chromatography/mass spectrometry (LC/MS) in conjunction with stable isotope tracing, we determined that Mitis group streptococci synthesize PC via the rare host metabolite scavenging pathway, the glycerophosphocholine (GPC) pathway, which is largely uncharacterized in bacteria. Our work demonstrates that Mitis group streptococci including *S. pneumoniae* remodel their membrane in response to the major human metabolites GPC and lysoPC.

Importance

We lack fundamental information about the composition of the cellular membrane even for the best studied pathogens of critical significance for human health. The Mitis group streptococci are closely linked to humans in health and disease, yet their membrane biology is poorly understood. Here, we demonstrate that these streptococci scavenge major human metabolites and use them to synthesize the membrane phospholipid phosphatidylcholine. Our work is significant because it identifies a mechanism by which the major human pathogen *S. pneumoniae* and the primary human oral colonizers *S. mitis* and *S. oralis* remodel their membrane in response to host metabolites.

Introduction

The Mitis group streptococci are Gram-positive bacteria that natively inhabit the human oral cavity, nasopharynx, and gastrointestinal tract (1). They include the species *S. mitis* and *S. oralis*, which are among the first colonizers of the human oral cavity from birth and facilitate host-microbe-microbe interactions by creating anchors for biofilm formation with other oral microbiota (2, 3). *S. mitis* and *S. oralis* are also opportunistic pathogens causing bacteremia and infective endocarditis (4–7). The Mitis group streptococci also include the major human pathogen, *S. pneumoniae. S. pneumoniae* has >99% 16S rRNA sequence identity (8–10), exchanges capsule biosynthesis and antibiotic resistance genes (11, 12), and shows antibody cross-reactivity (13) with *S. mitis* and *S. oralis.*

We, and others, recently reported that certain Mitis group streptococci have unusual membrane physiology in that they can proliferate while lacking the major anionic phospholipids phosphatidylglycerol (PG) and cardiolipin (CL) (14, 15). More specifically, *S. mitis* and *S. oralis* can tolerate deletion or inactivation of *cdsA*, the gene encoding phosphatidate cytidylyltransferase (14, 15). CdsA catalyzes synthesis of cytidine diphosphate-diacylglycerol (CDP-DAG), a key intermediate in the synthesis of major phospholipids (Fig 1). Deletion or inactivation of *cdsA* in Mitis group streptococci, and the corresponding loss of PG and CL, confers high-level (MIC >256 μg/mL) resistance to daptomycin, a last-line lipopeptide antibiotic (14–16).

Unexpectedly, lipidomic analysis of *S. mitis* and *S. oralis* by normal phase liquid

chromatography-electrospray ionization/mass spectrometry (NPLC-ESI/MS) revealed

the presence of phosphatidylcholine (PC) in wild-type strains and in *cdsA*-null mutants

(14). To our knowledge, PC in streptococci had not been previously described. Overall,

the lipid compositions of streptococci are under-studied and poorly characterized.

Previous studies analyzing lipids of streptococci primarily used thin layer

chromatography, whose limitations in analytical sensitivity and molecular specificity

prohibit comprehensive lipidomic identification; these studies did not detect PC (17–22).

PC is a biologically significant lipid. As a zwitterionic phospholipid, PC promotes bilayer

formation (23), reduces the rate of protein folding to allow for correct protein

configurations (24, 25), aids in resistance to antimicrobials targeting prokaryotic

membranes (26), aids in surviving environmental fluctuations such as temperature shifts

(27), and is a major component of eukaryotic membranes. There is evidence that PC

plays important roles in host-microbe interactions. *Legionella* lacking functional PC

biosynthesis exhibit decreased virulence because of poor recognition by host

macrophages and reduced motility (28). *Brucella abortus* and *Agrobacterium*

tumefaciens also exhibit diminished virulence when PC biosynthesis is inactivated (29–

31). Alternatively, *Pseudomonas aeruginosa* strains lacking PC show no detectable

alteration in virulence (32).

Because PC may impact how Mitis group streptococci interact with the human host, in

this study we investigated the mechanism for PC biosynthesis in these organisms.

There are four experimentally confirmed PC biosynthesis pathways in bacteria (Fig 1),

two of which are widespread and well-characterized, namely the
phosphatidylethanolamine (PE) methylation pathway (23, 33), and the
phosphatidylcholine synthase (Pcs) pathway (34, 35). The PE methylation pathway
uses PE as a starting substrate, which is methylated via phospholipid *N*-
methyltransferase (Pmt) in three subsequent steps to form monomethylPE (MMPE),
dimethylPE (DMPE), and finally PC, using *S*-adenosylmethionine as the methyl group
donor (33). The PE methylation pathway is utilized by mammalian liver cells (23, 36)
and bacteria including *Rhodobacter sphaeroides* and *Sinorhizobium meliloti* (23). The
Pcs pathway is exclusive to prokaryotes and is a one-step reaction in which a
phosphatidylcholine synthase (Pcs) enzyme condenses choline with CDP-DAG to form
PC (34). The presence of either *pmt* or *pcs* genes has been used to identify bacterial
taxa likely to produce PC (23).

A third pathway, the CDP-choline pathway (referred to as the Kennedy pathway in
eukaryotes), was recently identified in the Gram-negative human oral colonizer,
Treponema denticola (37). In this pathway, choline is scavenged from the environment
and activated to CDP-choline via the LicAC enzymes. Many host-associated bacteria
possess LicAC and utilize host-derived choline to decorate a wide range of extracellular
structures (23) including the Type IV lipoteichoic acid (LTA) of *S. pneumoniae, S. mitis*,
and *S. oralis* (38–42). In the CDP-choline pathway, CDP-choline is condensed with
diacylglycerol (DAG) by a 1,2-diacylglycerol choline phosphotransferase (CPT) to form
PC.

A fourth pathway, the glycerophosphocholine (GPC) pathway, has been reported for

only two organisms: the Gram-negative plant pathogen *Xanthomonas campestris* (43)

and yeast (44). In eukaryotic cells, GPC is a breakdown product of choline-containing

membrane phospholipids. Yeast can utilize GPC as the source for glycerol-3-

phosphate, choline, or phosphate, depending on the environmental conditions (45).

GPC is a major human metabolite present in saliva and blood (46, 47). In the GPC

pathway, GPC is scavenged from the environment and acylated twice to form the

intermediate lysoPC and then PC. The genetics underlying the GPC pathway in *X.*

campestris have not been fully elucidated. Moser *et al.* identified two *X. campestris*

acyltransferases that performed the second acylation event from lysoPC to PC (43).

Yeast possess a fully elucidated GPC pathway (44, 48).

Here we use NPLC-ESI/MS and other biochemical and genetic approaches to

investigate PC biosynthesis in Mitis group streptococci using type strains and an

infective endocarditis isolate (Table 1). We determined that these organisms synthesize

PC by the rare GPC pathway via scavenging the host metabolites GPC and lysoPC.

Results

PC biosynthesis by model *S. mitis* and *S. oralis* strains used in this study. We

previously reported lipidomic analysis by NPLC-ESI/MS of three Mitis group infective

endocarditis isolates cultured in the rich, undefined laboratory growth medium Todd

Hewitt broth (THB). We determined that these organisms possess PC in their

membranes ((14) and Table 1). We confirmed these results using the *S. mitis* and *S.*

oralis type strains ATCC 49456 and ATCC 35037, respectively (Table 1).

In this study, we used one of the infective endocarditis isolates, referred to as SM43, for

most of our mechanistic studies of PC biosynthesis. Phylogenetic assignments within

the Mitis group are difficult due to variable phenotypes and highly conserved 16S rRNA

sequences (8, 9). SM43 was initially assigned to the *S. mitis* species using standard

biochemical techniques (49). We analyzed a complete SM43 genome sequence using

an average nucleotide identity (ANI) calculator (50, 51). ANI values are used for

molecular species definitions (52). Two bacterial strains with an ANI of ≥95% in their

shared genes are considered to be the same species and with an ANI of >70% are

considered to be the same genus (53, 54). SM43 possesses 94.3% ANI with *S. oralis*

ATCC 35037 and 86.9% ANI with *S. mitis* ATCC 49456. Based on these data showing a

close phylogenetic relationship between SM43 and *S. oralis*, we refer to SM43 as a

Mitis group *Streptococcus* in this study (Table 1).

PC biosynthesis in SM43 is not by the PE methylation or Pcs pathway. The well-

characterized PE methylation pathway for PC biosynthesis is catalyzed by the enzyme

Pmt (Fig 1). This pathway is excluded as SM43 does not synthesize PE (14), nor does it

encode *pmt*.

The Pcs pathway requires CDP-DAG and choline as substrates (Fig 1). *cdsA* loss-of-

function mutants arising spontaneously under daptomycin selection in Mitis group

streptococci do not synthesize CDP-DAG but still synthesize PC (14). These results

indicate that PC biosynthesis in Mitis group streptococci is by a CDP-DAG-independent

pathway. To confirm these results, we generated a *cdsA* deletion in SM43. PC was

detected in both wild-type and Δ*cdsA* SM43 cultured in THB. Fig 2a shows the relative

abundance of PC in the SM43 membrane during mid-exponential phase. The ESI/MS

spectra of the major PC species, PC (30:1) at *m/z* 704, PC (32:1) at *m/z* 732, PC (34:1)

at *m/z* 760, and PC (36:2) at *m/z* 786 are shown for SM43 (Fig 2b) and *S. mitis* type

strain ATCC 49456 (Fig S1a). The chemical structure of PC (34:1) shown in Fig 2c is

supported by MS/MS (Fig S1b). The ESI/MS spectra of PC in SM43Δ*cdsA* is shown in

Fig 2d. Based on these results and the absence of a *pcs* ortholog in the SM43 and

ATCC 49456 genomes (Fig 1), the *pcs* pathway is excluded.

SM43 encodes a partial CDP-choline pathway. The CDP-choline pathway requires

activation of choline to CDP-choline by the LicA and LicC enzymes. This is followed by

condensation of CDP-choline with DAG by a 1,2-diacylglycerol choline

phosphotransferase (CPT) enzyme to form PC (Fig 1). *S. mitis* and *S. oralis* encode

licABC for the activation of exogenous choline to CDP-choline, which is required for

choline decoration of Type IV LTA in these organisms and *S. pneumoniae* (38–42).

Given that SM43 encodes *licABC*, we assessed the possibility that the CDP-choline

pathway is used for PC biosynthesis in this organism.

The CPT of *T. denticola* possesses a CDP-alcohol phosphatidyltransferase domain

(Superfamily accession: cl00453) (37). Only one SM43 predicted protein,

phosphatidylglycerophosphate synthase (PgsA), possesses this domain. PgsA

catalyzes the addition of glycerolphosphate to CDP-DAG to form

phosphatidylglycerophosphate (PG-P) (55) which is required for subsequent PG and CL

synthesis (Fig 1).

To investigate if PgsA has CPT activity in SM43, a 284 bp region of *pgsA* encoding the

catalytic domain was replaced with an erythromycin resistance cassette. The *pgsA*

mutant has a significant growth defect, with a doubling time almost twice that of the wild

type (Fig S2a). The defect is likely due to PgsA interaction with RodZ in membrane

homeostasis, as has been reported for *Bacillus subtilis* (56). Similar to SM43Δ*cdsA*, PC

is present in SM43Δ*pgsA* cultured in THB (Fig 2e). As expected, the SM43Δ*pgsA*

mutant lacks PG and CL (Table 1) and is high-level daptomycin-resistant (MIC >256

µg/mL), in agreement with experiments by Tran *et al*. utilizing *S. mitis-oralis* strains (57).

We conclude that the sole candidate for CPT activity in SM43, PgsA, does not catalyze

PC biosynthesis.

To conclusively exclude the CDP-choline pathway, we performed stable isotope-

labeling experiments. SM43 was cultured in THB supplemented with 2 mM deuterated

choline (choline-*d9*), in which all nine hydrogen (1 Da) atoms on the three methyl

groups are replaced with deuterium (2 Da) atoms, thereby increasing the mass of

choline by 9 Da. If SM43 utilizes the CDP-choline pathway for PC biosynthesis, an *m/z*

shift of 9 Da would be observed for CDP-choline and PC species in choline-*d9-*

supplemented SM43 cultures. We observed choline-*d9* incorporation into CDP-choline

via a shift of its M$^+$ ion from *m/z* 489 (Fig 3a; no choline-*d9* supplementation) to *m/z* 498

(Fig 3b, with choline-*d9* supplementation). The identification of CDP-choline and CDP-

choline-*d9* was confirmed by MS/MS (Fig S3). In contrast, no *m/z* shift was observed for

PC species between control (Fig 3c) and choline-*d9*-supplemented (Fig 3d) cultures.

These data definitively eliminate the CDP-choline pathway as the SM43 PC

biosynthesis pathway.

SM43 utilizes the GPC scavenge pathway for PC biosynthesis. GPC is a major

human metabolite, present in blood and saliva at concentrations up to 40 µM and 10 µM

respectively (46). Using reverse-phase LC/MS, we detected GPC (*m/z* 258.1) in THB

(Fig S4), likely originating from the heart infusion component of the medium. Therefore,

GPC is available for scavenge in the medium used for routine SM43 cultures.

To determine whether SM43 utilizes the GPC pathway for PC biosynthesis, we used

stable isotope-labeled GPC to trace the conversion of GPC into PC. SM43 was cultured

in THB with and without 0.13 mM GPC-*d9* supplementation. A *m/z* shift of 9 Da was

observed for all PC species (Fig 4a), demonstrating that SM43 uses the GPC pathway

for PC biosynthesis. To further confirm this result, SM43 and *S. mitis* ATCC 49456 were

cultured in a chemically defined medium containing 0.5 mM choline (DM) (58, 59), with

and without 0.13 mM GPC supplementation. SM43 and ATCC 49456 synthesize PC

only when GPC is present in the defined medium (Fig 5 and Table 2). In summary,

GPC-*d9* isotope tracking and defined medium experiments independently confirm that

SM43 utilizes the GPC pathway for PC biosynthesis. Moreover, these results are not

strain-specific, as the *S. mitis* type strain also synthesizes PC only when GPC is present

in the growth environment.

PC is not an essential component in the lipid membrane for SM43 and ATCC 49456,

evidenced by their abilities to grow in DM lacking GPC. To assess the impact that PC

has on growth dynamics, SM43 and ATCC 49456 were cultured in THB and in DM with

and without GPC supplementation. GPC presence or absence had no impact on growth

(Fig S2b,c).

Since lysoPC is an intermediate in the GPC pathway, we hypothesized that SM43 could

also scavenge lysoPC from its environment. LysoPC is present in human blood at up to

200 µM (60). We supplemented THB with lysoPC (20:0), which has an acyl chain that is

not usually observed in bacterial membranes. SM43 readily scavenged exogenous

lysoPC (20:0) and acylated it to form PC (Fig 4b). The PG species in the SM43

membrane remain unchanged in the presence of lysoPC (20:0), indicating low

transacylation activity (Fig S5). We conclude that SM43 scavenges both GPC and

lysoPC from its environment to synthesize PC (Fig 6).

GPC-dependent PC synthesis by *S. pneumoniae and S. oralis*. *S. pneumoniae* is a

major human pathogen and a close relative of *S. mitis*. Surprisingly, there are only a few

reports on lipid analysis of *S. pneumoniae* (17–19), for which thin layer chromatography

was used as the analytical technique. We applied LC-ESI/MS, with much higher

sensitivity and specificity, to investigate the lipidome of *S. pneumoniae*. PC is present in

the membranes of *S. pneumoniae* D39 (Fig 7a,b) and TIGR4 (Fig 7c,d) cultured in THB

supplemented with yeast extract (also see Table 1). Fig 7a and c shows the positive ion

TIC and the relative abundance of PC in the D39 and TIGR4, respectively, in early

stationary phase. The ESI/MS spectra of the major PC species, PC (30:1) at m/z 704,

PC (32:1) at m/z 732, PC (34:2) at m/z 758, and PC(36:2) at m/z 786 are shown for D39

(Fig 7b) and TIGR4 (Fig 7d). We identified PC presence in infective endocarditis *S.*

oralis isolates 1647 and 1648 in a previous study (14).

To determine if *S. pneumoniae* and *S. oralis* utilize the GPC pathway for PC

biosynthesis, *S. pneumoniae* D39 and *S. oralis* ATCC 35037 were cultured in defined

media with and without GPC supplementation. When GPC is not available, *S.*

pneumoniae and *S. oralis* do not synthesize PC (Fig 8a,c). PC is only present when

GPC is available in the medium (Fig 8b,d). We conclude that *S. pneumoniae* and *S.*

oralis also scavenge GPC to synthesize PC.

Discussion

The lipid membrane is a dynamic site of interaction between microbial pathogens and

their hosts. However, for many pathogens, the composition of the membrane is poorly

understood. In this study, we characterized the lipidomes of selected species of Mitis

group streptococci and investigated the mechanistic basis for biosynthesis of the

phospholipid PC. We found that Mitis group streptococci remodel their membrane lipid

compositions in response to the host metabolites GPC and lysoPC. To our knowledge,

this is the first description of PC in *S. pneumoniae,* a major human pathogen which has

been studied for over a century and yet its membrane lipid composition remains poorly

understood. There have been very few lipidomic studies performed in *S. pneumoniae*

(17–19), and little is known about how *S. pneumoniae* remodels its membrane in

response to changing environments inside and outside the host. Here, we reported the

first identification of PC in *S. pneumoniae* and demonstrated that *S. pneumoniae*

synthesizes PC only when GPC, a major human metabolite, is available for scavenge.

Very little is known about the GPC in bacteria; however, a complete GPC pathway has

been characterized in yeast (44, 48). It includes a dual substrate transporter, Git1, for

the uptake of glycerophosphoinositol and GPC; a GPC acyltransferase referred to as

Gpc1; and the acyltransferase Ale1 performing lysoPC acylation. To fully elucidate the

GPC pathway in streptococci, identification of transporters for GPC and lysoPC as well

as the acyltransferases are required. However, no orthologs of the yeast GPC pathway

components were identified in the SM43 genome. There are two acyltransferases

annotated in the *S. mitis* genome, PlsY and PlsC, which are responsible for

phosphatidic acid biosynthesis in other organisms and may play a role in acylation of

either GPC or lysoPC. In *Mycobacterium tuberculosis,* a sugar-binding ABC transporter,

UgpABCE, transports GPC (61). The UgpB substrate binding domain is flexible in

substrate affinity, binding to phosphate-containing substrates such as *sn*-glycerol 3-

phopshate, glycerol 2-phosphate, and GPC (61–63). Given the reduced size of

streptococcal genomes, it is possible that GPC uptake is also conferred by a transport

system with flexible substrate specificity.

Is PC biosynthesis important for Mitis group streptococcal virulence? We expect that PC levels in the membrane will impact membrane charge, for example, which could in turn impact biofilm formation and interactions with the host immune system. Moreover, GPC and lysoPC levels vary at different sites within the human body and in health and disease (≤ 500 µM and ≤ 200 µM, respectively (46, 60)), which could impact the relative ratio of the zwitterionic PC to the anionic phospholipids PG and CL in Mitis group streptococci colonizing these sites. Due to our limited understanding of the genes underlying the GPC pathway, at present our only method to control this pathway is by altering the *in vitro* growth medium. For this reason *in vivo* studies are not currently feasible. However, by culturing Mitis group streptococci in defined media with and without GPC, in future studies we can assess the biophysical impact of PC on streptococcal membranes in terms of lipid/microdomain organization, charge, rigidity, and protein composition, which would be informative from a basic science perspective.

Overall, our work highlights the importance of utilizing laboratory culture media that mimic the *in vivo* nutritional environments where pathogens are found. Our identification of PC in the membrane of Mitis group streptococci including the major human pathogen *S. pneumoniae*, and their utilization of the rare GPC pathway, justifies further investigation into streptococcal membrane biology, about which little is known.

Materials and methods

Bacterial strains, media, and growth conditions. Strains and plasmids used in this study are shown in Table S1. Streptococcal strains were grown in THB in 5% CO_2 at 37°C unless otherwise stated. *S. pneumoniae* THB cultures were supplemented with 0.5% yeast extract. Streptococcal chemically defined medium (58) was diluted from stock as described (59) supplemented with 0.5 mM choline (referred to as DM), slightly modified from (64) unless otherwise stated. Media were supplemented with 130 µM glycerophosphocholine (Sigma-Aldrich) where stated. Erythromycin was used at 20 µg/mL for SM43. Daptomycin susceptibilities were assessed using daptomycin Etest strips (bioMérieux) on Mueller Hinton Agar plates, per CLSI standards (65).

Genome sequencing and assembly. Genomic DNA was isolated using the Qiagen DNeasy Blood and Tissue kit per the manufacturer's protocol with the exception that cells were pre-treated with 180 µL 50 mg/mL lysozyme, 25 µL 2500 U/mL mutanolysin, and 15 µL 20 mg/mL pre-boiled RNase A and incubated at 37°C for 2 h. Pacific Biosciences single molecule real time (SMRT) sequencing was performed by the Johns Hopkins Genome Core. The SM43 whole genome was assembled using the Unicyler assembly pipeline (66) combining SMRT long reads generated in this study and Illumina reads we previously generated for SM43 (accession PRJNA354070; (14)). Sequencing of SM43Δ*pgsA* was performed by Molecular Research DNA (Shallowater, Texas) using Illumina Hi-Seq 2x150 paired end reads.

Gene deletions in SM43. Primers used in this study are shown in Table S2. The SM43 *cdsA* deletion construct was designed essentially as previously described (67, 68). See

Text S1. For *pgsA* deletion, the protocol was the same as for *cdsA* deletion, except an

erythromycin resistance marker with its native promoter was amplified from pG⁺host4

(69) and inserted between the homologous flanking arms to allow for selection-based

screening of putative *pgsA* mutants. The resulting construct deleted 284 bp encoding

the catalytically active site of *pgsA* and replaced them with the erythromycin resistance

marker. The replacement of *pgsA* was confirmed by whole genome sequencing.

Natural transformation. The protocol was as previously described (68) with minor

modifications. Briefly, 100 µL exponential phase pre-cultures (OD_{600} ~0.5) in THB were

frozen with an equal volume of 10% glycerol. Precultures were thawed at room

temperature and diluted in 900 µL THB. This was further diluted 1:50 in pre-warmed 5

mL THB and incubated at 37°C for 45 minutes. 500 µL culture was aliquoted along with

1 µL of 1 mg/mL competence stimulating peptide (DWRISETIRNLIFPRRK) and 1 µg/mL

linear DNA construct. Transformation reactions were cultured for 2 h at 37°C in

microcentrifuge tubes before plating on THB agar with selection as appropriate.

Lipidomics. Unless otherwise noted, lipidomics analyses were performed on overnight

cultures in stationary phase. Centrifugation was performed using a Sorvall RC6+

centrifuge. Cultures were pelleted at 4,280 x *g* for 5 min at room temperature. The

supernatants were removed and stored at -80°C until acidic Bligh-Dyer lipid extractions

were performed as described (70) with minor modifications. See Text S1.

Normal-phase LC was performed on an Agilent 1200 quaternary LC system equipped with an Ascentis silica high-performance liquid chromatography (HPLC) column (5m; 25 cm by 2.1 mm; Sigma-Aldrich) as described previously (70, 71). Data analysis was performed using Analyst TF1.5 software (Sciex, Framingham, MA). See Text S1.

Metabolite extractions. Cultures were pelleted at 4,280 x g in a Sorvall RC 6+ floor centrifuge at room temperature, washed once with 1X PBS, and transferred to 1.5 mL microcentrifuge tubes. Cells were pelleted and frozen at -80°C until use. Metabolite extraction was performed as previously described (72) with minor modifications. See Text S1.

Reverse phase LC-ESI/MS analysis. RPLC-ESI/MS analysis of water-soluble metabolites was performed using a Shimadzu LC system (comprising a solvent degasser, two LC-10A pumps and a SCL-10A system controller) coupled to a TripleTOF5600 mass spectrometer (Sciex, Framingham, MA). See Text S1 for detailed information of LC flow rate and mass spectrometer settings. Data acquisition and analysis were performed using the Analyst TF1.5 software (Sciex, Framingham, MA).

Deuterated isotope (*d9*) and lysoPC tracking. Deuterated (*d9*) isotope tracking was performed by addition of 2 mM choline-*d9* (Sigma-Aldrich) in 50 mL THB. 3.7 mM GPC-*d9* (Toronto Research Chemicals) was added to 15 mL THB, to yield a final concentration of 130 µM. Cultures were grown in 5% CO_2 at 37°C for 18 h before 10 mL of choline-*d9* culture or 5 mL of GPC-*d9* culture was removed for metabolite extraction

and the remaining culture was pelleted for lipid extraction. LysoPC (20:0) was obtained

from Avanti Polar Lipids. Cultures were supplemented with 2 mg lysoPC per 15 mL

THB, unless otherwise stated, and incubated overnight in 5% CO_2 at 37°C. Lipid

extractions were performed as described above.

Growth curves. Individual colonies were incubated overnight in 5 mL DM. Cultures

were diluted to a starting OD_{600nm} of 0.05 in pre-warmed 20 mL DM, DM with 130 μM

GPC, and THB. OD_{600nm} was monitored every hour using a Thermo Scientific Genesys

30 spectrophotometer. Growth curves were performed in biological triplicates for each

strain.

ANI analysis. The ANI calculator (50, 51) was used with default parameters to analyze

the following genomes: SM43, *S. oralis* ATCC 35037 (PRJNA38733) and *S. mitis* ATCC

49456 (PRJNA173).

Accession numbers. The SM43 whole genome sequence generated in this study has

been deposited in Genbank under the accession number CP040231. Illumina and

SMRT sequence reads generated in this study have been deposited in the Sequence

Read Archive under the accession number PRJNA542100.

Acknowledgments

This work was supported by R01AI116610, R21AI130666, and the Cecil H. and Ida

Green Chair in Systems Biology Science to K.P. and GM069338 and EY023666 to Z.G.

We gratefully acknowledge Dr. Michael Federle and Jennifer Chang at the University of

Illinois at Chicago for providing *S. pneumoniae* cell pellets from THB+Y for our original

analysis and for hosting L.J. for *S. pneumoniae* culture work.

References

1. Doern CD, Burnham CAD. 2010. It's not easy being green: The Viridans Group Streptococci, with a focus on pediatric clinical manifestations. J Clin Microbiol 48:3829–3835.

2. Mitchell J. 2011. *Streptococcus mitis*: Walking the line between commensalism and pathogenesis. Mol Oral Microbiol 26:89–98.

3. Kolenbrander PE, London J. 1993. Adhere today, here tomorrow: Oral bacterial adherence. J Bacteriol 175:3247–3252.

4. Murdoch DR, Corey GR, Hoen B, Miró JM, Fowler VG, Bayer AS, Karchmer AW, Olaison L, Pappas PA, Moreillon P, Chambers ST, Chu VH, Falcó V, Holland DJ, Jones P, Klein JL, Raymond NJ, Read KM, Tripodi MF, Utili R, Wang A, Woods CW, Cabell CH, Investigators TI. 2009. Clinical presentation, Etiology and Outcome of Infective Endocarditis in the 21st Century: The International Collaboration on Endocarditis-Prospective Cohort Study. Arch Intern Med 169:463–473.

5. Marín M, Gudiol C, Garcia-Vidal C, Ardanuy C, Carratalà J. 2014. Bloodstream infections in patients with solid tumors: Epidemiology, antibiotic therapy, and outcomes in 528 episodes in a single cancer center. Medicine (Baltimore) 93:143–149.

6. Gudiol C, Bodro M, Simonetti A, Tubau F, González-Barca E, Cisnal M, Domingo-Domenech E, Jiménez L, Carratalà J. 2013. Changing aetiology, clinical features,

437 antimicrobial resistance, and outcomes of bloodstream infection in neutropenic

438 cancer patients. Clin Microbiol Infect 19:474–479.

439 7. Shelburne SA, Sahasrabhojane P, Saldana M, Yao H, Su X, Horstmann N,

440 Thompson E, Flores AR. 2014. *Streptococcus mitis* strains causing severe clinical

441 disease in cancer patients. Emerg Infect Dis 20:762–771.

442 8. Kawamura Y, Hou X-G, Sultana F, Miura H, Ezaki T. 1995. Determination of 16S

443 rRNA Sequences of *Streptococcus mitis* and *Streptococcus gordonii* and

444 Phylogenetic Relationships among Members of the Genus *Streptococcus*. Int J

445 Syst Bacteriol 45:406–408.

446 9. Scholz CFP, Poulsen K, Kilian M. 2012. Novel Molecular Method for Identification

447 of *Streptococcus pneumoniae* Applicable to Clinical Microbiology and 16S rRNA

448 Sequence-Based Microbiome Studies. J Clin Microbiol 50:1968–1973.

449 10. Kilian M, Riley DR, Jensen A, Brüggemann H, Tettelin H. 2014. Parallel Evolution

450 of *Streptococcus pneumoniae* and *Streptococcus mitis* to Pathogenic and

451 Mutualistic Lifestyles. MBio 5:e01490-14.

452 11. Skov Sørensen UB, Yao K, Yang Y, Tettelin H, Kilian M. 2016. Capsular

453 Polysaccharide Expression in Commensal *Streptococcus* Species: Genetic and

454 Antigenic Similarities to *Streptococcus pneumoniae*. MBio 7:e01844-16.

455 12. Jensen A, Valdórsson O, Frimodt-Møller N, Hollingshead S, Kilian M. 2015.

456 Commensal Streptococci Serve as a Reservoir for β-Lactam Resistance Genes in

457 *Streptococcus pneumoniae*. Antimicrob Agents Chemother 59:3529–3540.

13. Shekhar S, Khan R, Ferreira DM, Mitsi E, German E, Rørvik GH, Berild D,

Schenck K, Kwon K, Petersen F. 2018. Antibodies reactive to commensal

Streptococcus mitis show cross-reactivity with virulent *Streptococcus pneumoniae*

serotypes. Front Immunol 9:747.

14. Adams HM, Joyce LR, Guan Z, Akins RL, Palmer KL. 2017. *Streptococcus mitis*

and *S. oralis* Lack a Requirement for CdsA, the Enzyme Required for Synthesis of

Major Membrane Phospholipids in Bacteria. Antimicrob Agents Chemother

61:e02552-16.

15. Mishra NN, Tran TT, Seepersaud R, Miro JM, Rybak MJ, Bayer AS, Arias CA,

Sullam PM. 2017. Perturbations of Phosphatidate Cytidylyltransferase (CdsA)

Mediate Daptomycin Resistance in *Streptococcus mitis/oralis* by a Novel

Mechanism. Antimicrob Agents Chemother 61:1–13.

16. Humphries RM, Pollett S, Sakoulas G. 2013. A current perspective on daptomycin

for the clinical microbiologist. Clin Microbiol Rev 26:759–780.

17. Meiers M, Volz C, Eisel J, Maurer P, Henrich B, Hakenbeck R. 2014. Altered lipid

composition in *Streptococcus pneumoniae cpoA* mutants. BMC Microbiol 14:1–

12.

18. Trombe M-C, Lanéelle M-A, Lanéelle G. 1979. Lipid composition of aminopterin-

resistant and sensitive strains of *Streptococcus pneumoniae*. Biochim Biophys

Acta - Lipids Lipid Metab 574:290–300.

19. Brundish DE, Shaw N, Baddiley J. 1967. The phospholipids of Pneumococcus I-

192R, A.T.C.C. 12213. Biochem J 104:205–11.

20.   Custer JE, Goddard BD, Matter SF, Kaneshiro ES. 2014. The relative proportions

of different lipid classes and their fatty acid compositions change with culture age

in the cariogenic dental pathogen *Streptococcus mutans* UA159. Lipids 49:543–

554.

21.   Chiu T-H, Hung S-A. 1979. Effect of age on the membrane composition of

Streptococcus sanguis. Biochim Biophys Acta 558:267–272.

22.   Goldfine H, Guan Z. 2015. Lipidomic Analysis of Bacteria by Thin-Layer

Chromatography and Liquid Chromatography/Mass Spectrometry, p. 125–139. *In*

Hydrocarbon and Lipid Microbiology Protocols.

23.   Sohlenkamp C, López-Lara IM, Geiger O. 2003. Biosynthesis of

phosphatidylcholine in bacteria. Prog Lipid Res 42:115–162.

24.   Bogdanov M, Sun J, Kaback HR, Dowhan W. 1996. A phospholipid acts as a

chaperone in assembly of a membrane transport protein. J Biol Chem

271:11615–11618.

25.   Bogdanov M, Umeda M, Dowhan W. 1999. Phospholipid-assisted Refolding of an

Integral Membrane Protein. J Biol Chem 274:12339–12345.

26.   Lysenko ES, Gould J, Bals R, Wilson JM, Weiser JN. 2000. Bacterial

Phosphorylcholine Decreases Susceptibility to the Antimicrobial Peptide LL-

37/hCAP18 Expressed in the Upper Respiratory Tract. Infect Immun 68:1664–

1671.

27.   Geiger O, López-Lara IM, Sohlenkamp C. 2013. Phosphatidylcholine biosynthesis

and function in bacteria. Biochim Biophys Acta - Mol Cell Biol Lipids 1831:503–

513.

28.   Conover GM, Martinez-Morales F, Heidtman MI, Luo Z-Q, Tang M, Chen C,

Geiger O, Isberg RR. 2008. Phosphatidylcholine synthesis is required for optimal

function of *Legionella pneumophila* virulence determinants. Cell Microbiol 10:514–

528.

29.   Comerci DJ, Altabe S, De Mendoza D, Ugalde RA. 2006. *Brucella abortus*

synthesizes phosphatidylcholine from choline provided by the host. J Bacteriol

188:1929–1934.

30.   Aktas M, Danne L, Moller P, Narberhaus F. 2014. Membrane lipids in

Agrobacterium tumefaciens: biosynthetic pathways and importance for

pathogenesis. Front Plant Sci 5:109.

31.   Aktas M, Wessel M, Hacker S, Klüsener S, Gleichenhagen J, Narberhaus F.

2010. Phosphatidylcholine biosynthesis and its significance in bacteria interacting

with eukaryotic cells. Eur J Cell Biol 89:888–894.

32.   Malek AA, Wargo MJ, Hogan DA. 2012. Absence of membrane

phosphatidylcholine does not affect virulence and stress tolerance phenotypes in

the opportunistic pathogen *Pseudomonas aeruginosa*. PLoS One 7:e30829.

33.   Kaneshiro T, Law JH. 1964. Phosphatidylcholine Synthesis in *Agrobacterium*

tumefaciens. J Biol Chem 239:1705–1713.

34. De Rudder KEE, Sohlenkamp C, Geiger O. 1999. Plant-exuded choline is used

for rhizobial membrane lipid biosynthesis by phosphatidylcholine synthase. J Biol

Chem 274:20011–20016.

35. Sohlenkamp C, De Rudder KEE, Röhrs V, López-Lara IM, Geiger O. 2000.

Cloning and characterization of the gene for phosphatidylcholine synthase. J Biol

Chem 275:18919–18925.

36. DeLong CJ, Shen Y-J, Thomas MJ, Cui Z. 1999. Molecular distinction of

phosphatidylcholine synthesis between the CDP-choline pathway and

phosphatidylethanolamine methylation pathway. J Biol Chem 274:29683–29688.

37. Vences-Guzmán MÁ, Paula Goetting-Minesky M, Guan Z, Castillo-Ramirez S,

Córdoba-Castro LA, López-Lara IM, Geiger O, Sohlenkamp C, Christopher Fenno

J. 2017. 1,2-Diacylglycerol choline phosphotransferase catalyzes the final step in

the unique *Treponema denticola* phosphatidylcholine biosynthesis pathway. Mol

Microbiol 103:896–912.

38. Gisch N, Schwudke D, Thomsen S, Heß N, Hakenbeck R, Denapaite D. 2015.

Lipoteichoic acid of *Streptococcus oralis* Uo5: a novel biochemical structure

comprising an unusual phosphorylcholine substitution pattern compared to

Streptococcus pneumoniae. Sci Rep 5:16718.

39. Bergström N, Jansson PE, Kilian M, Skov Sørensen UB. 2000. Structures of two

cell wall-associated polysaccharides of a *Streptococcus mitis* biovar 1 strain: A

unique teichoic acid-like polysaccharide and the group O antigen which is a C-

polysaccharide in common with pneumococci. Eur J Biochem 267:7147–7157.

40.   Seong Seo H, Cartee RT, Pritchard DG, Nahm MH. 2008. A new model of

544   pneumococcal lipoteichoic acid structure resolves biochemical, biosynthetic, and

545   serologic inconsistencies of the current model. J Bacteriol 190:2379–2387.

546   41.   Denapaite D, Brückner R, Hakenbeck R, Vollmer W. 2012. Biosynthesis of

547   Teichoic Acids in *Streptococcus pneumoniae* and Closely Related Species:

548   Lessons from Genomes. Microb Drug Resist 18:344–358.

549   42.   Fischer W. 1997. Pneumococcal lipoteichoic and teichoic acid. Microb Drug

550   Resist 3:309–325.

551   43.   Moser R, Aktas M, Narberhaus F. 2014. Phosphatidylcholine biosynthesis in

552   *Xanthomonas campestris* via a yeast-like acylation pathway. Mol Microbiol

553   91:736–750.

554   44.   Stålberg K, Neal AC, Ronne H, Ståhl U. 2008. Identification of a novel GPCAT

555   activity and a new pathway for phosphatidylcholine biosynthesis in *S. cerevisiae*.

556   J Lipid Res 49:1794–1806.

557   45.   Fisher E, Almaguer C, Holic R, Griac P, Patton-vogt J. 2005.

558   Glycerophosphocholine-dependent Growth Requires Gde1p (YPL110c) and

559   Git1p in *Saccharomyces cerevisiae*. J Biol Chem 280:36110–36117.

560   46.   Wishart DS, Feunang YD, Marcu A, Guo AC, Liang K, Vázquez-Fresno R, Sajed

561   T, Johnson D, Li C, Karu N, Sayeeda Z, Lo E, Assempour N, Berjanskii M,

562   Singhal S, Arndt D, Liang Y, Badran H, Grant J, Serra-Cayuela A, Liu Y, Mandal

563   R, Neveu V, Pon A, Knox C, Wilson M, Manach C, Scalbert A. 2018. HMDB 4.0:

564   The human metabolome database for 2018. Nucleic Acids Res 46:D608–D617.

565 47. Guo L, Milburn M V., Ryals JA, Lonergan SC, Mitchell MW, Wulff JE, Alexander

566   DC, Evans AM, Bridgewater B, Miller L, Gonzalez-Garay ML, Caskey CT. 2015.

567   Plasma metabolomic profiles enhance precision medicine for volunteers of normal

568   health. Proc Natl Acad Sci 112:E4901–E4910.

569 48. Głab B, Beganovic M, Anaokar S, Hao MS, Rasmusson AG, Patton-Vogt J,

570   Banaś A, Stymne S, Lager I. 2016. Cloning of glycerophosphocholine

571   acyltransferase (GPCAT) from fungi and plants: A novel enzyme in

572   phosphatidylcholine synthesis. J Biol Chem 291:25066–25076.

573 49. Akins RL, Katz BD, Monahan C, Alexander D. 2015. Characterization of high-level

574   daptomycin resistance in viridans group streptococci developed upon in vitro

575   exposure to daptomycin. Antimicrob Agents Chemother 59:2102–2112.

576 50. Rodriguez-R LM, Konstantinidis KT. 2016. The enveomics collection: a toolbox for

577   specialized analyses of microbial genomes and metagenomes. PeerJ Prepr

578   4:e1900v1.

579 51. Goris J, Konstantinidis KT, Klappenbach JA, Coenye T, Vandamme P, Tiedje JM.

580   2007. DNA-DNA hybridization values and their relationship to whole-genome

581   sequence similarities. Int J Syst Evol Microbiol 57:81–91.

582 52. Rodriguez-R LM, Konstantinidis KT. 2014. Bypassing Cultivation To Identify

583   Bacterial Species. Microbe 9:111–118.

584 53. Konstantinidis K, Tiedje T. 2005. Genomic insights that advance the species

585   definition for prokaryotes. PNAS 102:2567–2572.

586   54.   Ciufo S, Kannan S, Sharma S, Badretdin A, Clark K, Turner S, Brover S, Schoch

587   CL, Kimchi A, DiCuccio M. 2018. Using average nucleotide identity to improve

588   taxonomic assignments in prokaryotic genomes at the NCBI. Int J Syst Evol

589   Microbiol 68:2386–2392.

590   55.   Gopalakrishnan AS, Chen Y-C, Temkin M, Dowhan W. 1986. Structure and

591   expression of the gene locus encoding the phosphatidylglycerophosphate

592   synthase of *Escherichia coli*. J Biol Chem 261:1329–1338.

593   56.   van Beilen J, Blohmke CJ, Folkerts H, de Boer R, Zakrzewska A, Kulik W, Vaz

594   FM, Brul S, Ter Beek A. 2016. RodZ and PgsA Play Intertwined Roles in

595   Membrane Homeostasis of *Bacillus subtilis* and Resistance to Weak Organic Acid

596   Stress. Front Microbiol 7:1633.

597   57.   Tran TT, Mishra NN, Seepersaud R, Diaz L, Rios R, Dinh AQ, Garcia-de-la-Maria

598   C, Rybak MJ, Miro JM, Shelburne SA, Sullam PM, Bayer AS, Arias CA. 2019.

599   Mutations in cdsA and pgsA Correlate with Daptomycin Resistance in

600   *Streptococcus mitis/oralis*. Antimicrob Agents Chemother 63.

601   58.   Van De Rijn I, Kessler RE. 1980. Growth characteristics of group A streptococci in

602   a new chemically defined medium. Infect Immun 27:444–448.

603   59.   Chang JC, LaSarre B, Jimenez JC, Aggarwal C, Federle MJ. 2011. Two group A

604   streptococcal peptide pheromones act through opposing rgg regulators to control

605   biofilm development. PLoS Pathog 7:e1002190.

60.   Ojala PJ, Hirvonen TE, Hermansson M, Somerharju P, Parkkinen J. 2007. Acyl

chain-dependent effect of lysophosphatidylcholine on human neutrophils. J

Leukoc Biol 82:1501–1509.

61.   Jiang D, Zhang Q, Zheng Q, Zhou H, Jin J, Zhou W, Bartlam M, Rao Z. 2014.

Structural analysis of *Mycobacterium tuberculosis* ATP-binding cassette

transporter subunit UgpB reveals specificity for glycerophosphocholine. FEBS J

281:331–341.

62.   Großhennig S, Schmidl SR, Schmeisky G, Busse J, Stülke J. 2013. Implication of

glycerol and phospholipid transporters in *Mycoplasma pneumoniae* growth and

virulence. Infect Immun 81:896–904.

63.   Yang K, Wang M, Metcalf WW. 2009. Uptake of Glycerol-2-Phosphate via the

ugp-Encoded Transporter in *Escherichia coli* K-12. J Bacteriol 191:4667–4670.

64.   Gupta R, Shah P, Swiatlo E. 2009. Differential gene expression in *Streptococcus*

pneumoniae in response to various iron sources. Microb Pathog 47:101–109.

65.   Clinical and Laboratory Standards Institute. 2016. Performance standards for

antimicrobial susceptibility testing, 26th ed. CLSI supplement M100SClinical and

Laboratory Standards Institute, Wayne, PA.

66.   Wick RR, Judd LM, Gorrie CL, Holt KE. 2017. Unicycler: Resolving bacterial

genome assemblies from short and long sequencing reads. PLoS Comput Biol

13:e1005595.

67.   Morrison DA, Khan R, Junges R, Åmdal HA, Petersen FC. 2015. Genome editing

by natural genetic transformation in *Streptococcus mutans*. J Microbiol Methods

119:134–141.

68. Salvadori G, Junges R, Morrison DA, Petersen FC. 2016. Overcoming the barrier

of low efficiency during genetic transformation of *Streptococcus mitis*. Front

Microbiol 7:1009.

69. Biswas I, Gruss A, Ehrlich SD, Maguin E. 1993. High-efficiency gene inactivation

and replacement system for gram-positive bacteria. J Bacteriol 175:3628–3635.

70. Tan BK, Bogdanov M, Zhao J, Dowhan W, Raetz CRH, Guan Z. 2012. Discovery

of a cardiolipin synthase utilizing phosphatidylethanolamine and

phosphatidylglycerol as substrates. Proc Natl Acad Sci 109:16504–16509.

71. Li C, Tan BK, Zhao J, Guan Z. 2016. In vivo and in vitro synthesis of

phosphatidylglycerol by an *Escherichia coli* cardiolipin synthase. J Biol Chem

291:25144–25153.

72. Lu W, Kimball E, Rabinowitz JD. 2006. A high-performance liquid

chromatography-tandem mass spectrometry method for quantitation of nitrogen-

containing intracellular metabolites. J Am Soc Mass Spectrom 17:37–50.

Figures

Fig 1. Phospholipid pathways in bacteria, including pathways for PC

biosynthesis. Shown are the methylation pathway (1), phosphatidylcholine synthase

pathway (2), CDP-choline pathway (3), and the glycerophosphocholine pathway (4).

Lipids detected in THB-cultured SM43 are shown in red. Genes encoded by SM43 are

shown in green. Abbreviations are PA, phosphatidic acid; CDP-DAG, CDP-

diacylglycerol; PS, phosphatidylserine; PE, phosphatidylethanolamine; G3-P, glycerol-3-

phosphate; PGP, phosphatidylglycerol-3-P; PG, phosphatidylglycerol; CL, cardiolipin;

LPG, lysly-phosphatidylglycerol; PIP, phosphatidylinositolphosphate; PI,

phosphatidylinosotiol; PC, phosphatidylcholine, LPC, lysophosphatidylcholine; GPC,

glycerophosphocholine; LPA, lysophosphatidic acid.

Fig 2. PC species detected in SM43, SM43Δ*cdsA* and SM43Δ*pgsA*. Shown are the positive mode total ion chromatogram of SM43 lipids (A), the ESI-MS of SM43 PC species (B), chemical structure of PC (34:1) (C), the ESI-MS of PC species in SM43Δ*cdsA* (D), and (E) the ESI-MS of PC species in SM43Δ*pgsA*. The mass spectra shown are averaged from spectra acquired by NPLC-ESI/MS during the 20–21 min window. PC species are detected by positive ion ESI/MS as the M$^+$ ions.

Fig 3. Exogenous deuterated choline (choline-*d9*) is used by SM43 to synthesize CDP-choline, but not PC. CDP-choline and PC were detected in the soluble-metabolite extract and lipid extract, respectively. Soluble metabolites were analyzed by reverse phase LC/ESI-MS in positive ion mode. Lipids were analyzed by normal phase LC/ESI-MS in positive ion mode. Shown are CDP-choline (M$^+$ ion at *m/z* 489.1) present in SM43 cultured in THB (A), and CDP-choline-*d9* (M$^+$ ion at *m/z* 498.1) present in SM43 cultured in THB supplemented with 2 mM choline-*d9* (B). Note the expected mass shift (9 Da) between CDP-choline and CDP-choline-*d9.* Also shown are PC species detected in SM43 cultured in THB (C), and PC species detected in SM43 cultured in THB supplemented with choline-*d9* (D). No corresponding mass shift (9 Da) is detected in the PC species, excluding the possibility of SM43 using the CDP-choline pathway for PC synthesis.

Fig 4. PC species of SM43 cultured in THB supplemented with GPC-*d9* and lysoPC (20:0). A) ESI/MS detection of PC species in SM43 cultured with GPC-*d9*. Blue dots indicate PC species normally detected in THB-grown SM43, and red dots indicate GPC-*d9*-originating PC species. B) ESI/MS detection of PC species in SM43 when cultured in the presence of lysoPC (20:0). Blue dots indicate PC species normally detected in THB-grown SM43. Red dots are lysoPC (20:0)-originating species. Incorporation of GPC-*d9* and lysoPC (20:0) into PC indicates that the GPC pathway is utilized by SM43 for PC biosynthesis.

Fig 5. PC species of SM43 and *S. mitis* ATCC 49456 cultured in defined media (DM) with and without GPC. PC species (red dots) detected in SM43 (A) and ATCC 49456 (C) cultured in DM, and SM43 (B) and ATCC 49456 (D) cultured in DM supplemented with GPC are shown. PC is only detected when GPC is present in the culture medium.

Fig 6. Proposed model for the GPC pathway in Mitis group streptococci.
Exogenous GPC and lysoPC are transported into the cell via unidentified transporters.
GPC is sequentially acylated to form lysoPC and PC. Acyl chain lengths vary in the lipid
membrane; representative chain lengths are shown.

Fig 7. Positive ion total ion chromatogram (TIC) and PC mass spectrum of *S.* *pneumoniae* D39 and TIGR4 strains in early stationary phase. Shown are the TIC of *S. pneumoniae* D39 lipids (A), the ESI/MS of major PC species in *S. pneumoniae* D39 (B), the TIC of *S. pneumoniae* TIGR4 lipids (C), and the ESI/MS of major PC species in *S. pneumoniae* TIGR4 (D). The mass spectra shown are averaged from spectra acquired by NPLC-ESI/MS during the 19.5 – 21 min window. PC species are detected by positive ion ESI/MS as the M^+ ions, while the co-eluting undecaprenyl phosphate (C_{55}-P) is detected as the $[M+NH_4]^+$ ion (*m/z* 864.7).

Fig 8. GPC pathway confirmation in *S. pneumoniae* D39 and *S. oralis* ATCC 35037.

Shown are PC species (red dots) detected for D39 (A) and ATCC 35037 (C) cultured in

DM and D39 (B) and ATCC 35037 (D) cultured in DM supplemented with GPC. PC is

only detected when GPC is present in the culture medium.

718 **Table 1.** Summary of major glycolipids and phospholipids detected in Mitis group

719 streptococci in routine laboratory media[1].

Streptococcal species	Strain	Source	DAG	MHDAG	DHDAG	PA	PG	CL	C55-P	PC	L-PG	Growth medium[2]
S. mitis	ATCC 49456	Type strain	+	+	+	+	+	+	+	+	−	THB
S. oralis	ATCC 35037	Type strain	+	+	+	+	+	+	+	+	−	THB
	1647[3]	Endocarditis	+	+	+	+	+	+	+	+	−	THB
	1648[3]	Endocarditis	+	+	+	+	+	+	+	+	−	THB
S. pneumoniae	D39	Historical strain	+	+	+	+	+	+	+	+	−	THB+Y
	TIGR4	Bacteremia	+	+	+	+	+	+	+	+	−	THB+Y
Mitis group	1643 (SM43)[3]	Endocarditis	+	+	+	+	+	+	+	+	−	THB
	SM43ΔcdsA	This study	+	+	+	+	-	-	+	+	−	THB
	SM43ΔpgsA	This study	+	+	+	+	-	-	+	+	−	THB

[1]Abbreviations: DAG, diacylglycerol; MHDAG, monohexosyldiacylglycerol; DHDAG,

dihexosyldiacylglycerol; PA, phosphatidic acid; PG, phosphatidylglycerol; CL,

cardiolipin; C55-P, undecaprenyl phosphate; PC, phosphatidylcholine; L-PG, lysyl-

phosphatidylglycerol

[2]Todd-Hewitt Broth (THB); THB supplemented with 0.5% yeast extract (THB+Y).

[3]Lipid profiles previously published by Adams *et al* 2017 (14).

Table 2. Summary of major glycolipids and phospholipids detected in Mitis group

streptococci cultured in DM with or without GPC[1].

Streptococcal species	Strain	Source	DAG	MHDAG	DHDAG	PA	PG	CL	C55-P	PC	L-PG	Growth medium[2]
S. mitis	ATCC 49456	Type strain	+	+	+	+	+	+	+	-	−	DM
	ATCC 49456		+	+	+	+	+	+	+	+	−	DM+GPC
S. oralis	ATCC 35037	Type strain	+	+	+	+	+	+	+	-	−	DM
	ATCC 35037		+	+	+	+	+	+	+	+	−	DM+GPC
S. pneumoniae	D39	Historical strain	+	+	+	+	+	+	+	-	−	DM
	D39		+	+	+	+	+	+	+	+	−	DM+GPC
Mitis group	1643 (SM43)	Endocarditis	+	+	+	+	+	+	+	-	−	DM
	1643 (SM43)		+	+	+	+	+	+	+	+	−	DM+GPC

[1]Abbreviations: DAG, diacylglycerol; MHDAG, monohexosyldiacylglycerol; DHDAG,

dihexosyldiacylglycerol; PA, phosphatidic acid; PG, phosphatidylglycerol; CL,

cardiolipin; C55-P, undecaprenyl phosphate; PC, phosphatidylcholine; L-PG, lysyl-

phosphatidylglycerol

[2]Defined media (DM); Defined media supplemented with glycerophosphocholine

(DM+GPC).

Supplemental information

Phosphatidylcholine biosynthesis in Mitis group streptococci via host metabolite

scavenging

Luke R. Joyce[a], Ziqiang Guan[b]#, and Kelli L. Palmer[a]#

[a]Department of Biological Sciences, The University of Texas at Dallas, Richardson,

Texas, USA

[b]Department of Biochemistry, Duke University Medical Center, Durham, NC[2]

Running title: Phosphatidylcholine biosynthesis in streptococci

#Corresponding authors:

Ziqiang Guan: Ziqiang.Guan@duke.edu

Kelli Palmer: Kelli.Palmer@utdallas.edu

Text S1: Supplementary Materials and Methods.

Acidic Bligh-Dyer extractions. Centrifugation was performed using a Sorvall RC6+ centrifuge. Cultures were pelleted at 4,280 x *g* for 5 min at room temperature. The supernatants were removed and stored at -80°C until acidic Bligh-Dyer lipid extractions were performed as described (1) with minor modifications. Cell pellets were resuspended in 1X PBS (Sigma-Aldrich) and transferred to Coring Pyrex glass tubes with PTFR-lined caps (VWR), followed by 1:2 vol:vol chloroform:methanol addition. Single phase extractions were vortexed periodically and incubated at room temperature for 15 minutes before 500 x *g* centrifugation for 10 min. A two-phase Bligh-Dyer was achieved by addition of 100 µl 37% HCL, 1 mL CHCl$_3$, and 900 µl of 1X PBS, which was then vortexed and centrifuged for 5 min at 500 x *g.* The lower phase was removed to a new tube and dried under nitrogen before being stored at -80°C prior to lipidomic analysis.

Normal phase Liquid Chromatography/ Mass Spectrometry. Normal-phase LC/MS was performed on an Agilent 1200 quaternary LC system equipped with an Ascentis silica high-performance liquid chromatography (HPLC) column (5m; 25 cm by 2.1 mm; Sigma-Aldrich) as described previously (1, 2). Briefly, mobile phase A consisted of chloroform-methanol-aqueous ammonium hydroxide (800:195:5, vol/vol), mobile phase B consisted of chloroform-methanol-water-aqueous ammonium hydroxide (600: 340:50:5, vol/vol), and mobile phase C consisted of chloroform-methanol-water-aqueous ammonium hydroxide (450:450:95:5, vol/vol/vol/vol). The elution program

consisted of the following: 100% mobile phase A was held isocratically for 2 min, then

linearly increased to 100% mobile phase B over 14 min, and held at 100% mobile phase

B for 11 min. The LC gradient was then changed to 100% mobile phase C over 3 min,

held at 100% mobile phase C for 3 min, and, finally, returned to 100% mobile phase A

over 0.5 min and held at 100% mobile phase A for 5 min. The LC eluent (with a total

flow rate of 300 ml/min) was introduced into the ESI source of a high-resolution

TripleTOF5600 mass spectrometer (Sciex, Framingham, MA). Instrumental settings for

negative-ion ESI and MS/MS analysis of lipid species were as follows: IS = -4,500 V,

CUR = 20 psi, GSI = 20 psi, DP = -55 V, and FP = -150V. The MS/MS analysis used

nitrogen as the collision gas. Data analysis was performed using Analyst TF1.5 software

(Sciex, Framingham, MA).

Metabolite extractions. Briefly, cell pellets were resuspended in 300 µL super-chilled

methanol:water (vol:vol 4:1) via vortexing and pipetting, and stored on dry ice for 15

min. Suspensions were centrifuged at 17,136 x g at 4°C for 5 min in a table top

centrifuge. The supernatant was removed to a pre-chilled Corning Pyrex glass tube and

kept on dry ice. A further 2 resuspensions in 200 µL methanol:water (vol:vol 4:1) were

performed. Pooled supernatants were dried under nitrogen at room temperature and

stored at -80°C prior to analysis.

Reverse phase LC-ESI/MS analysis. RPLC-ESI/MS analysis of water-soluble

metabolites was performed using a Shimadzu LC system (comprising a solvent

degasser, two LC-10A pumps and a SCL-10A system controller) coupled to a

TripleTOF5600 mass spectrometer (Sciex, Framingham, MA). LC was operated at a

flow rate of 200 µl/min with a linear gradient as follows: 100% of mobile phase A was

held isocratically for 2 min and then linearly increased to 100% mobile phase B over 14

min and held at 100% B for 4 min. Mobile phase A consisted of

methanol/acetonitrile/aqueous 1mM ammonium acetate (60/20/20, v/v/v). Mobile phase

B consisted of 100% ethanol containing 1 mM ammonium acetate. A Zorbax SB-C8

reversed-phase column (5 µm, 2.1 x 50 mm) was obtained from Agilent (Palo Alto, CA).

The LC eluent was introduced into the ESI source of the mass spectrometer. Instrument

settings for negative ion ESI/MS and MS/MS analysis of lipid species were as follows:

Ion spray voltage (IS) = -4500 V; Curtain gas (CUR) = 20 psi; Ion source gas 1 (GS1) =

20 psi; De-clustering potential (DP) = -55 V; Focusing potential (FP) = -150 V. Data

acquisition and analysis were performed using the Analyst TF1.5 software (Sciex,

Framingham, MA).

Deletion of SM43 *cdsA*. Primers used in this study are shown in Table S2. The *cdsA*

deletion construct was designed essentially as previously described (3, 4).

Approximately 2 Kb flanking regions on either side of the gene were amplified using

Phusion polymerase (Thermo Fisher). PCR products were digested with restriction

enzymes (New England Biolabs) and ligated using T4 DNA ligase (New England

Biolabs). Ligations were amplified by PCR utilizing the 5' and 3' most primers with

Phusion polymerase to obtain a linear product. The PCR product was analyzed on a 1%

agarose gel and extracted using the QIAquick Gel Extraction Kit per the manufacturer's

protocol. The linear construct was transformed into SM43 via natural transformation,

described below. Transformation plates were incubated overnight. Putative transformant

colonies were inoculated into THB and screened via PCR for the clean deletion of *cdsA*.

The *cdsA* regions of putative mutants were sequenced for confirmation (Massachusetts

General Hospital DNA Core).

A

B

Fig S1. Phosphatidylcholine (PC) species in *S. mitis* ATCC 49456. A) *S. mitis* ATCC 49456, B) MS/MS daughter ion fragmentation of PC (34:1) *m/z* 760.6 in SM43. The mass spectra shown are averaged from spectra acquired by NPLC-ESI/MS during the 20–21 min window. PC species are detected by positive ion ESI/MS as the M[+] ions.

Fig S2. **Growth curves.** A) Growth curves for SM43 and SM43Δ*pgsA* in THB. B)

Growth curves of SM43 in THB, and in defined medium (DM) with and without GPC

supplementation and C) Growth curves of ATCC49456 in THB, and in defined medium

(DM) with and without GPC supplementation. OD_{600nm} readings were taken every hour,

102   with a starting inoculum of OD_{600nm} 0.05. Growth curves were performed as biological

103   triplicate. Error bars indicate standard deviation.

Fig S3. MS/MS fragmentation of CDP-choline and CDP-choline-*d9*. A) MS/MS of CDP-choline (*m/z* 489) B) MS/MS of CDP-choline-*d9* (*m/z* 498). Note the mass shift of 9-*da* for all major MS/MS peaks in B.

MS/MS *m/z* 258.1
GPC

109
110 **Fig S4. MS detection of GPC in THB.** The soluble-metabolite extract of THB was

111 analyzed by reverse phase LC/MS. A) GPC is detected by positive ion ESI/MS as the

112 M$^+$ ion at *m/z* 258.108 and [M-H+Na]$^+$ ion at *m/z* 280.091. B) MS/MS spectrum of the M$^+$

113 ion (*m/z* 258.1) of GPC. The chemical structures of major fragments are shown.

Fig S5. ESI/MS of phosphatidylglycerol (PG) species of SM43 cultured in THB with

and without lysoPC (20:0) addition. A) ESI/MS detection of PG species in SM43

cultured in THB. B) ESI/MS detection of PG species in SM43 cultured in THB in the

presence of lysoPC (20:0). In sharp contrast to what is observed for PC, there is no

appreciable incorporation of the 20:0-acyl chain into PG in the presence of lysoPC

(20:0). The lack of incorporation of C20:0 into PG indicates the insignificant

transacylation activity of SM43. PG species are detected by negative ion ESI/MS as the

[M-H]⁻ ions.

123 **Table S1.** Strains and plasmids used in this study.

Organism	Strain	Description	Ref
Mitis group streptococci	1643 (SM43)	Wild-type infective endocarditis isolate	(5)
	1643ΔcdsA	cdsA deletion strain	This work
	1643ΔpgsA::ermB	pgsA deletion strain, replaced with ermB. Erm^r	This work
S. mitis	ATCC 49456 (NCTC 12261)	Wild-type S. mitis type strain from ATCC	(6)
S. oralis	ATCC 35037 (NCTC 11427)	Wild-type S. oralis type strain from ATCC	(7)
S. pneumoniae	D39	Wild type, historical strain	(8)
	TIGR4	Wild type, bloodstream isolate	(9)
Plasmid	**Description**		**Ref**
pG$^+$host 4	Encodes ermB		(10)

126 **Table S2.** Primers used in this study.

Primer	Sequence (5'-3')	Use
L_Updel_F	TTTACGTGAATATATCGGT	SM43 upstream of *cdsA* region amplification
L_Updel_R PstI	ACGTCACTGCAGATCCTTGGTCATATCTTCTC	SM43 upstream of *cdsA* region amplification
L_Dwndel_F PstI	ACGTCACTGCAGCCAATGATGCACTTATTC	SM43 downstream of *cdsA* region amplification
L_Dwndel_R	CTTGATTTCTTCTTGCAC	SM43 downstream of *cdsA* region amplification
L1_R	CGTACATATCTTCGACTGTC	*cdsA* knockout sequencing with L_Updel_F
L2_F	CCTTCGTACCATGATTGAG	*cdsA* knockout sequencing
L2_R	GTCAGATTTTCCCATTTTTC	*cdsA* knockout sequencing
L3_F	TTTAGAGAATGAGGACCG	*cdsA* knockout sequencing
L3_R	CATATCTTCTGTGATGT	*cdsA* knockout sequencing
L4_F	ATGGTGGTCGTGCTGAA	*cdsA* knockout sequencing
L4_R	GATACTGTACATCCAAAGG	*cdsA* knockout sequencing
L5_F	CAGTAGATCACGATGCAAC	*cdsA* knockout sequencing
L5_R	GTCGCATAATGTCATTCC	*cdsA* knockout sequencing
L6_F	GTGTAGTTATCATGGTTG	*cdsA* knockout sequencing with L_Dwndel_R
L7F	CAAACGGCTCTTCCTATGC	*cdsA* knockout sequencing
L7R	CTGATAACTTGAGCATCG	*cdsA* knockout sequencing
SM43cdsA_reseq_F	TCAGGTCTATATGCATCTG	*cdsA* upstream resequence
SM43cdsA_reseq_R	GCTTCATCAAAATCAGG	*cdsA* upstream resequence
PgsA_updel_F	CTTTACTCCCTAAGCAAG	SM43 upstream of *pgsA* region amplification
PgsA_updel_R_PstI	ACGTCACTGCAGTCTACCAATAGTTAATACATTAGG	SM43 upstream of *pgsA* region amplification
PgsA_dwndel_F_XmaI	ACGTCACCCGGGTTGTTGCTTGTTGAGACG	SM43 downstream of *pgsA* region amplification
PgsA_dwndel_R	GATCAAGAAGAACATGGA	SM43 downstream of *pgsA* region amplification
Erm_F_XmaI	ACGTCACCCGGGTAACGATCACTCATCATG	*erm* amplification from pG⁺host4 plasmid
Erm_R_PstI	ACGTCACTGCAGCAAGTTAAGGGATGCAGT	*erm* amplification from pG⁺host4 plasmid

References

1. Tan BK, Bogdanov M, Zhao J, Dowhan W, Raetz CRH, Guan Z. 2012. Discovery
    of a cardiolipin synthase utilizing phosphatidylethanolamine and
    phosphatidylglycerol as substrates. Proc Natl Acad Sci 109:16504–16509.

2. Li C, Tan BK, Zhao J, Guan Z. 2016. In vivo and in vitro synthesis of
    phosphatidylglycerol by an *Escherichia coli* cardiolipin synthase. J Biol Chem
    291:25144–25153.

3. Morrison DA, Khan R, Junges R, Åmdal HA, Petersen FC. 2015. Genome editing
    by natural genetic transformation in *Streptococcus mutans*. J Microbiol Methods
    119:134–141.

4. Salvadori G, Junges R, Morrison DA, Petersen FC. 2016. Overcoming the barrier
    of low efficiency during genetic transformation of *Streptococcus mitis*. Front
    Microbiol 7:1009.

5. Akins RL, Katz BD, Monahan C, Alexander D. 2015. Characterization of high-level
    daptomycin resistance in viridans group streptococci developed upon in vitro
    exposure to daptomycin. Antimicrob Agents Chemother 59:2102–2112.

6. Kilian M, Mikkelsen L, Henrichsen J. 1989. Taxonomic Study of Viridans
    Streptococci: Description of *Streptococcus gordonii* sp. nov. and Emended
    Descriptions of *Streptococcus sanguis* (White and Niven 1946), *Streptococcus*
    oralis (Bridge and Sneath 1982), and *Streptococcus mitis* (Andrewes and Horder
    1906). Int J Syst Bacteriol 39:471–484.

7.   Bridge PD, Sneath PH. 1982. *Streptococcus gallinarum* sp. nov. and

Streptococcus oralis sp. nov. Int J Syst Bacteriol 32:410–415.

8.   Lanie JA, Ng WL, Kazmierczak KM, Andrzejewski TM, Davidsen TM, Wayne KJ,

Tettelin H, Glass JI, Winkler ME. 2007. Genome sequence of Avery's virulent

serotype 2 strain D39 of *Streptococcus pneumoniae* and comparison with that of

unencapsulated laboratory strain R6. J Bacteriol 189:38–51.

9.   Tettelin H, Nelson KE, Paulsen IT, Eisen JA, Read TD, Peterson S, Heidelberg J,

Deboy RT, Haft DH, Dodson RJ, Durkin AS, Gwinn M, Kolonay JF, Nelson WC,

Peterson JD, Umayam LA, White O, Salzberg SL, Lewis MR, Radune D,

Holtzapple E, Khouri H, Wolf AM, Utterback TR, Hansen CL, Mcdonald LA,

Feldblyum T V, Angiuoli S, Dickinson T, Hickey EK, Holt IE, Loftus BJ, Yang F,

Smith HO, Venter JC, Dougherty BA, Morrison DA, Hollingshead SK, M. FC.

2001. Complete Genome Sequence of a Virulent Isolate of *Streptococcus*

pneumoniae. Science (80-) 293:498–507.

10.   Biswas I, Gruss A, Ehrlich SD, Maguin E. 1993. High-efficiency gene inactivation

and replacement system for gram-positive bacteria. J Bacteriol 175:3628–3635.