Modeling microbial diversity with metabolic trade-offs

Zhiyuan Li[1,2], Bo Liu[3], Sophia Hsin-Jung Li[4], Christopher G. King[5], Zemer Gitai[4], and Ned S. Wingreen*[4,6]

[1.] Center for the Physics of Biological Function, Princeton University.
[2.] Princeton Center for Theoretical Science, Princeton University.
[3.] Yuanpei College, Peking University, Beijing, China.
[4.] Department of Molecular Biology, Princeton University.
[5.] Department of Physics, Princeton University.
[6.] Lewis-Sigler Institute for Integrative Genomics, Princeton University, Princeton, NJ, USA

ABSTRACT

Nature exhibits much higher biodiversity than predicted by theories of competition. One solution for reconciling this "paradox of the plankton" is to imposes metabolic trade-offs, where species need to allocate limited cellular resources into multiple functions. However, two questions exist for metabolic models: first, as many such models have been proposed with diverse assumptions and different results, can we find a universal language to summarize various models into one unified framework? Second, under the pressure of evolution, will there be a single optimal metabolic strategy that finally dominates over all others? In this work, we address these two questions by constructing a generalizable framework to describe the species-environment feedback in chemostat-type resource-competition models. Employing this framework, a fitness landscape based on the strategy-growth rate relationship can be constructed. Species are capable of creating their own fitness landscape by shaping their nutrient environment, which allows for dynamic fitness landscapes and rich ecological behaviors, and is crucial for biodiversity in all the models we examined. A non-invasible strategy corresponds to a species creating a fitness landscape that places itself at the top. Under certain conditions, more than one species is required to complete this task, which leads to evolutionarily stable coexistence. Our approach facilitates quantitative understanding of chemostat experiments, and provides insight into the competitive-exclusion paradox.

INTRODUCTION

In the natural world, species are in constant competition. So why doesn't the fittest species outcompete the others and become the sole survivor? This question, captured by the "paradox of the plankton" (Hutchinson, 1961), has perplexed community ecologists for nearly a century. On the basis of simple resource-competition models, it has been argued that the number of stably coexisting species cannot exceed the number of resources, leading to the so-called competitive exclusion principle (Armstrong & McGehee, 1980; Hardin, 1960; Levin, 1970; McGehee & Armstrong, 1977). Nevertheless, tremendous biodiversity manifests in the real world, from environmental surveys to controlled lab experiments (Friedman, Higgins, & Gore, 2017; Goldford et al., 2018; Maharjan, Seeto, Notley-McRobb, & Ferenci, 2006). Even in well-mixed aquatic environments, hundreds of species of phytoplankton coexist on a few types of abiotic nutrients (Cermeño, Teixeira, Branco, Figueiras, & Marañón, 2014; Moore et al., 2013).

A multitude of hypotheses have been proposed to circumvent the competitive exclusion principle (Palmer, 1994; Roy & Chattopadhyay, 2007; Scheffer, Rinaldi, Huisman, & Weissing, 2003). Some introduce additional factors such as cooperative or antagonistic interactions between species (Bairey, Kelsic, & Kishony, 2016; Freilich et al., 2011; Kerr, Riley, Feldman, & Bohannan, 2002; Murdoch & Oaten, 1975; Venturelli et al., 2018; Wintermute & Silver, 2010), spatially structured habitats (Amarasekare, 2003; Geyrhofer & Brenner, 2018; Huisman, van Oostveen, & Weissing, 1999; D. Tilman, 1994), temporal fluctuations of the environment (Amarasekare, 2003; Behar, Brenner, & Louzoun, 2014; Descamps-Julien & Gonzalez, 2005; Huisman et al., 1999; D. Tilman, 1994), or complex life-histories of species (Huisman, Johansson, Folmer, & Weissing, 2001). Even in spatially homogeneous models with constant nutrient supply and without direct species interactions, rich dynamics have been uncovered under various conditions. In a model where species compete for essential resources, different nutrient requirements can produce intrinsically oscillatory or even chaotic dynamics that allows for increased diversity (Huisman & Weissing, 1999, 2001). Alternatively, cross-feeding (Goldford et al., 2018; Pfeiffer & Bonhoeffer, 2004), preferential nutrient utilization (A. Goyal, Dubinkina, & Maslov, 2018), or trade-offs (Beardmore, Gudelj, Lipson, & Hurst, 2011; Taillefumier, Posfai, Meir, & Wingreen, 2017) can promote stable coexistence. Recently, a simple model with a trade-off in nutrient uptake, was shown to self-organize to a state of unlimited coexistence (Posfai, Taillefumier, & Wingreen, 2017). This large variety of models and the richness of possible behaviors raises the question of unification: is there a simple framework that consolidates this diverse group of models into one easily understandable picture?

A key challenge to producing such a framework is that fitness landscapes are not static. Not only can extrinsic environmental factors fluctuate in space and time (Mustonen & Lässig, 2009), but species can also actively reshape their habitats (Laland, Matthews, & Feldman, 2016; Leibold, 1995; Odling-Smee, Laland, & Feldman, 2003). The feedback loop between species and their environment produces an intrinsically dynamic fitness landscape in which the action of one species can influence the fitness of all species. A

profound example is the Great Oxygenation Event, when cyanobacteria created an
oxygen-rich atmosphere (Kasting & Siefert, 2002), causing a massive extinction of
anaerobic bacteria but also stimulating an explosion of biodiversity (Schirrmeister, de
Vos, Antonelli, & Bagheri, 2013). Today, species continue shaping their habitats on all
scales: from humans inducing the sixth mass extinction (Ceballos et al., 2015) to
microbes consuming nutrients, releasing wastes, and producing toxins (Callahan,
Fukami, & Fisher, 2014).
85
Resource-competition models provide a simple context to explore the interaction
between species and their environment (Smith & Waltman, 1995). In such models,
species interact only indirectly, via consumption (and sometimes production) of a
common pool of nutrients. A steady state can be reached if the species present can
shape the nutrient concentration to support a growth rate equal to their dilution or death
rate (David Tilman, 1982). Resource-competition models underpin many ecosystem
theories including contemporary niche theory as pioneered by MacArthur (MacArthur,
1970), popularized by Tilman (David Tilman, 1980, 1982), and extended by Chase and
Leibold (Chase & Leibold, 2003). A central component of contemporary niche theory is
a graphical approach, generally consisting of three components: zero net growth
isoclines (ZNGIs) in nutrient space, an impact vector representing a species' nutrient
consumption, and a supply point to described the external resource supply (Koffel,
Daufresne, Massol, & Klausmeier, 2016). This graphical approach is a powerful and
intuitive way of evaluating the outcome of competition and community assembly, but
has not been commonly utilized to understand models of coexistence (Letten, Ke, &
Fukami, 2017), especially coexistence beyond the limit of competitive exclusion
(Huisman & Weissing, 1999; Posfai et al., 2017).
103
In this work, we utilize and extend the graphical tools of resource-competition theory to
relate and unify multiple models for microbial diversity, emphasizing the consequences
of species creating their own environment. The nutrient environment shaped by one
species through growth and consumption may be inviting or prohibiting to another
species. We represent this species-environment interplay via an intuitive geometric
visualization of a "rule of invasion". Under various model assumptions, the species-
environment feedback allows intransitivity of fitness, in which there is no strict
competition hierarchy and therefore no single best species or group of species
(Soliveres et al., 2015). We demonstrate how such intransitivity can lead to rich
ecosystem dynamics, including mutual invasion, multistability, and oscillations, and how
all of these behaviors can be simply related via our graphical representation combined
with the rule of invasion.
116
We extend our investigation of coexistence to encompass evolution. As species evolve
to adapt to their environment, an ongoing threat to diversity is that mutation/selection
may produce a supreme winner that takes over the habitat. To quantify the impact of
evolution on biodiversity, we focus on models with metabolic trade-offs: with limited
cellular resources, the growth rate of cells cannot increase without bound. Rather
evolution optimizes over cells' internal resource allocation strategies (S. Goyal, Yuan,
Chen, Rabinowitz, & Wingreen, 2010; Liebermeister et al., 2014). Different metabolic

strategies lead to different growth rates in different environments, driving a dynamic
fitness landscape. From the perspective of species-environment feedback, we can
define non-invasible/optimal metabolic strategies – namely, one or more species that
construct a fitness landscape which places themselves on the top. When multiple
species are indispensable in co-creating such an environment, their coexistence
becomes evolutionarily stable.

RESULTS

Nutrient limitations, resource allocation strategies, and the chemostat

The growth of microbes is limited by the availability of external nutrients and by their own internal resources, including proteins and energy. Facing different stringencies, cells need to adjust the allocation of their limited internal resources toward a variety of cellular functions (nutrient import and assimilation, energy production, reproduction, maintenance, etc.) to achieve efficient growth. For example, in *E. coli* a large fraction of cellular resources is allocated to synthesize ribosomes, which is reflected in the RNA-to-Protein ratio (RP ratio). In a previous work, we used chemostats (Fig 1A) to quantitatively control growth rate and nutrient supply and showed that the RP ratio in *E. coli* is significantly lower under Phosphorus (P)-limitation than under Carbon (C)- or Nitrogen (N)-limitation at the same growth rates (Li et al., 2018), indicating cells adopt different proteome allocation strategies in response to different nutrient limitations. This observation raised further questions: How can we achieve fine control of nutrient limitation? Is the limiting nutrient determined by the absolute concentrations of supplied nutrients, or the relative abundance of one nutrient compared to the others?

To answer these questions, we ran chemostats at multiple growth rates and supply conditions, using RP ratio as a metric for P or C/N limitation. We started from a supply designed for P-limitation, i.e. with carbon and nitrogen supplied in excess, then kept the phosphate concentration in the supply unchanged but decreased the nitrogen concentration. If the nutrient limitation is controlled by the absolute concentration in the supply, cells should retain their RP ratio regardless of nitrogen supply. However, we observed that as the concentration of supplied nitrogen decreased, the RP ratios at all measured growth rates shifted up, from values reflecting P-limitation to values more characteristic of N-limitation, even though the supply of phosphorus remained at the initial "limiting" concentration (Fig 1B). Similar results were obtained for Carbon limitation (Fig 1C), where the RP-ratio curve was shifted to values resembling C-limitation over a series of dilutions of the carbon supply starting from P-limitation. These observed shifts of RP ratios indicate that nutrient limitation in a chemostat depends on the relative difference between supplied nutrients, rather than on the absolute concentration of any single supplied nutrient. Also, these observations highlight that non-intuitive behaviors can occur even in a system as simple as chemostat, highlighting the need for quantitative modeling.

The convergence towards steady state makes the chemostat an ideal experimental system to culture microorganisms and investigate their physiology with a constant environment and growth rate (Wides & Milo, 2018; Ziv, Brandt, & Gresham, 2013). Despite its simplicity, the chemostat captures an important property of ecosystems, namely that species create their own environment. For these reasons, many resource-competition models are based on chemostat-type dynamics. In the following sections, we present graphical representations that facilitate the interpretation of chemostat models and experiments, and provide intuitive understanding of a variety of resource-competition models via the lens of species-environment feedback.

A graphical representation of a chemostat model

In an idealized model of a chemostat (Fig 1A), p types of nutrients are supplied at rate d

and concentrations $\vec{c}_{\text{supply}} = (c_{1,\text{supply}}, c_{2,\text{supply}}, \dots c_{p,\text{supply}})$, meanwhile cells and medium

are diluted at the same rate d. However, the environment that directly impacts cells is

the nutrient concentration inside the chemostat, $\vec{c} = (c_1, c_2, \dots c_p)$, which influences the

intracellular metabolite concentrations $\vec{q}$ and the growth rate g of each species.

Accordingly, the biomass density m of each species in the chemostat obeys:

$$\frac{dm}{dt} = m \cdot (g(\vec{c}, \vec{q}) - d). \tag{1}$$

184 The concentration c_i of the i-th nutrient is a variable, influenced by its rate of

185 consumption by cells. For a single species with import rate I_i per cell volume, the

186 concentration c_i satisfies

$$\frac{dc_i}{dt} = d \cdot (c_{i,\text{supply}} - c_i) - \frac{m}{r} \cdot I_i(\vec{c}, \vec{q}), \tag{2}$$

where r is a constant representing the biomass per cell volume. (If the volume of the

chemostat is $V_{\text{chemostat}}$ and the total volume of cells is V_{cells}, the import flux of the i-th

nutrient $V_{\text{cells}} \cdot I_i$ implies a rate of change of concentration inside cells of I_i and a

corresponding rate of change of the concentration in the chemostat of $V_{\text{cells}}/V_{\text{chemostat}} \cdot$

$I_i = (m \cdot V_{\text{chemostat}}/r)/V_{\text{chemostat}} \cdot I_i = m/r \cdot I_i$.)

In this manuscript, we define $\vec{c}$ as the "nutrient environment", and all possible values of

$\vec{c}$ constitute the "nutrient space". Within a cell, the concentration of metabolites is

influenced by intake rate, and influences the growth rate. Different metabolic models

assume different forms for such influences, and we use $\vec{f}(\vec{I}(\vec{c}, q), \vec{q})$ to represent the

rate of change of $\vec{q}$:

$$\frac{d\vec{q}}{dt} = \vec{f}(\vec{I}(\vec{c}, \vec{q}), \vec{q}). \tag{3}$$

Eqs. (1)-(3) represent a chemostat model in general. The simplicity of the chemostat

has inspired many theoretical studies of resource competition, and different model

assumptions about how species grow and consume nutrients have produced a variety

of intriguing behaviors and conclusions. However, the origins of these differences are

not always simple to discern. To provide a unified view, we present a graphical

representation that allows ready visualization of the rich behaviors that emerge from the

feedback between species and the environment in a chemostat.

We focus on steady-state behavior, where Eq. (3) is equal to zero and $\vec{q}$ can be solved

for as a function of $\vec{c}$, reducing $g(\vec{c}, \vec{q})$ and $I_i(\vec{c}, \vec{q})$ to functions fully dependent on $\vec{c}$,

namely $g(\vec{c})$ and $I_i(\vec{c})$. A graphical representation of the steady state created by a

single species consists of three components (details in Methods):

1. *The growth contour* reflects how the nutrient environment determines cell growth. The
growth-rate function $g(\vec{c})$ maps different points in nutrient space onto growth rates
(background color in Fig 1D). At steady state, the relation $dm/dt = 0$ (Eq. (1)) requires
the growth rate to be exactly equal to the dilution rate (assuming nonzero cell density).
Therefore, the contour in nutrient space satisfying $g(\vec{c}) = d$ indicates all possible
environments that could support the steady state of the species (red curve in Fig 1D).
219
2. *The flux-balance curve and the supply line* reflect how cells shape the nutrient
environment for a given supply condition. At steady state, nutrient influx, dilution, and
consumption need to be balanced such that $dc_i/dt = 0$ in Eq. (2). Flux balance can be
expressed in two ways, depending on whether the unknown is the nutrient environment
or the supply condition: First, given a supply condition, different values of cell density m
lead to different steady-state nutrient concentrations (Eq. (S5)), constituting a one-
dimensional "flux-balance curve" in nutrient space (purple, cyan, and blue curves in Fig
1D). Alternatively, given a specified steady-state nutrient concentration, different values
of cell density m lead to a straight line in the space of supply conditions, which we call
the "supply line" (see Methods for details). Despite the fact that the supply space and
the nutrient space are distinct, they share the same units of concentration in each
dimension. Therefore, for ease of visualize we typically show supply lines along with
other features in the nutrient space (Eq. (S6), black dashed line in Fig 1D).
233
234 3. *The steady-state nutrient environment* $\vec{c}_{\mathrm{ss}}$ created by the species is the intersection of
235 the growth contour and the flux-balance curve (Fig 1D, red dot). Due to consumption by
236 the cells, $\vec{c}_{\mathrm{ss}}$ is less than $\vec{c}_{\mathrm{supply}}$ for all nutrients.
237

238 ### Subtleties in the control of nutrient limitation
239 The ability of cells to create their own nutrient environment brings complexity to
240 chemostat experiments. In this section, we present two examples of subtleties in the
241 control of nutrient limitation, and show how a graphical representation can assist in
242 interpreting the experiments.
243
First, as all the $\vec{c}_{\mathrm{supply}}$ on one supply line correspond to a single steady-state $\vec{c}_{\mathrm{ss}}$, the
resulting nutrient limitation in a chemostat depends on the difference between nutrient
supplies, not on the absolute concentration of any single supplied nutrient. Different
supply conditions (such as the purple, cyan, and blue open circles in Fig 1D), as long as
they fall on the same supply line, can lead to identical steady-state nutrient
environments (red dot in Fig 1D). Therefore, these supply conditions will lead to
identical nutrient limitations and thus the same cellular response. This phenomenon is
distinct from ratio-sensing, in which cells process dissimilar environmental inputs into
similar responses (Escalante-Chong et al., 2015; Wang & Tang, 2017). The graphical
result that the steady-state nutrient concentration and thus the nutrient limitation
depends on the relative supply of different nutrients is consistent with our experimental
observation (Fig 1B and C) that decreasing the nitrogen or carbon supply from a P-
limited condition induces a similar proteome allocation as that induced by an N/C-limited
supply with P in excess.
258

Interestingly, changing the dilution rate d alone has the potential to switch the limiting
nutrient. As shown in Fig 1E, if the supply concentrations (blue open circle) are chosen
to produce the blue flux-balance curve, this curve intersects with the yellow, orange,
and deep red growth contours on the horizontal, horizontal, and vertical edges,
respectively. Therefore, as the dilution rate and growth rate increase, the species will
transition from nutrient b-limited to nutrient a-limited growth, even though the supply
concentrations $\vec{c}_{\mathrm{supply}}$ are kept unchanged.
266
Metabolic trade-off and strategies
As demonstrated by our R/P ratio measurements of *E. coli*, microorganisms allocate
their limited internal resources according to the nutrient environment they perceive. In
our models, We use α_j to denote the fraction of internal resources allocated to the j-th
cellular function, with $\vec{\alpha} = (\alpha_1, \alpha_2 \dots)$ representing a metabolic strategy. An exact
metabolic trade-off is assumed, such that $\sum_j \alpha_j = 1$. For example, Figure 2A shows a
simple metabolic model with two substitutable nutrients a and b, such as glucose and
galactose (see Methods for details), that contribute linearly to biomass increase. Since a
substantial investment of protein and energy is required for nutrient intake, the model
assumes a trade-off between the allocation of internal resources to import either
nutrient. Specifically, a fraction α_a of resources is allocated to import a and a fraction α_b
(= $1 - \alpha_a$) to import b. All values of α_a from 0 to 1 define a continuous spectrum of
metabolic strategies. How then shall we evaluate these strategies given that a single
species adopting any one of these strategies will grow at exactly the same rate as
dilution in a steady-state chemostat?
282
283 ***Rule of invasion***
284 We use the outcome of competition between species to evaluate metabolic strategies,
285 assuming each species adopts a given strategy. In particular, we focus on invasion: the
286 introduction of a small number of an "invader" species to a steady-state chemostat
287 already occupied by an "indigenous" species.
288
In the graphical representation of species-environment interaction, the outcome of an
invasion can be summed up by a simple geometric rule, as demonstrated in Fig 2B and
C. The growth contour of the invader (species *Red*) separate the nutrient space into
two regions: an "invasion zone" where the invader grows faster than dilution (green-
colored region in Fig 2B and C), and "no-invasion zone" where the invader has a growth
rate lower than dilution. If the steady-state environment constructed by the indigenous
species (species *Blue*) is located within the invasion zone of the invader, the invader will
initially grow faster than dilution. Therefore, the invader will expand its population and
the invasion will be successful (Fig 2B). By contrast, if the steady-state nutrient
environment created by the indigenous species lies outside of the invasion zone, the
invasion will be unsuccessful (Fig 2C, same species but different supply condition
therefore different steady state). (See Methods for details.)
301
Mutual invasion, flat fitness landscape, and unlimited coexistence
Such a rule of invasion does not guarantee transitivity of competitiveness. That is, if
species *Red* can invade species *Blue*, that does not mean *Blue* cannot invade *Red*.

Figure 2D shows an example of mutual invasibility. With a supply condition different
from those in Fig 2B and C, while the steady-state environment created by *Blue* is
located within the invasion zone of *Red*, the steady-state environment created by *Red* is
also located within the invasion zone of *Blue*. According to the rule of invasion, each
species can therefore invade the steady-state environment created by the other. In the
face of such successful invasions, the only possible stable nutrient environment for this
system is at the intersection of two growth contours, where the two species can coexist.
312
This mutual invasion can be readily understood within a "fitness-landscape" picture.
Given an environment, we define the fitness landscape as the relation between the
instantaneous growth rate and the metabolic strategy of any invader (Eq. (S8)-(S9), see
Methods for details). Different environments give rise to different fitness landscapes. In
the steady-state environment created by *Red* ($\alpha_a = 0.6$), strategies with smaller α_a have
higher fitness (Fig 2D, upper inset, red curve). In the steady-state environment created
by *Blue* ($\alpha_a = 0.2$), strategies with larger α_a have higher fitness (Fig 2D, upper inset,
blue curve). Therefore, each species creates an environment that is more suitable for its
competitor, which leads to coexistence.
322
For the environment co-created by *Blue* and *Red* (Fig 2D, purple dot), the fitness
landscape becomes flat (Fig 2D, upper inset, purple curve): species with any metabolic
strategy will grow at the same rate as dilution in this environment. Therefore, in this
system, once a pair of species with a mutual-invasion relationship construct the nutrient
environment together, all species become neutral and can coexist indefinitely (see
Methods for details). This graphical approach to mutual invasion and the flat fitness
landscape provide an intuitive representation of species self-organizing to a state of
unlimited coexistence beyond competitive exclusion, as first reported by Posfai et al.
(Posfai et al., 2017).
332
Rock-paper-scissor invasion loop and oscillation
Resource-competition models focusing on various aspects of cellular metabolism vary
in their assumptions regarding $g(\vec{c}, \vec{x}, \vec{\alpha})$, $I(\vec{c}, \vec{x}, \vec{\alpha})$, and $f(\vec{c}, \vec{x}, \vec{\alpha})$, and can lead to
diverse results for community structure and coexistence. However, the above general
"rule of invasion" allows us to treat these divergent resource-competition models in a
unified framework. In the following example, we utilized a metabolic model slightly
different from that in Fig 2, to show that a dynamic fitness landscape is indispensable
for coexistence.
341
In the metabolic model shown in Fig 3A, three substitutable nutrients, *a*, *b*, and *c*,
contribute additively to cell growth. In this three-dimensional nutrient space, the growth
contour for each species is a two-dimensional surface (Fig 3B). In addition to requiring
enzymes to import the raw forms of these nutrients as in the model of Fig 2A, enzymes
are also required to convert the imported raw materials into biomass. In this model, a
six-element $\vec{\alpha}$ is required to describe the metabolic strategy, and there is the possibility
of "mismatches" in the fraction of internal resources allocated to import and to convert
the same nutrient. Such mismatches can produce a "rock-paper-scissor" invasion loop
(Fig S1A): In the environment created by species 1, species 2 has a higher fitness but

not species 3; therefore species 2 can invade species 1 and establish its own
environment; however, this environment lies within the invasion zone of species 3 (Fig
3B) but not of species 1, therefore species 3 subsequently invades; then in turn, species
3 create an environment where species 1 has the highest fitness. Such a loop of
invasions leads to oscillatory population dynamics (Fig 3C, upper panel), with an ever-
changing fitness landscape (Fig 3C, lower panel).
357
358 Oscillation and even chaos in resource-competition model have been demonstrated by
359 Huisman et al. (Huisman et al., 1999), and shown to allow dynamical coexistence
360 beyond competitive exclusion. The simple model presented here illustrates how
361 oscillation can be understood as a loop of invasion creating an ever-changing fitness
362 landscape.
363
Multi-stability, chain of invasion, and non-invasible strategies
When species create environments that are more favorable for their competitors,
mutual-invasion and oscillations can occur. Can species create environments that are
hostile to their competitors, and if so what will be the consequences?
Fig 4A shows a simple metabolic model with two essential nutrients *a* and *b*, such as
nitrogen and phosphorus (see Methods for details). Similar to the model in Fig 2A, the
model assumes a trade-off between the allocation of internal resources to import
nutrients, so that a resource allocation strategy is fully characterized by the fraction of
resources α_a allocated to import nutrient *a*. The growth rate is taken to be the minimum
of the two input rates (Odum & Barrett, 1971). As shown in Fig 4B, two species, *Red*
and *Blue*, each creates a nutrient environment outside of the invasion zone of each
other. According to the rule of invasion, neither can be invaded by the other. Therefore,
the steady state of the community depends on initial conditions – whichever species
occupies the chemostat first will dominate indefinitely.
378
From the perspective of the strategy-growth relationship (Fig 4B, inset), species *Red*
($\alpha_a = 0.65$) creates a fitness landscape where small α_a is disfavored. Symmetrically,
species *Blue* ($\alpha_a = 0.35$) creates a fitness landscape where large α_a is disfavored.
However, neither *Red* nor *Blue* sits on the top of the fitness landscape each one creates
(Fig 4C). In the fitness landscape created by *Blue*, a slightly larger α_a (green diamond in
Fig 4C) has the highest growth rate. Consequently, species adopting the *Green* strategy
can invade *Blue*. Nevertheless, species *Green* is not on the top of its own fitness
landscape as an even larger α_a (yellow diamond in Fig 4C) maximizes the growth rate
in the environment created by *Green*. A series of replacements by the fastest-growing
species in the environment created by the former species creates a chain of invasion.
389
It is worth noting that in this model after four steps of replacement, bistability appears.
The species with α_a marked by *Deep Purple*, which is reached by the chain of invasion
going from *Blue*, to *Green*, to *Yellow*, to *Deep Green*, cannot invade the original species
Blue. A similar relationship holds between *Cyan* and *Red*. This phenomenon highlights
the difference between ecological stability and evolutionary stability: Ecologically, as
both *Blue* and *Deep Purple* create a fitness landscape where the other species grows
slower than dilution, they constitute a bistable system. However, evolutionarily, mutants

with slightly larger α_a can invade *Blue,* bringing the system towards *Deep Purple* until
bistability collapses.
399
In this model, with symmetric parameters, the only evolutionarily stable strategy is $\alpha_a =$
0.5 (black diamond in Fig 4C). This is the only strategy that locates itself on the top of
the fitness landscape it creates, and therefore cannot be invaded by any other species.
This simple model demonstrates a general definition of evolutionarily stable (aka
optimal or non-invasible) strategies: those strategies that create a fitness landscape
which places themselves on the top (Eq. (S10)).
406
A nutrient environment defines a fitness landscape, and the steady-state nutrient
environment created by species is influenced by supply condition, dilution rate, and cell
metabolism. Therefore, different chemostat parameters and different metabolic models
lead to different optimal strategies. In the following, we described a generally applicable
protocol for obtaining the non-invasible strategies, using the metabolic model in Fig 4A
as the example (Fig 4D, details in Methods):
First, under a nutrient environment $\vec{c}$, the maximal growth rate $g_{\max}(\vec{c})$ (background
color in Fig 4D) and the corresponding resource allocation strategy $\vec{\alpha}_{\max}(\vec{c})$ can be
obtained analytically or via numerical search through the strategy space (Eq. (S11)).
$g_{\max}(\vec{c})$ and $\vec{\alpha}_{\max}(\vec{c})$ are independent of the chemostat parameters $\vec{c}_{\text{supply}}$ and d.
Second, the maximal growth contour for dilution rate d is defined as all nutrient
environments $\vec{c}$ that support a maximal growth rate of d (Eq. (S12)). Different
maximizing strategies $\vec{\alpha}_{\max}(\vec{c})$ exist at different points of the maximal growth contour, as
shown by the colors of the curve in Fig 4D. By definition, the maximal growth contour
envelops the growth contour of any single strategy, and nutrient environments on the
maximal growth contour are outside of the invasion zone of any strategy. Therefore, if a
species is able to create a steady-state environment on the maximal growth contour, it
cannot be invaded. Finally, different $\vec{c}_{\text{supply}}$ form different maximal flux-balance curves
(Eq. (S14)), which intersect with the maximal growth contour at one point $\vec{c}_{\text{opt}}$. Species
$\vec{\alpha}_{\max}(\vec{c}_{\text{opt}})$ that adopt the maximizing strategy at $\vec{c}_{\text{opt}}$ create the environment $\vec{c}_{\text{opt}}$, and
are therefore immune to invasion. Under different $\vec{c}_{\text{supply}}$, different species become
non-invasible (orange, green, and blue growth contours in Fig 4D), and the supply lines
emanating from different points on the maximal growth contour indicate the supply
conditions for which the corresponding strategies are evolutionarily stable.
431
Evolutionarily stable coexistence
Given d and $\vec{c}_{\text{supply}}$, the maximal growth contour and the maximal flux-balance curve
are unique, therefore there is only one $\vec{c}_{\text{opt}}$. Does the uniqueness of $\vec{c}_{\text{opt}}$ imply a single
evolutionarily stable species? Or is coexistence still possible even in the face of
evolution? In a recent work (Taillefumier et al., 2017), this question was addressed by
modeling a population of microbes competing for steadily supplied resources. Though
in-silico evolution and network analysis, the authors found that multiple species with
distinct metabolic strategies can coexist as evolutionarily-stable co-optimal consortia,
which no other species can invade.
441

Using a simplified version of Taillefumier et al.'s model (Fig 5A), we employ the
graphical approach to help identify the requirements for such evolutionarily-stable
coexistence and the role of each species in supporting the consortium. In this model, at
the cost of producing the necessary enzymes, cells are not only able to import external
nutrients, but can also convert any one of the internal nutrients into any other.
Meanwhile, nutrients passively diffuse in and out of the cell. The internal concentrations
of nutrient *a* and nutrient *b* are both essential for cell growth (see Methods for detail).
Therefore, metabolic trade-offs in this system have four elements: the fraction of internal
resources allocated to import nutrient *a* (α_a) or nutrient *b* (α_b) and/or convert one
nutrient into another (α_{ab} converts internal *b* into *a,* and α_{ba} converts internal *a* into *b*).
Each species is defined by its internal resource allocation strategy $\vec{\alpha} = (\alpha_a, \alpha_b, \alpha_{ab}, \alpha_{ba})$.
453
Following the general protocol described in the previous section, we first identified the
maximal growth rates $g_{\max}(\vec{c})$ and the corresponding strategy or strategies $\vec{\alpha}_{\max}(\vec{c})$ at
each point $\vec{c}$ in the nutrient space, and generated maximal growth contours for different
dilution rates (Fig 5B). The maximal growth contours are not smoothly continuous, nor
are the corresponding strategies. In nutrient space, three distinct sectors of maximizing
strategies appear (Fig 5B, Fig S3A): When nutrient *a* is very low compared to *b*, the
maximizing strategy is a "*b-a* converter" which imports *b* and converts it into *a* (blue
sector, only α_b and α_{ab} are non-zero). Symmetrically, when *a* is comparatively high, the
optimal strategy is a "*a-b* converter" (green sector, only α_a and α_{ba} are non-zero).
Otherwise, the maximizing strategy is an "importer" which imports both nutrients without
conversion (red sector, only α_a and α_b are non-zero). On the border between sectors,
the maximal growth contour has a discontinuous slope.
466
Evolutionarily stable coexistence occurs at these discontinuous points. If an
environment point $\vec{c}_0$ is located in a continuous region of the maximal growth contour,
only one maximizing strategy $\vec{\alpha}_{\max}(\vec{c}_0)$ exists for that environment (maximizing
strategies along the maximal growth contour are indicated by colored squares in Fig
5C). Supply conditions that make $\vec{\alpha}_{\max}(\vec{c}_0)$ the optimal strategy (i.e. allow $\vec{\alpha}_{\max}(\vec{c}_0)$ to
create the steady-state environment $\vec{c}_0$) constitute the supply line for $\vec{c}_0$ and $\vec{\alpha}_{\max}(\vec{c}_0)$.
However, at the discontinuous points of the maximal growth contour, where two classes
of strategies meet, two different strategies are both maximizing. For example, at the
purple dot in Fig 5C a strategy belonging to the "*b-a* converter" class (species *Blue*) and
one belonging to the "importer" class (species *Red*) are both maximizing strategies.
Each strategy derives a supply line from the purple dot (black dashed line, Fig 5C). The
two supply lines span a gray region where no supply line from any single strategy
enters. Correspondingly, for any supply conditions inside the gray region, no single
species can alone create an environment on the maximal growth contour. For example,
under the supply condition shown by the black open circle in the gray region, species
Blue and species *Red* both create nutrient environments that lie within the maximal
growth contour (blue and red dots, Fig 5C), and are thus subject to invasion by other
species. Nevertheless, the species-specific growth contours of *Blue* and of *Red*
intersect at the purple point on the maximal growth contour. Therefore, only when *Blue*
and *Red* coexist can they co-create an environment on the maximal growth contour,
and thus resistant to invasion from any other species. Indeed, when we simulate

multiple species with different maximizing strategies under the supply condition
indicated by the open black circle, species *Blue* and species *Red* are the only two that
survive (Fig 5C, inset).
491
The optimal coexistence of species *Blue* and species *Red* can be understood intuitively
from the dynamic fitness landscape. Given a nutrient environment, the relation between
α_a and growth rate of importer (red curve) or *a-b* converter (green curve), and that
between α_b and growth rate of *b-a* converter (green curve) constitute the fitness
landscape of species adopting different possible maximizing strategies (Fig 5D). In the
environment created by species *Blue* (blue dot in Fig 5C), not only will some importers
grow faster than *Blue*, species *Blue* (strategy marked by blue diamond) is not even on
the fitness peak of its own class (Fig 5D, upper box). Similarly, in the environment
created by species *Red*, the strategy of *Red* is not at the top of the fitness landscape
(Fig 5D, middle box). By contrast, in the environment co-created by species *Blue* and
Red (purple dot in Fig 5C), their strategies are at the top of the fitness landscapes of
their own classes and at equal height. For all supply conditions in the gray region,
species *Blue* and species *Red* jointly drive the nutrient concentrations to the
discontinuous point of the optimal growth contour, and thereby achieve evolutionarily
stable coexistence.
507
Species creating a new nutrient dimension
As discussed in the introduction, one possible solution to the paradox of plankton is the
creation of new nutrient "dimensions" by species secreting metabolites that can be
utilized by other species. For example, *E. coli* secretes acetate as a by-product of
glucose metabolism. Accumulation of acetate impedes the growth of *E. coli* on glucose
(Luli & Strohl, 1990), but the acetate can be utilized as a carbon source by mutant
strains that emerge in long-term evolution experiments (D'Souza et al., 2018;
Rosenzweig, Sharp, Treves, & Adams, 1994).
516
To explore the possibilities of evolutionarily stable coexistence when species create
new nutrients, we used a simplified model to represent multi-step energy generation
with a dual-role intermediate metabolite (Fig 6A). A single chemical energy source S is
supplied into the chemostat. The pathway for processing S consists of four relevant
reactions driven by designated enzymes: External S can be imported and converted into
intermediate I_{int} to generate ATP (with corresponding fraction of enzyme budget
α_{ATP1}).The intermediate has a dual role in energy production: on the one hand, it
positively contributes to ATP production via a downstream reaction (with fraction of
enzyme budget α_{ATP2}); on the other hand, it negatively contributes to ATP production
through product inhibition of the first energy-producing reaction. To deal with the
negative effect of internal intermediate, cells may synthesize transporters to export
intermediate out into environment, where it becomes external intermediate I_{ext} (fraction
of enzyme budget α_{exp}). By this reaction, cell can increase the dimension of nutrient
space from one (S) into two (S and I_{ext}). Cells can also import I_{ext} into I_{int} (fraction of
enzyme budget α_{imp}), then use I_{int} as an energy source via the second reaction. (See
Methods for details.)
533

The metabolic strategy $\vec{\alpha}$ in this model has four components: $\vec{\alpha} = (\alpha_{\mathrm{ATP1}}, \alpha_{\mathrm{ATP2}}, \alpha_{\exp},$
$\alpha_{\mathrm{imp}})$. When we examine the maximizing strategies and maximal growth rates in the
nutrient space, three distinctive classes of strategy emerge (Fig 6B). When S is
abundant and I_{ext} is low, the maximizing strategies have only two non-zero components,
α_{ATP1} and $\alpha_{\exp}$ (Fig S3B), meaning this class of species only imports S for the first
energy-generating reaction then exports intermediate as waste. Therefore, we call
strategies in this class "polluters" (blue section in Fig 6B, Fig S3C). When I_{ext} is high
while S is low, the maximizing strategies have two different non-zero components, α_{ATP2}
and α_{imp} (Fig S3B), meaning this class of species solely relies on I_{ext} as its energy
source. We call these strategies "cleaners" as they clean up the I_{ext} in the environment,
which is detrimental to the polluters (green section in Fig 6B, Fig S3C). When there are
comparable amounts of S and I_{ext} present, a third class of maximizing strategies
appears: these cells neither export nor import intermediates, but rather allocate all
enzyme budget to α_{ATP1} and α_{ATP2} to carry out both energy-producing reactions. We
call species in this class "generalists" (red section in Fig 6B, Fig S3C).
549
As shown in Fig 6B, on the borders between classes of strategies in nutrient space, the
maximal growth contours turn discontinuously. These points of discontinuity, as in the
previous section, are nutrient environments corresponding to evolutionarily stable
coexistence of species from distinct metabolic classes. The classes of optimally
coexisting species change with dilution rate. When the dilution rate is low ($d = 0.4$, Fig
6C), at the discontinuous point of the maximal growth contour, the corresponding two
maximizing strategies are one polluter (species *Blue*) and one cleaner (species *Green*).
Their supply lines span a gray region where both species *Blue* and species *Green* are
required to create a steady-state environment on the maximal growth contour. As we
are only supplying the system with S, the supply condition always lies on the *x*-axis of
concentration space. For the supply condition shown by the black open circle in Fig 6C,
polluter *Blue* creates a nutrient environment (blue dot) far from the maximal growth
contour. When the cleaner *Green* is added to the system, not only does the biomass of
Blue increase (inset), but also the steady-state nutrient environment moves to the
discontinuous point of the maximal growth contour (cyan dot), where both *Blue* and
Green occupies the peaks of their fitness landscapes (Fig 6D). This result is consistent
with the long-term evolution experiment of *E. coli* and also intuitive: polluter *Blue* and
cleaner *Green* form a mutually beneficial relationship by, respectively, providing
nutrients and cleaning up waste for each other, thereby reaching an optimal cooperative
coexistence.
570
A quite different coexistence occurs at higher dilution rate ($d = 0.6$, Fig 6E). Growth
contours at this dilution rate show two turning points, but neither are between the
polluter and the cleaner class. One discontinuous point is between the cleaner class
(green squares) and the generalist class (red squares), but the gray region spanned by
the corresponding supply lines does not cover the *x*-axis and so does not represent an
attainable coexistence when only S is supplied. The other discontinuous point is
between the generalist class and the polluter class (blue squares). The gray region
spanned by the supply lines of the corresponding two maximizing strategies of
generalist class (species *Red*) and polluter class (species *Blue*) does cover the *x*-axis.

Therefore, a supply condition with only S within the gray region (e.g., the black open
circle) leads to the optimal coexistence of generalist *Red* and polluter *Blue* on the
discontinuous point (purple dot), despite the fact that they do not directly benefit each
other. Indeed, when the generalist *Red* is added to a system with polluter *Blue* and a
cleaner *Green*, the cleaner *Green* goes extinct and the biomass of the polluter *Blue*
decreases (inset). Nevertheless, the steady-state nutrient environment is moved from a
cyan dot lying inside the maximal growth contour to the purple dot lying on the maximal
growth contour. In the environment of the cyan dot created by cleaner *Green* and
polluter *Blue*, *Blue* is not on the top of the fitness landscape of the polluter class (Fig 6F,
upper box). By contrast, for the fitness landscape created by polluter *Blue* and
generalist *Red* (Fig 6F, lower box), despite being lower in biomass, *Blue* occupies the
top of the landscape. Therefore, the optimal coexistence of this polluter and this
generalist does not arise from direct cooperation, but rather from collaborating to defeat
other competitors.

DISCUSSION

The phrase "survival of the fittest" used to describe natural selection can be applied both to competition within species and to species competing in the same environment. One doctrine governing resource competition among species is the competitive exclusion principle: theoretically, there should be no more surviving species than the number of resources. However, the enormous diversity of coexisting species in the natural world seems to contradict the competitive exclusion principle. This so-called "paradox of the plankton" has stimulated many theoretical models of resource competition, each with its own assumptions and different conclusions. In this work, we examined a range of models for metabolic competition among microbes within a unified framework, using the species-environment feedback as an organizing principle and the geometric "rule of invasion" and dynamic fitness landscapes as common tools. Under this unified framework, it becomes apparent how metabolic tradeoffs promote diversity by allowing a dynamic fitness landscape without a fittest peak. The implications of non-static landscapes have been under discussion (de Visser, Elena, Fragata, & Matuszewski, 2018) ever since the introduction of fitness landscapes by Wright (Wright, 1932). Nonetheless, in most of these studies, variability in fitness is introduced by externally-imposed environmental fluctuations (Mustonen & Lässig, 2009; Zhang, 2012). In our work, we focused on how fitness landscapes can be intrinsically dynamic due to species' own actions on their environment.

Graphical representations of resource-competition models are not new. The school of contemporary niche theory dates back almost 50 years (MacArthur, 1970; David Tilman, 1980, 1982), and the growth contours in our work reduce to the zero-net growth isoclines (ZNGI) introduced by this school in the particular case of two growth-promoting resources. Our framework, nonetheless, differs in several aspects: First, we focused on metabolic models with trade-offs, for which there are not only different ZNGIs for different species but a continuous family of growth contours, and the envelope of all growth contours is the maximal growth contour. Given these resource allocation trade-offs, the growth contours of any pair of species must intersect, clearly demonstrating why metabolic trade-offs prevent a single species from unconditional dominance. Moreover, the definition of growth contours is not limited by the number of resources, nor constrained by whether external chemical concentrations contribute positively or negatively to growth, making the approach suitable to address more realistic metabolic models. Second, the introduction of the flux-balance curve, in additional to the supply line, makes it easier to determine the species-specific environment for a given supply condition, which is particularly useful for determining the non-invasible environment as the intersection between maximal growth contour and the maximal flux-balance curve. In brief, our new graphical approach is well suited to our goals of understanding coexistence from the perspective of species-environment feedback, demonstrating how the fitness landscape is changed by the species present, and identifying evolutionarily stable strategies.

Our work is not aimed at adding another solution to the paradox of the plankton. Rather we provide a graphic tool to unify several approaches, and suggest how different

proposed solutions to the paradox can emerge intrinsically from competition for
resources. When nutrients are substitutable, resource competition among species with
metabolic trade-offs has been shown to lead to emergent neutrality (Posfai et al., 2017),
as the fitness landscape is made flat by the competing species. Other commonly
invoked solutions to the paradox of the plankton are extrinsic temporal and/or spatial
heterogeneity. In this work, we showed that both types of heterogeneity can also
emerge intrinsically from species-environment feedback. When the rule of invasion
allows non-transitive loops, oscillations and chaos can occur, which have been shown
to allow coexistence beyond competitive exclusion (Huisman & Weissing, 1999, 2001).
In addition, when multiple nutrients are all essential, the ability of each species to create
an environment that favors itself allows for the spontaneous emergence of spatial
heterogeneity in an extended system (Fig S2).
651
Yet, even if fixed species can coexist in an ecosystem, will coexistence survive the
ceaseless process of mutation and adaptation? Our approach provides a general
protocol to determine non-invasible/evolutionarily stable metabolic strategies, which we
demonstrate in the context of three different metabolic models. In these examples, there
is a unique non-invasible solution at the intersection of the maximal growth contour and
the maximal flux-balance curve. Nevertheless, other models for species competition
suggest multiple or, in some cases, zero evolutionarily stable outcomes. For example, in
a metabolic model with multiple essential nutrients, Goyal et al. (Goyal, Dubinkina et al.
2018) found multistability in a discrete strategy space, with each steady state non-
invasible. By contrast, a study based on a Lotka-Volterra model (in which resource
competition is taken to contribute to direct interactions between species (Letten et al.,
2017)) found that evolution via continuous introduction of new species drives the system
into an chaotic state with a large number of species, i.e. coexistence without a steady
state (Ackland & Gallagher, 2004). In the future it will be worthwhile to employ graphical
approaches to investigate what types of metabolic models with various shapes of
growth contours and flux-balance curves can lead to such non-unique non-invasible
outcomes of resource competition.
669
Another advantage of our graphical approach, besides providing an intuitive picture of
species competition, is that it can help understand and control nutrient limitation in
chemostat experiments. The capacity of species to shape their own environment, even
in a system as simple as a chemostat, presents challenges to controlling which nutrient
or nutrients are limiting. By traditional definition, if increasing a certain nutrient leads to
an increase of a cell's growth rate, that nutrient is considered "limiting". However,
growth rate is invariant in a chemostat, being set experimentally by the dilution rate, so
inferring nutrient limitation requires special attention. For example, if one sees the same
cellular responses under different nutrient supplies, what can one conclude? Cells may
be creating the same nutrient environment out of different supply conditions (*cf*. Fig 1D),
or alternatively cells may be transducing different nutrient environments into the same
physiological response through mechanisms such as "ratio sensing" (Escalante-Chong
et al., 2015). Our graphical approach combined with direct measurements of steady-
state nutrient concentrations in the chemostat can precisely define and help control
nutrient limitation (Boer, Crutchfield, Bradley, Botstein, & Rabinowitz, 2010)(Boer et al.,

2010). As described above, changes in supply concentrations shift the flux-balance
curve, but do not change the shape of the growth contour. Therefore, by experimentally
varying the supply conditions and measuring the nutrient environment created by cells,
the shape of the growth contour can be obtained. The resulting slope of the growth
contour provides information on nutrient limitation even in the absence of detailed
knowledge about a cell's metabolism. For example, in the nutrient a - nutrient b plane, a
near-horizontal growth contour indicates b-limited growth while a near-vertical growth
contour means a-limited growth, and an intermediate slope implies that the two nutrients
are co-limiting.
694
Many future directions can follow this work. From the perspective of experiment, our
framework can assist in analyzing and interpreting results of microbial evolution in the
lab (Van den Bergh, Swings, Fauvart, & Michiels, 2018), where the continual
emergence of new mutants under defined experimental conditions suggests an
intrinsically dynamic fitness landscape. From the perspective of theory, we do not yet
have a rigorous mathematical theorem about the conditions for discontinuity of the
maximal growth contour, nor proof that discontinuity necessarily leads to evolutionarily
stable coexistence. Theoretical developments paralleling those on the general existence
of ecologically stable states (De Leenheer, Levin, Sontag, & Klausmeier, 2006;
Marsland III, Cui, & Mehta, 2019) would bring a more comprehensive understanding of
evolutionarily optimal states in metabolic models. Besides, the metabolic models
considered in this work are highly simplified. Going forward, more detailed and
experimentally-based models can be examined using the same graphical framework.

METHODS

Programs for this work is coded in MATLAB R2018a. A repository of all tools used to
generate results in this program can be found at:
https://github.com/zhiyuanli1987/Qbiotoolbox.git
712
713 ***Total RNA and total protein measurements***
714 The method for total RNA and protein measurements is described in (Li et al., 2018).
715
716 ***Supplemental Table 1: Symbols***

Chemostat parameters	
$\vec{c}_{\text{supply}} = \left(c_{1,\text{ supply}}, c_{2,\text{ supply}}, \dots c_{p,\text{ supply}}\right)$	Nutrient supply. $c_{i,\text{ supply}}$ is the concentration of the i-th nutrient in the supply.
d	Dilution rate (same as supply influx rate to keep volume fixed).
Chemostat variables	
$\vec{c}_k = \left(c_{1,k}, c_{2,k}, \dots c_{p,k}\right)$	Nutrient environment inside the k-th chemostat. $c_{i,k}$ is the concentration of the i-th nutrient within the medium of the k-th chemostat. All possible $\vec{c}$ constitute the "nutrient space".
$m_{\sigma,k}$	Biomass density of species σ in the k-th chemostat.
Species-specific quantities	
$\vec{\alpha}_\sigma = (\alpha_{1,\sigma}, \alpha_{2,\sigma} \dots)$	Resource allocation strategy of species σ. $\alpha_{j,\sigma}$ is the fraction of resources allocated to the j-th cellular function by species σ.
$\vec{q}_\sigma$	Intracellular concentrations of growth-related metabolites for species σ.
$g(\vec{c}, \vec{q}, \vec{\alpha})$	Growth rate as a function of $\vec{c}$, $\vec{q}$, and $\vec{\alpha}$.
$I_i(\vec{c}, \vec{q}, \vec{\alpha})$	Intake rate per biomass of the i-th nutrient as a function of $\vec{c}$, $\vec{q}$, and $\vec{\alpha}$. $I_i(\vec{c}, \vec{q}, \vec{\alpha})$ can be negative to describe cells exporting secondary metabolites.
r	Biomass concentration within a cell, taken to be a constant that always equal to 100.
$\vec{f}(\vec{I}(\vec{c}, \vec{q}, \vec{\alpha}), \vec{q}, \vec{\alpha})$	Functions defining the changing rate of intracellular metabolite concentrations $\vec{q}$, as a function of $\vec{c}$, $\vec{q}$, and $\vec{\alpha}$.
GC_σ	Growth-rate contour of species σ.
FB_σ	Flux-balance curve of species σ.
$\vec{c}_{\sigma,\text{ ss}}$	The steady-state environment created by one species σ.

$SL_\sigma(\vec{c})$	The supply line for species σ in environment $\vec{c}$.
$\{\sigma^*\}$	A set of species stably surviving in chemostat. A set can contain one or more species.
$\vec{c}_{\{\sigma^*\},ss}$	The steady-state environment created by a set of species $\{\sigma^*\}$.

718 *Metabolic model and resource allocation strategy*
719 In modeling population dynamics in a chemostat, multiple assumptions need to be
720 made concerning how cells sense the environment, import nutrients, export metabolites,
721 utilize resources, and grow in biomass. Different assumptions result in different
722 metabolic models. Some metabolic models focus on trade-offs in resource allocation, as
723 the amount of resources "owned" by a cell, including proteins and energy, is limited.
724 Cells need to allocate these limited resources into different cellular functions, such as
725 metabolism, gene expression, reproduction, motility, maintenance, etc. We use $\alpha_{j,\sigma}$ to
726 represent the fraction of resources allocated to the j-th cellular function of species σ,
727 with $\vec{\alpha}_\sigma = (\alpha_{1,\sigma}, \alpha_{2,\sigma} \ldots)$ representing the resource allocation strategy of species σ. For
728 simplicity, we assume each species has a fixed resource allocation strategy.

730 *Dynamic equations for a single species in a chemostat*
731 In a chemostat with nutrient supply $\vec{c}_{\text{supply}}$, dilution rate d and a single species σ with
732 fixed strategy $\vec{\alpha}_\sigma$ and intracellular metabolite concentration $\vec{q}_\sigma$, the cell biomass density
733 m_σ and the chemostat nutrient concentrations $\vec{c}$ are generally described by the following
734 equations:

$$\frac{dm_\sigma}{dt} = m_\sigma \cdot (g(\vec{c}, \vec{q}_\sigma, \vec{\alpha}_\sigma) - d), \tag{S1}$$

$$\frac{d\vec{c}}{dt} = d \cdot (\vec{c}_{\text{supply}} - \vec{c}) - m_\sigma/r \cdot \vec{I}(\vec{c}, \vec{q}_\sigma, \vec{\alpha}_\sigma). \tag{S2}$$

735 In considering the details of cellular metabolism, one may choose to incorporate the
736 dynamics of intracellular metabolites that originate from nutrient import and influence
737 cell growth. We make the assumption that the biomass concentration r, e.g. protein
738 concentration, is constant for cells under all growth conditions. Thus, an increase of
739 total cell mass induces a linear increase of total cell volume. m_σ is the cell mass per
740 volume in the chemostat, and r is the cell mass per volume within a cell. For a
741 chemostat-to-cell flux of mass J, the concentration of the metabolite in chemostat
742 decrease by $J/V_{chemsotat}$ while the concentration in cell increase by J/V_{cell}. As a result,
743 the metabolites imported into cells are enriched by a factor of r, and metabolites
744 secreted by cells are diluted by $1/r$. Also, all metabolites are diluted by cellular growth,
745 which is generally a slow process compared to metabolic reactions and can be ignored
746 in most cases. We use a function $\vec{f}(\vec{I}(\vec{c}, \vec{q}, \vec{\alpha}), \vec{q}, \vec{\alpha})$ to represent the rate of change of
747 intracellular metabolite $\vec{q}_\sigma$:

$$\frac{d\vec{q}_\sigma}{dt} = \vec{f}(\vec{I}(\vec{c}, \vec{q}_\sigma, \vec{\alpha}_\sigma), \vec{q}_\sigma, \vec{\alpha}_\sigma). \tag{S3}$$

748 Where Eq. (S2) represents p equations for p types of nutrients, and Eq. (S3) represents
749 h equations for h growth-related intracellular metabolites.

Species-specific steady state
In the steady state of chemostat, Eqs. (S1)- (S3) should be all equal to zero.
For intracellular metabolites, as $\vec{f}\left(\vec{I}(\vec{c}, \vec{q}_\sigma, \vec{\alpha}_\sigma), \vec{q}_\sigma, \vec{\alpha}_\sigma\right) = 0$ as a result of Eq. (S3)=0,
given an environment $\vec{c}$, the steady state of $\vec{q}_\sigma$ can be expressed as a function of $\vec{c}$:
$\vec{q}_\sigma^* = \vec{f}^{-1}(\vec{c}, \vec{\alpha}_\sigma)$.
Growth contour (GC): From the perspective of the environment influencing species, at
each constant environment, the steady-state growth rate is fully determined by $\vec{c}$:
$g^*(\vec{c}, \vec{q}_\sigma) = g\left(\vec{c}, \vec{f}^{-1}(\vec{c}, \vec{\alpha}_\sigma)\right)$. If the biomass of a species is non-zero ($m \neq 0$), Eq. (S1)
requires $g^* = d$. In the p-dimensional nutrient space, this requirement defines a $(p - 1)$-
dimensional surface, constituted by all environments $\vec{c}$ that support an equal-to-dilution
growth rate. This surface reduces to the zero-growth isoclines in contemporary niche
theory when the nutrient space is two-dimensional and the growth rate g solely relies on
$\vec{c}$ monotonically, but is not necessarily limited by the nutrient dimension or the form of
the growth function. For convenience, we name this surface the "growth contour" (GC)
for species σ:

$$GC_\sigma := \{\vec{c} \mid g\left(\vec{c}, \vec{f}^{-1}(\vec{c}, \vec{\alpha}_\sigma), \vec{\alpha}_\sigma\right) = d\}. \qquad \textbf{(S4)}$$

766 An example of growth contours is shown in Fig 1D.
767
Flux-balance curve (FB): Eq. (S2) describes how species act on the environment. In
steady state, the influx, out-flux, and consumption by species should be balanced for
each nutrient, which enables calculation of the biomass density-to-dilution ratio for every
i: $\frac{m_\sigma}{d \cdot r} = \frac{c_{i,\,\text{supply}} - c_i}{I_i\left(\vec{c}, \vec{f}^{-1}(\vec{c}, \vec{\alpha}_\sigma), \vec{\alpha}_\sigma\right)}$. For a p-dimensional nutrient space, there are p equations for the
same value of $\frac{m}{d \cdot r}$. This leads to a one-dimensional curve in the nutrient space, which we
name the "flux-balance curve" (FB), defined as:

$$FB := \{\vec{c} \mid \frac{c_{i,\,\text{supply}} - c_i}{I_i\left(\vec{c}, \vec{f}^{-1}(\vec{c}, \vec{\alpha}_\sigma), \vec{\alpha}_\sigma\right)} = \frac{m_\sigma}{d \cdot r} \text{ AND } c_i < c_{i,\text{supply}}\}. \qquad \textbf{(S5)}$$

For example, for two nutrients a and b, the flux balance curve is: $\frac{c_{a,\,\text{supply}} - c_a}{I_a\left(\vec{c}, \vec{f}^{-1}(\vec{c}, \vec{\alpha}_\sigma), \vec{\alpha}_\sigma\right)} -$
$\frac{c_{a,\,\text{supply}} - c_a}{I_a\left(\vec{c}, \vec{f}^{-1}(\vec{c}, \vec{\alpha}_\sigma), \vec{\alpha}_\sigma\right)} = 0$, as demonstrated in Fig 1D.
In nutrient space, the steady-state environment ($\vec{c}_{\sigma,\text{ss}}$) with non-zero biomass of species
σ will be located at the intersection of the growth contour and the flux-balance curve.
This environment is constructed by the species σ via its consumption of nutrients. If $\vec{c}_{\sigma,\text{ss}}$
exists, this species can survive in the chemostat. Otherwise, this species will be washed
out by dilution even without competition from other species. For the following
discussion, we only consider species that can survive when alone in a chemostat.
782
783 Supply line (SL): The flux-balance curve is determined by the supply condition $\vec{c}_{\text{supply}}$. In
784 many cases, it is helpful to derive the supply conditions that enable a species σ to
785 construct a steady-state environment $\vec{c}_{\sigma,\text{ss}}$. All possible values of $\vec{c}_{\text{supply}}$ that can
786 produce a given $\vec{c}_{\sigma,\text{ss}}$, form a straight line in the space of supply concentrations,
787 described by:

$$SL := \{\vec{c}_{\text{supply}} \mid \vec{c}_{\text{supply}} = \frac{m_\sigma}{d \cdot r} \cdot \vec{I}\left(\vec{c}_{\sigma,\text{ss}}, \vec{f}^{-1}(\vec{c}_{\sigma,\text{ss}}, \vec{\alpha}_\sigma), \vec{\alpha}_\sigma\right) + \vec{c}_{\sigma,\text{ss}}\}, \qquad \textbf{(S6)}$$

788   with varying non-negative values of m_σ/d. An example of a supply line is shown in
789   Fig 1D.
790
Dynamic equations for multiple species in a chemostat
In nutrient competition models, multiple species ($\sigma = 1 \dots n$) each with biomass density
m_σ compete for resources. They have species-specific growth rates $g(\vec{c}, \vec{q}_\sigma, \vec{\alpha}_\sigma)$ and
import rates $\vec{I}(\vec{c}, \vec{q}_\sigma, \vec{\alpha}_\sigma)$, yet all experience the same nutrient environment $\vec{c}$. Therefore,
Eq. (S1) and Eq. (S3) remain the same for each species, while the rate of change of
chemostat nutrient concentrations is influenced by the summed action of all species:

$$\frac{d\vec{c}}{dt} = d \cdot \left(\vec{c}_{\text{supply}} - \vec{c} \right) - \sum_{\sigma=1}^{n} m_\sigma/r \cdot \vec{I}(\vec{c}, \vec{q}_\sigma, \vec{\alpha}_\sigma). \tag{S7}$$

Multiple species steady state
Multiple species, even if each alone can survive in chemostat, do not generally coexist
when competing together. For a system starting with n different species, the stable
steady state contains n^* ($1 \leq n^* \leq n$) species with non-zero biomass. We define these
n^* surviving species as a stable species set $\{\sigma^*\}$, and mark the steady-state
environment created by this set as $\vec{c}_{\{\sigma^*\},\text{ss}}$. If $n > 1$, according to Eq. (S1), $\vec{c}_{\{\sigma^*\},\text{ss}}$ must be
located at the common intersection of growth contours formed by every species in $\{\sigma^*\}$.
805

Invasion
Invasion is defined as introducing a small number of invaders (with biomass density
m_{inv}) to a steady-state chemostat occupied by a set of local species. At the time of
introduction, if the invader can increase in biomass ($\frac{dm_{\text{inv}}}{dt} > 0$), the invasion is
successful; otherwise if the invader decreases in biomass ($\frac{dm_{\text{inv}}}{dt} < 0$), the invasion is
unsuccessful. If the biomass stays constant ($\frac{dm_{\text{inv}}}{dt} = 0$), the species is neutral with
respect to the local species.
In evaluating invasion by a species σ with strategy $\vec{\alpha}_\sigma$ of any environment $\vec{c}$, we make
two assumptions:
1. The biomass of the invader is so small that it does not disturb the environment at the
time of introduction.
2. There is a separation of timescales such that the concentrations of intracellular
metabolites reach equilibrium instantaneously at the time of introduction of the invader,
therefore Eq. (S3) is always equal to zero and $\vec{q}_\sigma = \vec{f}^{-1}(\vec{c}, \vec{\alpha}_\sigma)$ holds.
Therefore, we define the "invasion growth rate" of a species σ with strategy $\vec{\alpha}_\sigma$
introduced into environment $\vec{c}$ as:

$$g_{\text{inv}}(\vec{\alpha}|\vec{c}) = g\left(\vec{c}, \vec{f}^{-1}(\vec{c}, \vec{\alpha}), \vec{\alpha}\right). \tag{S8}$$

<u>Invasion zone</u>: By definition, the growth contour of the invader GC_{inv} divides the nutrient
space into two regions: an "invasion zone" that includes all environments where the
invader has an invasion growth rate higher than dilution, and "no-invasion zone" where
the invader has an invasion growth rate lower than dilution. If the steady-state
environment constructed by local species $\vec{c}_{\{\text{local}\},\text{ss}}$ is located within the invasion zone of

the invader, $g_{\mathrm{inv}}(\vec{\alpha}_{\mathrm{inv}}|\vec{c}_{\{\mathrm{local}\},\mathrm{ss}}) > d$, therefore $\frac{dm_{\mathrm{inv}}}{dt} > 0$ by Eq. (S1), and the invasion is
successful; otherwise, if $\vec{c}_{\{\mathrm{local}\},\mathrm{ss}}$ is located outside of the invasion zone of the invader,
$g_{\mathrm{inv}}(\vec{\alpha}_{\mathrm{inv}}|\vec{c}_{\{\mathrm{local}\},\mathrm{ss}}) < d$, and the invasion is unsuccessful. If $\vec{c}_{\{\mathrm{local}\},\mathrm{ss}}$ locate exactly on
the growth contour, it is neutral.
Two examples of this rule of invasion are presented in Fig 2A and 2B.
If the growth rate monotonically increases with the concentration of each nutrient, it can
be proven that the invasion zone is always above the growth contour of the invader (an
environment $\vec{c}_+$ "above" the growth contour GC_{inv} is defined as $\exists \vec{c}_0 \in GC_{\mathrm{inv}}, \mathrm{s.t.}\ c_{i,+} \geq$
$c_{i,0} \forall i$). If the growth rate is not monotonically increasing with nutrient concentrations,
identifying the invasion zone requires more model-specific analysis.
837
838 **Fitness landscape**
839 We quantified the fitness landscapes in the chemostat via the relationship between
840 metabolic strategies $\vec{\alpha}$ and the invasion growth rates of an invader adopting strategy $\vec{\alpha}$
841 in a given nutrient environment $\vec{c}$. Specifically,
842

$$Fitness\ landscape \coloneqq g_{\mathrm{inv}}(\vec{\alpha}|\vec{c}). \tag{S9}$$

843 Each environment $\vec{c}$ defines a fitness landscape. A set of species $\{\sigma^*\}$ constructs a
844 steady-state environment $\vec{c}_{\{\sigma^*\},\mathrm{ss}}$ and a corresponding fitness landscape $g_{\mathrm{inv}}(\vec{\alpha}|\vec{c}_{\{\sigma\},\mathrm{ss}})$.
845 Some examples of fitness landscapes are shown in Figs 2D, 3C, 4B-C, 5D and 6D-F.
846
847 **Non-invasible /optimal/ evolutionarily stable strategies**
848 A set of species $\{\sigma^*\}_{\mathrm{opt}}$ is non-invasible, aka optimal or evolutionarily stable, if no other
849 species can invade the steady-state environment constructed by $\{\sigma^*\}_{\mathrm{opt}}$:

$$g_{\mathrm{inv}}\left(\vec{\alpha}_\sigma|\vec{c}_{\{\sigma^*\}_{\mathrm{opt}},\mathrm{ss}}\right) < d\ , \forall\ \sigma \notin \{\sigma^*\}_{\mathrm{opt}}. \tag{S10}$$

850 Equivalently, Eq. (S10) can be expressed as "a set of species $\{\sigma^*\}_{\mathrm{opt}}$ construct a fitness
851 landscape which places themselves on the top", according to Eq. (S9).
852
853 The steady state constructed by $\{\sigma^*\}$ is influenced by the supply $\vec{c}_{\mathrm{supply}}$ and the dilution
854 rate d. For different chemostat parameters, the non-invasible species set $\{\sigma^*\}_{\mathrm{opt}}$ may
855 be different. In the following steps, we described a generally applicable protocol for
856 obtaining the non-invasible strategies:
857
858 1. Maximal growth rates and maximizing strategies: In a metabolic model with trade-offs
859 in resource allocation, the maximizing resource allocation strategy $\vec{\alpha}_{\mathrm{max}}$ under a given
860 environment $\vec{c}$ is defined as the strategy that maximizes invasion growth rate:

$$g_{\mathrm{max}}(\vec{c}) \coloneqq \max_{\vec{\alpha}}\left(g_{\mathrm{inv}}(\vec{\alpha}_\sigma|\vec{c})\right) \tag{S11}$$
$$\vec{\alpha}_{\mathrm{max}}(\vec{c}) \coloneqq \arg\max_{\vec{\alpha}}\left(g_{\mathrm{inv}}(\vec{\alpha}_\sigma|\vec{c})\right).$$

861
862 2. Maximal growth contour: For a given dilution rate d, all environments that support a
863 maximal growth rate of d constitute the "maximal growth contour":

$$GC_{\max} := \{\vec{c}_0 \mid g_{\max}(\vec{c}_0) = d\}. \tag{S12}$$

$GC_{\max}$ is generally formed by many species, with each species adopting the maximizing
strategies $\vec{\alpha}_{\max}(\vec{c}_0)$ corresponding to one environment $\vec{c}_0$ on the maximal growth
contour.
$GC_{\max}$ is outside of the invasion zone for any species σ. (Otherwise, if a species σ could
invade an environment $\vec{c}_0$ on $GC_{\max}$, $g_{\mathrm{inv}}(\vec{\alpha}_\sigma \mid \vec{c}_0 \in GC_{\max}) > d$, this would directly violate
the requirement by Eqs. (S11) and (S12) that $\max_{\vec{\alpha}}\big(g_{\mathrm{inv}}(\vec{\alpha}_\sigma \mid \vec{c}_0)\big) = d$.) Therefore, the
necessary and sufficient condition for a set of species to be evolutionarily stable, is to
construct a steady-state environment on the maximal growth contour:

$$\vec{c}_{\{\sigma^*\}_{\mathrm{opt,ss}}} \in GC_{\max}. \tag{S13}$$

872   Therefore, a strategy belonging to the non-invasible set must be a maximizing strategy.
873   An example of maximal growth contour is shown in Fig 4D.
874
3. Non-invasible strategy: Nevertheless, adopting one of the maximizing strategies
along the maximal growth contour does not guarantee that a species will satisfy Eq.
(S13) and become non-invasible, as a maximizing strategy for environment $\vec{c}_1$ may end
up constructing a different environment $\vec{c}_2$. To identify a non-invasible species for supply
condition $\vec{c}_{\mathrm{supply}}$, the flux-balance condition needs to be considered, with the strategies
maximized at each environment. This requirement forms a "maximal flux-balance curve"
in the nutrient space:

$$FB_{\max} := \left\{ \vec{c} \mid \frac{c_{i,\,\mathrm{supply}} - c_i}{I_i\big(\vec{c}, \vec{f}^{-1}(\vec{c}, \vec{\alpha}_{\max}(\vec{c})), \vec{\alpha}_{\max}(\vec{c})\big)} = \frac{m}{d \cdot r} \text{ AND } c_i < c_{i,\mathrm{supply}} \right\}. \tag{S14}$$

882   If the intersection of the maximal growth contour and the maximal flux-balance curve
883   exists, it is the evolutionarily stable environment under dilution rate d and supply
884   condition $\vec{c}_{\mathrm{supply}}$, $\vec{c}_{\mathrm{opt}}$. The maximizing strategy for this environment, $\vec{\alpha}_{\mathrm{opt}} = \vec{\alpha}_{\max}(\vec{c}_{\mathrm{opt}})$,
885   constructs the environment $\vec{c}_{\mathrm{opt}}$, and is evolutionarily stable.
886
887   4. Evolutionarily stable coexistence at the discontinuous points of the maximal growth
888   contour: Inversely, for each environment $\vec{c}_0$ on the maximal growth contour, all supply
889   conditions that enables the maximizing strategy of $\vec{c}_0$ to become the non-invasible
890   strategy can be calculated from the supply line according to Eq. (S6):

$$SL(\vec{c}_0) := \left\{ \vec{c}_{\mathrm{supply}} \mid \vec{c}_{\mathrm{supply}} = x \cdot \vec{I}\big(\vec{c}_0, \vec{f}^{-1}(\vec{c}_0, \vec{\alpha}_{\max}(\vec{c}_0)), \vec{\alpha}_{\max}(\vec{c}_0)\big) + \vec{c}_0 \right\}, \tag{S15}$$

for any non-zero value of x. Some examples are shown in Fig 4D.
When there are discontinuous points on the maximal growth contour, there can be
"gaps" in the nutrient supply space, where no single strategy on the maximal growth
contour satisfies Eq. (S14) . Under this condition, more than one strategy are required
to co-create an environment on the conjunctions of discontinuous points of the maximal
growth contour. Therefore, discontinuous points of the maximal growth contour permit
evolutionarily stable coexistence, where $\{\sigma^*\}_{\mathrm{opt}}$ contains more than one species. Two
examples of such discontinuities and coexistence are shown in Fig 5 and Fig 6.
899
Metabolic models

Different assumptions can be made regarding the metabolic models $\vec{f}(\vec{c}, \vec{x}, \vec{\alpha})$, $g(\vec{c}, \vec{x}, \vec{\alpha})$,
and $\vec{I}(\vec{c}, \vec{x}, \vec{\alpha})$, focusing on various aspects of cellular growth. Different assumptions lead
to distinct classes of metabolic models with various results. Nevertheless, our analysis
schemes, including the invasion geometry, fitness landscape, and evolutionary stable
strategies, are generally applicable for various metabolic models. In this work, we used
five metabolic models to illustrate multiple aspects of the species-environment
feedback:
908
1. Metabolic model with two essential nutrients
When two nutrients are both essential for growth, such as nitrogen and phosphorus,
and both require a substantial allocation of resources for import, the system can be
abstractly modeled as shown in Fig 4A. In this metabolic model, we assume an exact
trade-off between the allocation of limited resources to import nutrient *a* or nutrient *b*.
The fraction of resources allocated to import nutrient *a* is represented by α_a, thus
leaving a fraction $\alpha_b = 1 - \alpha_a$ to import nutrient *b*. The import rate of nutrient i is
assumed to follow the Monod equation as a function of nutrient concentration, and is
proportional to α_i:

$$I_i(\vec{c}) = \alpha_i \cdot \frac{c_i}{c_i + K_i} \text{ for } i = a, b. \tag{S16}$$

918 Import of both nutrients is required for cell growth:

$$g(\vec{c}) = \gamma \cdot \min(I_a(\vec{c}), I_b(\vec{c})). \tag{S17}$$

919 For this model, for simplicity we do not explicitly consider intracellular metabolites.
920 Rather, import directly determines growth.
921 In this model, a "species" is defined by its value of α_a.
922 Nutrient limitation can be clearly quantified in this system: if $I_a(\vec{c}) > I_b(\vec{c})$, the system is
923 limited by nutrient *b*; if $I_a(\vec{c}) < I_b(\vec{c})$, the system is limited by nutrient *a*.
924
925 A species with the following parameters was used to generate Fig 1D and E, focusing
926 on how supply conditions and dilution rate influence nutrient limitation:

K_a	K_b	γ	α_a
0.7	1.3	10	0.3

In Fig 1D, to demonstrate how species construct the same environment out of different
supply conditions, the chemostat dilution rate was set to $d = 1$, and three supply
conditions were used: $\vec{c}_{\text{supply}} = [0.6, 0.3546]$ (purple), $\vec{c}_{\text{supply}} = [0.8, 0.5273]$ (cyan), and
$\vec{c}_{\text{supply}} = [1, 0.7]$ (blue).
In Fig 1E, to demonstrate how dilution rates may switch the nutrient, we used the same
supply condition as the blue condition in Fig 1D ($\vec{c}_{\text{supply}} = [1, 0.7]$), and three dilution
rates: 0.5 (yellow), 1 (red), and 1.6 (deep red).
934
935 Species with following parameters were used to generate Fig 4B-D:

K_a	K_b	γ
0.5	0.5	10

The strategy α_a varies for different species. In Fig 4B, Species *Blue* has $\alpha_a = 0.35$,
species *Red* has $\alpha_a = 0.65$. In Fig 4C, we started with species *Blue* and species *Red*.

We then generated the fitness landscape for each species at the steady-state
environment it constructed, then chose the strategy that maximized invasion growth rate
for this fitness landscape to generate a new species, and iterated this process five
times. The species *Black* has $\alpha_a = 0.5$.
In generating Fig 4D, we followed the protocols described in section "Non-invasible /
evolutionarily stable strategies".
944
2. Metabolic model with substitutable nutrients
When two nutrients are mutually substitutable for growth, such as glucose and
galactose, the system can be described by metabolic model as shown in Fig 2A. The
trade-off and import functions are taken to be the same as in Model 1: *metabolic model*
with two essential nutrients. However, import of the two nutrients contributes additively
toward growth rate:

$$g(\vec{c}) = \gamma \cdot (I_a(\vec{c}) + I_b(\vec{c})). \tag{S18}$$

951   A species is defined by its value of α_a.
952   For this model, all growth contours intersect at one point. The growth contour of species
953   σ satisfies the equation: $\alpha_a \cdot \frac{c_a}{c_a + K_a} + (1 - \alpha_a) \cdot \frac{c_b}{c_b + K_b} = d/\gamma$. Regardless of the value of
954   α_a, the environment $[\frac{K_a}{\frac{\gamma}{d}-1}, \frac{K_b}{\frac{\gamma}{d}-1}]$ is always on the growth contour.
955   Species with the following parameters were used to generate Fig 2B-E:

K_a	K_b	γ
1.2	0.8	3

956   The strategy α_a varies for different species. In Fig 2B-D, Species *Blue* has $\alpha_a = 0.2$,
957   species *Red* has $\alpha_a = 0.6$. Supply conditions are different among the three figures: in
958   Fig 2B, $\vec{c}_{\text{supply}} = [0.5, 1]$; in Fig 2C, $\vec{c}_{\text{supply}} = [1, 0.5]$; in Fig 2D, $\vec{c}_{\text{supply}} = [1, 1]$.
959   All conditions in Fig 2E are the same as in Fig 2D, other than that five additional species
960   are added to the system. Their strategies are indicated by the legend at the right.
961
3. Metabolic model with substitutable nutrients that require assimilation
In cells, the assimilation of imported raw material, such sugars, into biomass such as
proteins, takes multiple steps and enzymes and consumes a considerable amount of
energy. When the resources allocated to nutrient assimilation are considered, a cell's
strategy becomes more complex. A mathematical model involving three substitutable
nutrients *a, b, c* that need assimilation is shown in Fig 3A, with α_{i1} represents the
fraction of resources allocated to importing nutrient *i* into internal metabolite, and α_{i2}
represents the fraction of resources allocated to assimilate the internal *i* into biomass. In
this model, the import rate has a similar form to the previous two models,

$$I_i(c_i) = V \cdot \alpha_{i1} \cdot \frac{c_i}{c_i + K_i}. \tag{S19}$$

971   The internal metabolite concentration $c_{i,\text{internal}}$ has an influx of $r \cdot I_i(c_i)$, meanwhile, it is
972   diluted by cell growth in the rate of g. We assume all nutrients are substitutable
973   therefore the internal pools contributes via summation to growth, it is converted into
974   biomass at a rate $k \cdot \alpha_{i2} \cdot c_{i,\text{internal}}$:

$$\frac{dc_{i,\text{internal}}}{dt} = I_i(c_i) - g(\vec{c}_{\text{internal}}) \cdot c_{i,\text{internal}} - k \cdot \alpha_{i2} \cdot c_{i,\text{internal}}. \tag{S20}$$

975 Therefore, the mass converted into biomass per unit time per unit volume is:
976 $\sum_i (k \cdot \alpha_{i2} \cdot c_{i,\text{internal}})$, and the growth rate defined as the relative gain of total biomass M
977 is:

$$g(\vec{c}_{\text{internal}}) = \frac{\frac{dM}{dt}}{M} = \frac{k}{r} \cdot \sum_i (\alpha_{i2} \cdot c_{i,\text{internal}}).$$ **(S21)**

978
979 In generating Fig 4B-C, the chemostat parameters were: $\vec{c}_{\text{supply}} = [1,1,1]$, and $d = 1$,
980 and the species parameters were:

V	K_i $(i = a, b, c)$	k
1000	0.5	11

981 The three species allocate their resources differently:

Strategies	α_{a1}	α_{a2}	α_{b1}	α_{b2}	α_{c1}	α_{b2}
Red	0.15	0.2	0.1	0.25	0.26	0.04
Green	0.26	0.04	0.15	0.2	0.1	0.25
Blue	0.1	0.25	0.26	0.04	0.15	0.2

982
983 To generate the Fig S1B, all other parameters are the same, other than $k = 10$.
984
985
4. Metabolic model with essential nutrients that can be interconverted
If two nutrients are both essential for growth, and a cell is able to convert one nutrient
into another albeit at a certain cost, as shown in Fig 5A, metabolic trade-offs involve the
following four elements of the allocation strategy $\vec{\alpha}$:
α_a: Fraction of resources allocated to import nutrient *a*.
α_b: Fraction of resources allocated to import nutrient *b*.
α_{ab}: Fraction of resources allocated to convert internal *b* into *a*.
α_{ba}: Fraction of resources allocated to convert internal *a* into *b*.
To implement trade-offs, the sum of elements of $\vec{\alpha} = (\alpha_a, \alpha_b, \alpha_{ab}, \alpha_{ba})$ is taken to be
equal to 1.
In this metabolic model, cells internalize nutrient *a* and nutrient *b* from the chemostat to
supply internal concentration of nutrients, $c_{a,\text{internal}}$ and $c_{b,\text{internal}}$. Meanwhile, the
internal nutrients can be converted into each other. Nutrients also diffuse in and out of
the cell passively with rate β. Cell growth requires both internal nutrients, and depletes
them in a fixed proportion.
In this model, the growth rate of a cell is taken to be:

$$g(\vec{c}_{\text{internal}}) = \frac{\gamma}{\frac{K_a}{c_{a,\text{internal}}} + \frac{K_b}{c_{b,\text{internal}}}}.$$ **(S22)**

1002 The net import rate, including passive diffusion, is:

$$I_i = (\alpha_i + \beta) \cdot c_i - \beta \cdot c_{i,\text{internal}}, \quad i = a, b.$$ **(S23)**

1003 Therefore, the dynamical equations for the internal nutrients are:

$$\frac{dc_{a,\text{internal}}}{dt} = I_a + \alpha_{ab} \cdot c_{b,\text{internal}} - \alpha_{ba} \cdot c_{a,\text{internal}} - g/K_a,$$ **(S24)**

$$\frac{dc_{b,\text{internal}}}{dt} = I_b + \alpha_{ba} \cdot c_{a,\text{internal}} - \alpha_{ab} \cdot c_{b,\text{internal}} - g/K_b. \tag{S25}$$

A species is defined by its value of $\vec{\alpha}$.

This metabolic model was used to demonstrate how to obtain locally optimal strategies and cartels, as shown in Fig 5. The parameter values used to generate the plots in Fig 5B-D were:

γ	$K_i \ (i = a, b)$	β
1	1	0.2

In generating Fig 5B, we searched for the maximizing strategies in the nutrient space, and classified them by their non-zero values. Maximal growth contours for four dilution rates: 0.1, 0.2, 0.3, 0.4 are shown from black to gray and white colors.
In generating Fig 5C, the chemostat parameters were set to $\vec{c}_{\text{supply}} = [0.5,1]$, and $d = 0.2$. The maximal growth contours for $d = 0.2$ were drawn, along with maximizing strategies along the contour shown as squares with colors corresponding to their sub-classes. At the discontinuous point of the maximal growth contour where the "converter" and the "importer" converge, the distinct two maximizing strategies are denoted species *Red* and species *Blue*. In generating the competition dynamics in inset, additional to the species *Red* and species *Blue*, ten other maximizing strategies along the maximal growth contours were chosen.

5. Metabolic model with multiple energy generating steps
Cell growth is also tightly coupled with energy production. For example, with a single carbon supply as the energy source, cells employ multi-step reactions to generate multiple ATP molecules. Each step requires dedicated enzymes. The reaction intermediates, such as acetate, usually have dual roles: on the one hand, they positively contribute to ATP production via downstream reactions; on the other hand, they negatively contribute to ATP production by hampering upstream reactions. To deal with the negative effects of intermediates, cells may transport them out into the environment, generally with some metabolic cost for transporters. On the other hand, cells can also uptake such intermediates and use them as an energy source.
We abstract such a process by the model shown in Fig 6A. A single chemical energy source *S* is supplied into the chemostat, which can be converted into intermediate *I* by cells. Four reactions are possible in this model, each mediate by a specific enzyme:
1. Import the resource *S* into the cell and convert it into internal intermediate I_{int} to extract energy (e.g. ATP). The fraction of the model enzyme budget allocated to this reaction is α_{ATP1}. We assume the reaction is reversible, with the concentration *S* contributing positively to the reaction rate while the concentration I_{int} contributes negatively:

$$J_1 = \alpha_{\text{ATP1}} \cdot V_1 \cdot \frac{[S] - \frac{[I_{\text{int}}]}{K_3}}{K_1 + [S] + \frac{[I_{\text{int}}]}{K_5}}. \tag{S26}$$

2. Process I_{int} via a downstream reaction to obtain more energy. The fraction of enzymes being allocated to this reaction is α_{ATP2}. For this model system, it does not

1042 qualitatively influence the final results whether this reaction is product inhibited. For
1043 simplicity, we assume this reaction has Michaelis–Menten form:

$$J_2 = \alpha_{\text{ATP2}} \cdot V_2 \cdot \frac{[I_{\text{int}}]}{K_2 + [I_{\text{int}}]}. \tag{S27}$$

1044 3. Export the internal intermediate out into the environment by diffusion, with a fraction
1045 of proteins α_{exp} allocated to channels that allow the excretion of the intermediate into
1046 the environment to become external intermediate I_{ext}.

$$J_3 = \alpha_{\text{exp}} \cdot k \cdot ([I_{\text{int}}] - [I_{\text{ext}}]). \tag{S28}$$

1047 4. Import the external intermediate into cells, with a fraction of proteins α_{imp} allocated to
1048 the import process. In reflect the property of the internal intermediate in inhibiting this
1049 transport reaction, the rate for this process is also product-inhibited:

$$J_4 = \alpha_{\text{imp}} \cdot V_4 \cdot \frac{[I_{\text{ext}}] - \frac{[I_{\text{int}}]}{K_6}}{K_4 + [I_{\text{ext}}] + \frac{[I_{\text{int}}]}{K_7}}. \tag{S29}$$

1050 Under this model, the rate of change for the energy source concentration in the
1051 chemostat is:

$$\frac{d[S]}{dt} = d \cdot ([S_{\text{supply}}] - [S]) - m/r \cdot J_1 \tag{S30}$$

1052 The rate of change of the external intermediate concentration in the chemostat is:

$$\frac{d[I_{\text{ext}}]}{dt} = d \cdot (-[I_{\text{ext}}]) - m/r \cdot (J_4 - J_3). \tag{S31}$$

1053
1054 The concentration for the intercellular metabolite I_{int} follows the equation:

$$\frac{d[I_{\text{int}}]}{dt} = J_1 - J_2 - J_3 + J_4. \tag{S32}$$

1055
1056 The growth rate is a weighted sum of the ATP produced by J_1 and J_2:

$$g = n_{\text{ATP1}} \cdot J_1 + n_{\text{ATP2}} \cdot J_2. \tag{S33}$$

1057
1058 In generating plots in Fig 6B-F, the species parameters were:

V_1	V_2	k	V_4	K_1	K_2	K_3	K_4	K_5	K_6	K_7	n_{ATP1}	n_{ATP2}
5	1	8	10	0.5	0.5	0.5	0.1	0.5	15	10	1	1

1059
1060 Maximal growth contours for dilution rates 0.2, 0.4, and 0.6 are shown in Fig 6B.
1061 For Fig 6C-D, the chemostat parameters are: $S_{\text{supply}} = 1, d = 0.4$.
1062 For Fig 6E-F, the chemostat parameters are: $S_{\text{supply}} = 1.8, d = 0.6$.
1063
Dynamic equations for multiple species in a chain of chemostats
Real ecosystems seldom exist in isolation. We modeled interconnected ecosystems via
a chain of chemostats labeled $k = 1$ to k_{tot} (Fig S2A). Each chemostat exchanges
medium and cells at leakage rate l with its two neighboring chemostats (if $k = 1$ or $k = $
k_{tot}, there is only one neighbor). The chemostat parameters $\vec{c}_{\text{supply}}$ and d are taken to
be identical for all chemostats.

1070

1071 For the k-th chemostat, the dynamical equations for the biomass density of species σ
1072 and the concentration of the i-th nutrient are:

$$\frac{dm_{\sigma,k}}{dt} = m_{\sigma,k} \cdot (g_\sigma(\vec{c}_k) - d) + l \cdot (m_{\sigma,k-1} + m_{\sigma,k+1} - 2 \cdot m_{\sigma,k}), \quad \textbf{(S34)}$$

$$\frac{dc_{i,k}}{dt} = d \cdot (c_{i,\text{ supply}} - c_{i,k}) - \sum_{\sigma=1}^{n} m_{\sigma,k} \cdot I_{i,\sigma}(\vec{c}_k) + l \cdot (c_{i,k+1} + c_{i,k-1} - 2 \cdot c_{i,k}). \quad \textbf{(S35)}$$

A steady-state solution to these equations is shown in Fig S2, using the same growth
and import models and parameters as in Fig 4, with the leakage rate set to be $l = 1$.

ACKNOWLEDGEMENTS

We thank Simon Levin for insightful discussions. Zhiyuan Li was supported by Princeton
Center for Theoretical Science and Center for the Physics of Biological Function. This
work was supported in part by NSF grant PHY-1607612, the National Science
Foundation Physics Frontier Center grant through the Center for the Physics of
Biological Function (PHY-1734030), and National Institutes of Health Grant (R01
GM082938).
1082

1083   **COMPETING INTERESTS**

1084   The authors declare that they have no conflict of interest.
1085
1086
1087

REFERENCES

Ackland, G., & Gallagher, I. (2004). Stabilization of large generalized Lotka-Volterra foodwebs by evolutionary feedback. *Physical Review Letters, 93*(15), 158701.

Amarasekare, P. (2003). Competitive coexistence in spatially structured environments: a synthesis. *Ecology Letters, 6*(12), 1109-1122.

Armstrong, R. A., & McGehee, R. (1980). Competitive exclusion. *The American Naturalist, 115*(2), 151-170.

Bairey, E., Kelsic, E. D., & Kishony, R. (2016). High-order species interactions shape ecosystem diversity. *Nat Commun, 7*, 12285. doi:10.1038/ncomms12285

Beardmore, R. E., Gudelj, I., Lipson, D. A., & Hurst, L. D. (2011). Metabolic trade-offs and the maintenance of the fittest and the flattest. *Nature, 472*(7343), 342-346. doi:10.1038/nature09905

Behar, H., Brenner, N., & Louzoun, Y. (2014). Coexistence of productive and non-productive populations by fluctuation-driven spatio-temporal patterns. *Theoretical Population Biology, 96*, 20-29.

Boer, V. M., Crutchfield, C. A., Bradley, P. H., Botstein, D., & Rabinowitz, J. D. (2010). Growth-limiting intracellular metabolites in yeast growing under diverse nutrient limitations. *Mol Biol Cell, 21*(1), 198-211. doi:10.1091/mbc.E09-07-0597

Callahan, B. J., Fukami, T., & Fisher, D. S. (2014). Rapid evolution of adaptive niche construction in experimental microbial populations. *Evolution, 68*(11), 3307-3316. doi:10.1111/evo.12512

Ceballos, G., Ehrlich, P. R., Barnosky, A. D., Garcia, A., Pringle, R. M., & Palmer, T. M. (2015). Accelerated modern human-induced species losses: Entering the sixth mass extinction. *Sci Adv, 1*(5), e1400253. doi:10.1126/sciadv.1400253

Cermeño, P., Teixeira, I. G., Branco, M., Figueiras, F. G., & Marañón, E. (2014). Sampling the limits of species richness in marine phytoplankton communities. *Journal of Plankton Research, 36*(4), 1135-1139.

Chase, J. M., & Leibold, M. A. (2003). *Ecological niches: linking classical and contemporary approaches*: University of Chicago Press.

D'Souza, G., Shitut, S., Preussger, D., Yousif, G., Waschina, S., & Kost, C. (2018). Ecology and evolution of metabolic cross-feeding interactions in bacteria. *Nat Prod Rep, 35*(5), 455-488. doi:10.1039/c8np00009c

De Leenheer, P., Levin, S. A., Sontag, E. D., & Klausmeier, C. A. (2006). Global stability in a chemostat with multiple nutrients. *Journal of Mathematical Biology, 52*(4), 419-438. doi:10.1007/s00285-005-0344-4

de Visser, J., Elena, S. F., Fragata, I., & Matuszewski, S. (2018). The utility of fitness landscapes and big data for predicting evolution. *Heredity (Edinb), 121*(5), 401-405. doi:10.1038/s41437-018-0128-4

Descamps-Julien, B., & Gonzalez, A. (2005). Stable coexistence in a fluctuating environment: an experimental demonstration. *Ecology, 86*(10), 2815-2824.

Escalante-Chong, R., Savir, Y., Carroll, S. M., Ingraham, J. B., Wang, J., Marx, C. J., & Springer, M. (2015). Galactose metabolic genes in yeast respond to a ratio of galactose and glucose. *Proc Natl Acad Sci U S A, 112*(5), 1636-1641. doi:10.1073/pnas.1418058112

Freilich, S., Zarecki, R., Eilam, O., Segal, E. S., Henry, C. S., Kupiec, M., . . . Ruppin, E. (2011). Competitive and cooperative metabolic interactions in bacterial communities. *Nat Commun, 2*, 589. doi:10.1038/ncomms1597

Friedman, J., Higgins, L. M., & Gore, J. (2017). Community structure follows simple assembly rules in microbial microcosms. *Nature ecology & evolution, 1*(5), 0109.

Geyrhofer, L., & Brenner, N. (2018). Coexistence and cooperation in structured habitats. *BioRxiv*, 429605.

Goldford, J. E., Lu, N., Bajić, D., Estrela, S., Tikhonov, M., Sanchez-Gorostiaga, A., . . . Sanchez, A. (2018). Emergent simplicity in microbial community assembly. *Science, 361*(6401), 469-474.

Goyal, A., Dubinkina, V., & Maslov, S. (2018). Multiple stable states in microbial communities explained by the stable marriage problem. *The ISME Journal*. doi:10.1038/s41396-018-0222-x

Goyal, S., Yuan, J., Chen, T., Rabinowitz, J. D., & Wingreen, N. S. (2010). Achieving optimal growth through product feedback inhibition in metabolism. *PLoS computational biology, 6*(6), e1000802.

Hardin, G. (1960). The competitive exclusion principle. *Science, 131*(3409), 1292-1297.

Huisman, J., Johansson, A. M., Folmer, E. O., & Weissing, F. J. (2001). Towards a solution of the plankton paradox: the importance of physiology and life history. *Ecology Letters, 4*(5), 408-411.

Huisman, J., van Oostveen, P., & Weissing, F. J. (1999). Species dynamics in phytoplankton blooms: incomplete mixing and competition for light. *The American Naturalist, 154*(1), 46-68.

Huisman, J., & Weissing, F. J. (1999). Biodiversity of plankton by species oscillations and chaos. *Nature, 402*(6760), 407-410. doi:10.1038/46540

Huisman, J., & Weissing, F. J. (2001). Biological conditions for oscillations and chaos generated by multispecies competition. *Ecology, 82*(10), 2682-2695.

Hutchinson, G. E. (1961). The paradox of the plankton. *The American Naturalist, 95*(882), 137-145.

Kasting, J. F., & Siefert, J. L. (2002). Life and the evolution of Earth's atmosphere. *Science, 296*(5570), 1066-1068. doi:10.1126/science.1071184

Kerr, B., Riley, M. A., Feldman, M. W., & Bohannan, B. J. M. (2002). Local dispersal promotes biodiversity in a real-life game of rock–paper–scissors. *Nature, 418*(6894), 171.

Koffel, T., Daufresne, T., Massol, F., & Klausmeier, C. A. (2016). Geometrical envelopes: Extending graphical contemporary niche theory to communities and eco-evolutionary dynamics. *Journal of Theoretical Biology, 407*, 271-289.

Laland, K., Matthews, B., & Feldman, M. W. (2016). An introduction to niche construction theory. *Evol Ecol, 30*, 191-202. doi:10.1007/s10682-016-9821-z

Leibold, M. A. (1995). The Niche Concept Revisited: Mechanistic Models and Community Context. *Ecology, 76*(5), 1371-1382. doi:10.2307/1938141

Letten, A. D., Ke, P. J., & Fukami, T. (2017). Linking modern coexistence theory and contemporary niche theory. *Ecological Monographs, 87*(2), 161-177. doi:10.1002/ecm.1242

Levin, S. A. (1970). Community equilibria and stability, and an extension of the competitive exclusion principle. *The American Naturalist, 104*(939), 413-423.

Li, S. H.-J., Li, Z., Park, J. O., King, C. G., Rabinowitz, J. D., Wingreen, N. S., & Gitai, Z. (2018). Escherichia coli translation strategies differ across carbon, nitrogen and phosphorus limitation conditions. *Nature microbiology, 3*(8), 939.

Liebermeister, W., Noor, E., Flamholz, A., Davidi, D., Bernhardt, J., & Milo, R. (2014). Visual account of protein investment in cellular functions. *Proceedings of the National Academy of Sciences, 111*(23), 8488-8493.

Luli, G. W., & Strohl, W. R. (1990). Comparison of growth, acetate production, and acetate inhibition of Escherichia coli strains in batch and fed-batch fermentations. *Applied and environmental microbiology, 56*(4), 1004-1011.

MacArthur, R. (1970). Species packing and competitive equilibrium for many species. *Theoretical Population Biology, 1*(1), 1-11.

Maharjan, R., Seeto, S., Notley-McRobb, L., & Ferenci, T. (2006). Clonal adaptive radiation in a constant environment. *Science, 313*(5786), 514-517. doi:10.1126/science.1129865

Marsland III, R., Cui, W., & Mehta, P. (2019). The Minimum Environmental Perturbation Principle: A New Perspective on Niche Theory. *arXiv preprint arXiv:1901.09673*.

McGehee, R., & Armstrong, R. A. (1977). Some mathematical problems concerning the ecological principle of competitive exclusion. *Journal of Differential Equations, 23*(1), 30-52.

Moore, C., Mills, M., Arrigo, K., Berman-Frank, I., Bopp, L., Boyd, P., . . . Jaccard, S. (2013). Processes and patterns of oceanic nutrient limitation. *Nature Geoscience, 6*(9), 701.

Murdoch, W. W., & Oaten, A. (1975). Predation and population stability.

Mustonen, V., & Lässig, M. (2009). From fitness landscapes to seascapes: non-equilibrium dynamics of selection and adaptation. *Trends in Genetics, 25*(3), 111-119. doi:10.1016/j.tig.2009.01.002

Odling-Smee, F. J., Laland, K. N., & Feldman, M. W. (2003). *Niche construction: the neglected process in evolution*: Princeton university press.

Odum, E. P., & Barrett, G. W. (1971). *Fundamentals of ecology* (Vol. 3): Saunders Philadelphia.

Palmer, M. W. (1994). Variation in species richness: Towards a unification of hypotheses. *Folia Geobotanica et Phytotaxonomica, 29*(4), 511-530. doi:10.1007/BF02883148

Pfeiffer, T., & Bonhoeffer, S. (2004). Evolution of cross-feeding in microbial populations. *The American Naturalist, 163*(6), E126-E135.

Posfai, A., Taillefumier, T., & Wingreen, N. S. (2017). Metabolic Trade-Offs Promote Diversity in a Model Ecosystem. *Physical Review Letters, 118*(2). doi:10.1103/PhysRevLett.118.028103

Rosenzweig, R. F., Sharp, R., Treves, D. S., & Adams, J. (1994). Microbial evolution in a simple unstructured environment: genetic differentiation in Escherichia coli. *Genetics, 137*(4), 903-917.

Roy, S., & Chattopadhyay, J. (2007). Towards a resolution of 'the paradox of the plankton': A brief overview of the proposed mechanisms. *Ecological complexity, 4*(1-2), 26-33.

Scheffer, M., Rinaldi, S., Huisman, J., & Weissing, F. J. (2003). Why plankton communities have no equilibrium: solutions to the paradox. *Hydrobiologia, 491*(1-3), 9-18.

Schirrmeister, B. E., de Vos, J. M., Antonelli, A., & Bagheri, H. C. (2013). Evolution of multicellularity coincided with increased diversification of cyanobacteria and the Great Oxidation Event. *Proc Natl Acad Sci U S A, 110*(5), 1791-1796. doi:10.1073/pnas.1209927110

Smith, H. L., & Waltman, P. (1995). *The theory of the chemostat: dynamics of microbial*
competition (Vol. 13): Cambridge university press.
Soliveres, S., Maestre, F. T., Ulrich, W., Manning, P., Boch, S., Bowker, M. A., . . . Allan, E.
(2015). Intransitive competition is widespread in plant communities and maintains their
species richness. *Ecology Letters, 18*(8), 790-798. doi:10.1111/ele.12456
Taillefumier, T., Posfai, A., Meir, Y., & Wingreen, N. S. (2017). Microbial consortia at steady
supply. *eLife, 6*, e22644.
Tilman, D. (1980). Resources: a graphical-mechanistic approach to competition and predation.
The American Naturalist, 116(3), 362-393.
Tilman, D. (1982). *Resource competition and community structure*: Princeton university press.
Tilman, D. (1994). Competition and Biodiversity in Spatially Structured Habitats. *Ecology,*
75(1), 2-16. doi:Doi 10.2307/1939377
Van den Bergh, B., Swings, T., Fauvart, M., & Michiels, J. (2018). Experimental design,
population dynamics, and diversity in microbial experimental evolution. *Microbiol. Mol.*
Biol. Rev., 82(3), e00008-00018.
Venturelli, O. S., Carr, A. C., Fisher, G., Hsu, R. H., Lau, R., Bowen, B. P., . . . Arkin, A. P.
(2018). Deciphering microbial interactions in synthetic human gut microbiome
communities. *Mol Syst Biol, 14*(6), e8157. doi:10.15252/msb.20178157
Wang, X., & Tang, C. (2017). Optimal growth of microbes on mixed carbon sources. *arXiv*
preprint arXiv:1703.08791.
Wides, A., & Milo, R. (2018). Understanding the dynamics and optimizing the performance of
chemostat selection experiments. *arXiv preprint arXiv:1806.00272*.
Wintermute, E. H., & Silver, P. A. (2010). Emergent cooperation in microbial metabolism. *Mol*
Syst Biol, 6, 407. doi:10.1038/msb.2010.66
Wright, S. (1932). *The roles of mutation, inbreeding, crossbreeding, and selection in evolution*
(Vol. 1): na.
Zhang, X. S. (2012). Fisher's geometrical model of fitness landscape and variance in fitness
within a changing environment. *Evolution: International Journal of Organic Evolution,*
66(8), 2350-2368.
Ziv, N., Brandt, N. J., & Gresham, D. (2013). The Use of Chemostats in Microbial Systems
Biology. *Jove-Journal of Visualized Experiments*(80). doi:UNSP e50168
10.3791/50168
1254
1255

    FIGURES

Figure 1

1257
1258
1259   **Figure 1 - Chemostat behavior can be represented in the space of nutrient**
1260   **concentrations.**
1261
1262   A. Schematic diagram of a chemostat occupied by a single microbial species. In the
1263   well-mixed medium (pale blue) of a chemostat, cells (orange ellipses) consume
1264   nutrients and grow. An influx of nutrients with fixed concentrations (blue and green
1265   arrows) is supplied at the same rate as dilution, keeping the medium volume constant.
1266
B. Nutrient supply shifts the relationship between RNA/Protein ratio and growth rate of
E. coli cultured in chemostats from phosphorus limitation (P-limited, green open circles
and dotted line) to nitrogen limitation (N-limited, blue open circles and dotted line). Data
for dilution of supplied nitrogen by 2, 5, and 10-fold starting from the P-limited condition
are shown as solid dots and corresponding best-fit lines. Each measurement was
repeated three times and standard errors are shown by bars.
1273

1274   C. Same as (B), but for phosphorus and carbon limitation instead of phosphorus and
1275   nitrogen limitation. Data for dilution of supplied carbon by 2, 5, and 10-fold starting from
1276   the P-limited condition are shown as solid dots and corresponding best-fit lines.
1277
D. Visual representation of how a species creates its own chemostat environment.
Background color indicates the growth rate of cells as a function of nutrient
concentrations c_a and c_b, with the growth contour shown by the red curve. The supply
line for the steady-state environment (purple dot) is shown as a dotted black line.
Different supply concentrations ($c_{a,supply}$ and $c_{b,supply}$) along the supply line are marked by
purple, cyan, and blue circles, with the corresponding flux-balance curves shown in the
same colors.
1285
1286   E. Dilution rate can flip nutrient limitation. The external supply condition is marked by a
1287   blue circle, and the flux-balance curve for this supply is shown in the same color. Three
1288   growth contours with increasing dilution rates are shown from yellow to deep red, and
1289   the corresponding steady-state environments are shown in colored dots.
1290
1291
1292

Figure 2

1293
1294 **Figure 2 – Invasion and coexistence in a chemostat.**
1295
1296 A. Example of a metabolic model with a trade-off in allocation of resources for import of
1297 two substitutable nutrients, with both nutrients contributing additively to growth. Species
1298 *Red* and species *Blue* allocate resources differently (indicated by parameter α_a, see
1299 Methods).
1300
B. Example of successful invasion of species *Blue* by species *Red*. A small amount of
species *Red* is introduced to a steady-state chemostat of species *Blue*. Growth contours
and steady-state environments of species *Blue* and species *Red* are shown as curves
and dots in the corresponding colors (colored background indicates the "invasion zone"
of *Red*, and represents the growth rate of *Red* in this zone). The supply condition is
marked by black circle. Black curves with arrows show the time trajectory of the
invasion in nutrient space. Inset: biomass of species in chemostat over time course of
invasion.
1309

1310   C. Same as (B), except that because the supply condition (black circle) is different, the
1311   attempted invasion by species *Red* is unsuccessful.
1312
D. Growth contours and the steady-state environments created by *Red* or *Blue* alone,
under the supply condition shown by the black circle. Black curve with arrows shows a
trajectory in nutrient space. Purple dot indicates the steady-state environment created
by *Red* and *Blue* together. Lower inset: time course of species biomass. Upper inset:
the fitness landscape created by *Red* alone (color red, with the strategy of *Red* marked
by red diamond), created by *Blue* alone (color blue, with the strategy of *Blue* marked by
blue diamond), and created by both species (color purple). The colors correspond to the
steady-state environments shown by colored dots in (D).
1321
1322   E. Growth contours and the species-specific steady-state environments for seven
1323   different species alone, under the supply condition shown by the black circle. Black
1324   curve with arrows shows a trajectory in nutrient space. Lower inset: time course of
1325   species biomass.
1326

Figure 3

A Import   Conversion

α_{a1} α_{a2}

Growth

α_{b1} α_{b2}

$$g = \frac{k}{r}\sum_i \alpha_i \cdot c_{i,internal}$$

α_{c1} α_{c2}

B

c_c

Species

— 1
— 2
— 3

c_b c_a

C

Biomass

Species

Fitness

Time

Figure 3 - Rock-paper-scissors oscillations.

A. Example of a metabolic model with a trade-off in allocation of resources for importing of three substitutable nutrients and assimilating them into biomass, with all three nutrients contributing additively to growth. Species *Red*, species *Blue*, and species *Green* allocate resources differently (see Methods).

B. Growth contours (surfaces), flux-balance curves (lines), and steady-state nutrient concentrations (dots) for the three species in a three-dimensional nutrient space. Black curves with arrows show the system's limit-cycle trajectory.

C. The top panel shows the time course of species biomass for the limit cycle in (B). The bottom panel shows how the fitness landscape changes with time over one period of the oscillation.

Figure 4

Figure 4 – Multi-stability, chain of invasion, and non-invasible strategy.

A. Example of bistability for a metabolic model with a trade-off in allocation of resources for import of two essential nutrients, with the lower of the two import rates determining growth rate. Species *Red* and species *Blue* allocate resources differently (indicated by parameter α_a, see Methods).

B. Bistability of the system in (A) shown in nutrient space. Black curves with arrows show the trajectories of simulations with different initial conditions. Inset: the fitness landscape created by species *Red* or *Blue* alone, with colors corresponding to the steady-state environments shown by colored dots in (B).

C. A chain of invasion. Fitness landscape created by species with different resource allocation strategies (marked by diamond shapes). Starting from species *Blue*, the species having the highest growth rate in the fitness landscape created by the "former" species is chosen. This creates a chain of invasion from *Blue* to *Light Green*, *Yellow*, *Deep Green*, *Deep Purple*, all the way (intermediate processes omitted) to the species *Black*, which places itself on the peak of its fitness landscape. The same procedure is also performed starting with species *Red*.

D. Depiction of non-invasible strategies under different supply conditions. Black-white background indicates the maximal growth rate of model in (A) under each environment, and the contour of maximal growth rates contains different strategies (represented by red-to-blue color). Growth contours of three species adopting one of the "maximizing strategies" are colored by their strategies. The supply conditions allowing these strategies to be "non-invasible" are marked by dashed black lines.

Figure 5

1374
1375
1376 **Figure 5 - Non-invasible cartels.**
1377
1378 A. Metabolic model with a trade-off in allocation of resources for import of two nutrients
1379 plus their interconversion, with both nutrients necessary for growth.
1380
1381 B. Three subclasses of maximizing metabolic strategies in nutrient space are indicated
1382 by background color, and circles with arrows illustrate the metabolic strategies of each
1383 subclass. The maximal growth contours for four growth rates (0.1, 0.2, 0.3, 0.4) are
1384 marked by gray colors.
1385
C. Two maximizing strategies co-creating a non-invasible steady state. At dilution rate
0.2, the maximal growth contour and the corresponding maximizing strategies are
shown as colored squares. At a discontinuous point of the growth contour, the supply

lines of two distinct metabolic strategies (*Red* and *Blue*) span a gray region, where any
supply condition (e.g. black circle) requires the two maximizing strategies to co-create
the environment on the discontinuous point. Red and blue dots mark the environments
created by species *Red* and species *Blue* alone, and the purple dot marks the
environment co-created by *Red* and *Blue*. Black curve with arrows shows a trajectory in
nutrient space. Inset is the competition dynamics of the species *Red* and species *Blue*
together with 10 other maximizing species with different strategies.
1396
1397   D. The fitness landscapes for the three environments in (C) indicated by corresponding
1398   box colors. For class Green and Red, the strategy is represented by α_a, for class Blue,
1399   the strategy is represented by α_b.
1400

Figure 6

1402 **Figure 6 – Species creating new nutrient dimensions and achieving evolutionarily**
1403 **stable coexistence.**
1404
1405 A. Metabolic model with a single supplied nutrient S. Cells allocate enzymes to convert
1406 S into internal intermediate I_{int} and produce energy (denoted as "ATP"), export internal
1407 intermediate into the chemostat to become I_{ext}, import external intermediate, or
1408 consume I_{int} to produce ATP. The growth rate is the sum of ATP production (see
1409 Methods).
1410
1411 B. Three subclasses of maximizing metabolic strategies in nutrient space are indicated
1412 by background color, and circles with arrows illustrate the metabolic strategies of each
1413 subclass. The maximal growth contours for three growth rates (0.2, 0.4, 0.6) are marked
1414 by black-to-white colors.
1415
C. At dilution rate 0.4, two maximizing strategies co-create a non-invasible environment.
The maximal growth contour and the corresponding maximizing strategies are shown as
colored squares. At a discontinuous point of the growth contour, the supply lines of two
distinct metabolic strategies (*Green* and *Blue*) span a gray region, where any supply
condition (e.g. black circle) requires two maximizing strategies to co-create the
environment at the discontinuous point. Blue dot marks the environment created by
species *Blue* alone, and the cyan dot marks the environment co-created by *Blue* and
Green. Black curve with arrows shows a trajectory in nutrient space. Inset shows the
time course of species biomass, with species *Green* added to the chemostat at time
100.
1426
1427 D. The fitness landscapes for two environments in (C) indicated by corresponding box
1428 colors, reflecting the relationship between instantaneous growth rate and resource
1429 allocation strategy. For class Blue and Red, the strategy is represented by α_{ATP1}; for
1430 class Green the strategy is represented by α_{imp}.
1431
1432 E. Same as (C), except that the dilution rate is 0.6, and the inset shows the time course
1433 of species biomass, starting with *Blue* and *Green*, with species *Red* added to the
1434 chemostat at time 100.
1435
1436 D. Same as (D), except that it is for the two steady-state environments shown in (E).
1437
1438

1439 **SUPPLEMENTAL FIGURES**
1440

Figure S1

A

B

C

Figure S1- Rock-paper-scissor fitness landscape and heteroclinic cycle

A. The fitness of Species 1, 2, and 3 in the steady-state environment constructed by species 1, 2, and 3 for the model in Fig 3.

B. Growth contours (surfaces), flux-balance curves (lines), and steady-state nutrient concentrations (dots) for three species in a three-dimensional nutrient space, with a different conversion speed ($k = 10$) than in Fig. 3 ($k = 1$) (see Methods). Black curves with arrows show the system's oscillatory trajectory.

C. The top panel shows the time course of species biomass for the limit cycle in (B). The bottom panel shows how the fitness landscape changes with time.

D. Time course of species biomass over a long duration.

Figure S2

A

B

C

D

1457 **Figure S2- Steady-state spatial heterogeneity for linked chemostats.**
1458
1459 With initial seeding of two species, one at each of the two ends of a chain of
1460 chemostats, a steady-state gradient of species biomass density spontaneously emerges
1461 accompanied by a gradient of nutrient concentrations, even though the supply
1462 conditions and dilution rates are identical for all the chemostats.
1463
1464 A. Schematic of k_{tot} linked chemostats exchanging medium and cells via leakage,
1465 described by Eqs. S34-S35. The two species in the chemostats (*Blue* and *Red*) are the
1466 same bistable pair as in Fig 4B and the leakage rate is $l = 1$.
1467
1468 B. The species composition along 20 linked chemostats for the system in (A). Species
1469 colors correspond to those in Fig 4B, with species *Blue* having $\alpha_a = 0.35$, species *Red*
1470 having $\alpha_a = 0.65$. The dashed black curve shows the sum of the two biomass densities.
1471 The initial condition was cell-free chemostats with a small amount of *Blue* added to
1472 Chemostat 1 and small amount of *Red* added to Chemostat 20.
1473
1474 C. Concentrations along the 20 chemostats for nutrient *a* (green) and nutrient *b* (cyan)
1475 for system in (A).
1476
1477 D. The fitness landscape along the chain of chemostats. The *x*-axis is the 20 linked
1478 chemostats, and the *y*-axis is the metabolic strategy represented by α_a. Color indicates
1479 the growth rate of species adopting the given strategy in the *k*-th chemostat.
1480
1481

Figure S3

Figure S3 – Maximizing strategies in nutrient space.

A. For each environment in the nutrient space, the maximizing resource allocation strategies that maximizes growth rates for the model in Fig 5A. Each strategy is represented by the four elements $[\alpha_a, \alpha_b, \alpha_{ab}, \alpha_{ba}]$, and values for each element are shown by a heatmap. Black-to-white curves are the maximal growth contours for $d = 0.1, 0.2, 0.3, 0.4$.

B. For each environment in the nutrient space, the maximizing resource allocation strategies that maximizes growth rates for the model in Fig 6A. Each strategy is represented by the four elements $[\alpha_{\mathrm{ATP1}}, \alpha_{\mathrm{ATP2}}, \alpha_{\mathrm{exp}}, \alpha_{\mathrm{imp}}]$, and values for each element are shown by a heatmap. Black-to-white curves are the maximal growth contours for $d = 0.2, 0.4, 0.6$.

C. Schematic representations of the three classes of maximizing strategies appearing in (B).