Diversity of fall armyworm, *Spodoptera frugiperda* and their gut bacterial community in Kenya

Joseph Gichuhi[1,2], Sevgan Subramanian[1], Fathiya M. Khamis[1], Johnnie Van den Berg[2], Hannalene du
Plessis[2], Sunday Ekesi[1], and Jeremy K Herren[1,3]

[1] International Centre of Insect Physiology and Ecology (ICIPE), Kasarani, Nairobi, Kenya

[2] Unit for Environmental Sciences and Management, North-West University, Potchefstroom, South Africa,

[3] MRC-University of Glasgow Centre for Virus Research, Henry Wellcome Building, Glasgow, UK

For correspondence: jherren@icipe.org Tel: +254716660160

Abstract

The invasive fall armyworm (FAW), *Spodoptera frugiperda* (J.E. Smith) is a polyphagous pest that causes
widespread damage particularly to maize and sorghum in Africa. The microbiome associated with *S.
frugiperda* could play a role in their success and adaptability. However, these bacterial communities remain
poorly studied, especially for *S. frugiperda* in Africa. We investigated the composition, abundance and
diversity of microbiomes associated with larval and adult specimens of *S. frugiperda* collected from four
maize growing regions in Kenya through high throughput sequencing of bacterial 16S rRNA gene. We
identified Proteobacteria and Firmicutes as the most dominant phyla and lesser proportions of
Bacteroidetes and Actinobacteria. We also observed differences in bacterial microbiome diversity between
larvae and adults that are a likely indication that some prominent larval bacterial groups are lost during
metamorphosis. Several bacterial groups were found in both adults and larvae suggesting that they are
transmitted across developmental stages. Reads corresponding to several known entomopathogenic
bacterial clades as well as the non-bacterial entomopathogen, *Metarhizium rileyi* (Farl.) Kepler, Rehner &
Humber (2014)*,* were observed. Mitochondrial DNA haplotyping of the *S. frugiperda* population in Kenya
indicated the presence of both 'Rice' and 'Corn' strains, with a higher prevalence of the 'Rice' strain. Insights
into the microbiota may ultimately provide alternative avenues for controlling of this pest.

Keywords: Fall armyworm, *Spodoptera frugiperda*, gut bacteria, 16S sequencing, corn strain, rice strain,
mtDNA haplotype

Introduction

Invasions by exotic pests can have major detrimental effects on agricultural production and natural
resources (Huber et al. 2002). The fall armyworm, *Spodoptera frugiperda* (J. E. Smith) (Lepidoptera:
Noctuidae) is a polyphagous pest that is native to tropical regions of the western hemisphere, where it is
known for its ability to cause economic damage to several crop species. In 2016, it was first detected in
West Africa (Goergen et al. 2016), and since then this pest has rapidly spread across the continent
(Goergen et al. 2016; Tindo et al. 2016; Day et al. 2017; Nagoshi et al. 2017, 2018; Cock et al. 2017;
Rwomushana et al. 2018; Uzayisenga et al. 2018; Jacobs et al. 2018). By 2018, *S. frugiperda* was reported

present in all countries in Sub-Saharan Africa except Djibouti and Lesotho (Rwomushana et al. 2018).

Furthermore, *S. frugiperda* has now also reached the Asian continent (Deole and Paul 2018; Sisodiya et

al. 2018). Maize and other economically important food crops in these regions are extensively damaged by

S. frugiperda larvae (Day et al. 2017) causing extensive economic losses threatening food security.

There is a lack of information about *S. frugiperda* - host plant interactions and other factors that may be

leading to the rapid spread of *S. frugiperda* in the geographic regions that have recently been invaded.

Many of the control measures used in the western hemisphere (e.g. transgenic maize, chemical

insecticides) might not be readily available and economically viable for subsistence farmers in Africa.

Further, the use of highly hazardous pesticides is not considered a sustainable long term control measure

for any pest (FAO 2018). In addition, *S. frugiperda* is known to readily develop resistance to most chemical

insecticides (e.g. pyrethroids, organophosphates and carbamates) and to transgenic maize that are used

in its control (Yu 1991; Jakka et al. 2016; Banerjee et al. 2017; Flagel et al. 2018; Botha et al. 2019). In light

of this, there is a great need for alternative, cost-effective control strategies for *S. frugiperda* (FAO 2018).

A recent survey in Ethiopia, Kenya and Tanzania indicated that *S. frugiperda* has established interactions

with indigenous parasitoid species (Sisay et al. 2018) that could be harnessed for biological control. A study

on *S. frugiperda* host plant interactions in East Africa has also suggested a climate adapted push-pull

system (Midega et al. 2018) and maize-legume intercropping (Hailu et al., 2018) for management of pests

including fall armyworm on maize farms. However, many factors related to *S. frugiperda* rapid spread, host

plant interactions, bio-ecology and insect-microbiome interactions in the African region remain poorly

understood.

Insect microbiomes can have important consequences for the outcome of insect pest-natural enemies- host

plant interactions (Ferrari et al. 2011). Strategies that involve modifying insect microbiomes are currently

being evaluated for control and management of pests and vectors of plant diseases (Crotti et al. 2012;

Perilla-henao and Casteel 2016; Arora and Douglas 2017; Beck and Vannette 2017). Insect microbiomes

play a key role in the adaptation of insect to their environment and are therefore a major and often poorly

understood determinant of the host plant and geographic range of insect pests (Su et al. 2013). In general,

a greater diversity of microbial symbionts exist within the insect's gut lumen, while few others exist inside

cells of the host, or on the cuticle (Douglas 2016). Gut microbial symbionts are known to influence their

host's nutrition, usually by promoting digestion and availability of nutrients (Douglas 2009). These

symbionts can also modulate the immune response and accessibility of the host to invading organisms,

and therefore have direct or indirect effects on host susceptibility to parasites and pathogens (Dillon et al.

2005; Dong et al. 2009; Garcia et al. 2010; Vorburger et al. 2010; Narasimhan et al. 2014; Mclean and

Godfray 2015; Vorburger and Rouchet 2016; Ubeda et al. 2017). Previous studies have also identified

important roles of bacterial symbionts in the interactions between phytophagous insects and host plants

(Frago et al. 2012; Biere and Bennett 2013; Brady and White 2013). In addition, microbial symbionts can

break down complex molecules such as insecticides and promote insecticide resistance (Kikuchi et al.

2012; Xia et al. 2018). It is also notable that pathogenic bacteria can reside in host guts, only initiating or

facilitating pathogenesis under certain conditions (Wei et al. 2017). Studying the gut microbiome is not only

important from the standpoint of understanding mutualistic relationships but also for the potential

development of microbial biocontrol agents.

There are an increasing number of studies examining the microbial diversity of lepidopterans. While in some

of the assessed species, consistent bacterial communities have been observed in both field and laboratory

collected populations as well as in insects reared on different diets (Broderick et al. 2004; Xiang et al. 2006;

Pinto-Tomás et al. 2011), other studies reported no host specific resident communities that occurred,

regardless of the insect diet (Hammer et al. 2017). It is possible that lepidopterans are less prone to forming

robust 'core' microbiomes due to several factors: 1) very high pH in the midgut, 2) low retention time of

food, 3) lack of microbe housing structures in the intestinal tract, and 4) continual replacement of the

peritrophic matrix (Hammer et al. 2017). Nevertheless, bacterial communities do continually associate with

lepidopterans and influence a variety of important host processes (Broderick et al. 2006; Anand et al. 2010;

Wang et al. 2017).

Few studies have assessed the *Spodoptera*-associated gut microbiome. In a recent study, the microbial

diversity of *Spodoptera exigua* (Hübner) (Lepidoptera: Noctuidae) was examined by 16S rDNA sequence

profiling (Gao et al. 2018). In *Spodoptera exigua,* the dominant bacterial clades are Proteobacteria and

Firmicutes, with the predominant genus in larvae being *Enterococcus*. In *S. frugiperda*, previous studies

have isolated several bacterial strains using culture-dependent methods (De Almeida et al. 2017; Acevedo

et al. 2017). In this study, we used 16S rDNA sequence profiling to characterize the diversity of bacteria

associated with populations of *S. frugiperda* in Kenya and assessed the prevalence of the corn and rice

strains in these populations using Mitochondrial COI gene sequences.

Materials and methods

Insect collection

Spodoptera frugiperda larvae were collected from infested maize fields in Kenya between June and

December 2017 at the following locations: Ngeria (N00.37024 E035.9862) and Burnt Forest (N00.22505

E035.42479) in Uasin Gishu County; Msamia, Kitale (N00.98009 E034.97170) in Trans Nzoia County;

Shimba Hills (S04.33228 E039.34361) in Kwale County and Chala Irrigation Scheme (S03.27338

E037.13816) and Wundanyi (S03.337538 E038.33612) in Taita Taveta County. Part of the field collected

insects from each sampled region in Kenya were reared on fresh maize leaves in ventilated cages to

pupation and eclosion at 27 °C and 60% humidity, while the rest were stored in absolute ethanol at -20°C.

DNA extraction and 16S rDNA sequencing

Guts from 9 live stage 5-6 larvae and 9 one-day old emerging adults from the Kenya collected samples were dissected separately in phosphate buffered saline (PBS) following surface sterilization and used for DNA extraction. Insects were surface sterilized in 70% ethanol, in 5% v/v sodium hypochlorite solution followed by 3 washes in PBS for 3 minutes in each solution. DNA was extracted using the ISOLATE II Genomic DNA Kit (Bioline, London, UK) according to the manufacturer's instructions. DNA extracted from gut samples was submitted for high throuput sequencing targeting the v4 region of the bacterial 16s rRNA gene using the Illumina Miseq platform (Center for Integrated Genomics, University of Lausanne, Switzerland). Sequence reads were checked for quality using FastQC v 0.11.28 (Andrews 2010) and pre-processed to remove adapters and sequencing primers using Cutadapt v1.18 (Martin 2011). Forward and reverse reads were imported into the QIIME2-2018.11 (Boylen et al. 2018). The deblur plugin (Amir et al. 2017) was used to further filter the reads based on per base quality scores, merge the paired-end-reads and cluster reads into operational taxonomic units (OTUs). A total of 457501 sequence reads were retained after removal of spurious reads and all reads shorter than 220 nucleotides in length for further analysis. These sequences clustered into 1796 OTUs. Of these, 197 OTUs survived low count and interquartile range-based variance filtering to eliminate OTUs that could arise from sequencing errors and contamination. Taxonomic assignment was done using the blast classifier against the Silva132 reference database (Quast et al. 2013) at a 99% identity cut-off. OTUs initially characterized as "*Candidatus Hamiltonella*" by comparison to the Silva132 reference database were re-analyzed by homology searches against the NCBI nr nucleotide database through blast (Altschul et al. 1990) and found to be *Pseudomonas*, highlighting a potential incorrect assignment in the reference database. OTU prevalence and variance based filtering as well as alpha and beta diversity measures were applied to the data in the Microbial Analyst Marker Data Profiling (Dhariwal et al. 2017). Shannon diversity indices were applied along with Mann-Whitney and analysis of variance statistics in profiling alpha diversity between sets of samples. Beta diversity was evaluated using Bray-Curtis and unweighted Unifrac distances. Significance testing was done using analysis of group similarities (ANOSIM) and non-metric multidimensional scaling (NMDS) used for ordination. The empirical analysis of digital gene expression data in R (edgeR) algorithm (Robinson et al. 2009) was used to evaluate differential abundance of bacterial genera reads between sample groups. All sequence reads were archived in the Sequence Read Archive (SRA) under the BioProject: PRJNA521837.

mtDNA haplotyping

DNA was extracted from surface-sterilized whole insects using the ISOLATE II Genomic DNA Kit (Bioline, London, UK) according to the manufacturer's instructions. Mitochondrial COI gene sequences were amplified from insect DNA by PCR using the primer LCO1490 and HCO2198 (Folmer et al. 1994). Reactions were set up in total volumes of 10 µl each, containing 5× MyTaq reaction buffer (5 mM dNTPs, 15 mM MgCl2, stabilizers and enhancers) (Bioline, London, UK), 2 µM of each primer, 0.25 mM MgCl2 (Thermo Fischer Scientific, Massachusetts, USA), 0.125 µl MyTaq DNA polymerase (Bioline, London, UK), and 7.5 ng/µl of DNA template. These reactions were set up in a Master cycler Nexus gradient thermo-

cycler (Thermo Fischer Scientific, Massachusetts, USA) using the following cycling conditions: initial
denaturation for 2 min at 95 °C, followed by 40 cycles of 30 s at 95 °C, 45 s at 50.6 °C and 1 min at 72 °C,
then a final elongation step of 10 min at 72 °C. PCR products were run through 1% agarose gel
electrophoresis and visualized by ethidium bromide staining and UV trans-illumination. Direct sequencing
was done for all host mtCOI gene and the sequences deposited in the GenBank.

Results

We profiled the bacterial microbiome for 18 samples from 4 different locations in Kenya. In addition,
samples were collected from these 4 sites plus two additional sites for mtDNA haplotyping (Fig. 1).

Fig. 1 Sites from which *Spodoptera frugiperda* larvae were collected in Kenya

The most abundant bacterial Phyla observed across the fall armyworm gut samples were Proteobacteria,

Firmicutes, Bacteroidetes and a small proportion of Actinobacteria (Fig.10, Supplementary material). OTUs

clustering in the orders Enterobacteriales and Pseudomonadales were predominant in the majority of the

samples (Fig. 2).

Fig. 2 Composition of bacterial OTUs at Order level in screened *Spodoptera frugiperda* samples

We note that despite the high genus-level diversity between samples (Fig. 3), there were some similarities

based on developmental stage and location. For example, there was a very high proportion of: 1)

Pseudomonas in the two adult male samples from Chala, 2) *Citrobacter* in two larval samples from Kwale,

3) *Lysinibacillus* in two male samples from Kitale and 4) *Enterococcus* in two larval samples from Ngeria.

It was noted that *Stenotrophomonas*, *Sphingobacterium*, *Serratia*, *Pseudomonas*, *Morganella*,

Enterococcus and *Delftia* were observed in both larvae and adult samples.

Fig. 3 Genus level composition of (% of OTUs) in the different samples of *S. frugiperda*. The relative abundances of the 19 most abundant genera are represented here

In one of the larval samples from the Ngeria site (Ngeria-l2), we observed an excessive number of non-bacterial reads. Upon closer inspection, these were found to be closely related to *Metarhizium rileyi* (Farl.) Kepler, Rehner & Humber (2014) (formerly *Nomuraea rileyi*)*,* an entomopathogenic fungi that is known to infect *S. frugiperda* (Fig. 4).

171   0.9

Fig. 4 Neighbor-joining tree of fungus OTU detected in *S. frugiperda* sample (Ngeria-l2; in bold) and GenBank accessions of small subunit ribosomal RNA gene sequences from related fungi. Sequences are labelled by their GeneBank Accessions followed by genus, species and strain where available. Bootstrap values are indicated above the branches. Branches with a bootstrap value less than 50 are collapsed. A sequence from a species in the genus *Rozella* is included as an out-group

The bacterial OTU richness appeared to be higher in *S. frugiperda* larvae than adults, however this difference was not statistically significant (p-value: 0.062526; [Mann-Whitney] statistic: 19) using Shannon diversity metrics (Fig. 5a). In addition, no significant variation in OTU richness and abundance was observed between larvae from different sampling sites (p-value: 0.32834; [ANOVA] F-value: 1.3486) (Fig. 5b).

Fig. 5 A comparison of the Shannon diversity indices for: **(a)** adult and larval samples from all sites and **(b)** larvae collected from different sites. The Shannon diversity index (H') was calculated based on the OTU-level of classification. The boxplots show the distribution of H' values across all samples

The composition of bacterial OTUs between larvae and adult *S. frugiperda* was observed to overlap, with no significant dissimilarity ([ANOSIM] R=0.17365; p-value < 0.081 stress=0.14876) (Fig. 6a). However, OTU composition was observed to vary significantly among larval samples from different sites ([ANOSIM] R: 0.45679; p-value < 0.017 stress=0.05711) (Fig. 6b).

Fig. 6a Non-metric multidimensional scaling (NMDS) ordination based on Bray-Curtis dissimilarities in bacterial communities detected in the *S. frugiperda* samples. Samples are colored according to their developmental stage and sex as indicated on the legend where F=female, L=larvae and M=male

Fig. 6b Non-metric multidimensional scaling (NMDS) ordination based on Weighted Unifrac distance
dissimilarities in bacterial OTU composition between larval sample pairs from different sites

A significant differential abundance was observed for 3 bacterial genera between larvae and adult *S.*
frugiperda samples using the EdgeR algorithm at an adjusted p-value of 0.05. Two of these: *Citrobacter*
(log2FC=4.4178, p value=3.6E-6, FDR=7.218E-5) and *Sphingobacterium* (log2FC=3.625, p value=1.01E-
4, FDR=0.0010118) were more abundant in larvae whereas the third: *Lysinibacillus* (log2FC=-3.2247, p
value= 4.4E-3, FDR=0.029375) was more abundant in adults (Fig. 7).

202

203 **Fig. 7** Comparative abundance of A) *Citrobacter,* B) *Lysinibacillus* and C) *Sphingobacterium* between adults
204 and larvae of *S. frugiperda.* Abundance is shown on a log transformed scale of original counts

205 Based on mtDNA sequences, the *S. frugiperda* strains detected in this study were identical to strains from

206 Canada, USA and Brazil, as well as strains that were recently reported in Kenya and other parts of Africa

207 and India (Fig. 8).

KJ634291_S_exigua
MH639006_India
R-strain_Kenya
GU439148_Canada
JQ547900_Costa Rica
KY472249_Northern-region_Ghana
MF593251_South_Africa
MH753330_Badravathi_India
HM136593_USA
KX580616_Ibadan_Nigeria
KY472253_Volta-region_Ghana
MH190444_KEPH_A_Kenya
KY472250_Northern-region_Ghana
MG993205_Malawi
U72977_H1_USA
98 KX580618_Ibadan_Nigeria
KY472240_Brong-Ahafo_Ghana
MF197867_Uganda
MH190445_KEPH_E1_Kenya
HM136602_USA
MH639004_India
KY472245_Brong-Ahafo_Ghana
MF278659_Tanzania
MH639005_India
KY472242_Brong-ahafo_Ghana
MH639007_India
MF278657_Tanzania
MH190446_KEPH_C_Kenya
64 JF854745_Brazil
JF854746_Canada
MH819361_East_Godovari
63 MH819360_Ujjain_India
MH819359_Pune_India
MH819358_Anakapalli_India
MH819357_India
MH819356_West-Godovari_India
MH819355_Hyderabad_India
MH819354_Jammikunta_India
MH819353_India
MH753333_Haniyuru_India
MH819352_Bengaluru_India
U72976_H3_USA
78 MH190448_KEPH_N4_Kenya
MH190447_KEPH_E2_Kenya
MF593241_South_Africa
MF197868_Uganda
KY472251_Volta-region_Ghana
KY472248_Northern-region_Ghana
KX580615_ Porto_Allegre_Sao_Tome
KX580614_Mesquita_Sao_Tome
KF624877_Roraima_Brazil
HQ964527_ USA
HM136586_USA
C-strain_Kenya
GU094754_Canada

0.02

Fig 8 Neighbor-joining tree based on mtCOI sequences of *S. frugiperda* from the GenBank and
representative haplotypes from this study (in bold). Bootstrap values are indicated above branches.
Branches with bootstrap values less than 50 are collapsed. A sequence from *Spodoptera exigua is* included
as an out-group. Sequences are labelled with their GenBank accession numbers, collection site where
available and country of collection

All the samples clustered in two major clades widely referred to as either the 'Rice' or the 'Corn' strain

(hereafter referred to as R- strain and C- strain). We investigated the frequency of mtDNA haplotypes of *S.*

frugiperda samples collected at several sites in Kenya. Overall, 90% of the samples (n=85) clustered as R-

strain, whereas 10% (n=9) clustered as C-strain. Proportions of the R-strain in populations at the different

sites were 100% (n=6) for Burnt Forest, 83% (n=6) for Chala, 86% (n=7) for Wundanyi, 82% (n=11) for

Kitale, 91% (n=35) for Kwale and 82% (n=17) for Ngeria (Fig. 9).

Fig. 9 Mitochondrial COI haplotype map of the *S. frugiperda* samples collected at various sites in Kenya.
Node size is proportional to number of samples and individual samples are represented as fractions of the
nodes. A sequence of *Spoladea recurvalis* is included as an out-group. Sequences for all samples are
accessible from GenBank using the accessions: MK492929-MK493010

Discussion

We found that the gut bacterial communities of most *S. frugiperda* samples were dominated by

Proteobacteria. This observation is similar to proportions reported in other phytophagous insects, in

particular lepidopterans (Belda et al. 2011; Xia et al. 2013, 2017; Landry et al. 2015; Ramya et al. 2016;

Snyman et al. 2016; Strano et al. 2017; Chen et al. 2018). Only three samples, two adult males from Kitale

(Kitale-m2 and Kitale-m3) and one larvae from Ngeria (Ngeria-l2) were dominated by Firmicutes. Four

genera of bacteria, *Pseudomonas*, *Delftia*, *Enterococcus* and *Serratia* that were recorded in this study have

previously been isolated from *S. frugiperda* (De Almeida et al. 2017; Acevedo et al. 2017). Surprisingly,
Staphylococcus, *Microbacterium*, *Arthrobacter* and *Leclercia* that were previously isolated from *S.*
frugiperda in Brazil (De Almeida et al. 2017) were not found in any of the samples we profiled in Kenya.
Similarly, *Pantoea*, *Enterobacter*, *Raoultella* and *Klebsiella* previously identified in oral secretions of *S.*
frugiperda in Pennsylvania, USA (Acevedo et al. 2017) were not found in the profiled Kenyan samples.

We observed significant differences in OTU composition between larvae from different sites. This was most
likely caused by complex biological and environmental factors in the diverse agro-ecological zones that
were sampled. Diet is known to strongly influence the microbiome of lepidopterans (Strano et al. 2017;
Sittenfeld et al. 2002; Priya et al. 2012; Montagna et al. 2016), however in this study all samples were
collected from maize plants. Hence, the observed compositional differences are not likely to be caused
solely by diet. We observed differences in bacterial OTU composition between larvae and adults, however
with a relatively low number of samples these differences were not statistically significant. It is interesting
that many of the detected bacterial genera such as *Stenotrophomonas*, *Sphingobacterium*, *Serratia*,
Pseudomonas, *Morganella*, *Enterococcus* and *Delftia* are found in both life stages, which suggests that gut
bacterial community members are transmitted across developmental stages. Bacteria that are continually
transmitted across developmental stages (and across generations) may evolve a closer, mutualistic
relationship with their hosts (Moran 2006). Future studies should investigate the effects of these microbes
on host fitness and investigate the extent to which they are vertically transmitted from parents to offspring.
In contrast, *Citrobacter* and *Sphingobacterium* were observed to be differentially abundant in larvae than in
adults, a likely indicator that these two genera may be part of the fraction of bacterial communities that are
lost during transition of *S. frugiperda* into the adult stage. *Lysinibacillus*, on the other hand, was more
abundant in adults than in larvae and therefore could have an adult-specific function.

Notably, we identified *Serratia*, *Lysinibacillus* (formerly *Bacillus*) and *Pseudomonas*, species of which have
been reported to have entomopathogenic properties (Castagnola and Stock 2014). In addition, one sample
had a high number of reads attributed to a relative of a non-bacterial entomopathogen, *Metarhizium rileyi,*
which has been previously isolated and tested for efficiency against *S. frugiperda* (Maniania and Fargues
1985; Mallapur et al. 2018). It will be worthwhile to explore the pathogenicity of these microbes for *S.*
frugiperda and to determine if they could be incorporated into biological pest management strategies (Ruiu
et al. 2015).

Based on the mtCOI gene sequence, we observed two mtDNA haplotypes in Kenya (C- and R- strains).
These findings confirm that both haplotypes are present in Kenya, as has been demonstrated for other
countries in Africa (Rwomushana et al. 2018). The majority of the *S. frugiperda* samples collected were
characterized as R-strain suggesting that this strain is dominant in *S. frugiperda* populations in Kenya.
These observations are in agreement with a previous study (Goergen et al. 2016) that observed C- and R-
strains appear to have an East-West axis alignment in the African region with the Eastern Africa having

progressively lower frequencies of the mtCOI C-strain (Goergen et al. 2016). We observed that some

variants of the rice strain have been reported in other places such as Ghana and India but those were not

detected in this study. It is interesting to note that in addition to a similar rice strain as the one detected in

Kenya, a variant differing by a single nucleotide polymorphism has been recorded from various locations in

India (Fig. 8). This variant has however not been reported in Africa. It is therefore possible that the invasion

into India may not have come directly from the African continent, or invasion could have included strains

from Africa and elsewhere.

Symbiotic bacteria play a key role in the biology of insects. We characterized the gut bacterial communities

in *S. frugiperda* larvae and adult samples collected from several locations in Kenya, finding some important

differences and similarities across samples and in relation to other studies on this species (Acevedo et al.

2017; De Almeida et al. 2017). Understanding the gut microbial symbionts of this pest species may facilitate

the development of novel, cost-effective control strategies.

Acknowledgements

The authors gratefully acknowledge the financial support for this research by the following organizations
and agencies: European Union Funded project "Integrated pest management" strategy to counter the
threat of invasive fall armyworm to food security in Eastern Africa (FAW-IPM) (FOOD/2018/402-634); UK
Aid from the UK Government; Swedish International Development Cooperation Agency (Sida); the Swiss
Agency for Development and Cooperation (SDC); and the Kenyan Government. Financial assistance for
sequencing at the Center for Integrated Genomics, Lausanne, was generously provided by the Global
Health Institute of the École Polytechnique Fédérale de Lausanne. The views expressed herein do not
necessarily reflect the official opinion of the donors. Authors also acknowledge the technical assistance of
Mr. Peter Malusi during sample collection.

Author contributions

SE, SS, JH and FMK conceived and designed the research, JG conducted experiments and analyzed
data, JG and JH wrote the manuscript, JVB and HP contributed materials. All authors read and approved
the manuscript

Conflict of Interest

The authors declare that they have no conflict of interest.

References

Acevedo FE, Peiffer M, Tan C-W, Stanley BA, Stanley A, Wang J, Jones AG, Hoover K, Rosa C, Luthe D, Felton G (2017) Fall armyworm-associated gut bacteria modulate plant defense responses. Mol Plant-Microbe Interact 30:127–137. doi: 10.1094/MPMI-11-16-0240-R

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ (1990) Basic local alignment search tool. J Mol Biol 215:403–410

Amir A, McDonald D, Navas-Molina, Jose A. Kopylova E, Morton JT, Xu ZZ, Kightley EP, Thompson LR, Hyde ER, Gonzalez A, Rob K (2017) Deblur rapidly resolves single-nucleotide community sequence patterns. mSystems 2:1–7

Anand AAP, Vennison SJ, Sankar SG, Prabhu DIG, Vasan PT, Raghuraman T, Geoffrey CJ, Vendan SE (2010) Isolation and characterization of bacteria from the gut of *Bombyx mori* that degrade cellulose, xylan, pectin and starch and their impact on digestion. J Insect Sci 10:1–20. doi: 10.1673/031.010.10701

Andrews S (2010) FastQC: a quality control tool for high throughput sequence data. http://www.bioinformatics.babraham.ac.uk/projects/fastqc

Arora AK, Douglas AE (2017) Hype or opportunity? Using microbial symbionts in novel strategies for insect pest control. J Insect Physiol 103:10–17. doi: 10.1016/j.jinsphys.2017.09.011

Banerjee R, Hasler J, Meagher R, Nagoshi R, Hietala L, Huang F, Narva K, Jurat-Fuentes JL (2017) Mechanism and DNA-based detection of field-evolved resistance to transgenic Bt corn in fall armyworm (*Spodoptera frugiperda*). Sci Rep 7:1–10. doi: 10.1038/s41598-017-09866-y

Beck JJ, Vannette RL (2017) Harnessing insect-microbe chemical communications to control insect pests of agricultural systems. J Agric Food Chem 65:23–28. doi: 10.1021/acs.jafc.6b04298

Belda E, Pedrola L, Peretó J, Martínez-Blanch JF, Montagud A, Navarro E, Urchueguía J, Ramón D, Moya A, Porcar M (2011) Microbial diversity in the midguts of field and lab-reared populations of the European corn borer *Ostrinia nubilalis*. PLoS One 6:e21751. doi: 10.1371/journal.pone.0021751

Biere A, Bennett AE (2013) Three-way interactions between plants, microbes and insects. Funct Ecol 27:567–573. doi: 10.1111/1365-2435.12100

Bolyen, E., Rideout, J.R., Dillon, M.R., Bokulich, N.A., Abnet, C., Al-Ghalith, G.A., Alexander, H., Alm, E.J., Arumugam, M., Asnicar, F., Bai, Y., Bisanz, J.E., Bittinger, K., Brejnrod, A., Brislawn, C.J., Brown, C.T., Callahan, B.J., Caraballo-Rodríguez, A.M., Chase, J., Cope, E., Da Silva, R., Dorrestein, P.C., Douglas, G.M., Durall, D.M., Duvallet, C., Edwardson, C.F., Ernst, M., Estaki, M., Fouquier, J., Gauglitz, J.M., Gibson, D.L., Gonzalez, A., Gorlick, K., Guo, J., Hillmann, B., Holmes, S., Holste, H., Huttenhower, C., Huttley, G., Janssen, S., Jarmusch, A.K., Jiang, L., Kaehler, B., Kang, K.B., Keefe, C.R., Keim, P., Kelley, S.T., Knights, D., Koester, I., Kosciolek, T., Kreps, J., Langille, M.G.I., Lee, J., Ley, R., Liu, Y.-X., Loftfield, E., Lozupone, C., Maher, M., Marotz, C., Martin, B., McDonald, D., McIver, L.J., Melnik, A.V., Metcalf, J.L., Morgan, S.C., Morton, J., Naimey, A.T., Navas-Molina, J.A., Nothias, L.F., Orchanian, S.B., Pearson, T., Peoples, S.L., Petras, D., Preuss, M.L., Pruesse, E., Rasmussen, L.B., Rivers, A., Robeson, II, M.S., Rosenthal, P., Segata, N., Shaffer, M., Shiffer, A., Sinha, R., Song, S.J., Spear, J.R., Swafford, A.D., Thompson, L.R., Torres, P.J., Trinh, P., Tripathi, A., Turnbaugh, P.J., Ul-Hasan, S., van der Hooft, J.J.J., Vargas, F., Vázquez-Baeza, Y., Vogtmann, E., von Hippel, M., Walters, W., Wan, Y., Wang, M., Warren, J., Weber, K.C., Williamson, C.H.D., Willis, A.D., Xu, Z.Z., Zaneveld, J.R., Zhang, Y., Knight, R., Caporaso, J.G.: (2018) QIIME 2: reproducible, interactive, scalable, and extensible microbiome data science. PeerJ. doi: 10.7287/peerj.preprints.27295

Brady CM, White JA (2013) Cowpea aphid (*Aphis craccivora*) associated with different host plants has different facultative endosymbionts. Ecol Entomol 1–5. doi: 10.1111/een.12020

Broderick AN, Raffa KF, Handelsman J (2006) Midgut bacteria required for *Bacillus thuringiensis* insecticidal activity. PNAS 103:15196–15199. doi: 10.1371/journal.pone.0170933

Broderick N, Raffa K, Goodman R, Handelsman J (2004) Census of the bacterial community of the gypsy moth larval midgut by using culturing and culture-independent methods. Appl Environ Microbiol 70:293–300. doi: 10.1128/AEM.70.1.293

Castagnola A, Stock SP (2014) Common virulence factors and tissue targets of entomopathogenic bacteria for biological control of Lepidopteran pests. Insects 5:139–166. doi:

10.3390/insects5010139

Chen B, Du K, Sun C, Vimalanathan A, Liang X, Li Y, Wang B, Lu X, Li L, Shao Y (2018) Gut bacterial
and fungal communities of the domesticated silkworm (*Bombyx mori*) and wild mulberry-feeding
relatives. ISME J 12:2252–2262. doi: 10.1038/s41396-018-0174-1

Cock MJW, Beseh PK, Buddie AG, Cafá G, Crozier J (2017) Molecular methods to detect *Spodoptera*
frugiperda in Ghana, and implications for monitoring the spread of invasive species in developing
countries. Sci Rep 7:1–10. doi: 10.1038/s41598-017-04238-y

Crotti E, Balloi A, Hamdi C, Sansonno L, Marzorati M, Gonella E, Favia G, Cherif A, Bandi C, Alma A,
Daffonchio D (2012) Review microbial symbionts: a resource for the management of insect-related
problems. Microb Biotechnol 5:307–317. doi: 10.1111/j.1751-7915.2011.00312.x

Day R, Abrahams P, Bateman M, Beale T, Clottey V, Cock M, Colmenarez Y, Corniani N, Early R,
Godwin J, Gomez J, Moreno PG, Murphy ST, Birgitta O-M, Phiri N, Pratt C, Silvestri S, Witt A (2017)
Fall armyworm: impacts and implications for Africa. Outlooks Pest Manag 2016:196–201. doi:
10.1564/v28

De Almeida LG, De Moraes LAB, Trigo JR, Omoto C, Cônsoli FL (2017) The gut microbiota of insecticide-
resistant insects houses insecticide-degrading bacteria: a potential source for biotechnological
exploitation. PLoS One 12:1–19. doi: 10.1371/journal.pone.0174754

Deole S, Paul N (2018) First report of fall army worm, *Spodoptera frugiperda* (J. E. Smith), their nature of
damage and biology on maize crop at Raipur, Chhattisgarh. J Entomol Zool Stud 6:219–221

Dhariwal A, Chong J, Habib S, King IL, Agellon LB, Xia J (2017) MicrobiomeAnalyst: a web-based tool for
comprehensive statistical, visual and meta-analysis of microbiome data. Nucleic Acids Res
45:W180–W188. doi: 10.1093/nar/gkx295

Dillon RJ, Vennard CT, Buckling A, Charnley AK (2005) Diversity of locust gut bacteria protects against
pathogen invasion. Ecol Lett 8:1291–1298. doi: 10.1111/j.1461-0248.2005.00828.x

Dong Y, Manfredini F, Dimopoulos G (2009) Implication of the mosquito midgut microbiota in the defense
against malaria parasites. PLOS Pathog 5:e1000423. doi: 10.1371/journal.ppat.1000423

Douglas AE (2009) The microbial dimension in insect nutritional ecology. Funct Ecol 23:38–47. doi:
10.1111/j.1365-2435.2008.01442.x

Douglas AE (2016) Multiorganismal insects: diversity and function of resident microorganisms. Annu Rev
Entomol 60:17–34. doi: 10.1146/annurev-ento-010814-020822

FAO (2018) Sustainable management of the fall armyworm in Africa: a framework for partnership. Rome,
Italy. http://www.fao.org/3/I9160EN/i9160en.pdf

Ferrari J, Vavre F, Lyon D (2011) Bacterial symbionts in insects or the story of communities affecting
communities. Philos Trans R Soc Boyal Soc B 366:1389–1400. doi: 10.1098/rstb.2010.0226

Flagel L, Lee YW, Wanjugi H, Swarup S, Brown A, Wang J, Kraft E, Greenplate J, Simmons J, Adams N,
Wang Y, Martinelli S, Haas JA, Gowda A, Head G (2018) Mutational disruption of the ABCC2 gene
in fall armyworm, *Spodoptera frugiperda*, confers resistance to the Cry1Fa and Cry1A.105
insecticidal proteins. Sci Rep 8:1–11. doi: 10.1038/s41598-018-25491-9

Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R (1994) DNA primers for amplification of mitochondrial
cytochrome c oxidase subunit I from diverse metazoan invertebrates. Mol Mar Biol Biotechnol
3:294–299. doi: 10.1371/journal.pone.0013102

Frago E, Dicke M, Godfray HCJ (2012) Insect symbionts as hidden players in insect-plant interactions.
Trends Ecol Evol 27:705–711. doi: 10.1016/j.tree.2012.08.013

Gao X, Li W, Luo J, Zhang L, Ji J, Zhu X, Wang L, Cui J (2018) Biodiversity of the microbiota in
Spodoptera exigua (Lepidoptera: Noctuidae). J Appl Microbiol 126:1199–1208. doi:
https://doi.org/10.1111/jam.14190

Garcia ES, Castro DP, Figueiredo MB, Azambuja P (2010) Immune homeostasis to microorganisms in
the guts of triatomines (Reduviidae) - a review. Mem Inst Oswaldo Cruz 105:605–610

Goergen G, Kumar PL, Sankung SB, Togola A, Tamò M (2016) First report of outbreaks of the fall
armyworm *Spodoptera frugiperda* (J E Smith) (Lepidoptera, Noctuidae), a new alien invasive pest in
West and Central Africa. PLoS One 11:1–9. doi: 10.1371/journal.pone.0165632

Hammer TJ, H. Janzen D, Hallwachs W, Jaffe SP, Fierera N (2017) Caterpillars lack a resident gut
microbiome. PNAS 114:9641–9646. doi: 10.1073/pnas.1707186114

Huber DM, Hugh-Jones ME, Rust MK, Sheffield SR, Simberloff D, Taylor CR (2002) Invasive pest
species: impacts on agricultural production, natural resources and the environment. Counc Agric Sci
Technol 20:1–18

Jacobs A, van Vuuren A, Rong IH (2018) Characterisation of the fall armyworm (*Spodoptera frugiperda* J.E. Smith) (Lepidoptera: Noctuidae) from South Africa. African Entomol 26:45–49. doi: 10.4001/003.026.0045

Jakka SRK, Gong L, Hasler J, Banerjee R, Sheets JJ, Narva K, Blanco CA, Jurat-Fuentes JL (2016) Field-evolved mode 1 resistance of the fall armyworm to transgenic Cry1Fa-expressing corn associated with reduced Cry1Fa toxin binding andmidgut alkaline phosphatase expression. Appl Environ Microbiol 82:1023–1034. doi: 10.1128/aem.02871-15

Kikuchi Y, Hayatsu M, Hosokawa T, Nagayama A, Tago K (2012) Symbiont-mediated insecticide resistance. Proc Natl Acad Sci 109:8618–8622. doi: 10.1073/pnas.1200231109/-/DCSupplemental.www.pnas.org/cgi/doi/10.1073/pnas.1200231109

Landry M, Comeau AM, Derome N, Cusson M, Levesque RC (2015) Composition of the spruce budworm (*Choristoneura fumiferana*) midgut microbiota as affected by rearing conditions. PLoS One 10:1–11. doi: 10.1371/journal.pone.0144077

Mallapur CP, Naik AK, Hagari S, Praveen T, Patil RK (2018) Potentiality of *Nomuraea rileyi* (Farlow) Samson against the fall armyworm, *Spodoptera frugiperda* (J E Smith) infesting maize. J Entomol Zool Stud 6:1062–1067

Maniania NK, Fargues J (1985) Susceptibility of the fall armyworm, *Spodoptera frugiperda*, to the fungal pathogens *Paecilomyces fumosoroseus* and *Nomuraea rileyi*. Florida Entomol 68:178–183

Martin M (2011) Cutadapt removes adapter sequences from high-throughput sequencing reads. EMBnet.journal 17:10. doi: 10.14806/ej.17.1.200

Mclean AHC, Godfray HCJ (2015) Evidence for specificity in symbiont-conferred protection against parasitoids. Proc R Soc B Biol Sci 282:1–8

Midega CAO, Pittchar JO, Pickett JA, Hailu GW, Khan ZR (2018) A climate-adapted push-pull system effectively controls fall armyworm, *Spodoptera frugiperda* (J E Smith), in maize in East Africa. Crop Prot 105:10–15. doi: 10.1016/j.cropro.2017.11.003

Montagna M, Mereghetti V, Gargari G, Guglielmetti S, Faoro F, Locatelli D, Limonta L (2016) Evidence of a bacterial core in the stored products pest *Plodia interpunctella*: the influence of different diets. Environ Microbiol 18:4961–4973

Moran NA (2006) Symbiosis. Curr Biol 16:R866-871

Nagoshi RN, Goergen G, Tounou KA, Agboka K, Koffi D, Meagher RL (2018) Analysis of strain distribution, migratory potential, and invasion history of fall armyworm populations in northern Sub-Saharan Africa. Sci Rep 8:3710. doi: 10.1038/s41598-018-21954-1

Nagoshi RN, Koffi D, Agboka K, Tounou KA, Banerjee R, Jurat-Fuentes JL, Meagher RL (2017) Comparative molecular analyses of invasive fall armyworm in Togo reveal strong similarities to populations from the eastern United States and the Greater Antilles. PLoS One 12:1–15. doi: 10.1371/journal.pone.0181982

Narasimhan S, Rajeevan N, Liu L, Zhao YO, Heisig J, Pan J, Eppler-epstein R, Deponte K, Fish D, Fikrig E (2014) Gut microbiota of the tick vector *Ixodes scapularis* modulate colonization of the lyme disease spirochete. Cell Host Microbe 15:58–71. doi: 10.1016/j.chom.2013.12.001

Perilla-henao LM, Casteel CL (2016) Vector-borne bacterial plant pathogens: interactions with Hemipteran insects and plants. Front Plant Sci 7:1–15. doi: 10.3389/fpls.2016.01163

Pinto-Tomás AA, Sittenfeld A, Uribe-Lorío L, Chavarría F, Mora M, Janzen DH, Goodman RM, Simon HM (2011) Comparison of midgut bacterial diversity in tropical caterpillars (Lepidoptera: Saturniidae) fed on different diets. Environ Entomol 40:1111–1122. doi: 10.1016/S0140-6736(02)87727-3

Priya NG, Ojha A, Kajla MK, Raj A, Rajagopal R (2012) Host plant induced variation in gut bacteria of *Helicoverpa armigera*. PLoS One 7:1–10. doi: 10.1371/journal.pone.0030768

Quast C, Prusse EP, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glockner FO (2013) The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. Nucleic Acids Res 41:D590–D596. doi: 10.1093/nar/gks1219

Ramya SL, Venkatesan T, Srinivasa Murthy KS, Jalali SK, Verghese A (2016) Detection of carboxylesterase and esterase activity in culturable gut bacterial flora isolated from diamondback moth, *Plutella xylostella* (Linnaeus), from India and its possible role in indoxacarb degradation. Brazilian J Microbiol 47:327–336. doi: 10.1016/j.bjm.2016.01.012

Robinson MD, McCarthy DJ, Smyth GK (2009) edgeR: A Bioconductor package for differential expression analysis of digital gene expression data. Bioinformatics 26:139–140. doi: 10.1093/bioinformatics/btp616

Ruiu L, Agraria D, Vegetale P (2015) Insect pathogenic bacteria in integrated pest management. Insects
6:352–367. doi: 10.3390/insects6020352
Rwomushana I, Bateman M, Beale T, Beseh P, Cameron K, Chiluba M, Clottey V, Davis T, Day R, Early
R, Godwin J, Gonzalez-Moreno P, Kansiime M, Kenis M, Makale F, Mugambi I, Murphy S, Nunda
W, Phiri N, Pratt C, Tambo J (2018) Fall armyworm: impacts and implications for Africa evidence note
update, October 2018
Sisay B, Simiyu J, Malusi P, Likhayo P, Mendesil E, Elibariki N, Wakgari M, Ayalew G, Tefera T (2018)
First report of the fall armyworm, *Spodoptera frugiperda* (Lepidoptera : Noctuidae), natural enemies
from Africa. J Appl Entomol 142:800–804. doi: 10.1111/jen.12534
Sisodiya DB, Raghunandan BL, Bhatt NA, Verma HS, Shewale CP, Timbadiya BG, Borad PK (2018) The
fall armyworm, *Spodoptera frugiperda* (J. E. Smith) (Lepidoptera: Noctuidae); first report of new
invasive pest in maize fields of Gujarat, India. J Entomol Zool Stud 6:2089–2091
Sittenfeld A, Uribe-lorío L, Mora M, Nielsen V, Arrieta G, Janzen DH (2002) Does a polyphagous
caterpillar have the same gut microbiota when feeding on different species of food plants? Rev Biol
Trop 50:547–560
Strano CP, Malacrinò A, Campolo O, Palmeri V (2017) Influence of host plant on *Thaumetopoea*
pityocampa gut bacterial community. Microb Ecol 75:487–494. doi: 10.1007/s00248-017-1019-6
Su Q, Zhou X, Zhang Y (2013) Symbiont-mediated functions in insect hosts. Commun Integr Biol
6:e23804 1-7. doi: https://doi.org/10.4161/cib.23804
Tindo M, Tagne A, Tigui A, Kengni F, Atanga J, Bila S, Abega R (2016) First report of the fall army worm,
Spodoptera frugiperda (Lepidoptera, Noctuidae) in Cameroon. Cameroon J Biol Biochem Sci 25:30–
32
Ubeda C, Djukovic A, Isaac S (2017) Roles of the intestinal microbiota in pathogen protection. Clin Transl
Immunol 6:1–10. doi: 10.1038/cti.2017.2
Uzayisenga B, Waweru B, Kajuga J, Karangwa P, Uwumukiza B, Edgington S, Thompson E, Offord L,
Cafá G, Buddie A (2018) First record of the fall armyworm, *Spodoptera frugiperda* (JE Smith,
1797)(Lepidoptera: Noctuidae), in Rwanda. African Entomol 26:244–246
Vorburger C, Gehrer L, Rodriguez P (2010) A strain of the bacterial symbiont *Regiella insecticola* protects
aphids against parasitoids. Biol Lett 6:109–111
Vorburger C, Rouchet R (2016) Are aphid parasitoids locally adapted to the prevalence of defensive
symbionts in their hosts? BMC Evol Biol 16:1–11. doi: 10.1186/s12862-016-0811-0
Wang J, Peiffer M, Hoover K, Rosa C, Zeng R, Felton GW (2017) *Helicoverpa zea* gut-associated
bacteria indirectly induce defenses in tomato by triggering a salivary elicitor(s). New Phytol
214:1294–1306. doi: 10.1111/nph.14429
Wei G, Lai Y, Wang G, Chen H, Li F, Wang S (2017) Insect pathogenic fungus interacts with the gut
microbiota to accelerate mosquito mortality. PNAS 114:5994–5999. doi: 10.1073/pnas.1703546114
Xia X, Gurr GM, Vasseur L, Zheng D, Zhong H, Qin B, Lin J, Wang Y, Song F, Li Y, Lin H, You M (2017)
Metagenomic sequencing of diamondback moth gut microbiome unveils key holobiont adaptations
for herbivory. Front Microbiol 8:1–12. doi: 10.3389/fmicb.2017.00663
Xia X, Sun B, Gurr GM, Vasseur L, Xue M, You M (2018) Gut microbiota mediate insecticide resistance in
the diamondback moth, *Plutella xylostella* (L.). Front Microbiol 9:1–10. doi:
10.3389/fmicb.2018.00025
Xia X, Zheng D, Zhong H, Qin B, Gurr GM, Vasseur L, Lin H, Bai J, He W, You M (2013) DNA
sequencing reveals the midgut microbiota of diamondback moth, *Plutella xylostella* (L.) and a
possible relationship with insecticide resistance. PLoS One 8:1–8. doi:
10.1371/journal.pone.0068852
Xiang H, Wei G-F, Jia S, Huang J, Miao X-X, Zhou Z, Zhao L-P, Huang Y-P (2006) Microbial communities
in the larval midgut of laboratory and field populations of cotton bollworm (*Helicoverpa armigera*).
Can J Microbiol 52:1085–1092. doi: 10.1139/W06-064
Yu SJ (1991) Insecticide resistance in the fall armyworm, *Spodoptera frugiperda*. Pestic Biochem Physiol
39:84–91

521 **Supplementary material**

523 Fig. 10 Relative abundance of bacterial phyla detected in *S. frugiperda* samples