Zebrafish *mbnl* mutants model molecular and physical phenotypes of human

***MBNL* loss of function disorders**

Melissa N. Hinman[1], Jared I. Richardson[4,6], Rose A. Sockol[1], Eliza D Aronson[1], Sarah

J. Stednitz[2], Katrina N. Murray[3], J. Andrew Berglund[4,6], and Karen Guillemin[1,5*]

[1]Institute of Molecular Biology, [2]Institute of Neuroscience, [3]Zebrafish International

Resource Center, University of Oregon, Eugene, Oregon, [4]RNA Institute, State

University of New York at Albany, Albany, New York, [5]Humans and the Microbiome

Program, CIFAR, Toronto, Ontario M5G 1Z8, Canada, [6]Center for NeuroGenetics,

University of Florida, Gainesville, Florida

*Corresponding author

Corresponding author email address: kguillem@uoregon.edu

Corresponding author ORCID identifier: https://orcid.org/0000-0001-6004-9955

Summary Statement

21 Zebrafish *mbnl* mutants that were generated to model myotonic dystrophy and Fuchs

22 endothelial corneal dystrophy exhibited disease-relevant phenotypes including

23 alternative splicing changes, decreased body size, and impaired movement.

Abstract

The muscleblind RNA binding proteins (MBNL1, MBNL2, and MBNL3) are highly conserved across vertebrate species and are important regulators of alternative splicing and other aspects of RNA metabolism. Loss of MBNL protein function through sequestration by CUG or CCUG RNA repeats is partially responsible for the phenotypes of two important human genetic disorders, myotonic dystrophy (DM) and Fuchs endothelial corneal dystrophy (FECD). We generated the first stable zebrafish (*Danio rerio*) models of *MBNL* loss of function disorders through CRISPR-based mutation of the three zebrafish *mbnl* genes. In contrast to mouse models, zebrafish double and triple homozygous *mbnl* mutants were viable to adulthood. Zebrafish *mbnl* mutants displayed many DM- and FECD-associated alternative splicing changes in disease relevant tissues. In general, *mbnl1* and *mbnl2* mutants displayed more pronounced alternative splicing defects than *mbnl3* mutants, and compound *mbnl* mutants exhibited larger splicing changes than single *mbnl* mutants. Splicing changes in zebrafish *mbnl* mutants were greatest in adult heart and skeletal muscle. Zebrafish *mbnl* mutants displayed disease relevant physical phenotypes including decreased sized and impaired movement. The high fecundity and larval optical transparency of these new zebrafish models will make them useful for studying DM- and FECD- related phenotypes and how individual Mbnl proteins contribute to them.

Introduction

The muscleblind (MBNL) family of RNA binding proteins is highly conserved in structure and function across multicellular species, and vertebrates have three family members: MBNL1, MBNL2, and MBNL3 (Oddo et al., 2016). MBNL proteins use zinc finger motifs (two pairs in the case of vertebrates) to recognize and bind to consensus YGCY sequences (where Y is a pyrimidine) in target RNAs and regulate multiple aspects of RNA metabolism, the most well-characterized of which is alternative splicing (Ashwal-Fluss et al., 2014; Batra et al., 2014; Goers et al., 2010; Grammatikakis et al., 2011; Ho et al., 2004; Rau et al., 2011; Wang et al., 2012; Warf and Berglund, 2007). In general, MBNL proteins promote exon skipping when bound to YGCY sequences in the intron upstream of an alternative exon, and promote exon inclusion when bound downstream of an alternative exon (Du et al., 2010; Goers et al., 2010; Wang et al., 2012). Regulation of alternative splicing by MBNL proteins is critical for the molecular function of many target genes. For example, MBNL proteins promote skipping of exon 5 in the *MBNL1* and *MBNL2* mRNAs by binding to highly conserved YGCY elements in the upstream intron, affecting the subcellular localization and alternative splicing regulatory activity of the resulting proteins (Gates et al., 2011; Terenzi and Ladd, 2010; Tran et al., 2011).

The regulation of RNA metabolism by MBNL proteins is crucial for the proper functioning of many body systems, as shown by the multi-systemic phenotypes in human genetic disorders that involve *MBNL* loss of function. In myotonic dystrophy (DM), expression of an expanded CTG repeat in the 3' UTR of the *DMPK* gene (DM1) or an expanded CCTG repeat in an intron of the *CNBP* gene (DM2) leads to the formation of RNA stem loop structures that contain dozens to thousands of YGCY motifs that bind and sequester MBNL proteins, preventing them from performing their normal functions (Brook et al., 1992; Fardaei et al., 2002; Fu et al., 1992; Liquori et al., 2001; Mahadevan et al., 1992; Miller et al., 2000) (Fig. 1A). DM is most well-known for its skeletal muscle phenotypes, including weakness, myotonia, atrophy, and pain. However, individuals with DM experience a wide variety of symptoms in many tissues that can be as debilitating as muscle phenotypes, including but not limited to cataracts,

breathing difficulties, behavioral and psychological disorders, sleep disorders, insulin

resistance, heart conduction abnormalities, cardiomyopathy, and alterations in the

motility and microbiota of the gut (Bellini et al., 2006; Hilbert et al., 2017; Tarnopolsky et

al., 2010; Thomas et al., 2018; Tieleman et al., 2008; Wenninger et al., 2018).

Misregulation of specific alternative splicing events was shown to contribute to disease-

related phenotypes (Thomas et al., 2018). For example, missplicing of *RYR1* and

ATP2A1 contributed to altered calcium homeostasis in DM1 muscle (Kimura et al.,

2005; Zhao et al., 2015). Fuchs endothelial corneal dystrophy (FECD) leads to corneal

edema and loss of vision, and a subtype of this common disorder was recently shown to

be caused by an expanded CUG repeat that is expressed from an intron of the *TCF4*

gene. Signs of FECD were also reported in a subset of individuals with DM1 (Mootha et

al., 2017; Mootha et al., 2015; Wieben et al., 2017; Winkler et al., 2018). As in DM,

MBNL protein sequestration and missplicing were demonstrated in FECD, but how

MBNL loss of function contributes to the disease phenotype is not understood (Du et al.,

2015; Wieben et al., 2017).

Mouse *Mbnl* mutants greatly improved our understanding of the biological roles

of individual MBNL proteins and how they may contribute to human MBNL loss of

function disorders. Mouse *Mbnl1* is widely expressed and *Mbnl1$^{-/-}$* mice exhibited

changes in alternative splicing, myotonia, myopathy, and cataracts, but did not have

other DM-related phenotypes such as muscle weakness, suggesting that loss of *Mbnl1*

function alone was not sufficient to cause all DM-related phenotypes (Kanadia et al.,

2003; Lee et al., 2013). *Mbnl2* expression is enriched in the brain, and *Mbnl2$^{-/-}$* mice

exhibited DM-like CNS phenotypes including missplicing in the brain, abnormal sleep

patterns, and spatial memory deficits (Charizanis et al., 2012). Mouse *Mbnl3* is

expressed primarily during embryonic development and injury-induced adult skeletal

muscle regeneration. *Mbnl3$^{-/-}$* mice exhibited delays in injury-induced muscle

regeneration, accelerated onset of age-related pathologies, and gene expression

changes, but displayed minimal changes in alternative splicing, suggesting that *Mbnl3*

played a more modest role in DM-related missplicing (Choi et al., 2016; Poulos et al.,

2013). Compound *Mbnl* mutant mice had more severe phenotypes than single mutant

mice, suggesting that Mbnl proteins had overlapping functions in mice. $Mbnl1^{-/-};Mbnl2^{-/-}$

mice were embryonic lethal, while $Mbnl1^{-/-};Mbnl2^{+/-}$ mice were viable but had severe

missplicing, decreased body weight, myotonia, progressive muscle weakness, and

reduced lifespan (Lee et al., 2013). Muscle-specific double ($Mbnl1^{-/-}; Mbnl2^{C/C};$

$Mbnl3^{+/Y}; Myog-Cre^{+/-}$) and triple ($Mbnl1^{-/-}; Mbnl2^{C/C}; Mbnl3^{C/Y}; Myog-Cre^{+/-}$) mutant mice

were reduced in size, had dramatic and widespread alternative splicing changes,

muscle weakness, and most died in the neonatal period due to respiratory distress

(Thomas et al., 2017). Compound $Mbnl$ mutant mice recapitulated many of the

phenotypes associated with severe forms of DM, but their utility in future experiments is

limited because they are difficult to generate in large numbers.

We sought to expand the understanding of MBNL function using another

vertebrate model organism, the zebrafish. Like mice, zebrafish have one ortholog of

each of the human $MBNL$ genes (Liu et al., 2008). There are several advantages of

using zebrafish over mice to study Mbnl function. Zebrafish embryos develop rapidly

and externally, making studies of developmental phenotypes easier than in mice.

Zebrafish embryos are optically transparent, which allows for direct analysis of DM-

related phenotypes such as altered gut motility and abnormal heart rhythm. Finally,

zebrafish produce hundreds of embryos at once, and adults can be maintained much

more cheaply than mice, which enables experiments to be performed with large

numbers of animals and improves statistical power when studying subtle or variable

phenotypes. Two zebrafish models of $MBNL$ loss of function disorders were generated

previously, but both had severe limitations. Zebrafish embryos that were injected with

CUG repeat RNA had subtle developmental abnormalities, but did not exhibit alternative

splicing changes, and the injected RNA was degraded by two days post fertilization

(dpf) (Todd et al., 2014). This model did not allow for the study of phenotypes beyond

early development, and could not distinguish between the effects of loss of individual

Mbnl proteins. Another model employed morpholinos, but these are also transient and

can have off-target effects. Morpholino-based knockdown of $mbnl2$ in zebrafish

embryos resulted in profound morphological abnormalities that were inconsistent with

the much milder phenotypes observed in $Mbnl2^{-/-}$ mice (Charizanis et al., 2012;

Machuca-Tzili et al., 2011).

In the current study, we used CRISPR-Cas9 based mutagenesis to make

homozygous zebrafish *mbnl1*, *mbnl2,* and *mbnl3* loss of function mutants, which were

crossed to generate double (*mbnl1/mbnl2*, *mbnl2/mbnl3,* and *mbnl1/mbnl3*) and triple

(*mbnl1/mbnl2/mbnl3*) homozygous mutants, all of which were viable to adulthood. We

analyzed alternative splicing in these mutants, and found that they exhibited many of

the same changes that were present in human *MBNL* loss of function disorders and

mouse *Mbnl* mutants. These alternative splicing changes occured both in larval and

adult fish, but were most dramatic in adult skeletal and heart muscle. As in mice,

double and triple homozygous zebrafish *mbnl* mutants exhibited larger alternative

splicing changes than single mutants and had decreased body size. *mbnl* mutants also

had decreased swim test performance. Thus, zebrafish *mbnl* mutants had molecular

and physical phenotypes consistent with those present in human *MBNL* loss of function

disorders, and are a powerful new vertebrate model for studies of *MBNL* function.

Results

Generation of a comprehensive set of zebrafish *mbnl* mutants

To create stable zebrafish models of *mbnl* loss of function disorders, we used

CRISPR/Cas9 technology and mutated zebrafish *mbnl1*, *mbnl2,* and *mbnl3*. We

targeted constitutively included exons in each zebrafish *mbnl* gene (Liu et al., 2008),

and established two independent *mbnl1* mutant lines ($1^{A/A}$ and $1^{B/B}$), one *mbnl2* mutant

line ($2^{A/A}$), and three independent *mbnl3* mutant lines ($3^{A/A}$, $3^{B/B}$, $3^{C/C}$) (Fig. 1B, Table

S1, and Table S2). The *mbnl* mutations were predicted to result in a frameshift leading

to a premature termination codon (1^A, 2^A, 3^A, and 3^B), insertion of an early stop codon

(3^C), or loss of the start codon leading to usage of a downstream out-of-frame start

codon (1^B) (Fig. 1B and Table S2). Each mutation was also predicted to result in loss of

function of one or more of the zinc fingers that are important for RNA binding (Fig. 1B)

(Grammatikakis et al., 2011). Homozygous larval mutants exhibited decreased ($2^{A/A}$,

$3^{A/A}$, and $3^{B/B}$) or unchanged ($1^{A/A}$, $1^{B/B}$, and $3^{C/C}$) levels of the mRNA expressed from

the mutated gene, with minimal effect on the expression of the other *mbnl* family

members (Fig. S1).

All homozygous *mbnl1, mbnl2,* and *mbnl3* mutants survived to adulthood in

roughly Mendelian ratios (Table S3), were fertile, and did not exhibit the dramatic

morphological phenotypes that were previously observed in *mbnl2* morpholino-injected

larvae (Machuca-Tzili et al., 2011). We crossed single *mbnl* mutant fish to generate

double *mbnl* mutants ($1^{B/B};2^{A/A}$, $1^{B/B};3^{C/C}$, and $2^{A/A};3^{C/C}$). All double *mbnl* mutants were

also viable to adulthood and fertile (Table S3). This contrasted with mouse models, in

which *mbnl*$^{-/-}$;*mbnl2*$^{-/-}$ mutants were embryonic lethal, and to our knowledge represented

the first time that *mbnl1/mbnl3* and *mbnl2/mbnl3* double homozygous mutant animals

were generated (Lee et al., 2013). We further crossed double homozygous animals to

generate triple homozygous mutants ($1^{B/B};2^{A/A};3^{C/C}$). Surprisingly, these fish were viable

to adulthood, albeit in lower than expected numbers (Table S3), but they were unable to

produce embryos except through *in vitro* fertilization. In summary, we created the first

complete panel of vertebrate *mbnl1, mbnl2,* and *mbnl3* single, double, and triple

homozygous mutants for studying the functions of Mbnl proteins.

Larval zebrafish *mbnl* mutants displayed disease-associated changes in

alternative splicing

We first asked whether zebrafish *mbnl* mutants modeled the alternative splicing

changes observed in humans with DM and FECD and in mouse *Mbnl* mutants. Our

initial studies used the extensively characterized suppression of *mbnl1* exon 5 and

mbnl2 exon 5 inclusion by Mbnl proteins as a model system (Gates et al., 2011). The

intronic sequence immediately upstream of exon 5, which contains putative YGCY Mbnl

protein binding sites, is highly conserved between human and zebrafish, suggesting

functional importance (Fig. S2).

We harvested total RNA from larval zebrafish and analyzed exon inclusion

through RT-PCR analysis. Inclusion of both *mbnl1* exon 5 and *mbnl2* exon 5 was

elevated in zebrafish *mbnl* mutants (Fig. 2A,B). The effect was very modest in single

homozygous *mbnl* mutants, with only one of two *mbnl1* homozygous mutants showing

slightly increased inclusion of both *mbnl1* exon 5 and *mbnl2* exon 5, and $2^{A/A}$ mutants

displaying increased inclusion of *mbnl2* exon 5 only (Fig. 2A,B). Exon 5 inclusion was

elevated in double homozygous *mbnl* mutants above that in single homozygous

mutants, with the highest level of exon inclusion observed in $1^{B/B};2^{A/A}$ mutants (Fig.

2A,B). This suggested that zebrafish Mbnl proteins had overlapping functions in

splicing regulation, similar to findings in mice (Lee et al., 2013; Thomas et al., 2017).

mbnl3 contributed minimally to autoregulation of *mbnl* splicing in larval fish, as none of

the three *mbnl3* single mutants showed elevated *mbnl1* or *mbnl2* exon 5 inclusion, and

$1^{B/B};2^{A/A};3^{C/C}$ mutants did not show increased exon 5 inclusion compared to $1^{B/B};2^{A/A}$

mutants (Fig. 2A,B). This was consistent with mouse data, in which *mbnl3* contributed

minimally to splicing regulation (Choi et al., 2016; Poulos et al., 2013).

Adult zebrafish *mbnl* mutant skeletal muscle showed robust alternative splicing

changes

A well-characterized pattern in DM is a tendency to switch from the production of

adult-specific RNA isoforms to the production of isoforms that are normally found in

development (Kalsotra et al., 2008; Lin et al., 2006). Therefore, the magnitude of DM-

associated splicing changes tends to be larger in adult tissues than in embryos. We

asked whether this was also the case in zebrafish *mbnl* mutants, focusing on skeletal

muscle because it is one of the most severely affected tissues in DM. The mRNA levels

of *mbnl1, mbnl2*, and *mbnl3* were all higher in adult skeletal muscle than in whole larvae

(Fig. S3). The splicing patterns of *mbnl1* and *mbnl2* exon 5 changed much more

dramatically in adult *mbnl* mutant skeletal muscle than in whole larvae, from

predominant exon skipping in WT fish to predominant inclusion in $1^{B/B};2^{A/A}$ mutants (Fig.

2C,D). In mouse and DM patient tissues, some regulated exons were much more

sensitive than others to changes in overall levels of free MBNL proteins (Wagner et al.,

2016). It appears that zebrafish also followed this pattern, as in skeletal muscle *mbnl1*

mutation alone was sufficient to increase *mbnl1* exon 5 inclusion, whereas loss of

function of both *mbnl1* and *mbnl2* was required to increase *mbnl2* exon 5 inclusion (Fig.

2C,D). *mbnl3* played a small role in *mbnl* splicing autoregulation in skeletal muscle, as

$1^{B/B};2^{A/A};3^{C/C}$ mutants showed slightly elevated *mbnl2* exon 5 inclusion compared to

$1^{B/B};2^{A/A}$ mutants (Fig. 2D).

Alternative splicing changes occurred across tissues in zebrafish *mbnl* mutants

We next asked whether adult zebrafish *mbnl* mutants exhibited alternative

splicing changes in other tissues relevant to DM and FECD pathology. Zebrafish *mbnl1,*

mbnl2, and *mbnl3* mRNAs were all expressed broadly across tissues, with the highest

mbnl mRNA levels in skeletal muscle and brain (Fig. S3). Splicing patterns for both

mbnl1 exon 5 and *mbnl2* exon 5 in the heart were very similar to those in skeletal

muscle (Figs 2C,D and 3A,B). The brain and cornea exhibited *mbnl* splicing patterns

quite similar to one another, but the magnitude of splicing changes between WT and

mutants was slightly lower than those observed in muscle (Fig. 3C,D,E,F). *mbnl*

splicing changes in the intestine were small in magnitude, and appeared to be driven

primarily by *mbnl1* mutation (Fig. 3G,H). Therefore, zebrafish *mbnl* mutants

recapitulated alternative splicing changes in disease relevant tissues.

Misregulation of alternative splicing was widespread in zebrafish *mbnl* mutants

We expanded our analysis to determine whether our findings related to

misregulation of *mbnl1* exon5 and *mbnl2* exon 5 alternative splicing in *mbnl* mutant fish

applied more broadly to other disease-associated missplicing events. We investigated

five additional exons that were conserved between human and zebrafish, two of which

exhibited increased inclusion in human *MBNL* loss of function disorders and/or mouse

models (*ank3b* exon 36 and *tpd52l2b* exon 4) and three of which exhibited decreased

inclusion (*atp2a1l* exon 24*, ryr1b* exon 72*,* and *csnk1da* exon 9) (Bachinski et al., 2014;

Freyermuth et al., 2016; Goodwin et al., 2015; Hino et al., 2007; Kimura et al., 2005).

Four showed splicing changes in the same direction as that observed in human *MBNL*

loss of function disorders and/or mouse models, while one, *csnk1da*, had a splicing

change in the opposite direction (Fig. 4). We noticed similarities between regulation of

these splicing events and auto-regulation of *mbnl1* exon 5 and *mbnl2* exon 5 inclusion.

For example, in all cases $1^{B/B};2^{A/A}$ mutants showed larger changes in exon inclusion

compared to WT than either of the other two double homozygous mutants, suggesting

that *mbnl1* and *mbnl2* played the largest roles in alternative splicing regulation (Fig. 4).

mbnl3 appeared to play a lesser role in the regulation of these splicing events, as none

of the three single homozygous *mbnl3* mutants showed significant changes in inclusion

of any of the exons tested, and triple homozygous *mbnl* mutants did not show more

robust splicing changes than $1^{B/B};2^{A/A}$ mutants (Fig. 4). Consistent with the idea that

some exons were more sensitive to overall *mbnl* levels than others, some of the

alternative exons responded to loss of *mbnl1* function alone (*ank3b* exon 36, *atp2a1l*

exon 24, and *ryr1b* exon 72, Fig. 4A,B,C), while others required loss of both *mbnl1* and

mbnl2 function for a splicing change to occur (*tpd52l2b* exon 4 and *csnk1da* exon 9, Fig.

4D,E).

Mutation of <u>mbnl</u> genes led to decreased zebrafish size

Mouse compound *Mbnl* loss of function mutants, but not single mutants, were

significantly smaller than WT mice (Lee et al., 2013; Thomas et al., 2017). To determine

whether loss of *mbnl* function had a similar effect in zebrafish, we raised WT, double

homozygous *mbnl* mutant, and triple homozygous *mbnl* mutant fish together in the

same tank and measured their lengths in early adulthood. WT fish were significantly

larger than $1^{B/B};3^{C/C}$ fish, which in turn were significantly larger than $1^{B/B};2^{A/A}$ fish (Fig.

5A). $2^{A/A};3^{C/C}$ fish were slightly smaller than WT fish, but the difference was not

statistically significant (Fig. 5A). Triple homozygous *mbnl* mutant fish were similar in

size to $1^{B/B};2^{A/A}$ fish (Fig. 5A). Interestingly, mutation of *mbnl1*, *mbnl2,* or *mbnl3* alone

was not sufficient to decrease adult zebrafish size (Fig. S4A,B,C). Taken together,

these results suggested that loss of function of more than one Mbnl protein was

required to reduce fish size, and that *mbnl1* had the strongest effect on fish size, *mbnl2*

played a smaller role, and *mbnl3* contributed minimally to adult fish size.

To assess whether zebrafish *mbnl* mutants had a developmental size phenotype,

we measured 7 dpf larval fish whose nutrition was derived from the yolk rather than

feeding. All double homozygous *mbnl* mutant larvae were similar in size to WT, except

for $2^{A/A};3^{C/C}$ larvae, which were slightly smaller than WT (Fig. 5B). $1^{B/B};2^{A/A};3^{C/C}$ larvae

were similar in size to their $1^{+/B};2^{A/A};3^{C/C}$ and $2^{A/A};3^{C/C}$ clutch mates, and all single

homozygous *mbnl* mutant larvae were either similar in size or slightly larger than WT

(Figs 5C and S4D). These results suggest that the reduced size phenotype in zebrafish

mbnl mutants arose later in development, after feeding began.

Histological analysis of adult skeletal muscle was performed by an expert

zebrafish pathologist to determine whether the fish genotypes with the most profound

size and splicing phenotypes, $1^{B/B};2^{A/A}$ and $1^{B/B};2^{A/A};3^{C/C}$, also had changes in adult

muscle fiber morphology. We did not observe obvious signs of DM-related muscle

histopathology in the *mbnl* mutant fish (Fig S5), such as myofiber atrophy or centralized

nuclei as has been described in compound *mbnl* mutant mice and in individuals with DM

(Thomas et al., 2018; Thomas et al., 2017), but we cannot rule out the possibility that

the mutant fish had subtle alterations in muscle structure or function.

Zebrafish *mbnl* mutants had altered movement

Given that individuals with DM have impaired motor function, we asked whether

zebrafish *mbnl* mutants exhibited altered swimming behavior. We performed a test in

which individual adult fish were introduced to a novel tank and their position within the

tank was tracked for five minutes (Fig. 6A,B). A similar test was previously performed in

DM patients, in which the distance walked in six minutes was used as a measure of

motor function (Kierkegaard and Tollback, 2007; Park et al., 2018). While all single

homozygous mutants swam equal or greater distance than WT, two of the double

homozygous *mbnl* mutants ($1^{B/B};2^{A/A}$ and $1^{B/B};3^{C/C}$) and the $1^{B/B};2^{A/A};3^{C/C}$ mutants swam

significantly decreased distances compared to WT (Fig. 6C). One explanation for this

result is impaired motor function, although this phenotype needs to be explored further.

While performing these tests, we noticed a striking difference in the amount of time

mbnl mutants and WT fish spent near the bottom of the tank. We quantified this by

measuring the distance from the bottom of the tank in each frame for each fish and

averaging this over the recording period. Except for $2^{A/A};3^{C/C}$ mutants, all *mbnl* mutants

spent significantly more time toward the bottom of the tank than WT fish (Fig. 6D).

Avoiding the top of the tank is a well-known indicator of anxiety in zebrafish, and future

tests will explore whether *mbnl* mutants exhibit other signs of anxiety or perhaps have a

morphological or motor deficit that explains their lower swimming position (Cachat et al.,

2010). Overall, these results indicated that loss of *mbnl* function resulted in altered

movement in fish, but the exact mechanisms underlying these changes are unknown.

Discussion

For the first time, we generated a complete panel of vertebrate *mbnl* single,

double, and triple homozygous loss of function mutants. These mutants will be

invaluable for investigating the functions of vertebrate Mbnl proteins both individually

and in concert on an organismal level. Existing *Mbnl* mutant mice have provided

important insights into the biological functions of individual Mbnl proteins (Charizanis et

al., 2012; Choi et al., 2016; Kanadia et al., 2003; Lee et al., 2013; Poulos et al., 2013;

Thomas et al., 2017). However, the compound *Mbnl* mutant mouse models that most

closely mimic DM phenotypes are challenging to breed in large numbers and most die

during the neonatal period, making studies of adult phenotypes difficult. In contrast, all

double and triple homozygous *mbnl* mutant zebrafish are viable to adulthood, and fertile

double homozygous mutants can be incrossed to generate hundreds of mutants for

experiments. Even triple homozygous *mbnl* mutants, which do not mate on their own,

can be generated in large numbers through *in vitro* fertilization or through incrosses of

$1^{+/B};2^{A/A};3^{C/C}$ fish. The ability to study hundreds of zebrafish mutants at once provides

the statistical power to analyze subtle or variable phenotypes.

We propose a few explanations for why zebrafish $1^{B/B};2^{A/A}$ and $1^{B/B};2^{A/A};3^{C/C}$

mutants were viable to adulthood. Most muscle-specific double and triple *Mbnl* mutant

mice died during the neonatal period when they transitioned abruptly between receiving

oxygen through the placenta and breathing on their own. The small fraction of muscle-

specific compound *Mbnl* mutant mice that were able to make it through this transition

survived to adulthood (Thomas et al., 2018). In contrast, larval fish underwent a much

more gradual transition between receiving oxygen through diffusion to receiving oxygen

through the gills, and thus may have been better able to overcome this developmental

hurdle. Another possibility is that one or more of the zebrafish *mbnl* CRISPR mutations

did not result in a complete loss of function. In particular, $2^{A/A}$ mutants had the potential

to produce a partially functional protein containing one of the two sets of zinc finger

motifs (Fig. 1B), although we are confident that $2^{A/A}$ mutants represented at least a

partial loss of function given their splicing phenotypes and reduced *mbnl2* mRNA levels

(Figs 2,3,S1). Larvae injected with *mbnl2* morpholinos were reported to have dramatic

morphological phenotypes (Machuca-Tzili et al., 2011), although we were unable to

recapitulate these results (data not shown). The fact that our $2^{A/A}$ fish did not exhibit

similar phenotypes could be explained by partially retained Mbnl2 function in $2^{A/A}$ fish.

We suspect, however, that off target effects of the *Mbnl2* morpholinos contributed to the

observed morphological defects, given the mild phenotypes of $Mbnl2^{-/-}$ mice(Charizanis

et al., 2012).

Zebrafish *mbnl* mutants exhibited many of the same alternative splicing changes

that were observed in DM, FECD, and mouse DM models, suggesting that the splicing

targets of Mbnl proteins were conserved across species (Figs 2,3,4) (Bachinski et al.,

2014; Freyermuth et al., 2016; Goodwin et al., 2015; Hino et al., 2007; Kimura et al.,

2005). Similar to previous findings in mice, *mbnl1* and *mbnl2* played a larger role in

zebrafish splicing regulation than *mbnl3* for the exons tested (Figs 2,3,4) (Choi et al.,

2016; Poulos et al., 2013). Also consistent with mouse studies was the finding that

splicing changes tended to be larger in double homozygous zebrafish mutants than in

single homozygous mutants, suggesting partially overlapping functions of mbnl proteins

(Figs 2,3,4) (Lee et al., 2013; Thomas et al., 2017). Splicing changes in zebrafish *mbnl*

mutants occured in both larvae and adults, and across many DM-relevant tissues

(skeletal muscle, heart, brain, and intestine) as well as in the cornea, the tissue affected

in FECD.

Our data indicated that, just like other model organisms, zebrafish Mbnl-mediated

alternative splicing regulation was quite complex. Mbnl proteins suppressed inclusion of

some exons (*mbnl1* exon 5, *mbnl2* exon 5, *ank3b* exon 36, *tpd52l2b* exon 4, and

csnk1da exon 9), while they promoted inclusion of others (*ryr1b* exon 72 and *atp2a1l*

exon 24) (Figs 2,4). Similar to observations in other systems, some zebrafish exons

were more sensitive to changes in overall *mbnl* concentration than others (Wagner et

al., 2016). For example, in skeletal muscle, *mbnl2* exon 5 required the loss of both

mbnl1 and *mbnl2* function for increased inclusion, whereas *mbnl1* exon 5 only required

the loss of *mbnl1* function (Fig. 2C,D).

An immediate future goal of ours is to perform a genome wide analysis of RNA

processing and expression changes in our complete panel of zebrafish *mbnl* mutants.

This will allow us to gain an understanding of how each Mbnl protein contributes to RNA

processing on a global level. It will identify candidate RNA processing events to

investigate further for relevance to disease phenotypes. A global analysis will also

enable us to determine the extent to which changes in RNA processing overlap

between zebrafish *mbnl* mutants and human patients with *MBNL* loss of function

disorders. While we demonstrated that the misregulation of several disease-relevant

alternative exons was conserved between human DM and FECD patients and zebrafish

models, not all missplicing events were conserved between species. For example,

zebrafish do not appear to have an ortholog of the *CLCN1* exon whose misregulation

contributed to myotonia in DM, so the zebrafish is unlikely to be an appropriate model

for studying myotonia, although it can be used to study many other important disease

phenotypes (Lueck et al., 2007a; Lueck et al., 2007b).

In addition to their molecular phenotypes, zebrafish *mbnl* mutants had physical

phenotypes that were relevant to human disease and mouse models. Adult $1^{B/B};2^{A/A}$,

$1^{B/B};3^{C/C}$, and $1^{B/B};2^{A/A};3^{C/C}$ fish were significantly smaller than their WT tank mates (Fig.

5A). Neonatal muscle-specific compound mutant mice likewise were substantially

smaller than their littermates (Thomas et al., 2017). In the case of fish, the size

phenotype did not arise in early in development, as 7 dpf larval *mbnl* mutant fish were

not smaller than WT fish (Figs 5B,C and S4D). We do not know the mechanism

underlying the size phenotype in *mbnl* mutant fish, but one possibility is impaired

swimming ability, leading to decreased ability to compete with tank mates for access to

food. Consistent with this idea, we found that $1^{B/B};2^{A/A}$, $1^{B/B};3^{C/C}$, and $1^{B/B};2^{A/A};3^{C/C}$ fish

swam a significantly decreased distance compared to WT fish during a five-minute swim

test (Fig. 6B,C). Interestingly, human DM patients exhibited decreased performance on

a similar 6-minute walk test due to muscle weakness and impaired motor function,

leading us to hypothesize that *mbnl* mutant fish also had muscle weakness

(Kierkegaard and Tollback, 2007; Park et al., 2018). Although basic histological

analysis did not indicate obvious signs of muscle fiber pathology in *mbnl* mutant fish

(Fig. S5), more in-depth analyses are needed to determine whether subtle changes in

muscle structure or function are present that could lead to muscle weakness. We also

found that all genotypes of *mbnl* mutant fish stayed closer to the bottom of the tank on

average than WT fish (Fig. 6B,D), which could also decrease access to food as fish

were fed from the top of the tank. The *mbnl* mutant fish could have stayed at the

bottom of the tank due to motor dysfunction or a morphological defect, but this

phenotype is also a well-characterized indicator of anxiety in fish (Cachat et al., 2010).

Individuals with DM often show signs of avoidant personality disorder with anxiety, and

an intriguing possibility is that *mbnl* mutant fish also have increased anxiety (Delaporte,

1998; Minier et al., 2018). Our future studies will explore in depth the mechanisms

underlying decreased size and altered swimming behavior in *mbnl* mutant fish.

In summary, we successfully generated a complete panel of zebrafish single,

double, and triple homozygous *mbnl* mutants for modeling human *mbnl* loss of function

disorders. These fish exhibited molecular and physical phenotypes similar to those

observed in humans with FECD and DM. The new models will be highly useful in future

studies of the functions of individual Mbnl proteins and how they contribute to disease

relevant phenotypes. Strengths of using zebrafish over other vertebrate model systems

are that large numbers of fish can be generated to study subtle or variable phenotypes,

and the fact that transparent larvae can be used to directly study disease-related

phenotypes, such as altered gut motility and heart abnormalities, in live animals.

Materials and Methods

<u>Generation of mutant zebrafish</u>

All zebrafish experiments were performed with the guidance and approval of the

University of Oregon Institutional Animal Care and Use Committee. Guide RNAs

(gRNAs) targeting zebrafish (*Danio rerio*) *mbnl1*, *mbnl2*, and *mbnl3* were designed

using the Chop Chop website (http://chopchop.cbu.uib.no). DNA templates for the

gRNAs were generated by a template-free Phusion polymerase (New England Biolabs)

PCR reaction using a common scaffold primer (gRNA scaffold, Table S1) and a gene-

specific primer (*mbnl1* gRNA1, *mbnl1* gRNA2, *mbnl2* gRNA, *mbnl3* gRNA1, or *mbnl3*

gRNA2, Table S1), then cleaned using the QIAquick PCR Purification Kit (Qiagen).

gRNAs were transcribed from DNA templates using a MEGAscript kit (Ambion) and

purified by phenol-chloroform extraction and isopropanol precipitation. Cas9 RNA was

generated by linearizing the pT3TS-nls-zCas9-nls plasmid (Jao et al., 2013) with XbaI,

purifying it using the QIAquick Gel Extraction Kit (Qiagen), performing an in vitro

transcription reaction using the T3 mMESSAGE kit (Invitrogen), and purifying the RNA

using the RNeasy Mini kit (Qiagen). AB zebrafish embryos were microinjected at the

one cell stage with 1-2 nL of a mixture containing 100ng/μL Cas9 mRNA, 50 ng/μL

gRNA, and phenol red, and raised to adulthood. Mosaic mutants were identified by

PCR amplification and Sanger sequencing of fin clip DNA using primers specific to the

targeted region (Tables S1,S2), and outcrossed to wildtype AB zebrafish to generate

heterozygotes. Fish with predicted loss of function mutations were identified by Sanger

sequencing (Table S2), and further crossed to generate single, double, and triple *mbnl1*,

mbnl2, and *mbnl3* homozygotes. Fish were genotyped by restriction fragment length

polymorphism analysis using the primers and restriction enzymes indicated in Table S2.

Sperm from all zebrafish mutant lines were cryopreserved and are available upon

request from the corresponding author.

<u>Measurement of zebrafish size</u>

Young adult fish (2-4 months old) (Figs 5A,S4A,B,C) were anesthetized in 168

mg/L tricaine methane sulfonate, photographed against a white background with a ruler

using a tablet computer, finclipped, and genotyped. To ensure identical density and

feeding conditions, fish were compared with others from the same clutch (Fig. S4A,B,C)

or different clutches that had been raised together in the same tank (Fig. 5A). For

measurement of larval fish, unfed embryos of different genotypes were raised to 7 days

post fertilization (dpf) in separate dishes at a density of one fish per mL of embryo

medium, with the exception of the larvae in Fig. 5C, which were grown in the same dish

and genotyped after measurement. Fish were anesthetized in 168 mg/mL tricaine

methane sulfonate, laid out on a microscope slide in 3% methylcellulose, and imaged

using a Leica M165FC microscope. An investigator that was blinded to genotype

measured the distance from snout to caudal peduncle or posterior tip of the notochord

(standard length, SL) of each fish (Parichy et al., 2009). Data were analyzed by

ordinary one-way ANOVA with multiple comparison correction.

<u>Alternative splicing analysis</u>

Fish were euthanized by tricaine methane sulfonate overdose (larvae) or

hypothermic shock (adults). Tissues of interest were dissected and flash frozen in 1 mL

of Trizol (Ambion), thawed, and homogenized with a mortar and pestle (larvae) or Bullet

Blender Storm 24 (adult tissues). Chloroform (200 μL) was added to each tube followed

by mixing, centrifugation at 12,000 g for 10 minutes at 4°C, transfer of the aqueous

phase to a separate tube, addition of 200 μL ethanol, and binding of sample to an

RNeasy mini kit column (Qiagen). RNA was washed and eluted according to the

manufacturer's instructions and the concentration was measured using a NanoDrop

2000 (Thermo Scientific).

RNA (20-200 ng) was reverse transcribed with Superscript II Reverse

Transcriptase (Invitrogen) according to the manufacturer's instructions using gene-

specific reverse (R) primers located in the exon downstream of the regulated exon of

interest (Table S1). The resulting cDNA was amplified by 28-31 cycles of PCR using

Taq polymerase and the forward (F) and reverse (R) primers indicated in Table S1.

Samples were separated by electrophoresis on a 6% *bis*-Acrylamide (19:1) gel that was

stained overnight with 1X SYBR green I nucleic acid gel stain (Invitrogen).   The gel was

imaged and quantified using an AlphaImagerHP (Alpha Innotech). The background-

corrected sum of each band was measured and the percent exon inclusion was

calculated using the following formula: ((exon included sum) /(exon included sum +

exon excluded sum)) X 100. Data were analyzed by ordinary one-way ANOVA with

multiple comparison correction.

RT-qPCR

Total RNA was prepared from 5 dpf larvae and adult tissues using the same

procedure as for alternative splicing analysis. RNA was treated with TURBO DNase

(Ambion) according to manufacturer's instructions then reverse transcribed with an

oligo(dT)$_{20}$ primer using the Superscript III cDNA First Strand Synthesis kit (Invitrogen).

The qPCR reaction was set up using the KAPA SYBR FAST ABI Prism kit (KAPA

Biosystems) according to manufacturer's instructions and run on a Quant Studio 3

System (ThermoFisher) using the default settings for SYBR Green reagents and the

fast run mode in the QuantStudio Design & Analysis Software v.1.4.2. The comparative

C_T ($\Delta\Delta C_T$) method was used to calculate relative mRNA levels. Five biological samples

of each genotype or tissue type were run in triplicate and data were normalized to the

expression of the housekeeping gene *eef1a1l1*(Vanhauwaert et al., 2014). Primers

used for RT-qPCR are shown in Table S1.

<u>Zebrafish behavior assays</u>

Age-matched single adult zebrafish were placed into a novel environment

consisting of a custom glass aquarium measuring 18 cm deep X 25.4 cm long X 7.6 cm

wide (Fig. 6A). The fish were monitored with a Logitech camera during a 5 minute time

frame to characterize basic exploratory swimming behavior. The raw tracking data was

analyzed with custom software (DaniOPEN, https://github.com/stednitzs/daniopen) to

measure swim distance and distance from the bottom of the tank(Stednitz et al., 2018).

Following the test period, the fish were returned to their home tanks.

<u>Histology</u>

Ten-month-old WT and mutant adult zebrafish were euthanized by hypothermic

shock and fixed in Bouin's solution (Sigma), then washed in 70% ethanol. Fixed fish

were bisected parasagitally or transversely, processed for paraffin embedding,

sectioned at 7 μm thickness, and stained with hematoxylin and eosin by a histologist

from the University of Oregon Institute of Neuroscience. Muscle morphology was

analyzed by a zebrafish pathologist from the Zebrafish International Resource Center

who was blinded to genotype. Representative DIC images of muscle transverse

sections were taken using a Leica DMLB microscope.

Acknowledgements

We thank the staff of the University of Oregon Zebrafish Facility for maintaining fish and

providing expertise in performing zebrafish procedures, and Dr. Poh Kheng Loi for

histological services. We thank Ellie Melancon, Camila Morales Fenero, Michelle

Massaquoi, and Sam Peterson for advice in performing experiments and data analysis,

and members of the Guillemin, Berglund, and Eisen labs for helpful discussions.

Competing Interests

No competing interests declared.

Funding

Research reported in this publication was supported by the National Institute of General

Medical Sciences of the National Institutes of Health under award numbers

1P50GM098911 and 1P01GM125576 to K.G. and 5R01GM121862 to J.A.B., and by

the National Institute of Diabetes and Digestive and Kidney Diseases of the National

Institutes of Health under award number F32DK107318 to M.N.H. It was also

supported by grants from the Muscular Dystrophy Association (516314 to J.A.B.) and

the Myotonic Dystrophy/Wyck Foundation (Wyck-FF-2014-0013 to M.N.H.). The

content is solely the responsibility of the authors and does not necessarily represent the

official views of the National Institutes of Health, the Muscular Dystrophy Association, or

the Myotonic Dystrophy/Wyck Foundation.

References

Ashwal-Fluss, R., Meyer, M., Pamudurti, N. R., Ivanov, A., Bartok, O., Hanan, M., Evantal, N., Memczak, S., Rajewsky, N. and Kadener, S. (2014). circRNA biogenesis competes with pre-mRNA splicing. *Mol Cell* **56**, 55-66.

Bachinski, L. L., Baggerly, K. A., Neubauer, V. L., Nixon, T. J., Raheem, O., Sirito, M., Unruh, A. K., Zhang, J., Nagarajan, L., Timchenko, L. T. et al. (2014). Most expression and splicing changes in myotonic dystrophy type 1 and type 2 skeletal muscle are shared with other muscular dystrophies. *Neuromuscul Disord* **24**, 227-40.

Batra, R., Charizanis, K., Manchanda, M., Mohan, A., Li, M., Finn, D. J., Goodwin, M., Zhang, C., Sobczak, K., Thornton, C. A. et al. (2014). Loss of MBNL leads to disruption of developmentally regulated alternative polyadenylation in RNA-mediated disease. *Mol Cell* **56**, 311-322.

Bellini, M., Biagi, S., Stasi, C., Costa, F., Mumolo, M. G., Ricchiuti, A. and Marchi, S. (2006). Gastrointestinal manifestations in myotonic muscular dystrophy. *World J Gastroenterol* **12**, 1821-8.

Brook, J. D., McCurrach, M. E., Harley, H. G., Buckler, A. J., Church, D., Aburatani, H., Hunter, K., Stanton, V. P., Thirion, J. P., Hudson, T. et al. (1992). Molecular basis of myotonic dystrophy: expansion of a trinucleotide (CTG) repeat at the 3' end of a transcript encoding a protein kinase family member. *Cell* **68**, 799-808.

Cachat, J., Stewart, A., Grossman, L., Gaikwad, S., Kadri, F., Chung, K. M., Wu, N., Wong, K., Roy, S., Suciu, C. et al. (2010). Measuring behavioral and endocrine responses to novelty stress in adult zebrafish. *Nat Protoc* **5**, 1786-99.

Charizanis, K., Lee, K. Y., Batra, R., Goodwin, M., Zhang, C., Yuan, Y., Shiue, L., Cline, M., Scotti, M. M., Xia, G. et al. (2012). Muscleblind-like 2-mediated alternative splicing in the developing brain and dysregulation in myotonic dystrophy. *Neuron* **75**, 437-50.

Choi, J., Dixon, D. M., Dansithong, W., Abdallah, W. F., Roos, K. P., Jordan, M. C., Trac, B., Lee, H. S., Comai, L. and Reddy, S. (2016). Muscleblind-like 3 deficit results in a spectrum of age-associated pathologies observed in myotonic dystrophy. *Sci Rep* **6**, 30999.

Delaporte, C. (1998). Personality patterns in patients with myotonic dystrophy. *Arch Neurol* **55**, 635-40.

Du, H., Cline, M. S., Osborne, R. J., Tuttle, D. L., Clark, T. A., Donohue, J. P., Hall, M. P., Shiue, L., Swanson, M. S., Thornton, C. A. et al. (2010). Aberrant alternative splicing and extracellular matrix gene expression in mouse models of myotonic dystrophy. *Nat Struct Mol Biol* **17**, 187-93.

Du, J., Aleff, R. A., Soragni, E., Kalari, K., Nie, J., Tang, X., Davila, J., Kocher, J. P., Patel, S. V., Gottesfeld, J. M. et al. (2015). RNA toxicity and missplicing in the common eye disease fuchs endothelial corneal dystrophy. *J Biol Chem* **290**, 5979-90.

Fardaei, M., Rogers, M. T., Thorpe, H. M., Larkin, K., Hamshere, M. G., Harper, P. S. and Brook, J. D. (2002). Three proteins, MBNL, MBLL and MBXL, co-localize in vivo with nuclear foci of expanded-repeat transcripts in DM1 and DM2 cells. *Hum Mol Genet* **11**, 805-14.

Freyermuth, F., Rau, F., Kokunai, Y., Linke, T., Sellier, C., Nakamori, M.,
Kino, Y., Arandel, L., Jollet, A., Thibault, C. et al. (2016). Splicing misregulation of
SCN5A contributes to cardiac-conduction delay and heart arrhythmia in myotonic
dystrophy. *Nat Commun* **7**, 11067.
Fu, Y. H., Pizzuti, A., Fenwick, R. G., Jr., King, J., Rajnarayan, S., Dunne, P.
W., Dubel, J., Nasser, G. A., Ashizawa, T., de Jong, P. et al. (1992). An unstable
triplet repeat in a gene related to myotonic muscular dystrophy. *Science* **255**, 1256-8.
Gates, D. P., Coonrod, L. A. and Berglund, J. A. (2011). Autoregulated splicing
of muscleblind-like 1 (MBNL1) Pre-mRNA. *J Biol Chem* **286**, 34224-33.
Goers, E. S., Purcell, J., Voelker, R. B., Gates, D. P. and Berglund, J. A.
(2010). MBNL1 binds GC motifs embedded in pyrimidines to regulate alternative
splicing. *Nucleic Acids Res* **38**, 2467-84.
Goodwin, M., Mohan, A., Batra, R., Lee, K. Y., Charizanis, K., Fernandez
Gomez, F. J., Eddarkaoui, S., Sergeant, N., Buee, L., Kimura, T. et al. (2015). MBNL
Sequestration by Toxic RNAs and RNA Misprocessing in the Myotonic Dystrophy Brain.
Cell Rep **12**, 1159-68.
Grammatikakis, I., Goo, Y. H., Echeverria, G. V. and Cooper, T. A. (2011).
Identification of MBNL1 and MBNL3 domains required for splicing activation and
repression. *Nucleic Acids Res* **39**, 2769-80.
Hilbert, J. E., Barohn, R. J., Clemens, P. R., Luebbe, E. A., Martens, W. B.,
McDermott, M. P., Parkhill, A. L., Tawil, R., Thornton, C. A., Moxley, R. T., 3rd et al.
(2017). High frequency of gastrointestinal manifestations in myotonic dystrophy type 1
and type 2. *Neurology* **89**, 1348-1354.
Hino, S., Kondo, S., Sekiya, H., Saito, A., Kanemoto, S., Murakami, T.,
Chihara, K., Aoki, Y., Nakamori, M., Takahashi, M. P. et al. (2007). Molecular
mechanisms responsible for aberrant splicing of SERCA1 in myotonic dystrophy type 1.
Hum Mol Genet **16**, 2834-43.
Ho, T. H., Charlet, B. N., Poulos, M. G., Singh, G., Swanson, M. S. and
Cooper, T. A. (2004). Muscleblind proteins regulate alternative splicing. *EMBO J* **23**,
3103-12.
Jao, L. E., Wente, S. R. and Chen, W. (2013). Efficient multiplex biallelic
zebrafish genome editing using a CRISPR nuclease system. *Proc Natl Acad Sci U S A*
110, 13904-9.
Kalsotra, A., Xiao, X., Ward, A. J., Castle, J. C., Johnson, J. M., Burge, C. B.
and Cooper, T. A. (2008). A postnatal switch of CELF and MBNL proteins reprograms
alternative splicing in the developing heart. *Proc Natl Acad Sci U S A* **105**, 20333-8.
Kanadia, R. N., Johnstone, K. A., Mankodi, A., Lungu, C., Thornton, C. A.,
Esson, D., Timmers, A. M., Hauswirth, W. W. and Swanson, M. S. (2003). A
muscleblind knockout model for myotonic dystrophy. *Science* **302**, 1978-80.
Kierkegaard, M. and Tollback, A. (2007). Reliability and feasibility of the six
minute walk test in subjects with myotonic dystrophy. *Neuromuscul Disord* **17**, 943-9.
Kimura, T., Nakamori, M., Lueck, J. D., Pouliquin, P., Aoike, F., Fujimura, H.,
Dirksen, R. T., Takahashi, M. P., Dulhunty, A. F. and Sakoda, S. (2005). Altered
mRNA splicing of the skeletal muscle ryanodine receptor and

sarcoplasmic/endoplasmic reticulum Ca2+-ATPase in myotonic dystrophy type 1. *Hum*
Mol Genet **14**, 2189-200.

Lee, K. Y., Li, M., Manchanda, M., Batra, R., Charizanis, K., Mohan, A.,
Warren, S. A., Chamberlain, C. M., Finn, D., Hong, H. et al. (2013). Compound loss
of muscleblind-like function in myotonic dystrophy. *EMBO Mol Med* **5**, 1887-900.

Lin, X., Miller, J. W., Mankodi, A., Kanadia, R. N., Yuan, Y., Moxley, R. T.,
Swanson, M. S. and Thornton, C. A. (2006). Failure of MBNL1-dependent post-natal
splicing transitions in myotonic dystrophy. *Hum Mol Genet* **15**, 2087-97.

Liquori, C. L., Ricker, K., Moseley, M. L., Jacobsen, J. F., Kress, W., Naylor,
S. L., Day, J. W. and Ranum, L. P. (2001). Myotonic dystrophy type 2 caused by a
CCTG expansion in intron 1 of ZNF9. *Science* **293**, 864-7.

Liu, Y. F., Liu, H. Y., Tu, L. C., Lin, C. W., Hsiao, K. M. and Pan, H. (2008).
Zebrafish muscleblind-like genes: identification, structural features and expression.
Comp Biochem Physiol B Biochem Mol Biol **151**, 118-24.

Lueck, J. D., Lungu, C., Mankodi, A., Osborne, R. J., Welle, S. L., Dirksen, R.
T. and Thornton, C. A. (2007a). Chloride channelopathy in myotonic dystrophy
resulting from loss of posttranscriptional regulation for CLCN1. *Am J Physiol Cell*
Physiol **292**, C1291-7.

Lueck, J. D., Mankodi, A., Swanson, M. S., Thornton, C. A. and Dirksen, R.
T. (2007b). Muscle chloride channel dysfunction in two mouse models of myotonic
dystrophy. *J Gen Physiol* **129**, 79-94.

Machuca-Tzili, L. E., Buxton, S., Thorpe, A., Timson, C. M., Wigmore, P.,
Luther, P. K. and Brook, J. D. (2011). Zebrafish deficient for Muscleblind-like 2 exhibit
features of myotonic dystrophy. *Dis Model Mech* **4**, 381-92.

Mahadevan, M., Tsilfidis, C., Sabourin, L., Shutler, G., Amemiya, C., Jansen,
G., Neville, C., Narang, M., Barcelo, J., O'Hoy, K. et al. (1992). Myotonic dystrophy
mutation: an unstable CTG repeat in the 3' untranslated region of the gene. *Science*
255, 1253-5.

Miller, J. W., Urbinati, C. R., Teng-Umnuay, P., Stenberg, M. G., Byrne, B. J.,
Thornton, C. A. and Swanson, M. S. (2000). Recruitment of human muscleblind
proteins to (CUG)(n) expansions associated with myotonic dystrophy. *EMBO J* **19**,
4439-48.

Minier, L., Lignier, B., Bouvet, C., Gallais, B. and Camart, N. (2018). A
Review of Psychopathology Features, Personality, and Coping in Myotonic Dystrophy
Type 1. *J Neuromuscul Dis* **5**, 279-294.

Mootha, V. V., Hansen, B., Rong, Z., Mammen, P. P., Zhou, Z., Xing, C. and
Gong, X. (2017). Fuchs' Endothelial Corneal Dystrophy and RNA Foci in Patients With
Myotonic Dystrophy. *Invest Ophthalmol Vis Sci* **58**, 4579-4585.

Mootha, V. V., Hussain, I., Cunnusamy, K., Graham, E., Gong, X., Neelam,
S., Xing, C., Kittler, R. and Petroll, W. M. (2015). TCF4 Triplet Repeat Expansion and
Nuclear RNA Foci in Fuchs' Endothelial Corneal Dystrophy. *Invest Ophthalmol Vis Sci*
56, 2003-11.

Oddo, J. C., Saxena, T., McConnell, O. L., Berglund, J. A. and Wang, E. T. (2016). Conservation of context-dependent splicing activity in distant Muscleblind homologs. *Nucleic Acids Res* **44**, 8352-62.

Parichy, D. M., Elizondo, M. R., Mills, M. G., Gordon, T. N. and Engeszer, R. E. (2009). Normal table of postembryonic zebrafish development: staging by externally visible anatomy of the living fish. *Dev Dyn* **238**, 2975-3015.

Park, D., Lee, S. H., Shin, J. H. and Park, J. S. (2018). Lower limb muscle magnetic resonance imaging in myotonic dystrophy type 1 correlates with the six-minute walk test and CTG repeats. *Neuromuscul Disord* **28**, 29-37.

Poulos, M. G., Batra, R., Li, M., Yuan, Y., Zhang, C., Darnell, R. B. and Swanson, M. S. (2013). Progressive impairment of muscle regeneration in muscleblind-like 3 isoform knockout mice. *Hum Mol Genet* **22**, 3547-58.

Rau, F., Freyermuth, F., Fugier, C., Villemin, J. P., Fischer, M. C., Jost, B., Dembele, D., Gourdon, G., Nicole, A., Duboc, D. et al. (2011). Misregulation of miR-1 processing is associated with heart defects in myotonic dystrophy. *Nat Struct Mol Biol* **18**, 840-5.

Stednitz, S. J., McDermott, E. M., Ncube, D., Tallafuss, A., Eisen, J. S. and Washbourne, P. (2018). Forebrain Control of Behaviorally Driven Social Orienting in Zebrafish. *Curr Biol* **28**, 2445-2451 e3.

Tarnopolsky, M. A., Pearce, E., Matteliano, A., James, C. and Armstrong, D. (2010). Bacterial overgrowth syndrome in myotonic muscular dystrophy is potentially treatable. *Muscle Nerve* **42**, 853-5.

Terenzi, F. and Ladd, A. N. (2010). Conserved developmental alternative splicing of muscleblind-like (MBNL) transcripts regulates MBNL localization and activity. *RNA Biol* **7**, 43-55.

Thomas, J. D., Oliveira, R., Sznajder, L. J. and Swanson, M. S. (2018). Myotonic Dystrophy and Developmental Regulation of RNA Processing. *Compr Physiol* **8**, 509-553.

Thomas, J. D., Sznajder, L. J., Bardhi, O., Aslam, F. N., Anastasiadis, Z. P., Scotti, M. M., Nishino, I., Nakamori, M., Wang, E. T. and Swanson, M. S. (2017). Disrupted prenatal RNA processing and myogenesis in congenital myotonic dystrophy. *Genes Dev* **31**, 1122-1133.

Tieleman, A. A., van Vliet, J., Jansen, J. B., van der Kooi, A. J., Borm, G. F. and van Engelen, B. G. (2008). Gastrointestinal involvement is frequent in Myotonic Dystrophy type 2. *Neuromuscul Disord* **18**, 646-9.

Todd, P. K., Ackall, F. Y., Hur, J., Sharma, K., Paulson, H. L. and Dowling, J. J. (2014). Transcriptional changes and developmental abnormalities in a zebrafish model of myotonic dystrophy type 1. *Dis Model Mech* **7**, 143-55.

Tran, H., Gourrier, N., Lemercier-Neuillet, C., Dhaenens, C. M., Vautrin, A., Fernandez-Gomez, F. J., Arandel, L., Carpentier, C., Obriot, H., Eddarkaoui, S. et al. (2011). Analysis of exonic regions involved in nuclear localization, splicing activity, and dimerization of Muscleblind-like-1 isoforms. *J Biol Chem* **286**, 16435-46.

Vanhauwaert, S., Van Peer, G., Rihani, A., Janssens, E., Rondou, P., Lefever, S., De Paepe, A., Coucke, P. J., Speleman, F., Vandesompele, J. et al.

(2014). Expressed repeat elements improve RT-qPCR normalization across a wide
range of zebrafish gene expression studies. *PLoS One* **9**, e109091.
Wagner, S. D., Struck, A. J., Gupta, R., Farnsworth, D. R., Mahady, A. E.,
Eichinger, K., Thornton, C. A., Wang, E. T. and Berglund, J. A. (2016). Dose-
Dependent Regulation of Alternative Splicing by MBNL Proteins Reveals Biomarkers for
Myotonic Dystrophy. *PLoS Genet* **12**, e1006316.
Wang, E. T., Cody, N. A., Jog, S., Biancolella, M., Wang, T. T., Treacy, D. J.,
Luo, S., Schroth, G. P., Housman, D. E., Reddy, S. et al. (2012). Transcriptome-wide
regulation of pre-mRNA splicing and mRNA localization by muscleblind proteins. *Cell*
150, 710-24.
Warf, M. B. and Berglund, J. A. (2007). MBNL binds similar RNA structures in
the CUG repeats of myotonic dystrophy and its pre-mRNA substrate cardiac troponin T.
RNA **13**, 2238-51.
Wenninger, S., Montagnese, F. and Schoser, B. (2018). Core Clinical
Phenotypes in Myotonic Dystrophies. *Front Neurol* **9**, 303.
Wieben, E. D., Aleff, R. A., Tang, X., Butz, M. L., Kalari, K. R., Highsmith, E.
W., Jen, J., Vasmatzis, G., Patel, S. V., Maguire, L. J. et al. (2017). Trinucleotide
Repeat Expansion in the Transcription Factor 4 (TCF4) Gene Leads to Widespread
mRNA Splicing Changes in Fuchs' Endothelial Corneal Dystrophy. *Invest Ophthalmol*
Vis Sci **58**, 343-352.
Winkler, N. S., Milone, M., Martinez-Thompson, J. M., Raja, H., Aleff, R. A.,
Patel, S. V., Fautsch, M. P., Wieben, E. D. and Baratz, K. H. (2018). Fuchs'
Endothelial Corneal Dystrophy in Patients With Myotonic Dystrophy, Type 1. *Invest*
Ophthalmol Vis Sci **59**, 3053-3057.
Zhao, Y., Ogawa, H., Yonekura, S., Mitsuhashi, H., Mitsuhashi, S., Nishino,
I., Toyoshima, C. and Ishiura, S. (2015). Functional analysis of SERCA1b, a highly
expressed SERCA1 variant in myotonic dystrophy type 1 muscle. *Biochim Biophys Acta*
1852, 2042-7.
736

Figure 1 Zebrafish models of human *MBNL* loss of function disorders were generated by mutating *mbnl1, mbnl2,* and *mbnl3*. A. Primary molecular mechanism of the human MBNL loss of function disorders myotonic dystrophy (DM) and Fuchs endothelial corneal dystrophy (FECD). In unaffected individuals, free MBNL1, MBNL2, and MBNL3 proteins bind to target pre-mRNAs and regulate the inclusion of alternative exons in mRNAs. MBNL proteins suppress alternative exon inclusion in the example shown here, but they promote inclusion of other target alternative exons. In individuals with DM or FECD, MBNL proteins are sequestered by long CUG or CCUG repeat RNAs, which decreases their availability to bind to target pre-mRNAs and alters mRNA isoform production and downstream protein function. **B.** Diagram of WT and mutant zebrafish *mbnl1*, *mbnl2,* and *mbnl3* predicted coding sequences. Mutant sequences are shown in Table S2. *mbnl1* mutant alleles are denoted as 1^A and 1^B, *mbnl2* as 2^A, and *mbnl3* as $3^{A/A}$, $3^{B/B}$, and $3^{C/C}$.

Figure 2 Misregulation of *mbnl* alternative splicing was conserved in zebrafish *mbnl* mutants. A & B. RT-PCR analysis showing inclusion of **(A)** *mbnl1* exon 5 or **(B)** *mbnl2* exon 5 in WT and *mbnl* mutant whole larval zebrafish at 5 days post fertilization (dpf). Each dot represents RNA pooled from 5 larvae. **C & D.** RT-PCR analysis showing inclusion of **(C)** *mbnl1* exon 5 or **(D)** *mbnl2* exon 5 in WT and *mbnl* mutant adult zebrafish skeletal muscle. Each dot represents RNA from one fish. Error bars indicate standard error of the mean. For all graphs, *mbnl1* mutant alleles are denoted as 1^A and 1^B, *mbnl2* alleles as 2^A, and *mbnl3* alleles as 3^A, 3^B, and 3^C. Representative RT-PCR gels are shown above each graph. Dividing lines indicated samples run on separate gels. Data were analyzed by ordinary one-way ANOVA with Tukey's multiple comparisons test. Data bars that do not share a letter above them are significantly different from one another.

Figure 3 Alternative splicing of *mbnl1* and *mbnl2* was misregulated across tissues in zebrafish *mbnl* mutants. A-H. RT-PCR analysis showing inclusion of *mbnl1* exon 5 **(A, C, E, & G)** or *mbnl2* exon 5 **(B, D, F, & H)** in WT and *mbnl* mutant adult zebrafish heart **(A & B)**, brain **(C & D)**, cornea **(E &F)**, and intestine **(G & H)**. For all graphs, *mbnl1* mutant alleles are denoted as *1^A* and *1^B*, *mbnl2* alleles as *2^A*, and *mbnl3* alleles as *3^A*, *3^B*, and *3^C*. Each dot represents RNA from one fish. Error bars indicate standard error of the mean. Representative RT-PCR gels are shown above each graph. Dividing lines indicated samples run on separate gels. Data were analyzed by ordinary one-way ANOVA with Tukey's multiple comparisons test. Data bars that do not share a letter above them are significantly different from one another.

Figure 4 Misregulation of many myotonic dystrophy-associated alternative splicing events was conserved in zebrafish *mbnl* mutants. **A-E.** RT-PCR analysis showing inclusion of **(A)** *ank3b* exon 36, **(B)** *atp2a1l* exon 24, **(C)** *ryr1b* exon 72, **(D)** *tpd52l2b* exon 4, and **(E)** *csnk1da* exon 9 in WT and *mbnl* mutant adult zebrafish skeletal muscle. For all graphs, homozygous *mbnl1* mutant alleles are denoted as 1^A and 1^B, *mbnl2* alleles as 2^A, and *mbnl3* alleles as 3^A, 3^B, and 3^C. Each dot represents RNA from one fish. Error bars indicate standard error of the mean. Representative RT-PCR gels are shown above each graph. Dividing lines indicated samples run on separate gels. Data were analyzed by ordinary one-way ANOVA with Tukey's multiple comparisons test. Data bars that do not share a letter above them are significantly different from one another.

Figure 5 Double and triple homozygous *mbnl* mutant zebrafish were decreased in length. A. Standard length of young adult 60 days post fertilization (dpf) WT and *mbnl* mutant zebrafish that were raised in the same tank.   On the right are representative images of fish of each genotype taken at 76 dpf. **B.** Standard length of 7 dpf WT and double *mbnl* mutant zebrafish. **C.** Standard length of 7 dpf clutchmates from an incross of *mbnl1*[+/B]*;mbnl2*[A/A]*;mbnl3*[C/C] fish. For all graphs, homozygous *mbnl1* mutant alleles are denoted as *1*[A] and *1*[B], *mbnl2* alleles as *2*[A]*,* and *mbnl3* alleles as *3*[A]*, 3*[B]*,* and *3*[C]*.* Each dot represents one fish and error bars indicate standard deviation. Data were analyzed by ordinary one-way ANOVA with Tukey's multiple comparisons test. Data bars that do not share a letter above them are significantly different from one another.

Figure 6 Zebrafish *mbnl* mutants exhibited altered swimming behavior. A.
Example still image taken from video of swim tests. Software was used to track the
center of mass of each fish in the X and Y directions over 5 minutes. **B.** Representative
traces of individual fish of each genotype taken during the swim tests. X and Y axes are
the same as in A. **C.** Total distance that adult fish swam during a five-minute swim
test. The Y axes are the same for all three graphs. **D.** Average relative distance from
the bottom of the tank of fish during a five-minute swim test. Zero represents the
bottom of the tank and 100 represents the top of the tank. The Y axes are the same for
all three graphs. For all sub-figures, homozygous *mbnl1* mutant alleles are denoted as
1^A and 1^B, *mbnl2* alleles as 2^A, and *mbnl3* alleles as 3^A, 3^B, and 3^C. Each dot represents
one fish and error bars indicate standard error. Single and double mutant data were
analyzed by ordinary one-way ANOVA with Tukey's multiple comparisons test, while
triple mutant data were analyzed by a Student's t-test. Data bars that do not share a
letter above them are significantly different from one another.