Use of a mechanistic growth model in evaluating post-restoration habitat quality for juvenile salmonids

Carlos M. Polivka[1†], Joseph R. Mihaljevic[2‡], Greg Dwyer[2],

1 Pacific Northwest Research Station, USDA Forest Service, Wenatchee, WA 98801 USA
2 Department of Ecology and Evolution, University of Chicago, Chicago, IL 60637 USA

†Corresponding author: carlos.polivka@usda.gov
‡Current Address: School of Informatics, Computing, and Cyber Systems, Northern Arizona University, Flagstaff, AZ 86001, joseph.mihaljevic@nau.edu

Abstract

Individual growth data are useful in assessing relative habitat quality, but this approach is less common when evaluating the efficacy of habitat restoration. Furthermore, available models describing growth are infrequently combined with computational approaches capable of handling large data sets. We apply a mechanistic model to evaluate whether selection of restored habitat can affect individual growth. We used mark-recapture to collect size and growth data on sub-yearling Chinook salmon and steelhead in restored and unrestored habitat in five sampling years (2009, 2010, 2012, 2013, 2016). Modeling strategies differed for the two species: For Chinook, we compared growth patterns of individuals recaptured in restored habitat over 15-60 d with those not recaptured regardless of initial habitat at marking. For steelhead, we had enough recaptured fish in each habitat type to use the model to directly compare habitats. The model generated spatially explicit growth parameters describing size of fish over the growing season in restored vs. unrestored habitat. Model parameters showed benefits of restoration for both species, but that varied by year and time of season, consistent with known patterns of habitat partitioning among them. The model was also supported by direct measurement of growth rates in steelhead and by known patterns of spatio-temporal partitioning of habitat between these two species. Model parameters described not only the rate of growth, but the timing of size increases, and is spatially explicit, accounting for habitat differences, making it widely applicable across taxa. The model usually supported data on density differences among habitat types in Chinook, but only in a couple of cases in steelhead. Modeling growth can thus prevent overconfidence in distributional data, which are commonly used as the metric of restoration success.

Introduction 1

An important advance in restoration ecology has been the development of methods of 2
evaluating restoration treatments. The most common approach is to rely on 3
observations of relative density in restored habitat and unrestored habitat, but such 4
data often have high uncertainty [1] or show weak or no effects of restoration [2,3]. 5
Another option is mechanistic model fitting, often used at the whole-population scale, 6
which may involve the estimation of demographic parameters and survival through 7
stages of the life cycle [4–6]. Because of the development of robust statistical tools [7], 8

model fitting has become a vital tool in conservation biology, but the models again rely heavily on population numbers or densities and correlative relationships between density and environmental variables [8].

In many cases, however, growth can provide a more robust indication of habitat quality than numbers or densities [9–11], especially given that growth is often correlated with survival and reproduction [12,13]. This can be important in restoration ecology when differences in habitat quality drive heterogeneity in growth and development across individuals, leading to life-history variation and trait evolution [14]. Growth rates and other life-history traits are nevertheless used only rarely to assess habitat restoration efforts [15,16]. There is no shortage of models that quantify effects of habitat quality on growth, ranging from calculation of suitability indices [17] to individual-based behavioral models [18]. Growth models are also useful for providing descriptions of growth patterns when there are multiple ecological inputs [19–22]. In cases when growth data have been collected, it should therefore be possible to combine models and data to provide more accurate assessments of restoration efforts. Such an approach could augment, and increase confidence in, distribution and abundance studies.

In contrast to demographic studies, what is often missing in growth studies is the use of robust statistical methods of fitting the models to data. In growth studies, comparison of growth rates between habitats is usually based only on mean growth rates [23]. Moreover, when mechanistic models [21,22] are fit to the data, they are usually still sufficiently simple that they can be transformed to the point at which linear fitting methods can be used [24]. This restriction on model complexity likely reduces our ability to make inferences about growth. Modern methods of non-linear fitting therefore provide an opportunity to quantify the effects of habitat restoration on growth in a statistically robust fashion. With the proliferation of habitat restoration efforts in species conservation, the the use of mechanistic models to describe growth data also has the potential to expand the theoretical basis of restoration ecology [25].

Growth and development are key features of salmonid life-cycle models, which are often able to accurately predict population trajectories [4]. Studies of restoration effectiveness, however, only occasionally quantify salmonid growth or survival [3,26,27], requiring that related parameters in life cycle models be estimated from correlative data in earlier research. The extent to which these life history traits are changed by restoration is nevertheless of high importance in the use of life cycle models to predict and confirm population-level responses [8].

Here we present the results of a multi-year study of growth in sub-yearling anadromous salmonids. We use our data to illustrate the usefulness of mechanistic growth models in restoration ecology. We collected size data, as well as growth over time, via fish marked and recaptured through the entire rearing season in each of five years to determine whether growth of young-of-the-year Chinook and steelhead is improved in restored habitat. Because these fish grow more or less continuously during the growing season, we were able to fit mechanistic growth models to both types of data. We used Bayesian model fitting techniques to estimate the parameters of our growth functions, and we compared growth rates between individuals with different habitat selection behaviors. Salmonid growth rates are rarely constant; therefore, we allowed for the possibility that growth-rate parameters would change over time [28,29]. Steelhead recaptures were sufficient in both habitat types, so we were able to use raw growth data to determine the extent to which the model accurately reproduces growth rates in the field. We then compared our estimates of growth rates with previous observations of relative density [30] to determine the frequency with which increased abundance in restored habitat is associated with increased growth rates. The model demonstrates relatively consistently improved growth associated with restoration in Chinook, but that this was the case in only two study years for steelhead. Nevertheless, key model

parameters describing growth rate were supported by steelhead raw growth data. Thus, our model has a strong application to evaluation of salmon conservation efforts.

Materials and methods

Study System

Restoration of fish habitats in rivers involves a large set of approaches and can sometimes target more than one species [27, 31]. A common stream restoration technique is the installation of engineered log jams (ELJs) in streams, which create pools that serve as primary habitat for target species. In the Pacific Northwest, USA, use of ELJs is a common conservation strategy for threatened and endangered salmonids [2, 31, 32]. In the interior Pacific Northwest, ELJs are targeted toward sub-yearling anadromous Chinook salmon (*Oncorhynchus tschawytscha*), which grow and develop in streams for one year, and steelhead trout (*O. mykiss*) which grow and develop in streams for 1-3 years, before migrating to the marine environment. Seaward migration occurs at the smolt stage and, in principle, ELJs should improve growth and survival to this stage. However, distribution patterns of salmonids vary spatio-temporally, and differ between species [33], exacerbating the difficulty of assessing restoration efficacy with distribution and abundance studies [27, 30, 34–36].

The restoration project studied here was implemented in the Entiat River sub-basin (49.6567° N, 120.2244° W), a tributary of the interior Columbia River. Restoration efforts in the Entiat River spanned the years 2008-2016, and during this period, ELJs were installed in multiple reaches of the river. We compared a single restored reach that was 4.5-4.9 km from the confluence with the Columbia River, and an unrestored reach that was 0.3 km upstream of the restored reach (Fig. 1). The restored reach was the only one with ELJs at the beginning of the study in 2009. ELJ pools in the restored reach ("restored pools"; N = 11) were compared with pools in the unrestored reach (N = 11) that lacked major log and rock structures. Both types of pools occurred on the stream margins where nearly all sub-yearling habitat use occurs beechie2005classification. Unrestored pools were 20-40% smaller and shallower than restored pools, but [30, 37] showed that the unrestored pools had a similar rate of increase in fish density with increasing area to the restored pools. We therefore concluded that the unrestored pools served as reasonable controls for the restored pools.

Fig 1. Map of study area in the Entiat River. Locations (river km relative to the confluence with the Columbia River) of the study reaches with restored pools and with unrestored pools are indicated.

Fish Capture and Marking

We collected growth data using mark and recapture of young-of-the-year Chinook and steelhead in restored and unrestored pools during the growing seasons of 2009, 2010, 2012, 2013 and 2016. In each set of pools, capture and recapture were carried out every 10-14 days throughout the growing season, and recapture intervals therefore ranged from 15-65 days [38]. We collected no growth data in 2011, 2014 or 2015 because periods of high water (2011) and post-fire sediment deposition (2014, 2015) prevented mark-recapture studies.

Fish were captured using a 3 m × 1.5 m × 3 mm mesh seine. The substrate and flow conditions made it impossible to pull the seine through the water, and so two field crew members instead stood at the downstream end of the pool and held the seine open as

two other crew members, snorkeling in the water, used large hand nets to capture fish $_{103}$ individually or to coerce fish into the seine. Visibility in the Entiat River is 4-5 m, so $_{104}$ the two snorkelers could see the entire sampled area, ensuring that all visible fish were $_{105}$ captured. Captured fish were transferred to insulated, aerated buckets for enumeration, $_{106}$ marking and recording of size data (standard length, SL, in mm and mass in g). $_{107}$

Mild anaesthetization using MS-222 (<0.1 g·l^{-1}) exposure for 2-3 minutes made it $_{108}$ possible to measure, weigh and apply an identifying mark to each fish. We avoided PIT $_{109}$ tagging because it would have caused too much handling stress [39] to allow for robust $_{110}$ mark and recapture, especially given the small size (< 50 mm SL) of some fish at the $_{111}$ start of field sampling each year. We therefore marked fish with a subcutaneous $_{112}$ injection of visual implant elastomer (VIE, Northwest Marine Inc.), a minimally $_{113}$ invasive procedure. Individuals were identified by varying VIE color combinations and $_{114}$ positions on the body. We marked ~ 1000 fish each year, with fewer than 2-3 deaths $_{115}$ per year from handling/marking procedures. Following handling, fish were held in an $_{116}$ aerated bucket for at least 10 min, or until they displayed a full righting response and $_{117}$ normal activity, and were then released into the pool in which they were captured. $_{118}$

After 24 hrs field crews returned to the same pools as part of a separate short-term $_{119}$ behavioral study (Polivka in review) and captured all fish using the same methods. $_{120}$ Recaptured fish were released to the pool and newly captured fish on that day were $_{121}$ given marks as described above. The pools were sampled again 10-14 days later, with $_{122}$ size data recorded on recaptured fish and marks applied to newly captured fish that had $_{123}$ immigrated into the pool during the sampling interval. $_{124}$

Data Analysis $_{125}$

To analyze our data, we constructed mechanistic growth models that described fish $_{126}$ growth over each growing season, and we fit these models to data on fish size. Size data $_{127}$ included observations both of individuals marked and recaptured, and of individuals $_{128}$ that were marked but never recaptured. We then compared best-fit growth models in $_{129}$ restored and unrestored habitats. In these species, individual growth data are typically $_{130}$ collected via mark-recapture studies, but the high mobility of Chinook [40] often limits $_{131}$ their recapture rates, especially in unrestored habitats [38], and fitting growth models $_{132}$ allowed us to nevertheless use these data. To do this, we combined the repeated size $_{133}$ measurements on the recaptured individuals with individuals captured only once, $_{134}$ regardless of the habitat of initial capture. The repeated measurements of recaptured $_{135}$ fish ("recaptures") required use of a random effect in the model specification (see $_{136}$ below), whereas the individuals captured only once ("others") only required fitting the $_{137}$ model to the sizes observed during the course of the season. This analysis for Chinook $_{138}$ is more conservative, because at least some of the "others" occupied restored habitat for $_{139}$ some portion of the interval of time between sampling events. If there is a benefit to $_{140}$ restoration some portion of the "others" will therefore be more comparable to $_{141}$ "recaptures," and may affect the likelihood of detection of growth differences with the $_{142}$ model. For steelhead in contrast, we had a large number of recaptures in unrestored $_{143}$ habitat (N = 179), and so we fit models to the data only for recaptured fish and $_{144}$ compared habitat types. $_{145}$

Model Specification $_{146}$

We constructed our growth model to describe the growth of young-of-the-year Chinook $_{147}$ salmon and steelhead in streams during the growing season. Size-over-time models in $_{148}$ fisheries, e.g., [19] usually describe growth over the lifetime of the organism. One $_{149}$ important consideration was that inspection of our data showed that there were $_{150}$ inflection points; therefore, we used a generalized version of a logistic growth $_{151}$

model [20–22]. This model allows for accelerating growth in the early stages of the growing season, followed by an inflection point, after which growth decelerates, leading to a maximum size at the end of the first season. The use of a size asymptote is consistent with patterns evident in our data, and in size data reported from several years of downstream migration of smolts in the spring [41]. Our model is:

$$l_t = Y' + \frac{L_\infty - Y'}{1 + e^{\hat{\alpha} - at}}. \tag{1}$$

Here, l_t is the length at time t, Y' is the lower bound on size, effectively a minimum, and a is the rate of increase in the size curve. L_∞ is the maximum size that a fish can reach during the growing season, so that $Y' \leq l_t \leq L_\infty$ for all time t. The inflection point of the curve occurs at $\hat{t} = \frac{\hat{\alpha}}{a}$.

Importantly, both $\hat{\alpha}$ and a can have profound effects on the dynamics during the growing season, potentially resulting in substantial differences in l_t at any given time t. As the value of $\hat{\alpha}$ decreases, with the other parameters held constant, fish reach larger sizes earlier in the season. The a parameter describes the steepness of of the curve at small sizes, so that increasing values of a result in more rapid increases in size over time (Fig. 2).

Fig 2. Growth curves generated from Eq. 1. over a 100 day growing period. A: Effect of $\hat{\alpha}$ on the timing of the inflection point. B: Effect of a on the slope of the growth curve. Panels C and D: Growth curves for values of the two parameters that are closer to those estimated from our data. For the curves in C in particular, the inflection point occurs close to time $t = 0$, so that the intercepts vary between parameter values. In these cases, fish are close to the lower bound on the curve Y' when time is less than zero, which is to say, before sampling begins. In all panels $Y' = Y_\infty = 35$.

To avoid confusion with other models such as the Von Bertalanffy model, we re-parameterized the model according to $Y_\infty = L_\infty - Y'$, so that Y_∞ is the total amount by which a fish increases in size during the growing season. The model is then:

$$l_t = Y' + \frac{Y_\infty}{1 + e^{\hat{\alpha} - at}}, \tag{2}$$

When non-linear models are fit to data, a standard approach is to transform the model into a linear form to accommodate linear least-squares fitting routines and their assumptions, such as normally distributed residuals. For our data, however, it turned out that the residuals are normal even without transformation of the model, and the variance in the residuals for the untransformed model was roughly constant. We therefore did not transform the model. Instead we simply added residual variation to the non-linear model, according to:

$$l_{it} = Y' + \frac{Y_\infty}{1 + e^{\hat{\alpha} - at}} + \epsilon_{it}, \tag{3}$$

Here i indicates individual fish i. We then assume that the residual ϵ_{it} for fish i at time t follows a normal distribution with mean zero, and variance $\sigma^2_{residual}$, which we estimated from the data. We thus have $\epsilon_{i,t} \sim N(0, \sigma^2_{residual})$, so that $\sigma^2_{residual}$ is the residual variation in length. Thus, Y' and Y_∞ describe growth bounds averaged over individuals in the population.

We then extended the model to allow for fixed treatment effects, and random year and individual effects. First, to include habitat treatment effects, restored vs. unrestored, we allowed the four model parameters (Y', Y_∞, $\hat{\alpha}$, and a) to vary by habitat type, as fixed effects. Second, to account for the statistical properties of our sampling

design, we included random effects of year, to allow for annual variability in the parameters. Third, in the case of individuals that were recaptured during a season, we accounted for repeated measurements by including random effects of individual. Given these considerations, the model becomes:

$$l_{ihyt} = (Y'_h + \epsilon^1_{hy}) + \frac{(Y_{\infty h} + \epsilon^2_{hy})}{1 + e^{\left[(\hat{\alpha}_h + \epsilon^3_{hy}) - (a_h + \epsilon^4_{hy})t\right]}} + \epsilon^5_{ih} + \epsilon^6_{iht} \tag{4}$$

Here i is again the individual, h is the habitat type, restored or unrestored, y is the sampling year, and t is time within a growing season. The ϵ's represent the different types of random effects, as follows.

The random effects of year on each of the four model parameters, $\epsilon^{1\ldots4}_{hy}$, vary between the two habitats, such that, for example, $\epsilon^1_{hy} \sim N(0, \sigma^2_{Y'})$, where $\sigma^2_{Y'}$ is the random variation in initial size Y' measured across habitats h and years y. For the random effect of recaptured individuals, $\epsilon^5_{ih} \sim N(0, \sigma^2_{individual_h})$, where $\sigma^2_{individual_h}$ is the variance in fish length among recaptured individuals from a given habitat type. This random effect of individual affects only the overall length, rather than the rate of growth. Among other things, this means that we can account for the possibility of observing fish that start off larger than average being more likely to continue being larger than average throughout the growth period pfister2002genesis. To be conservative, we began by assuming that no Chinook in the unrestored pools were recaptured, and so the random effects of individual Chinook are only relevant for recaptures in the restored habitats.

Finally, the residual variation is allowed to differ between the two habitats, where $\epsilon^6_{ht} \sim N(0, \sigma^2_{residual_h})$. This is particularly important for the Chinook data set, because some of the variation among the recaptured individuals in restored habitats can be explained by the random individual effects. In the unrestored habitats in contrast, no individuals were recaptured, and therefore more variation was left unexplained. Because of this, it turned out to be the case that $\sigma^2_{residual_{unrestored}} > \sigma^2_{residual_{restored}}$.

Model fitting

We fit our models using Hamiltonian Monte Carlo (HMC) with the open-source software, *Stan* [42] and **R** [43], utilizing the package *rstan* [44]. The HMC algorithm provides an efficient method of fitting nonlinear models, and the software platforms ensure that our methods are open and reproducible (see Supporting Information).

In general, we used vague priors for our parameters, but to aid HMC performance, we centered the prior distributions of Y' and Y_∞ on realistic values, based on previous work with these two species. Also, we constrained $\hat{\alpha}$ and a to fall within realistic ranges, namely 0-10 and 0-1, respectively, again based on previous work. Sensitivity analysis showed that these priors did not strongly influence our posterior inferences, and that the posterior was clearly dominated by the likelihood, rather than by the prior distributions. For each model, we ran three HMC chains for 9000 iterations, using the first 2000 iterations as a warm up. We then thinned by 7 iterations to produce a total of 1000 samples per chain. We evaluated HMC chain convergence based on Gelman and Rubin's potential scale reduction factor, $\hat{R}$ [45]. We visually inspected chain mixing using traceplots, running mean plots, and marginal posterior density plots [45], and we saw no evidence of pathological MCMC behavior. Although we thinned our chains to avoid auto-correlation, our effective sample size was close to the total number of iterations, and thinning was likely unnecessary [46].

To compare parameter estimates between the restored and unrestored habitats, we calculated differences between paired parameter values for the restored and unrestored habitats for each sample in the model's joint posterior. For example, to calculate differences in the shape parameter between restored habitats r and unrestored habitats

u, we calculated $\hat{\alpha}_r - \hat{\alpha}_u$ for each of the 3000 samples in the posterior (our 3 Markov chains each produced 1000 samples). We then calculated the 95% credible interval (CI) of that set of differences. This approach allowed for the possibility of correlations in parameters across the samples in the posterior, and is therefore a more robust method than calculating parameter differences by making independent draws of each parameter from the posterior [47].

Note that, in the case of Chinook, the u group consisted of all uncategorized fish, meaning transient individuals that were not recaptured, regardless of habitat of original capture. Accordingly, when we tested for interactions between habitat and year for Chinook, we took into account the random year effects. For example, in testing for differences in the shape parameter between years for Chinook, we calculated $(\hat{\alpha}_r + \epsilon_{r,y}^1) - (\hat{\alpha}_u + \epsilon_{u,y}^1)$ for each posterior draw, where $\epsilon_{r,y}^1$ and $\epsilon_{u,y}^1$ are the random year effects in year y.

We considered the parameter estimates for the two habitats to be meaningfully different if the 95% CIs of their differences did not overlap zero. For most parameters, however, the fraction of differences that was above or below zero was reasonably consistent across years, even if the 95% CI overlapped zero. Because this consistency provides at least modest additional support for some of our arguments, we report the fraction of differences that were above or below zero in each case.

Our best-fit models suggested that fish almost always exceeded the minimum size Y' before sampling began. To estimate fish size at the beginning of sampling, we therefore used the best-fit models to back calculate median fish length at time $t = 0$ when observations began. This also allowed us to confirm that there were no differences in fish size among habitat types at the start of the season. Such initial differences can lead to growth advantages that are independent of environmental factors such as habitat quality [48].

Model Comparison With Direct Measurement of Growth in Steelhead

Because we recaptured steelhead in both restored and unrestored pools, we were able to directly compare observed growth rates for recaptured steelhead between habitat types, in addition to comparing our estimates of growth that came from fitting models to the data. In carrying out analysis of measured growth rates, a key consideration is that, in many organisms, individual growth rates are strongly size dependent [48–50]. Because we encountered a wide range of initial sizes (40-75 mm SL), it was important to allow for the possibility of size-dependent growth when comparing steelhead growth rates between habitats.

We generated a predicted growth vs. size regression for steelhead for each year, by carrying out a log-linear regression of change in size (mm $\cdot$ day^{-1} = $\frac{SL_{recaptured} - SL_{marked}}{days}$) on initial standard length, for each recaptured individual. To compare growth in restored vs. unrestored habitat in these regressions, we kept track of the habitat in which each individual was recaptured. We then used the residual for each point as an indicator of growth rate relative to the population average for that size, where the average for the size was calculated from the regression line of growth on size. A positive residual indicates that an individual has a higher than average growth rate, adjusted for size, whereas a negative residual indicates that an individual has a lower than average growth rate, again adjusted for size (Fig. 3). In comparing size-dependent growth rates between habitats, we compared the mean residual from restored habitat to the mean residual from unrestored habitat for each year of the study, using two-sample t-tests. Because growth rate in mm $\cdot$ day^{-1} corresponds to the a term in Equations 1-4, we compared differences among habitats in mean residuals with differences among habitats in the a parameter.

Fig 3. Use of residual values from a size vs. growth rate regression to compare growth in different habitats. The arrow shows the distance from the points to the line. Mean residual values $\bar{R}$ are calculated for all individuals in each habitat, and if $\bar{R}_r > \bar{R}_u$, then we conclude that growth is higher in restored than in unrestored habitats.

Concordance of Growth and Habitat Occupancy Patterns

From a management perspective, our goal was to determine whether observations of density differences between habitats are supported by growth differences. We therefore compared our model output first to the fish density data from [30] for each of the study study years 2009, 2010, 2012, and 2013. For 2016, we had density data available from initial captures during these mark-recapture studies.

Results

Chinook Salmon

Across the five sampling years (2009, 2010, 2012, 2013, and 2016), we collected 3,568 length records from Chinook. In the restored habitats, we recaptured 238 Chinook, for a total of 481 length records, with five fish that were recaptured more than once. The best-fit growth curves in Fig. 4 show that Chinook recaptured in restored habitats were generally larger early in the season compared to unrecaptured fish. This visual impression is confirmed by a lower value of $\hat{\alpha}$ for individuals recaptured in restored habitats compared to $\hat{\alpha}$ values for unrecaptured fish (Fig. 5). Because $\hat{\alpha}$ is an inverse measure of size early in the season, the difference in $\hat{\alpha}$ values means that the recaptured fish were larger than unrecaptured fish early in the season. Negative values of the difference in $\hat{\alpha}$ values occurred in 98.5% of sample differences for all years combined, and in 98.3%-100% of samples for 2009, 2010, and 2013 (Fig. 5). In 2012 and 2016, the 95% CIs overlapped zero, but negative values occurred in $> 90\%$ of samples, consistent with the differences for the other years and with the differences for all years together.

Fig 4. Growth curves based on estimated model parameters (mean of the posterior) for Chinook salmon recaptured in treated habitat ("recaps," solid lines) vs. those captured in either habitat type, but not recaptured ("others," dashed lines). The left panel shows the data and the habitat-specific models for all years combined. The small panels to the right show the data and the habitat-specific models for individual years. Parameter differences (see Fig. 5) are shown in the upper left corner of each panel.

Fig 5. The difference (Δ; recaptures $-$ others) in estimates of all model parameters $\pm$ the 95% credible interval between the two capture types of young-of-the-year Chinook salmon. Recaptures in restored habitat; others = fish marked and not recaptured, regardless of capture origin. Year-specific estimates incorporate the random effect of year, and combined differences are based on the average parameter values among years. The number in parentheses represents the fraction of posterior draws for which the difference was above or below zero, as appropriate. Δ Length at time = 0 was substituted for model parameter Y'.

In 2009 and 2010, however, values of the growth rate a were larger for unrecaptured fish than for recaptured fish. This observation, in combination with the larger $\hat{\alpha}$ values

for unrecaptured fish describing small early-season size, indicates that unrecaptured fish experienced a rapid increase in size later in the season (Fig. 4) that compensated for the early season disadvantage. The difference in those two years was indicated in >99% of differences in sample values drawn from the posterior distributions of a (Fig. 5). Although the 95% CI for the differences in a for all years together overlapped zero, they were below zero in 91.1% of the draws, consistent with the trend observed in 2009 and 2010.

All else equal, smaller $\hat{\alpha}$ values lead to earlier inflection points $\hat{\alpha}/a$. To test whether the smaller values of $\hat{\alpha}$ for Chinook recaptured in restored habitat in all years combined, and in 2009, 2010, and 2013, led to earlier inflection points for those fish, we drew pairs of values of the shape parameter $\hat{\alpha}$ and the growth parameter a for recaptured and unrecaptured Chinook. We then calculated differences in inflection points $\hat{\alpha}/a$ for recaptured and unrecaptured fish in each year and for all years combined, as we did for the other parameters.

This procedure showed that inflection points for recaptured fish did indeed occur earlier than for recaptured fish in 2009, with 99.8% of sampled differences being less than zero. In 2013, there was a similar trend, with 94.6% of sampled differences less than zero. Differences in the other years, and for the combined data, were not meaningful, likely because independent variation in the two parameters obscured differences in the inflection points. We therefore conclude that there is at least modest evidence that restoration is associated with larger Chinook size earlier in the season as indicated by the shape parameter $\hat{\alpha}$, but that variation in the two parameters prevents a consistently earlier inflection point.

Steelhead

We recorded the lengths of 2,871 steelhead across the five sampling years. In the restored habitats, we recaptured 306 individuals for a total of 689 observations, whereas in the unrestored habitats, we recaptured 179 individuals for a total of 413 observations. The number of steelhead recaptured from unrestored habitats was thus sufficient to fit the model directly to recaptured individuals in each habitat type, in contrast with Chinook. In 2013, however, there were no recaptures of steelhead in unrestored habitat for recapture intervals of longer than 24h. We therefore excluded the 2013 data from our analyses.

For all years combined, there were no differences in growth parameters in steelhead recaptured in restored versus unrestored habitats (Fig. 6), but there were differences in some parameters in 2009 and 2010 (Fig. 7). In those two years, both the shape parameter $\hat{\alpha}$ and the growth rate parameter a were higher for fish in restored habitat in 99-100% of samples from the posterior distributions. Over all years, almost 90% of differences in $\hat{\alpha}$ were above zero, and almost 93% of differences in a were above zero, supporting the meaningful differences seen in 2009 and 2010.

Fig 6. Growth curves based on estimated model parameters (mean of the posterior) for steelhead recaptured in restored habitat (solid lines) vs. those recaptured in unrestored habitat (dashed lines). Left panel shows data and fitted curves for all years combined; additional figures show individual years. Parameter differences (see Fig. 7) are indicated in the lower right corner of each panel. The analysis omits 2013 owing to lack of long-term steelhead recaptures in unrestored habitats.

The higher values of a for fish in restored habitat mean that growth rates were higher for steelhead in those two years, but the higher values of $\hat{\alpha}$ mean that fish in restored habitat reached larger size later in the season and caught up to the fish in

unrestored habitats that were larger earlier. The lack of differences in total growth Y_∞ supports this conclusion. The inflection point was later for steelhead in restored habitats in 2010 only (99.3% of sampled differences above zero), again suggesting that the inflection point is not a specific indicator of growth differences among habitats. Nevertheless, mid-season occupancy of restored pools was associated with rapid growth.

Fig 7. The difference in estimates of all model parameters $\pm$ the 95% credible interval between the two capture types of young-of-the-year steelhead (recaptures in restored habitat - recaptures in unrestored habitat). Year-specific estimates incorporate the random effect of year, and combined differences are based on the average parameter values among years. The number in parentheses represents the fraction of posterior draws for which the difference was above or below zero, depending on trend of the data. For all panels except the lower right, this represents the fraction of posterior draws above zero. Δ Length at time $= 0$ was substituted for model parameter Y'. The analysis omits 2013 owing to lack of long-term steelhead recaptures in unrestored habitats.

Direct Observations of Growth in Steelhead

Residuals from the log-linear regression of growth (mm $\cdot$ day^{-1}) on initial size were more positive, on average, for individuals marked and recaptured in restored habitat than in unrestored habitat, indicating higher growth in unrestored habitat. This trend, however, held for 2009 and 2010 but not 2012 or 2016 (Fig. 8). Inspection of the regression plots for 2009 and 2010 indicated that the largest differences in residuals between habitats in those two years were among individuals 55-60 mm in length. Separate residual analyses on individuals < 60 mm, and individuals > 60mm, produced significant differences in residuals only in individuals < 60 mm (not shown).

Fig 8. Mean residuals from a log-linear regression of growth rate versus size, in restored vs. unrestored habitat, for each study year. The analysis omits 2013 owing to lack of long-term steelhead recaptures in unrestored habitats.

Concordance of growth and density

In each year of our study, Chinook salmon were more abundant in restored habitat than in unrestored habitat whereas parameters from the growth model only showed a growth benefit in three of the five years (2009, 2010 and 2013; Table 1). For all years combined, however, density and growth were concordant for Chinook. For steelhead, 2009 was the only year in which there was concordance between observed fish abundance and growth patterns. In 2010, when there was no significant difference in density among habitats, there was nevertheless more rapid growth in restored pools, as indicated above by the a term in the model and by the positive mean residuals from the growth vs. size curve (Fig. 8 and Table 1).

Discussion

Mechanistic ecological models play an important role in restoration ecology [51], because their description of observed data increases the chances that species conservation efforts will be successful. Perhaps the best known set of tools comes from population viability analysis. Population viability analysis has been widely used in the

Table 1. Concordance of occupancy differences and growth-parameter differences. Recall that large $\hat{\alpha}$ means that the inflection point comes later in the season, indicating less favorable growth conditions (NS = No difference/not significant).

Year	Density higher in ...	Growth more favorable in...	Parameters
a) *Chinook Salmon*			
2009	Restored	Restored	$\hat{\alpha}_{\text{restored}} < \hat{\alpha}_{\text{unrestored}}$ $a_{\text{restored}} < a_{\text{unrestored}}$
2010	Restored	Restored	$\hat{\alpha}_{\text{restored}} < \hat{\alpha}_{\text{unrestored}}$ $a_{\text{restored}} < a_{\text{unrestored}}$
2012	Restored	NS	
2013	Restored	Restored	$\hat{\alpha}_{\text{restored}} < \hat{\alpha}_{\text{unrestored}}$
2016	Restored	NS	
Years Combined	Restored	Restored	$\hat{\alpha}_{\text{restored}} < \hat{\alpha}_{\text{unrestored}}$
b) *Steelhead*			
2009	Restored	Restored	$a_{\text{restored}} > a_{\text{unrestored}}$
2010	NS	Restored	$a_{\text{restored}} > a_{\text{unrestored}}$
2012	Restored	NS	
2013	NS	NS	
2016	NS	NS	
Years Combined	Restored	NS	

development of recovery criteria, and in calculations of the amount of habitat that needs to be restored [52]. Species distribution analysis is a related approach that is used to identify the habitat characteristics that are important for conservation [53, 54]. Like population viability analysis, species distribution analysis is used to predict how species respond to management actions or environmental change [55]. Predator-prey theory has been used to construct more complex models, again to analyze the outcomes of conservation actions [56].

All these methods, however, assume that there are correlations between abundance and habitat characteristics, but the support for such correlations is often mixed [57]. Growth data in contrast are infrequently used in restoration ecology. Surveys of relative abundance cannot easily identify the processes that determine ecological differences between habitat types [9, 58, 59]. Data sets on growth thus provide opportunities to expand the use of mechanistic models in describing specific effects of habitat restoration in conservation efforts. By comparing growth between restored and unrestored habitat, we have shown that restoration enhances growth rather than simply redistributing organisms among habitats or increasing numbers. Compared to observations of abundance, fitting mechanistic growth models to data can thus provide stronger evidence in favor of restoration.

Growth is an important life history trait because of its correlation with fitness in many species. Growth models can therefore help identify the potential fitness advantages of a given habitat type. Accordingly, although we designed our analysis to focus on a specific study system, our over-arching goal is to to provide an adaptable tool for a wide range of studies. Moreover, by using state-of-the-art Bayesian methods, we have shown how growth models can be used to make statistically robust inferences in restoration ecology. The development of Bayesian analytical methods has similarly allowed for the expansion of the applicability of ecological models in other areas of restoration ecology. For species distribution analysis, for example, Bayesian methods have provided useful examples of the potential of advanced statistical computing in restoration ecology [60–62].

Although our model describes a specific life history trait for our study species, in the

form of growth during early development, we constructed our model by adapting models $_{400}$
that describe growth over the lifetime of an organism. Our model could thus be $_{401}$
expanded to describe longer segments of a species' life history. Moreover, our model $_{402}$
offers two advantages over previous models. First, our model includes parameters $_{403}$
describing not only the rate of growth, represented by a, but also the timing of size $_{404}$
increases, represented by the shape parameter, $\hat{\alpha}$. Second, the model allows for spatial $_{405}$
variation, in that its parameters and residuals were allowed to vary by habitat type, $_{406}$
thus allowing for direct description of habitat differences in the data. $_{407}$

Our estimates of the model parameters for sub-yearling Chinook salmon and $_{408}$
steelhead demonstrate that habitat restoration can make growth conditions more $_{409}$
favorable. Our estimates of the shape parameter $\hat{\alpha}$ showed that Chinook reached large $_{410}$
size earlier in three of the five study years, and across all years combined, as a result of $_{411}$
remaining in restored pools. In two of the study years, the growth rate parameter, a, $_{412}$
was higher among individuals that moved among habitat types, but this occurred later $_{413}$
in the season. The interpretation of model parameters therefore can be specific to the $_{414}$
study system. $_{415}$

After growth to the parr stage (late August/early September), Chinook begin $_{416}$
migrating downstream to overwintering habitat [40,63]. As Chinook migrate $_{417}$
downstream, slower growing, later hatching, or previously more transient Chinook can $_{418}$
immigrate into the pools being vacated by earlier occupants. This can lead to $_{419}$
compensatory growth because of release from competition [64]. Chinook previously at a $_{420}$
growth disadvantage may then increase their growth once their competitors leave, and $_{421}$
may ultimately grow faster than their competitors once did [65]. This effect also likely $_{422}$
depends on the overall population size and distributional differences between habitat $_{423}$
types. The largest difference in Chinook abundance between habitats was in 2009 and $_{424}$
2010 [30]. Thus, fish in those two years were more likely to have been originally $_{425}$
captured in restored pools. By mid-season, individuals captured in restored pools could $_{426}$
have occupied a pool for up to 10-14 days, the typical interval between sampling events, $_{427}$
even if not recaptured later. This could explain the higher growth rate a in $_{428}$
unrecaptured Chinook initially captured during mid-season of 2009 and 2010 (Fig. 5). $_{429}$

As an explanation for observed habitat differences in steelhead growth in 2009 and $_{430}$
2010, competitive release is also consistent with estimated growth parameters. In this $_{431}$
study system, steelhead are less abundant overall and overlap with Chinook, $_{432}$
particularly in restored pools [30]. When steelhead are less abundant than other $_{433}$
salmonids, they usually respond by adopting a more generalist habitat use pattern [66]. $_{434}$
Indeed, steelhead recapture rates in unrestored pools indicate higher residency time $_{435}$
there compared with Chinook [38], despite the fact that they are usually relatively more $_{436}$
abundant in restored pools [30,67]. If early season growth by steelhead in restored pools $_{437}$
is slowed by competition with Chinook, then steelhead might do better in unrestored $_{438}$
pools, especially in early season. $_{439}$

Moreover, the competitive pressure exerted by Chinook is probably more severe $_{440}$
when Chinook are at high density. Further evidence that competition affects steelhead $_{441}$
growth then comes from the observation that the $\hat{\alpha}$ parameter for steelhead was larger $_{442}$
in restored habitat in 2009 and 2010, when Chinook densities were highest. In those two $_{443}$
years, the $\hat{\alpha}$ parameter indicated that steelhead attained larger size earlier in unrestored $_{444}$
habitat. The opportunity to grow rapidly in restored habitat with less competition $_{445}$
becomes available mid-season when Chinook begin downstream migration. Because our $_{446}$
best-fit models also showed that transient Chinook showed higher growth during this $_{447}$
time, our model-fitting approach has apparently revealed the effects of both intra- and $_{448}$
interspecific competition. $_{449}$

Direct measurements of growth in in steelhead supported the inferences we made by $_{450}$
model-fitting. Although direct measurements can in some cases provide more obvious $_{451}$

answers, our model-fitting approach has important advantages. First, our model-fitting ²⁴⁵²
approach is capable of not only comparing habitats directly, but also of using time series ⁴⁵³
of size to compare recaptured and non-recaptured individuals (as in the case of ⁴⁵⁴
Chinook). Thus it also integrates over the effects of environmental variables, which are ⁴⁵⁵
difficult to include as individual correlates in other approaches, across the whole rearing ⁴⁵⁶
season. Second, model-fitting allowed us to test for effects of different habitats on ⁴⁵⁷
Chinook, even though the Chinook recapture rate in unrestored habitat type was nearly ⁴⁵⁸
zero [38]. Third, our approach demonstrated that subtle differences in growth exist ⁴⁵⁹
between early- and mid-season for both species. ⁴⁶⁰

Because our data are a time series, mark-recapture intervals can be different and can ⁴⁶¹
span different parts of the growing season. This further increases the difficulty of ⁴⁶²
correlating growth with environmental variables. Continuous monitoring of depth, ⁴⁶³
current velocity, and other physical variables is costly and, when fish are marked and ⁴⁶⁴
recaptured at different time points in the season, the measured growing interval can ⁴⁶⁵
vary among individual fish. Thus, relating physical data to fish behavior is not easy. ⁴⁶⁶
Indeed, exploratory multiple regression analyses attempting to predict growth data with ⁴⁶⁷
variables such as temperature and current velocity yielded no discernible pattern [38]. ⁴⁶⁸

Our work has implications for the use of life cycle models to predict population-level ⁴⁶⁹
responses of salmonids to restoration [4,68], but there are also substantial uncertainties. ⁴⁷⁰
First, although we found that there was higher growth in restored habitat, and although ⁴⁷¹
changes in habitat capacity are allowed for in life cycle models [4], such changes have ⁴⁷²
not yet been detected in nearby sub-basins [8]. Second, positive growth in restored ⁴⁷³
habitat is evidence of capacity increase, but whether this is important for the entire life ⁴⁷⁴
cycle remains unclear. Salmonid growth at one life stage is sometimes a strong predictor ⁴⁷⁵
of growth at a subsequent stage [69], but not always [70]. More broadly, observations of ⁴⁷⁶
increased growth at a few sites of course does not mean that growth would be increased ⁴⁷⁷
over larger areas. ⁴⁷⁸

Model fitting nevertheless allowed us to show that growth can be reliably quantified ⁴⁷⁹
as a biological response to habitat quality differences, a rare accomplishment in fish ⁴⁸⁰
restoration. Model fitting also provided at least partial confirmation that differences in ⁴⁸¹
growth between habitats are concordant with differences in relative density between ⁴⁸²
habitats. It is also important to note that, in 2010, higher steelhead growth was not ⁴⁸³
associated with higher abundance, emphasizing that there is more to restoration than ⁴⁸⁴
abundance. Finally, the model strongly suggests that phenology and competition may ⁴⁸⁵
be modulated by habitat restoration. Growth models may thus provide a tool of general ⁴⁸⁶
usefulness in restoration ecology. ⁴⁸⁷

Supporting information ⁴⁸⁸

All of our data and model-fitting code are available in an open-source repository: ⁴⁸⁹
`https://bitbucket.org/jrmihalj/mechanistic_growth_salmon` ⁴⁹⁰

Acknowledgments ⁴⁹¹

Early portions of this work (2009-2010) were funded by Bonneville Power ⁴⁹²
Administration (Project No. 2003-017-00), and by the American Recovery and ⁴⁹³
Re-investment Act enacted by President B. Obama. The later activities (2012-2016) ⁴⁹⁴
were funded by the U.S. Bureau of Reclamation. JRM was funded by a US Department ⁴⁹⁵
of Agriculture (USDA) National Institute of Food and Agriculture (NIFA) Postdoctoral ⁴⁹⁶
Fellowship (2014-67012-22272). The authors have no conflict of interest. We thank ⁴⁹⁷
Cascadia Conservation District, S. Eichler, R. Logan, A. Bushy (supported by the ⁴⁹⁸

References

1. Taylor BL, Wade PR. "Best" abundance estimates and best management: why they are not the same. Quantitative methods for conservation biology Springer-Verlag, New York, New York, USA. 2000; p. 96–108.

2. Irwin LL, Rock DF, Rock SC, Loehle C, Van Deusen P. Forest ecosystem restoration: Initial response of spotted owls to partial harvesting. Forest Ecology and Management. 2015;354:232–242.

3. Sievers M, Hale R, Morrongiello JR. Do trout respond to riparian change? A meta-analysis with implications for restoration and management. Freshwater Biology. 2017;62(3):445–457.

4. Scheuerell MD, Hilborn R, Ruckelshaus MH, Bartz KK, Lagueux KM, Haas AD, et al. The Shiraz model: a tool for incorporating anthropogenic effects and fish–habitat relationships in conservation planning. Canadian Journal of Fisheries and Aquatic Sciences. 2006;63(7):1596–1607.

5. Jorgensen JC, Honea JM, McClure MM, Cooney TD, Engie K, Holzer DM. Linking landscape-level change to habitat quality: an evaluation of restoration actions on the freshwater habitat of spring-run Chinook salmon. Freshwater Biology. 2009;54(7):1560–1575.

6. James JJ, Svejcar TJ, Rinella MJ. Demographic processes limiting seedling recruitment in arid grassland restoration. Journal of Applied Ecology. 2011;48(4):961–969.

7. Morris WF, Doak DF, et al. Quantitative conservation biology. Sinauer, Sunderland, Massachusetts, USA. 2002;.

8. Honea JM, Jorgensen JC, McClure MM, Cooney TD, Engie K, Holzer DM, et al. Evaluating habitat effects on population status: influence of habitat restoration on spring-run Chinook salmon. Freshwater Biology. 2009;54(7):1576–1592.

9. Van Horne B. Density as a misleading indicator of habitat quality. The Journal of Wildlife Management. 1983; p. 893–901.

10. Mosser A, Fryxell JM, Eberly L, Packer C. Serengeti real estate: density vs. fitness-based indicators of lion habitat quality. Ecology Letters. 2009;12(10):1050–1060.

11. Davies AB, Marneweck DG, Druce DJ, Asner GP. Den site selection, pack composition, and reproductive success in endangered African wild dogs. Behavioral Ecology. 2016;6(1):1869–1879.

12. Tarkan AS, Almeida D, Godard MJ, Gaygusuz Ö, Rylands M, Sayer CD, et al. A review and meta-analysis of growth and life-history traits of a declining European freshwater fish, crucian carp *Carassius carassius*. Aquatic Conservation: Marine and Freshwater Ecosystems. 2016;26(1):212–224.

13. Street GM, Rodgers AR, Avgar T, Fryxell JM. Characterizing demographic parameters across environmental gradients: a case study with Ontario moose (*Alces alces*). Ecosphere. 2015;6(8):1–13.

14. LaRue EA, Chambers SM, Emery NC. Eco-evolutionary dynamics in restored communities and ecosystems. Restoration Ecology. 2017;25(1):19–26.

15. Bjorndal KA, Bolten AB, Chaloupka MY. Green turtle somatic growth model: evidence for density dependence. Ecological Applications. 2000;10(1):269–282.

16. Höckendorff S, Tonkin JD, Haase P, Bunzel-Drüke M, Zimball O, Scharf M, et al. Characterizing fish responses to a river restoration over 21 years based on species' traits. Conservation Biology. 2017;31(5):1098–1108.

17. Theuerkauf SJ, Lipcius RN. Quantitative validation of a habitat suitability index for oyster restoration. Frontiers in Marine Science. 2016;3:64.

18. Railsback SF, Harvey BC, Hayse JW, LaGory KE. Tests of theory for diel variation in salmonid feeding activity and habitat use. Ecology. 2005;86(4):947–959.

19. Von Bertalanffy L. A quantitative theory of organic growth (inquiries on growth laws. II). Human biology. 1938;10(2):181–213.

20. Richards F. A flexible growth function for empirical use. Journal of Experimental Botany. 1959;10(2):290–301.

21. Schnute JT, Richards LJ. A unified approach to the analysis of fish growth, maturity, and survivorship data. Canadian Journal of Fisheries and Aquatic Sciences. 1990;47(1):24–40.

22. Birch CP. A new generalized logistic sigmoid growth equation compared with the Richards growth equation. Annals of Botany. 1999;83(6):713–723.

23. Croak BM, Webb JK, Shine R. The benefits of habitat restoration for rock-dwelling velvet geckos, *Oedura lesueurii*. Journal of Applied Ecology. 2013;50(2):432–439.

24. Oele DL, Gaeta JW, Rypel AL, McIntyre PB. Growth and recruitment dynamics of young-of-year northern pike: Implications for habitat conservation and management. Ecology of Freshwater Fish. 2019;28(2):285–301.

25. Young TP, Petersen D, Clary J. The ecology of restoration: historical links, emerging issues and unexplored realms. Ecology Letters. 2005;8(6):662–673.

26. Rosenfeld J. Assessing the habitat requirements of stream fishes: an overview and evaluation of different approaches. Transactions of the American Fisheries Society. 2003;132(5):953–968.

27. Roni P, Hanson K, Beechie T. Global review of the physical and biological effectiveness of stream habitat rehabilitation techniques. North American Journal of Fisheries Management. 2008;28(3):856–890.

28. Karkach A. Trajectories and models of individual growth. Demographic Research. 2006;15:347–400.

29. Tjørve KM, Tjørve E. Shapes and functions of bird-growth models: how to characterise chick postnatal growth. Zoology. 2010;113(6):326–333.

30. Polivka KM, Steel EA, Novak JL. Juvenile salmon and steelhead occupancy of stream pools treated and not treated with restoration structures, Entiat River, Washington. Canadian Journal of Fisheries and Aquatic Sciences. 2015;72(2):166–174.

31. Roni P, Beechie TJ, Bilby RE, Leonetti FE, Pollock MM, Pess GR. A review of stream restoration techniques and a hierarchical strategy for prioritizing restoration in Pacific Northwest watersheds. North American Journal of Fisheries Management. 2002;22(1):1–20.

32. Sivakoff FS, Morris WF, Aschehoug ET, Hudgens BR, Haddad NM. Habitat restoration alters adult butterfly morphology and potential fecundity through effects on host plant quality. Ecosphere. 2016;7(11).

33. Beechie TJ, Liermann M, Beamer EM, Henderson R. A classification of habitat types in a large river and their use by juvenile salmonids. Transactions of the American Fisheries Society. 2005;134(3):717–729.

34. Smokorowski K, Pratt T. Effect of a change in physical structure and cover on fish and fish habitat in freshwater ecosystems–a review and meta-analysis. Environmental Reviews. 2007;15:15–41.

35. Whiteway SL, Biron PM, Zimmermann A, Venter O, Grant JW. Do in-stream restoration structures enhance salmonid abundance? A meta-analysis. Canadian Journal of Fisheries and Aquatic Sciences. 2010;67(5):831–841.

36. Stranko SA, Hilderbrand RH, Palmer MA. Comparing the fish and benthic macroinvertebrate diversity of restored urban streams to reference streams. Restoration Ecology. 2012;20(6):747–755.

37. Polivka CM, Volking RA, Claeson SM, Hosman RD. Scale of monitoring influences interpretation of stream habitat restoration results for juvenile Chinook salmon. Pacific Northwest Research Station, USDA Forest Service. 2019;PNW-RN-579.

38. Polivka KM. Population ecology and effectiveness monitoring of small-scale instream habitat restoration structures in the Entiat River. In: Upper Columbia Regional Technical Team 2010 Analysis Workshop Synthesis Report. Edited by MB Ward, J. Morgan, and C. Baldwin. Prepared for the Upper Columbia Salmon Recovery Board by the Upper Columbia Regional Technical Team and Terraqua, Inc., Wenatchee, Wash., UCSRB; 2010.

39. Tiffan KF, Perry RW, Connor WP, Mullins FL, Rabe CD, Nelson DD. Survival, growth, and tag retention in age-0 Chinook Salmon implanted with 8-, 9-, and 12-mm PIT tags. North American Journal of Fisheries Management. 2015;35(4):845–852.

40. Hillman TW, Griffith J, Platts W. Summer and winter habitat selection by juvenile Chinook salmon in a highly sedimented Idaho stream. Transactions of the American Fisheries Society. 1987;116(2):185–195.

41. Grote J, Desgroseillier T. Juvenile Salmonid Out-migration Monitoring on the Entiat River, 2015. US Fish and Wildlife Service, Leavenworth, Washington. 2016;.

42. Stan Development Team. Stan Modeling Language Users Guide and Reference Manual; 2017. Available from: `http://mc-stan.org/`.

43. R Core Team. R: A Language and Environment for Statistical Computing; 2017. Available from: `https://www.R-project.org/`.

44. Stan Development Team. RStan: the R interface to Stan; 2017. Available from: `http://mc-stan.org/`.

45. Gelman A, Carlin JB, Stern HS, Dunson DB, Vehtari A, Rubin DB. Bayesian Data Analysis. 3rd ed. Chapman and Hall/CRC; 2014.

46. Link WA, Eaton MJ. On thinning of chains in MCMC. Methods in Ecology and Evolution. 2012;3(1):112–115. doi:10.1111/j.2041-210X.2011.00131.x.

47. Kruschke JK. Bayesian estimation supersedes the t test. Journal of Experimental Psychology: General. 2013;142(2):573.

48. Pfister CA, Stevens FR. The genesis of size variability in plants and animals. Ecology. 2002;83(1):59–72.

49. Elliott J. The growth rate of brown trout (*Salmo trutta* L.) fed on maximum rations. The Journal of Animal Ecology. 1975;44(3):805–821.

50. Beckman BR, Larsen DA, Lee-Pawlak B, Dickhoff WW. Relation of fish size and growth rate to migration of spring Chinook salmon smolts. North American Journal of Fisheries Management. 1998;18(3):537–546.

51. Lake PS. On the maturing of restoration: linking ecological research and restoration. Ecological Management & Restoration. 2001;2(2):110–115.

52. Schultz CB, Crone EE. Using ecological theory to develop recovery criteria for an endangered butterfly. Journal of Applied Ecology. 2015;52(5):1111–1115.

53. Elith J, Graham CH, Anderson RP, Ferrier S, Guisan A, Hijmans RJ, et al. Novel methods improve prediction of species' distributions from occurrence data. Ecography 29: 129-151. In: Peterson AT, Phillips SJ, Richardson KS, Scachetti-Pereira R, Schapire RE, Soberón J., Williams S, Wisz MS, Zimmermann NE. Citeseer; 2006.

54. Elith J, Leathwick JR. Species distribution models: ecological explanation and prediction across space and time. Annual Review of Ecology, Evolution, and Systematics. 2009;40:677–697.

55. Wood KA, Stillman RA, Hilton GM. Conservation in a changing world needs predictive models. Animal Conservation. 2018;21(2):87–88.

56. Johnson CJ, Mumma MA, St-Laurent MH. Modeling multispecies predator–prey dynamics: predicting the outcomes of conservation actions for woodland caribou. Ecosphere. 2019;10(3):e02622.

57. Dibner RR, Doak DF, Murphy M. Discrepancies in occupancy and abundance approaches to identifying and protecting habitat for an at-risk species. Ecology and Evolution. 2017;7(15):5692–5702.

58. Rosenzweig ML. Habitat selection and population interactions: the search for mechanism. The American Naturalist. 1991;137:S5–S28.

59. Polivka KM. Resource matching across habitats is limited by competition at patch scales in an estuarine-opportunist fish. Canadian Journal of Fisheries and Aquatic Sciences. 2005;62(4):913–924.

60. Falke JA, Flitcroft RL, Dunham JB, McNyset KM, Hessburg PF, Reeves GH. Climate change and vulnerability of bull trout (*Salvelinus confluentus*) in a fire-prone landscape. Canadian Journal of Fisheries and Aquatic Sciences. 2014;72(2):304–318.

61. Flitcroft RL, Falke JA, Reeves GH, Hessburg PF, McNyset KM, Benda LE. Wildfire may increase habitat quality for spring Chinook salmon in the Wenatchee River subbasin, WA, USA. Forest Ecology and Management. 2016;359:126–140.

62. MacPherson MP, Webb EB, Raedeke A, Mengel D, Nelson F. A review of Bayesian belief network models as decision-support tools for wetland conservation: Are water birds potential umbrella taxa? Biological conservation. 2018;226:215–223.

63. Everest FH, Chapman D. Habitat selection and spatial interaction by juvenile Chinook salmon and steelhead trout in two Idaho streams. Journal of the Fisheries Board of Canada. 1972;29(1):91–100.

64. Nicieza AG, Metcalfe NB. Growth compensation in juvenile Atlantic salmon: responses to depressed temperature and food availability. Ecology. 1997;78(8):2385–2400.

65. Ali M, Nicieza A, Wootton RJ. Compensatory growth in fishes: a response to growth depression. Fish and Fisheries. 2003;4(2):147–190.

66. Young KA. Asymmetric competition, habitat selection, and niche overlap in juvenile salmonids. Ecology. 2004;85(1):134–149.

67. Polivka CM, Claeson SM. Beyond redistribution: In-stream habitat restoration increases capacity for young-of-the-year salmon and trout in the Entiat River, WA. North American Journal of Fisheries Management. in review;.

68. Bartz KK, Lagueux KM, Scheuerell MD, Beechie T, Haas AD, Ruckelshaus MH. Translating restoration scenarios into habitat conditions: an initial step in evaluating recovery strategies for Chinook salmon (*Oncorhynchus tshawytscha*). Canadian Journal of Fisheries and Aquatic Sciences. 2006;63(7):1578–1595.

69. Achord S, Zabel RW, Sandford BP. Migration timing, growth, and estimated parr-to-smolt survival rates of wild Snake River spring–summer Chinook salmon from the Salmon River basin, Idaho, to the lower Snake River. Transactions of the American Fisheries Society. 2007;136(1):142–154.

70. Marco-Rius F, Caballero P, Morán P, de Leaniz CG. And the last shall be first: heterochrony and compensatory marine growth in sea trout (*Salmo trutta*). PloS ONE. 2012;7(10):e45528.

Restored: RK 4.5 - 4.9

Unrestored: RK 5.2 - 5.5