Vaccination strategies to control Ebola epidemics in the context of variable household inaccessibility levels

G. Chowell[*1], A. Tariq[1], M. Kiskowski[2]

[1]Department of Population Health Sciences, School of Public Health, Georgia State University, 30303, Atlanta, GA, USA

[2]Department of Mathematics and Statistics, University South Alabama, 36688, Mobile, AL, USA

[*]Corresponding author

Abstract

Despite a very effective vaccine, active conflict and community distrust during the ongoing DRC Ebola epidemic are undermining control efforts, including a ring vaccination strategy that require the prompt immunization of close contacts of infected individuals. However, 20% or more of close contacts cannot be reached or refuse vaccination [1], and it is predicted that the ring vaccination strategy would not be effective with such a high level of inaccessibility. In May 2019, the DRC expanded their control efforts to include a community vaccine strategy. To assess the impact of vaccination strategies for controlling Ebola epidemics in the context of variable levels of community resistance, we employed an individual-level stochastic transmission model that incorporates four sources of heterogeneity: a proportion of the population is inaccessible for contact tracing and vaccination due to lack of confidence in interventions or geographic inaccessibility, two levels of population mixing resembling household and community transmission, two types of vaccine doses with different time periods until immunity, and transmission rates that depend on spatial distance. Our results indicate that a ring vaccination strategy alone would not be effective for containing the epidemic in the context of significant delays to vaccinating contacts even for low levels of household inaccessibility and affirm the positive impact of a supplemental community vaccination strategy. Our key results are that as levels of inaccessibility increase, there is a qualitative change in the effectiveness of the vaccination strategy. For higher levels of vaccine access, the probability that the epidemic will end steadily increases over time, even if probabilities are lower than they would be otherwise with full community participation. For levels of vaccine access that are too low, however, the vaccination strategies are not expected to be successful in ending the epidemic even though they help lower incidence levels, which saves lives, makes the epidemic easier to contain and reduces spread to other communities. This qualitative change occurs for both types of vaccination strategies: ring vaccination is effective for containing an outbreak until the levels of inaccessibility exceeds approximately 10% in the context of significant delays to vaccinating contacts, a combined ring and community vaccination strategy is effective until the levels of inaccessibility exceeds approximately 50%. More broadly, our results underscore the need to enhance community engagement to public health interventions in order to enhance the effectiveness of control interventions to ensure outbreak containment.

Keywords: Mathematical modeling, individual-based model, Ebola, vaccination strategies, ring vaccination, community vaccination, community accessibility, community resistance, epidemic containment, endemic state, epidemic wave.

Author summary

Containing Ebola outbreaks hinges on educating local communities. In the context of the ongoing Ebola epidemic in DRC, active conflict and community distrust are undermining control efforts, including vaccination strategies. In this paper, we employed an individual-level stochastic structured transmission model to assess the impact of vaccination strategies on epidemic control in the context of variable levels of household inaccessibility. We found that a ring vaccination strategy would not be effective for containing the epidemic in the context of significant delays to vaccinating contacts even for low levels of household inaccessibility and evaluate the impact of a supplemental community vaccination strategy. For lower levels of inaccessibility, the probability of epidemic containment increases over time. For higher levels of inaccessibility, vaccination strategies are not expected to contain the epidemic even though they help lower incidence levels, which saves lives, makes the epidemic easier to contain and reduces spread to other communities. We found that ring vaccination is effective for containing an outbreak until the levels of inaccessibility exceeds approximately 10%, a combined ring and community vaccination strategy is effective until the levels of inaccessibility exceeds approximately 50%. Our findings underscore the need to enhance community engagement to public health interventions.

Introduction

A key lesson from past Ebola outbreaks indicates that their containment hinges on

building community trust of response teams and educating local communities in order

to fight misinformation relating to the transmission of the virus in the population and

the role of control interventions [2, 3]. These educational activities are critical for the

successful deployment of contact tracing and surveillance efforts, treatment and

management of Ebola cases, and promoting behaviors that mitigate transmission rates

[4]. However, the ongoing Ebola outbreak in the Democratic Republic of Congo (DRC),

which has been active in the region for about 10 months (August 2018 to May 2019),

has become the most complex Ebola epidemic to date, threatening to spread to

neighboring countries. In particular, this unprecedented epidemic is unfolding in a

climate of community distrust in the interventions [5, 6] in an active conflict zone [7],

where over one hundred attacks have deliberately targeted healthcare workers and

treatment centers involved in the Ebola response efforts, allowing sustained

transmission in the region. Consequently, the DRC Ebola epidemic has accumulated

2084 cases including 1405 deaths as of June 11, 2019 and has spilled over to Uganda

where 3 cases have been confirmed as of June 12, 2019 [8, 9]. A recent surge in Ebola

incidence coincided with an increasing trend in the number of violent attacks on health

centers and health teams fighting the epidemic in the region [10].

Although control efforts now employ a highly effective emergency vaccine, the

ongoing Ebola epidemic in the DRC is the first to occur in an active conflict zone.

Deliberate violent attacks and threats to health workers on a scale not seen in previous

Ebola outbreaks are directly undermining active control efforts in the region [7]. A ring

vaccination strategy [11, 12] that was highly effective in the capital [13] was

introduced rapidly on August 8[th], 2018 within 8 days of the outbreak declaration in the

provinces of North Kivu and Ituri [14]. However, it has been challenging to implement

an effective ring vaccination strategy in a climate of violence targeting healthcare

workers and decreasing accessibility to a mobile, fractionated population as it relies on

the identification of contacts and contacts of contacts [15]. This issue is compounded

by a large fraction of Ebola cases unconnected to known chains of transmission [16]. In

order to boost the impact of the vaccine in a challenging transmission setting with a

large fraction of hidden or underground transmission, the vaccination strategy is now

incorporating community-level vaccination. The community-level vaccination strategy

will immunize members of communities even if they are not known contacts of Ebola

cases using a lower vaccine dose, which will take longer to elicit protective antibody

levels in vaccinated individuals [17, 18].

Here we sought to evaluate the impact of ongoing Ebola vaccination campaigns by

extending an individual-based model with household-community mixing [19, 20],

which has been previously used to analyze the effects of lower versus higher rates of

population mixing on Ebola transmission dynamics. This model has been able to fit the

growth patterns of the 2014-15 Western Africa Ebola epidemic (e.g., the growth

pattern of the epidemic in Guinea, versus Sierra Leone or Liberia) by calibrating the

extent of community mixing [19]. This structured transmission model predicts

outbreaks that propagate through the population as spatial waves with an endemic

state [20] and has been useful to gain insight on the level of control that would be

required to contain Ebola epidemics [20]. One of the less intuitive results of the

community model was that even a low daily incidence can indicate an epidemic that is

difficult to extinguish if saturation effects decrease incidence and are masking a more

robust transmission rate. Saturation effects occur when contacts of infectious

individuals are already infected by other members of the community, such as family

members or members of other close groups, and decrease the incidence when

community mixing is low.

Assessing the effect of targeted vaccination efforts requires mathematical models that

capture the contact structure of the community network [21-25]. In our paper, we

aimed to investigate the effect of ring [21, 23, 26-28] and community vaccination

strategies on outbreak control. For this purpose, our baseline stochastic model, which

has been calibrated using data for the Western African Ebola epidemic [19, 20], was

adapted to incorporate key features of the Ebola epidemic in the DRC. Specifically, we

evaluate the effectiveness of control strategies in the context of a fraction of the

population that is accessible to vaccination teams, tied to the success of contact

tracing and vaccination efforts. Our baseline network transmission model also adapts

for the ring vaccination strategy by integrating heterogeneity in community

transmission rates that scale with the distance between an infectious individual and

each member of that individual's community. As the distance between an infectious

individual and a contact increases, the transmission rate decreases exponentially (or by

any other function). Ring and community vaccination strategies are compared by

accounting for the two different spatial scales for the radius of contacts that are

vaccinated (either as a ring or community-wide) and the two different vaccine doses (a

full or half dose), that result in different time periods for the immunization of

vaccinated contacts.

Methods

Household-Community Model for Ebola Spread and Two Types of Vaccination with

Accessible and Inaccessible Households

We previously described our model for disease spread [19, 20] and vaccination [29] in

a contact network with household-community structure. A new component of our

model is that we add a second type of household that is inaccessible to vaccination

teams. These households may not participate for a variety of reasons including Ebola

cases that are not identified, lack of confidence in the vaccination program and

geographic inaccessibility. The inaccessible households do not participate in

vaccination and do not provide contacts lists. We also modify the model to include two

vaccine types. A larger vaccine dose is used for ring vaccination with faster

immunization occurring 10 days later, and a smaller vaccine dose is used for

community vaccination with immunization occurring 28 days later. This corresponds to

the dosing and predicted timing of the ring and community vaccination regimens of 0.5

ml and 0.2 ml doses, respectively, recently described for the DRC Ebola epidemic [15,

17].

<u>**Disease and Vaccination States**</u>

The progression of Ebola disease, vaccination and immunization are modeled with

seven epidemiological states (S, E, I, R, Svr, Svc, M); including four SEIR states for Ebola

disease progression (susceptible S, exposed E, infectious I and refractory R) and three

states for vaccination progression (two susceptible but vaccinated states S_{VR} and S_{VC}

and an immune state M) (see Figure 1). The states Svc and Svr are assigned to

individuals vaccinated by community or ring vaccination. Individuals with these states

remain susceptible to Ebola exposure. After a delay of 10 days for Svr states or 28 days

for Svc states, the individuals become immune by transitioning to state M.

Individuals in three epidemiological states (S, Svc and Svr) are at risk of contracting

Ebola from infectious individuals (I), and transitioning to the latent period (exposed

state E). Transition rates from any of these three susceptible states to the exposed

state depend on the contact network and increase with the number of infectious

contacts. In our model, individuals in the exposed state E (latency) are no longer

available for effective vaccination and immunization. Thus, if an individual is indicated

to be vaccinated by either the ring or community vaccination programs, they may be

approached by a vaccination team and given the vaccine since their exposure status is

unknown, but once in the exposed class, they may not change states to the susceptible

states Svc or Svr. Once exposed, nodes transition from the exposed (E) to infectious

class (I) with probability $\frac{1}{\gamma}$ per day where $\gamma = 9$ days is the average incubation period

[30-34] and once infectious, nodes transition from the infectious (I) to refractory class

(R) with probability $\frac{1}{\lambda}$ per day, where $\lambda = 5.6$ days is the average infectious period [30-

33]. Parameter definitions and values used in simulations and their sources are given

in Table 1.

Contact Network with Household and Community Structure

As in refs. [19, 20, 29], we model the spread of Ebola on a contact network that

consists of households of size H that are organized in communities of size C

households. Households are arranged as a continuous line of households within an

LxH grid where L is the number of households in the total contact network N.

Households are indexed by their position in the line $\{h_i, h_{i+1}, ...\}$ and a network

distance η between two households h_i and h_j is defined as the number of households

separating them ($\eta=|i-j|$). Each household has its own "community" that overlaps with

the communities of other nearby households. In particular, the i_{th} community is

centered at the ith household and includes all of the households within a radius of the

i_{th} household (for a community containing C households, the community radius is

Rc=(C-1)/2). This contact network is highly mutually connected, the extent of overlap

between the i_{th} and j_{th} communities depend on the network distance of the i_{th} and j_{th}

households in that communities of nearby households overlap and share most

households. Due to this high mutual connectivity and linear arrangement of the

contact network, saturation effects build up quickly and disease progresses spatially as

1-dimensional wave through the lattice network. The size of the contact network and

the size of the simulated population is unlimited since the lattice size LxH is dynamic

and extends as needed as the disease and immunity extends through the population

by adding households to the ends of the array.

In this version of the model, each household is designated as accessible or inaccessible

as illustrated in Figure 2. This assignment is random when the household is created

and once a household is labeled, this assignment does not change. During the initial

Contact Network with Household and Community Structure

As in refs. [19, 20, 29], we model the spread of Ebola on a contact network that

consists of households of size H that are organized in communities of size C

households. Households are arranged as a continuous line of households within an

LxH grid where L is the number of households in the total contact network N.

Households are indexed by their position in the line $\{h_i, h_{i+1}, ...\}$ and a network

distance η between two households h_i and h_j is defined as the number of households

separating them ($\eta=|i-j|$). Each household has its own "community" that overlaps with

the communities of other nearby households. In particular, the i_{th} community is

centered at the ith household and includes all of the households within a radius of the

i_{th} household (for a community containing C households, the community radius is

Rc=(C-1)/2). This contact network is highly mutually connected, the extent of overlap

between the i_{th} and j_{th} communities depend on the network distance of the i_{th} and j_{th}

households in that communities of nearby households overlap and share most

households. Due to this high mutual connectivity and linear arrangement of the

contact network, saturation effects build up quickly and disease progresses spatially as

1-dimensional wave through the lattice network. The size of the contact network and

the size of the simulated population is unlimited since the lattice size LxH is dynamic

and extends as needed as the disease and immunity extends through the population

by adding households to the ends of the array.

In this version of the model, each household is designated as accessible or inaccessible

as illustrated in Figure 2. This assignment is random when the household is created

and once a household is labeled, this assignment does not change. During the initial

construction of the contact network, and as the array is dynamically extended,

households are labeled inaccessible with probability β, otherwise the household is

accessible. Here we compare results as a function of the fraction of inaccessible

households between 0 and 50%. (This range includes the estimate of 30% non-

participating individuals that was estimated for a vaccination program when it is

expanded to younger age groups in [35]; and indeed the program was approved for

expansion to younger age groups in the first half of 2019 [36] This range also includes

estimates of 25% for the number of cases that are missed for the ongoing DRC

epidemic[37].

Distance Dependent Transmission Rates

At the onset of an epidemic, before the accumulation of saturation effects when all

nodes of the network are susceptible except for a single infectious case, the household

reproductive number $R0_H$ is the average number of secondary infections within the

household of the infected individual and the community reproductive number $R0_C$ is

the average number of secondary infections within the community. As in ref. [20], we

set $R0_H$ =2.0 infections within the household and $R0_C$ =0.7 infections within the

community, based on total R_0 and how they were observed to distribute within

household and communities in historical Ebola epidemics [4, 19, 30, 38].

Transmission rates for homogenous network transmission: In the simplest case of

homogeneous transmission rates, all members of a household have an equal

probability of infection from an infectious contact, and all members of the broader

community have an equal probability of infection from an infectious contact.

Transmission rates within the household and within the community are normalized so

that the reproductive number of household and community infections are always $R0_H$

and $R0_C$, respectively. Since there are H-1 household contacts within the household of

a single infectious individual, and $(CH - H)$ community contacts within the

community of a single infectious individual, for a fixed infectious period λ, household

and community transmission rates for each infectious-susceptible contact are given by:

251
$$t_H := \frac{R_{0H}}{\lambda\,(H-1)}, \; t_C := \frac{R_{0C}}{\lambda\,(C \cdot H - H)}.$$

252 However, as described in refs. [39, 40], for exponentially distributed infectious periods

253 λ, household and community transmission rates for each infectious-susceptible contact

254 would be:

255
$$t_H := \frac{R_{0H}}{\lambda\,(H-1-R_{0H})}, \; t_C := \frac{R_{0C}}{\lambda\,(C \cdot H - H - R_{0C})}.$$

Since there are $(CH - H)$ community contacts, the total rate of community

transmission in the network (for a single infectious individual with all other nodes

susceptible) would be $(CH - H) \cdot t_C$.

<u>Transmission rates for distance-dependent network transmission</u>: In ref. [29], we

introduced distance-dependent transmission so that within the community, the

transmission rate decreases with the distance between the infectious and susceptible

individuals. For any distance function f(η), transmission rates on the network may be

scaled to yield a given set of reproductive numbers $R0_H$ and $R0_C$. This is accomplished

by assigning values α·f(η) to each community contact and choosing α so that the sum

of transmission rates (summed along every edge between an individual and every

other member of the community) is equal to $(CH - H) \cdot t_C$. Initial transmission rates

when all contacts are susceptible are $R0_H$ infections per household and $R0_C$ infections

within the community. As the number of exposed nodes accumulate, actual

transmission rates decrease due to saturation effects that depend on the radial profile

of the transmission function. Thus, the spread of an infection depends on this

transmission profile, as well as the household and community reproductive numbers

and the network structure itself.

Here we consider relatively simple distance-dependent transmission profiles in which

transmission rates decrease exponentially with distance using the function $f(\eta) =$

$e^{-k\eta}$ that we used in ref. [29] or inversely with distance using the function $g(\eta) = \frac{1}{\eta^k}$

(the intuitive "gravity model" from transportation theory [41, 42] , applied to the case

of households of equal size as we have here). The normalized transmission profiles are

shown in Figure 3. Other transmission functions that depend on distance may be more

complex exponential forms (e.g. beta-pert distributed as in ref. [43]).

283   Community and Ring Vaccination

We model two types of vaccination. Ring vaccination is the vaccination of household

members and other close contacts that are members of the community within a radius

R_{vR}. Community vaccination is the vaccination of members of the entire community.

Both types of vaccination occur on a daily basis, for flexible start and end times.

<u>Ring vaccination</u>: On the day D that an infectious individual within an accessible

household becomes infectious, the infectious individual provides a list of their close

contacts (household members and members of the community within radius R_{vR}). Ring

vaccination occurs after delay of τ days so that vaccination occurs on D+τ, with a value

that ranges from $\tau = 4$ to $\tau = 9$ days [44]. A baseline delay of $\tau = 6$ days is consistent

with an estimate of 6 days (IQR: 4–9 days) for the median time-from symptom onset to

effective isolation based on field data of the ongoing outbreak in DRC [44]. For

reference, in prior outbreaks this delay has been shorter and estimated at 2 days

based on data from the Guinea ring vaccination trial [35]. Ring vaccination begins on a

number of days Vstart after a simulation begins with one infectious case. If ring

vaccination has begun by day D, then members added to contact lists on day D are

vaccinated on day D+τ (their state changes from S to S_{VR}) if they are susceptible, not

yet vaccinated, and located within an accessible household. The radius $R_{VR}=5$

households was chosen based on the household size of H=5 to include 55 individuals.

This was based on the identification of 50 contacts per infectious case during the

Guinea ring vaccination trial [11]. In sensitivity analyses, we explored two additional

values for the radius of ring vaccination (3 and 7 households).

<u>Community vaccination</u>: If community vaccination is occurring on a given day D, all

community contacts of infectious individuals will be vaccinated with rate r_{VC} if they are

susceptible, not yet vaccinated, and located within accessible households. Vaccination

rate of 5% and 10% were studied, corresponding to a program in which most

accessible members of the community are vaccinated in 20 or 10 days.

We do not model exponentially distributed immunity rates. Rather, immunization

occurs for individuals with state S_{VR} after 10 days and for individuals with state Svc

after 28 days. It is possible that an individual is vaccinated with community vaccination

and then discovers they are a close contact of an infectious individual, where ring

vaccination is indicated. Because of this, a spike may occur when community

vaccination is initiated, due to the comparable delay of protection for these

individuals. We optimistically model that the individual can be ring vaccinated as well,

either with a supplemental or additional vaccine dose, will be immunized by whichever

vaccine would mature first.

Simulation Description, and Fitting of the Community Size C

For a typical simulation, the simulation is initiated on Day 1 with a single infectious

individual centered within a single community. With each newly exposed individual,

the contact network is dynamically extended to include the community of every

exposed individual. As the simulation is run, disease and vaccination states accumulate

in the network and the algorithm keeps track of the state of each node for each day.

Disease and/or vaccination states progress as a linear wave through the contact

network. Outbreaks die out spontaneously when all members of the contact network

are susceptible, refractory or immune.

The average growth of an epidemic per day is computed as the average number of

cases versus day for 500 simulations. The average number of cases for the nth day is

averaged only for the number of simulations that have not yet extinguished by day n.

While $R0_H$ and $R0_C$ are parameters that we consider intrinsic to the virology of Ebola,

the community size C is an abstract quantity that is meant to capture the net effect of

community interactions that describe the accumulation of saturation effects. In ref.

[19], we found that different community sizes would result in different steady state

levels for the number of cases per day, so that regions with a larger growth rate have a

larger community size. The community size C must also be calibrated for each

transmission profile. We choose the community size for each transmission profile by

assuming that an outbreak in North Kivu would be similar in size to the 2014-15 Ebola

outbreak in Liberia, which was not affected by vaccination.

Model Limitations

We have made several simplifying assumptions. We assume that the day an individual

becomes infectious is also the day when an individual becomes symptomatic. If the

individual belongs to an accessible household, they will be identified as an Ebola case

on that day. We assumed that already exposed individuals could not benefit from

vaccination before becoming infectious. However, there is evidence that vaccination

after exposure can reduce Ebola symptoms and would likely decrease transmission of

the infectious individual and for this reason in ref. [35] researchers estimated a slighter

lower time to immunization of 7 rather than 10 days for the 0.5 ml ring vaccine dose.

We assumed that the vaccines are 100% effective, and that they are effective after a

fixed rather than random time interval. Evidence indicates that the vaccine efficacy is

close to 100% [11]. By assuming a 100% vaccine effectiveness, we assume that the

effects of non-participation in vaccination or unvaccinated community contacts are a

much larger effect than a small percentage of vaccine failures.

There are many unknowns regarding the parametrization of the household community

structure. We chose a community size that would match the incidence observed in

Liberia during the West African Ebola epidemic, in the absence of vaccination, but this

community size depends on the transmission profile as a function of distance that is

unknown. The community size and the transmission profile together should be

interpreted as a summary of complex network interactions that result in an observed

incidence and are not constrained by the currently available data. We model a random

distribution of inaccessible households, whereas they may be clustered (also as in ref.

[35]). Finally, we simulate epidemics in the context of community transmission in the

absence of long-range links, so we cannot capture the effects of missed contacts that

seed outbreaks in new locations (and the initial faster growth of those outbreaks in

naïve communities).

Table 1. Parameter values used in simulations**.** Description of each model parameter,
the value or range that is used, and the reference source for the value that is used if
applicable.
377

Parameter	Description	Parameter Value (Range)	Source
H	Household Size	5	[19, 20] and [29]
C	Community Size	401	⯑
B	Fraction of households that are inaccessible.	0.0-0.6	⯑
R_{OH}	Household reproductive number	2.0	[20]
R_{OC}	Community reproductive number	0.7	[20]
Γ	Average incubation period	9 days	[34]
Λ	Average infectious period	5.6 days	[4, 31]
R_{VR}	Radius of ring vaccination (number of households)	5 (3, 7)	[11]
r_{VC}	Rate of community vaccination	0.05, 0.1	⯑
τ	Days to vaccinate an individual with the ring vaccination program	6 (4, 9)	[35, 44]
d_{VR}	Days till immunity for 0.5 ml vaccine	10	[15, 17]
d_{VC}	Days till immunity for 0.2 ml vaccine	28	[15, 17]
$Vstart_r$	Start of the ring vaccination campaign	0-9 months	
$Vstart_c$	Start of the community vaccination campaign	0-9 months	

Results

<u>Incidence and extinction with no vaccine interventions</u>

In the absence of vaccination, our calibrated model predicts that case incidence grows

during the first 10 serial intervals and then surviving epidemic realizations reach a

stable case incidence around 40-43 cases per day (Figure 4). A significant fraction of

epidemic realizations dies out; about 22-30% of the epidemic realizations

spontaneously extinguish early on, within the first 30 days of the simulation (Figure 4).

Once epidemic realizations have persisted several serial intervals, they become robust

to extinction with a very low extinction rate.

<u>Incidence and extinction with ring vaccination</u>

Our results indicate that the impact of a ring vaccination strategy depends on the

fraction of inaccessible households. Using our baseline parameter values (Table 1), in

the best-case scenario when all households are accessible (β=0), the probability of

epidemic control increases over time with a ring vaccination strategy that starts 30

days after epidemic onset (Figure 5). Since epidemic realizations are likely to end

spontaneously within the first 30 days, the effect of ring vaccination applied after 30

days shows the effect of ring vaccination on the fraction of remaining epidemics that

would otherwise likely persist. Data from the curves in Figure 5 showing this increase

in the probability of epidemic extinction over time are summarized in Figure 6. Over

12 and 18 months, the supplemental probability of epidemic containment with no

further cases of Ebola with ring vaccination for a fully accessible population is 22-32%

and 33-44%, respectively. At a 10% level of household inaccessibility, the probability

of epidemic control decreases relative to the case of full accessibility (curves in Figure 5, Panel A) but the probability that the epidemic is contained with no further cases of Ebola still slowly accumulates over time. At higher (>10%) levels of household inaccessibility, the pattern is qualitatively different. The probability of epidemic control no longer increases with the application of ring vaccination and rapidly saturates around 19-30%, just as it would without the application of ring vaccination. However, the trajectory of outbreak incidence conditional on non-extinction stabilizes around 8-22 cases per day, which is markedly lower than the case of no ring vaccination (Figure 5).

These results for vaccination applied at 30 days show that for even low fractions of inaccessible households and a significant delay to vaccinating individuals ($\tau = 6$ days), ring vaccination has little effect on ending an epidemic but substantially decreases the size of the epidemic. This pattern is robust to the timing of start of the ring vaccination program: when ring vaccination is applied to an established epidemic wave after 9 months, there is no effect on the lifetime of the epidemic if the fraction of accessible households is ≥20%, but the steady state incidence decreases (Figure 7). Ring vaccination applied after 9 months decreases steady state incidence to the same values as ring vaccination applied earlier, after 30 days (Figure 5).

Results from sensitivity analyses indicate that increasing the radius of ring vaccination from 5 households to 7 significantly enhances the probability of epidemic control when levels of household inaccessibility are below 20% (Figure 6). However, decreasing the delay to vaccinating contacts from 6 to 4 days only increases the probability of

epidemic control by about 10% for the best case scenario when all households are

accessible (Figure 6), but it does not influence the probability of epidemic control even

for low levels of household inaccessibility (as low as 10%).

Community vaccination that supplements a ring vaccination strategy can significantly

increase the probability of epidemic control.

For the scenario resembling the DRC Ebola epidemic where a ring vaccination strategy

starts one week after epidemic onset and is subsequently supplemented with

community vaccination 9 months later, we found that the community vaccination

strategy substantially increases the probability of extinction, after a delay, and can

substantially decrease the endemic state (Figure 8). However, the probability of

achieving epidemic containment again appears to not be improved when the fraction

of inaccessible household is too high (e.g., 50% inaccessible; Figure 8). Figure 9

illustrates the effects of the community vaccination rates on case incidence and the

probability of epidemic control. The supplemental effect of community vaccination on

the probability of epidemic extinction for populations with different levels of

inaccessible households ranging from 0 to 50% are summarized in Figure 10. After 1.5

years of disease transmission, our model-based results indicate that the probability of

epidemic control is at 60-71% for a 30% level of household inaccessibility and 30-41%

for a 40% household inaccessibility level based on a community vaccination rate of

10% per day.

As shown in Figures 9 and 10, supplemental community vaccination has a substantial

effect on decreasing the daily incidence of cases and increasing the rate of extinction.

While community vaccination has a near-immediate effect on decreasing case

incidence, the effect of community vaccination on the rate of extinction occurs with a

delay, from 3 months to as long as a year, that varies with the fraction of accessible

households and the daily rate of immunization (Figures 9 and 10). For the longest delay

of one year for the case of 40% inaccessible households, the increase in the rate of

extinction is striking for showing an abrupt rise after such a long delay. For all

conditions evaluated in Figures 9 and 10, the rate of extinction becomes non-negligible

and the probability of extinction begins increasing only when the case incidence drops

to a small number of cases (2-4 cases). This provides an explanation for the year-long

delay observed for the case of 40% inaccessible households : the daily incidence drops

at a slower rate so that it takes a year for the incidence to fall to a sufficiently low level

of cases. For populations where the fraction of inaccessible households is higher and

the probability of containment does not increase at all; the daily incidence decreases

but never decreases to such low levels before reaching a steady state number of cases.

Discussion

In this paper we have employed an individual-level stochastic transmission model to

evaluate ring and community vaccination strategies for containing Ebola epidemics in

the context of varying levels of community accessibility to public health interventions.

Our individual-based model, which was calibrated based on the transmission dynamics

of the 2014-15 Western African Ebola epidemic in the absence of vaccination,

incorporates three sources of heterogeneity including a proportion of the population

that is inaccessible for effective contact tracing and vaccination efforts due to lack of

confidence in interventions or geographic inaccessibility, two levels of population

mixing resembling household and community transmission, and spatial dependence on

transmission rates. Our findings indicate that ring vaccination, which targets a radius

of contacts for each infectious individual, is an effective intervention to contain Ebola

epidemics at low levels of household inaccessibility (<10%) in the presence of

significant delays to vaccinating contacts. At higher levels of household inaccessibility,

ring vaccination could be useful for reducing the disease endemicity level, but it is no

longer an effective intervention to ensure outbreak containment in the absence of any

other interventions. We also found that community vaccination strategies that

supplement a ring vaccination strategy can speed up and enhance the probability of

epidemic containment but only for moderate levels of household inaccessibility (e.g.,

<50%). For instance, our results for the scenario motivated by the DRC Ebola epidemic

predict a probability of epidemic containment at about 60-71% after 1.5 years of

disease transmission for a 30% household inaccessibility level. More generally, our

results highlight the critical need to enhance community engagement to public health

interventions while offering a safe and secure environment to the population in order

to increase the effectiveness of control interventions to ensure outbreak containment.

Active local conflict and community mistrust of the government and public health

authorities hamper containment of epidemics driven by person-to-person transmission

[5]. In this context, it is particularly important that health communication and

community education efforts are coordinated and implemented in collaboration with

the affected communities [2]. Findings from a recent survey in DRC indicate that about

25-32% of the population is skeptical about the reality of the Ebola epidemic and/or

distrust local public health authorities [45]. Importantly, there is some evidence that

Ebola outbreak in DRC is being perceived in the region as a political tool against certain

groups while some other groups benefit politically and financially from the ongoing

epidemic [46] . This underscores the need to ensure that the Ebola response is entirely

isolated from political biases to avoid exacerbating resource inequality and distrust in

the institutional authorities. Indeed, there is significant evidence that supports the

implementation of integrative communal approaches to prevent rumor spreading,

distrust and fear about Ebola in the communities and help improve epidemiological

surveillance and enhance the adoption of Ebola preventive measures [2, 47-49].

In contrast to the simple SIR compartmental transmission models based on

homogenous mixing assumptions that support bell-shaped epidemic trajectories [50,

51] [52, 53], our spatially structured stochastic model employed in this study has been

able to successfully capture stationary disease waves where the virus moves through

the host population over time. This transmission pattern is qualitatively similar to

those of the ongoing Ebola epidemic in DRC [54, 55]. In particular, they achieve

relatively steady incidence levels as Ebola spreads from one community to another,

and the total incidence can be interpreted as the super-positioning of the cases from a

discrete number of communities involved rather than explosive exponential growth.

However, our model does not predict sporadic increases in case incidence, as in the

most recent surge in case incidence [56], which is not surprising as our model does not

incorporate time-dependent changes in the effectiveness of control interventions tied

to violent attacks to healthcare workers and public health infrastructure. From an

epidemic modeling perspective, our results underscore the need to capture an

appropriate spatial structure in models of disease transmission [19, 23-25, 57, 58].

Such considerations may be more important for infectious diseases that are

transmitted via close contact such as Ebola and HIV.

In summary, we found that the role of vaccination strategies in containing Ebola

epidemics significantly depends on the level of community inaccessibility using an

individual-level stochastic transmission model that successfully captures stationary

disease waves that are qualitatively similar to those of the ongoing Ebola epidemic in

DRC. For lower levels of inaccessibility, the probability of containment increases over

time. For higher levels, vaccination strategies alone are not expected to contain the

epidemic, but they help reduce incidence levels, which saves lives, makes the epidemic

easier to contain and reduces spread to other communities. This qualitative change

occurs for both types of vaccination strategies: ring vaccination is effective for

containing an outbreak until the level of inaccessibility exceeds approximately 20%, a

combined ring and community vaccination strategy is effective until the level of

inaccessibility exceeds approximately 50%. In order to enhance the effectiveness of

control interventions, it is crucial to ensure community engagement to public health

interventions.

Conflict of interest

551   The authors declare no conflicts of interest.

References

1.   Cohen, J., *Ebola outbreak continues despite powerful vaccine. Available from:* *https://science.sciencemag.org/content/364/6437/223.* Science, 2019.
2.   Marais, F., et al., *A community-engaged infection prevention and control approach to Ebola.* Health Promot Int, 2016. **31**(2): p. 440-9.
3.   Nyenswah, T.G., et al., *Ebola and Its Control in Liberia, 2014-2015.* Emerg Infect Dis, 2016. **22**(2): p. 169-77.
4.   Chowell, G. and H. Nishiura, *Transmission dynamics and control of Ebola virus disease (EVD): a review.* BMC Med, 2014. **12**(1): p. 196.
5.   Vinck, P., et al., *Institutional trust and misinformation in the response to the 2018-19 Ebola outbreak in North Kivu, DR Congo: a population-based survey.* Lancet Infect Dis, 2019. **19**(5): p. 529-536.
6.   Nguyen, V.K., *An Epidemic of Suspicion - Ebola and Violence in the DRC.* N Engl J Med, 2019. **380**(14): p. 1298-1299.
7.   Claude, K.M., J. Underschultz, and M.T. Hawkes, *Ebola virus epidemic in war-torn eastern DR Congo.* Lancet, 2018. **392**(10156): p. 1399-1401.
8.   *WHO, Ebola Virus Disease, Democratic Republic of the Congo, External Situation Report 45. 2019: World Health Organization. Available from: https://apps.who.int/iris/bitstream/handle/10665/325242/SITREP_EVD_DRC_UGA_20190612-eng.pdf?ua=1 (accessed on 12 June 2019).*
9.   *WHO. Ebola virus disease – Democratic Republic of the Congo Disease outbreak news. 2019 June 14, 2019]; Available from: https://www.who.int/csr/don/13-june-2019-ebola-drc/en/. (accessed on 15 June 2019).*
10. *Kivu Security Tracker. Available from: https://kivusecurity.org/map#.*
11. Henao-Restrepo, A.M., et al., *Efficacy and effectiveness of an rVSV-vectored vaccine expressing Ebola surface glycoprotein: interim results from the Guinea ring vaccination cluster-randomised trial.* Lancet, 2015. **386**(9996): p. 857-66.
12. Merler, S., et al., *Containing Ebola at the Source with Ring Vaccination.* PLoS Negl Trop Dis, 2016. **10**(11): p. e0005093.
13. Wells, C.R., et al., *Ebola vaccination in the Democratic Republic of the Congo.* Proc Natl Acad Sci U S A, 2019. **116**(20): p. 10178-10183.

588   14.   *WHO. Ebola virus disease – Democratic Republic of the Congo. 2018 [cited*
589   *2018 August 19]; Available from: https://www.who.int/csr/don/17-august-2018-*
590   *ebola-drc/en/.*

591   15.   *The report of the SAGE April 2019 will be published in the WHO Weekly*
592   *Epidemiological Record (www.who.int/wer/en/) on 30 May 2019.*

593   16.   *Maxmen, A. Ebola detectives race to identify hidden sources of infection as*
594   *outbreak spreads. Available from: https://www.nature.com/articles/d41586-018-*
595   *07618-0 (accessed on 23 May 2019).*

596   17.   Cohen, J., *DRC expands Ebola vaccine campaign as cases mount rapidly.*
597   Science.

598   18.   *World Health Organization. Interim Recommendations on Vaccination against*
599   *Ebola Virus Disease (EVD). Available from:*
600   *https://www.who.int/immunization/policy/position_papers/interim_ebola_recom*
601   *mendations_may_2019.pdf?ua=1 (accessed on 20 May 2019).*

602   19.   Kiskowski, M., *Three-Scale Network Model for the Early Growth Dynamics of*
603   *2014 West Africa Ebola Epidemic. .* PLOS Currents Outbreaks, 2014. **doi:**
604   **10.1371/currents.outbreaks.b4690859d91684da963dc40e00f3da81.**

605   20.   Kiskowski, M. and G. Chowell, *Modeling household and community*
606   *transmission of Ebola virus disease: epidemic growth, spatial dynamics and*
607   *insights for epidemic control.* Virulence, 2015. **7**(2): p. 63-73.

608   21.   Tildesley, M.J., et al., *Optimal reactive vaccination strategies for a foot-and-*
609   *mouth outbreak in the UK.* Nature, 2006. **440**(7080): p. 83-6.

610   22.   Ferguson, N.M., C.A. Donnelly, and R.M. Anderson, *The foot-and-mouth*
611   *epidemic in Great Britain: pattern of spread and impact of interventions.*
612   Science, 2001. **292**(5519): p. 1155-60.

613   23.   Browne, C., H. Gulbudak, and G. Webb, *Modeling contact tracing in outbreaks*
614   *with application to Ebola.* J Theor Biol, 2015. **384**: p. 33-49.

615   24.   Lau, M.S.Y., et al., *A mechanistic spatio-temporal framework for modelling*
616   *individual-to-individual transmission-With an application to the 2014-2015*
617   *West Africa Ebola outbreak.* PLoS Comput Biol, 2017. **13**(10): p. e1005798.

618   25.   Yang, W., et al., *Transmission network of the 2014-2015 Ebola epidemic in*
619   *Sierra Leone.* J R Soc Interface, 2015. **12**(112).

620   26.   Greenhalgh, D., *Optimal control of an epidemic by ring vaccination.*
621   Communications in Statistics. Stochastic Models, 1986. **2**(3): p. 339--363.

622   27.   Muller, J., B. Schonfisch, and M. Kirkilionis, *Ring vaccination.* J Math Biol,
623   2000. **41**(2): p. 143-71.

28. Kretzschmar, M., et al., *Ring vaccination and smallpox control.* Emerg Infect
Dis, 2004. **10**(5): p. 832-41.
29. Chowell, G. and M. Kiskowski, *Modeling ring-vaccination strategies to control*
Ebola virus disease epidemics, in *Mathematical modelling for emerging and*
reemerging infectious diseases, G. Chowell and J.M. Hyman, Editors. 2016,
Springer.
30. Chowell, G., et al., *The basic reproductive number of Ebola and the effects of*
public health measures: the cases of Congo and Uganda. J Theor Biol, 2004.
229(1): p. 119-26.
31. Legrand, J., et al., *Understanding the dynamics of Ebola epidemics.* Epidemiol
Infect, 2007. **135**(4): p. 610-21.
32. Lekone, P.E. and B.F. Finkenstadt, *Statistical inference in a stochastic epidemic*
SEIR model with control intervention: Ebola as a case study. Biometrics, 2006.
62(4): p. 1170-7.
33. Eichner, M., S.F. Dowell, and N. Firese, *Incubation period of ebola*
hemorrhagic virus subtype zaire. Osong Public Health Res Perspect, 2011. **2**(1):
p. 3-7.
34. Team, W.H.O.E.R., *Ebola Virus Disease in West Africa - The First 9 Months of*
the Epidemic and Forward Projections. N Engl J Med, 2014. **371**(16): p. 1481-
95.
35. Kucharski, A.J., et al., *Effectiveness of Ring Vaccination as Control Strategy for*
Ebola Virus Disease. Emerg Infect Dis, 2016. **22**(1): p. 105-8.
36. *ReliefWeb Report. UNICEF DR Congo Ebola Situation Report North Kivu and*
Ituri - 03 February 2019. Available from:
https://reliefweb.int/report/democratic-republic-congo/unicef-dr-congo-ebola-
situation-report-north-kivu-and-ituri-03.
37. *Branswell, H. WHO sees progress in Ebola response, but others see a grimmer*
reality. StatNews. Available from: https://www.statnews.com/2019/06/06/who-
sees-progress-in-ebola-response-but-others-see-a-grimmer-reality/ (accessed on
06/06/2019). 2019.
38. Althaus, C.L., *Estimating the reproduction number of Zaire ebolavirus (EBOV)*
during the 2014 outbreak in West Africa. PLOS Currents Outbreaks. Edition 1.
doi: 10.1371/currents.outbreaks.91afb5e0f279e7f29e7056095255b288., 2014.
39. Lloyd AL, V.S., Cintrón-Arias A., *Infection dynamics on small-world networks.*
Contemp Math. , 2006. **410**: p. 209.

40. Keeling, M.J. and B.T. Grenfell, *Individual-based perspectives on R(0).* J Theor
Biol, 2000. **203**(1): p. 51-61.

41. Xia, Y.C., O.N. Bjørnstad, and B.T. Grenfell, *Measles metapopulation*
dynamics: a gravity model for epidemiological coupling and dynamics. Am Nat
2004. **164**: p. 267-81.

42. Viboud, C., et al., *Synchrony, waves, and spatial hierarchies in the spread of*
influenza. Science, 2006. **312**(5772): p. 447-51.

43. Durr, S. and M.P. Ward, *Development of a Novel Rabies Simulation Model for*
Application in a Non-endemic Environment. PLoS Negl Trop Dis, 2015. **9**(6): p.
e0003876.

44. *WHO Outbreak News: 06/06/2019. Ebola virus disease – Democratic Republic*
of the Congo. Available from: [https://www.who.int/csr/don/06-june-2019-ebola-](https://www.who.int/csr/don/06-june-2019-ebola-drc/en/)
[drc/en/](https://www.who.int/csr/don/06-june-2019-ebola-drc/en/) *(accessed on 06/15/2019).*

45. Schnirring, L. *As Ebola rages in DRC, survey highlights wide mistrust, in*
CIDRAP News.Available from: [http://www.cidrap.umn.edu/news-](http://www.cidrap.umn.edu/news-perspective/2019/03/ebola-rages-drc-survey-highlights-wide-mistrust)
[perspective/2019/03/ebola-rages-drc-survey-highlights-wide-mistrust](http://www.cidrap.umn.edu/news-perspective/2019/03/ebola-rages-drc-survey-highlights-wide-mistrust) *(accessed*
on 05 May 2019).

46. Aizenman, N. *Why The Ebola Outbreak In The Democratic Republic Of Congo*
Keeps Getting Worse. NPR News. Available from:
[https://www.npr.org/2019/05/23/725789837/why-the-ebola-outbreak-in-the-](https://www.npr.org/2019/05/23/725789837/why-the-ebola-outbreak-in-the-democratic-republic-of-congo-keeps-getting-worse)
[democratic-republic-of-congo-keeps-getting-worse](https://www.npr.org/2019/05/23/725789837/why-the-ebola-outbreak-in-the-democratic-republic-of-congo-keeps-getting-worse) *(accessed on 23 May 2019).*

47. Tambo, E., E.C. Ugwu, and J.Y. Ngogang, *Need of surveillance response*
systems to combat Ebola outbreaks and other emerging infectious diseases in
African countries. Infect Dis Poverty, 2014. **3**: p. 29.

48. Thiam, S., et al., *Challenges in controlling the Ebola outbreak in two*
prefectures in Guinea: why did communities continue to resist? Pan Afr Med J,
2015. **22 Suppl 1**: p. 22.

49. Buseh, A.G., et al., *The Ebola epidemic in West Africa: challenges,*
opportunities, and policy priority areas. Nurs Outlook, 2015. **63**(1): p. 30-40.

50. Anderson, R.M. and R.M. May, *Infectious diseases of humans.* 1991, Oxford:
Oxford University Press.

51. Hethcote, H.W., *The mathematics of infectious diseases.* SIAM review, 2000.
42(4): p. 599-653.

52. Brauer, F., *Some simple epidemic models.* Math Biosci Eng, 2006. **3**(1): p. 1-15.

53. Kermack, W.O. and A.G. McKendrick, *Contributions to the mathematical*
theory of epidemics: IV. Analysis of experimental epidemics of the virus disease
mouse ectromelia. J Hyg (Lond), 1937. **37**(2): p. 172-87.
54. *Ebola virus disease – Democratic Republic of the Congo. Disease outbreak*
news: Update for 23 April 2019. Available from:
https://www.who.int/csr/don/23-may-2019-ebola-drc/en/ (accessed on 24 May
2019).
55. Tariq, A., et al., *Assessing reporting delays and the effective reproduction*
number: The Ebola epidemic in DRC, May 2018-January 2019. Epidemics,
2019. **26**: p. 128-133.
56. *WHO. (2019). Ebola Virus Disease, Democratic Republic of Congo, External*
Situation Reports. April 24, 2019.   Retrieved April 24, 2019. Available
from:https://apps.who.int/iris/bitstream/handle/10665/312080/SITREP-EVD-
DRC-20192404-eng.pdf?ua=1.
57. Sattenspiel, L. and K. Dietz, *A structured epidemic model incorporating*
geographic mobility among regions. Math Biosci, 1995. **128**(1-2): p. 71-91.
58. Merler, S., et al., *Spatiotemporal spread of the 2014 outbreak of Ebola virus*
disease in Liberia and the effectiveness of non-pharmaceutical interventions: a
computational modelling analysis. Lancet Infect Dis, 2015. **15**(2): p. 204-11.
712
713

Figures

Figure 1. The seven epidemiological states for individuals in the population and the
transitions between states. Susceptible (S) and individuals vaccinated with either the
ring or community dose of vaccine (Svr, SVc) may become exposed (E). Once exposed,
individuals eventually become infectious and then refractory. Vaccinated individuals
that do not become exposed eventually become immunized (M).

S	Susceptible to Ebola
Sv_c	Susceptible to Ebola but vaccinated with community dose of the vaccine
Sv_r	Susceptible to Ebola but vaccinated with ring dose of the vaccine
E	Exposed to Ebola
I	Infectious
R	Refractory after an Ebola infection
M	Immunized by the vaccine and no longer susceptible to Ebola

722
723
724
725

Figure 2. Description of the contact network. A) The contact network is initialized with
a single infectious individual (red node) and that individual's community. For this
illustrative schematic, the household size H=5 and the community size is small with
C=25. The accessibility (white nodes) or inaccessibility (gray nodes) of a household is
assigned randomly during the construction of the contact network. The index case
happens to be in an inaccessible household: the first case of the simulation does not
supply a contact list, and no contacts are vaccinated. B) After several timesteps, the
infectious index case exposes another member of their household (blue node in the
h_{ith} household) and a member of their community (blue node at a distance of 8
households). When the community member is exposed in the h_{i+8th} household, the
total contact network is extended by 8 households to include their community (the
h_{i+8th} community is outlined with a black rectangle). C) The h_{i+8th} household is
accessible, so once the exposed individual becomes infectious a contact list is provided
(green nodes) of households within a radius Rv=2. One household within this radius is
excluded since it is inaccessible. The members of this household are not included on
contact lists, do not participate in vaccination, and/or are not geographically accessible
to the vaccination teams, etc.

744
745
746
747

Figure 3: Community transmission profiles and steady states in the absence of
vaccination. A) Transmission rates as a function of distance from an infectious
individual for an inverse transmission profile (left) and exponential transmission profile
(right) with community reproductive number R_{0C}=0.7 for both profiles. The inverse
distance function is $g(\eta) = \frac{1}{\eta^{0.93}}$ and the exponential distance function is $f(\eta) =$
$e^{-0.02\eta}$ for the community size C=401. B). For this community size and transmission
profile parameters, a steady state incidence comparable to that of the 2014-15 Ebola
epidemic in Liberia is achieved when simulating epidemics in the absence of
vaccination (see <mark>Figure 4</mark>).
757

758
759

Figure 4. Probability of epidemic extinction and the baseline mean daily incidence
curves in the absence of interventions for the exponential and gravity transmission
profiles that define transmission rates as a function of the distance from an infectious
individual (see also Figure 3). For each transmission profile, the model was calibrated
by choosing the community size for each transmission profile by assuming that the
ongoing Ebola outbreak in North Kivu would be similar in size to the 2014-15 Ebola
outbreak in Liberia in the absence of vaccination. Outbreaks propagate through the
population as spatial waves with an endemic state. The average number of cases for
the n_{th} day is averaged only for the number of simulations that have not yet
extinguished by day n. The error bars show the 95% confidence intervals for the mean
daily incidence.

772
773
774

 Figure 5. The impact of ring vaccination that starts 30 days after epidemic onset on

the probability of outbreak extinction and the mean daily incidence curves for various

percentage levels of inaccessible households, which do not participate in vaccination

and do not provide contacts lists as explained in the text. These simulations were

generated using the exponential transmission profile, but similar results were obtained

using the gravity transmission profile described in the main text. The vertical dashed

line indicates the timing of start of the ring vaccination program.

Figure 6. The probability of epidemic control for a ring vaccination strategy as a

function of the radius of the ring and the delay to vaccinating contacts. The baseline

parameter value for the radius of ring vaccination is the intermediate value of $R_{VR}=5$,

and the baseline parameter value for the delay to vaccination is the intermediate value

of $\tau=6$ days. Baseline parameter values for other parameters not varied in this figure

are given in Table 1.

Figure 7. The impact of ring vaccination on an established epidemic wave (after a delay

of 270 days) on the probability of outbreak extinction and the mean daily incidence

curves for various percentage levels of household inaccessibility. These simulations

were generated using the exponential transmission profile, but similar results were

obtained using the gravity transmission profile described in the main text. The vertical

dashed line indicates the timing of start of the ring vaccination program.

Figure 8. Mean daily case incidence when a community vaccination rate of 10% per

day supplements a ring vaccination strategy 9 months later for different levels of

household inaccessibility. The vertical dashed line indicates the timing of start of the

supplemental community vaccination efforts. These simulations were generated using

the exponential transmission profile, but similar results were obtained using the

gravity transmission profile described in the main text.

Figure 9. Probability of epidemic extinction over time and the mean daily case

incidence when different community vaccination rates supplement a ring vaccination

strategy for a 30% level of household inaccessibility and community vaccination starts

9 months after epidemic onset. The vertical dashed line indicates the timing of start of

community vaccination. These simulations were generated using the exponential

transmission profile, but similar results were obtained using the gravity transmission

profile described in the main text.

Figure 10. Supplemental effect on community vaccination on the probability of

epidemic extinction for populations with different household inaccessibility levels

ranging from 0 to 50%. Bars show the probability of epidemic extinction after 1.5 years

for increasing vaccination measures: i) ring vaccination alone (blue bars), ii) ring

vaccination with supplemental community vaccination with a rate of 5% community

vaccination per day (light blue bars) and iii) ring vaccination with supplemental

community vaccination with a rate of 10% vaccination per day. The horizontal line

indicates the mean probability of extinction for the baseline scenario in the absence of

vaccination. Timing was chosen to resemble the scenario for the DRC Ebola epidemic,

with ring vaccination applied after 7 days and community vaccination applied after 9

months. These simulations were generated using the exponential transmission profile,

but qualitatively similar results were obtained using the gravity transmission profile

described in the main text.