The rapid divergence of the Antarctic crinoid species *Promachocrinus kerguelensis*

Running title: Phylogeography of Antarctic crinoids

Yacine Ben Chehida[1,2*], Marc Eléaume[1], Cyril Gallut[1], Guillaume Achaz[1,3]

[1] Institut de Systématique, Évolution, Biodiversité (ISYEB), Muséum national d'Histoire naturelle, CNRS, Sorbonne Université, EPHE, Université des Antilles. 57 rue Cuvier, CP 50, 75005, Paris, France;
[2] University of Groningen: Groningen Institute for Evolutionary Life Sciences (GELIFES), University of Groningen, PO Box 11103 CC, Groningen, The Netherlands;
[3] Stochastic Models for the Inference of Life Evolution, CIRB (UMR 7241 CNRS), Collège de France, Paris, France.

* **Correspondence**: Yacine Ben Chehida (h.y.ben.chehida@rug.nl). University of Groningen: Groningen Institute for Evolutionary Life Sciences (GELIFES), University of Groningen, PO Box 11103 CC, Groningen, The Netherlands.

Keywords: Crinoids; species delimitation; Southern Ocean; cryptic species; *Florometra mawsoni*; speciation; *Promachocrinus kergulensis*; "apparent" drift; recent divergence

Abstract

Climatic oscillations in Antarctica caused a succession of expansion and reduction of the ice sheets covering the continental shelf of the Southern Ocean. For marine invertebrates, these successions are suspected to have driven allopatric speciation, endemism and the prevalence of cryptic species, leading to the so-called Antarctic 'biodiversity pump' hypothesis. Here we took advantage of the recent sampling effort influenced by the International Polar Year (2007-8) to test for the validity of this hypothesis for 1,797 samples of two recognized crinoid species: *Promachocrinus kerguelensis* and *Florometra mawsoni*. Species delimitation analysis identified seven phylogroups. As previously suggested, *Promachocrinus kerguelensis* forms a complex of six cryptic species. Conversely, despite the morphological differences, our results show that *Florometra mawsoni* is a lineage nested within *Promachocrinus kerguelensis*. It suggests that *Florometra mawsoni* and *Promachocrinus kerguelensis* belong to the same complex of species. Furthermore, this study indicates that over time and space the different sectors of the Southern Ocean show a remarkable rapid turn-over in term of phylogroups composition and also of genetic variants within phylogroups. We argue that strong "apparent" genetic drift causes this rapid genetic turn-over. Finally, we dated the last common ancestor of all phylogroups at less than 1,000 years, raising doubts on the relevance of the Antarctic "biodiversity pump" for this complex of species. This work is a first step towards a better understanding of how life is diversifying in the Southern Ocean.

Introduction

During the last decade following the last International Polar Year (2007-8) a huge effort has been made to better understand Antarctic marine ecosystems and one of the major outcomes has been the creation of a reference barcode database (BOLD) that reconciles morphologically recognized species with a unique COI fragment. This opened to a new era of discoveries that allowed reassess some of the main known characteristics of marine Antarctic ecosystems, including eurybathy, excess in brooding species, high rate of endemism, rapid diversification and cryptic speciation as a consequence of the biodiversity pump (Clarke & Crame, 1989). Recent biodiversity assessments using molecular tools have revealed an increasing number of cryptic species in teleost fishes as well as Echinodermata, Mollusca, Arthropoda, or Annelida (see (Cornils & Held, 2014; Griffiths, 2010; Rogers, 2007) for comprehensive reviews). Cryptic species have been shown to be homogeneously distributed among taxa and biogeographic regions (Pfenninger & Schwenk, 2007) and Antarctica may well be no exception.

The Southern Ocean is known to have undergone several glaciation events (Clarke & Crame, 1989; Clarke, Crame, Stromberg, & Barker, 1992; Thatje, Hillenbrand, Mackensen, & Larter, 2008). Massive ice sheet advance and retreat seem to have bulldozed the entire continental shelf several times during the last 25 Mya. These events are thought to be driven by Milankovitch cycles. These cycles may be among the strongest evolutionary forces that have shaped Antarctic terrestrial and marine biodiversity (Clarke et al., 1992; Thatje et al., 2008; Thatje, Hillenbrand, & Larter, 2005). Thatje *et al.* (2005; 2008) hypothesized that vicariant speciation could have occurred during the glacial periods on the Antarctic continental shelf, within multiple ice-free refugia like polynya. As lineages evolved independently during the glacial period, they accumulated genetic differences that lead to reproductive isolation and probably to the formation of cryptic species. During interglacial periods, barriers to gene flow may have been removed allowing for secondary contact between the vicarious lineages (Heimeier, Lavery, & Sewell, 2010; Thatje et al., 2005; 2008; Thornhill, Mahon, Norenburg, & Halanych, 2008). An alternative hypothesis is based on the idea that Antarctic continental shelf may be understood as a species flock generator (Eastman & McCune, 2000; Lecointre et al., 2013).

Here, we have analyzed COI sequences of 1,797 individuals of two crinoid species:

Promachocrinus kerguelensis (Carpenter, 1888) and *Florometra mawsoni* (AH Clark, 1937),

endemic to the Southern Ocean. These species represent the most abundant crinoid species in

the Southern Ocean (Eléaume, Hemery, Roux, & Améziane, 2014; Speel & Dearborn, 1983).

They are morphologically distinct (but see (Eléaume, 2006) for counter arguments) and

genetically close (Hemery, Améziane, & Eléaume, 2013a). These species are thought to have

a reproduction cycle that involves external fertilization that could result in a large dispersal

potential. *P. kerguelensis* produces positively buoyant ovocytes that, after fertilization, may

remain in the plankton for weeks or even months (McClintock & Pearse, 1987). Some adults

of *P. kerguelensis* have also been observed swimming (Eléaume, unpublished observations),

and some adults of *F. mawsoni*, though never observed swimming in situ, have shown this

ability in tanks (Eléaume, unpublished observations).

During the last decade, numerous specimens of crinoids have been collected from all around

Antarctica and sub-Antarctic islands. Over 3,000 specimens attributed to 45 species have been

sampled (Eléaume et al., 2014). Species such as *Promachocrinus kerguelensis* and

Florometra mawsoni are represented by a large number of well distributed specimens.

Knowlton (1993; 2000) predicted for Southern Ocean organisms that an increase in sampling

effort and the application of genetic tools, would reveal cryptic species. The analysis of

P. kerguelensis COI barcode fragment suggested that this species may constitute such an

example of a complex of unrecognized species. Wilson *et al.* (2007) identified six lineages in

the Atlantic sector whereas Hemery *et al.* (2012) extending the analysis of Wilson to the

entire Southern Ocean, identified seven lineages. As no obvious morphological character have

been shown to distinguish between these lineages (Eléaume, 2006), they may represent true

cryptic species. However, Hemery *et al.* (2012) using nuclear markers have shown that the six

identified COI lineages may only represent three distinct entities. All of the six or seven

lineages are circumpolar in distribution, sympatric and eurybathic but show various levels of

connectivity that depend on the lineage and the geographical area.

Here we analyze the pooled COI datasets of *P. kerguelensis* and *F. mawsoni* of Wilson *et al.*

(2007) and Hemery *et al.* (2012) using different approaches to separate different sources of

genetic variation, i.e. differences due to diversification (clade), time (year collected), space

(geographical origin of samples). We first analyze the phylogenetic relationships within

P. kerguelensis sensu largo using different species delineation methods. We then measure the

genetic turn-over, within each phylogroup sampled at a given location. Using a population

genetics coalescent framework, we estimated an extraordinary small effective population size

that corresponds to a mutation rate that is larger than previously reported before. We then

discuss our results in the light of the climate history of the Southern Ocean.

Materials and methods

Sequences

We included 13 sequences of *P. kerguelensis* from Wilson *et al.* (2007), for which we had precise information about the place and the date of sampling. We have also included 1303 sequences of *P. kerguelensis* from Hemery *et al.* (2012) together with 479 sequences of *F. mawsoni* (Hemery *et al.,* in preparation). A total of 1,797 COI sequences were used for this study. All specimens were collected between 1996 and 2010 in seven geographical regions that were described in Hemery *et al.* (Hemery et al., 2012): Kerguelen Plateau (KP), Davis Sea (DS), Terre Adélie (TA), Ross Sea (RS), Amundsen Sea (AS), West Antarctic Peninsula (WAP), East Weddell Sea (EWS). We however further split the Scotia Arc into Scotia Arc East (SAE) and Scotia Arc West (SAW) as there can be as much as 2,000 km between them. The counts of all sequences of all locations are reported in Table S1.

DNA extraction, PCR and sequencing

For this analysis, no additional sequences have been produced. For DNA extraction and PCR procedures see (Hemery et al., 2012; Wilson et al., 2007) and 2013. A total of 1797 554-bp sequences of the barcode region of cytochrome c oxidase subunit I (COI) were amplified using the Folmer *et al.* (1994) primers and other specific primers described in Hemery *et al*. (2012). All COI sequences from Hemery *et al.* (2012) have been made easily available through a data paper article (Hemery et al., 2013a).

Species delimitation

We used three independent species delimitation methods: Generalized Mixed Yule Coalescent (GYMC, (Pons et al., 2006)), Automatic Barcode Gap Discovery (ABGD, (Puillandre, Lambert, Brouillet, & Achaz, 2011)) and Poisson Tree Process (PTP, (Zhang, Kapli, Pavlidis, & Stamatakis, 2013)) to estimate the number of phylogroups in our sample. For all these methods, we used all the 199 unique haplotypes of the dataset to have legible results and a faster computation time. The phylogenetic tree used for PTP was constructed by Neighbor Joining method the BioNJ implementation (Gascuel, 1997) in *Seaview* software (Gouy, Guindon, & Gascuel, 2010) version 3.2. PTP analysis was conducted online (http://species.h-its.org/ptp/). The ultrametric tree needed for GYMC method was constructed using *BEAST* software (Bayesian Evolutionary Analysis Sampling Trees, (Drummond, Suchard, Xie, &

151 Rambaut, 2012) version 1.7). GYMC analysis was conducted online (http://species.h-
152 its.org/gmyc/). ABGD analysis was conducted online
153 (http://wwwabi.snv.jussieu.fr/public/abgd/abgdweb.html).

Assessing the temporal structure within phylogroups

155 We characterized the temporal structures of our samples in pairwise comparisons using both
156 fixation index F_{ST} as well as a non-parametric structure test based on $K*s$ by Hudson *et al.*
157 (1992). All p-values were estimated using permutations as described in Hudson *et al.* (1992).
158 A web-interface for the latter can be found at http://wwwabi.snv.jussieu.fr/public/mpweb/.

Estimation of the mutation rate and the population effective size

As described by Fu (2001) one can use temporal data to estimate the mutation rate (μ). In
summary, the method assumes that the population is sampled in at least two time points (say
T_1 and then later T_2) with multiple sequences in T_1. The expected average pairwise differences
between sequences taken from the two time points (K_{12}) equals the average pairwise
differences of sequences within the past time point (K_{11}) plus the difference accumulated
since then. Mathematically, it is expressed as $E[K_{12}] = E[K_{11}] + (T_2-T_1)\mu$. Because the time
difference between the samples (T_2-T_1) is known, the mutation rate can be estimated from the
observed pairwise differences: $\mu = (K_{12}-K_{11})/(T_2-T_1)$. This idea can be expanded when more
data are available in other time points (Fu, 2001).

Once μ is known, the average pairwise difference can be used to estimate the effective
population size (Ne). Under the standard neutral model, the pairwise difference within groups
of the same sample equals $2Ne\mu$, when μ is expressed as a rate by generations. Here, we
assumed one generation every three year to estimate Ne (Clark, 1921).

Estimation of the time of divergence between phylogroups

174 For visualization purposes, an ultrametric tree was constructed with the unique sequences by
175 UPMGA (Michener & Sokal, 1957) as implemented in the *Phylip* package (Felsenstein, 1989)
176 version 3.6. Using the estimated mutation rate, we derived a time of divergence within and
177 between phylogroups from the average sequence differences.

Polymorphisms analysis

We used *Dnasp* software version 5.10 (Librado & Rozas, 2009) and the program used to study genetic structure to study the SFS (Site Frequency Spectrum) within phylogroups as well as two of its summary statistics: nucleotide diversity (π) and number of polymorphic sites (S). We then tested neutrality using Tajima's D (Tajima, 1989), Fu and Li's F* and D* (Fu & Li, 1993), Achaz's Y* (Achaz, 2008), that is immune to sequencing errors, and the Strobeck (Strobeck, 1987) haplotype test.

Assessment of migration effect on Ne estimation through simulations

To assess the potential effect of migrants from a distant population on the estimation of μ and *Ne*, we used coalescent simulations where two samples of 20 sequences each are taken 3N generations apart in a focal population of size *N*. We assume a large unsampled ghost population of size *10N* that exchange migrants with the focal one at rate *M=4Nm*, where *m* is the migration rate per generation. For values of *M* ranging from 0.001 (very low) to 1000 (very large), we applied the Fu (Fu, 2001) method to estimate *Ne*, with 10^3 replicates for each *M* value. The program is based on a C++ library developed by G. Achaz that is available upon request.

Results

P. kerguelensis and F. mawsoni are composed of seven phylogroups

We first reinvestigated the species delimitation of the sample using three independent approaches (GMYC, ABGD and PTP) that cluster the unique sequences by phylogroup. We adopt in this manuscript a nominalist definition of species. However, *F. mawsoni*, a nominal species, is nested within *P. kerguelensis*, another nominal species, lineages. To avoid confusion, we have decided to use the term "phylogroup" in the rest of the manuscript to designate any well-supported cluster of sequences derived from species delineation approaches.

Two of the three methods estimate a similar number of phylogroups: PTP estimates that there are nine phylogroups in the sample whereas ABGD reports 5-7 phylogroups, depending on the chosen prior value. Looking at the proposed partitions, we have chosen to consider seven different phylogroups for this study that are reported as phylogroups A-G in Fig. 1. Interestingly, these phylogroups correspond to the A-F groups proposed by Wilson *et al.* (2007), plus a 'new' group (G) that corresponds to the *F. mawsoni* species. The two extra sub-splits suggested by the PTP method are indicated as C1/C2 and E1/E2 on Fig. 1. Note that PTP only seeks monophyletic groups and thus must split the C group into the C1/C2 subgroups.

In contrast with the two previous methods, depending on the sequence evolution model we used and the *BEAST* parameters, GMYC reports between 3 and 51 phylogroups with non-overlapping grouping between individuals. As GMYC is very sensitive in our case to the chosen parameters, it is here unreliable and we chose to not consider the GMYC partitions.

Genetic distance between phylogroups are similar to the distance of each phylogroup to the G group. However, G group corresponds to a nominal species (*F. mawsoni)* which suggests that the A-G phylogroups could correspond to seven different yet undescribed species.

A very rapid local turn-over

Relative abundance of phylogroups: Two of the analyzed locations (Ross Sea and East Weddell Sea) were densely sampled (at least 50 sequences) at two (or more) different time points. We therefore took advantage of these time series to evaluate the turnover of the resident crinoids from year to year. The analysis of the relative abundance of the various

phylogroups shows that for both locations the genetic composition is highly unstable through

time (Fig. 2a and 2c). Phylogroup composition in EWS is stable from 1996 to 2004, and

changes drastically in 2005 when phylogroup C has not been sampled. This pattern leads to a

highly significant difference between all years (Fisher exact test, P = 4.7 10^{-10}). Similarly, the

composition of the RS location is also highly significantly different between 2004 and 2008

(Fisher exact test, P = 3.2 10^{-11}); in this case, phylogroup A is replaced by phylogroups B and

E.

Other locations display a single sampling event (AS, DS, TA) or reduced number of

sequences over several sampling events (e.g. SAW: 2 sequences in 2000, 11 in 2002, 7 in

2006 and 68 in 2009), and cannot be used to robustly assess the stability of phylogroup

relative abundance.

Turn-over within the phylogroups: We then measured the replacement of genetic variants

within the phylogroups. To do so, we considered samples with at least 10 sequences of the

same phylogroup in the same location at two different time points. The pairwise comparisons

for such groups are all reported in Table 1 and we provide two graphical representations for

phylogroup D in the RS location (Fig. 2b) and phylogroup C in the EWS location (Fig. 2d). In

two cases (phylogroup C in EWS and phylogroup D in RS), we found a significant difference

in the genetic composition within each phylogroups even when they are separated by only few

years, again suggesting an unexpectedly high turn-over of the genetic pool of these crinoids.

Interestingly, the temporal differentiation of phylogroup E within EWS is already F_{ST}=4% in

four years, that is not significant because of the small sample size (10 and 11 respectively).

Only the two samples of phylogroup G in EWS, that are separated by a single year, show no

sign of temporal differentiation (F_{ST}=1%).

Estimation of the mutation rate and the effective population size

Using the robust yet elegant approach devised by Fu (2001), we estimated the yearly mutation

rates (μ) from the same sample comparisons (i.e. phylogroups with two or more samples of 10

or more sequences within the same location). Furthermore, we also estimated the

corresponding effective population size (Ne) assuming a generation time of three years.

Results (Table 2) shows that all estimated mutation rates are 10^{-5}-10^{-4} /bp/year, with a mean

of 8.10^{-5} /bp/year. Results also show that the effective sizes are very low, on the order of 2-7

(Table 2). The very low effective size corresponds to the unexpected high turn-over of the

genetic pool that we observe within the phylogroup at the same location.

Polymorphism analysis

For samples of 10 sequences or more (with the same year and same location), we have
analyzed the characteristics of segregating polymorphisms using four neutrality tests:
Tajima's D (Tajima, 1989), Fu and Li's F* and D* (Fu & Li, 1993), Achaz's Y* (Achaz,
2008) and Strobeck P-values (Strobeck, 1987). Results (Fig. 3) show almost no deviation
from the standard neutral model, except for the Fu and Li's F* and D*. This statistics points
to a clear excess of singletons. However, singletons likely contain sequencing errors that
strongly skew these statistics (Achaz, 2008). The Y statistics that ignores singletons exhibits a
distribution that is very close to the one expected under the standard neutral model. We
therefore conclude that the observed polymorphisms suggest that basic assumptions of the
standard neutral model (i.e. constant population size, no migration, no structure and no
selection) cannot be rejected for these crinoid phylogroups.

Assessing the effect of migration using simulations

To test whether migration could accelerate the genetic turn-over of a species on a given
location, we studied a simple model where the sampled location exchanges migrants with a
large pool of panmictic individuals. One expects that a constant arrival of migrants from this
large pool could interfere with the estimation of effective size. We thus simulated a model
where the sampled population has a size N and the pool $10N$. We studied the impact of the
migration rate on the estimation of Ne using the same method than above (2001). Results (Fig.
4) show that at low migration rate, the estimator is unaffected whereas it is inflated up to $11N$
at high migration rate. This shows that the existence of a large pool from where migrants can
come into the sampled population can only inflate the estimation of *Ne* and therefore cannot
explain the low *Ne* we have estimated.

Phylogroups divergence

We then computed an estimate for the times of split between the seven phylogroups using our
population genetics estimation of the mutation rate (8.10^{-5} mutations /bp /year). Our estimates
are reported in an ultrametric tree (Fig. 5) where the oldest split between the phylogroups is
estimated around 400 years. This very short time scale is in sharp contrast with the millions of
years of previous estimations based on other estimated mutation rates (3.10^{-8} mutations /bp
/year) that were based on echinoderms thought to be separated by the formation of the
panama isthmus (Lessios, Kessing, Robertson, & Paulay, 1999).

Discussion

Speciation in Antarctica: the classical view

Due to its extreme environmental conditions, it has been hypothesized that climate driven evolution in Southern Ocean has been more important than in other ecosystems where ecological interactions play a larger role (Rogers, 2012). The diversity and the distribution of many marine taxa in Antarctica has been strongly shaped by climate variation and continental drift leading to dispersal, vicariance and extinction since the break-up of Gondwana (Williams, Reid, & Littlewood, 2003). Indeed, the global climate change at the end of the Eocene related to the continental drift is characteristic of the transition from a temperate climate to the current polar climate in Antarctica (Clarke et al., 1992; Clarke & Crame, 1989). The physical separation of Antarctica, South America and Australia resulted in the cooling of the Southern Ocean, which in turn initiated the physical and climate isolation of this region of the world (Tripati, Backman, Elderfield, & Ferretti, 2005). This isolation was followed by an extensive continental glaciations and the initiation of the Antarctic Circumpolar Current in the Miocene (Clarke et al., 1992; Clarke & Crame, 1989; Thatje et al., 2008). These two processes exacerbated the environmental and biological isolation of the Southern Ocean and are classically recognized as the main explanation to the high level of endemism observed in Antarctica (Clarke & Crame, 1989; Griffiths, Barnes, & Linse, 2009).

The Quaternary was marked by the oscillation of glaciation and interglacial periods, which strongly affected the ice cover present in Antarctica (Davies, Hambrey, Smellie, Carrivick, & Glasser, 2012). These cycles led to severe temperature oscillations and to the succession of large extensions of the ice sheets on the continental shelf followed by retreats of the ice sheets. They have been one of the main drivers of the diversification of species in Antarctica. Indeed, the isolation of populations in ice free refugia during the glacial era, such as polynya or in the deep sea, has been suggested as a mechanism of fragmentation of populations leading to allopatric speciation in the Southern Ocean. This process is called the Antarctic 'biodiversity pump' (Clarke et al., 1992; Clarke & Crame, 1989). The Antarctic 'biodiversity pump' has been proposed to be the main mechanism driving speciation and diversification in Antarctica.

Our knowledge on evolutionary history of Antarctic fauna was classically derived from studies of the systematics and distribution of marine animals relying heavily on the morphology of present and past (fossil) organisms. Such approaches are limited for two main

reasons. First, species morphology may not reflect the true evolutionary relationships between taxa (DeBiasse & Hellberg, 2015). Second, several periods of time lack a reliable fossil record. Today, the emergence of molecular approaches provides a powerful tool to circumvent these limitations. To this regard, the increasing number of phylogeographic studies allow to get a better understanding of the effects of past changes on the current diversity and spatial distribution of organisms (Allcock & Strugnell, 2012). They also allow to accurately assess the distribution of genetic diversity within and among species. Such information is extremely valuable to evaluate and mitigate the impacts of human-induced activities on the biodiversity of the Southern Ocean (Chown et al., 2015). This study take place in this context and by expanding the results of two previous studies (Hemery et al., 2012; Wilson et al., 2007), we aimed to determine the drivers of the diversity of two crinoid taxa in the Southern Ocean, *P. kerguelensis* and *F. mawsoni*.

The concept of species: is Promachocrinus kerguelensis a complex of multiple cryptic species?

The analysis of 1797 sequences using recent species delimitation approaches allowed to identify seven operational taxonomic units. We confirmed the presence of six phylogroups (A-F) identified in two previous study (Hemery et al., 2012; Wilson et al., 2007), but only the PTP method supports the further split between E1 and E2 suggested by Hemery *et al.* (2012). Furthermore, despite their strong morphological differences and monophyly (Fig. 1), our results strongly suggest that *F. mawsoni* (phylogroup G) belongs to the *P. kerguelensis* complex. Indeed, the reciprocal monophyly of G and F and the proximity of E to G/F compare to A-D confirm the affiliation of G as a member of *P. kerguelensis* complex (Fig. 1). Therefore, we corroborated Hemery *et al.* (2012) results showing that *P. kerguelensis* consists in a complex of several cryptic species. We also confirmed Eléaume (2006) conclusion that despite striking exterior morphological differences phylogroup G is also a member of this complex of species, and Hemery et al. (2013b) results based on a multi-markers phylogenetic reconstruction.

Morphological similarities between cryptic species often reflect a recent speciation event (Bickford et al., 2007). In the marine realm, speciation is less associated to morphology than to other phenotypic aspects such as chemical recognition systems (Palumbi, 1994). Knowlton (1993) argued that marine habitats are likely to be filled with cryptic species although rarely

recognized due to limited access to marine habitats. Molecular approaches can be particularly

powerful in detecting cryptic species on the basis of their molecular divergence, and recent

molecular studies helped reveal the prevalence of cryptic species (See for example (Brasier et

al., 2016; Grant, Griffiths, Steinke, Wadley, & Linse, 2010)). This is particularly the case in

the Southern Ocean where numerous molecular studies have suggested the presence of cryptic

species (Baird, Miller, & Stark, 2011; Brasier et al., 2016; Grant et al., 2010; Wilson, Schrödl,

& Halanych, 2009) with areas such as the Scotia Arc representing potential hotspots of

cryptic diversity (Linse, Cope, Lörz, & Sands, 2007). The prevalence of invertebrate cryptic

species in the Southern Ocean is probably due to the cyclical variation of ice sheets extent in

Antarctica during repeated glacial and interglacial periods which could have fostered the

separation of population, promoting genetic divergence and allopatric cryptic speciation

(Clarke & Crame, 1989). During glacial periods, ice sheets extension is thought to have

forced marine species inhabiting the continental shelf to take refuge in the deep sea or in shelf

refugia, such as areas where permanent polynyas (Thatje et al., 2008) occurred. During glacial

maxima the decrease of gene flow between populations would have promoted reproductive

isolation and increased genetic variation between populations. Under glacial maximum

extreme environmental conditions, a potential increase of the diverging selective pressures on

behavioral and physiological characters rather than morphology (which could be under strong

stabilizing selection), could lead to a reduction of the morphological changes usually

associated with speciation (Fišer, Robinson, & Malard, 2018). In such a scenario, high levels

of divergence are expected on neutrally evolving genes as observed in this study. This process

might explain the high number of cryptic species reported in the Southern Ocean.

The concept of species is controversial in biology as there is no unique definition that can be

applied to all species under all circumstances (Mayden, 1997). Here, the application of

species delimitation methods (based on the genetic concept of species) using molecular tools

identified six highly divergent lineages indiscernible on the basis of their morphology and a

last morphologically divergent lineage. Therefore, this study show how the use of molecular

data can provide new insights into the nature of the genetic boundaries between species (Pante

et al., 2015). However, in the light of these findings we can wonder what do the lineages

highlighted in this study represent? Are they genuine different species? Here the use of

different criteria to delineate species based on reproductive isolation (biological concepts of

species) or on morphology (morphological concept of species) will probably lead to the

identification of a different number of species (Fišer et al., 2018). This is a good illustration of

the "species problem" and stress out the fact that the notion of species itself is an abstract

concept that should be questioned (De Queiroz, 2007). It has also important consequences in

term of the protection of the biodiversity because disparate groupings of species will in turn

result in different economic implications and conservation decisions (Frankham et al., 2012).

A remarkably fast turn over and low effective population sizes

This study indicates that over time the different sectors of the Southern Ocean show a

remarkable turn over in term of phylogroup composition. For example, in less than a decade,

the Ross and East Weddell Sea phylogroups composition changed substantially (Fig 2a and

2c). A, B and F may or may not be sampled from one year to the next, whereas C, D and G

are always present but their proportions change drastically (Fig 2a and 2c). This observation

suggests that within the same geographical region, the real number of individual per

phylogroup varies considerably over time. One hypothesis that could explain these patterns is

the presence of strong marine currents (such as the ACC, Ross Gyre and Weddell Gyre)

mixing individuals and thus changing continuously their distribution across the Southern

Ocean over time. A similar mechanism has been evoked to explain the current diversity and

the distribution of sponges across the Southern Ocean (Downey, Griffiths, Linse, & Janussen,

2012).

Furthermore, within phylogroups, in a few years, the genetic makeup can change significantly.

For instance, the genetic composition of the phylogroup C and D changed totally between

1996 and 2008, respectively in the East Weddell Sea and Ross Sea (Fig. 2b, d). Such spatio-

temporal variations in term of genetic composition could also be attributed to marine currents

that constantly shuffle individuals between populations. Over time, the constant arrival of

individuals stemming from different populations continually modify the observed genetic

composition within the same geographical location.

Alternatively, a rapid divergence of the genetic pool by genetic drift, or any other population

processes, could also explain this rapid genetic turn over. The low effective population sizes

highlighted in this study ($Ne < 10$; Table 2) are also consistent with this pattern. Indeed, lower

values of Ne are associated with increased amount of "apparent" genetic drift. Furthermore,

isolated populations are expected to diverge inversely proportional to their effective

population sizes. Therefore, when the effective population sizes are extremely low,

populations can significantly diverge from each other in just a few generations (Hudson &
Coyne, 2002).

The simulations conducted in this study (Fig. 4) show that gene flow from an unsampled
reservoir should inflate the *Ne* values estimated with the method of Fu (Fu, 2001). Therefore,
due to the extremely low value of *Ne* observed, the hypothesis of a rapid divergence by
"apparent" genetic drift is favored here over the hypothesis of a rapid divergence mediated by
marine current continually modifying the genetic pool of the populations.

It is also important to bear in mind that our estimates of *Ne* depend directly on the number of
generations per year (Fu, 2001). In Table 2, we assumed a generation time of three year,
however if for example we assumed a generation time of 30 years, our estimates of *Ne* would
increase tenfold. In our case, depending on the assumed value of the generation time, the
interpretation of the *Ne* could change accordingly.

Overall, the *Ne* values estimated in this study for the *P. kerguelensis* and *F. mawsoni* are low.
However, they are highly abundant in the Southern Ocean (Hemery et al., 2013a; McClintock
& Pearse, 1987) which is indicative of a high census size. Such a discrepancy between the
census (*Nc*) and effective (*Ne*) population size is commonly observed in nature, especially for
marine species with large size (Filatov, 2019). However, the reasons behind it are not always
clear (Filatov, 2019; Palstra & Fraser, 2012). At least four biological phenomena could cause
Nc and *Ne* to be different in *P. kerguelensis* complex (with *Ne* << *Nc*). First, an unequal sex
ratio, i.e. an uneven number of males and females contributing to the next generation, lead the
rarer sex to have a greater effect on genetic drift which decreases *Ne* compare to *Nc*. Although
this hypothesis is plausible, to this day the sex ratio of *P. kerguelensis* and *F. mawsoni*
remains unknown. Furthermore, the sex-ratio distortion would need to be extreme to reach
such a low *Ne*. Second, a fluctuation of *Nc* over time can affect the ratio between *Ne* and *Nc*.
Indeed, an estimate of *Ne* over several generations is the harmonic mean of *Nc* over time.
Therefore, if *Nc* changes substantially over time, *Ne* is typically smaller than *Nc*. Fig. 2
indicates that the proportion of *P. kerguelensis* and *F. mawsoni* lineages can vary radically
from year to year and thus the fluctuation of *Nc* over time for each lineage can change
consequently leading *Ne* to be smaller than *Nc*. Third, the reproductive strategy of *P.*
kerguelensis and *F. mawsoni* can participate to reduce *Ne* compared to *Nc*. Indeed, when
individuals contribute unequally to the progeny of the next generation, a great proportion of
the next generation comes from a small number of individuals which reduces *Ne*. Due to its

mode of reproduction, *P. kerguelensis* is particularly incline to have a very reduced number of
individuals contributing disproportionately to the progeny. Reproduction occurs
synchronously around the months of November and December where males and females
spawn and due to the dispersion of gametes by currents, a very small proportion of individuals
have their ovocytes fertilized (McClintock & Pearse, 1987). Therefore, only a very small part
of the produced larvae contributes effectively to the gene pool. This phenomenon can
drastically reduce *Ne* compare to *Nc* and has been reported in different spawning species
(Lind, Evans, Knauer, Taylor, & Jerry, 2009). This mode of reproduction can also explain the
spatio-temporal genetic turn over mentioned previously. Lastly, due to its haploid and uni-
parental nature, the *Ne* of the mtDNA is known to be four times smaller than that of nuclear
DNA. Moreover, this effect can be exacerbated by the action of natural selection. Indeed,
natural selection can reduce *Ne* and the mtDNA is known to be under positive and negative
selection (Meiklejohn, Montooth, & Rand, 2007). Positive natural selection, for example, is
expected to reduce the genetic diversity leading to a low *Ne* (Gossmann, Woolfit, & Eyre-
Walker, 2011). Indeed, recurrent selective sweeps, related to positive mutations, will reduce
the genetic diversity by hitchhiking resulting in a lower *Ne* compared to *Nc*. In this regard,
Piganeau & Eyre-Walker (2009) found a strong negative correlation between *Ne* and the
strength of the natural selection operating on the mtDNA on non-synonymous mutations. In
addition, the non-recombining nature of the mtDNA is expected to make this effect even
stronger. There are accumulating evidences suggesting that positive natural selection is acting
on echinoderm mitogenomes (Castellana, Vicario, & Saccone, 2011). For example, the strong
bias of codon usage observed in the mtDNA of the crinoid species *Florometra serratissima*
compared to many other echinoderms, is also in favor of the idea that some sort of natural
selection is occurring, though the mechanism behind it is still not fully understood (Scouras &
Smith, 2001).

Tempo of speciation: Promachocrinus kerguelensis challenges the biodiversity pump hypothesis.

The mutation rate estimated in this study is approximatively equal to 10^{-5} mutations/site/year
and is based on population genetics data. Previous estimate of the mutation rate based also on
the COI applied to closely related species (Lessios et al., 1999) used a phylogeny based
approach and yielded a mutation rate of roughly 3 order of magnitude lower ($\approx 10^{-8}$

mutations/site/year). Such a discrepancy between mutation rates derived from population data

and phylogeny has been already documented (Burridge, Craw, Fletcher, & Waters, 2008;

Madrigal et al., 2012) and is called "time dependency of molecular rates" (Ho et al., 2011; Ho,

Phillips, Cooper, & Drummond, 2005). For example, in human, the mitochondrial D-loop

mutation rate derived from pedigree approaches produced an average value of 8.10^{-7}

mutations/site/year whereas phylogenetic estimates yielded value of 2.10^{-8} mutations/site/year

(Santos et al., 2005). Likewise, in fishes, Burridge (Burridge et al., 2008) observed a mutation

rate of $\approx10^{-7}$ using a population based approach and $\approx10^{-8}$ mutations/site/year from a

phylogenetic estimates. The potential causes of the "time scale dependency" are reviewed in

Ho *et al.* (2011), but one of the main reason behind it is that population genetics based

calibrations lead to estimates reflecting the spontaneous mutation rates, whereas phylogeny

based calibrations reflect the substitution rates (i.e. fixed mutations). Therefore, because

purifying selection remove the vast majority of the spontaneous deleterious mutations, which

constitute a large proportion of them, the spontaneous mutation rate is higher than the

substitution rate. Consequently, the rates of molecular evolution decrease with the age of the

calibration used to estimate them and it is invalid to assume that a unique molecular clock

applies over all time scales (Ho et al., 2005; 2011). A major implication of time dependency

is that estimates of recent population divergence times will require re-estimation (Burridge et

al., 2008). Therefore, many studies suggesting population isolation related to the Pleistocene

glaciations might need to convert these time estimates to more recent ones (i.e. last glacial

maximum or Holocene).

Furthermore, the large variation of the mutation rates across the mtDNA regions and between

different lineages (Nabholz, Glemin, & Galtier, 2009), leads the mtDNA to strongly deviate

from the molecular clock assumption, which is generally assumed in phylogenetic

reconstruction. Therefore, estimates of mtDNA mutation rates relying on phylogenetic

approach should be taken with caution (Nabholz et al., 2009; Nabholz, Glemin, & Galtier,

2008). Conversely, the mutation rate estimated in this study are quite robust as they are

model-free ((Fu, 2001) ; See also Material and Methods). In the case of *P. kerguelensis* and *F.*

mawsoni, the different mutation rates estimated have important consequences on the age of

the separation of the different lineages (Fig. 4). Indeed, the estimate from Lessios *et al.* (1999)

suggest that the split between the different lineages would have occurred in the last million

year (Fig. 4), which is congruent with the hypothesis of the biodiversity pump suggesting that

most of the speciation events would have occurred during the Quaternary cycles (Clarke et al.,

1992; Clarke & Crame, 1989). Conversely, with our estimate the separation would have

occurred in the last 1000 years, implying an extremely recent and rapid diversification for *P.*

kerguelensis and *F. mawsoni*.

Explosive radiations have been reported for several taxa (Mahler, Ingram, Revell, & Losos,

2013; Moore & Robertson, 2014; Muschick, Indermaur, & Salzburger, 2012) and are, in

general, associated with the Pleistocene glacial cycles (Hawlitschek et al., 2012). In contrast,

using the molecular rate estimated in this study, in our case the diversification would have

happened during the Holocene. Such a remarkably rapid diversification has been rarely

recorded in nature (but see Muschick et al., 2012; Peccoud, Simon, McLaughlin, & Moran,

2009). Furthermore, most of rapid radiations reported so far are adaptive and associated to

morphological novelties that allow taxa to exploit separate niches. Taxa involved in rapid

diversification are generally ecologically/morphologically highly differentiated (Losos &

Miles, 2002). Here, most of the lineages display no obvious morphological differences

(except for *Florometra*) or niche differentiation. As a consequence, the radiation within the *P.*

kerguelensis complex is probably mainly nonadaptive (Rundell & Price, 2009) or adaptive but

related to characters that are challenging to distinguish/observe (physiology or behavior). In

this context, it is worth noting that structural genomic changes could also drive taxa to rapid

adaptive (Kirkpatrick & Barton, 2006) and nonadaptive (Rowe, Aplin, Baverstock, & Moritz,

2011; Rundell & Price, 2009) radiation and may represent the mechanism underlying the

diversification of the *P. kerguelensis* complex.

Finally, the low effective population sizes estimated (Table 2) are congruent with the rapid

diversification rate observed. Indeed, Hudson and Coyne (2002) showed that for mtDNA,

under the isolation model, two lineages would take between 2 and 3*Ne* generations to become

reciprocally monophyletic by lineage sorting. Therefore, using the low effective population

size estimated in this study (Table 2), it would take a few hundreds of years for two isolated

taxa to reach complete reciprocal monophyly. This result is concordant with the divergence

observed in this study and contributes to invalidate the Antarctic "biodiversity pump" for the

Promachocrinus complex.

Acknowledgments

The following persons deserve our special thanks for having collected, curated and made available specimens from all around Antarctica: Ty Hibberd (AAD, Hobart), Owen Anderson, David Bowden, Sadie Mills, Kareen Schnabel and Peter Smith (NIWA, Wellington), Stefano Schiaparelli (University of Genoa), Jens Bohn and Eva Lodde (ZSM, Munich), David Barnes, Katrin Linse and Chester Sands (BAS, Cambridge). Our thanks also go to crew and scientists on board various cruises: CEAMARC (IPY project 53), POKERII, TAN08 and AMLR2009 cruises. Funding parties also include three Actions Transversales du MNHN: "Biodiversité actuelle et fossile; crises, stress, restaurations et panchronisme: le message systématique", "Taxonomie moléculaire: DNA Barcode et gestion durable des collections" and "Biominéralisation"; the French Polar Institute IPEV (travel grants to LGH and ME on REVOLTA); This work was supported by the Consortium National de Recherche en Génomique, and the Service de Systématique Moléculaire (SSM) at the MNHN (USM 2700). Part of the molecular work was also supported by collaboration between the Census of Antarctic Marine Life, the Marine Barcode of Life (MarBOL) project and the Canadian Centre for DNA Barcoding (CCDB). DS was supported by funding of the Alfred P. Sloan Foundation to MarBOL. Laboratory analyses on sequences generated at the CCDB were funded by the Government of Canada through Genome Canada and the Ontario Genomics Institute (2008-OGI-ICI-03). We also gratefully thank Lucile Perrier, Charlotte Tarin, Jose Ignacio Carvajal III Patterson (students) and Céline Bonillo (SSM) for their invaluable help in the molecular lab. This work was also funded by the University of Groningen through a PhD fellowship allocated to Yacine Ben Chehida. We also thank Michael C. Fontaine for the logistical support provided during the PhD of Yacine Ben Chehida.

Literature cited:

Achaz, G. (2008). Testing for neutrality in samples with sequencing errors. *Genetics*, *179*(3), 1409–1424. http://doi.org/10.1534/genetics.107.082198

Allcock, A. L., & Strugnell, J. M. (2012). Southern Ocean diversity: new paradigms from molecular ecology. *Trends in Ecology & Evolution*, *27*(9), 520–528. http://doi.org/10.1016/j.tree.2012.05.009

Baird, H. P., Miller, K. J., & Stark, J. S. (2011). Evidence of hidden biodiversity, ongoing speciation and diverse patterns of genetic structure in giant Antarctic amphipods. *Molecular Ecology*, *20*(16), 3439–3454. http://doi.org/10.1111/j.1365-294X.2011.05173.x

Bickford, D., Lohman, D. J., Sodhi, N. S., Ng, P. K. L., Meier, R., Winker, K., et al. (2007). Cryptic species as a window on diversity and conservation. *Trends in Ecology & Evolution*, *22*(3), 148–155. http://doi.org/10.1016/j.tree.2006.11.004

Brasier, M. J., Wiklund, H., Neal, L., Jeffreys, R., Linse, K., Ruhl, H., & Glover, A. G. (2016). DNA barcoding uncovers cryptic diversity in 50% of deep-sea Antarctic polychaetes. *Royal Society Open Science*, *3*(11), 160432. http://doi.org/10.1098/rsos.160432

Burridge, C. P., Craw, D., Fletcher, D., & Waters, J. M. (2008). Geological dates and molecular rates: fish DNA sheds light on time dependency. *Molecular Biology and Evolution*, *25*(4), 624–633. http://doi.org/10.1093/molbev/msm271

Castellana, S., Vicario, S., & Saccone, C. (2011). Evolutionary patterns of the mitochondrial genome in Metazoa: exploring the role of mutation and selection in mitochondrial protein coding genes. *Genome Biology and Evolution*, *3*(3), 1067–1079. http://doi.org/10.1093/gbe/evr040

Chown, S. L., Clarke, A., Fraser, C. I., Cary, S. C., Moon, K. L., & McGeoch, M. A. (2015). The changing form of Antarctic biodiversity. *Nature*, *522*(7557), 431–438. http://doi.org/10.1038/nature14505

Clark, A. H. (1921). A monograph of the existing crinoids. *Bulletin of the United States National Museum*, *1*(82), 795–949. http://doi.org/10.5479/si.03629236.82.2

Clarke, A., & Crame, J. A. (1989). The origin of the Southern Ocean marine fauna. *Geological Society, London, Special Publications*, *47*(1), 253–268. http://doi.org/10.1144/GSL.SP.1989.047.01.19

Clarke, A., Crame, J. A., Stromberg, J. O., & Barker, P. F. (1992). The Southern Ocean Benthic Fauna and Climate Change: A Historical Perspective [and Discussion]. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *338*(1285), 299–309. http://doi.org/10.1098/rstb.1992.0150

Cornils, A., & Held, C. (2014). Evidence of cryptic and pseudocryptic speciation in the *Paracalanus parvus* species complex (Crustacea, Copepoda, Calanoida). *Frontiers in Zoology*, *11*(1), 19. http://doi.org/10.1186/1742-9994-11-19

Davies, B. J., Hambrey, M. J., Smellie, J. L., Carrivick, J. L., & Glasser, N. F. (2012). Antarctic Peninsula Ice Sheet evolution during the Cenozoic Era. *Quaternary Science Reviews*, *31*, 30–66. http://doi.org/10.1016/j.quascirev.2011.10.012

De Queiroz, K. (2007). Species Concepts and Species Delimitation. *Systematic Biology*, *56*(6), 879–886. http://doi.org/10.1080/10635150701701083

DeBiasse, M. B., & Hellberg, M. E. (2015). Discordance between morphological and molecular species boundaries among Caribbean species of the reef sponge Callyspongia. *Ecology and Evolution*, *5*(3), 663–675. http://doi.org/10.1002/ece3.1381

Downey, R. V., Griffiths, H. J., Linse, K., & Janussen, D. (2012). Diversity and distribution patterns in high southern latitude sponges. *PloS One*, *7*(7), e41672.

612   http://doi.org/10.1371/journal.pone.0041672

613 Drummond, A. J., Suchard, M. A., Xie, D., & Rambaut, A. (2012). Bayesian phylogenetics
614   with BEAUti and the BEAST 1.7. *Molecular Biology and Evolution*, *29*(8), 1969–1973.
615   http://doi.org/10.1093/molbev/mss075

616 Eastman, J. T., & McCune, A. R. (2000). Fishes on the Antarctic continental shelf: evolution
617   of amarine species flock?*. *Journal of Fish Biology*, *57*(sa), 84–102.
618   http://doi.org/10.1111/j.1095-8649.2000.tb02246.x

619 Eléaume, M. (2006). Approche morphométrique de la variabilité phénotipique : conséquences
620   systématiques et évolutives : application aux crinoïdes actuels (Crinoidea :
621   Echinodermata). *Www.Theses.Fr*.

622 Eléaume, M., Hemery, L. G., Roux, M., & Améziane, N. (2014). Southern Ocean Crinoids. In
623   *Biogeographic Atlas of the Southern Ocean* (pp. 208–212). Cambridge.

624 Felsenstein, J. (1989). PHYLIP - Phylogeny Inference Package (version 3.2). *Cladistics*, (5),
625   164–166.

626 Filatov, D. A. (2019). Extreme Lewontin's Paradox in Ubiquitous Marine Phytoplankton
627   Species. *Molecular Biology and Evolution*, *36*(1), 4–14.
628   http://doi.org/10.1093/molbev/msy195

629 Fišer, C., Robinson, C. T., & Malard, F. (2018). Cryptic species as a window into the
630   paradigm shift of the species concept. *Molecular Ecology*, *27*(3), 613–635.
631   http://doi.org/10.1111/mec.14486

632 Folmer, O., Black, M., Hoeh, W., Lutz, R., & Vrijenhoek, R. (1994). DNA primers for
633   amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan
634   invertebrates. *Molecular Marine Biology and Biotechnology*, *3*(5), 294–299.

635 Frankham, R., Ballou, J. D., Dudash, M. R., Eldridge, M. D. B., Fenster, C. B., Lacy, R. C., et
636   al. (2012). Implications of different species concepts for conserving biodiversity.
637   *Biological Conservation*, *153*, 25–31. http://doi.org/10.1016/j.biocon.2012.04.034

638 Fu, Y. X. (2001). Estimating mutation rate and generation time from longitudinal samples of
639   DNA sequences. *Molecular Biology and Evolution*, *18*(4), 620–626.
640   http://doi.org/10.1093/oxfordjournals.molbev.a003842

641 Fu, Y. X., & Li, W. H. (1993). Statistical tests of neutrality of mutations. *Genetics*, *133*(3),
642   693–709.

643 Gascuel, O. (1997). BIONJ: an improved version of the NJ algorithm based on a simple
644   model of sequence data. *Molecular Biology and Evolution*, *14*(7), 685–695.
645   http://doi.org/10.1093/oxfordjournals.molbev.a025808

646 Gossmann, T. I., Woolfit, M., & Eyre-Walker, A. (2011). Quantifying the variation in the
647   effective population size within a genome. *Genetics*, *189*(4), 1389–1402.
648   http://doi.org/10.1534/genetics.111.132654

649 Gouy, M., Guindon, S., & Gascuel, O. (2010). SeaView version 4: A multiplatform graphical
650   user interface for sequence alignment and phylogenetic tree building. *Molecular Biology*
651   *and Evolution*, *27*(2), 221–224. http://doi.org/10.1093/molbev/msp259

652 Grant, R. A., Griffiths, H. J., Steinke, D., Wadley, V., & Linse, K. (2010). Antarctic DNA
653   barcoding; a drop in the ocean? *Polar Biology*, *34*(5), 775–780.
654   http://doi.org/10.1007/s00300-010-0932-7

655 Griffiths, H. J. (2010). Antarctic marine biodiversity--what do we know about the distribution
656   of life in the Southern Ocean? *PloS One*, *5*(8), e11683.
657   http://doi.org/10.1371/journal.pone.0011683

658 Griffiths, H. J., Barnes, D. K. A., & Linse, K. (2009). Towards a generalized biogeography of
659   the Southern Ocean benthos. *Journal of Biogeography*, *36*(1), 162–177.
660   http://doi.org/10.1111/j.1365-2699.2008.01979.x

661 Hawlitschek, O., Hendrich, L., Espeland, M., Toussaint, E. F. A., Genner, M. J., & Balke, M.

662 (2012). Pleistocene climate change promoted rapid diversification of aquatic invertebrates
663 in Southeast Australia. *BMC Evolutionary Biology*, *12*(1), 142.
664 http://doi.org/10.1186/1471-2148-12-142
665 Heimeier, D., Lavery, S., & Sewell, M. A. (2010). Molecular species identification of
666 *Astrotoma agassizii* from planktonic embryos: further evidence for a cryptic species
667 complex. *The Journal of Heredity*, *101*(6), 775–779. http://doi.org/10.1093/jhered/esq074
668 Hemery, L. G., Améziane, N., & Eléaume, M. (2013a). Circumpolar dataset of sequenced
669 specimens of Promachocrinus kerguelensis (Echinodermata, Crinoidea). *ZooKeys*,
670 *315*(60), 55–64. http://doi.org/10.3897/zookeys.315.5673
671 Hemery, L. G., Eléaume, M., Roussel, V., Améziane, N., Gallut, C., Steinke, D., et al. (2012).
672 Comprehensive sampling reveals circumpolarity and sympatry in seven mitochondrial
673 lineages of the Southern Ocean crinoid species *Promachocrinus kerguelensis*
674 (Echinodermata). *Molecular Ecology*, *21*(10), 2502–2518. http://doi.org/10.1111/j.1365-
675 294X.2012.05512.x
676 Hemery, L. G., Roux, M., Améziane, N., & Eléaume, M. (2013b). High-resolution crinoid
677 phyletic inter-relationships derived from molecular data. *Cahiers De Biologie Marine*,
678 *54*(4), 511–525.
679 Ho, S. Y. W., Lanfear, R., Bromham, L., Phillips, M. J., Soubrier, J., Rodrigo, A. G., &
680 Cooper, A. (2011). Time-dependent rates of molecular evolution. *Molecular Ecology*,
681 *20*(15), 3087–3101. http://doi.org/10.1111/j.1365-294X.2011.05178.x
682 Ho, S. Y. W., Phillips, M. J., Cooper, A., & Drummond, A. J. (2005). Time dependency of
683 molecular rate estimates and systematic overestimation of recent divergence times.
684 *Molecular Biology and Evolution*, *22*(7), 1561–1568.
685 http://doi.org/10.1093/molbev/msi145
686 Hudson, R. R., & Coyne, J. A. (2002). Mathematical consequences of the genealogical
687 species concept. *Evolution*, *56*(8), 1557–1565.
688 Hudson, R. R., Boos, D. D., & Kaplan, N. L. (1992). A statistical test for detecting geographic
689 subdivision. *Molecular Biology and Evolution*, *9*(1), 138–151.
690 http://doi.org/10.1093/oxfordjournals.molbev.a040703
691 Kirkpatrick, M., & Barton, N. (2006). Chromosome inversions, local adaptation and
692 speciation. *Genetics*, *173*(1), 419–434. http://doi.org/10.1534/genetics.105.047985
693 Knowlton, N. (1993). Sibling Species in the Sea. *Annual Review of Ecology and Systematics*,
694 *24*(1), 189–216. http://doi.org/10.1146/annurev.ecolsys.24.1.189
695 Knowlton, N. (2000). Molecular genetic analyses of species boundaries in the sea.
696 *Hydrobiologia*, *420*(1), 73–90. http://doi.org/10.1023/A:1003933603879
697 Lecointre, G., Améziane, N., Boisselier, M.-C., Bonillo, C., Busson, F., Causse, R., et al.
698 (2013). Is the species flock concept operational? The Antarctic shelf case. *PloS One*, *8*(8),
699 e68787. http://doi.org/10.1371/journal.pone.0068787
700 Lessios, H. A., Kessing, B. D., Robertson, D. R., & Paulay, G. (1999). Phylogeography of the
701 Pantropical Sea Urchin Eucidaris in Relation to Land Barriers and Ocean Currents.
702 *Evolution*, *53*(3), 806. http://doi.org/10.2307/2640720
703 Librado, P., & Rozas, J. (2009). DnaSP v5: a software for comprehensive analysis of DNA
704 polymorphism data. *Bioinformatics*, *25*(11), 1451–1452.
705 http://doi.org/10.1093/bioinformatics/btp187
706 Lind, C. E., Evans, B. S., Knauer, J., Taylor, J. J. U., & Jerry, D. R. (2009). Decreased genetic
707 diversity and a reduced effective population size in cultured silver-lipped pearl oysters
708 (*Pinctada maxima*). *Aquaculture*, *286*(1-2), 12–19.
709 http://doi.org/10.1016/j.aquaculture.2008.09.009
710 Linse, K., Cope, T., Lörz, A.-N., & Sands, C. (2007). Is the Scotia Sea a centre of Antarctic
711 marine diversification? Some evidence of cryptic speciation in the circum-Antarctic

bivalve *Lissarca notorcadensis* (Arcoidea: Philobryidae). *Polar Biology*, *30*(8), 1059–
1068. http://doi.org/10.1007/s00300-007-0265-3
Losos, J. B., & Miles, D. B. (2002). Testing the hypothesis that a clade has adaptively
radiated: iguanid lizard clades as a case study. *The American Naturalist*, *160*(2), 147–157.
http://doi.org/10.1086/341557
Madrigal, L., Posthumously, L. C., Melendez-Obando, M., Villegas-Palma, R., Barrantes, R.,
Raventos, H., et al. (2012). High mitochondrial mutation rates estimated from deep-
rooting Costa Rican pedigrees. *American Journal of Physical Anthropology*, *148*(3), 327–
333. http://doi.org/10.1002/ajpa.22052
Mahler, D. L., Ingram, T., Revell, L. J., & Losos, J. B. (2013). Exceptional convergence on
the macroevolutionary landscape in island lizard radiations. *Science*, *341*(6143), 292–295.
http://doi.org/10.1126/science.1232392
Mayden, R. L. (1997). A hierarchy of species concepts: the denouement in the saga of the
species problem.
McClintock, J. B., & Pearse, J. S. (1987). Reproductive biology of the common antarctic
crinoid *Promachocrinus kerguelensis* (Echinodermata: Crinoidea). *Marine Biology*, *96*(3),
375–383. http://doi.org/10.1007/BF00412521
Meiklejohn, C. D., Montooth, K. L., & Rand, D. M. (2007). Positive and negative selection
on the mitochondrial genome. *Trends in Genetics : TIG*, *23*(6), 259–263.
http://doi.org/10.1016/j.tig.2007.03.008
Michener, C. D., & Sokal, R. R. (1957). A quantitative approach to a problem in classification.
Evolution, *11*(2), 130–162. http://doi.org/10.1111/j.1558-5646.1957.tb02884.x
Moore, W., & Robertson, J. A. (2014). Explosive adaptive radiation and extreme phenotypic
diversity within ant-nest beetles. *Current Biology : CB*, *24*(20), 2435–2439.
http://doi.org/10.1016/j.cub.2014.09.022
Muschick, M., Indermaur, A., & Salzburger, W. (2012). Convergent evolution within an
adaptive radiation of cichlid fishes. *Current Biology : CB*, *22*(24), 2362–2368.
http://doi.org/10.1016/j.cub.2012.10.048
Nabholz, B., Glemin, S., & Galtier, N. (2008). Strong variations of mitochondrial mutation
rate across mammals--the longevity hypothesis. *Molecular Biology and Evolution*, *25*(1),
120–130. http://doi.org/10.1093/molbev/msm248
Nabholz, B., Glemin, S., & Galtier, N. (2009). The erratic mitochondrial clock: variations of
mutation rate, not population size, affect mtDNA diversity across birds and mammals.
BMC Evolutionary Biology, *9*(1), 54. http://doi.org/10.1186/1471-2148-9-54
Palstra, F. P., & Fraser, D. J. (2012). Effective/census population size ratio estimation: a
compendium and appraisal. *Ecology and Evolution*, *2*(9), 2357–2365.
http://doi.org/10.1002/ece3.329
Palumbi, S. R. (1994). Genetic Divergence, Reproductive Isolation, and Marine Speciation.
Annual Review of Ecology and Systematics, *25*(1), 547–572.
http://doi.org/10.1146/annurev.ecolsys.25.1.547
Pante, E., Abdelkrim, J., Viricel, A., Gey, D., France, S. C., Boisselier, M. C., & Samadi, S.
(2015). Use of RAD sequencing for delimiting species. *Heredity*, *114*(5), 450–459.
http://doi.org/10.1038/hdy.2014.105
Peccoud, J., Simon, J.-C., McLaughlin, H. J., & Moran, N. A. (2009). Post-Pleistocene
radiation of the pea aphid complex revealed by rapidly evolving endosymbionts.
Proceedings of the National Academy of Sciences of the United States of America,
106(38), 16315–16320. http://doi.org/10.1073/pnas.0905129106
Pfenninger, M., & Schwenk, K. (2007). Cryptic animal species are homogeneously
distributed among taxa and biogeographical regions. *BMC Evolutionary Biology*, *7*(1),
121. http://doi.org/10.1186/1471-2148-7-121

Piganeau, G., & Eyre-Walker, A. (2009). Evidence for variation in the effective population
size of animal mitochondrial DNA. *PloS One*, *4*(2), e4396.
http://doi.org/10.1371/journal.pone.0004396
Pons, J., Barraclough, T. G., Gomez-Zurita, J., Cardoso, A., Duran, D. P., Hazell, S., et al.
(2006). Sequence-based species delimitation for the DNA taxonomy of undescribed
insects. *Systematic Biology*, *55*(4), 595–609.
Puillandre, N., Lambert, A., Brouillet, S., & Achaz, G. (2011). ABGD, Automatic Barcode
Gap Discovery for primary species delimitation. *Molecular Ecology*, *21*(8), 1864–1877.
http://doi.org/10.1111/j.1365-294X.2011.05239.x
Rogers, A. D. (2007). Evolution and biodiversity of Antarctic organisms: a molecular
perspective. *Philosophical Transactions of the Royal Society B: Biological Sciences*,
362(1488), 2191–2214. http://doi.org/10.1098/rstb.2006.1948
Rogers, A. D. (2012). Evolution and Biodiversity of Antarctic Organisms. In *Antarctic
Ecosystems* (Vol. 187, pp. 417–467). Chichester, UK: John Wiley & Sons, Ltd.
http://doi.org/10.1002/9781444347241.ch14
Rowe, K. C., Aplin, K. P., Baverstock, P. R., & Moritz, C. (2011). Recent and Rapid
Speciation with Limited Morphological Disparity in the Genus Rattus. *Systematic Biology*,
60(2), 188–203. http://doi.org/10.1093/sysbio/syq092
Rundell, R. J., & Price, T. D. (2009). Adaptive radiation, nonadaptive radiation, ecological
speciation and nonecological speciation. *Trends in Ecology & Evolution*, *24*(7), 394–399.
http://doi.org/10.1016/j.tree.2009.02.007
Santos, C., Montiel, R., Sierra, B., Bettencourt, C., Fernandez, E., Alvarez, L., et al. (2005).
Understanding differences between phylogenetic and pedigree-derived mtDNA mutation
rate: a model using families from the Azores Islands (Portugal). *Molecular Biology and
Evolution*, *22*(6), 1490–1505. http://doi.org/10.1093/molbev/msi141
Scouras, A., & Smith, M. J. (2001). A novel mitochondrial gene order in the crinoid
echinoderm *Florometra serratissima*. *Molecular Biology and Evolution*, *18*(1), 61–73.
http://doi.org/10.1093/oxfordjournals.molbev.a003720
Speel, J. A., & Dearborn, J. H. (1983). Comatulid Crinoids From R/V Eltanin Cruises In The
Southern Ocean. In *Biology of the Antarctic Seas XIII* (Vol. 38, pp. 1–60). Washington, D.
C.: American Geophysical Union. http://doi.org/10.1029/AR038p0001
Strobeck, C. (1987). Average number of nucleotide differences in a sample from a single
subpopulation: a test for population subdivision. *Genetics*, *117*(1), 149–153.
Tajima, F. (1983). Evolutionary relationship of DNA sequences in finite populations.
Genetics, *105*(2), 437–460.
Tajima, F. (1989). Statistical method for testing the neutral mutation hypothesis by DNA
polymorphism. *Genetics*, *123*(3), 585–595.
Thatje, S., Hillenbrand, C.-D., & Larter, R. (2005). On the origin of Antarctic marine benthic
community structure. *Trends in Ecology & Evolution*, *20*(10), 534–540.
http://doi.org/10.1016/j.tree.2005.07.010
Thatje, S., Hillenbrand, C.-D., Mackensen, A., & Larter, R. (2008). Life hung by a thread:
endurance of Antarctic fauna in glacial periods. *Ecology*, *89*(3), 682–692.
Thornhill, D. J., Mahon, A. R., Norenburg, J. L., & Halanych, K. M. (2008). Open-ocean
barriers to dispersal: a test case with the Antarctic Polar Front and the ribbon worm
Parborlasia corrugatus (Nemertea: Lineidae). *Molecular Ecology*, *17*(23), 5104–5117.
http://doi.org/10.1111/j.1365-294X.2008.03970.x
Tripati, A., Backman, J., Elderfield, H., & Ferretti, P. (2005). Eocene bipolar glaciation
associated with global carbon cycle changes. *Nature*, *436*(7049), 341–346.
http://doi.org/10.1038/nature03874
Williams, S. T., Reid, D. G., & Littlewood, D. T. J. (2003). A molecular phylogeny of the

Littorininae (Gastropoda: Littorinidae): unequal evolutionary rates, morphological
parallelism, and biogeography of the Southern Ocean. *Molecular Phylogenetics and*
Evolution, *28*(1), 60–86. http://doi.org/10.1016/S1055-7903(03)00038-1
Wilson, N. G., Hunter, R. L., Lockhart, S. J., & Halanych, K. M. (2007). Multiple lineages
and absence of panmixia in the "circumpolar" crinoid *Promachocrinus kerguelensis* from
the Atlantic sector of Antarctica. *Marine Biology*, *152*(4), 895–904.
http://doi.org/10.1007/s00227-007-0742-9
Wilson, N. G., Schrödl, M., & Halanych, K. M. (2009). Ocean barriers and glaciation:
evidence for explosive radiation of mitochondrial lineages in the Antarctic sea slug *Doris*
kerguelenensis (Mollusca, Nudibranchia). *Molecular Ecology*, *18*(5), 965–984.
http://doi.org/10.1111/j.1365-294X.2008.04071.x
Zhang, J., Kapli, P., Pavlidis, P., & Stamatakis, A. (2013). A general species delimitation
method with applications to phylogenetic placements. *Bioinformatics*, *29*(22), 2869–2876.
http://doi.org/10.1093/bioinformatics/btt499
826

827 # **Tables and figures**

830 Below diagonal: Fst. Above diagonal: probability of pamixia using the Hudson test.

		EWS									RS	
		C				E			G		D	
n=	39	37	24			10	11		14	44		27	26
	1996	2000	2004			2000	2004		2004	2005		2004	2008
1996	-	0.035	0.004	2000	-	0.32	2004	-	0.40	2004	-	0.0002
2000	0.05	-	$<10^{-4}$	2004	0.04	-	2005	0.01	-	2008	0.21	-
2004	0.19	0.47	-									

833 **Table 2: Estimation of mutation rate and effective population size**

834 Summary table of effective population sizes (*Ne* in numbers of individuals) and mutation

835 rates (µ per base pair per year) for each phylogroup estimated by the Fu (2001) method. We

836 also report the years and the number sequences used for each estimation.

Location	EWS								RS	
Phylogroup	C			E		G			D	
n=	39	37	24	10	11	14	44		27	26
Dates	1996	2000	2004	2000	2004	2004	2005		2004	2008
µ	$1.9 \cdot 10^{-5}$			$2.7 \cdot 10^{-5}$		$3.5 \cdot 10^{-4}$			$4.2 \cdot 10^{-4}$	
Ne	4.0			7.0		1.7			2.6	

Fig. 1: Phylogenetic representation of the 1797 COI sequences from *P. kerguelensis* and

F. mawsoni. The tree was inferred using BIONJ on the unique sequences. Each square

represents one sequence. The phylogroups are the ones used in the analysis (mostly ABGD

groups) and both supplementary splits proposed by PTP are indicated as C1/C2 and E1/E2.

The colors represent the different phylogroups considered in this study. The boxes and the

numbers refer to the number of individuals carrying the haplotype.

Fig. 2: Genetic turn over at the RS (a and b) and EWS (c and d) locations. The pie charts (a

and c) represent the number of phylogroups at different years in the locations, with the same

colors code as in Figure 1. The heatmaps (b and d) represents P-values of the Hudson et *al.*

(1992) tests for genetic structure within a phylogroup of each location.

Fig. 3: Distribution of p-values for different neutrality tests on the 1,624 sequences tested for

group with at least 10 sequences. From left to right: Tajima D (1983), Fu and Li D* (1993),

Fu and Li F* (1993), Achaz Y* (2008) and Strobeck S (1987).

Fig. 4: Simulation of migration effect on Ne estimation.

Fig. 5: Divergence time of the different phylogroup of the *P. kerguelensis* complex. The

depth of the triangles represents the maximum diversity within each phylogroup. The time is

indicated in years before present. The timescale on the top is assuming a mutation rate of

8.10^{-5} mutation/bp/year. The timescale on the bottom is assuming a mutation rate of 8.10^{-3}

mutation/bp/year.

a)

Ross Sea

2004
n=123

2008
n=174

b)

within
phylogroup D

2008

2004

2004
(n=26)

2008
(n=27)

c)

East Weddle
Sea

1996
n=59

2002
n=56

2004
n=55

2005
n=55

d)

within
phylogroup C

2004

2000

1996

1996
(n=24)

2000
(n=37)

2004
(n=39)

phylogroup

A B C D F E G

P-value

≤0.001 ≤0.01 ≤0.05 ≤0.1 ≤0.25 ≤0.5 ≤1

Population genetics estimate
($\mu = 8 \cdot 10^{-5}$)

| 400 | 300 | 200 | 100 | 0 |

Lessios (1999) estimate
($\mu = 3 \cdot 10^{-8}$)

~10^6 ~$5 \cdot 10^5$ 0

Years