1

Neuroblast sensory quiescence depends of vascular cytoneme contacts and sensory neuronal differentiation requires initiation of blood flow

Laura Taberner[1], Aitor Bañón[1] and Berta Alsina[1]*

1-Developmental Biology Unit, Department of Experimental and Health Sciences, Universitat Pompeu Fabra-Parc de Recerca Biomèdica de Barcelona, Dr. Aiguader 88, 08003 Barcelona, Spain

*Author for correspondence, berta.alsina@upf.edu

Running Title: Neurovascular interaction in the sensory niche

Keywords: sensory neurons, vascular, niche, cytoneme, Dll4/Notch, blood flow, zebrafish

Highlights

♦ Vasculature is part of the cranial sensory ganglia niche and regulates neurogenesis.

♦ Cytoneme contacts between endothelial cells and sensory neuroblasts are required for neuroblast quiescence.

♦ Endothelial Dll4 and neuroblast Notch1 signal to regulate the growth of cranial sensory ganglia.

♦ Initiation of blood flow triggers a transcriptional metabolic switch and sensory neuronal differentiation.

Summary

In many organs, stem cell function depends on the communication with their niche partners. Cranial sensory neurons develop in close proximity to blood vessels, however whether vasculature is an integral component of their niches is yet unknown. Here, two separate, novel roles for vasculature in cranial sensory neurogenesis in zebrafish are uncovered. The first involves precise spatiotemporal endothelial-neuroblast cytoneme contacts and Dll4-Notch signalling to restrain neuroblast proliferation. Secondly, we find that blood flow onset triggers a transcriptional response to modify neuroblast metabolic status and is required for sensory neuron differentiation. In contrast, no role of sensory neurogenesis in vascular development is found, suggesting a unidirectional signalling from vasculature to sensory neuroblasts. Altogether, we demonstrate that the cranial vasculature constitutes a hitherto unrecognized niche component of the sensory ganglia that regulates the pace of their growth and differentiation dynamics.

Introduction

Proper interaction and understanding of our external world depend on specialized sensory organs and neurons from the peripheral nervous system (PNS). The cranial PNS consists of several ganglia with sensory bipolar neurons that transmit external sensory information to the brain. Sensory neurogenesis is achieved by a coordinated sequence of specification, delamination, transit-amplification of progenitors and terminal differentiation events (Alsina

et al., 2003; Begbie et al., 2002; Maier et al., 2014; Schlosser, 2010). Several intrinsic factors

such as Igf1, FGFs or neurotrophins participate in the proliferation, innervation and function

of cranial ganglia during embryogenesis and adulthood (Bilak et al., 2003; Camarero et al.,

2003; Fantetti and Fekete, 2012; Murdoch and Roskams, 2012; Vemaraju et al., 2012).

Mutations in some of these factors, therefore result in sensory pathologies, i.e. Igf1 mutations

are associated with hearing loss in rodents and humans (Bonapace et al., 2003; Riquelme et

al., 2010). The rate of sensory neurogenesis is determined by the balance between

multipotent stem cells (SCs) and differentiated sensory neurons and the combination of

intrinsic factors and extrinsic cues of their microenvironment.

Recent evidences indicate that multipotent SC function *in vivo* is tightly linked to

communication with the components of the microenvironment in which they reside, the so

called niche (reviewed in Spradling et al., 2001). Blood vessels have emerged as key

participants of SC function in several niches, such as the adult neural, hematopoietic and

spermatogonia SC niches. Depending on the context, blood vessels regulate SC quiescence,

proliferation and/or differentiation (Palmer et al., 2000; Rafii et al., 2016; Shi and Gronthos,

2003; Yoshida et al., 2007).

The adult brain SC niche is one of the niches in which the blood vessel-neuronal interaction

was early revealed and, thus, the signals and mechanisms are well understood. There, two

types of neurovascular interactions take place, one mediated by direct cell-cell contacts

through neural SCs (NSCs) endfeet enwrapping endothelial cells (ECs) (Kacem et al., 1998)

and the other mediated by secreted signals from ECs (Shen et al., 2008; Tavazoie et al.,

2008)(Mirzadeh et al., 2008; Palmer et al., 2000; Shen et al., 2008; Tavazoie et al., 2008). Cell-

cell contact signalling by EphrinB2 and Jagged1 expressed in ECs and, Eph and Notch receptors

in NSCs, regulates NSCs quiescence (Ottone et al., 2014). On the other hand, endothelia

secrete VEGFs, Angiopoietins, BDNF, NT-3 and PEDF that control NSC self-renewal, progenitor

cell proliferation or differentiation (Androutsellis-Theotokis et al., 2009; Calvo et al., 2011;

Delgado et al., 2014; Jin et al., 2002a; Louissaint et al., 2002; Ramírez-Castillejo et al.,

2006)(Androutsellis-Theotokis et al., 2009; Calvo et al., 2011; Delgado et al., 2014; Jin et al.,

2002b; Louissaint et al., 2002; Ramírez-Castillejo et al., 2006). Blood vessels not only

communicate through secreted factors, but also by the perfusion of oxygen and nutrients.

Oxygen tension mediates the metabolic switch from glycolysis to Oxidative Phosphorylation

(OxPhos), which has been shown recently to control NSC fate (Ito and Suda, 2014; Llorens-

Bobadilla et al., 2015). In the embryo, vasculogenesis and neurogenesis are spatiotemporally

coupled in the central nervous system (Hogan et al., 2004; Ulrich et al., 2011; Vasudevan et

al., 2008), but only few studies have addressed the molecular mechanisms underlaying this neurovascular interaction (Lange et al., 2016; Tan et al., 2016; Tata et al., 2016).

Cranial ganglia in vertebrates lie in very close proximity to head blood vessels (Taberner et al., 2018), suggesting that vasculature might be an essential component of their niche. Notwithstanding, this question and the mechanisms involved in their communication remain to be addressed in the sensory PNS.

In this study, by taking advantage of an avascular model and double labelling of ECs and sensory neurons in zebrafish, we identify two distinct mechanisms by which vasculature regulates cranial sensory neurogenesis. *In vivo* imaging of the developing sensory ganglia reveals that ECs and neuroblasts establish direct contacts through filopodial protrusions. Inhibition of the filopodial contacts show that they exert signalling functions over sensory neuroblast. Thus, the work establishes first evidences for neurovascular interaction mediated by cytonemes. Genetic and pharmacological perturbations show that cytoneme contacts and Dll4 signalling inhibit neuroblasts cell cycle entry and regulate quiescence at early timepoints. Later, blood flow initiation activates a transcriptional program linked to oxygen sensing by Egln2 and a metabolic switch to OxPhos, together with the promotion of cranial ganglia differentiation. Our results uncover, for the first time, that the cranial vasculature is an essential component of the sensory ganglia niche and can temporally regulate neuroblast quiescence and differentiation by using distinct signalling mechanisms.

Results

Vasculature regulates the number of SAG neuroblasts by restraining cell cycle entry of neuroblasts

We focused on neurovascular interactions in the statoacoustic ganglion (SAG), which comprise the sensory neurons that innervate mechanosensory hair cells of the inner ear and the brainstem. During neurogenesis, *neurog1*[+] neuronal precursors (blue) transit into neurod[+] neuroblasts (green) that delaminate and coalesce into the SAG. Later they differentiate into Islet2[+] neurons (orange) (Figure 1A). The developing zebrafish SAG lies between three main vessels, the Primordial Hindbrain Channel (PHBC) that runs between the hindbrain and the otic vesicle (OV), the Lateral Dorsal Aorta (LDA) beneath the OV and the Primary Head Sinus (PHS) surrounding laterally the OV (Taberner et al., 2018) (Figure 1B).

To functionally evaluate if cranial vasculature regulates the development of cranial sensory neurons in SAG, we analysed SAG growth dynamics in *cloche* (*clo*) homozygous mutants in the double *TgBAC(neurod:EGFP) [nl1];(Kdrl:ras-mCherry) [s896]* background, in which eGFP marks

sensory neurons and mCherry marks ECs. *Cloche* encodes *npas4l*, a PAS-domain-containing

bHLH transcription factor required for the formation of endothelial and hematopoietic cells

(Reischauer et al., 2016; Stainier et al., 1995) and *clo* mutants have extensively been used as

an avascular model (Liao et al., 1997; Sumanas et al., 2005). In *clo* mutant embryos, the SAG

appeared larger compared to siblings (sib) at 36 and 72 hpf (Figure 1C). Quantification of SAG

volumes revealed a statistically significant increase in *clo* mutant embryos (Figure 1D).

Moreover, the SAG displayed an abnormal shape and a significant expansion along its

mediolateral axis but not along the anteroposterior or dorsoventral axis (Figures 1C and 1E).

The increased volume in *clo* mutant embryos could reflect a loss of confinement of the SAG

cells by taking the space left by the lack of blood vessels or, alternatively, a higher number of

Neurod[+] cells. To distinguish between both possibilities, we quantified in serial transverse

sections the nuclei within the GFP-labelled SAG cells between 30 to 72 hpf in the *clo* mutant

and sibling embryos (Figures 1F and S1). In siblings, there is a period of steep SAG growth

between 30 to 36 hpf, (sib 30 hpf: 89 ± 5; sib 36 hpf: 128 ± 5). From 36 to 72 hpf, the mean

number of Neurod[+] cells is 105 ± 8, suggesting that after 36 hpf this population does not grow.

Remarkably, we found that the number of Neurod[+] cells was significantly higher in *clo* mutant

embryos compared to siblings at 36 hpf (sib: 128 ± 5 vs *clo*[-/-]: 177 ± 5) and this increase

persisted at all the following timepoints (Figure 1F). Several mechanisms could account for

the increased number of Neurod[+] cells in *clo* mutant embryos: i) increased specification of

neurog1[+] cells within the otic epithelium and thereby increased delamination of Neurod[+] cells,

ii) reduced differentiation of Neurod[+] cells into Islet2[+] neurons or iii) increased proliferation

of the transit-amplifying Neurod[+] population. We did not find a higher number of *neurog1*[+]-

expressing cells in the otic epithelium in *clo* mutant embryos compared to siblings (Figures 1G

and 1G') nor significant differences in the number of differentiated Islet2[+] cells in the SAG

(Figures 1H and 1H'). Instead, in *clo* mutant embryos we detected a significantly higher

number of proliferating Neurod[+] cells after a BrdU pulse from 24 to 30 hpf (expressed as

BrdU[+]/Neurod[+] ratio) (Figures 1I and 1I').

Hence, our results show that at early timepoints, cranial vasculature regulates the number of

SAG neuroblasts by negatively controlling cell proliferation of Neurod[+] cells and therefore

constitutes an essential component of the neurosensory niche during development.

Cranial blood vessels and sensory neuroblasts establish direct contacts through filopodia

In both the embryonic and adult brain, NSCs establish contacts with ECs through endfeet

projections (Ottone et al., 2014; Tan et al., 2016; Tavazoie et al., 2008). Therefore, we

wondered if neuroblasts from the SAG also establish direct contacts with the adjacent vessels.

To investigate this in detail, we live-imaged sensory neuroblasts from the SAG and ECs from

the PHBC in double *TgBAC(neurod:EGFP)* [nl1]*;(Kdrl:ras-mCherry)* [s896] reporter lines, at high

spatiotemporal resolution. At 30 hpf, we did not observe neuronal extensions enwrapping

ECs of the PHBC, but instead observed dynamic, thin filopodia extending from ECs to Neurod[+]

cells, as well as similar filopodia from Neurod[+] cells extending to ECs (Figures 2A, 2B, S2, S3

and Supplemental Movies S1-4). The dynamics of filopodia were varied; in some instances, we

observed filopodia extending and retracting in seconds (Figures 2A and 2B, cyan and magenta

arrowheads respectively), in others, filopodia remained stable for several minutes (Figures 2A

and 2B, yellow arrowheads). Interestingly, some PHBC filopodia did not remain at the surface

of the SAG but contacted deeper neuroblast somas (Supplemental Movie S4). Moreover,

stable filopodia-filopodia contacts between neuroblasts and ECs were also seen in some cases

(Figures 2C and S3). As observed in transverse optical confocal reconstructions of the vessels

and the SAG, filopodia were highly directional: PHBC filopodia extended only ventrally in the

direction of sensory Neurod[+] cells (Figures 2D and 2E). Interestingly, filopodia were present

from 24 to 36 hpf but not later, suggesting an early role of filopodia in vessel-neuroblast

communication (Figure 2F). Altogether, highly directed and temporally regulated filopodia-

mediated cell contacts are established between both EC and sensory neuroblasts.

Sensory neurons are not required for cranial vascular patterning

Communication between neurons and vessels in the CNS can take place from vessels to

neurons, vice-versa and bidirectionally (Barber et al., 2018; Makita et al., 2008; Mukouyama

et al., 2002). Thus, we next investigated if sensory neuron development has a role in vascular

patterning near sensory ganglia. In this case, we analysed homozygous *neurog1*[hi1059] mutant

embryos -which fail to specify sensory neuronal precursors (Ma et al., 1998)- in the double

transgenic *TgBAC(neurod:EGFP)* [nl1]*;(Kdrl:ras-mCherry)* [s896] reporter line. Strikingly, despite the

total absence of SAG and other cranial ganglia in *neurog1* mutant embryos, development of

the PHBC and LDA displayed no obvious patterning defects at 30 hpf (Figure 3A). The central

arteries (CtAs) ingrowth into the hindbrain also did not display any delay nor obvious

alteration at 48 hpf (Figure 3A). We quantified the width of PHBC and the number of ECs at

48 hpf, and again found no significant differences between *neurog1* mutants and siblings

(Figures 3B and 3C). In conclusion, our data indicate that cranial vascular development does

not require signals from sensory neuroblasts and suggest that primarily the signalling direction

goes from vessels to neurons.

Since imaging revealed directed filopodial contacts from cranial vessels to sensory neurons

and vice-versa (Figure 2A), we wondered about possible growth interdependency. We imaged

neuronal filopodia dynamics in the absence of vessels and, endothelial filopodia in the

absence of neurons. In *clo* mutants, the number of neuronal filopodia was drastically reduced

(Figures 3D and 3D'). In contrast, in *neurog1* mutant embryos the number of ECs filopodia did

not change, though they were shorter (Figures 3E and 3E''). These data indicate that signals

from ECs are required for neuronal filopodia formation.

While sensory neurons in SAG do not appear to be required for vascular development, our

data indicate that vasculature is required for SAG neurons to proliferate during a temporal

window that coincides with the period of filopodial contacts. Moreover, neuroblasts only

extend filopodia when ECs are present. Altogether, the results support the hypothesis that

endothelial-neuronal filopodia contacts are required for vessels to signal to neuroblasts in the

SAG to restrict growth.

Cell signalling mediated by cytonemes regulates the number of SAG neuroblasts

Cell signalling through filopodia has emerged as a novel mechanism of signalling between

adjacent cells in different embryonic tissues (Kornberg, 2017; Stanganello and Scholpp, 2016;

Stanganello et al., 2015). Signalling filopodia have been named cytonemes (Ramírez-Weber

and Kornberg, 1999). In order to assess the possible role of filopodial signalling in SAG growth,

filopodia formation was inhibited with LatrunculinB (LatB), a toxin that binds to actin and has

been used to impair filopodia function *in vivo* (Morton et al., 2000; Phng et al., 2013). Imaging

and quantification of filopodia in embryos treated with LatB for 6 hours indicated that the

treatment was effective, the number of filopodia in ECs and in neuroblasts was reduced

(Figures 4A and 4B). Interestingly, inhibition of filopodia caused an increased number of

Neurod[+] cells in LatB-treated embryos (Figure 4C) and this phenotype was also due to a higher

number of Neurod[+] cells incorporating BrdU (Figures 4D and 4E). No additional effects were

found in treated *clo* mutant embryos, indicating specific effects of neuroblasts (Figure 4C).

Hence, filopodia formation inhibition at early stages recapitulated the SAG growth phenotype

found previously in the avascular mutant. These data demonstrate that the endothelial-

neuronal filopodial contacts are cytonemes exerting a signalling role onto SAG neuroblast

proliferation.

Dll4 inhibition increases the Neurod[+] population

To identify signals that could mediate the neurovascular crosstalk we revisited the published vascular signals regulating CNS neurogenesis for evidence of ligand expression in cranial vasculature and their receptors in cranial sensory neurons (Howe et al., 2013). In the Ventricular-Subventricular Zone (V-SVZ) of the adult brain, endothelial Jagged1 ligand restrains NSC proliferation (Ottone et al., 2014). Cranial vessels do not express Jagged1 but do express Delta-like 4 (Dll4) (Fujita et al., 2011), another Notch ligand, which is essential for proper vascular remodelling and sprouting (Bussmann et al., 2011; Leslie et al., 2007; Siekmann and Lawson, 2007). We therefore hypothesized that Dll4 could be signalling to SAG neurons, which express Notch1 at the same early timepoints (Nikolaou et al., 2009). Moreover, it has been postulated that pigmentation patterning in zebrafish is regulated by DeltaC-Notch signalling through filopodia (Hamada et al., 2014).

To test this hypothesis, we inhibited *dll4* through the well-studied *dll4* morpholino (MO) (Bussmann et al., 2011; Siekmann and Lawson, 2007) and assessed its effects on SAG growth. Remarkably, *dll4* morphant embryos presented an expansion of the Neurod[+] population similar to the one observed previously in *clo* mutant embryos (Figure 5A). This expansion was reflected both in SAG volume (Fig 5B) as well as in Neurod[+] cell number (Figure 5C). As a further control, *dll4* morpholino was also injected in *clo* mutant embryos and the number of Neurod[+] cells did not change, suggesting that the results are due to loss of *dll4* expressed by the vasculature (Figures 5A-5C). These findings reveal that Dll4 is required to regulate Neurod[+] cell number.

In angiogenic sprouts, the number of filopodia increases after *dll4* downregulation (Suchting et al., 2007). When we assessed the number of filopodia in the developed PHBC, we detected a small increase in filopodial number and no differences in their length (Figures 5D-5F). Thus, the increase of Neurod[+] cells in *dll4* morphants cannot be attributed to a lack of filopodial processes but to a specific lack of signal. Finally, the number of ECs did not change in *dll4* morphant embryos compared to embryos injected with a random morpholino (Figures 5G and 5H). Therefore, the increase in Neurod[+] cell number after *dll4* down-regulation is not due to vascular developmental changes but to Dll4-Notch1 signalling.

Together, these results suggest that Dll4 signalling constitutes one of the key molecular mechanisms by which the vasculature regulates cranial sensory neurogenesis.

Absence of vasculature impairs sensory neuronal differentiation

Since more neuroblasts form in the SAG in the absence of vasculature, we next assessed if

these neuroblasts differentiate, resulting in a higher number of Islet2[+] post-mitotic cells at

later stages. Unexpectedly we found, after plotting the number of immunostained Islet2[+]

nuclei from 30 to 96 hpf, that there was a significantly lower number of Islet2[+] cells in *clo*

mutants from 54 hpf onwards (Figures 6A and 6B). The differentiation curves indicate that,

while at early timepoints sensory differentiation occurs partially independently of vasculature,

from 54 hpf onwards, cranial vasculature is required to promote differentiation of Neurod[+]

cells to Islet2[+] cells. As expected from the immunostaining data, the fluorescence from sensory

axons was also reduced in *clo* mutant embryos, labelled by the *Tg(isl2:GFP)* [zc7] reporter line

which marks *in vivo* the cytoplasm of differentiated sensory neurons (Pittman et al., 2008)

(Figure 6C).

To determine if cranial vasculature regulates neurogenesis in other sensory ganglia besides

the SAG, we assessed for filopodia contacts and quantified the number of Neurod[+] and Islet2[+]

cells in different ganglia. Here, we did not perform a temporal analysis, but we chose discrete

timepoints. At 30 hpf, we could also detect filopodia from the PHBC to the trigeminal and

anterior lateral line ganglia (Figure S4C). At 36 hpf, we found a significantly increased number

of Neurod[+] cells in the trigeminal ganglion of *clo* mutants (Figures S4A and S4B), whereas at

72 hpf there was a reduced number of Islet2[+] cells in the trigeminal and vagal ganglia (Figures

S3D-S3G). Therefore, the neurovascular interaction appears to extend to other sensory

ganglia.

To complete the analysis, we assessed if vasculature could regulate the development of other

cell types such as the hair cells of the inner ear. For this aim, we generated *clo* homozygous

mutants in the *Tg(brn3c:mGFP)* [s356t] background that labels the membranes of hair cells and

their kinocilia (Xiao et al., 2005). Serial confocal stacks taken at 72 hpf and quantification of

hair cells did not reveal significant differences in hair cell number in cristae nor maculae of *clo*

mutants compared to siblings (Figures 6D-6F). Therefore, vasculature is specifically affecting

neuronal differentiation of the inner ear but not hair cell number.

Initiation of blood flow and oxygen sensing triggers sensory differentiation

It has been reported that in zebrafish blood flow begins due to the completion of anastomosis

of cranial blood vessels around 48 to 60 hpf (Siekmann et al., 2009). The coincidence of this

timing with the one of sensory differentiation defects in *clo* mutant embryos prompted us to

explore whether the inducing signal of sensory differentiation was linked to blood flow. To

this aim, we incubated embryos with Nifedipine, a drug that inhibits heartbeat and blood flow

(Li et al., 2008). As expected from blood flow inhibition, we observed a drastic vessel diameter

reduction in Nifedipine-treated embryos (Figures 7A and 7B). In embryos treated with

Nifedipine from 54 to 72 hpf, the number of Islet2$^+$ cells decreased, recapitulating the defects

found in *clo* mutant embryos (Figures 7C-7E). Treatment in the avascular model did not further

enhance the phenotype, suggesting that the differentiation arrest is mostly attributed to the

absence of blood flow in *clo* mutant embryos and not associated to side effects of Nifepidine

(Figure 7E). The same results were obtained when treatment was done with BDM, another

drug used to inhibit blood flow (Anderson et al., 2008) (Figure S5C and S5D). To narrow down

the time window of blood flow requirement, we incubated embryos with Nifedipine from 60

to 72 hpf and from 48 to 60 hpf. Incubation from 60 to 72 hpf did not inhibit differentiation

(Figures 7F and 7G), whereas incubation from 48 to 60 hpf did (Figures S5A and S5B). These

results indicate that blood flow induction of sensory differentiation takes place in a narrow

time window and once signalling is initiated, blood flow is no longer required.

To obtain insight into the molecular mechanisms underlying the arrest in sensory neuron

differentiation in the absence of vasculature, we conducted a transcriptional profiling of

freshly sorted Neurod GFP$^+$ SAG cells from *clo* mutant and sibling embryos at 54 hpf. RNA-seq

analysis revealed a transcriptional profile response in the absence of vasculature with 641

genes significantly downregulated and 15 upregulated in clo mutant compared to control

embryos (Figure 7H). This suggest that in absence of blood flow some signalling pathways are

shut down rather a new transcriptional program being activated, what matches with a cell

differentiation arrest. Whereas we expected pathways related to differentiation signals or

cell-cycle control being down-regulated, most differentially expressed genes found where

genes related with oxygen sensing (egln2), reduction-oxidation (redox) process (rdh8a) and

mitochondria metabolism (rmnd1 and abat) (Figures 7H ans 7I). Thus, the first transcriptional

response to blood flow onset seems to be related to oxygen sensing and switch from glycolysis

to OxPhos that ultimately drives sensory neuronal differentiation, as observed in other

contexts (Hawkins et al., 2016; Homem et al., 2014; Kida et al., 2015; Lange et al., 2016; Ryall

et al., 2015).

Altogether, the results define for the first time a prominent role of cranial vasculature in

sensory neurogenesis that can be segregated into two sequential and independent

mechanisms. A first one, in which Dll4-Notch1 cytoneme signalling from ECs to neurons

regulates neuroblast quiescence, and a second one in which initiation of blood flow and

oxygen sensing induces a metabolic switch and sensory neuron differentiation.

Discussion

The SAG and other cranial sensory ganglia reside in a vascularized milieu in embryos and adults. A fundamental question in developmental neurobiology is whether or not vasculature, in addition to transport of nutrients and oxygen, provides instructive cues for sensory neurogenesis. This is relevant to understand hearing loss and other sensory neuropathies linked to vascular defects (Jensen et al., 2004; Lago et al., 2018; Paul et al., 2016). Our detailed temporal analysis reveals that cranial ganglia indeed require vascular signals for proper growth and differentiation. We identified two segregated mechanisms: an early cytoneme-based, cell-cell contact dependent mechanism regulating neuroblast proliferation and a later blood flow-dependent mechanism controlling neuronal differentiation.

In the V-SVZ, quiescent NSCs physically contact blood vessels with specialized endfeet (Kacem et al., 1998). By taking advantage of the imaging and cell reporter tools available in the zebrafish model we resolve, for the first time, that cell-cell communication between vasculature and neuroblasts takes place through dynamic cytoneme extensions that project between them. It is still possible that in other contexts, dynamic contacts also exist, but imaging in fixed tissue might have hampered their identification, as suggested by Obernier and colleagues (Obernier et al., 2018). Filopodia in angiogenic tip cells and migrating ECs have been described (Fantin et al., 2015; Gerhardt et al., 2003; Ochsenbein et al., 2016; Phng et al., 2013). To our knowledge, filopodia from endothelial stalk cells contacting neuronal cells have not been reported before. One could envision cell-cell contact with only ECs extending filopodia to neuronal somas but, in our system, both cell types project directional filopodia. Signalling filopodia or cytonemes in *Drosophila* and vertebrates have been demonstrated, extending the cell-cell signalling distance of influence (Kornberg, 2017; Stanganello and Scholpp, 2016; Stanganello et al., 2015). This is relevant in the facial mesenchyme where vasculature does not grow inside the sensory ganglia but adjacent. Different types of cytoneme contacts have been described in previous studies, in some cases cytonemes contact the cell body of their target cell (Huang and Kornberg, 2015; Stanganello et al., 2015), while in other cases cytoneme-cytoneme contacts are observed (Du et al., 2016; González-Méndez et al., 2017; Huang and Kornberg, 2015; Roy et al., 2014; Sanders et al., 2013). It is difficult to tell if in our system cytoneme-cell body contacts occur, but we have clearly observed cytoneme-cytoneme contacts. These cytonemes have distinct behaviours, either highly transient or more stable, the latter possibly being actively signalling.

Moreover, the study highlights the requirement of signals from ECs for neuronal cytoneme growth. By blocking filopodia projections, we demonstrate that they have a signalling function

in the negative control of neuroblast cell proliferation. Most likely only a subset of neuroblasts

are contacted by endothelial cytonemes, allowing a scattered pattern of cell cycle-arrested

neuroblasts between others with proliferative capacities. Cytonemes have been described to

participate in stem cell maintenance in *Drosophila* (Fuwa et al., 2015; Inaba et al., 2015;

Mandal et al., 2007; Rojas-Ríos et al., 2012), but not yet described to participate in a NSC niche

in vertebrates.

In contrast to other contexts in which neurons control vascular patterning (Himmels et al.,

2017), our results suggest that in the context of sensory ganglia and surrounding vessels, an

absence of cranial ganglia does not have a major effect on early vasculogenesis.

During adult neurogenesis, cell contact-dependent neurovascular signalling is mediated by

ephrinB2 and Jagged1 present in ECs that suppress cell-cycle entry in NSCs (Ottone et al.,

2014). We propose that in the PNS Dll4, but not Jagged1, plays a similar role. Maintenance of

a quiescent and undifferentiated population of neural progenitors is key for life-long

regeneration. Thus, the role we have uncovered for cranial vasculature and Dll4 in the cranial

sensory niche might be relevant in regenerative studies upon neuronal damage.

Several studies have highlighted the role of blood flow initiation in remodelling the

vasculature, VEGF expression, angiogenesis and stimulation of hematopoietic stem cell

formation (Adamo et al., 2009; Bussmann et al., 2011; Nicoli et al., 2010). Blood flow can

exert tangential and perpendicular forces on ECs leading to actomyosin rearrangements, YAP-

TAZ nuclear translocation and the induction of the mechano-sensitive transcription factor

Klf2a (Nakajima et al., 2017; Nicoli et al., 2010; Le Noble et al., 2008). Ultimately, blood flow

triggers gene transcriptional changes in ECs and vessel maturation (Chiu and Chien, 2011).

On the other hand, blood vessels can exert their roles on neural cells via delivery of nutrients

or oxygen (Cleaver and Dor, 2012; Ramasamy et al., 2015). The transcriptome analysis of SAG

neuroblasts in clo mutant vs control embryos at the timepoint of blood flow initiation allowed

us to identify *egln2* (*Egl nine homolog 2*) as one of the early-response genes more differentially

expressed. Egln2 is cellular oxygen sensor involved in the degradation of the transcription

factor hypoxia-inducible factor (*HIF1*) (Epstein et al., 2001), thus in wild-type conditions

embryos sense the passage from hypoxic to normoxic conditions. Additionally, genes related

to redox state and mitochondria metabolism are also differentially expressed. Our data

suggest that SAG sensory precursors switch from glycolytic to OxPhos to induce their

differentiation. Numerous populations of multipotent SC, i.e hemaepoietic SC, intestinal crypt

SC and neural SC, use a glycolytic program in their niche and switch to OxPhos upon

regeneration and differentiation (Cliff and Dalton, 2017; Lange et al., 2016; Mohrin et al.,
2015; Shi et al., 2015).

Our current work identifies novel mechanisms for head vasculature in the balance between
sensory precursor and differentiation fates and shall contribute to our basic understanding of
the cranial sensory niche in which head vasculature enters in scene.

Acknowledgments

415 We thank D Stainer for the *Tg(cloche)* ^{m378} and *Tg(kdrl:ras-mCherry)* ^{s896} lines, the PRBB Imaging
416 Facility and the UPF Genomics Core and Flow Cytometry Facilities for technical support, T.
417 Schilling, A. Bigas and I. Fariñas for comments on the manuscript and lab members for
418 discussions. The work was supported by BFU2014-53203 and BFU2017-82723P from MCINN
419 to BA and the Unidad de Excelencia Maria de Maeztu (MDM-2014-0370).

Author Contributions

422 LT and BA designed research, LT and AB performed experiments, LT, AB and BA analysed data
423 and TL and BA wrote the manuscript.

Declaration of Interests

426 The authors declare that they have no conflict of interest.

Figure Legends

Figure 1 In the absence of vessels, the Neurod$^+$ population in the SAG is expanded due to
increased proliferation but not to effects on specification or differentiation.

(A) Schematic drawing of the sequential steps of otic neurogenesis. Coloured graphs

representing effects in specific cell types will follow this scheme: neurog1⁺ neuronal

committed progenitors in blue, Neurod⁺ neuroblasts in green, Islet1/2⁺ neurons in orange and

blood vessels in magenta.

(B) Lateral (left) and transversal (right) 3D drawing of the OV (dotted line), the SAG and the adjacent vessels: PHBC, LDA and PHS.

(C) Representative anterior and posterior transverse images of the SAG in sib and clo$^{-/-}$ *TgBAC(neurod:egfp)* ^{nl1} embryos of cryostat sections, at 36 and 72 hpf. Neurod$^+$ cells: green, neuroblasts; DAPI: blue, nuclei.

(D) Quantification of SAG volumes in sib and clo$^{-/-}$ embryos at 36 and 72 hpf (n=8).

(E) Graph showing measurements of the anteroposterior (AP), dorsoventral (DV) or mediolateral (ML) axis length of the SAG in sib and clo$^{-/-}$ embryos (n=8-10), at 72 hpf.

(F) Quantification of Neurod$^+$ nuclei in the SAG visualized through the *TgBAC(neurod:egfp)* ^{nl1} of sib and clo$^{-/-}$ embryos at different timepoints from 30 to 72 hpf (n = 6-10).

(G) Transversal images of *neurog1 in situ* hybridization in the OV epithelium in sib and clo$^{-/-}$ embryos, at 24 hpf; and quantified in G' (n = 7).

(H) Confocal lateral images of immunostained Islet 1/2$^+$ nuclei in the SAG (dotted line, visualized by *TgBAC(neurod:egfp)* ^{nl1}) in sib and clo$^{-/-}$ embryos, at 36 hpf; and quantified in H' (sib n = 10, clo$^{-/-}$ n = 8).

(I) Transversal images of BrdU incorporation (magenta) in the SAG (continuous line, visualized by *TgBAC(neurod:egfp)* ^{nl1}) in sib and clo$^{-/-}$ embryos, at 30 hpf; and quantified in I' (n = 6). DAPI: blue, nuclei.

Scale bar, 50 µm. Error bars, mean ± SEM for D,E,G',H',I'; and mean ± SD for F. Unpaired two-tailed Student's *t*-test. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$. ns, non-significant. OV, otic vesicle; SAG, statoacoustic ganglion; PHBC, primordial hindbrain channel; LDA, lateral dorsal aorta; PHS, primary head sinus; HB, hindbrain; sib, siblings.

Figure 2 Endothelial cells and SAG neuroblasts contact through directional filopodia.

(A,B) Timelapse of lateral images of *TgBAC(neurod:egfp)* [nl1]; *(Kdrl:ras-mCherry)* [s896] embryos

showing filopodia extending from the PHBC to the SAG and from the SAG to the PHBC at 30

and 36 hpf (videos are representative of 5 experiments). Cyan, magenta and yellow

arrowheads point to examples of growing, reducing and stable filopodia, respectively.

(C) 4 μm thick z-projection exemplifying a filopodia-filopodia contact remaining stable for

more than 45 seconds, 30 hpf.

(D) Single plane images in lateral view and transversal optic reconstruction showing a

filopodial process from the PHBC to the SAG, at 30 hpf. Arrowhead indicates point for

transverse optical reconstruction.

(E) Rose plot of PHBC filopodia frequency of angle distribution and length (0º, lateral; 90º,

dorsal; 180º, medial; 270º, ventral (n=76 filopodia, 15 embryos), at 30 hpf.

(F) Histogram showing the number of filopodia in the PHBC (magenta) and the SAG (green) at

different timepoints from 24 to 48 hpf (n = 5).

Scale bars, 20 μm in A,B,D and 10 μm in C. SAG, statoacoustic ganglion; PHBC, primordial

hindbrain channel.

Figure 3 Lack of neurogenesis does not affect cranial vasculature patterning nor filopodia formation, while loss of vasculature prevents neuroblasts filopodia formation.

(A) Representative lateral images of cranial vasculature in sib and *neurog1⁻/⁻* homozygous mutant embryos in *TgBAC(neurod:egfp) ᵘˡˢ¹, (Kdrl:ras-mCherry) ˢ⁸⁹⁶* background, at 30 hpf and 48 hpf.

(B) Quantification of EC numbers in sib and *neurog1⁻/⁻* (n = 12), at 48 hpf.

(C) Quantification of PHBC width in sib and *neurog1⁻/⁻* (n = 10), at 48 hpf.

(D) Confocal lateral images showing neuroblasts filopodial processes visualized through the *TgBAC(neurod:egfp) ⁿˡ¹* in sib and clo ⁻/⁻, at 30 hpf.

(D') Graph of SAG neuroblasts filopodia number in sib and clo ⁻/⁻ (n = 8), at 30 hpf.

(E) Confocal lateral images showing PHBC filopodial processes visualized through the *Tg(Kdrl:ras-mCherry) ˢ⁸⁹⁶* in sib and *neurog1⁻/⁻* embryos, at 30 hpf. Arrowheads indicate filopodia in D and E.

(E') Graph of PHBC filopodia number in sib and *neurog1⁻/⁻* (n = 18), at 30 hpf.

(E'') Graph of PHBC filopodia length in sib (n = 50) and *neurog1*[-/-] (n = 48), at 30 hpf.

Scale bars, 50 µm in A and 20 µm in D,E. Error bars, mean ± SEM. Unpaired two-tailed

Student's *t*-test for B, C, D', E' and unpaired one-tailed Student's *t*-test for E''. * p < 0.05, ** p

< 0.01, *** p < 0.001. ns, non-significant. PHBC, primordial hindbrain channel; LDA, lateral

dorsal aorta; AAs, aortic arches; PHS, primary head sinus; CtAs, central arteries; SAG,

statoacoustic ganglion; EC, endothelial cells; sib, siblings.

Figure 4 Inhibition of cytoneme processes by Latrunculin B results in expansion of

neuroblasts.

(A) Confocal lateral images showing filopodial processes in vessels and neuroblasts visualized

through *TgBAC(neurod:egfp)* [nl1] and *(Kdrl:ras-mCherry)* [s896] respectively, after 6h treatment in

DMSO or LatB, at 30 hpf. Arrowheads indicate filopodia.

(B) Quantification of filopodial processes number in vessels (n = 13-16) and neuroblasts (n =

12) after 6 hours incubation with LatB compared with control (DMSO), at 30 hpf.

(C) Graph showing the number of Neurod+ nuclei at 36 hpf, using the *TgBAC(neurod:egfp)* [nl1]

background and DAPI staining, in sib and clo [-/-] embryos treated with DMSO or LatB from 24-

30 hpf and washed until 36 hpf (n = 12).

(D) Lateral images of BrdU incorporation (magenta) in Neurod+ cells of the SAG (dotted line),

visualized through *TgBAC(neurod:egfp)* [nl1], in 6 hours DMSO and LatB treated embryos, at 30

hpf.

(E) Graph showing the ratio of BrdU+ cells in the Neurod+ population after 6 hours DMSO (n =

9) and LatB (n = 8) treated embryos, at 30 hpf.

Scale bars, 20 μm in A and 50 μm in D. Error bars, mean ± SEM. In B,E unpaired two-tailed

Student's *t*-test was performed. In C the data was analysed using one-way ANOVA, followed

by post-hoc test. * p < 0.05, ** p < 0.01, *** p < 0.001. PHBC, primordial hindbrain channel;

sib, siblings.

Figure 5 Down-regulation of Dll4 signalling induces neuroblast expansion.

(A) Representative transverse sections showing the SAG in sib and clo-/- *TgBAC(neurod:egfp)*

nl1 embryos injected with of random or *dll4* splicing morpholino. Neurod+ cells: green,

neuroblasts; DAPI: blue, nuclei.

(B) Graph showing SAG volume quantification in sib and clo-/- embryos injected with random

or *dll4* splicing morpholino (n = 10), at 36 hpf.

(C) Graph showing quantification of Neurod$^+$ nuclei in sib and clo$^{-/-}$ embryos injected with random or *dll4* splicing morpholino (n = 14), at 36 hpf.

(D) Confocal lateral images of PHBC filopodial processes in random and *dll4* morphants visualized with the *Tg(Kdrl:ras-mCherry)*^{s896}, at 30 hpf. Arrowheads indicate filopodia.

(E,F) Quantification of PHBC filopodia number (n = 14) (E) and length (n = 14) (F), in random and *dll4* morphants, at 30 hpf.

(G) Confocal lateral images showing EC membranes and cytoplasm using the *Tg(Kdrl:ras-mCherry)*^{s896} and *Tg(flk1:egfp)*^{s843} respectively, in the PHBC, at 30 hpf.

(H) Quantification of EC number in random and *dll4* morphants (n = 14), at 30 hpf.

Scale bars, 50 μm in A, and 20 μm in D,G. Error bars, mean ± SEM. In B,C the data was analysed using one-way ANOVA, followed by post-hoc test. In E the non-parametric Mann-Whitney *U* test was performed. In F,H unpaired two-tailed Student's *t*-test was used. * p < 0.05, ** p < 0.01, *** p < 0.001. HB, hindbrain; OV, otic vesicle; SAG, statoacoustic ganglion; PHBC, primordial hindbrain channel; EC, endothelial cell.

Figure 6 Sensory neuronal differentiation requires blood vessels.

(A) Graph showing the number of Islet 2$^+$ nuclei in the SAG at different timepoints from 30 to 96 hpf (n = 4-10).

(B) Confocal lateral images of immunostained Islet2$^+$ nuclei in the SAG (dotted line, visualized by *TgBAC(neurod:egfp)*^{nl1}) in sib and clo$^{-/-}$ embryos at 72 hpf. Arrowheads indicate pm.

(C) Transverse sections showing axon density (arrowhead) in the *Tg(isl2:gfp)* ^{zc7} in sib and clo$^-$

$^{/-}$ embryos, of cryostat sections, at 72 hpf. Dotted line indicates OV and continuous line the

SAG.

(D,E) Graph showing quantification of hair cell number in cristae (D) and maculae (E) in sib (n

= 8) and clo$^{-/-}$ (n = 9) embryos.

(F) Representative lateral images of hair cells in cristae and maculae using the *Tg(brn3c:mGFP)*

^{s356t}, at 72 hpf. Dotted line indicates otic vesicle. Non-labelled hair cells are neuromasts.

Scale bars, 50 μm. Error bars, mean ± SD for A; and mean ± SEM for D,E. Unpaired two-tailed

Student's *t*-test. * p < 0.05, ** p < 0.01, *** p < 0.001. ns, non-significant. OV, otic vesicle;

SAG, statoacoustic ganglion; ac, anterior crista; lc, lateral crista; pc, posterior crista; am,

anterior macula; pm, posterior macula; sib, siblings.

Figure 7   Blood flow triggers transcriptional changes related to oxygen sensing and

metabolism together with sensory neuronal differentiation in a restricted temporal

window.

(A) Representative images of the PHBC of control and 48-54 hpf Nifedipine treated *Tg(Kdrl:ras-*

mCherry) [s896] embryos, at 54 hpf, in lateral (right) and transversal optic reconstructions (left)

in the point marked by the arrowhead.

(B)   Analysis of PHBC width in control and Nifedipine treated *Tg(Kdrl:ras-mCherry)* [s896]

embryos, at 54, 60 and 72 hpf with treatment starting at 48 hpf (n=15).

(C-G) Confocal lateral images showing immunostained Islet2[+] nuclei in the SAG (dotted line,

visualized by *Tg(neurod:egfp)* [nl1]) in sib and clo[-/-] embryos at 72 hpf in control (untreated) (C),

54-72 hpf Nifedipine treated embryos (D), 60-72 hpf Nifedipine treated embryos (F), and

quantified in E,G. 72 hpf sib (n = 10), 72 hpf clo$^{-/-}$ (n = 9), 54-72 hpf sib Nife (n = 9), 54-72 clo$^{-/-}$

Nife (n = 6), 60-72 hpf sib Nife (n = 9), 60-72 hpf clo$^{-/-}$ Nife (n = 9).

(H)   Volcano plot of differentially expressed genes (DEGs) in the SAG at 54 hpf. Red and blue

plots represent significant downregulated and upregulated DEGs, respectively (P<0.05; FC<-

1; FC>1), while black plots represented insignificant DEGs (P≥0.05; FC≥-1; FC≤1). Genes of

special interest are highlighted in green.

(I) Histogram showing the fold change of DEGs of special interest in clo$^{-/-}$ versus sibling SAGs

at 54 hpf.

Scale bars, 50 μm for A,B,D,F, and 20 μm for H. Error bars, mean ± SEM. Unpaired two-tailed

Student's *t*-test. * p < 0.05, ** p < 0.01, *** p < 0.001. ns, non-significant. pm, posterior

macula; Nife, Nifedipine; sib, siblings; PHBC, primordial hindbrain channel.

STAR Methods

Zebrafish Strains and Genotyping

Zebrafish (*Danio rerio*) were maintained as previously described (Westerfield, 2000) at the

aquatic facility of the Parc de Recerca Biomèdica de Barcelona (PRBB). The following

transgenic lines in different combinations were used: *Tg(neurog1)* ^{hi1059} (Golling et al., 2002)

mutant to inhibit neurogenesis and *Tg(cloche)* ^{m378} (Stainier et al., 1996) as an avascular model.

Five different reporter lines were used: *Tg(Kdrl:ras-mCherry)* ^{s896} (Chi et al., 2008) and

Tg(flk1:EGFP) ^{s843} (Jin et al., 2005) for vessels , *TgBAC(neurod:EGFP)* ^{nl1} (Obholzer et al., 2008)

for neuroblasts, *Tg(isl2:gfp)* ^{zc7} (also called Isl3) (Pittman et al., 2008) for sensory differentiated

neurons and *Tg(brn3c:mGFP)* ^{s356t} (Xiao et al., 2005) for hair cells .

Embryos were staged as previously described (Kimmel et al., 1995). Adult heterozygous for

neurog1 ^{hi1059} allele were identified by PCR genotyping by fin-clip genomic DNA(Golling et al.,

2002). Homozygous *neurog1* mutant embryos were recognized by the significant reduction of

GFP signal in *Tg(neurog1)* ^{hi1059}, *BAC(neurod:EGFP)* ^{nl1}, *(Kdrl:ras-mCherry)* ^{s896} background.

Avascular mutants were identified by the lack of vasculature in the reporter line for

endothelial cells *Tg(Kdrl:ras-mCherry)* ^{s896} or *Tg(flk-1:EGFP)* ^{s843}, lack of blood cells and

circulation (visible under scope from 30-36hpf) and atrium enlargement, as described by

Stainer et al. (Stainier et al., 1995).

In both cases, wild-type and heterozygous-deficient siblings were undistinguishable and both

have been used and referred as siblings (sib). Finally, since *clo* mutant embryos die around

7dpf (Stainier et al., 1996), we studied embryos until 4 dpf to exclude vessel directed effects

from embryonic morbidity.

Live imaging, confocal microscopy and quantifications

Images of cryosections were acquired with a Leica DFC 7000 T microscope, and 40x oil

immersion (0.75 NA).

Whole-mount embryos imaging was performed on a Leica SP8 confocal microscope, with 20x

glycerol immersion lens (0.7 NA). Embryos were mounted in 1% low-melting agarose

(Ecogene) in PBS or Danieau's solution, for fixed and life embryos respectively, and placed

onto a glass-bottom dish. For life embryos, 0.1% Tricaine (Sigma) was also added. The 488-

and 561-nm laser lines were employed at a scan speed of 600 Hz.

For time-lapse imaging of filopodial, the resonant mode (8000 Hz) was used to achieve a high

spatiotemporal resolution. Frames were captured simultaneously with 488 nm and 561 nm

lasers at a time interval of 15 seconds. Z-size of 2 µm was used for videos and 0.5 µm for static

stacks and transverse optical reconstructions. Confocal stacks and movies were flattened by

maximum projections using FIJI (Image J).

Measurements were done manually with FIJI (Image J) for SAG and PHBC filopodia number,

length and orientation using one single static image, from a defined region of interest (ROI).

PHBC width was measured from maximum projection confocal images always at the same

point, between CtA 2 and 3. PHBC EC number was calculated by perinuclear green cytosolic

signalling from *Tg(flk-1:EGFP)* [s843] and EC contours from *Tg(Kdrl:ras-mCherry)* [s896] in a defined

ROI. When ROIs were used, the same area was used to compare samples. SAG anteroposterior

(AP) and dorsoventral (DV) axes length were measured from maximum projection confocal

lateral images. The mediolateral (ML) SAG axis was calculated from 5 serial cryosections, in

which the biggest measurement was used.

The number of Neurod[+] nuclei was counted manually using DAPI staining in the

TgBAC(neurod:egfp) [nl1] labelled cells. *neurog1[+]* cells were also counted manually after an ISH

and DAPI staining. The number of HC was calculated using the *Tg(brn3c:mGFP)* [s356t] and

counting kinocilia number.

The volume of the SAG was measured by calculating the area the 5 serial cryosections that

include all SAG and multiplying them with each section thickness (20µm).

Statistical Analysis

All the data in the work were first tested for normal distribution using the Kolmogorov-Smirnov test and the Levene's test for homogeneity of variances. For two-group comparison two-tailed Student's *t*-test was used or Mann-Whitney *U* test for non-parametric tests. When more than two samples were being compared, one-way ANOVA followed by post-hoc test was performed. Values are expressed as median ± SEM. Graphs were performed with PRISM 7 (Graph-Pad) software. Error bars represent SEM, but for Fig. 2d and Fig. 5a, were shaded error bars represent SD. ns= non-significant, *$p \leq 0.05$, **$p \leq 0.01$, ***$p \leq 0.001$.

Immunostaining and *in situ* Hybridization

Embryos at the desired stage were fixed with 4% paraformaldehyde (PFA) in PBS 0,1% Tween-20 (PBT) for 3 hours at RT or overnight at 4ºC. Fixed embryos were dehydrated, kept in 100% MeOH -20ºC 1h or overnight (O/N), and rehydrated. Embryos were permeabilized with Proteinase K 1:1000 in PBT, or cold Acetone for mouse anti-Islet1/2, and incubated with blocking solution (BS) for 2h at room temperature (RT) and with primary antibody O/N at 4ºC. Primary antibody was washed 5 times 10min, and secondary antibody was applied for 2h RT or O/N 4ºC.

Embryos were cryoprotected in 15% sucrose and embedded in 7.5% gelatine/15% sucrose. Blocks were frozen in 2-methylbutane (Sigma) to improve tissue preservation. Sagittal and transversal sections were cut at 20µm thickness in a Leica CM 1950 cryostat.

For BrdU incorporation experiments, dechorionated live embryos were incubated with 10mM BrdU, 10% DMSO in Danieau's solution for 6 hours. After this they were washed twice in clean Danieau's solution for 5 min prior 4% PFA fixation for 3 hours at RT. BrdU staining was performed in cryosections after 2N HCl DNA denaturation 1h RT. Anti-BrdU (bdbiosciences; mouse Cat: 555627, 1:100). Anti-GFP was used to enhance GFP signal from transgenic lines (Torrey Pines BioLabs Inc; rabbit Cat: TP401, 1:400). Anti-Islet1/2 was used to detect differentiated neurons (Cat: TP401; mouse Cat: 39.4D5, 1:200). Alexa-fluor 488 and 594 anti-rabbit and anti-mouse (Invitrogen, 1:400) were used as secondary antibodies. DAPI staining for cryosections was done at 1:10000 PBS for 5min, and 2x5min wash in PBS followed by mowiol mounting.

Whole-mount *in situ* hybridization was performed as previously described (Thisse et al., 2004) using *neurog1* digoxigenin-labelled probe.

Inhibitor treatments

To block actin polymerization in filopodia, dechorionated live embryos were incubated in 0,1µg/mL Latrunculin B (LatB) (Merk, Millipore), 0,4% DMSO in Danieau's solution for 6 hours (Phng et al., 2013); while control embryos were raised at 0,4% DMSO only. After treatment, embryos were washed twice in clean Danieau's solution for 5min followed by immediate confocal imaging for filopodia visualization or 4% PFA fixation. In other cases, embryos were washed in Danieau's solution until 36 hpf. It has been previously demonstrated that the concentration used of LatB specifically inhibits filopodial formation while it does not affect other cellular behaviours for up to 17 hours of treatment (Phng et al., 2013).

Blood flow was blocked through the use of two different drugs. Dechorionated live embryos were incubated for 6, 12 and 24 hours in 40µM Nifedipine or 20mM BDM (2, 3-butanedione-2-monoxime) (Sigma-Aldrich) in Danieau's solution, washed twice 5min and fixed in 4% PFA at the desired stage (Anderson et al., 2008; Li et al., 2008).

Morpholino knockdown experiments

To inhibit protein translation of Dll4, 10ng of random and *dll4* splicing morpholino antisense oligonucleotides (Gene Tools, Philomath, OR) were injected in the yolk of 1-2 cell stage embryos as previously described (Siekmann and Lawson, 2007). Efficiency of the knockdown was assessed by RT-PCR (data not shown).

FACS purification of GFP⁺ cells

Embryos of the transgenic line *Tg(neurod:gfp) [nl1]* were grown until 54 hpf, stage at which they were anesthetized with tricaine. The SAG was dissected cutting the embryos head after the eye and in the posterior end of the otic vesicle. The hindbrain was also removed to prevent the sorting of any neurod⁺ cells also present in the CNS. The presence of the SAG was confirmed under a fluorescent scope. The dissected tissue was kept in Medium 199, on ice, until we summed around 60 embryos per each condition siblings and *clo*. Cell were disaggregated using a protease solution (10x Trypsin, 0,5M EDTA, 1x PBS) for 20 min with mechanical help by both vortexing and doing up&down with a 200 µL pipete. Cell disaggregation was stopped with STOP solution (30% FBS, 2M CaCl2, 1x PBT). Cells were recovered by centrifugation and re-suspended in cell suspension medium (20% BSA, 0,5M EDTA, 1x PBS). Neurod⁺ cells were FACS purified based on GFP fluorescence utilizing a INFLUX Sorter. Dead cells were eliminated by DAPI staining (1:1000). Cells were collected in collection medium (1:1, PBS:FBS) and immediately used to RNA extraction.

Four replicates were used, always preparing one sample for siblings and another for *clo*

mutants each time, coming from the same breed.

RNA extraction and sequencing

RNA-seq was performed on FACS-sorted neuroblasts from the *Tg(neurod:egfp)* [nl1] zebrafish

embryos at 54 hpf. RNA from these cells was extracted using the PicoPure™ RNA Isolation

Kit (Thermo Fisher, KIT0214).

Cells from around 60 embryos were pooled to obtain enough RNA for library preparation.

cDNA was prepared with SMART-Seq® v4 Ultra® Low Input RNA Kit for Sequencing (Takara,

634888) and validated using Bioanalyzer High Sensitivity DNA Kit (Agilent, Cat. No. 5067-

4626). Between 66-306 pg of these cDNAs were used to prepare libraries using Nextera XT

DNA Library Prep Kit (Illumina, FC-131-1024). Libraries were validated using Bioanalyzer High

Sensitivity DNA Kit (Agilent, Cat. No. 5067-4626). Then, libraries were normalized and pooled

for sequencing using the NextSeq High Output (Illumina, 20024907) with 75 bp paired ends

run with 50 million expected reads per sample.

Mapped reads with the Zebrafish genome (version GRCz11) were obtained using the STAR

software (v2.5.2b, https://github.com/alexdobin/STAR), and the reads' counts within exons

with the HTSeq software (v0.9.1, https://htseq.readthedocs.io/en/release_0.11.1/). The DE

analysis   was   done   with   the   limma   package   in   R

(v3.38.3, https://bioconductor.org/packages/release/bioc/html/limma.html).

Supplemental Figures and Video Legends

736
737   **Figure S1 Protocol used for quantification of Neurod+ cell number.**

738
(A-C) For a series of latero-medial in sagittal or antero-posterior transversal (data not shown)
cryostat sections, the perimeter of the SAG was drawn using the green signal of

TgBAC(neurod:egfp) nl1 (B), subsequently green light was hidden and blue light from DAPI
staining was visualized (C) to manually count the number of cells (D).

Figure S2 EC from the PHBC and neuroblasts from the SAG extend filopodia towards each
other from 24 to 36 hpf.

747
(A-C) Still images of time-lapse videos of *TgBAC(neurod:egfp) nl1*; *(Kdrl:ras-mCherry) s896*
showing filopodia extending from the PHBC to neuroblasts and from neuroblasts to PHBC at
24 hpf (A), 30 hpf (B), and 36 hpf (C) (time-lapses are representative of 5 experiments), in a
lateral view.

Cyan, magenta and yellow arrowheads point to examples of growing, reducing and stable
filopodia, respectively.
754
755 Scale bar 50 µm. PHBC, primordial hindbrain channel.
756
757 **Figure S3 Filopodia from vasculature and neuroblasts contact each other at the same plane.**

758
759 (A) Confocal lateral image of a single plane showing a filopodia-filopodia contact between EC
760 and neuroblasts. Square shows magnified image in the right, and arrowhead the mentioned
761 contact.
762
763 (B) Confocal lateral image of a single plane and transverse optical reconstruction also in single
764 plane showing a filopodial process from the neuroblasts contacting the EC. Arrowhead
765 indicates point for transverse optical reconstruction.
766
767 Scale bar 20 µm. PHBC, primordial hindbrain channel; SAG, statoacoustic ganglion; HB,
768 hindbrain.
769
770
771 **Figure S4 Filopodial contacts and defects on sensory neurogenesis in absence of vasculature**
772 **in other sensory ganglia.**

773
774 (A) Confocal lateral images of Tg and aLL ganglia in sib and clo[-/-] *TgBAC(neurod:egfp)* [nl1]
775 embryos, at 36 hpf. Neurod[+] cells: green, neuroblasts; DAPI: blue, nuclei.
776
777 (B) Quantification of Neurod[+] cells in the Tg of sib and clo[-/-] embryos, at 36 hpf (n = 10).
778
779 (C) Confocal lateral images of a single plane showing filopodial processes from the PHBC
780 contacting neuroblasts of the Tg and aLL ganglia. Arrowheads indicate filopodia.
781
782 (D-G) Confocal lateral images showing the number of Islet2[+] nuclei in the Tg (c) and Xg ganglia
783 (e) through Islet 1/2 antibody staining and quantified in E and G respectively (n = 10), at 72
784 hpf.
785
786 Scale bars 50 μm for A, B, D-G, and 20 μm for C. Error bars, mean ± SEM. Unpaired two-tailed
787 Student's t-test. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$. ns, non-significant. Tg, Trigeminal
788 ganglion; aLL, anterior Lateral Line, OV, Otic Vesicle; Xg, Vagus ganglion; sib, siblings.

791 **Figure S5   Blood flow arrest by Nifepidine at 60-72 hpf does not affect neuronal**
792 **differentiation, while BDM at 54-72 hpf does.**

793
(A,B)   Confocal images showing immunostained Islet2$^+$ nuclei in the SAG (dotted line,
visualized by *TgBAC(neurod:egfp)* ^{nl1}) at 60 hpf in control (untreated) and 48-60 hpf Nifedipine
treated sib and clo$^{-/-}$ embryos; and quantified in I. 60 hpf sib (n = 7), 60 hpf clo$^{-/-}$ (n = 10), 48-
60 hpf sib (n = 7), 48-60 hpf clo$^{-/-}$ (n = 9).

(C)   Confocal lateral images showing immunostained Islet2$^+$ nuclei in the SAG (dotted line,
visualized by *TgBAC(neurod:egfp)* ^{nl1}) in sib and clo$^{-/-}$ embryos at 72 hpf in control (untreated)
and 54-72 hpf BDM treated embryos.

(D)   Graph showing the number of Islet 1/2$^+$ cells in sib (n = 10), clo-/- (n = 9), 54-72 hpf sib
BDM (n = 10), and 54-72 hpf clo$^{-/-}$ BDM (n = 10).

Scale bar 50 µm. Error bars mean ± SEM. Unpaired two-tailed Student's t-test. * $p < 0.05$, **
$p < 0.01$, *** $p < 0.001$. ns, non-significant. pm, posterior macula; sib, siblings.

Movie S1   Filopodial processes of the endothelial cells and neuroblasts contact each other
at 30 hpf.

Z-projection time-lapse video of *TgBAC(neurod:egfp)* ^{nl1}; *(Kdrl:ras-mCherry)* ^{s896} embryos
showing filopodia extending from the PHBC to the SAG and from the SAG to the PHBC at 30
hpf. Note thickenings in some neuroblasts filopodial. Lateral view. Scale bar 10 µm.

Movie S2   Filopodial processes of the endothelial cells and neuroblasts contact each other
at 30 hpf.

820
821 Z-projection time-lapse video of *TgBAC(neurod:egfp) [nl1]; (Kdrl:ras-mCherry) [s896]* embryos
822 showing filopodia extending from the PHBC to the SAG and from the SAG to the PHBC at 30
823 hpf. Note dynamism of filopodial processes. Lateral view. Scale bar 10 μm.
824
825 **Movie S3 Filopodial processes of the endothelial cells and neuroblasts contact each other**
826 **at 36 hpf.**
827
828 Z-projection time-lapse video of *TgBAC(neurod:egfp) [nl1]; (Kdrl:ras-mCherry) [s896]* embryos
829 showing filopodia extending from the PHBC to the SAG and from the SAG to the PHBC at 36
830 hpf. Note dynamics of filopodial processes. Lateral view. Scale bar 10 μm.
831
832
833 **Movie S4 Filopodial processes of the endothelial cells and neuroblasts contact each other**
834 **at 36 hpf.**
835
836 Z-projection time-lapse video of *TgBAC(neurod:egfp) [nl1]; (Kdrl:ras-mCherry) [s896]* embryos
837 showing filopodia extending from the PHBC to the SAG and from the SAG to the PHBC at 36
838 hpf. Note length of EC filopodia. Lateral view. Scale bar 10 μm.
839

References

Adamo, L., Naveiras, O., Wenzel, P.L., McKinney-Freeman, S., Mack, P.J., Gracia-Sancho, J.,

Suchy-Dicey, A., Yoshimoto, M., Lensch, M.W., Yoder, M.C., et al. (2009). Biomechanical

forces promote embryonic haematopoiesis. Nature *459*, 1131–1135.

Alsina, B., Giraldez, F., and Varela-Nieto, I. (2003). Growth Factors and Early Development of

Otic Neurons: Interactions between Intrinsic and Extrinsic Signals. Curr. Top. Dev. Biol. *57*,

177–206.

Anderson, M.J., Pham, V.N., Vogel, A.M., Weinstein, B.M., and Roman, B.L. (2008). Loss of

unc45a precipitates arteriovenous shunting in the aortic arches. Dev. Biol. *318*, 258–267.

Androutsellis-Theotokis, A., Rueger, M.A., Park, D.M., Mkhikian, H., Korb, E., Poser, S.W.,

Walbridge, S., Munasinghe, J., Koretsky, A.P., Lonser, R.R., et al. (2009). Targeting neural

precursors in the adult brain rescues injured dopamine neurons. Proc. Natl. Acad. Sci. *106*,

13570–13575.

Barber, M., Andrews, W.D., Memi, F., Gardener, P., Ciantar, D., Tata, M., Ruhrberg, C., and

Parnavelas, J.G. (2018). Vascular-derived VEGFA promotes cortical interneuron migration

and proximity to the vasculature in the developing forebrain. Cereb. Cortex *28*, 2577–2593.

Begbie, J., Ballivet, M., and Graham, A. (2002). Early Steps in the Production of Sensory

Neurons by the Neurogenic Placodes. Mol. Cell. Neurosci. *21*, 502–511.

Bilak, M.M., Hossain, W.A., and Morest, D.K. (2003). Intracellular fibroblast growth factor

produces effects different from those of extracellular application on development of avian

cochleovestibular ganglion cells in vitro. J. Neurosci. Res. *71*, 629–647.

Bonapace, G., Concolino, D., Formicola, S., and Strisciuglio, P. (2003). A novel mutation in a

patient with insulin-like growth factor 1 (IGF1) deficiency. J Med Genet. *40*, 913–917.

Bussmann, J., Wolfe, S.A., and Siekmann, A.F. (2011). Arterial-venous network formation

during brain vascularization involves hemodynamic regulation of chemokine signaling.

Development *138*, 1717–1726.

Calvo, C.F., Fontaine, R.H., Soueid, J., Tammela, T., Makinen, T., Alfaro-Cervello, C., Bonnaud,

F., Miguez, A., Benhaim, L., Xu, Y., et al. (2011). Vascular endothelial growth factor receptor

3 directly regulates murine neurogenesis. Genes Dev. *25*, 831–844.

Camarero, G., Leon, Y., Gorospe, I., De Pablo, F., Alsina, B., Giraldez, F., and Varela-Nieto, I.

(2003). Insulin-like growth factor 1 is required for survival of transit-amplifying neuroblasts

and differentiation of otic neurons. Dev. Biol. *262*, 242–253.

Chi, N.C., Shaw, R.M., Val, S. De, Kang, G., Jan, L.Y., Black, B.L., and Stainier, D.Y.R. (2008).

Foxn4 directly regulates tbx2b expression and atrioventricular canal formation. Genes Dev.

734–739.

Chiu, J.-J., and Chien, S. (2011). Effects of Disturbed Flow on Vascular Endothelium:

Pathophysiological Basis and Clinical Perspectives. Physiol. Rev. *91*, 327–387.

Cleaver, O., and Dor, Y. (2012). Vascular instruction of pancreas development. *2843*, 2833–

2843.

Cliff, T.S., and Dalton, S. (2017). Metabolic switching and cell fate decisions : implications for

pluripotency , reprogramming and development. Curr. Opin. Genet. Dev. *46*, 44–49.

Delgado, A.C., Ferrón, S.R., Vicente, D., Porlan, E., Perez-Villalba, A., Trujillo, C.M., D'Ocón,

P., and Fariñas, I. (2014). Endothelial NT-3 Delivered by Vasculature and CSF Promotes

Quiescence of Subependymal Neural Stem Cells through Nitric Oxide Induction. Neuron *83*,

572–585.

Du, Y., Zhang, S., Yu, T., Du, G., Zhang, H., and Yin, Z. (2016). Wnt3a is critical for endothelial

progenitor cell-mediated neural stem cell proliferation and differentiation. Mol. Med. Rep.

14, 2473–2482.

Epstein, A., Gleadle, J., McNeill, L., Hewitson, K., O'Rourke, J., Mole, D., Mukherji, M.,

Metzen, E., Wilson, M., Dhanda, A., et al. (2001). C . elegans EGL-9 and Mammalian

Homologs Define a Family of Dioxygenases that Regulate HIF by Prolyl Hydroxylation The

Henry Wellcome Building of Genomic Medicine. Cell *107*, 43–54.

Fantetti, K.N., and Fekete, D.M. (2012). Members of the BMP, Shh, and FGF morphogen

families promote chicken statoacoustic ganglion neurite outgrowth and neuron survival in

vitro. Dev. Neurobiol. *72*, 1213–1228.

Fantin, A., Lampropoulou, A., Gestri, G., Raimondi, C., Senatore, V., Zachary, I., and

Ruhrberg, C. (2015). NRP1 Regulates CDC42 Activation to Promote Filopodia Formation in

Endothelial Tip Cells. Cell Rep. *11*, 1577–1590.

Fuwa, T.J., Kinoshita, T., Nishida, H., and Nishihara, S. (2015). Reduction of T antigen causes

loss of hematopoietic progenitors in Drosophila through the inhibition of filopodial

extensions from the hematopoietic niche. Dev. Biol. *401*, 206–219.

Gerhardt, H., Golding, M., Fruttiger, M., Ruhrberg, C., Lundkvist, A., Abramsson, A., Jeltsch,

M., Mitchell, C., Alitalo, K., Shima, D., et al. (2003). VEGF guides angiogenic sprouting

utilizing endothelial tip cell filopodia. J. Cell Biol. *161*, 1163–1177.

Golling, G., Amsterdam, A., Sun, Z., Antonelli, M., Maldonado, E., Chen, W., Burgess, S.,

Haldi, M., Artzt, K., Farrington, S., et al. (2002). Insertional mutagenesis in zebrafish rapidly

identifies genes essential for early vertebrate development. Nat. Genet. *31*, 135–140.

González-Méndez, L., Seijo-Barandiarán, I., and Guerrero, I. (2017). Cytoneme-mediated cell-

cell contacts for Hedgehog reception. Elife *6*, 1–24.

Hamada, H., Watanabe, M., Lau, H.E., Nishida, T., Hasegawa, T., Parichy, D.M., and Kondo, S.

(2014). Involvement of Delta/Notch signaling in zebrafish adult pigment stripe patterning.

Development *141*, 318–324.

Hawkins, K., Joy, S., Delhove, J., Kotiadis, V., Fernandez, E., Fitzpatrick, L., Whiteford, J., King,

P., Bolanos, J., Duchen, M., et al. (2016). NRF2 Orchestrates the Metabolic Shift during

Induced Pluripotent Stem Cell Reprogramming Report NRF2 Orchestrates the Metabolic Shift

during Induced Pluripotent Stem Cell Reprogramming. Cell Rep. *14*, 1883–1891.

Himmels, P., Paredes, I., Adler, H., Karakatsani, A., Luck, R., Marti, H.H., Ermakova, O.,

Rempel, E., Stoeckli, E.T., and Ruiz De Almodóvar, C. (2017). Motor neurons control blood

vessel patterning in the developing spinal cord. Nat. Commun. *8*, 1–16.

Hogan, K.A., Ambler, C.A., Chapman, D.L., and Bautch, V.L. (2004). The neural tube patterns

vessels developmentally using the VEGF signaling pathway. Development *131*, 1503-13.

Homem, C.C.F., Steinmann, V., Burkard, T.R., Jais, A., and Esterbauer, H. (2014). Ecdysone

and Mediator Change Energy Metabolism to Terminate Proliferation in Drosophila Neural

Stem Cells. Cell *158*, 874–888.

Howe, D., Bradford, Y., Conlin, T., Eagle, A., Fashena, D., Frazer, K., Knight, J., Mani, P.,

Martin, R., Moxon, S., et al. (2013). ZFIN, the Zebrafish Model Organism Database: increased

support for mutants and transgenics.

Huang, H., and Kornberg, T.B. (2015). Myoblast cytonemes mediate Wg signaling from the

wing imaginal disc and Delta-Notch signaling to the air sac primordium. Elife *4*, 1–22.

Inaba, M., Buszczak, M., and Yamashita, Y.M. (2015). Nanotubes mediate niche-stem-cell signalling in the Drosophila testis. Nature *523*, 329–332.

Ito, K., and Suda, T. (2014). Metabolic requirements for the maintenance of self-renewing stem cells. Nat. Rev. Mol. Cell Biol. *15*, 243–256.

Jensen, R.W., Chuman, H., Trobe, J.D., and Deveikis, J.P. (2004). Facial and Trigeminal Neuropathies in Cavernous Sinus Fistulas. J. Neuro-Ophthalmology *24*, 34–38.

Jin, K., Zhu, Y., Sun, Y., Mao, X.O., Xie, L., and Greenberg, D.A. (2002a). Vascular endothelial growth factor (VEGF) stimulates neurogenesis in vitro and in vivo. Proc. Natl. Acad. Sci. *99*, 11946–11950.

Jin, K., Mao, X.O., Sun, Y., Xie, L., and Greenberg, D.A. (2002b). Stem cell factor stimulates neurogenesis in vitro and in vivo. J. Clin. Invest. *110*, 311–319.

Jin, S.-W., Beis, D., Mitchell, T., Chen, J.-N., and Stainier, D.Y.R. (2005). Cellular and molecular analyses of vascular tube and lumen formation in zebrafish. Development *132*, 5199–5209.

Kacem, K., Lacombe, P., Seylaz, J., and Bonvento, G. (1998). Structural Organization of the Perivascular Astrocyte Endfeet and Their Relationship With the Endothelial Glucose Transporter : A Confocal Microscopy Study. Glia *23*, 1–10.

Kida, Y., Kawamura, T., Wei, Z., Sogo, T., Jacinto, S., Shigeno, A., Kushige, H., Yoshihara, E., Liddle, C., Ecker, J., et al. (2015). ERRs Mediate a Metabolic Switch Required for Somatic Cell Reprogramming to Pluripotency. Stem Cell *16*, 547–555.

Kimmel, C.B., Ballard, W.W., Kimmel, S.R., Ullmann, B., and Schilling, T.F. (1995). Stages of Embryonic Development of the Zebrafish. Dev. Dyn. *203*, 253–310.

Kornberg, T.B. (2017). Distributing signaling proteins in space and time: the province of cytonemes. Curr. Opin. Genet. Dev. *45*, 22–27.

Lago, M.R.R., Fernandes, L. da C., Lyra, I.M., Ramos, R.T., Teixeira, R., Salles, C., and Ladeia, A.M.T. (2018). Sensorineural hearing loss in children with sickle cell anemia and its association with endothelial dysfunction. Hematology *23*, 849–855.

Lange, C., Turrero Garcia, M., Decimo, I., Bifari, F., Eelen, G., Quaegebeur, A., Boon, R., Zhao, H., Boeckx, B., Chang, J., et al. (2016). Relief of hypoxia by angiogenesis promotes neural stem cell differentiation by targeting glycolysis. EMBO J. *35*, 924–941.

Leslie, J.D., Ariza-McNaughton, L., Bermange, A.L., McAdow, R., Johnson, S.L., and Lewis, J. (2007). Endothelial signalling by the Notch ligand Delta-like 4 restricts angiogenesis. Development *134*, 839–844.

Li, W.M., Webb, S.E., Chan, C.M., and Miller, A.L. (2008). Multiple roles of the furrow

deepening Ca2+ transient during cytokinesis in zebrafish embryos. Dev. Biol. *316*, 228–248.

Liao, W., Bisgrove, B., Sawyer, H., and Hug, B. (1997). The zebrafish gene cloche acts

upstream of a flk-1 homologue to regulate endothelial cell differentiation. Development

124, 381–389.

Llorens-Bobadilla, E., Zhao, S., Baser, A., Saiz-Castro, G., Zwadlo, K., and Martin-Villalba, A.

(2015). Single-Cell Transcriptomics Reveals a Population of Dormant Neural Stem Cells that

Become Activated upon Brain Injury. Cell Stem Cell *17*, 329–340.

Louissaint, A., Rao, S., Leventhal, C., and Goldman, S. (2002). Coordinated interaction of

neurogenesis and angiogenesis in the adult songbird brain. Neuron *34*, 945–960.

Ma, Q., Chen, Z., Barrantes, I. del B., Luis de la Pompa, J., and Anderson, D.J. (1998).

Neurogenin1 Is Essential for the Determination of Neuronal Precursors for Proximal Cranial

Sensory Ganglia. Neuron *20*, 469–482.

Maier, E.C., Saxena, A., Alsina, B., Bronner, M.E., and Whitfield, T.T. (2014). Sensational

placodes: Neurogenesis in the otic and olfactory systems. Dev. Biol. *389*, 50–67.

Makita, T., Sucov, H.M., Gariepy, C.E., Yanagisawa, M., and Ginty, D.D. (2008). Endothelins

are vascular-derived axonal guidance cues for developing sympathetic neurons. Nature *452*,

759–763.

Mandal, L., Martinez-Agosto, J.A., Evans, C.J., Hartenstein, V., and Banerjee, U. (2007). A

Hedgehog- and Antennapedia-dependent niche maintains Drosophila haematopoietic

precursors. Nature *446*, 320–324.

Mirzadeh, Z., Merkle, F.T., Soriano-Navarro, M., Garcia-Verdugo, J.M., and Alvarez-Buylla, A.

(2008). Neural Stem Cells Confer Unique Pinwheel Architecture to the Ventricular Surface in

Neurogenic Regions of the Adult Brain. Cell Stem Cell *3*, 265–278.

Mohrin, M., Shin, J., Liu, Y., Brown, K., Luo, H., Xi, Y., Haynes, C.M., and Chen, D. (2015). A

mitochondrial UPR-mediated metabolic checkpoint regulates hematopoietic stem cell aging.

Science (80-.). *347*, 1374-7.

Morton, W.M., Ayscough, K.R., and McLaughlin, P.J. (2000). Latrunculin alters the actin-

monomer subunit interface to prevent polymerization. Nat. Cell Biol. *2*, 376–378.

Mukouyama, Y., Shin, D., Britsch, S., Taniguchi, M., and Anderson, D.J. (2002). Sensory

Nerves Determine the Pattern of Arterial Differentiation and Blood Vessel Branching in the

Skin. Cell *109*, 693–705.

Murdoch, B., and Roskams, A.J. (2012). Fibroblast Growth Factor Signaling Regulates

Neurogenesis at Multiple Stages in the Embryonic Olfactory Epithelium. Stem Cells Dev. *22*,

525–537.

Nakajima, H., Yamamoto, K., Agarwala, S., Terai, K., Fukui, H., Fukuhara, S., Ando, K.,

Miyazaki, T., Yokota, Y., Schmelzer, E., et al. (2017). Flow-Dependent Endothelial YAP

Regulation Contributes to Vessel Maintenance. Dev. Cell *40*, 523-536.e6.

Nicoli, S., Standley, C., Walker, P., Hurlstone, A., Fogarty, K.E., and Lawson, N.D. (2010).

MicroRNA-mediated integration of haemodynamics and Vegf signalling during angiogenesis.

Nature *464*, 1196–1200.

Nikolaou, N., Watanabe-Asaka, T., Gerety, S., Distel, M., Koster, R.W., and Wilkinson, D.G.

(2009). Lunatic fringe promotes the lateral inhibition of neurogenesis. Development *136*,

2523–2533.

Le Noble, F., Klein, C., Tintu, A., Pries, A., and Buschmann, I. (2008). Neural guidance

molecules, tip cells, and mechanical factors in vascular development. Cardiovasc. Res. *78*,

232–241.

Obernier, K., Cebrian-Silla, A., Thomson, M., Parraguez, J.I., Anderson, R., Guinto, C., Rodas

Rodriguez, J., Garcia-Verdugo, J.M., and Alvarez-Buylla, A. (2018). Adult Neurogenesis Is

Sustained by Symmetric Self-Renewal and Differentiation. Cell Stem Cell *22*, 221-234.e8.

Obholzer, N., Wolfson, S., Trapani, J.G., Mo, W., Nechiporuk, A., Busch-Nentwich, E., Seiler,

C., Sidi, S., Sollner, C., Duncan, R.N., et al. (2008). Vesicular Glutamate Transporter 3 Is

Required for Synaptic Transmission in Zebrafish Hair Cells. J. Neurosci. *28*, 2110–2118.

Ochsenbein, A.M., Karaman, S., Proulx, S.T., Berchtold, M., Jurisic, G., Stoeckli, E.T., and

Detmar, M. (2016). Endothelial cell-derived semaphorin 3A inhibits filopodia formation by

blood vascular tip cells. Development *143*, 589–594.

Ottone, C., Krusche, B., Whitby, A., Clements, M., Quadrato, G., Pitulescu, M.E., Adams, R.H.,

and Parrinello, S. (2014). Direct cell-cell contact with the vascular niche maintains quiescent

neural stem cells. Nat. Cell Biol. *16*, 1045–1056.

Palmer, T.D., Willhoite, A.R., and Gage, F.H. (2000). Vascular Niche for Adult Hippocampal

Neurogenesis. J. Comp. Neurolgy *494*, 479–494.

Paul, A., Mazighi, M., Lenck, S., Bresson, D., Herman, P., and Hautefort, C. (2016). Isolated

intermittent bilateral hearing loss revealing a brain hemorrhage. J. Neurol. Sci. *370*, 18–20.

Phng, L.-K., Stanchi, F., and Gerhardt, H. (2013). Filopodia are dispensable for endothelial tip

cell guidance. Development *140*, 4031–4040.

Pittman, A.J., Law, M.-Y., and Chien, C.-B. (2008). Pathfinding in a large vertebrate axon

tract: isotypic interactions guide retinotectal axons at multiple choice points. Development

135, 2865–2871.

Rafii, S., Butler, J.M., and Ding, B. Sen (2016). Angiocrine functions of organ-specific

endothelial cells. Nature *529*, 316–325.

Ramasamy, S.K., Kusumbe, A.P., and Adams, R.H. (2015). Regulation of tissue morphogenesis by endothelial cell-derived signals. Trends Cell Biol. *25*, 148–157.

Ramírez-Castillejo, C., Sánchez-Sánchez, F., Andreu-Agulló, C., Ferrón, S.R., Aroca-Aguilar, J.D., Sánchez, P., Mira, H., Escribano, J., and Fariñas, I. (2006). Pigment epithelium-derived factor is a niche signal for neural stem cell renewal. Nat. Neurosci. *9*, 331–339.

Ramírez-Weber, F.-A., and Kornberg, T.B. (1999). Cytonemes: Cellular Processes that Project to the Principal Signaling Center in Drosophila Imaginal Discs. Cell *97*, 599–607.

Reischauer, S., Stone, O.A., Villasenor, A., Chi, N., Jin, S.W., Martin, M., Lee, M.T., Fukuda, N., Marass, M., Witty, A., et al. (2016). Cloche is a bHLH-PAS transcription factor that drives haemato-vascular specification. Nature *535*, 294–298.

Riquelme, R., Cediel, R., Contreras, J., Rodríguez-de La Rosa, L., Murillo-Cuesta, S., Hernandez, C., Zubeldia, J., Cerdan, S., and Varela-Nieto, I. (2010). A comparative study of age-related hearing loss in wild type and insulin-like growth factor I deficient mice. Front. Neuroanat. *4*, 27.

Rojas-Ríos, P., Guerrero, I., and González-Reyes, A. (2012). Cytoneme-mediated delivery of Hedgehog regulates the expression of bone morphogenetic proteins to maintain germline stem cells in Drosophila. PLoS Biol. *10*, e1001298.

Roy, S., Huang, H., Liu, S., and Kornberg, T.B. (2014). Cytoneme-mediated contact-dependent transport of the Drosophila decapentaplegic signaling protein. Science (80-.). *343*, 1244624.

Ryall, J.G., Cliff, T., Dalton, S., and Sartorelli, V. (2015). Metabolic Reprogramming of Stem Cell Epigenetics. Cell Stem Cell *17*, 651–662.

Sanders, T.A., Llagostera, E., and Barna, M. (2013). Specialized filopodia direct long-range transport of SHH during vertebrate tissue patterning. Nature *497*, 628–632.

Schlosser, G. (2010). Making senses development of vertebrate cranial placodes. Int Rev Cell Mol Biol *283*, 129–234.

Shen, Q., Wang, Y., Kokovay, E., Lin, G., Chuang, S.M., Goderie, S.K., Roysam, B., and Temple, S. (2008). Adult SVZ Stem Cells Lie in a Vascular Niche: A Quantitative Analysis of Niche Cell-Cell Interactions. Cell Stem Cell *3*, 289–300.

Shi, S., and Gronthos, S. (2003). Perivascular Niche of Postnatal Mesenchymal Stem Cells in Human Bone Marrow and Dental Pulp. J. Bone Miner. Res. *18*, 696–704.

Shi, X., Alves, T., Zeng, X.-L., Kibbey, R., Estes, M., and Guan, X. (2015). GLP-2 reprograms glucose metabolism in intestinal stem cells. FASEB J *29*, 851–854.

Siekmann, A.F., and Lawson, N.D. (2007). Notch signalling limits angiogenic cell behaviour in

developing zebrafish arteries. Nature *445*, 781–784.

Siekmann, A.F., Standley, C., Fogarty, K.E., Wolfe, S.A., and Lawson, N.D. (2009). Chemokine

signaling guides regional patterning of the first embryonic artery. Genes Dev. *23*, 2272–2277.

Spradling, A., Drummond-Barbosa, D., and Kai, T. (2001). Stem cells find their niche. Nature

414, 98–104.

Stainier, D.Y., Weinstein, B.M., Detrich, H.W., Zon, L.I., and Fishman, M.C. (1995). Cloche, an

early acting zebrafish gene, is required by both the endothelial and hematopoietic lineages.

Development *121*, 3141–3150.

Stainier, D.Y., Fouquet, B., Chen, J.N., Warren, K.S., Weinstein, B.M., Meiler, S.E., Mohideen,

M.A., Neuhauss, S.C., Solnica-Krezel, L., Schier, A.F., et al. (1996). Mutations affecting the

formation and function of the cardiovascular system in the zebrafish embryo. Development

123, 285–292.

Stanganello, E., and Scholpp, S. (2016). Role of cytonemes in Wnt transport. J. Cell Sci. *129*,

665–672.

Stanganello, E., Hagemann, A.I.H., Mattes, B., Sinner, C., Meyen, D., Weber, S., Schug, A.,

Raz, E., and Scholpp, S. (2015). Filopodia-based Wnt transport during vertebrate tissue

patterning. Nat. Commun. *6*, 1–14.

Suchting, S., Freitas, C., le Noble, F., Benedito, R., Breant, C., Duarte, A., and Eichmann, A.

(2007). The Notch ligand Delta-like 4 negatively regulates endothelial tip cell formation and

vessel branching. Proc. Natl. Acad. Sci. *104*, 3225–3230.

Sumanas, S., Jorniak, T., and Lin, S. (2005). Identification of novel vascular endothelial-

specific genes by the microarray analysis of the zebrafish cloche mutants. Blood *106*, 534–

541.

Taberner, L., Bañón, A., and Alsina, B. (2018). Anatomical map of the cranial vasculature and

sensory ganglia. J. Anat. *232*, 431–439.

Tan, X., Liu, W.A., Zhang, X.J., Shi, W., Ren, S.Q., Li, Z., Brown, K.N., and Shi, S.H. (2016).

Vascular Influence on Ventral Telencephalic Progenitors and Neocortical Interneuron

Production. Dev. Cell *36*, 624–638.

Tata, M., Wall, I., Joyce, A., Vieira, J.M., Kessaris, N., and Ruhrberg, C. (2016). Regulation of

embryonic neurogenesis by germinal zone vasculature. Proc. Natl. Acad. Sci. *113*, 13414–

13419.

Tavazoie, M., Van der Veken, L., Silva-Vargas, V., Louissaint, M., Colonna, L., Zaidi, B., Garcia-

Verdugo, J.M., and Doetsch, F. (2008). A Specialized Vascular Niche for Adult Neural Stem

Cells. Cell Stem Cell *3*, 279–288.

Thisse, B., Heyer, V., Lux, A., Alunni, V., Degrave, A., Seiliez, I., Kirchner, J., Parkhill, J.-P., and

Thisse, C. (2004). Spatial and Temporal Expression of the Zebrafish Genome by Large-Scale In

Situ Hybridization Screening. Methods Cell Biol. *77*, 505–519.

Ulrich, F., Ma, L.H., Baker, R.G., and Torres-Vázquez, J. (2011). Neurovascular development

in the embryonic zebrafish hindbrain. Dev. Biol. *357*, 134–151.

Vasudevan, A., Long, J.E., Crandall, J.E., Rubenstein, J.L.R., and Bhide, P.G. (2008).

Compartment-specific transcription factors orchestrate angiogenesis gradients in the

embryonic brain. Nat. Neurosci. *11*, 429–439.

Vemaraju, S., Kantarci, H., Padanad, M.S., and Riley, B.B. (2012). A Spatial and Temporal

Gradient of Fgf Differentially Regulates Distinct Stages of Neural Development in the

Zebrafish Inner Ear. PLoS Genet. *8*, e1003068.

Westerfield, M. (2000). The zebrafish book. A guide for the laboratory use of zebrafish

(Danio rerio). (Univ. of Oregon Press, Eugene.).

Xiao, T., Roeser, T., Staub, W., and Baier, H. (2005). A GFP-based genetic screen reveals

mutations that disrupt the architecture of the zebrafish retinotectal projection.

Development *132*, 2955–2967.

Yoshida, S., Sukeno, M., and Nabeshima, Y.I. (2007). A vasculature-associated niche for

undifferentiated spermatogonia in the mouse testis. Science (80-.). *317*, 1722–1726.