Deviant EEG resting-state large-scale brain network dynamics in euthymic bipolar disorder patients

Alena Damborská[1,2*], Camille Piguet[3], Jean-Michel Aubry[3,4], Alexandre G. Dayer[3,4], Christoph M. Michel [1,5], Cristina Berchio[1,3]

[1]Functional Brain Mapping Laboratory, Campus Biotech, Department of Basic Neurosciences, University of Geneva, Geneva, Switzerland
[2]Department of Psychiatry, Faculty of Medicine, Masaryk University and University Hospital Brno, Brno, Czech Republic
[3]Department of Psychiatry, Geneva University Hospital, Service of Psychiatric Specialties, Mood Disorders, Geneva, Switzerland
[4]Department of Psychiatry, University of Geneva, Geneva, Switzerland
[5]Lemanic Biomedical Imaging Centre (CIBM), Lausanne and Geneva, Switzerland

* Correspondence:
Alena Damborská
adambor@med.muni.cz

Keywords: EEG microstate, Large-scale brain networks, Resting state, Dynamic brain activity, Bipolar disorder, High-density EEG.

Abstract

Background: Neuroimaging studies provided evidence for disrupted resting-state functional brain network activity in bipolar disorder (BD). Electroencephalographic (EEG) studies found altered temporal characteristics of functional EEG microstates during depressive episode within different affective disorders. Here we investigated whether euthymic patients with BD show deviant resting-state large-scale brain network dynamics as reflected by altered temporal characteristics of EEG microstates.
Methods: We used high-density EEG to explore between-group differences in duration, coverage and occurrence of EEG microstates in 17 euthymic adults with BD and 17 age- and gender-matched healthy controls.
Results: Microstate analysis revealed five microstates (A-E) in global clustering across all subjects. In patients compared to controls, we found increased occurrence and coverage of microstate A that did not significantly correlate with anxiety scores.
Conclusion: Our results provide neurophysiological evidence for altered large-scale brain network dynamics in BD patients and suggest the increased presence of A microstate to be an electrophysiological trait characteristic of BD.

1   Introduction

Bipolar disorder (BD) is a common and severe psychiatric disorder, with an important personal and societal burden (Cloutier et al., 2018; Eaton et al., 2012). The prevalence of BD worldwide is considered to range between 1% and 3% (Merikangas et al., 2007). BD patients are frequently misdiagnosed and often identified at late stages of disease progression, which can lead to inadequate treatment (Hirschfeld, 2007) and worse functional prognosis (Vieta et al., 2018). A better

understanding of the underlying pathophysiology is needed to identify objective biomarkers of BD
that would improve diagnostic and/or treatment stratification of patients.

Potential candidates for neurobiological biomarkers could arise from functional brain network
abnormalities in BD patients. Evidence from brain imaging studies consistently points to
abnormalities in circuits implicated in emotion regulation and reactivity. Particularly, attenuated
frontal and enhanced limbic activations are reported in BD patients (Chen et al., 2011; Houenou et
al., 2011; Kupferschmidt and Zakzanis, 2011). Interestingly, regions implicated in the
pathophysiology of the disease, such as the inferior frontal gyrus, the medial prefrontal cortex
(mPFC), and the amygdala present altered activation patterns even in unaffected first-degree relatives
of BD patients (Piguet et al., 2015), pointing towards brain alterations that could underlie disease
vulnerability. Moreover, evidence from functional magnetic resonance imaging (fMRI) studies
showed aberrant resting-state functional connectivity between frontal and meso-limbic areas in BD
when compared to healthy controls (Vargas et al., 2013). A recently developed functional
neuroanatomic model of BD suggests, more specifically, decreased connectivity between ventral
prefrontal networks and limbic brain regions including the amygdala (Strakowski et al., 2012; Chase
and Philips, 2016). The functional connectivity abnormalities in BD in brain areas associated with
emotion processing were shown to vary with mood state. A resting-state functional connectivity
study of emotion regulation networks demonstrated that subgenual anterior cingulate cortex
(sgACC)-amygdala coupling is critically affected during mood episodes, and that functional
connectivity of sgACC plays a pivotal role in mood normalization through its interactions with the
ventrolateral PFC and posterior cingulate cortex (Rey et al., 2016). Nevertheless, although different
fMRI metrics allowed to report deviant patterns of large-scale networks and altered resting-state
functional connectivity (Rey et al., 2016; Wang et al., 2016) in BD, the precise temporal dynamics of
the functional brain networks at rest remain to be determined.

Large-scale neural networks dynamically and rapidly re-organize themselves to enable efficient
functioning (de Pasquale et al., 2018; Bressler and Menon, 2010). Fast dynamics of the resting-state
large-scale neural networks can be studied on sub-second temporal scales with EEG microstate
analysis (Pascual-Marqui et al., 1995; Van de Ville et al., 2010; Michel and Koenig, 2018). EEG
microstates are defined as short periods (60-120 ms) of quasi-stable electric potential scalp
topography (Lehmann et al., 1987; Koenig et al., 2002). Therefore, microstate analysis can cluster the
scalp's topographies of the resting-state EEG activity into the set of a few microstate classes
including the four canonical classes A-D (Michel and Koenig, 2018) and more recent additional ones
(Custo et al., 2017; Bréchet et al., 2019). Since each microstate class topography reflects a coherent
neuronal activity (Khanna et al., 2015; Michel and Koenig, 2018), the temporal characteristics, such
as duration, occurrence and coverage, may be linked to the expression of spontaneous mental states
and be representative of the contents of consciousness (Changeux and Michel, 2004; Lehmann et al.,
1990). Numerous studies reported abnormalities in temporal properties of resting-state EEG
microstates in neuropsychiatric disorders (for review see Khanna et al., 2015; Michel and Koenig,
2018). Evidence from microstate studies suggests that altered resting-state brain network dynamics
may represent a marker of risk to develop neuropsychiatric disorders (Tomescu et al., 2014, 2015;
Andreou et al., 2014), may predict clinical variables of an illness (Gschwind et al., 2016), or help to
assess the efficacy of a treatment (Atluri et al., 2018; Sverak et al., 2018). Only two studies
investigated resting-state EEG in BD patients (Strik et al., 1995; Damborská et al., 2019). These
studies examined patients during a depressive episode within different affective disorders. Adaptive
segmentation of resting-state EEG showed abnormal microstate topographies and reduced overall
average microstate duration in patients that met criteria for unipolar or bipolar mood disorders or for
dysthymia (Strik et al., 1995). Using a *k*-means cluster analysis, an increased   occurrence of

microstate A with depression as an effect related to the symptom severity was observed during a
period of depression in unipolar and bipolar patients (Damborská et al., 2019).

Trait markers of BD based on neurobiological findings can be considered as biomarkers of illness
(Piguet et al., 2016; Berchio et al., 2017). These trait markers of BD can be studied during the
periods of remission, or euthymia. No microstate study on spontaneous activity, however, has been
performed on euthymic BD patients to the best of our knowledge. Thus, the main goal of the current
study was to explore group differences between euthymic patients with BD and healthy controls in
terms of resting-state EEG microstate dynamics. We hypothesized that BD patients during remission
will show altered temporal characteristics of EEG microstates such as duration, coverage, and
occurrence.

98   2   Materials and Methods

99   2.1   Subjects

Data were collected from 17 euthymic adult patients with BD and 17 healthy control (HC) subjects.
The patients were recruited from the Mood Disorders Unit at the Geneva University Hospital. A
snowball convenience sampling was used for the selection of the BD patients. Control subjects were
recruited by general advertisement. All subjects were clinically evaluated using clinical structured
interview (DIGS: Diagnostic for Genetic Studies, (Nurnberger et al., 1994). BD was confirmed in the
experimental group by the usual assessment of the specialized program, an interview with a
psychiatrist, and a semi-structured interview and relevant questionnaires with a psychologist.
Exclusion criteria for all participants were a history of head injury, current alcohol or drug abuse.
Additionally, a history of psychiatric or neurological illness and of any neurological comorbidity
were exclusion criteria for controls and bipolar patients, respectively. Symptoms of mania and
depression were evaluated using the Young Mania Rating Scale (YMRS) (Young et al., 1978) and
the Montgomery-Åsberg Depression Rating Scale (MADRS) (Williams and Kobak, 2008),
respectively. Participants were considered euthymic if they scored < 6 on YMRS and < 12 on
MADRS at the time of the experiment, and were stable for at least 4 weeks before. All patients were
medicated, receiving pharmacological therapy including antipsychotics, antidepressants and mood
stabilizers, and had to be under stable medication for at least 4 weeks. The experimental group
included both BD I ($n = 10$) and BD II ($n = 7$) types.
To check for possible demographic or clinical differences between groups, subject characteristics
such as age, education or level of depression were compared between groups using independent t-
tests. Anxiety is highly associated with BD (Simon et al., 2004; 2007) and is a potential confounding
variable when investigating microstate dynamics at rest. For example, decreased duration of EEG
microstates at rest in patients with panic disorder has been reported (Wiedemann et al., 1998). To
check for possible differences in anxiety symptoms, all subjects were assessed with the State-trait
Anxiety Inventory (STAI) (Spielberger et al., 1970) and the scores were compared between patients
and controls using independent t-tests.
This study was carried out in accordance with the recommendations of the Ethics Committee for
Human Research of the Geneva University Hospital, with written informed consent from all subjects.
All subjects gave written informed consent in accordance with the Declaration of Helsinki. The
protocol was approved by the Ethics Committee for Human Research of the Geneva University
Hospital, Switzerland.

130   2.2   EEG recording and pre-processing

The EEG was recorded with a high density 256-channel system (EGI System 200; Electrical
Geodesic Inc., OR, USA), sampling rate of 1kHz, and Cz as acquisition reference. Subjects were
sitting in a comfortable upright position and were instructed to stay as calm as possible, to keep their
eyes closed and to relax for 6 minutes. They were asked to stay awake.

To remove muscular artifacts originating in the neck and face the data were reduced to 204 channels.
Two to four minutes of EEG data were selected based on visual assessment of the artifacts and band-
pass filtered between 1 and 40 Hz. Subsequently, in order to remove ballistocardiogram and oculo-
motor artifacts, infomax-based Independent Component Analysis (Jung et al., 2000) was applied on
all but one or two channels rejected due to abundant artifacts. Only components related to
physiological noise, such as ballistocardiogram, saccadic eye movements, and eye blinking, were
removed based on the waveform, topography and time course of the component. The cleaned EEG
recordings were down-sampled to 125 Hz and the previously identified noisy channels were
interpolated using a three-dimensional spherical spline (Perrin et al., 1989), and re-referenced to the
average reference. All the preprocessing steps were done using MATLAB and the freely available
Cartool Software 3.70 (https://sites.google.com/site/cartoolcommunity/home), programmed by Denis
Brunet.

2.3 EEG data analysis

To estimate the optimal set of topographies explaining the EEG signal, a standard microstate analysis
was performed using *k*-means clustering (see Supplementary Fig. 1). The polarity of the maps was
ignored in this procedure (Brunet et al., 2011; Murray et al, 2008; Pascual-Marqui et al., 1995). To
determine the optimal number of clusters, we applied a meta-criterion that is a combination of seven
independent optimization criteria (for details see Bréchet et al., 2019). In order to improve the signal-
to-noise ratio, only the data at the time points of the local maximum of the Global Field Power (GFP)
were clustered (Pascual-Marqui et al., 1995; Koenig et al., 2002; Britz et al, 2010, Tomescu et al.,
2014). The GFP is a scalar measure of the strength of the scalp potential field and is calculated as the
standard deviation of all electrodes at a given time point (Michel et al., 1993; Brunet et al., 2011;
Murray et al., 2008). The cluster analysis was first computed at the individual level yielding one set
of representative maps for each subject. Then each participant's representative maps were clustered
at global level across all participants (patients and controls) yielding one set of representative maps
for all participants.

In order to retrieve the temporal characteristics of the microstates, spatial correlation was calculated
between every map identified at the global level and the individual subject's topographical map in
every instant of the pre-processed EEG recording. Each continuous time point of the subject's EEG
(not only the GFP peaks) was then assigned to the microstate class of the highest correlation, again
ignoring polarity (Brunet et al., 2011; Bréchet et al., 2019; Michel and Koenig, 2018; Santarnecchi et
al., 2017). Temporal smoothing parameters (window half size = 3, strength (Besag Factor) = 10)
ensured that the noise during low GFP did not artificially interrupt the temporal segments of stable
topography (Brunet et al., 2011; Pascual-Marqui et al., 1995). For each subject, three temporal
parameters were then calculated for each of the previously identified microstates: (i) occurrence, (ii)
coverage, and (iii) duration. Occurrence indicates how many times a microstate class recurs in one
second. The coverage in percent represents the summed amount of time spent in a given microstate
class as a portion of the whole analyzed period. The duration in milliseconds for a given microstate
class indicates the amount of time that a given microstate class is continuously present. In order to
assess the extent to which the representative microstate topographies explain the original EEG data,
the global explained variance (GEV) was calculated as the sum of the explained variances of each
microstate weighted by the GFP. Microstate analysis was performed using the freely available
Cartool Software 3.70, (https://sites.google.com/site/cartoolcommunity/home), programmed by

Denis Brunet. Mann-Whitney U test was used to investigate group differences for temporal
parameters of each microstate. Multiple comparisons were corrected using the false discovery rate
(FDR) method (Benjamini, 2010).
Spearman's rank correlations were calculated between the MADRS, YMRS, STAI-state, and STAI-
trait scores and significant microstate parameters to check for possible relationships between
symptoms and microstate dynamics. Statistical evaluation was performed by the routines included in
the program package Statistica'13 (1984-2018, TIBCO, Software Inc, Version 13.4.0.14).

185 3 Results

186 3.1 Clinical and demographic variables

187 There were no significant differences in age and level of education between the patient and the
188 control groups. In both groups, very low mean scores on depression and mania symptoms were
189 observed, which did not significantly differ between the two groups. BD patients showed higher
190 scores on state and trait scales of the STAI. For all subject characteristics, see Table 1.

191 3.2 Microstate results

The meta-criterion used to determine the most dominant topographies revealed five resting-state
microstate maps explaining 82.2 % of the global variance (Fig. 1). The topographies resembled those
previously reported as A, B, C, and D maps (Khanna et al., 2015; Michel and Koenig, 2018; Koenig
et al., 2002; Britz et al., 2010) and one of the three recently identified additional maps (Custo et al.,
2017). We labeled these scalp maps from A to E in accordance with the previous literature on
microstates. The scalp topographies showed left posterior-right anterior orientation (map A), a right
posterior-left anterior orientation (map B), an anterior-posterior orientation (map C), a fronto-central
maximum (map D), and a parieto-occipital maximum (map E).
Since some microstate parameters showed a non-homogeneity of variances in the two groups
(Levene's tests for the microstate C coverage and microstates A and C duration; $p<0.01$), we decided
to calculate Mann-Whitney U test to investigate group differences for temporal parameters of each
microstate.
We found significant between-group differences for microstate classes A and B. Both microstates
showed increased presence in patients in terms of occurrence and coverage. The two groups did not
differ in any temporal parameter of microstates C, D, or E. The results of the temporal characteristics
of each microstate are summarized in Table 2 and Figure 2.

208 3.3 Clinical correlations

The results of Spearman's rank correlation revealed a significant positive association between the
coverage of the microstate B and the STAI-state ($r = 0.40$; $p<0.05$) and STAI-trait ($r = 0.54$; $p<0.05$)
scores. The results of Spearman's rank correlation revealed a significant positive association between
the occurrence of the microstate B and the STAI-trait ($r = 0.47$; $p<0.05$) scores. The results of
Spearman's rank correlation revealed no significant associations between the STAI-state or STAI-
trait scores and the occurrence or coverage of the microstate A (all absolute r-values < 0.35).
The results of Spearman's rank correlation revealed no significant associations between the MADRS
and YMRS scores and the occurrence or coverage of the microstate A and B (all absolute r-values <
0.30).

218 4 Discussion

Our study presents the first evidence for altered resting-state EEG microstate dynamics in euthymic
patients with BD. Patients were stable and did not significantly differ in their depressive or manic
symptomatology from healthy controls at the time of experiment. Despite this fact, they showed
abnormally increased presence of microstates A and B, the latter correlating with the anxiety level.
In an earlier combined fMRI-EEG study the microstate A was associated with the auditory network
(Britz et al., 2010). Moreover, generators of the functional EEG microstates were estimated in recent
studies, where sources of the microstate A showed left-lateralized activity in the temporal lobe,
insula, mPFC, and occipital gyri (Custo et al., 2017; Bréchet et al., 2019).
In the fMRI literature as well, resting-state functional connectivity alterations of the insula (Yin et al.,
2018), the auditory network (Reinke et al., 2013), and the mPFC (Gong et al., 2019) were reported in
BD patients. Verbal episodic memory deficits and language-related symptoms in BD patients were
suggested to be associated with a diminished functional connectivity within the auditory/temporal
gyrus and to be compensated by increased fronto-temporal functional connectivity (Reinke et al.,
2013). The mPFC was also identified as a major locus of shared abnormality in BD and
schizophrenia (Öngür et al., 2010), showing reduced default mode network connectivity from the
mPFC to the hippocampus and fusiform gyrus, as well as increased connectivity between the mPFC
and primary visual cortex in BD. Hypoconnectivity of the default mode network from the left
posterior cingulate cortex to the bilateral mPFC and bilateral precuneus, and reduced salience
connectivity of the left sgACC to the right inferior temporal gyrus in BD patients (Gong et al., 2019)
was observed in unmedicated BD patients. In euthymic BD subjects compared to healthy controls,
resting-state functional connectivity of the insula (Minuzzi et al., 2018) and amygdala (Li et al.,
2018) to other brain regions was reported to be increased and decreased, respectively. In summary,
the evidence from fMRI studies shows both hypoconnectivity (Gong et al., 2019; Öngür et al., 2010)
and hyperconnectivity (Minuzzi et al., 2018; Reinke et al., 2013; Öngür et al., 2010) pointing to
complex alterations of functional resting-state networks. Our findings of increased presence of the
microstate A in euthymic BD patients might be related to the hyperconnectivity of the underlying
networks that involve the temporal lobe, insula, mPFC, and occipital gyri.
Anxiety symptoms were previously associated with greater severity and impairment in BD (Simon et
al., 2004) and euthymic bipolar patients tend to present high residual level of anxiety (Albert et al.,
2008), as it was the case here. No significant correlation was found between the increased anxiety
scores and the increased occurrence or coverage of the microstate A. Our results therefore indicate
that this alteration of microstate dynamics might represent a characteristic feature of BD that is not
affected by anxiety. The demonstrated alterations in microstate A dynamics during clinical remission
might reflect (i) an impaired resting-state large-scale brain network dynamics as a trait characteristic
of the disorder and/or (ii) a compensatory mechanism needed for clinical stabilization of the disorder.
Our study is the first to examine EEG microstate dynamics of spontaneous activity in BD patients
during remission. The here demonstrated group difference of BD patients vs. controls, is not
congruent with the previously reported reduced duration of the EEG microstates during a depressive
episode (Strik et al., 1995). The experimental group in that study was not restricted to bipolar
patients, however, and included also patients who met the criteria for unipolar depression or
dysthymia. Moreover, authors examined the overall microstate duration and did not examine distinct
microstates separately. These and other aspects, such as different clustering methods used, make it
difficult to compare our findings with that early evidence of disrupted microstate dynamics in
depression. Nevertheless, in our recent resting-state study we showed positive associations of
depressive symptoms with the occurrence of microstate A in a heterogenous group of patients with
affective disorders (Damborská et al., 2019).

The microstate B was previously associated with the visual network (Britz et al., 2010; Custo et al.,
2014, 2017; Bréchet et al., 2019). In our group of BD patients, we found an abnormally increased

occurrence and coverage of microstate B that was associated with higher anxiety. Previous studies
also suggest that anxiety may influence visual processing (Phelps et al., 2006; Laretzaki et al., 2010)
and that connections between amygdala and visual cortex might underlie enhanced visual processing
of emotionally salient stimuli in patients with social fobia (Goldin et al., 2009). Our finding of
increased presence of microstate B positively associated with anxiety level in euthymic BD patients
is consistent with these observations. To the best of our knowledge, there are no other studies that
would aim to examine in detail the relationship between anxiety and resting-state EEG microstate
dynamics. An early microstate study reported decreased overall resting-state microstate duration in
panic disorder (Wiedemann et al., 1998). This study, however, did not assess temporal characteristics
of different microstates separately and it is therefore difficult to compare those findings with our
observations. Further evidence is needed to determine, whether the increased presence of microstate
B in our experimental group is a characteristic feature of BD or anxiety, or whether it is related to
both conditions.
Changes in microstate A and B have been reported in several psychiatric conditions such as
schizophrenia, dementia, narcolepsy, multiple sclerosis, panic disorder, etc. (for review see Michel
and Koenig, 2018). In schizophrenia patients, both increased (Andreou et al., 2014) and decreased
(Nishida et al., 2013; Lehmann et al., 2005) durations of microstate B as well as increased occurrence
of microstate A (Lehmann et al., 2005) were reported. Increases in duration and occurrence of
microstates A and B were also observed in patients with multiple sclerosis, moreover predicting
depression scores and other clincal variables (Gschwind, et al., 2016). It was suggested that multiple
sclerosis affects the "sensory" (visual, auditory) rather than the higher-order (salience, central
executive) functional networks (Michel and Koenig, 2018). Our findings of impaired dynamics in
microstates A and B suggest a similar interpretation for the BD. Evidence from fMRI studies points
to topographical dysbalances between the default mode and sensorimotor networks in BD patients
with opposing patterns in depression and mania (Martino et al., 2016). Cyclothymic and depressive
temperaments were associated with opposite changes in the sensorimotor network variability in the
resting state signal measured by fractional standard deviation of Blood-Oxygen-Level Dependent
signal (Conio et al., 2019). Our findings of altered microstates A and B dynamics is consistent with
this fMRI evidence of impaired sensorimotor network in affective disorders, and moreover suggests
that neural correlates of these deficits are prominent even during the euthymic state in BD patients.

In summary, results of the current study seem to indicate that dysfunctional activity of resting-state
brain networks underlying microstates A and B is a detectable impairment in BD during an euthymic
state. The presence of microstate A and B represent measures that might be implicated in clinical
practice if our findings are replicated in larger cohorts. Importantly, these parameters, whose changes
were observed during remission, could be potentially useful for early identification of BD that could
help better management of the disease. If future studies confirm the same pattern in prodromal or
vulnerable subjects, it could also help detection of at-risk subjects and therefore the possiblility for
early intervention. The present study has, however, some limitations. Our low sample size made it
impossible to examine any potential influence of medication on the microstate parameters by
comparing patients receiving a specific drug with those not receiving it. Possible effects of
medication on our results should be therefore taken into account. Due to the same reason, it was not
possible to examine any potential influence of subtypes of BD on microstate results.

5   Conclusions

Our study described altered EEG resting-state microstate temporal parameters in euthymic bipolar
patients. These findings provide an insight into the resting-state global brain network dynamics in
BD. The increased presence of microstate A might be considered as a candidate electrophysiological

non-specific trait marker of BD. Nevertheless, studies examining possible interactions between
microstate dynamics and BD symptoms are needed to better understand the dysfunction of large-
scale brain network resting-state dynamics in this affective disorder.

6   Table legends

Table 1. [a] Edinburgh inventory (Oldfield, 1971); [b] Education levels: 1 = no high school, 2 = high
school, 3 = university studies

7   Tables

Table 1. Subject characteristics

Characteristic	Healthy controls (n = 17)	Bipolar patients (n = 17)	*t*-value	p-value
Age: mean ± SD	36.6 ± 14.5	35.9 ± 11.9	-0.17	0.87
Gender: male, *n*	12	12		
Handedness[a]: right, *n*	14	14		
Education[b]: mean ± SD	2.3 ± 0.6	2.4 ± 0.5	0.63	0.53
MADRS: mean ± SD	1.4± 1.6	2.3 ±2.9	1.09	0.29
YMRS: mean ± SD	0.86 ±1.4	0.76 ± 1.4	-0.18	0.86
STAI-state: mean ± SD	26.7±4.8	36.9±15.2	-2.13	**0.04**
STAI-trait: mean ± SD	27.4±5.2	42.9±13.3	-3.7	**0.001**

321
322
323 Table 2. Mann-Whitney U test for group comparisons in the investigated microstate parameters

Microstate	A	B	C	D	E
Occurrence (s⁻¹)					
Patients (mean±s.d.)	4.5±1.1	3.9±1.1	5.2±1.1	3.9±2.2	1.9±1.2
Controls (mean±s.d.)	3.4±1.2	3.0±1.1	5.0±1.1	3.4±2.1	2.0±1.3
Z-value	2.51	2.51	-0.03	0.65	-0.14
uncorrected *p*-value	0.01	0.01	0.97	0.51	0.89
FDR corrected *p*-value	**0.03**	**0.03**	0.97	0.85	0.97
Coverage (%)					
Patients (mean±s.d.)	18.9±6.5	16.0±7.8	25.1±8.6	18.5±12.3	6.4±5.1
Controls (mean±s.d.)	13.3±4.9	11.4±6.0	34.7±16.8	17.0±12.0	7.0±5.3
Z-value	2.62	2.79	-1.69	0.17	-0.21
uncorrected *p*-value	0.009	0.005	0.09	0.86	0.84
FDR corrected *p*-value	**0.02**	**0.02**	0.15	0.86	0.86

Duration (ms)					
Patients (mean±s.d.)	26±4	25±4	29±4	26±6	21±4
Controls (mean±s.d.)	24±1	24±3	35±9	27±6	22±4
Z-value	1.38	1.03	-2.27	0.17	-0.69
uncorrected *p*-value	0.17	0.30	0.02	0.86	0.49
FDR corrected *p*-value	0.42	0.50	0.12	0.86	0.61

8 Figures

Supplementary Figure 1. Microstate analysis: **(A)** resting-state EEG from subsample of 16 out of
204 electrodes; **(B)** global field power (GFP) curve with the GFP peaks (vertical lines) in the same
EEG period as shown in (A); **(C)** potential maps at successive GFP peaks, indicated in (B), from the
first 1 s period of the recording; **(D)** set of five cluster maps best explaining the data as revealed by
K-means clustering of the maps at the GFP peaks; **(E)** the original EEG recording shown in (A) with
superimposed color-coded microstate segments. Note that each time point of the EEG recording was
labelled with the cluster map, shown in (D), with which the instant map correlated best. The duration
of segments, occurrence, and coverage for all microstates were computed on thus labeled EEG
recording.
337

338
339   Figure 1. The five microstate topographies identified in the global clustering across all subjects.
340
341
342

343
Figure 2. Temporal dynamics of EEG microstates in patients with bipolar disorder (BD) and in
healthy controls (HC). In each subplot, the raw data is plotted on top of a boxplot showing the mean
(□), 95% confidence interval (box plot area), 1 standard deviation (whiskers), and significant
differences (*). In all plots, x-axes represent the subject group; y-axes represent the occurrence
(upper plots) or coverage (lower plots). Note significantly increased occurrence and coverage of the
microstate A and B in the BD compared to HC group (FDR corrected p < 0.05).

9   Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial
relationships that could be construed as a potential conflict of interest.

10   Author Contributions

AD – designed the study, performed the analysis, and wrote the initial draft; JMA, AGD and CP –
were responsible for clinical assessment; CMM – served as an advisor; CB – collected the HD-EEG
data and was responsible for the overall oversight of the study. All authors revised the manuscript.

11   Funding

This study was supported by the European Union Horizon 2020 research and innovation program
under the Marie Skłodowska-Curie grant agreement No. 739939, by the Swiss National Science
Foundation (grant No. 320030_184677), and by the National Centre of Competence in Research
(NCCR) "SYNAPSY–The Synaptic Basis of Mental Diseases" (NCCR Synapsy Grant # "51NF40 –
185897). The funding sources had no role in the design, collection, analysis, or interpretation of the
study.

12   Acknowledgments

Special thanks go to Anne-Lise Kung, psychologist, for her involvement in clinical data collection.

13   References

Albert, U., Rosso, G., Maina, G., Bogetto, F., (2008). Impact of anxiety disorder comorbidity on
quality of life in euthymic bipolar disorder patients: differences between bipolar I and II subtypes. J.
Affective Disord. 105, 297-303.

Andreou, C., Faber, P.L., Leicht, G., Schoettle, D., Polomac, N., Hanganu-Opatz, I.L., et al. (2014).
Resting-state connectivity in the prodromal phase of schizophrenia: Insights from EEG microstates.
Schizophr. Res. 152, 513-520.

Atluri, S., Wong, W., Moreno, S., Blumberger, D.M., Daskalakis, Z.J., Farzan, F., (2018). Selective
modulation of brain network dynamics by seizure therapy in treatment-resistant depression.
NeuroImage Clin. 20, 1176-1190.

Benjamini Y., (2010). Discovering the false discovery rate. J. R. Stat. Soc. Ser. B Stat. Methodol.
72, 405-416.

Berchio C., Piguet C., Michel C.M., Cordera P., Rihs T.A., Dayer A.G., et al. (2017). Dysfunctional
gaze processing in bipolar disorder. NeuroImage Clin. 16, 545-556.

Bréchet, L., Brunet, D., Birot, G., Gruetter, R., Michel C.M., Jorge, J., (2019). Capturing the
spatiotemporal dynamics of self-generated, task-initiated thoughts with EEG and fMRI. Neuroimage,
194, 82-92.

Bressler, S.L., Menon, V., (2010) Large-scale brain networks in cognition: emerging methods and
principles. Trends Cogn. Sci. 14, 277–290.

Britz, J., Van De Ville D., Michel, C.M., (2010). BOLD correlates of EEG topography reveal rapid
resting-state network dynamics. NeuroImage, 52, 1162-1170.

Brunet, D., Murray, M,M., Michel, C.M., (2011). Spatiotemporal analysis of multichannel EEG:
CARTOOL. Comput. Intell. Neurosci. 813870

Changeux, J.P. and Michel, C.M., (2004). Mechanism of neural Integration at the Brain-scale Level.
In: Grillner, S., Graybiel, A.M. (Eds.), Microcircuits. MIT Press, Cambridge, pp. 347-370.

Chase, H.W., Phillips, M.L. (2016) Elucidating Neural Network Functional Connectivity
Abnormalities in Bipolar Disorder: Toward a Harmonized Methodological Approach. Biol.
Psychiatry Cogn. Neurosci. Neuroimaging 1, 288-298.

Chen, C.H., Suckling, J., Lennox, B.R., Ooi, C., Bullmore, E.T., (2011). A quantitative meta-analysis
of fMRI studies in bipolar disorder. Bipolar Disord. 13, 1-15.

Cloutier, M., Greene, M., Guerin, A., Touya, M., Wu, E., (2018). The economic burden of bipolar I
disorder in the United States in 2015. J. Affective Disord., 226, 45-51.

Custo, A., Vulliemoz, S., Grouiller, F., Van De Ville, D., Michel, C., (2014). EEG source imaging of
brain states using spatiotemporal regression. Neuroimage 96, 106-116

Custo, A., Van De Ville, D., Wells, W.M., Tomescu, M.I., Brunet, D., Michel, C.M., (2017).
Electroencephalographic Resting-State Networks: Source Localization of Microstates. Brain
Connectivity 7, 671-682.

Damborská, A., Tomescu, M.I., Honzírková, E., Barteček, R., Hořínková, J., Fedorová, S. et al.
(2019) EEG resting-state large-scale brain network dynamics are related to depressive symptoms.
bioRxiv doi: https://doi.org/10.1101/619031

Eaton, W.W., Alexandre, P., Bienvenu, O.J., Clarke, D., Martins, S.S., Nestadt, G., et al. (2012). The
Burden of Mental Disorders, in: Eaton W.W. (Ed.), Public Mental Health, Oxford University Press

Goldin, P.R., Manber, T., Hakimi, S., Canli, T., Gross, J.J. (2009). Neural bases of social anxiety
disorder: Emotional reactivity and cognitive regulation during social and physical threat. Arch. Gen.
Psychiatry 66, 170-180.

Gong J., Chen G., Jia Y., Zhong S., Zhao L., Luo X., et al. (2019). Disrupted functional connectivity
within the default mode network and salience network in unmedicated bipolar II disorder. Prog.
Neuro-Psychopharmacol. Biol. Psychiatry 88, 11-18.

Gschwind, M., Hardmeier, M., Van De Ville, D., Tomescu, M.I., Penner, I., Naegelin, Y., et al.
(2016). Fluctuations of spontaneous EEG topographies predict disease state in relapsing-remitting
multiple sclerosis. NeuroImage Clin. 12, 466-477.

Hirschfeld, R.M.A., (2007). Screening for bipolar disorder. Am. J. Managed Care 13, S164-S169.

Houenou, J., Frommberger, J., Carde, S., Glasbrenner, M., Diener, C., Leboyer, M., et al. (2011).
Neuroimaging-based markers of bipolar disorder: Evidence from two meta-analyses. J. Affective
Disord. 132, 344-355.
441
442 Jung, T., Makeig, S., Westerfield, M., Townsend, J., Courchesne, E.c Sejnowski, T. J., (2000).
443 Removal of eye activity artifacts from visual event-related potentials in normal and clinical subjects.
444 Clin. Neurophysiol. 111, 1745-1758.
445
446 Khanna, A., Pascual-Leone, A., Michel, C.M., Farzan F., (2015). Microstates in resting-state EEG:
447 Current status and future directions. Neurosci. Biobehav. Rev. 49, 105-113.
448
449 Koenig, T., Prichep, L., Lehmann, D., Sosa, P.V., Braeker, E., Kleinlogel, H., et al. (2002).
450 Millisecond by millisecond, year by year: Normative EEG microstates and developmental stages.
451 Neuroimage 16, 41-48.
452
453 Kupferschmidt, D.A., Zakzanis, K.K., (2011). Toward a functional neuroanatomical signature of
454 bipolar disorder: Quantitative evidence from the neuroimaging literature. Psychiatry Res.
455 Neuroimaging 193, 71-79.
456
457 Laretzaki, G., Plainis, S., Argyropoulos, S., Pallikaris, I.G., Bitsios, P. (2010). Threat and anxiety
458 affect visual contrast perception. J Psychopharmacol 24, 667-675.
459
460 Lehmann, D., Ozaki, H., Pal, I., (1987). EEG alpha map series: brain micro-states by space-oriented
461 adaptive segmentation. Electroencephalogr. Clin. Neurophysiol. 67, 271-288.
462
463 Lehmann D., (1990). Brain Electric Microstates and Cognition: The Atoms of Thought. In: John
464 E.R., Harmony T., Prichep L.S., Valdés-Sosa M., Valdés-Sosa P.A. (eds) Machinery of the Mind.
465 Birkhäuser, Boston, MA, pp. 209-224.
466
467 Lehmann, D., Faber, P.L., Galderisi, S., Herrmann, W.M., Kinoshita, T., Koukkou, M., et al. (2005).
468 EEG microstate duration and syntax in acute, medication-naïve, first-episode schizophrenia: A multi-
469 center study. Psychiatry Res. Neuroimaging, 138, 141-156.
470
471 Li, G., Liu, P., Andari, E., Zhang, A., Zhang, K., (2018). The role of amygdala in patients with
472 euthymic bipolar disorder during resting state. Front. Psychiatry 9(SEP).
473
474 Martino, M., Magioncalda, P., Huang, Z., Conio, B., Piaggio, N., Duncan, N.W., et al. (2016).
475 Contrasting variability patterns in the default mode and sensorimotor networks balance in bipolar
476 depression and mania. Proc. Natl. Acad. Sci. U S A 113, 4824-4829.
477
478 Merikangas, K.R., Akiskal, H.S., Angst, J., Greenberg, P.E., Hirschfeld, R.M.A., Petukhova, M., et
479 al. (2007). Lifetime and 12-month prevalence of bipolar spectrum disorder in the national
480 comorbidity survey replication. Arch. Gen. Psychiatry 64, 543-552.
481
482 Michel, C.M., Brandeis, D., Skrandies, W., Pascual, R., Strik, W.K., Dierks, T. et al. (1993). Global
483 field power: a 'time-honoured' index for EEG/EP map analysis. Int. J. Psychophysiol. 15, 1-2.
484

485 Michel, C.M. and Koenig, T., (2018). EEG microstates as a tool for studying the temporal dynamics
486 of whole-brain neuronal networks: A review. Neuroimage 180, 577-593.
487
488 Minuzzi, L., Syan, S.K., Smith, M., Hall, A., Hall, G.B.C., Frey, B.N., (2018). Structural and
489 functional changes in the somatosensory cortex in euthymic females with bipolar disorder. Aust.
490 New Zealand J. Psychiatry 52, 1075-1083.
491
492 Murray, M.M., Brunet, D., Michel, C.M., (2008). Topographic ERP analyses: A step-by-step tutorial
493 review. Brain Topogr. 20, 249-264.
494
495 Nishida, K., Morishima, Y., Yoshimura, M., Isotani, T., Irisawa, S., Jann, K., et al. (2013). EEG
496 microstates associated with salience and frontoparietal networks in frontotemporal dementia,
497 schizophrenia and Alzheimer's disease. Clin. Neurophysiol. 124, 1106-1114.
498
499 Nurnberger Jr., J.I., Blehar, M.C., Kaufmann, C.A., York-Cooler, C., Simpson, S.G., Harkavy-
500 Friedman, J., et al. (1994). Diagnostic interview for genetic studies: Rationale, unique features, and
501 training. Arch. Gen. Psychiatry 51, 849-859.
502
503 Oldfield, R.C., (1971). The assessment and analysis of handedness: The Edinburgh inventory.
504 Neuropsychologia 9, 97-113.
505
506 Öngür, D., Lundy, M., Greenhouse, I., Shinn, A.K., Menon, V., Cohen, B.M., et al. (2010). Default
507 mode network abnormalities in bipolar disorder and schizophrenia. Psychiatry Res Neuroimaging
508 183, 59-68.
509
510 Pascual-Marqui, R.D., Michel, C.M., Lehmann, D., (1995). Segmentation of Brain Electrical Activity
511 into Microstates; Model Estimation and Validation. IEEE Trans. Biomed. Eng. 42, 658-665.
512
513 de Pasquale. F., Corbetta, M., Betti. V., Della Penna, S., (2018). Cortical cores in network dynamics.
514 Neuroimage 180, 370-382.
515
516 Perrin, F., Pernier, J., Bertrand, O., Echallier, J. F., (1989). Spherical splines for scalp potential and
517 current density mapping. Electroencephalogr. Clin. Neurophysiol. 72, 184-187.
518
519 Phelps, E.A., Ling, S., Carrasco, M. (2006) Emotion facilitates perception and potentiates the
520 perceptual benefits of attention. Psychol. Sci. 17, 292-299.
521
522 Piguet, C., Dayer, A., Aubry, J.M., (2016). Biomarkers and vulnerability to bipolar disorders.
523 Schweiz Arch. Neurol. Psychiatr. 167, 57-67.
524
525 Piguet, C., Fodoulian, L., Aubry, J.M., Vuilleumier, P., Houenou, J., (2015). Bipolar disorder:
526 Functional neuroimaging markers in relatives. Neurosci. Biobehav. Rev. 57, 284-296.
527
528 Reinke, B., Van de Ven, V., Matura, S., Linden, D.E.J., Oertel-Knöchel, V. (2013). Altered intrinsic
529 functional connectivity in language-related brain regions in association with verbal memory
530 performance in euthymic bipolar patients. Brain Sci. 3, 1357-1373.
531

532   Rey, G., Piguet, C., Benders, A., Favre, S., Eickhoff, S.B., Aubry, J.M., et al. (2016). Resting-state
533   functional connectivity of emotion regulation networks in euthymic and non-euthymic bipolar
534   disorder patients. Eur. Psychiatry 34, 56-63.
535
536   Santarnecchi, E., Khanna, A.R., Musaeus, C.S., Benwell, C.S.Y., Davila, P., Farzan, F., et al. (2017).
537   EEG Microstate Correlates of Fluid Intelligence and Response to Cognitive Training. Brain Topogr.
538   30, 502-520.
539
540   Simon, N.M., Otto, M.W., Wisniewski, S.R., Fossey, M., Sagduyu, K., Frank, E., et al. (2004).
541   Anxiety disorder comorbidity in bipolar disorder patients: Data from the first 500 participants in the
542   Systematic Treatment Enhancement Program for Bipolar Disorder (STEP-BD). Am. J. Psychiatry
543   161, 2222-2229.
544
545   Simon, N.M., Zalta, A.K., Otto, M.W., Ostacher, M.J., Fischmann, D., Chow, C.W., et al. (2007).
546   The association of comorbid anxiety disorders with suicide attempts and suicidal ideation in
547   outpatients with bipolar disorder. J. Psychiatr. Res. 41, 255-264.

548
549   Spielberger, C.D., Gorsuch, R.L., Lushene, R.E., (1970). Manual for the State-Trait Anxiety
550   Inventory.
551
552   Strakowski, S.M., Adler, C.M., Almeida, J., Altshuler, L.L., Blumberg, H.P., Chang, K.D., et al.
553   (2012) The functional neuroanatomy of bipolar disorder: A consensus model. Bipolar Disord. 14,
554   313-325.

555
556   Strik, W.K., Dierks, T., Becker, T., Lehmann, D., (1995). Larger topographical variance and
557   decreased duration of brain electric microstates in depression. J. Neural Transmission, 99, 213-222.
558
559   Sverak, T., Albrechtova, L., Lamos, M., Rektorova, I., Ustohal, L., (2018). Intensive repetitive
560   transcranial magnetic stimulation changes EEG microstates in schizophrenia: A pilot study.
561   Schizophr. Res. 193, 451-452.

562
563   Tomescu, M.I., Rihs, T.A., Becker, R., Britz, J., Custo, A., Grouiller F., et al. (2014). Deviant
564   dynamics of EEG resting state pattern in 22q11.2 deletion syndrome adolescents: A vulnerability
565   marker of schizophrenia? Schizophr. Res. 157, 175-181.
566
567   Tomescu, M.I., Rihs, T.A., Roinishvili, M, Karahanoglu, F.I., Schneider, M., Menghetti, S., et al.
568   (2015). Schizophrenia patients and 22q11.2 deletion syndrome adolescents at risk express the same
569   deviant patterns of resting state EEG microstates: A candidate endophenotype of schizophrenia.
570   Schizophr. Res. Cogn. 2, 159-165.
571
572   Van De Ville, D., Britz, J., Michel, C.M., (2010). EEG microstate sequences in healthy humans at
573   rest reveal scale-free dynamics. Proc. Natl. Acad. Sci. 107, 18179–18184.
574
575   Vargas, C., López-Jaramillo, C., Vieta, E., (2013). A systematic literature review of resting state
576   network-functional MRI in bipolar disorder. J. Affective Disord. 150, 727-735.
577

578 Vieta, E., Salagre, E., Grande, I., Carvalho, A.F., Fernandes, B.S., Berk, M., et al. (2018). Early
579 intervention in Bipolar disorder. Am. J. Psychiatry 175, 411-426.
580
581 Wang, Y., Zhong, S., Jia, Y., Sun, Y., Wang, B., Liu, T., et al. (2016). Disrupted resting-state
582 functional connectivity in nonmedicated bipolar disorder. Radiology 280, 529-536.
583
584 Wiedemann, G., Stevens, A., Pauli, P., Dengler, W., (1998). Decreased duration and altered
585 topography of electroencephalographic microstates in patients with panic disorder. Psychiatry Res.
586 Neuroimaging 84, 37-48.
587
588 Williams, J.B.W. and Kobak K.A., (2008). Development and reliability of a structured interview
589 guide for the Montgomery Asberg depression rating scale (sigma). Br. J. Psychiatry 192, 52-58.
590
591 Yin, Z., Chang, M., Wei, S., Jiang, X., Zhou, Y., Cui, L., et al. (2018). Decreased functional
592 connectivity in insular subregions in depressive episodes of bipolar disorder and major depressive
593 disorder. Front. Neurosci. 12(NOV).
594
595 Young R.C., Biggs, J.T., Ziegler, V.E., Meyer, D.A., (1978). A rating scale for mania: Reliability,
596 validity and sensitivity. Br. J. Psychiatry 133, 429-435.
597

598 **14   Data Availability Statement**

599 The raw data supporting the conclusions of this manuscript will be made available by the authors,
600 without undue reservation, to any qualified researcher.