Evolutionary time best explains the global distribution of living freshwater fish diversity

Elizabeth Christina Miller*[1][2] & Cristian Román-Palacios[1]

[1]Department of Ecology and Evolutionary Biology, University of Arizona, Tucson, Arizona
85721, U.S.A.

[2]School of Aquatic and Fishery Sciences, University of Washington, Seattle, Washington 98105,
U.S.A.

*Correspondence to: emille3@uw.edu

ABSTRACT:

Aim: The evolutionary causes of the latitudinal diversity gradient are debated. Hypotheses have

ultimately invoked either faster rates of diversification in the tropics, or more time for

diversification due to the tropical origins of higher taxa. Here we perform the first test of the

diversification rate and time hypotheses in freshwater ray-finned fishes, a group comprising

nearly a quarter of all living vertebrates.

Location: Global.

Time period: 368–0 mya.

Major taxa studied: Extant freshwater ray-finned fishes.

Methods: Using a mega-phylogeny of actinopterygian fishes and a global database of

occurrence records, we estimated net diversification rates and regional colonization times of co-

occurring species in freshwater drainage basins. We used Generalized Additive Models to test

whether diversification rates and time were related to latitude. We then compared the influence

of diversification rates and time on species richness, and how these factors are related to each

other.

Results: While both diversification rates and the timing of regional colonization were related to

richness, time-for-speciation had 2–6 times more explanatory power than diversification rates.

Time was also more strongly related to latitude than diversification rates. These results held after

controlling for surface area. The most species rich freshwater basins (Amazon, Congo rivers)

were dominated by lineages with Mesozoic origins. The temperate groups dominant today

arrived near the K-Pg boundary, leaving comparatively less time to build richness.

Diversification rates and colonization times were inversely related: recently colonized basins had

the fastest rates, while ancient species-rich faunas had modest rates.

Main conclusions: We concluded that evolutionary time is the lead driver of latitudinal richness

disparities in freshwater fish faunas. We suggest that the most likely path to building very high

species richness is through diversification over long periods of time, rather than diversifying

quickly.

KEYWORDS

diversification rates, freshwater fishes, latitudinal diversity gradient, species richness, time-for-

speciation

1 | INTRODUCTION

Species richness decreases from the equator to the poles. The latitudinal biodiversity gradient has been called the Earth's first-order biodiversity pattern due to its pervasiveness across groups and geologic time (Hillebrand, 2004). There are only three processes that can directly change regional species richness: in-situ speciation, local extinction, and dispersal (Ricklefs, 1987; Roy & Goldberg, 2007). While numerous ecological and evolutionary hypotheses have been proposed to explain the latitudinal biodiversity gradient (Mittelbach et al., 2007), it is clarifying to first determine how these three core processes change with latitude. Other factors such as area or productivity may also influence richness (Tedesco et al., 2012), but these factors must change richness indirectly by acting on speciation, extinction, and/or dispersal.

There has been great interest in comparing diversification rates across phylogenies in recent years, due to the confluence of the construction of large time-calibrated molecular phylogenies (e.g. Jetz, Thomas, Joy, Hartmann, & Mooers, 2012; Rabosky et al., 2018) and the increasing complexity of models of diversification (Rabosky, 2014). A growing number of analyses across groups are revealing that speciation and/or net diversification rates are similar among latitudes (general review in Schluter & Pennell, 2017; Jansson, Rodríguez-Castañeda, & Harding, 2013; birds: Weir & Schluter, 2007; Jetz et al., 2012; Rabosky, Title, & Huang, 2015; mammals: Weir & Schluter, 2007; ants: Economo, Narula, Friedman, Weiser, & Guénard, 2018), or even higher in high latitudes (marine fishes: Rabosky et al., 2018; Miller, Hayashi, Song, & Wiens, 2018; deep-sea invertebrates: O'Hara, Hugall, Woolley, Bribiesca-Contreras, & Bax, 2019; angiosperms: Igea & Tanentzap, 2019). These studies raise the question: if diversification rates do not explain spatial differences in richness then what does?

A potential resolution to this question is to compare the relative importance of

colonization history and diversification rates for explaining species richness (Stephens & Wiens,

2003). This distinction is important because while we may identify in-situ speciation as the

dominant process generating biodiversity in a region (Tedesco et al., 2012), similar richness

disparities may be produced either through faster speciation or speciation over longer periods of

time (Figure 1; Wiens, 2012). For example, tropical marine fishes have modest speciation rates

(Rabosky et al., 2018), but the tropics still have high richness due to the combination of early and

frequent colonization (Miller et al., 2018). Similarly, time matters for explanations involving

asymmetrical dispersal rates. Frequent colonization may not lead to high richness if this trend is

recent, such that new colonists have not had much time to diversify (Miller et al., 2018). The

time-for-speciation effect may also imply spatial variation in extinction pressure. One reason that

a region may be dominated by young lineages is that older lineages went extinct (Miller &

Wiens, 2017). The influence of time on species richness is an idea with deep historical roots

(Wallace, 1878; Willis, 1922; Wiens & Donoghue, 2004; Fine & Ree, 2006; Jablonski, Roy, &

Valentine, 2006; Mittelbach et al., 2007). While researcher attention on time as an explanation

for the latitudinal diversity gradient has waxed and waned over the years (Stephens & Wiens,

2003), recent studies that have simultaneously inferred colonization and diversification history

have found a strong role for time in explaining richness patterns (Jansson et al., 2013; Miller et

al., 2018; Economo et al., 2018; O'Hara et al., 2019; Li & Wiens, 2019; Machac, 2020).

A related hypothesis, that regions with more species have greater ecological limits,

ultimately implies that speciation, extinction or dispersal rates change in association with a

carrying capacity of species (Rabosky, 2009; Hurlbert & Stegen, 2014). Thus, a carrying

capacity influences richness indirectly by acting on speciation, extinction or dispersal, just as

area or climate influences richness. Ecological limits are not inconsistent with either the rate or

time hypotheses. For example, regions with low carrying capacities can take longer to be

colonized in simulations, showing that ecological limits can modulate richness through the time-

for-speciation effect (Pontarp & Wiens, 2017). If a region was colonized early in the history of a

clade, it is generally expected to contain more species than recently colonized regions due to

greater time allowed for speciation, even if speciation rates have slowed through time (Pontarp &

Wiens, 2017). Species richness will increase over time while speciation rates are non-zero and

exceed extinction rates (Machac, 2020).

Freshwater fishes represent nearly a quarter of all vertebrate species (Cavin, 2017) and

are major components of ecosystems in both tropical and temperate latitudes (Berra, 2001). A

body of work has demonstrated correlations with freshwater fish species richness and factors

including productivity and area (Oberdorff, Guégan, & Hugueny, 1995; Smith, Badgley, Eiting,

& Larson, 2010), in-situ speciation (Tedesco et al., 2012) and recent historical events such as

Quaternary sea-level changes (Oberdorff, Guégan, & Hugueny, 1997; Tedesco, Oberdorff,

Lasso, Zapata, & Hugueny, 2005; Leprieur et al., 2011; Dias et al., 2014). At least three

questions remain about what drives species richness in freshwater habitats. First, while in-situ

speciation (cladogenesis) is clearly important as inferred from endemism in species-rich basins

(Tedesco et al. 2012), is high species richness caused by faster rates of speciation or more time

for speciation due to earlier colonization (Figure 1)? Second, since area cannot directly change

species richness, in what potential ways is the species-area relationship related to latitudinal

trends in speciation and colonization? Third, recent events seem to impact local richness within a

continent, but differences in richness among continents remain (Oberdorff et al., 1997; Tedesco

et al., 2005; Dias et al., 2014). How did these continental differences in richness form? We

attempt to answer these questions herein.

Here we test whether variation in diversification rates or time-for-diversification best

explains global diversity patterns in freshwater fishes, especially the latitudinal diversity

gradient. Freshwater fishes have features that are conducive to testing both diversification rates

and time hypotheses. First, living freshwater fishes represent a wide range of ages, with some

radiations diversifying during the Mesozoic or earlier (Briggs, 2005; Capobianco & Friedman,

2018) and others only during the most recent glaciation cycles (Seehausen & Wagner, 2014).

Second, freshwater fishes have low dispersal ability, and therefore their systematics likely retain

signatures of regional events (Lavoué, 2016; Cavin, 2017; Capobianco & Friedman, 2018). Our

study capitalizes on the aggregation of natural history observations and genetic data over many

years (Tedesco et al., 2017; Rabosky et al., 2018), allowing us to make comparisons at a broad

spatial and temporal scale under a common phylogenetic framework.

2 | METHODS

Additional details are given in the Supporting Information. A list of data sources is found in

Appendix 1.

2.1 | Occurrence and phylogenetic data

Expert-vetted occurrence records of freshwater actinopterygian species were assembled by

Tedesco et al. (2017). Records were available for 3,119 drainage basins covering 80% of the

Earth's land surface. Sampling of drainage basins ranged from 70% of the surface of the Indo-

Malay region to 90% of the surface of the Afrotropics (Tedesco et al., 2017). Species coverage

for major freshwater fish groups ranged from 61% of Anabantiformes to 93% of Characiformes.

We removed non-native and uncertain records. Altogether, occurrence records from 14,947

species of freshwater fishes were used to estimate species richness of basins. This dataset also

reported surface area (Figure S1), median latitude and longitude of each basin.

For biogeographic and diversification-rate analyses, we used the maximum likelihood

time-calibrated molecular phylogeny of actinopterygians constructed by Rabosky et al. (2018;

see also http://fishtreeoflife.org). This phylogeny includes 11,638 species with genetic data

(36.9% of known ray-finned fishes).

2.2 | Obtaining diversification rates and colonization times for basins

To estimate diversification rates for each drainage basin, we used three types of branch-

associated measures. We used tip-based net diversification rates calculated using BAMM v.2.5.0

(Rabosky, 2014): three independent runs under a constant-rate model of diversification, and three

runs under a time-varying model. We also used tip-based estimates of the DR statistic (Jetz et al.,

2012). Unlike BAMM, the DR statistic was calculated from phylogenies with all unsampled

species grafted using taxonomic constraints (n=31,516 species). We note that DR tip rates better

approximate speciation rates than net diversification rates in comparison to rates from BAMM

(Title & Rabosky, 2019). For each measure of tip-based rates, we took the mean of rates among

co-occurring species in each basin.

The "time-for-speciation" effect represents the time allowed for in-situ diversification

since a lineage colonized a region (Figure 1; Stephens & Wiens, 2003). Note that our preferred

terminology is to use "dispersal" to refer to the general and bi-directional process of movement

among regions, "colonization" as the addition of new lineages to a focal region as a result of

dispersal, and "time-for-speciation" as the time elapsed between colonization and the present

(Stephens & Wiens, 2003; Hua & Bromham, 2020). To measure time-for-speciation, we must

estimate the amount of time a lineage has been present in the location of interest (Figure 1,

Figure S2). One major challenge to estimating colonization times at the local scale is that the

computation time of biogeographic models scales exponentially with the number of possible

ranges (Matzke, 2014). Modelling dispersal among >3,000 drainage basins is unfeasible using

phylogenetic approaches at present. To overcome this challenge, we instead modelled dispersal

among continental regions. We then used the mean and median colonization time associated with

species present in each drainage basin. This approach to measuring spatial variation in time-for-

speciation is analogous to the grid-cell approach often used to detect spatial variation in

diversification rates (Jetz et al., 2012; Rabosky et al., 2018; Machac, 2020). By focusing on

regions, our colonization time estimates should also be more robust to past range shifts since

regions change less over time than individual rivers and lakes (Hoorn et al., 2010).

We first fit a dispersal-extinction-cladogenesis model (DEC; Ree & Smith, 2008) using

the R package 'BioGeoBEARS' v.1.1 (Matzke, 2014; additional details in Extended Methods,

Supporting Information). We used the maximum likelihood phylogeny including species with

genetic data only (Rabosky et al., 2018), because semi-random grafting of unsampled species is

inappropriate for comparative methods that model the evolution of traits associated with the tips

(Rabosky, 2015). We removed 139 tips that were unsuitable for biogeographic reconstructions,

leaving 11,499 species. Our analysis included six major continental regions of the world: the

Neotropics, Afrotropics, Indo-Malay, Australasia, Nearctic, and Palearctic. Boundaries of these

regions followed Tedesco et al. (2017). Species restricted to marine environments were coded as

occurring in a seventh "marine" region. Though not our focus, these species are needed to inform

the timing of colonization of freshwater regions from the marine realm (Betancur-R, Ortí, Stein,

Marceniuk, & Pyron 2012; Betancur-R, Ortí, & Pyron, 2015; Rabosky, 2020). Our model was

time-stratified to apply constraints on dispersal in accordance with changing connectivity of

continents. We applied the following constraints over six time bins spanning the root (~368 mya)

to the present (following Toussaint, Bloom, & Short, 2017): dispersal between adjacent regions

was not constrained (i.e. the probability of dispersal between adjacent regions was scaled by 1);

dispersal between regions separated by a small marine barrier was scaled by 0.75; dispersal

between regions separated by another landmass was scaled by 0.50; and dispersal between

regions separated by a large marine barrier was scaled by 0.25 (more details in Table S1).

After model fitting, in order to count individual colonization events and visualize

uncertainty in these dates, we simulated 100 biogeographic stochastic maps (Dupin et al., 2016).

Each individual simulation is a realized history that is possible given the model and data,

including the time and location on the branches for biogeographic events. Averaging over all of

these simulations will approximate the ancestral state probabilities calculated by the model. We

used these simulations to estimate the time-for-speciation associated with each drainage basin.

To do this, we traced each individual species back in time to the location on the branch

reconstructed as the colonization of the region(s) it inhabits. Note that it is possible for this event

to precede the crown age of recognizable clades (such as orders), especially at this large

phylogenetic scale, since dispersal can happen at any time along a phylogeny (Hua & Bromham,

2020). See Figure S2 for an illustration for how these times were obtained. For each species we

took the mean time of this event across the 100 stochastic maps. The amount of time-for-

speciation associated with each basin was estimated as the mean and median colonization time

among co-occurring species.

2.3 | Comparing predictors of local richness

To sum, we first tested how basin richness, diversification rates, time-for-speciation, and surface area change with latitude and longitude. Second, we tested whether diversification rates and time-for-speciation each separately predict local richness. Third, we compared the relative support for diversification rates and time-for-speciation for predicting richness, with and without area as a covariate. Fourth, we tested how diversification rates and time-for-speciation are related to each other. Fifth, we tested whether variation in area with latitude could explain our results.

We fit Generalized Additive Models (GAMs) to examine the change in species richness, net diversification rates, and time-for-speciation with latitude, longitude, and both (i.e. the interaction between longitude and latitude). We considered longitude as well as latitude because species richness also varies strongly within the tropics (Figure 3A). We fit univariate GAMs between predictor (latitude or longitude) and response variables (species richness, net diversification rates, or time-for-speciation). GAMs were made spatially explicit by including the interaction between longitude and latitude of each basin in the smooth term (i.e. s(long, lat)). This approach for explicitly accounting for geography in GAMs was first proposed by Brumback & Rice (1998), with further details presented in Kammann & Wand (2003), Hefley et al. (2016), and Wood (2017). Univariate GAMs were fitted using the gam function in the 'mgcv' package in R (base; R Core Team, 2008; Wood, 2011). To assess the direction of the relationships we performed Spearman's rank correlation tests between latitude and species richness, net diversification rates, time-for-speciation and surface area. We also fit a GAM to confirm whether surface area was related to species richness.

Next, we fit spatially explicit GAMs to understand the relative importance of

diversification rates and time-for-speciation for explaining local richness. Again, to account for

spatial autocorrelation, we included a smoother term that summarized the interaction between

latitude and longitude in each basin (e.g. s(long, lat) in GAMs). We first analyzed the

relationships between basin richness and each predictor alone. Then, we included both variables

as predictors of richness. We compared the fit of four models using AIC values: a null model that

assumed species richness to be constant ("null model"); a model where species richness

depended on diversification rates ("div model"); a model where species richness depended with

time-for-speciation ("time model"); and a model where species richness depended on both

diversification rates and time-for-speciation (full model). We then estimated the amount of

deviance in richness explained by each predictor (details in the Extended Methods,

Supplementary Information). We fit this set of four models for each combination of

diversification rate estimate (seven alternatives) and estimate of basin-level colonization times

(two alternatives). In addition, we repeated these analyses while also controlling for the effect of

surface area (i.e. species-area scaling). For each combination of diversification rate and

colonization times, we fit the same four models described above with the addition of surface area

as a covariate.

We examined how two predictors of richness (diversification rates, time-for-speciation)

interact with each other. For example, if diversification rates and time-for-speciation are jointly

responsible for producing the most-diverse faunas, then one would expect these basins to harbor

rapidly-diversifying lineages that colonized a long time ago (Figure 1). Alternatively, some

basins may have fast diversification rates while others were colonized early. We fit spatially

explicit GAMs with time-for-speciation as the predictor of diversification rates. We also

performed Spearman's rank correlation tests between diversification rates and time-for-

speciation to quantify the strength and direction of the association between these variables.

Finally, we fit spatially explicit GAMs to assess the possibility that a covariation between

surface area and latitude was responsible for trends in time-for-speciation and diversification

rates. We fit a set of three GAMs for each target variable (time-for-speciation and diversification

rates). First, we fit a full model including the additive effects of area and latitude in explaining

spatial patterns in either time-for-speciation or diversification rates. Next, we fit two additional

models with changes in the predictor being explained by either latitude or area alone. To

compare the relative importance of latitude and surface area for explaining diversification rates

and time-for-speciation, we estimated the change in deviance from the exclusion of each

predictor compared to the full model.

2.4 | Assessing sensitivity of colonization time estimates

Several freshwater clades are known to have extinct marine members (Betancur-R et al.,

2015). Colonization times associated with these groups may be overestimated without

accounting for marine ancestry erased by extinction. To assess this, we also performed ancestral

range reconstructions using an alternative time-calibrated phylogeny containing 1,582 living and

240 extinct ray-finned fishes (Betancur-R et al., 2015). We used the literature to assign

freshwater fossil species to continental regions (Table S7; Appendix 1). Biogeographic analyses

were performed as above (see also Extended Methods, Supporting Information). We compared

the mean and range of colonization times inferred from biogeographic stochastic mapping for

major freshwater fish clades between the two phylogenies. From this comparison, we found that

colonization times associated with early-diverging (non-teleost) clades were overestimated using

the Rabosky et al. (2018) phylogeny. We removed the four living non-teleost orders

(Polypteriformes, Acipenseriformes, Lepisosteiformes and Amiiformes) from our basin-level

dataset to quantify their impact on basin-specific diversification-rate and time-for-speciation

estimates.

3 | RESULTS

Richness of freshwater actinopterygian fishes among basins reflects the latitudinal diversity

gradient found in other major groups (Figure 2A; Spearman's rank correlation between local

richness and latitude: rho=-0.27, P<0.001). However, latitude alone only explains ~9% of the

variance in species richness across basins (GAM, R_2=0.092, P<0.001; Figure 2A; Table S2).

Longitude alone had similar explanatory power (Figure 2B; GAM, R_2=0.10, P<0.001). The

interaction between longitude and latitude explains a higher proportion of the variance in species

richness (R_2=0.26, P<0.001). These results appear to reflect the strikingly high richness in the

Neotropics relative to other tropical regions. For example, the Amazon basin contains about

twice as many known species as the world's second-most rich basin, the Congo (2,968 vs 1,554

species, respectively; Figure 3A). In sharp contrast, the most species-rich basin in the

Australasian tropics is the Ramu River of New Guinea with 179 species. Therefore, while we

found a latitudinal gradient in freshwater fish richness, much variation in species richness was

unrelated to latitude.

Next, we tested for latitudinal trends in diversification rates, time-for-speciation, and

surface area. Latitude was significantly related to diversification rates (rho=-0.11–0.33,

R_2=0.205–0.212, P<0.001; Figures 2A, 3B; Table S2). This trend may have been influenced by

outlier basins with very high rates (e.g. Lake Malawi and Lake Titicaca; Figure 3B). We also

found a latitudinal trend in time-for-speciation, in which ancient colonizations were typical of

low-latitude basins (median colonization time: R_2=0.253, rho=-0.289, P<0.001; mean

colonization time: R_2=0.271, rho=-0.232, P<0.001; Figures 2A, 3C; Table S2). As expected,

species richness was positively related to surface area of drainage basins (R_2=0.242, P<0.001;

Figure S1A). However, surface area was poorly related to latitude (R_2=0.054, P<0.001; Table

S2; Figure 2A; Figure S1B). Therefore, the latitudinal diversity gradient is potentially explained

by geographic trends in time-for-speciation and diversification rates, but is unlikely to be

explained by a relationship between present-day area and latitude.

We examined the individual effects of diversification rates and time-for-speciation for

explaining richness patterns in general (not just in association with latitude). Both diversification

rates and time-for-speciation were significantly related to richness and explained a similar

portion of its variance globally (diversification rates: R_2=0.338–0.364, all P<0.001; time-for-

speciation: R_2=0.317–0.326, all P<0.001; Table S3). Adding surface area as a covariate increased

the variation in richness explained by GAMs (diversification rates: R_2=0.534–0.554, all P<0.001;

time-for-speciation: R_2=0.527–0.534, all P<0.001; Table S4), reflecting the positive relationship

between area and richness (Figure S1).

Our finding that both diversification rates and time-for-speciation influence species

richness does not necessarily mean that these two processes have synergic effects in each basin

(Figure 1). We used GAMs to test for a relationship between basin-level diversification rates and

colonization times. We found that colonization times explain 39–42% of variance in

diversification rates (all P<0.001; Table S5). Importantly, we found a negative relationship

between diversification rates and colonization times, such that diversification rates tend to be

higher in recently-colonized basins (Figure 4; rho=-0.13–-0.10, all P<0.001; Table S5). These

results suggest that basins with fast diversification rates are distinct from those with early

colonizations (Figure 1).

We then compared the relative importance of diversification rates and time-for-speciation

for explaining species richness. We found that time-for-speciation contributed 2.3–6.1 times

more to species richness patterns than diversification rates (based on deviance values for

alternative GAMs; Table S3). Time-for-speciation contributed 1.4–4.6 times more than

diversification rates based on models that included surface area (Table S4).

Finally, we asked whether latitudinal trends in time-for-speciation and diversification

rates could be responding to a covariance between latitude and area. We found that latitude

contributes 15 times more to variation in diversification rates than does surface area (Table S6).

Latitude contributes 119 times more to variation in time-for-speciation than surface area. This

suggests that latitudinal trends in diversification rates and time-for-speciation (Tables S3, S4) are

unlikely to be explained by latitudinal trends in surface area of drainage basins.

To sum, diversification rates, time-for-speciation and surface area were each significantly

related to species richness. However, time-for-speciation contributes more to richness disparities

and was more strongly related to latitude than diversification rates or surface area. These results

are reflected by the spatial patterns in richness and colonization times presented in Figure 3.

While species-rich tropical basins such as the Amazon and Congo had a mean colonization time

during the Mesozoic, 95.4% of basins in the Nearctic and Palearctic had mean colonization times

after the K-Pg boundary (66 mya; Figure 3C; Table A4). Colonization times and diversification

rates were inversely related. This suggests that the most species-rich basins are so because of

more time for diversification, not faster diversification. Diversification rate differences were also

important for explaining richness patterns but are more relevant among recently colonized (and

relatively depauperate) basins. Note that these results were robust to alternative estimates of

diversification rates (BAMM time-constant, BAMM time-varying, and DR) and whether we

used the mean or median colonization time among co-occurring species (Tables S2–S5).

We compared our colonization time estimates to those using an alternative phylogeny

that also included fossils (Betancur-R et al., 2015). Aside from non-teleosts, colonization

patterns for major freshwater fish lineages were generally congruent between the two

phylogenies (Rabosky et al., 2018; Betancur-R et al., 2015). Despite uncertainty in colonization

times in the ancestral reconstructions, the order of events was consistent such that colonization of

tropical regions generally preceded colonization of extratropical regions (Extended Results 1;

Figure S3). For more details see Extended Results 1.

The addition of fossils, especially marine members of families now restricted to

freshwater, suggested that colonization times estimated for early-diverging fishes were

overestimated (Extended Results 1; Figure S3). However, removing non-teleosts had a negligible

effect on diversification rates globally (mean of 0.24% faster; Extended Results 2). Removing

these species resulted in slightly younger mean colonizations globally (mean of 1.80% younger).

Most basins affected (76%) were in the Nearctic and Palearctic. Results based on teleosts alone

would widen the difference between tropical and extratropical time-for-speciation, in line with

our conclusions based on all actinopterygians.

4 | DISCUSSION

The three processes that directly change species richness are in-situ speciation, extinction, and

dispersal. Similar species richness disparities may be formed by differences in either the rate or

timing of these processes (Figure 1). We examined whether the rate of diversification (speciation

minus extinction) or the length of time allowed for diversification (time since colonization) best

explained the global distribution of freshwater fish richness. Overall, our results suggest that

time-for-speciation is the lead driver of species richness disparities, especially latitudinal

patterns. We also show that diversification rates are highest among recently colonized basins,

suggesting that richness differences among this set of basins are driven by diversification rates.

Time-for-speciation and diversification rates are therefore both needed to explain diversity

patterns overall.

372
4.1 | Time best explains latitudinal richness patterns

We found a latitudinal trend in time-for-speciation associated with the K-Pg boundary

(Figures 3, 5). While dominant tropical lineages tend to have Mesozoic origins, the dominant

temperate lineages tend to have Cenozoic origins (Figure 3C; Figure 5). These patterns are

exemplified by two major groups of fishes: Otophysi and Percomorpha (Figure 5; Extended

Results 1). The crown of Otophysi (142 mya in the phylogeny of Rabosky et al., 2018) was most

often reconstructed as widespread in the Neotropics, Indo-Malay and Afrotropics. The Neotropic

Characiformes, Gymnotiformes, and some Siluriformes are descended from this initial

colonization. Several lineages nested within Otophysi colonized the Nearctic and Palearctic

independently (Figure 5). The Nearctic and Palearctic members of Cyprinidae most likely

arrived from Southeast Asia near the K-Pg boundary or shortly after during the Eocene (Figure

5; Figure S3). These patterns are consistent with past studies (Cavender, 1998; Briggs, 2005;

Chen, Lavoué, & Mayden, 2013), and also using the alternative phylogeny of Betancur-R et al.

(2015; Extended Results 1, Figure S3). In percomorphs, the colonization of the tropics also pre-

dated colonization of the extratropics (Figure 5; Figure S3). For example, in many stochastic

maps colonization of the Neotropics pre-dated the crown of Ovalentaria (108 mya), the clade that

includes the Cyprinodontiformes and Cichlidae (Extended Results 1; Figure S3). These two

lineages contain younger temperate groups (Cavender, 1998).

These results are consistent with a body of evidence from fossil and molecular data that

the modern Neotropical biota was assembled over long time scales rather than from recent

diversification (Hoorn et al., 2010; Albert & Reis, 2011; Antonelli et al., 2018; Albert,

Tagliacollo, & Dagosta, 2020). Richness also varies within the tropics, a pattern that can be

attributed (in part) to colonization and time-for-speciation. While the Afrotropics contain ancient

groups such as the Polypteriformes and Osteoglossiformes, communities today are dominated by

the younger Cichlidae (Figure 5; Figure S3). Due to their numerical dominance, cichlids

contribute more to the mean colonization times estimated for African basins than these ancient

groups (Figures 3, 5). While cichlids famously have fast diversification rates (McGee et al.,

2020), they have not diversified over as long a period of time as the major Neotropical groups,

and so the Afrotropics have comparatively lower richness today (Figure 5). The Australian

tropics are unusual in that they were colonized many times, but no single lineage is dominant

(Figure 5). The number of recent colonizations is consistent with greater ecological opportunity

in the relatively depauperate Australian tropics (Betancur-R et al., 2012). The high endemism

within Australian drainage basins suggests that the lack of dominant groups is related to the

inability of fishes to expand across the continent (Unmack, 2001).

4.2 | Why time drives diversity patterns

What does time-for-speciation imply about the differences between low and high latitudes? The

two hypotheses given below are not mutually exclusive and might have a combined effect in

driving diversity patterns. One hypothesis may be more relevant for the origin of the pattern

(barriers to colonization) and the other for its maintenance (extinction).

4.2.1 | Barriers to extratropical colonization

Our results for freshwater fishes align with the tropical conservatism model (Wiens &

Donoghue, 2004) and the out-of-the-tropics model (Jablonski et al., 2006). The overall pattern of

colonization conforms to the expectations of both models; that is, older tropical clades (e.g.

Otophysi) exported nested lineages to extratropical regions (e.g. Cyprinidae). A prediction of the

tropical conservatism model is that movement out of the tropics is limited by niche conservatism,

or the tendency for related taxa to share similar environmental tolerances (Wiens & Donoghue,

2004). Under this model, we would expect tropical-extratropical dispersal to preferentially occur

during warm periods in Earth's history. The out-of-the-tropics model implies instead that niches

of tropical clades are more evolutionarily labile than extratropical groups (Tomašových &

Jablonski, 2017). Although we did not examine climactic niches in this study, some observations

suggest that niche conservatism may indeed underlie colonization patterns in freshwater fishes.

Many tropical clades have few or no living species that reach temperate latitudes today,

including Characiformes, Gymnotiformes, and Cichlidae. In addition, major radiations found in

the Holarctic (such as Cypriniformes and Percidae) arrived during a period when the Earth was

much warmer overall compared to the present day (Figure 5; Mannion, Upchurch, Benson, &

Goswami, 2014). Future work may investigate whether colonization preceded niche evolution in

temperate fishes (Folk et al., 2019).

The Earth was in a greenhouse period for much of the evolution of modern freshwater

fishes, from the mid-Permian to the Neogene (~272 to 23 mya; Mannion et al., 2014). Why did

the Holarctic otophysans and percomorphs arrive so late, if temperature was not a barrier? In

addition to the low dispersal capabilities of freshwater fishes, there may have been other barriers

to colonization. Laurasia and Gondwana were separated by the Atlantic Ocean by the time

Otophysi and Percomorpha originated (142.1 and 122.7 mya respectively; Rabosky et al., 2018).

Laurasian landmasses were also flooded by epicontinental seas to a greater extent than

Gondwanan landmasses during the Cretaceous, and these seas began to recede during the early

Cenozoic (Ronov, 1994). The flooding of Laurasia and its separation from Gondwana may have

delayed its colonization by freshwater fishes, especially those with poor salt tolerance.

4.2.2 | Extinction and stability of the tropics

It has been traditionally thought that Quaternary glaciation cycles were important for explaining

the latitudinal diversity gradient, because lower latitudes were unaffected by glaciation (Bush,

1994; Mittelbach et al., 2007). In freshwater fishes, Quaternary climate cycles have left a

signature on beta diversity (community turnover) at high latitudes, suggesting that high latitudes

were re-colonized by a subset of species after glaciers receded (Leprieur et al., 2011).

Nonetheless, our reconstructions imply that the latitudinal diversity gradient in freshwater fishes

was already established by the Cretaceous (Figures 3C, 5). Again, we must look to older events

to fully explain the latitudinal diversity gradient.

The Osteoglossiformes, Characiformes, Cyprinodontiformes, and Channidae all show

fossil evidence of extinction in high latitudes during the Cenozoic (Lavoué, 2016; Capobianco &

Friedman, 2018). Meseguer & Condamine (2020) pointed out that if the latitudinal diversity

gradient was flatter during warm periods in Earth's past (Mannion et al., 2014; Saupe et al.,

2019), then temperate extinctions and range contractions must have played a role in the sharp

gradient we see today. They suggested that this asymmetric extinction and dispersal model is an

extension of the tropical conservatism hypothesis (Wiens & Donoghue, 2004): when warm

climates became restricted to low latitudes, warm-adapted temperate lineages went extinct.

Therefore, extinction, colonization and time-for-speciation can be related. Recent arrivals to the

temperate zone have had limited time to replace diversity lost from extinction and range

contraction (see also Miller & Wiens, 2017). Tropical groups were less affected by range

contraction, and so were allowed to diversify over longer periods of time.

464
465
4.3 | Relation to other hypotheses

While we focused on diversification rates and time in this manuscript, our results have

implications for other potential influences on species richness.

4.3.1 | Ecological limits

The presence of ecological limits is often tested with the correlation between species richness

and proxy variables such as productivity (e.g. Machac, 2020). Regions with low productivity are

thought to have smaller carrying capacities, and diversification rates should slow more quickly

than regions with higher carrying capacities (Rabosky, 2009). Our results are seemingly in

conflict with this prediction. We found an inverse relationship between time and diversification

rates: species-rich tropical basins had modest rates, while recently colonized but depauperate

basins had the fastest rates (Figure 3, 4). Basins with fast rates tended to be found in arid regions

(Figure 3; Smith et al., 2010). This inverse pattern is also seen in simulations (Hurlbert & Stegen,

2014) and empirically in marine fishes (Rabosky et al., 2018; Miller et al., 2018), marine

invertebrates (O'Hara et al., 2019) and birds (Machac, 2020), and even has precedents in older

literature (Briggs, 1966). These observations suggest that using environmental variables to test

for ecological limits may give misleading results (Buckley et al., 2010). Studies that use a

phylogenetic approach to trace the evolution of biogeographic ranges have great potential to

reveal how ecological limits act on diversification and colonization (Betancur-R et al., 2012;

Moen & Morlon, 2014).

Machac (2020) suggested that the ecological limits, diversification rates and time

hypotheses could be integrated to explain why old tropical faunas have depressed diversification

rates. As species richness increases, diversification rates slow but are not zero, so that richness

continues to increase. Species-poor regions should have fast diversification rates due to

ecological opportunity, but low richness due to the limited time allowed for diversification. Our

results can add to this integrated view. Basins with mean colonization times from ~12–50 mya

had the fastest rates, but basins with mean times from ~50–150 mya did not vary strongly in rates

(Figure 4). Lineages in the second set of basins are likely past the "exponential growth" phase

(Rabosky, 2009). Once diversification rates begin to slow it will take more time to add new

species, such that species richness variation among these mature faunas is strongly related to the

amount of time allowed for speciation.

4.3.2 | Dispersal frequency

Biased dispersal rates can contribute to species richness disparities without invoking differences

in diversification rates or time (Roy & Goldberg, 2007). We did not test for biased dispersal rates

because it is unfeasible to model dispersal among thousands of discrete areas using phylogenetic

approaches (Matzke, 2014). Future studies at a restricted spatial scale may directly assess the

influence of dispersal asymmetry on richness. For example, the Amazon is a biodiversity source

within the Neotropics, which suggests its high richness is not due to dispersal into Amazonia

(Antonelli et al., 2018).

Still, our results have implications for the role of dispersal rates on freshwater richness

patterns, at least at the regional scale. Much freshwater fish diversity is derived from only a few

colonization events. The first arrival of the Neotropics by the Otophysi accounts for at least

~40% of present-day richness in the region (Figure 5). Australasia was colonized most (310

times), but also has the lowest richness of all regions. Biogeographic patterns in freshwater fishes

contrast with those for marine fishes, where colonization frequency is a major driver of richness

(Miller et al., 2018). This suggests that in freshwater, where dispersal capabilities are low, high

species richness is more easily achieved though in-situ speciation (Figure 1) than from high

dispersal rates.

4.3.3 | Area

Area has traditionally been considered alongside time as an important predictor of species

richness (Willis, 1922). We found that surface area of drainage basins was related to species

richness (Figure S1) in congruence with Oberdorff et al. (1995). However, latitude was a poor

predictor of basin area, and time-for-speciation was more closely related to latitude than area.

This suggests that latitudinal trends in time-for-speciation are not a response to an area-latitude

relationship. One possibility is that past area is a better predictor of latitudinal diversity patterns

than present-day area (Fine & Ree, 2006). Continental flooding would have reduced the area

accessible to freshwater fishes in the Holarctic (Ronov, 1994), which may have delayed

colonization. In conjunction with time, this could help explain why some Holarctic basins are

less species rich than expected given their size today (e.g. the Mississippi; Figure 3A).

Speciation takes time to complete, and long periods of time might be needed to build richness

even if the area is large enough to support many species (Li & Wiens, 2019).

4.4 | Caveats and sources of error

4.4.1 | Sampling and divergence times

Fishes are less densely sampled for genetic information than other vertebrates (Jetz et al., 2012),

and knowledge of fish communities varies by region (Tedesco et al., 2017). We avoided

inferring trends in diversification rates through time, which are sensitive to sampling (Blackburn,

Giribet, Soltis, & Stanley, 2019) and model identifiability (Louca & Pennell, 2020). Note

however that the DR estimates used here were inferred from a phylogeny with missing species

imputed with taxonomic constraints (Rabosky et al., 2018). Our results based on DR estimates

were similar to those using BAMM rates inferred from the tree with genetic data only (Tables

S2–S5), suggesting that variation in the magnitude of diversification rates among fishes is not

likely to be driven by sampling biases.

We believe that the temporal patterns of colonization inferred here should be robust.

While the phylogeny used contains ~36% of living actinopterygian species, it includes 90.2% of

families and 100% of orders (Rabosky et al., 2018). This degree of sampling is sufficient to

capture the age of higher taxa in the tree (Sanderson, 1996) which are more relevant to our time-

for-speciation results than the most recent nodes (Extended Results 1). We compared

colonization times estimated from this phylogeny and an alternative with greatly reduced

sampling and different divergence time estimates (Betancur-R et al., 2015). The order of events

was generally common to both trees, with clades originating in the tropics and exporting nested

lineages to the temperate zone (Extended Results 1; Figure S3). Sampling and divergence time

estimates will both improve with more study, but as long as these nested relationships are

preserved then tropical colonizations will still tend to be older than temperate colonizations

(Cavender, 1998; Chen et al., 2013).

4.4.2 | Range estimates

The methods used here estimate ancestral ranges using the relationships and present-day

distribution of living species. Range contractions from the temperate zone are known from the

Cenozoic fossil record of fishes (Cavender, 1998; Lavoué, 2016; Capobianco & Friedman,

2018). Whether or not our results will be biased by the lack of recent fossils in our tree depends

on the phylogenetic relationships between the missing fossils and living taxa. If the missing

temperate fossils represent independent colonizations with no living descendants, then they will

have little bearing on the ancestral reconstructions among living groups. If these fossils are

instead placed near the stem branch of living temperate groups, then the timing of temperate

colonizations could be underestimated from extant data. Even so, there is strong evidence that

modern tropical lineages have been present since the Mesozoic, and temperate lineages tend to

be nested within tropical ones (Figure 5). Therefore, this potential underestimation should not

overturn the general trend for tropical colonizations to precede temperate colonizations. We

predict that the continued integration of fossils and molecular phylogenies will refine

colonization time estimates, but should not overturn our conclusions about the role of time on the

latitudinal diversity gradient (Extended Results 1 and 2, Figure S3). As mentioned, these range

contractions may actually support, not contradict, the importance of time-for-speciation (Miller

& Wiens, 2017; Meseguer & Condamine, 2020).

5 | CONCLUSIONS

We show that the latitudinal diversity gradient in freshwater fishes is primarily driven by earlier colonization of low-latitude regions, extending the timeline of diversification in the tropics compared to higher latitudes. More broadly, our results suggest that the most likely path to building very high species richness is through diversifying over long periods of time rather than diversifying very quickly. A remaining question is whether we observe younger temperate colonizations because the temperate zone has been harder to colonize, because new colonizations are replacing older ones that went extinct, or other scenarios. The time-for-speciation effect may manifest through colonization opportunity, niche conservatism, ecological limits, environmental stability, and/or past extinction (Stephens & Wiens, 2003; Wiens & Donoghue, 2004; Miller & Wiens, 2017; Pontarp & Wiens, 2017). A priority for future research is determining which of these factors are most relevant for generating the latitudinal diversity gradient, how they interact, and why they vary in space and time.

DATA AVAILABILITY STATEMENT

The data utilized in this study are available at https://figshare.com/s/552e7549303de8f2a823. Data sources are listed in the Supplementary Information and Appendix 1.

ACKNOWLEDGEMENTS: We thank John J. Wiens and Andrew Zaffos for discussion and support. This manuscript was improved with comments from Adam Tomašových, Ricardo Betancur-R, and an anonymous reviewer. Kathryn Chenard painted the fishes in Figure 5. Elizabeth Miller was supported by an NSF Graduate Research Fellowship (DGE-1143953) and NSF Postdoctoral Fellowship (DBI-1906574).

CONFLICT OF INTEREST STATEMENT: We have no conflicts of interest to declare.

REFERENCES

1. Albert, J. S., & Reis, R. E. (2011). *Historical biogeography of Neotropical freshwater fishes.* Los Angeles, CA: University of California Press.

2. Albert, J. S., Tagliacollo, V. A., & Dagosta, F. 2020. Diversification of Neotropical freshwater fishes. *Annual Review of Ecology, Evolution, and Systematics,* 51, 27–53.

3. Antonelli, A., Zizka, A., Carvalho, F. A., Scharn, R., Bacon, C. D., Silvestro, D., & Condamine, F. L. (2018). Amazonia is the primary source of Neotropical biodiversity. *Proceedings of the National Academy of Sciences*, 115, 6034–6039.

4. Berra, T. M. (2001). *Freshwater fish distribution*. Chicago, IL: The University of Chicago Press.

5. Betancur-R, R., Ortí, G., Stein, A. M., Marceniuk, A. P., & Pyron, R. A. 2012. Apparent signal of competition limiting diversification after ecological transitions from marine to freshwater habitats. *Ecology Letters,* 15, 822–830.

6. Betancur-R, R., Ortí, G., & Pyron, R. A. (2015). Fossil-based comparative analyses reveal ancient marine ancestry erased by extinction in ray-finned fishes. *Ecology Letters,* 18, 441–450.

7. Blackburn, D. C., Giribet, G., Soltis, D. E., & Stanley, E. L. (2019). Predicting the impact of describing new species on phylogenetic patterns. *Integrative Organismal Biology,* 1, obz028.

8. Briggs, J. C. (1966). Zoogeography and evolution. *Evolution,* 20, 282–289.

9. Briggs, J. C. (2005). The biogeography of otophysan fishes (Ostariophysi: Otophysi): a

new appraisal. *Journal of Biogeography,* 32, 287–294.

10. Brumback, B. A., & Rice, J. A. (1998). Smoothing spline models for the analysis of

nested and crossed samples of curves (with discussion). *Journal of the American*

Statistical Association, 93, 961–994.

11. Buckley, L. B., Davies, T. J., Ackerly, D. D., Kraft, N. J. B., Harrison, S. P., Anacker, B.

L., … Wiens, J. J. (2010). Phylogeny, niche conservatism and the latitudinal diversity

gradient in mammals. *Proceedings of the Royal Society B,* 277, 2131–2138.

12. Bush, M. B. (1994). Amazonian speciation: a necessarily complex model. *Journal of*

Biogeography, 21, 5–17.

13. Capobianco, A., & Friedman, M. (2018). Vicariance and dispersal in southern

hemisphere freshwater fish clades: a palaeontological perspective. *Biological Reviews,*

94, 662–699.

14. Cavender, T. M. (1998). Development of the North American Tertiary freshwater fish

fauna with a look at parallel trends found in the European record. *Italian Journal of*

Zoology, 65, 149–161.

15. Cavin, L. (2017). *Freshwater fishes: 250 million years of evolutionary history*.

Amsterdam: Elsevier.

16. Chen, J., Lavoué, S., & Mayden, R. L. (2013). Evolutionary origin and early

biogeography of Otophysan fishes (Ostariophysi: Teleostei). *Evolution,* 67, 2218–2239.

17. Dias, M. S., Oberdorff, T., Hugueny, B., Leprieur, F., Jézéquel, J. C., Brosse, S., …

Tedesco, P. A. (2014). Global imprint of historical connectivity on freshwater fish

biodiversity. *Ecology Letters,* 17, 1130–1140.

18. Dupin, J., Matzke, N. J., Särkinen, T., Knapp, S., Olmstead, R. G., Bohs, L., & Smith, S.

D. (2017). Bayesian estimation of the global biogeographical history of the

Solanaceae. *Journal of Biogeography*, 44, 887–899.

19. Economo, E. P., Narula, N., Friedman, N. R., Weiser, M. D., & Guénard, B. (2018).

Macroecology and macroevolution of the latitudinal diversity gradient in ants. *Nature*

Communications, 9, 1778.

20. Fine, P. V. A., & Ree, R. H. (2006). Evidence for a time-integrated species-area effect on

the latitudinal gradient in tree diversity. *American Naturalist,* 168, 796–804.

21. Folk, R. A., Stubbs, R. L., Mort, M. E., Cellinese, N., Allen, J. M., Soltis, P. S., ... &

Guralnick, R. P. (2019). Rates of niche and phenotype evolution lag behind

diversification in a temperate radiation. *Proceedings of the National Academy of*

Sciences, *116*, 10874-10882.

22. Hefley, T. J., Broms, K. M., Brost, B. M., Buderman, F. E., Kay, S. L., Scharf, H. R., …

Hooten, M. B. (2016). The basis function approach for modeling autocorrelation in

ecological data. *Ecology*, 98, 632–646.

23. Hillebrand, H. (2004). On the generality of the latitudinal diversity gradient. *American*

Naturalist, 163, 192–211.

24. Hoorn, C., Wesselingh, F. P., Ter Steege, H., Bermudez, M. A., Mora, A., Sevink, J., ...

& Jaramillo, C. (2010). Amazonia through time: Andean uplift, climate change,

landscape evolution, and biodiversity. *Science*, 330, 927–931.

25. Hua, X., & Bromham, L. (2020). Modeling colonization rates over time: Generating null

models and testing model adequacy in phylogenetic analyses of species assemblages.

Evolution, in press (https://onlinelibrary.wiley.com/doi/epdf/10.1111/evo.14086).

26. Hurlbert, A. H., & Stegen, J. C. (2014). On the processes generating latitudinal richness

gradients: identifying diagnostic patterns and predictions. *Frontiers in Genetics,* 5, 420.

27. Igea, J., & Tanentzap, A. J. (2019) Angiosperm speciation cools down in the tropics.

Ecology Letters, 23, 692–700.

28. Jablonski, D., Roy, K., & Valentine, J. W. (2006). Out of the tropics: evolutionary

dynamics of the latitudinal diversity gradient. *Science,* 314, 102–106.

29. Jansson, R., Rodríguez-Castañeda, G., & Harding, L. E. (2013). What can multiple

phylogenies say about the latitudinal diversity gradient? A new look at the tropical

conservatism, out of the tropics, and diversification rate hypotheses. *Evolution,* 67, 1741–

1755.

30. Jetz, W., Thomas, G. H., Joy, J. B., Hartmann, K., & Mooers, A. O. (2012). The global

diversity of birds in space and time. *Nature,* 491, 444–448.

31. Kammann, E. E., & Wand, M. P. (2003). Geoadditive models. *Journal of the Royal*

Statistical Society. Series C Applied Statistics, 52, 1–18.

32. Lavoué, S. (2016). Was Gondwanan breakup the cause of the intercontinental distribution

of Osteoglossiformes? A time-calibrated phylogenetic test combining molecular,

morphological, and paleontological evidence. *Molecular Phylogenetics and Evolution,*

99, 34–43.

33. Leprieur, F., Tedesco, P. A., Hugueny, B., Beauchard, O., Dürr, H. H., Brosse, S., &

Oberdorff, T. (2011). Partitioning global patterns of freshwater fish beta diversity reveals

contrasting signatures of past climate changes. *Ecology Letters*, 14, 325-334.

34. Li, H., & Wiens, J. J. (2019). Time explains regional richness patterns within clades more

often than diversification rates or area. *The American Naturalist,* 193, 514–529.

35. Louca, S., & Pennell, M. W. (2020). Extant timetrees are consistent with a myriad of

diversification histories. *Nature,* 580, 502–505.

36. Machac, A. (2020). The dynamics of bird diversity in the New World. *Systematic*

Biology, in press (https://doi.org/10.1093/sysbio/syaa028).

37. Mannion, P. D., Upchurch, P., Benson, R. B. J., & Goswami, A. (2014). The latitudinal

biodiversity gradient through deep time. *Trends in Ecology and Evolution,* 29, 42–50.

38. Matzke, N. J. (2014). Model selection in historical biogeography reveals that founder-

event speciation is a crucial process in island clades. *Systematic Biology,* 63, 951–970.

39. McGee, M. D., Borstein, S. R., Meier, J. I., Marques, D. A., Mwaiko, S., Taabu, A., …

Seehausen, O. (2020). The ecological and genomic basis of explosive adaptive radiation.

Nature, in press (https://doi.org/10.1038/s41586-020-2652-7).

40. Meseguer, A. S., & Condamine, F. L. (2020). Ancient tropical extinctions at high

latitudes contributed to the latitudinal diversity gradient. *Evolution,* in press

(https://doi.org/10.1111/evo.13967).

41. Miller, E. C., & Wiens, J. J. (2017). Extinction and time help drive the marine-terrestrial

biodiversity gradient: is the ocean a deathtrap? *Ecology Letters,* 20, 911–921.

42. Miller, E. C., Hayashi, K. T., Song, D., & Wiens, J. J. (2018). Explaining the ocean's

richest biodiversity hotspot and global patterns of fish diversity. *Proceedings of the Royal*

Society B, 285, 20181314.

43. Mittelbach, G. G., Schemske, D. W., Cornell, H. V., Allen, A. P., Brown, J. M., Bush, M.

B., ... & McCain, C. M. (2007). Evolution and the latitudinal diversity gradient:

speciation, extinction and biogeography. *Ecology letters*, 10, 315-331.

44. Moen, D., & Morlon, H. (2014). Why does diversification slow down? *Trends in Ecology*
and Evolution, 29, 190–197.

45. Oberdorff, T., Guégan, J. F., & Hugueny, B. (1995). Global scale patterns of fish species
richness in rivers. *Ecography,* 18, 345–352.

46. Oberdorff, T., Hugueny, B., & Guégan, J. F. (1997). Is there an influence of historical
events on contemporary fish species richness in rivers? Comparisons between Western
Europe and North America. *Journal of Biogeography.,* 24, 461–467.

47. O'Hara, T. D., Hugall, A. F., Woolley, S. N. C., Bribiesca-Contreras, G., & Bax, N. J.
(2019). Contrasting processes drive ophiuroid phylodiversity across shallow and deep
seafloors. *Nature,* 565, 636–639.

48. Pontarp, M., & Wiens, J. J. (2017). The origin of species richness patterns along
environmental gradients: uniting explanations based on time, diversification rate and
carrying capacity. *Journal of Biogeography,* 44, 722–735.

49. R Core Team (2008). R: A Language and Environment for Statistical Computing (R
Foundation for Statistical Computing, Vienna). Version 3.5.1. Available at www.r-
project.org. Accessed May 2019.

50. Rabosky, D. L. (2009). Ecological limits on clade diversification in higher taxa. *The*
American Naturalist, 173, 662–674.

51. Rabosky, D. L. (2014). Automatic detection of key innovations, rate shifts, and diversity-
dependence on phylogenetic trees. *PLoS One,* 9, e89543.

52. Rabosky, D. L. (2015). No substitute for real data: a cautionary note on the use of
phylogenies from birth-death polytomy resolvers for downstream comparative analyses.
Evolution, 69, 3207–3216.

53. Rabosky, D. L., Title, P. O., and Huang, H. (2015). Minimal effects of latitude on

present-day speciation rates in New World birds. *Proceedings of the Royal Society B,*

282, 20142889.

54. Rabosky, D. L., Chang, J., Title, P. O., Cowman, P. F., Sallan, L., Friedman, M., ... &

Alfaro, M. E. (2018). An inverse latitudinal gradient in speciation rate for marine

fishes. *Nature*, 559, 392–395.

55. Rabosky, D. L. (2020). Speciation rate and the diversity of fishes in freshwaters and the

oceans. *Journal of Biogeography,* 47, 1207–1217.

56. Ree, R. H., & Smith, S. A. (2008). Likelihood inference of geographic range evolution by

dispersal, local extinction, and cladogenesis. *Systematic Biology,* 57, 4–14.

57. Ricklefs, R. E. (1987). Community diversity: relative roles of local and regional

processes. *Science,* 235, 167–171.

58. Ronov, A. B. (1994). Phanerozoic transgressions and regressions on the continents: a

quantitative approach based on areas flooded by the sea and areas of marine and

continental deposition. *American Journal of Science,* 294, 777–801.

59. Roy, K., & Goldberg, E. E. (2007). Origination, extinction, and dispersal: integrative

models for understanding present-day diversity gradients. *The American Naturalist,* 170,

S71–S85.

60. Sanderson, M. J. 1996. How many taxa must be sampled to identify the root node of a

large clade? *Systematic Biology,* 45, 168–173.

61. Saupe, E. E., Farnsworth, A., Lunt, D. J., Sagoo, N., Pham, K. V., & Field, D. J. (2019).

Climatic shifts drove major contractions in avian latitudinal distributions throughout the

Cenozoic. *Proceedings of the National Academy of Sciences*, *116*, 12895-12900.

62. Seehausen, O., & Wagner, C. E. (2014). Speciation in freshwater fishes. *Annual Review*

of Ecology, Evolution and Systematics, 45, 621–651.

63. Schluter, D., & Pennell, M. W. (2017). Speciation gradients and the distribution of

biodiversity. *Nature,* 546, 48–55.

64. Smith, G. R., Badgley, C., Eiting, T. P., & Larson, P. S. (2010). Species diversity

gradients in relation to geological history in North American freshwater fishes. *Evolution*

and Ecology Research, 12, 693–726.

65. Stephens, P. R., & Wiens, J. J. (2003). Explaining species richness from continents to

communities: the time-for-speciation effect in emydid turtles. *American Naturalist,* 161,

112–128.

66. Tedesco, P. A., Oberdorff, T., Lasso, C. A., Zapata, M., & Hugueny, B. (2005). Evidence

of history in explaining diversity patterns in tropical riverine fish. *Journal of*

Biogeography, 32, 1899–1907.

67. Tedesco, P. A., Leprieur, F., Hugueny, B., Brosse, S., Durr, H. H., Beauchard, O., … &

Oberdorff, T. (2012). Patterns and processes of global riverine fish endemism. *Global*

Ecology and Biogeography, 21, 977–987.

68. Tedesco, P. A., Beauchard, O., Bigorne, R., Blanchet, S., Buisson, L., Conti, L., ... &

Jézéquel, C. (2017). A global database on freshwater fish species occurrence in drainage

basins. *Scientific Data*, 4, 170141.

69. Title, P. O., & Rabosky, D. L. (2019). Tip rates, phylogenies and diversification: What

are we estimating, and how good are the estimates? *Methods in Ecology and*

Evolution, 10, 821–834.

70. Tomašových, A., & Jablonski, D. (2017). Decoupling of altitudinal gradients in species

and genus geographic range size: a signature of clade range expansion. *Global Ecology*

and Biogeography, 26, 288–303.

71. Toussaint, E. F. A., Bloom, D., Short, A. E. Z. (2017). Cretaceous West Gondwana

vicariance shaped giant water scavenger beetle biogeography. *Journal of Biogeography,*

44, 1952–1965.

72. Unmack, P. J. (2001). Biogeography of Australian freshwater fishes. *Journal of*

Biogeography, 28, 1053–1089.

73. Wallace, A. R. (1878). *Tropical nature, and other essays*. London: Macmillan and

Company.

74. Weir, J. T., & Schluter, D. (2007). The latitudinal gradient in recent speciation and

extinction rates of birds and mammals. *Science,* 315, 1574–1576.

75. Wiens, J. J. (2012). Why biogeography matters: historical biogeography vs.

phylogeography and community phylogenetics for inferring ecological and evolutionary

processes. *Frontiers in Biogeography,* 4, 128–135.

76. Wiens, J. J., & Donoghue, M. J. (2004). Historical biogeography, ecology, and species

richness. *Trends in Ecology and Evolution,* 19, 639–644.

77. Willis, J. C. (1922). Age and area: a study in geographical distribution and origin of

species. Cambridge: Cambridge University Press.

78. Wood, S. N. (2011). Fast stable restricted maximum likelihood and marginal likelihood

estimation of semiparametric generalized linear models. *Journal of the Royal Statistical*

Society: Series B, 73, 3–36.

79. Wood, S. N. (2017). *Generalized additive models: an introduction with R*. Boca Raton,

801   FL: CRC Press.

803
804
805

FIGURE 1. Schematic illustrating alternative pathways to building species richness through in-

situ speciation. In all three examples, a lineage colonizes a region and diversifies with no further

immigration or emigration. In (**A**), the lineage colonizes the region relatively early and

diversifies at a modest rate. In (**B**), the lineage colonizes later, but diversifies at a faster rate and

achieves the same species richness as (**A**). In (**C**), the lineage colonizes relatively early and also

diversifies quickly, achieving high richness. While diversification rates are constant in these

simple examples, these scenarios are also applicable to cases where rates change through time

(Pontarp & Wiens, 2017).

FIGURE 2. Latitudinal (**A**) and longitudinal (**B**) gradients of species richness, surface area,

mean net diversification rates, and mean colonization times of freshwater drainage basins.

Species richness and surface area is derived from Tedesco et al. (2017). Diversification rates

shown here were estimated using BAMM under a time-constant rates model; values represent the

mean tip-associated values of species found in each basin (Rabosky et al., 2018). Colonization

timing of biogeographic regions was inferred from ancestral range reconstructions (Matzke,

2014); values represent the mean regional colonization time for species in each basin. For GAM

results see Table S2.

FIGURE 3. Results for spatially explicit GAMs for predictors of species richness of freshwater

drainage basins. Geographical distribution of global freshwater fish (**A**) richness, (**B**) net

diversification rates, and (**C**) time-for-speciation. Species richness of basins is based on

occurrences from Tedesco et al. (2017). BAMM-estimated rates of net diversification calculated

under a time-constant rates model (Rabosky et al., 2018) are shown here. Colonization times of

biogeographic regions were inferred from ancestral range reconstructions (Matzke, 2014; see

Extended Methods). Values of net diversification rates and colonization times represent the

means among co-occurring species in each basin. Species richness and diversification rates are

log-transformed. For full GAM results see Tables S3 and S4.

FIGURE 4. The negative relationship between net diversification rates and colonization times

(R^2=0.393, rho=-0.125, both P<0.001). Values represent the means among co-occurring species

in each drainage basin. BAMM-estimated rates of net diversification calculated under a time-

constant rates model (Rabosky et al., 2018) are shown here. Results under a time-varying rates

model were similar (Table S5). Colonization times of biogeographic regions were inferred from

ancestral range reconstructions (Matzke, 2014; see Extended Methods).

FIGURE 5. (**A**) Total regional species richness among three groups of ray-finned fishes:

Otophysi (red), Percomorpha (blue), and all other groups combined (green). Richness estimates

are from Tedesco et al. (2017). Exemplar members of each group are illustrated. Paintings by

Kathryn Chenard. (**B**) Visualization of the relative contribution of colonization events though

time. Each bubble represents a lineage that colonized independently; bubbles are scaled by the

proportion of living species in the region descended from that event. The total number of

colonizations in each region is noted. We used stochastic mapping (Dupin et al., 2016) and the

phylogeny of Rabosky et al. (2018) to identify independently colonizing lineages and their living

descendants. One example of a stochastic map is shown here; for variation among maps and with

fossil data see Figure S3 and Extended Results 1. Mean and median times among stochastic

maps were used in GAM analyses (Figures 2–4). Note that the colonization of a region can

precede the origin of focal clades. For example, the MRCA of ray-finned fishes was

reconstructed in the Afrotropic and Nearctic regions, and these ranges were inherited along the

tree. Clade names represent the descendants of biogeographic events. See Figure S2 for an

illustration of how this information was obtained from stochastic map results.