Full Title

"Aligning RNA-Seq reads to a sex chromosome complement informed reference genome

increases ability to detect sex differences in gene expression"

Short Title

"*Sex chromosome complement informed alignment*"

Authors

Kimberly C. Olney[1,2]¶, Sarah M. Brotman[1,4]¶, Valeria Valverde-Vesling[1], Jocelyn Andrews[1], and

Melissa A. Wilson[1,2,3]*

Affiliations

1. School of Life Sciences,

2. Center for Evolution and Medicine,

3. Center for Mechanisms of Evolution, The Biodesign Institute

Arizona State University, Tempe AZ 85282 USA

4. Department of Genetics, University of North Carolina, Chapel Hill NC 27599 USA

*Corresponding author

Melissa A. Wilson

School of Life Sciences | Arizona State University | PO Box 874501 | Tempe, AZ 85287-4501

mwilsons@asu.edu

Author contributions

¶ These authors contributed equally to this work

KCO: Supervision, Formal Analysis, Investigation, Visualization, Writing - Original Draft

Preparation, Writing - Review and Editing

SMB: Formal Analysis, Investigation, Writing - Original Draft Preparation, Writing - Review

and Editing

VVV: Investigation, Writing - Review and Editing

JA: Investigation, Writing - Review and Editing

MAW: Conceptualization, Supervision, Visualization, Resources, Project Administration,

Writing - Original Draft Preparation, Writing - Review and Editing, Funding Acquisition

Abstract

Background: Human X and Y chromosomes share an evolutionary origin and, as a consequence,

sequence similarity. We investigated whether sequence homology between the X and Y

chromosomes affects alignment of RNA-Seq reads, and estimates of differential expression. We

tested the effects of using reference genomes informed by the sex chromosome complement of the

sample's genome on measurements of RNA-Seq abundance and sex-differences in expression.

Results: The default genome includes the entire human reference genome (GRCh38), including

the entire sequence of the X and Y chromosomes. We created two sex chromosome complement

informed reference genomes. One sex chromosome complement informed reference genome was

used for samples that lacked a Y chromosome; for this reference genome version, we hard-masked

the entire Y chromosome. For the other sex chromosome complement informed reference genome,

to be used for samples with a Y chromosome, we hard-masked only the pseudoautosomal regions

of the Y chromosome, because these regions are duplicated identically in the reference genome on

the X chromosome. We analyzed transcript abundance in the brain cortex, and whole blood from

three genetic female (46, XX) and three genetic male (46, XY) samples, using both HISAT and

STAR read aligners. Each sample was aligned twice; once to the default reference genome and

then independently aligned to a reference genome informed by the sex chromosome complement

of the sample. We then quantified sex-differences in gene expression using featureCounts to get

the raw count estimates followed by Limma/Voom for normalization and differential expression.

Conclusions: We show that regardless of the choice of read aligner, using an alignment protocol

informed by the sex chromosome complement of the sample results in higher expression estimates

on the X chromosome in both genetic male and genetic female samples and an increased number

of unique genes being called as differentially expressed between the sexes.

Author summary

The human X and Y chromosomes share an evolutionary origin and sequence homology, including regions of 100% identity; this sequence homology can result in reads misaligning between the sex chromosomes, X and Y. We hypothesized that misalignment of reads on the sex chromosomes would confound estimates of transcript abundance if the sex chromosome complement of the sample is not accounted for during the alignment step. For example, because of shared sequence similarity, X-linked reads could misalign to the Y chromosome. This is expected to result in reduced expression for regions between X and Y that share high levels of homology. For this reason, we tested the effect of using a default reference genome versus a reference genome informed by the sex chromosome complement of the sample on estimates of transcript abundance in human RNA-Seq samples from the brain cortex and whole blood of three genetic female (46, XX) and three genetic male (46, XY) samples. We found that using a reference genome with the sex chromosome complement of the sample resulted in higher measurements of X-linked gene transcription for both male and female samples and more differentially expressed genes on the X and Y chromosomes. We recommend future studies requiring aligning reads to a reference genome should consider the sex chromosome complement of their samples prior to running default pipelines.

Background

Sex differences in aspects of human biology, such as development, physiology, metabolism, and disease susceptibility are partially driven by sex-specific gene regulation [1–4]. There are reported sex differences in gene expression across human tissues [5–7] and while some may be attributed to hormones and environment, there are documented genome-wide sex differences in expression based solely on the sex chromosome complement [8]. However, accounting for the sex chromosome complement of the sample in quantifying gene expression has been limited due to shared sequence homology between the sex chromosomes, X and Y, that can confound gene expression estimates.

The X and Y chromosomes share an evolutionary origin: mammalian X and Y chromosomes originated from a pair of indistinguishable autosomes ~180-210 million years ago that acquired the sex-determining genes [9–11]. The human X and Y chromosomes formed in two different segments: a) one that is shared across all mammals called the X-conserved region (XCR) and b) the X-added region (XAR) that is shared across all eutherian animals [10]. The sex chromosomes, X and Y, previously recombined along their entire lengths, but due to recombination suppression from Y chromosome-specific inversions [9,12], now only recombine at the tips in the pseudoautosomal regions (PAR) PAR1 and PAR2 [9–11]. PAR1 is ~2.78 million bases (Mb) and PAR2 is ~0.33 Mb; these sequences are 100% identical between X and Y [10,13,14] (Figure 1A). The PAR1 is a remnant of the XAR [10] and shared among eutherians, while the PAR2 is recently added and human-specific [14]. Other regions of high sequence similarity between X and Y include the X-transposed-region (XTR) with 98.78% homology [15] (Figure 1A). The XTR formed from an X chromosome to the Y chromosome duplication event following the human-chimpanzee divergence [10,16]. Thus, the evolution of the X and Y

chromosomes has resulted in a pair of chromosomes that are diverged, but still share some regions

of high sequence similarity.

To infer which genes or transcripts are expressed, RNA-Seq reads can be aligned to a

reference genome. The abundance of reads mapped to a transcript is reflective of the amount of

expression of that transcript. RNA-Seq methods rely on aligning reads to an available high quality

reference genome sequence, but this remains a challenge due to the intrinsic complexity in the

transcriptome of regions with a high level of homology [17]. By default, the GRCh38 version of

the human reference genome includes both the X and Y chromosomes, which is used to align

RNA-Seq reads from both male XY and female XX samples. It is known that sequence reads from

DNA will misalign along the sex chromosomes affecting downstream analyses [18]. However,

this has not been tested using RNA-Seq data and the effects on differential expression analysis are

not known. Considering the increasing number of human RNA-Seq consortium datasets (e.g., the

Genotype-Tissue Expression project (GTEx) [19], The Cancer Genome Atlas (TCGA) [20],

Geuvadis project [21], and Simons Genome Diversity Project [22]), there is an urgent need to

understand how aligning to a default reference genome that includes both X and Y may affect

estimates of gene expression on the sex chromosomes [1,23]. We hypothesize that regions of high

sequence similarity will result in misaligning of RNA-Seq reads and reduced expression estimates

(Figure 1A & B).

Here, we tested the effect of sex chromosome complement informed read alignment to the

quantified levels of gene expression and the ability to detect sex-biased gene expression. We

utilized data from the GTEx project, focusing on two tissues, whole blood and brain cortex, which

are known to exhibit sex differences in gene expression [24–26]. Many genes have been reported

to be differentially expressed between male and female brain samples [5–7]. Differential

expression in blood samples between males and females has also been documented [5,6]. We used

brain cortex and whole blood tissues from three genetic male (46, XY) and three genetic female

(46, XX) individuals for a total of twelve samples evenly distributed among tissues and genetic

sex. We aligned all samples to a default reference genome that includes both the X and Y

chromosomes and to a reference genome that is informed on the sex chromosome complement of

the genome: Male XY samples were aligned to a reference genome that includes both the X and

Y chromosome where the Y chromosome PAR1 and PAR2 are hard-masked with Ns (Figure 1C)

so that reads will align uniquely to the X PAR sequences. Conversely, female XX samples were

aligned to a reference genome where the entirety of the Y chromosome is hard-masked (Figure

1C). We tested two different read aligners, HISAT [27] and STAR [28], to account for variation

between alignment methods and measured differential expression using Limma/Voom [29]. We

found that using a sex chromosome complement informed reference genome for aligning RNA-

Seq reads increased X chromosome expression estimates in both male XY and female XX samples

and uniquely identified differentially expressed genes.

Figure 1. Homology between the human X and Y chromosomes where misaligning could occur. A) High sequence homology exists between the human X and Y chromosomes in three regions: 100% sequence identity for the pseudoautosomal regions (PARs), PAR1 and PAR2 and ~99% sequence homology in the X-transposed region (XTR). The X chromosome PAR1 is ~2.78 million bases (Mb) extending from X:10,001 to 2,781,479 and the X chromosome PAR2 is ~0.33

Mb extending from X:155,701,383 to 156,030,895. The X chromosome PAR1 and PAR2 are

identical in sequence to the Y chromosome PAR1 Y:10,001 - 2,781,479 and PAR2 Y:56,887,903

- 57,217,415. **B)** Using a standard alignment approach will result in reads misaligning between

regions of high sequence homology on the sex chromosomes. **C)** Using a reference genome that

is informed by the genetic sex of the sample may help to reduce misaligning between the X and Y

chromosomes. In humans, samples without evidence of a Y chromosome should be aligned to a

Y-masked reference genome and samples with evidence of a Y should be aligned to a YPARs-

masked reference genome.

Methods

Building sex chromosome complement informed reference genomes

All GRCh38.p10 unmasked genomic DNA sequences, including autosomes 1-22, X, Y,

mitochondrial DNA (mtDNA), and contigs were downloaded from ensembl.org release 92 [13].

The default reference genome here includes all 22 autosomes, mtDNA, the X chromosome, the Y

chromosome, and contigs. For the two sex chromosome complement informed reference

assemblies, we included all 22 autosomes, mtDNA, and contigs from the default reference and a)

one with the Y chromosome either hard-masked for the "Y-masked reference genome" or b) one

with the pseudoautosomal regions, PAR1 and PAR2, hard-masked on the Y chromosome for

"YPARs-masked reference genome" (Figure 1C). Hard-masking with Ns will force reads to not

align to those masked regions in the genome. Masking the entire Y chromosome for the sex

chromosome complement informed reference genome, Y-masked, was accomplished by changing

all the Y chromosome nucleotides [ATGC] to Ns using sed command in linux. YPARs-masked

was created by hard-masking the Y PAR1: 6001-2699520 and the Y PAR2: 154931044-

155260560 regions. The GRCh38.p10 Y PAR1 and Y PAR2 chromosome start and end location

was defined using Ensembl GRCh38 Y PAR definitions [13]. After creating the Y chromosome

PAR1 and PAR2 masked fasta files, we concatenated all the Y chromosome regions together to

create a YPARs-masked reference genome. After creating the GRCh38.p10 default reference

genome and the two sex chromosome complement informed reference genomes, we indexed the

reference genomes and created a dictionary for each using HISAT version 2.1.0 [27] hisat2-build

-f option and STAR version 2.5.2 [28] using option --genomeDir and --sjdbGTFfile. Reference

genome indexing was followed by picard tools version 1.119 CreateSequenceDictionary [30],

which created a dictionary for each reference genome (Pipeline available on GitHub,

https://github.com/SexChrLab/XY_RNAseq).

RNA-Seq samples

From the Genotyping-Tissue Expression (GTEx) Project data, we downloaded SRA files for brain

cortex and whole blood tissues from 3 genetic female (46, XX) and 3 genetic male (46, XY)

individuals [19,31] (Additional file 1). The GTEx data is described and available through dbGaP

under accession phs000424.v6.p1; we received approval to access this data under dbGaP accession

#8834. Although information about the genetic sex of the samples was provided in the GTEx

summary downloads, it was additionally investigated by examining the gene expression of select

genes that are known to be differentially expressed between the sexes or are known X-Y

homologous genes: DDX3X, DDX3Y, PCDH11X, PCDH11Y, USP9X, USP9Y, USP9Y, ZFX,

ZFY, UTX, UTY, XIST, and SRY (Figure 2).

Figure 2. Genetic sex of RNA-Seq samples. We investigated gene expression, $\log_2$(CPM+0.25/L), of XY homologous genes, XIST, and SRY in all samples analyzed here from genetic males (blue squares) and genetic females (orange circles) **A)** when aligned to a default reference genome, and **B)** when aligned to a sex chromosome complement informed reference genome, using HISAT as the read aligner.

RNA-Seq trimming and quality filtering

RNA-Seq sample data was converted from sequence read archive (sra) format to the paired-end

FASTQ format using the SRA toolkit [32]. Quality of the samples' raw sequencing reads was

examined using FastQC [33] and MultiQC [34]. Subsequently, adapter sequences were removed

using Trimmomatic version 0.36 [35]. More specifically, reads were trimmed to remove bases with

a quality score less than 10 for the leading strand and less than 25 for the trailing strand, applying

a sliding window of 4 with a mean PHRED quality of 30 required in the window and a minimum

read length of 40 bases.

RNA-Seq read alignment

Following trimming, paired RNA-Seq reads from all samples were aligned to the default reference

genome. Unpaired RNA-Seq reads were not used for alignment. Reads from the female (46, XX)

samples were aligned to the Y-masked genome and reads from male (46, XY) individuals were

aligned to the YPARs-masked reference genome. Read alignment was performed using HISAT

version 2.1.0 [27], keeping all parameters the same, only changing the reference genome used, as

described above. Read alignment was additionally performed using STAR version 2.5.2 [28],

where all samples were aligned to a default reference genome and to a reference genome informed

on the sex chromosome complement, keeping all parameters the same (Pipeline available on

GitHub, https://github.com/SexChrLab/XY_RNAseq).

Processing of RNA-Seq alignment files

Aligned RNA-Seq samples from HISAT and STAR were output in Sequence Alignment Map

(SAM) format and converted to Binary Alignment Map (BAM) using bamtools version 2.4.0 [36].

Summaries on the BAM files, including the number of reads mapped, duplicate reads and

unaligned reads were computed using bamtools version 2.4.0 package [36] (Additional file 2).

RNA-Seq BAM files were indexed, sorted, duplicates were marked, and read groups added using

bamtools, samtools, and Picard [36–38]. All RNA-Seq BAM files were indexed using the default

reference genome using Picard ReorderSam [38], this was done so that all samples would include

all chromosomes in the index files. Aligning XX samples to a Y-masked reference genome using

HISAT indexes would result in no Y chromosome information in the aligned bam and bam index

bai files. For downstream analysis, some tools will require that all samples have the same

chromosomes, this is why we hard-mask rather than delete. Reindexing the BAM files to the

default reference genome does not alter the read alignment, and thus does not alter our comparison

between default and sex chromosome complement informed alignment.

Gene expression level quantification

Read counts for each gene across all autosomes, sex chromosomes, mtDNA, and contigs were

generated using featureCounts [39] for all aligned and processed RNA-Seq BAM files. Female

XX samples when aligned to a sex chromosome complement informed reference genome will

show zero counts for Y-linked genes but will still include those genes in the raw counts file. This

is an essential step for downstream differential expression analysis between males and females to

keep the total genes the same between the sexes for comparison. Only rows that matched gene

feature type in Ensembl Homo_sapiens.GRCh38.89.gtf gene annotation [13] were included for

read counting. There are 2,283 genes annotated on the X chromosome and a total of 56,571 genes

across the entire genome for GRCh38 version of the human reference genome [13]. Only primary

alignments were counted and specified using the --primary option in featureCounts.

Differential expression

Differential expression analysis was performed using the limma/voom pipeline [29,40] which has

been shown to be a robust differential expression software package [41,42]. Quantified read counts

from each sample generated from featureCounts were combined into a count matrix, each row

representing a unique gene id and each column representing the gene counts for each unique

sample. This was repeated for each tissue type, brain cortex and whole blood, and was read into R

using the DGEList function in the R limma package [29,40]. A sample-level information file

related to the genetic sex of the sample, male or female, and the reference genome used for

alignment, default or sex chromosome complement informed, was created and corresponds to the

columns of the count matrix described above.

Using edgeR [43], raw counts were normalized to adjust for compositional differences

between the RNA-Seq libraries using the voom normalize quantile function which normalizes the

reads by the method of trimmed mean of values (TMM) [29]. Counts were then transformed to

$\log_2(CPM+0.25/L)$, where CPM is counts per million, L is library size, and 0.25 is a prior count to

avoid taking the log of zero [43]. For this dataset, the average library size is about 15 million,

therefore L is ~15. Thus, the minimum $\log_2(CPM+0.25/L)$ value for each sample, representing

zero transcripts, is $\log_2(0+0.25/15) = -5.90$.

A minimum of 1 CPM, or the equivalent of 0 in $\log_2(CPM+2/L)$, in at least 3 samples per

comparison was required for the gene to be kept for downstream analysis. A CPM value of 1 was

used in our analysis to separate expressed genes from unexpressed genes, meaning that in a library

size of ~15 million reads, there are at least 15 counts in that sample. After filtering for a minimum

CPM, 22,695 out of the 56,571 quantified genes were retained for the brain samples and 14,944

for whole blood, as differential expression between the sexes was run separately for each tissue.

A linear model was fitted to the DGEList-object, which contains the normalized gene counts for

each sample, using the limma lmfit function which will fit a separate model to the expression

values for each gene [29].

For differential expression analysis a design matrix containing the genetic sex of the sample

(male or female) and which reference genome the sample was aligned to (default or sex

chromosome complement informed) was created for each tissue type brain cortex and whole blood

for contrasts of pairwise comparisons between the sexes and between reference genomes used for

alignment. Pairwise contrasts were generated using limma makecontrasts function [29]. We

identified genes that exhibited significant expression differences defined using an adjusted p-value

cutoff that is less than 0.05 (5%) to account for multiple testing in pairwise comparisons between

conditions using limma/voom decideTests vebayesfit [29,40] (Pipeline available on GitHub,

https://github.com/SexChrLab/XY_RNAseq).

GO analysis

We examined differences and similarities in gene enrichment terms between the differentially

expressed genes obtained from the differential expression analyses of the samples aligned to the

default and sex chromosome complement informed reference genomes, to investigate if the

biological interpretation would change depending on the reference genome the samples were

aligned to (Additional file 5). We investigated gene ontology enrichment for lists of genes that

were identified as showing overexpression in one sex versus the other sex for brain cortex and

whole blood samples (adjusted p-value < 0.05). We used the GOrilla webtool, which utilizes a

hypergeometric distribution to identify enriched GO terms [44]. A modified Fisher exact p-value

cutoff < 0.001 was used to select significantly enriched terms [44].

Results

RNA-Seq reads aligned to autosomes do not vary much between reference genomes

We compared total mapped reads when reads were aligned to a default reference genome and for

when reads were aligned to a reference genome informed on the sex chromosome complement

(Additional file 2). Reads mapped across the whole genome, excluding the sex chromosomes,

either stayed the same or increased slightly when samples were aligned to a reference genome

informed on the sex chromosome complement. This was true regardless of the read aligner used,

HISAT or STAR, or of the sex of the sample, XY or XX. To test the effects of realignment on an

autosome, we selected chromosome 8, because it is of similar size to chromosome X. Reads

aligning to chromosome 8 didn't vary tremendously or at all between aligning to the default versus

sex chromosome complement informed reference for both brain cortex and whole blood tissues

(1,479,544 versus 1,479,584 average number of reads mapped in default versus sex chromosome

complement informed, respectively when aligned using HISAT; Additional file 2). A female XX

brain cortex sample showed the largest chromosome 8 mapped read increase (334 more reads with

HISAT; 608 more reads with STAR) when aligned to a sex chromosome complement informed

reference genome using either aligner (Additional file 2). The sample that showed the highest

decrease in mapped reads was a male XY brain cortex sample, which showed 5 fewer reads on

chromosome 8 when aligned to a reference genome informed on the sex chromosome complement

using HISAT read aligner. There was no decrease in chromosome 8 mapped reads when aligned

to a sex chromosome complement informed reference genome using STAR read aligner for male

or female whole blood samples. Overall, we observed little difference in chromosome 8 mapped

reads but were interested in how reads changed on the sex chromosomes, X and Y, when aligned

to a sex chromosome complement informed reference genome compared to aligning to a default

reference genome.

Reads aligned to the X chromosome increase in both XX and XY samples when using a sex

chromosome complement informed reference genome

We found that when reads were aligned to a reference genome informed by the sex chromosome

complement for both male XY and female XX brain cortex and whole blood samples, reads on the

X chromosome increased by ~0.13% (1,627 increase in reads out of an average 1,238,463 reads

for chromosome X) when aligned using HISAT. Reads on the Y chromosome decreased 100%

(46,852 reads on average) in female XX samples and by ~63.43% (52,327 reads on average) in

male XY samples for brain cortex and whole blood when aligned using HISAT (Additional file

2). Similar increases in X chromosome and decreases in Y chromosome reads when aligned to a

sex chromosome complement informed reference was observed for when STAR was used as the

read aligner for both male XY and female XX brain cortex and whole blood samples (Additional

file 2). While we observed tens of thousands of reads aligning to new locations when sex

chromosome complement was taken into account, we also investigated the effect on expression

estimates.

Aligning to a sex chromosome complement informed reference genome increases the X

chromosome PAR1 and PAR2 expression

We explored the effect of changes in read alignment on gene expression. There is an increase in X

chromosome PAR1 and PAR2 expression when reads were aligned to a reference genome

informed on the sex chromosome complement and this is true for both male XY and female XX

samples from the brain cortex (Figure 3) and whole blood samples using either HISAT or STAR

as the read aligner (Additional file 6). We found an average of 2.83 fold increase in expression

(1.21 $\log_2$ fold increase) in PAR1 expression for female XX brain cortex samples and 2.71 fold

increase in expression (0.95 $\log_2$ fold change increase) in PAR1 for male XY brain cortex samples

using HISAT read aligner (Figure 3, Additional file 6, Additional file 7). XTR in female XX brain

cortex samples showed a 1.49 fold increase in expression (0.22 $\log_2$ fold increase) and no change

in male XY brain cortex samples. PAR2 showed an average of 2.47 fold increase (0.81 $\log_2$ fold

change increase) for female XX brain cortex samples and 2.19 fold increase (0.58 $\log_2$ fold change

increase) in PAR2 for male XY brain cortex samples using HISAT read aligner with similar results

for STAR read aligner (Figure 3, Additional file 6, Additional file 7). Complete lists of the

$\log_2$(CPM+0.25/L) values for each X chromosomal gene and each gene within the whole genome

for male XY and female XX brain cortex and whole blood samples for when reads were aligned

using the different read aligners and reference genomes are in Additional file 3 and Additional file

4.

Figure 3. X chromosome RNA-Seq alignment differences. We plot $\log_2$ fold change (FC) across

A) the entire X chromosome and **B)** the first 5 million bases (Mb) and show **C)** average fold change

in large genomic regions on the X chromosome between aligning brain cortex using HISAT to the

default genome and aligning to a sex-chromosome complement informed reference genome. For

$\log_2$ FC, a value less than zero indicates that the gene showed higher expression when aligned to

a default reference genome, while values above zero indicate that the gene shows higher expression

when aligned to a reference genome informed by the sex chromosome complement of the sample.

Samples from genetic females are plotted in orange circles, while samples from males are plotted

in blue squares. Darker shades indicate which gene points are in PAR1, XTR, and PAR2 while

lighter shades are used for genes outside of those regions.

Regions outside the PARs and XTR show little difference in expression between reference genomes

Intriguingly, regions outside the PARs on the X chromosome, and across the genome, showed

little to no increase in expression when aligned to a sex chromosome complement informed

reference genome compared to aligning to a default reference genome (Additional file 7). X and

Y homologous genes showed little to no increase in expression when aligned to a sex chromosome

complement informed reference genome compared to aligning to a default reference genome with

the exception of PCDH11X gene (Additional file 8). PCDH11X showed a 1.50 fold increase (0.58

$\log_2$ fold increase) in female XX brain cortex samples with a similar increase, 1.47 fold, (0.55 $\log_2$

fold increase) in female XX whole blood samples using HISAT read aligner. A similar increase in

expression for PCDH11X was observed for female XX brain cortex and whole blood samples

when STAR was used as the read aligner (Additional file 8). PCDH11X showed no differences in

expression between reference genomes for male XY brain cortex and whole blood samples when

using either HISAT or STAR (Additional file 8). With noticeable increases in X chromosome gene

expression and decreases in Y chromosome expression when aligned to a sex chromosome

complement informed reference genome, we next investigated how this would affect gene

differential expression between the sexes.

A sex chromosome complement informed reference genome increases the ability to detect sex

differences in gene expression

Generally, when comparing gene expression differences between the sexes, we find more genes

are differentially expressed on the sex chromosomes, and fewer, or the same, are differentially

expressed on the autosomes when taking sex chromosome complement into account. At an

adjusted p-value of 0.05 and aligning with HISAT, we find 12 new genes (all on the Y

chromosome) that are only called as differentially expressed between the sexes in the brain cortex

when aligned to reference genomes informed on the sex chromosome complement (Figure 4).

Additionally, we find 49 genes (48 autosomal and 1 X-linked) that are called as differentially

expressed when using a default reference genome for aligning reads that are no longer called as

differentially expressed between the sexes in the brain cortex when aligning to a reference genome

informed by sex chromosome complement (Figure 4; Additional file 9). We observed similar

trends in changes for differential expression between male XY and female XX whole blood

samples using either HISAT or STAR as the aligner (Additional file 9, Additional file 10).

Figure 4. Sex chromosome complement informed alignment calls more sex-linked genes as being differentially expressed. A) Sex differences in gene expression, $\log_2(\text{CPM}+0.25/\text{L})$, between the three samples from genetic males and females are shown when aligning all samples to the default reference genome (left) and a reference genome informed on the sex chromosome complement (right). Each point represents a gene. Genes that are differentially expressed, adjusted p-value < 0.05 are indicated in black for autosomal genes, blue for Y-linked genes, and red for X-linked genes. **B)** We show overlap between genes that are called as differentially expressed when all samples are aligned to the default genome, and genes that are called as differentially expressed when aligned to a sex chromosome complement informed genome. When aligned to a default reference genome there were 95 genes that were differentially expressed between male XY and

female XX brain cortex samples. Of these 95 genes, 49 genes were uniquely called as differentially

expressed when aligned to the default reference genome but were not called as differentially

expressed when aligned to a sex chromosome complement informed reference genome. Of the 49

genes, 48 are autosomal and 1 is X-linked, RBM3. When samples were aligned to a reference

genome informed on the sex chromosome complement 58 genes were called as differentially

expressed between the sexes of which 12 were uniquely called in the sex chromosome complement

informed alignment. All 12 uniquely called differentially expressed genes when samples were

aligned to a sex chromosome complement informed reference genome are Y-linked.

Increased concordance between aligners when informed by sex chromosome complement

When using a default reference genome, and calling genes as differentially expressed between the

sexes, of all genes when aligned using either HISAT or STAR as the aligner, only 46% of genes

called as differentially expressed overlapped between the aligners, HISAT and STAR for brain

cortex. There is 67% concordance in genes being called as differentially expressed for whole blood

between aligners, HISAT and STAR, when samples were aligned to a default reference genome.

In contrast, when aligning samples to a reference genome informed by the sex chromosome

complement, we find increased concordance in all tissues for the set of genes called as

differentially expressed between the sexes when aligned using either HISAT or STAR (Additional

file 9).

Using sex-linked genes alone is inefficient for determining the sex chromosome complement of a

sample

The sex of each sample was provided in the GTEx manifest. We investigated the expression of

genes that could be used to infer the sex of the sample. We studied X and Y homologous genes,

(DDX3X/Y, PCDH11X/Y, USP9X/Y, ZFX/Y, UTX/Y), and XIST and SRY gene expression in

male XY and female XX brain cortex and whole blood (Figure 2). Both males XY and females

XX are expected to show expression for the X-linked homologs, whereas only XY samples should

show expression of the Y-linked homologs. Further, XIST expression should only be observed in

XX samples and SRY should only be expressed in samples with a Y chromosome. Using the

default reference genome for aligning male XY and female XX brain cortex and whole blood

samples, we observed a small number of reads aligning to the Y-linked genes in female XX

samples, but also observed clustering by sex for DDX3Y and XIST gene expression (Figure 2).

Male XY samples showed expression for DDX3X and DDX3Y (greater than 5 $\log_2$(CPM+2/L).

Female XX samples showed expression for XIST (greater than 2.5 $\log_2$(CPM+2/L) and male XY

samples showed little to no expression for XIST (less than 0 $\log_2$(CPM+2/L). In contrasts to the

default reference genome, when aligned to a sex chromosome complement informed reference

genome, samples cluster distinctly by sex for DDX3Y, PCDH11Y, USP9Y, ZFY, UTY, and XIST

all showing at least a 2.5 $\log_2$(CPM+2/L) difference between the sexes (Figure 2).

Discussion

The Ensembl GRCh38 human reference genome includes all 22 autosomes, mtDNA, the X

chromosome, the Y chromosome with the Y PARs masked, and contigs [13]. The Gencode hg19

human reference genome includes everything with nothing masked [45]. Neither Ensembl or

Gencode human reference genomes are correct for aligning both XX and XY samples. The sex

chromosome complement of the sample should be taken into account when aligning RNA-Seq

reads to reduce misaligning sequences.

Measurements of X chromosome expression increase for both male XY and female XX

brain cortex and whole blood samples when aligned to a sex chromosome complement informed

reference genome versus aligning to a default reference genome (Figure 3). There was a minimum

1.6 fold increase in expression for the genes in PAR1 and PAR2 for all samples (male and female)

when aligned to a sex chromosome complement informed reference genome (Figure 3; Additional

file 7). We see that the XTR has a minimum 1.35 fold increase in expression for female brain

cortex and whole blood samples, when aligned to a sex chromosome complement informed

reference genome compared to a default reference genome, but no change in males. This is because

XTR is not hard-masked in the YPARs-masked reference genome, which is used to align male XY

samples. The XTR shares 98.78% homology between X and Y but no longer recombines between

X and Y [15] (Figure 1A) and is therefore, because of this divergence, not hard-masked when

aligning male XY samples. A comprehensive table for mean expression values for regions of the

X chromosome and X chromosome gene expression in default and sex chromosome complement

informed alignment is in additional file 7 and additional file 3, respectively. Given striking

increases in measurements of X chromosome expression, we further investigated the effect of sex

chromosome complement informed alignment on differential gene expression between the sexes.

Differential expression results changed when using a sex chromosome complement

informed alignment compared to using a default alignment. When aligned to a default reference

genome, due to sequence similarity, some reads from female XX samples aligned to the Y

chromosome (Figure 2; Figure 4). However, when aligned to a reference genome informed by the

sex chromosome complement, female XX samples no longer showed Y-linked gene expression,

and more Y-linked genes were called as being differentially expressed between the sexes. This

suggests that if using a default reference genome for aligning RNA-Seq reads, one would miss

some Y-linked genes as differentially expressed between the sexes (Figure 4). Furthermore, these

Y-linked genes serve in various important biological processes, thus altering the functional

interpretation of the sex differences. GO enrichment analysis of genes that are more highly

expressed in brain cortex male samples than females, when samples were aligned to a default

reference genome, were genes involved in germline cell cycle switching and mitotic to meiotic

cell cycle (Additional file 5). However, when these samples were aligned to a sex chromosome

complement informed reference genome, genes upregulated in males were enriched for positive

regulation of transcription from RNA polymerase II promoter in response to heat stress and GO

component of specific granule lumen, which are secretory vesicles of the immune system [44].

The choice of read aligner has long been known to give slightly differing results of

differential expression due to the differences in the alignment algorithms [42,46]. When using a

default reference genome for alignment we find discordance between HISAT and STAR in which

genes called as differentially expressed between the sexes. However, we find increased

concordance between HISAT and STAR, when using a sex chromosome complement informed

reference genome (Additional file 9). Differences between HISAT and STAR could be contributed

to differences in default parameters for handling multi-aligning reads [27]. We show here that the

choice of read aligner, HISAT and STAR, will have less variance on differential expression results

if a sex chromosome complement informed reference genome is used for aligning RNA-Seq reads.

Ideally, one would use DNA to confirm presence or absence of the Y chromosome, but if

DNA sequence was not generated, one would need to determine the sex of the sample by assessing

expression estimates for Y-linked genes. To more carefully investigate the ability to use gene

expression to infer sex chromosome complement of the sample, we examined the gene expression

for a select set of X-Y homologous genes, as well as XIST, and SRY that are known to be

differentially expressed between the sexes (Figure 2, Additional file 11). SRY is predominantly

expressed in the testis [47,48] and typically one would expect SRY to show male-specific

expression. In our set, we did not observe SRY expressed in any sample, and so it could not be

used to differentiate between XX and XY samples (Figure 2, Additional file 11). In contrast, the

X-linked gene XIST was differentially expressed between genetic males and genetic females in

both genome alignments (default and sex chromosome complement informed) for the brain cortex

and whole blood samples. XIST expression is important in the X chromosome inactivation process

[49] and serves to distinguish samples with one X chromosome from those with more than one X

chromosome [23]. However, this does not inform about whether the sample has a Y chromosome

or not. For X-Y homologous genes, we do not find sex differences in read alignment with either

default or sex chromosome complement informed for the X-linked homolog. When aligned to a

default reference genome, female XX samples showed some expression for homologous Y-linked

genes, so only presence/absence of Y-linked reads alone is insufficient to determine sex

chromosome complement of the sample (Figure 2, Additional file 11). That said, the samples

broadly segregated by sex for Y-linked gene expression using default alignment. However, the

pattern was messy for each individual Y-linked gene. Thus, if inferring sex from RNA-Seq data,

we recommend using the estimated expression of multiple X-Y homologous genes and XIST to

infer the genetic sex of the sample. Samples should be aligned to a default reference genome first

to look at the expression for several Y-specific genes to determine if the sample is XY or XX.

Then samples should be realigned to the appropriate sex chromosome complement informed

reference genome. Independently assessing sex chromosome complement of samples becomes

increasingly important as karyotypically XY individuals are known to have lost the Y chromosome

in particular tissues sampled, as shown in Alzheimer Disease [50], age-related macular

degeneration [51], and in the blood of aging individuals [52]. Self-reported sex may not match the

sex chromosome complement of the samples, even in karyotypic individuals.

Conclusion

Here we show that aligning RNA-Seq reads to a sex chromosome complement informed reference

genome will change the results of the analysis compared to aligning reads to a default reference

genome. We have previously observed that a sex chromosome complement informed alignment is

important for DNA as well [53]. A sex chromosome complement informed approach is needed for

a sensitive and specific analysis of gene expression on the sex chromosomes [1]. A sex

chromosome complement informed reference alignment resulted in increased X chromosome

expression for both male XY and female XX samples. We further found different genes called as

differentially expressed between the sexes, and identified sex differences in gene pathways that

were missed when samples were aligned to a default reference genome. We additionally identified

that there is greater concordance between read aligners, HISAT and STAR, in the genes that are

called as differentially expressed between the sexes when samples were aligned to a sex

chromosome complement informed reference genome. The accurate alignment of the short RNA-

Seq reads to the reference genome is essential to drawing reliable conclusions from differential

expression data analysis on the sex chromosomes. We strongly urge future human RNA-Seq

analysis to carefully consider the genetic sex of the sample when aligning reads, and have provided

a framework for doing so (https://github.com/SexChrLab/XY_RNAseq).

Acknowledgments

This research was supported by startup funds from the School of Life Sciences and the Biodesign

Institute at Arizona State University to MAW, School of Life Sciences Undergraduate Research

(SOLUR) funding to SMB, IMSD funding to VVV, ARCS funding to KCO. This study was

supported by the National Institute of General Medical Sciences of the National Institutes of Health

under Award Number R35GM124827 to MAW. The content is solely the responsibility of the

authors and does not necessarily represent the official views of the National Institutes of Health.

We thank Heini Natri and Angela Taravella for comments on the manuscript.

Additional files

Additional file 1. Sample IDs. RNA-Seq brain cortex and whole blood tissue samples from 3

genetic female (46, XX) and 3 genetic male (46, XY) individuals were downloaded from the

Genotype-Tissue Expression (GTEx) project [19,31] for a total of 12 RNAseq tissue samples.

Additional file 2. Table of mapped read statistics for default and sex chromosome

complement informed alignment. Total reads mapped in brain cortex and whole blood female

XX and male XY samples for when reads were aligned to a default reference genome and for when

reads were aligned to a reference genome informed on the sex chromosome complement. Total

mapped reads for HISAT and STAR as the read aligner. The difference in reads mapped between

sex chromosome complement informed and default alignment is shown to the right of the mapped

reads statistics for each tissue and aligner used. Chromosome Y and chromosome X show the

highest degree of difference between default and sex chromosome complement informed

alignment were chromosome Y always decreased in total mapped reads and chromosome X always

increased in total mapped reads when using a sex chromosome complement informed alignment.

Additional file 3. X chromosome gene expression values per sample, aligner and reference

genome used for alignment. CPM values for male XY and female XX brain cortex and whole

blood samples when aligned to a default and sex chromosome complement informed reference

genome for chromosome X. A text format of CPM values are available on GitHub,

https://github.com/SexChrLab/XY_RNAseq.

Additional file 4. Whole genome gene expression values per sample, aligner and reference

genome used for alignment. CPM values for male XY and female XX brain cortex and whole

blood samples when aligned to a default and sex chromosome complement informed reference

genome for the whole genome (1-22, MT, X, Y and non-chromosomal). A text format of CPM

values are available on GitHub, https://github.com/SexChrLab/XY_RNAseq.

Additional file 5. Gene enrichment analysis of genes that are more highly expressed in one sex

verses the other sex for when samples were aligned to a default or sex chromosome complement

informed reference genome using either HISAT or STAR.

Additional file 6. X chromosome expression differences between default and sex chromosome

complement informed alignment. X chromosome gene expression differences between default

and sex chromosome complement informed alignment. Increase in expression when aligned to a

sex chromosome complement informed reference genome is a $\log_2$ fold change (FC) > 0. A

decrease in expression when aligned to a sex chromosome complement informed reference

genome is $\log_2$ FC < 0. Female XX samples are indicated by red and pink circles for PAR1, XTR,

and PAR2 genes and for all other X chromosome genes respectively. Blue and light blue squares

represent male XY samples. Blue squares indicate which gene points are in PAR1, XTR, and

PAR2 and light blue squares are for genes outside of those regions. Differences in X chromosome

expression between reference genomes for male XY and female XX samples aligned using HISAT

for the whole X chromosome and the first 5Mb are shown for the brain cortex (**A** and **B**,

respectively), and the whole blood (**C** and **D**, respectively). Differences in X chromosome

expression between reference genomes for male XY and female XX samples aligned using STAR

for the whole X chromosome and the first 5Mb are shown for the brain cortex (**E** and **F,**

respectively), and the whole blood (**G** and **H**, respectively).

Additional file 7. X chromosome regions mean and median expression values. X chromosome

regions PAR1, PAR2, XTR, XDG, XAR, XCR mean and median CPM expression for male XY

and female XX brain cortex and whole blood samples when aligned to a default or sex chromosome

complement informed reference genome for HISAT and STAR.

Additional file 8. Gene expression for XY homologous genes. X chromosome expression for 26

X and Y homologous genes. Difference in gene expression for when male XY and female XX

brain cortex and whole blood samples were aligned to a default and sex chromosome complement

informed reference genome. Little to no difference in gene expression between default and sex

chromosome complement informed reference genome alignment was observed for 25 of the 26 X

and Y homologous genes for both male XY and female XX brain cortex and whole blood samples

using either HISAT or STAR. PCDH11X showed a 1.50 and 1.47 fold increase in expression for

brain cortex and whole blood, respectively in female XX samples using HISAT read aligner with

similar results for STAR. XY male brain cortex and whole blood samples showed little to no

differences in gene expression between reference genomes for the 26 X and Y homologous genes

using either HISAT or STAR.

Additional file 9. Differentially expressed genes between the sexes that were uniquely and

jointly called between reference genomes. Genes that are differentially expressed between the

sexes, male XY and female XX, for brain cortex and whole blood samples. Differentially

expressed genes that are uniquely called when using either the default or sex chromosome

complement informed reference genome and differentially expressed genes that jointly called

between the reference genomes.

Additional file 10. Gene expression differences between male XY and female XX samples.

Sex differences in gene expression for brain cortex and whole blood samples for when samples

were aligned to a default reference genome and a to a reference genome informed on the sex

chromosome complement. Showing sex differences in gene expression between reference

genomes used for alignment and for when samples were aligned using HISAT and STAR.

Additional file 11. Genetic sex of RNA-Seq samples. Gene expression $\log_2$(CPM+0.25/L) for

select XY homologous genes and XIST and SRY for when reads were aligned to a default

reference genome **A)** and **C)** using HISAT and STAR, respectively and for when reads were

aligned to a sex chromosome complement informed reference genome **B)** and **D)** using HISAT

and STAR, respectively. Male XY brain cortex and whole blood samples are shown in blue squares

and female XX brain and blood samples shown in red circles.

References

1.   Khramtsova EA, Davis LK, Stranger BE. The role of sex in the genomics of human

complex traits. Nat Rev Genet. 2019;20: 173–190.

2.   Arnold AP, Chen X, Itoh Y. What a Difference an X or Y Makes: Sex Chromosomes, Gene

Dose, and Epigenetics in Sexual Differentiation. Handbook of Experimental Pharmacology.

2012. pp. 67–88.

3.   Traglia M, Bseiso D, Gusev A, Adviento B, Park DS, Mefford JA, et al. Genetic

Mechanisms Leading to Sex Differences Across Common Diseases and Anthropometric

Traits. Genetics. 2017;205: 979–992.

4.   Raznahan A, Parikshak NN, Chandran V, Blumenthal JD, Clasen LS, Alexander-Bloch AF,

et al. Sex-chromosome dosage effects on gene expression in humans. Proc Natl Acad Sci U

S A. 2018;115: 7398–7403.

5.   Goldstein JM, Holsen L, Handa R, Tobet S. Fetal hormonal programming of sex differences

in depression: linking women's mental health with sex differences in the brain across the

lifespan. Front Neurosci. 2014;8: 247.

6.   Gershoni M, Pietrokovski S. The landscape of sex-differential transcriptome and its

consequent selection in human adults. BMC Biol. 2017;15: 7.

7.   Shi L, Zhang Z, Su B. Sex Biased Gene Expression Profiling of Human Brains at Major

Developmental Stages. Sci Rep. 2016;6. doi:10.1038/srep21181

8.   Arnold AP, Chen X. What does the "four core genotypes" mouse model tell us about sex

658  differences in the brain and other tissues? Front Neuroendocrinol. 2009;30: 1–9.

659 9. Lahn BT, Page DC. Four evolutionary strata on the human X chromosome. Science.

660   1999;286: 964–967.

661 10. Ross MT, Grafham DV, Coffey AJ, Scherer S, McLay K, Muzny D, et al. The DNA

662   sequence of the human X chromosome. Nature. 2005;434: 325–337.

663 11. Charlesworth B. The evolution of sex chromosomes. Science. 1991;251: 1030–1033.

664 12. Pandey RS, Wilson Sayres MA, Azad RK. Detecting evolutionary strata on the human x

665   chromosome in the absence of gametologous y-linked sequences. Genome Biol Evol.

666   2013;5: 1863–1871.

667 13. Aken BL, Achuthan P, Akanni W, Amode MR, Bernsdorff F, Bhai J, et al. Ensembl 2017.

668   Nucleic Acids Res. 2017;45: D635–D642.

669 14. Charchar FJ, Svartman M, El-Mogharbel N, Ventura M, Kirby P, Matarazzo MR, et al.

670   Complex events in the evolution of the human pseudoautosomal region 2 (PAR2). Genome

671   Res. 2003;13: 281–286.

672 15. Veerappa AM, Padakannaya P, Ramachandra NB. Copy number variation-based

673   polymorphism in a new pseudoautosomal region 3 (PAR3) of a human X-chromosome-

674   transposed region (XTR) in the Y chromosome. Funct Integr Genomics. 2013;13: 285–293.

675 16. Skaletsky H, Kuroda-Kawaguchi T, Minx PJ, Cordum HS, Hillier L, Brown LG, et al. The

676   male-specific region of the human Y chromosome is a mosaic of discrete sequence classes.

677   Nature. 2003;423: 825–837.

17. Piskol R, Ramaswami G, Li JB. Reliable identification of genomic variants from RNA-seq

data. Am J Hum Genet. 2013;93: 641–651.

18. Webster TH, Couse M, Grande BM, Karlins E, Phung TN, Richmond PA, et al. Identifying,

understanding, and correcting technical biases on the sex chromosomes in next-generation

sequencing data [Internet]. doi:10.1101/346940

19. GTEx Consortium. Human genomics. The Genotype-Tissue Expression (GTEx) pilot

analysis: multitissue gene regulation in humans. Science. 2015;348: 648–660.

20. Cancer Genome Atlas Research Network, Weinstein JN, Collisson EA, Mills GB, Shaw

KRM, Ozenberger BA, et al. The Cancer Genome Atlas Pan-Cancer analysis project. Nat

Genet. 2013;45: 1113–1120.

21. Lappalainen T, Sammeth M, Friedländer MR, 't Hoen PAC, Monlong J, Rivas MA, et al.

Transcriptome and genome sequencing uncovers functional variation in humans. Nature.

2013;501: 506–511.

22. Mallick S, Li H, Lipson M, Mathieson I, Gymrek M, Racimo F, et al. The Simons Genome

Diversity Project: 300 genomes from 142 diverse populations. Nature. 2016;538: 201–206.

23. Tukiainen T, Villani A-C, Yen A, Rivas MA, Marshall JL, Satija R, et al. Landscape of X

chromosome inactivation across human tissues. Nature. 2017;550: 244–248.

24. Li R, Singh M. Sex differences in cognitive impairment and Alzheimer's disease. Front

Neuroendocrinol. 2014;35: 385–403.

25. de Perrot M, Licker M, Bouchardy C, Usel M, Robert J, Spiliopoulos A. Sex differences in

presentation, management, and prognosis of patients with non-small cell lung carcinoma. J

Thorac Cardiovasc Surg. 2000;119: 21–26.

26. Melé M, Ferreira PG, Reverter F, DeLuca DS, Monlong J, Sammeth M, et al. Human

genomics. The human transcriptome across tissues and individuals. Science. 2015;348:

660–665.

27. Kim D, Langmead B, Salzberg SL. HISAT: a fast spliced aligner with low memory

requirements. Nat Methods. 2015;12: 357–360.

28. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast

universal RNA-seq aligner. Bioinformatics. 2013;29: 15–21.

29. Law CW, Chen Y, Shi W, Smyth GK. voom: Precision weights unlock linear model

analysis tools for RNA-seq read counts. Genome Biol. 2014;15: R29.

30. Picard Toolkit [Internet]. Broad Institute, GitHub Repository; 2018. Available:

http://broadinstitute.github.io/picard/

31. GTEx Consortium. The Genotype-Tissue Expression (GTEx) project. Nat Genet. 2013;45:

580–585.

32. Sequence Read Archive. Downloading SRA data using command line utilities. National

Center for Biotechnology Information (US); 2011; Available:

https://www.ncbi.nlm.nih.gov/books/NBK158899/

33. Babraham Bioinformatics - FastQC A Quality Control tool for High Throughput Sequence

Data [Internet]. [cited 11 Sep 2017]. Available:

http://www.bioinformatics.babraham.ac.uk/projects/fastqc/

34. Ewels P, Magnusson M, Lundin S, Käller M. MultiQC: summarize analysis results for

multiple tools and samples in a single report. Bioinformatics. 2016;32: 3047–3048.

35. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence

data. Bioinformatics. 2014;30: 2114–2120.

36. Barnett DW, Garrison EK, Quinlan AR, Strömberg MP, Marth GT. BamTools: a C++ API

and toolkit for analyzing and managing BAM files. Bioinformatics. 2011;27: 1691–1692.

37. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence

Alignment/Map format and SAMtools. Bioinformatics. 2009;25: 2078–2079.

38. Picard Tools - By Broad Institute [Internet]. [cited 7 May 2019]. Available:

http://broadinstitute.github.io/picard/

39. Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for

assigning sequence reads to genomic features. Bioinformatics. 2014;30: 923–930.

40. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2 [Internet]. 2014. doi:10.1101/002832

41. Seyednasrollah F, Laiho A, Elo LL. Comparison of software packages for detecting

differential expression in RNA-seq studies. Brief Bioinform. 2015;16: 59–70.

42. Costa-Silva J, Domingues D, Lopes FM. RNA-Seq differential expression analysis: An

extended review and a software tool. PLoS One. 2017;12: e0190152.

43. Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for differential

expression analysis of digital gene expression data [Internet]. Bioinformatics. 2010. pp.

139–140. doi:10.1093/bioinformatics/btp616

44. Eden E, Navon R, Steinfeld I, Lipson D, Yakhini Z. GOrilla: a tool for discovery and

visualization of enriched GO terms in ranked gene lists. BMC Bioinformatics. 2009;10: 48.

45. Harrow J, Frankish A, Gonzalez JM, Tapanari E, Diekhans M, Kokocinski F, et al.

GENCODE: the reference human genome annotation for The ENCODE Project. Genome

Res. 2012;22: 1760–1774.

46. Conesa A, Madrigal P, Tarazona S, Gomez-Cabrero D, Cervera A, McPherson A, et al. A

survey of best practices for RNA-seq data analysis. Genome Biol. 2016;17: 13.

47. Albrecht KH, Young M, Washburn LL, Eicher EM. Sry expression level and protein

isoform differences play a role in abnormal testis development in C57BL/6J mice carrying

certain Sry alleles. Genetics. 2003;164: 277–288.

48. Turner ME, Ely D, Prokop J, Milsted A. Sry, more than testis determination? [Internet].

American Journal of Physiology-Regulatory, Integrative and Comparative Physiology.

2011. pp. R561–R571. doi:10.1152/ajpregu.00645.2010

49. Carrel L, Willard HF. X-inactivation profile reveals extensive variability in X-linked gene

expression in females. Nature. 2005;434: 400–404.

50. Dumanski JP, Lambert J-C, Rasi C, Giedraitis V, Davies H, Grenier-Boley B, et al. Mosaic

Loss of Chromosome Y in Blood Is Associated with Alzheimer Disease. Am J Hum Genet.

2016;98: 1208–1219.

51. Grassmann F, Kiel C, den Hollander AI, Weeks DE, Lotery A, Cipriani V, et al. Y

chromosome mosaicism is associated with age-related macular degeneration. Eur J Hum

Genet. 2019;27: 36–41.

52. Forsberg LA. Loss of chromosome Y (LOY) in blood cells is associated with increased risk

for disease and mortality in aging men. Hum Genet. 2017;136: 657–663.

53. Webster TH, Couse M, Grande BM, Karlins E, Phung TN, Richmond PA, et al. Identifying,

understanding, and correcting technical biases on the sex chromosomes in next-generation

sequencing data [Internet]. 2018. doi:10.1101/346940