The VTA modulates a frequency-specific gain of thalamocortical inputs in

infragranular layers of the gerbil auditory cortex

Authors: Michael GK Brunk[1], Katrina E Deane[1], Martin Kisse[1], Matthias Deliano[1], Frank W

Ohl[1,2,3], Michael T Lippert[1,2], Max FK Happel[1,3]

1 Department of Systems Physiology of Learning, Leibniz Institute for Neurobiology, 39118

Magdeburg, Germany

2 Center for Behavioral Brain Sciences (CBBS), 39106 Magdeburg, Germany

3 Institute for Biology, Otto-von-Guericke-University, 39120 Magdeburg, Germany

Correspondence should be addressed to:

Michael Brunk or Dr. Max Happel

Department of Systems Physiology of Learning

Leibniz Institute for Neurobiology

Brenneckestr. 6

D-39118 Magdeburg

Tel.: +49-(0)391-6263-94411

Fax: +49-(0)391-6263-95489

Mail:

mbrunk@lin-magdeburg.de

mhappel@lin-magdeburg.de

Author contributions:

FWO, MTL and MFKH designed research. MGKB and KED performed research. MGKB ,

KED, MK, MD and MFKH analyzed data. MGKB, KED and MD contributed analytic tools.

MGKB and MFKH prepared the figures. MGKB, KED, FWO, MTL and MFKH wrote the

manuscript. All authors reviewed and revised the manuscript.

Conflict of interest:

The authors declare no competing financial interests.

Acknowledgements:

We very much thank Silvia Vieweg for assistance with surgeries, histology and optimization of

immunostainings. We further thank Janet Stallmann and Kathrin Ohl for assistance with

surgeries, histology- and immunostaining of brain slices. We would like to thank PD Dr. Eike

Budinger and Dr. Julia Henschke for their help and assistance in anatomical questions. Many

thanks to our interns for their assistance with anatomical reconstructions: Tarik Drewes,

Katharina Eick, Vincent Koop, Jan-Niklas Löbner, Julia Rieger and Ceylan Steinecke. This

project and corresponding research have been funded by the Leibniz Association WGL by the

Leibniz Postdoctoral Network, LPN (to MTL and MFKH) and, the Deutsche

Forschungsgemeinschaft DFG by the SFB779 (to MFKH and FWO) and SPP1665 (to MTL

and FWO).

Abstract

In order to reinforce relevant behavior, reinforcement-evaluating brain structures interact with

circuits involved in sensory processing. During reward-based auditory learning, the ventral

tegmental area (VTA) conveys a dopaminergic teaching signal to the primary auditory cortex.

It has been shown that dopaminergic levels within the auditory cortex are constantly increased

during initial auditory acquisition learning (Stark and Scheich, 1997). It is currently unknown,

however, how the VTA circuitry thereby influences cortical frequency information processing

and spectral integration. In this study, we therefore investigated the temporal effects of direct

VTA stimulation on sensory processing in the auditory cortex of anesthetized male Mongolian

gerbils on a synaptic circuit level by current-source-density analysis. While auditory lemniscal

input predominantly terminates in the granular input layers III/IV, we found that reward-related

modulation of spectral processing is relayed by a different circuit, namely thalamic inputs to

the infragranular layers Vb/VIa under the control of the VTA. Activation of this circuit yields

a frequency-specific gain amplification of sensory input and enhances corticocortical

information transfer, especially in supragranular layers I/II. We further verified that the gain

modulation of early thalamocortical inputs increase local intracolumnar processing, while

supragranular amplification is due to strengthened corticocortical processing. Altogether, this

modulation manifests as a long-lasting influence transcending the signaling of a mere reward-

prediction error, as previously suggested by the aforementioned behavioral experiments. For

the first time, our findings demonstrate a cellular and circuit substrate substrate for reward-

mediated influences in the auditory cortex.

Keywords:

Dopamine, auditory cortex, Mongolian gerbil, current source density, ventral tegmental area,

optogenetics

Significance Statement

Phasic release of dopamine from the ventral tegmental area has long been associated with the

reward prediction error signal postulated by reinforcement learning theory. How dopamine

released by the VTA affects the plasticity that integrates sensory information and such learning-

related signals in sensory cortex, however, is unknown.

Here, we show that VTA-based dopamine effectuated a long-lasting gain modulation of sensory

thalamocortical inputs into the infragranular layers in gerbil auditory cortex. Our findings

therefore assign a specific function to this rather understudied cortical circuit: VTA-modulated

input to deep layer neurons in the sensory cortex mediates the integration of sensory and task-

related information even when they are not precisely temporally contingent.

Introduction

The sensory cortex receives both bottom-up input relaying stimulus information from the

sensory epithelia and top-down input from, for example, reinforcement-evaluating brain

structures (Ohl, 2014). Among the latter, the ventral tegmental area (VTA) is a key structure

associated with the coding of reward, reward prediction, and reward prediction error

(Bromberg-Martin et al., 2010). Especially in the framework of reward prediction error coding,

projections of dopamine neurons in the VTA to the striatum and prefrontal cortex have been

investigated in great detail (Schultz, 2015). In contrast, the anatomy of projections from the

VTA towards sensory cortices (Dinopoulos and Parnavelas, 1991; Budinger et al., 2008;

Gardner et al., 2018), and consequently its direct impact on cortical processing, has remained

rather elusive.

Dopamine released in the sensory cortex may complement bottom-up stimulus processing with

a behaviorally relevant representation of stimulus value and salience to support adaptive

behavior (Ohl and Scheich, 2005; Bromberg-Martin et al., 2010; Happel, 2016). Consistently,

for the case of the auditory cortex (ACx), intracortical dopamine levels (Stark and Scheich,

1997) and their experimental manipulation (Schicknick et al., 2008, 2012) were shown to affect

behavioral measures in auditory learning scenarios. Thus, dopamine appears to be involved in

several general behaviorally relevant functions, including auditory perceptual decision making,

prediction, and learning, which are increasingly realized to be supported by the ACx (King et

al., 2018). In agreement, we have previously shown that the pharmacological stimulation of

D1/D5 receptors influences sensory processing at the level of both local and wide-spread

circuits in auditory cortex (Happel et al., 2014).

In order to determine the contribution of VTA projections to these cortical effects of dopamine,

we optogenetically stimulated the projection neurons of the VTA in adult male Mongolian

gerbils (*Meriones unguiculatus*) and measured the layer-specific processing in the auditory

cortex by tone-evoked current-source density (CSD) analysis. We demonstrate that VTA

stimulation effectuated a sensory gain amplification via thalamocortical inputs in the deep

layers Vb/VIa, rather than via recurrent excitation in layer III/IV (Liu et al., 2007; Happel et

al., 2010). Our results demonstrate for the first time a functional diversification of the

anatomically distinct thalamocortical input systems in the sensory cortex. Reward-modulated

sensory input in deep layer neurons therefore might provide a cellular substrate for integrating

sensory and task-related information in the service of sensory-based decision-making and

reinforcement learning.

Materials and Methods

All experiments were carried out in adult male Mongolian gerbils (*Meriones unguiculatus*, age 4-8 months, body weight: 70-100 g; n= 26). The C1V1 group (n=12) consisted of animals that were expressing the viral construct, allowing for optogenetic stimulation of the VTA whilst recording. The control group (n=7) consisted of non-transfected animals to control for temporal stability of the signal. In order to investigate effects due to laser-stimulation itself, an opsin-less YFP group (n=7) was tested. Experiments were conducted in accordance with ethical animal research standards defined by the German Law and approved by an ethics committee of the State of Saxony-Anhalt.

Viral transfection of the VTA

Gerbils were anesthetized by intraperitoneal injection (0.06-0.08 ml/h) of 45% v/v ketamine (50 mg/ml, Ratiopharm GmbH), 5% v/v xylazine (Rompun, 2%, Bayer Vital GmbH) and 50% v/v of isotonic sodium-chloride solution (154 mmol/l, B. Braun AG). Animals were fixed in an automated stereotaxic injector (Neurostar) for virus transduction (700 nl; 100 nl/min; 5 min resting; C1V1: AAV5-CamKIIα-C1V1(E162T)-p2A-eYFP or YFP: AAV2-CamKIIα-eYFP 3E12 particles/ml, UNC Vector Core; target side: AP: -3.8-4.0; ML: 0.5; DV: -6.2 mm, bregma for reference) and fiber implantation with a planar setting of bregma and lambda. Custom made fiber implants (fiber implant length 8 mm, Ø 230 µm, NA 0.39) were implanted at target side (AP: - 3.8-4.0; ML: 0.5; DV: -5.8 mm). Fiber positions were determined by histological analysis and compared with the gerbil brain atlas (Fig. 1 D and E) (Radtke-Schuller et al., 2016). Animals were kept for 3 weeks to allow recovery and enough protein expression.

Intracranial self-stimulation using optogenetic VTA activation

Successful activation of the VTA through optogenetic stimulation was confirmed in an intracranial self-stimulation paradigm. Animals were trained for 10 consecutive days (at 20 min per day). By activating a lever with their nose, they could elicit brief laser stimulation (10 pulses, 25 Hz, 473 nm, 10 mW) delivered to the VTA through the implanted fiber (Fig. 1 A, B and C). Only animals pressing the lever more than 50 presses/minute were kept for the C1V1 group (Fig. 1 G) (Helbing et al., 2016; Brocka et al., 2018; Lippert et al., 2018).

Electrophysiological recordings

Surgical procedure for electrophysiological recordings have been described in detail elsewhere (Deliano et al., 2018). Briefly, animals have been anesthetized by intraperitoneal injection (0.06-0.08 ml/h) of 45% v/v ketamine (50 mg/ml, Ratiopharm GmbH), 5% v/v xylazine (Rompun, 2%, Bayer Vital GmbH) and 50% v/v of isotonic sodium-chloride solution (154 mmol/l, B. Braun AG). The right auditory cortex was exposed by trepanation and target location of AI was chosen based on vascular landmarks. Status of anesthesia was checked via paw withdrawal-reflex on a regular basis (10-20 min). Body temperature was kept stable at 34°C. Due to an epileptic seizure during the electrophysiological recordings we had to omit an animal from the YFP group.

Experiments have been carried out in a Faraday-shielded acoustic soundproof chamber. The animals head was fixed 1m away from the speaker (Tannoy arena satellite KI-8710-32) by screwing the head post to a custom-made head holder. Local field potentials (LFPs) have been recorded with a linear 32-channel shaft electrode (NeuroNexus A1x32-50-413), which was implanted perpendicular to the surface area of AI (Happel et al., 2010). LFPs have been pre-amplified 500-fold and band-pass filtered (0.7-300 Hz) with a PBX2 preamplifier (Plexon Inc.).

Data were digitized at a sampling frequency of 1 kHz with a multichannel recording system

(Multichannel Acquisition Processor, Plexon Inc.).

Pure tones spanning 8 octaves (frequency range between 125 Hz-32 kHz, tone duration: 200

ms, ISI: 0.6-0.8, 50 pseudorandomized repetitions, 65 dB SPL, 7.5 min per measurement) were

generated in MATLAB, converted into an analog signal by a data acquisition card (NI PCI-

BNC2110, National Instruments, Germany), rooted through a controllable attenuator (gPAH,

Guger, Technologies, Austria) and amplified by an audio amplifier (Thomas Tech Amp75). A

measurement microphone and conditioning amplifier were used to calibrate acoustic stimuli

(G.R.A.S. 26AM and B&K Nexus 2690-A, Bruel & Kjaer, Germany).

Electrophysiological measurements were performed according to the scheme of Fig. 2 A. After

implantation, laminar LFPs were measured for 45-75 min until evoked CSD profiles had been

stabilized (Pre-measurements). The last three recordings before laser treatment were taken as

the referencing Pre-measurements. For the C1V1and YFP groups, light fibers were additionally

connected to a laser for VTA stimulation. Laser stimulation (10 mW at the optical fiber of the

setup) was then coupled and synchronized with tone presentation for one measurement (25 Hz,

473 nm, 10 mW; 400 repetitions in total). This measurement was taken as the laser

measurement followed by a series of post-measurements (>60 min). After the experiment,

gerbils were perfused with 4% PFA as described previously (Saldeitis et al., 2014) and brains

were removed for anatomical and histological validation.

Current source density analysis

Based on the tone-evoked LFPs for each presented stimulus we calculated the second spatial

derivative, indicating the laminar evoked current-source density (CSD) distribution in AI

(Mitzdorf, 1985):

(1) $-CSD \approx \frac{\delta^2 \emptyset(z)}{\delta z^2} = \frac{\emptyset(z+n\Delta z) - 2\emptyset(z) + \emptyset(z-n\Delta z)}{(n\Delta z)^2}$

The formula reflects the relation of the field potential ($\emptyset$), the spatial coordinate of the cortical

laminae (z), the inter-channel distance (Δz, 50 µm) and the differentiation grid (n). Prior to

CSD calculation, LFPs were smoothed using a weighted average of 9 channels (Hamming

window, spatial filter kernel size of 400 µm; linear extrapolation of 3 channels at boundaries)

(Happel et al., 2010).

CSD profiles revealed patterns of current influx (sinks) and efflux (sources). Based on CSD

profiles of the Pre-measurements, cortical layers were assigned to the respective channels of

sink components. Early sink components were used as indicators of layers III/IV, Va, and

Vb/VIa, whereas later sink activities were assigned to layers I/II, and VIb (Fig. 2; Fig. 2-1).

CSD data was also transformed by rectifying and averaging the waveforms of each single

channel (n). This average rectified CSD (AVREC) provides a temporal waveform of the overall

local columnar transmembrane current flow (Givre et al., 1994; Schroeder et al., 1998).

(2) $AVREC = \frac{\sum_{i=1}^{n} |CSD_i|(t)}{n}$

We further quantified the residue of the CSD defined as the sum of the non-rectified magnitudes

divided by the rectified magnitudes for each channel. Thereby, the ResidualCSD quantifies the

balance of the transmembrane charge transfer along the recording axis (Harding, 1992) and

gives rise to the lateral corticocortical contribution to stimulus related activity (Happel et al.,

2010).

(3) $ResidualCSD = \frac{\sum_{i=1}^{n} CSD_i(t)}{n}$

In order to quantify the early (0-50 ms) and late (80-300 ms) contributions of the AVREC and

ResidualCSD, root mean square (RMS) amplitudes were calculated. Based on the previously

described layer indexing, averaged layer-dependent CSDs were calculated, which were used

for the semi-automatic sink detection (Fig. 2-1). Briefly, before detection of significant tone-

evoked activity, sources were omitted before calculating the mean of CSD traces within a

cortical layer. Then, the 1.5-fold standard deviation (SD) of the baseline (-200-0 ms before tone

presentation onset for all channels and stimuli) was used to calculate the intercepts of the

smoothed (10 ms gaussian sliding window) mean CSD trace. For early tone-evoked sink

activity in cortical layers III/IV, Va, and Vb/VIa, we used an onset of the first intercept between

0-50 ms after tone onset. For subsequent tone-evoked sink components in layers I/II and VIb

onsets were only taken >50 ms after tone-onset. Offsets of each sink were defined as the

crossings below the 1.5 SD baseline and for each cortical layer. Peak amplitudes of layer-

dependent CSD-traces were determined within time windows of on- and offsets from the

average of the corresponding CSD channels. For calculating the RMS value of AVREC and

ResidualCSD we used fixed time windows of 0-50 ms after tone-onset for early and 80-300 ms

for late activity.

Data sorting and analysis

Tuning curves of layer-specific CSD-traces, AVREC, and ResidualCSD were averaged

according to the best-frequency of each animal separately to plot an averaged group tuning

curve. For illustration, all non-BF frequencies above and below the BF were binned pairwise,

resulting into the grouping of near-BFs (± 1-2 octaves) and non-BFs (± 3-4 octaves) in 5

frequency bins [- non-BF, - near-BF, BF, + near-BF, + non-BF]. For further analysis of BF-,

near- and non-BF evoked responses, we observed data from time windows before, during, and

after laser stimulation. Evoked responses were normalized within each animal using the

corresponding mean values of the last three Pre-measurements before laser-stimulation (Fig. 2

A.

For statistical comparison, parameters of interest were analyzed on a single-trial level using

linear-mixed effect (LME) models (R Studio; R 3.5.1) (Chang et al., 2018) binned across

measurements as follows: pre 1-3 (-22.5 - -7.5 min), laser (0 min), post 1-3 (7.5-22.5 min), post

4-5 (30-37.5 min), post 6-7 (45-52.5 min). Peak amplitudes of single-trial data were determined

within the time windows from on- and offsets revealed on averaged CSD traces. LME models

were calculated and fitted using normalized data with the lmerTest-package (3.0-1) and a Wald-

chi-square-test was performed using the car-package (3.0-2). Animals were used as "by-

subject" random effects, to account for the repeated measurements during and across time

points whereas measurements and groups were used as fixed effects to explain physiological

data. As significance criterion for effects on single-trial level, the normalized signal needed to

i) exceeded a $\pm$ 10% criterion compared to Pre-measurement (based on the SEM of the average

jitter before laser treatment), and ii) show significant difference from control group data

revealed by significant fixed effects of the LME models. For slope estimations and comparison

between measurements, we used the R-package Ismeans (version 2.27-62) to calculate a least

square linear fit for non-, near- and BF data bins and check significant changes in tuning slope

properties between measurements.

Anatomical estimation of fiber position

PFA fixated brains of the C1V1 group (n =7) were cut in 50 µm thick slices around the region

of the VTA. Every second slice was used for Nissl staining to anatomically evaluate the fiber

position, whereas the remaining slices were used for immunostaining to determine the distance

from fiber tip to areas of TH- and YFP-co-fluorescence with FIJI ImageJ (1.51g). Pictures were

taken with a Leica microscope (Axioscop 2, Leica GmbH).

Immunostaining was performed as previously described (Brocka et al., 2018) with a primary

antibody against tyrosine hydroxylase (1:1000 rabbit anti-TH, Milipore) and a secondary Alexa

Fluor 546 goat-anti-rabbit (1:400 anti-rabbit, Molecular probes) using an antibody diluent (Zytomed Systems). Position of the fiber was estimated using the recently published gerbil brain atlas for reference (Radtke-Schuller et al., 2016).

Results

Optogenetic stimulation of the VTA evokes self-stimulation behavior

To selectively target the excitatory projection neurons of the VTA, we expressed the red-shifted opsin C1V1 under the CamKIIα-promotor (Guo et al., 2014) by a stereotactically guided microinjection (Fig. 1 A; 700 nl; AAV-CamKIIα-C1V1(E162T)-p2A-eYFP, 3e12 particles per ml, UNC Vector Core). Opsin expression in the VTA overlapped significantly with tyrosine hydroxylase (TH)-immunostaining, selective for dopaminergic cell populations. The connectivity of the transduced neurons with the auditory cortex was confirmed by histology (Fig. 1 A,B) in agreement with previous reports (Budinger et al., 2008; Budinger and Scheich, 2009). After viral transduction, we waited 2-3 weeks for sufficient transgene expression before subjects were trained over the course of 10 days in an optogenetic intracranial self-stimulation task (Fig. 1 C; Movie 1-1; 10 pulses, 25 Hz, 473 nm laser-light per nose-poke; 10 mW power

at fiber implant) (Lippert et al., 2018). Animals that showed more than 50 presses/minute were

chosen as the C1V1 group for electrophysiological experiments (Fig. 1 G) (Helbing et al.,

2016; Brocka et al., 2018). Anatomical validation of the fiber position relative to Bregma (AP:-

3.57 mm, ML: 584.14 µm; cf. Fig. 1 D and E), as well as the fiber distance towards the area of

opsin expression (Fig. 1 F, 146.8 ± 38.9 µm), was performed in a subset of animals (n=7) in the

C1V1 group (see Materials and Methods). In contrast to the opsin group, animals of the YFP-

control group showed no self-stimulation behavior (Fig. 1 G).

Effect of VTA-activation on tone-evoked columnar activity in the auditory cortex

After evaluation of self-stimulation behavior in the C1V1 and YFP group, we performed acute

laminar local field potential electrophysiological recordings from their auditory cortices under

ketamine-xylazine anesthesia. This approach allows the investigation of circuitry effects by the

stimulation of the VTA on sensory cortical processing. Additionally, we also recorded from a

control group of naïve animals, which underwent no implantation or laser stimulation, to

determine which effects were due to opsin activation or laser light. Pure tones were presented

in iso-octave steps in order to cover the main tonotopic range of the gerbil (Stimulation

frequencies ranged from 0.125 kHz – 32 kHz, duration: 200 ms, ISI 0.6-0.8s, 50 rep., 65 dB

SPL). After insertion of the recording electrode (NeuroNexus A1x32-50-413), sets of 50

repetitions per stimulus (~7.5 minutes) were recorded over 45-75 min to guarantee stable

recordings (Fig. 2 A; see Materials and Methods). We then paired the presentation of each pure

tone with laser-stimulation of the VTA (473 nm; 25 Hz; 10 ms pulse length, 200 ms duration).

Thereby, the VTA was stimulated 400 times within 7.5 minutes of the combined laser/tone

presentation (Fig. 2). After this paired measurement, consecutive tone-only measurements

(post) were performed over the course of 60 minutes. This procedure allowed us to observe

phasic and long-lasting effects of optogenetic VTA stimulation on auditory cortex tone-evoked

activity.

Fig. 2 B displays a representative example of pure-tone evoked CSD-profiles after stimulation

with the best frequency (BF, top) and an off-BF (bottom, 2 octaves apart) before (left), during

(center) and 30 minutes after (right) VTA- stimulation in a C1V1-transduced animal. Before

VTA stimulation, tone-evoked CSD profiles show the canonical feed-forward pattern of early

current sink activity in layers III/IV and Vb/VIa (Fig. 2 B, left) (Szymanski et al., 2009; Happel

et al., 2010; Happel and Ohl, 2017). Subsequent later supra- and infragranular sink activities

can be related to intracortical synaptic circuits in layers I/II, Va and VIb (see semi-automatic

detection of sink activity in Materials and Methods and Fig. 2 B and Fig. 2-1). Responses to

frequencies apart from the BF generally evoke less prominent tone-evoked activity across all

cortical layers. When pure tone stimulation is paired with VTA stimulation, we observed

prominent changes of amplitudes, but not spatiotemporal flow of synaptic activity (Fig. 2 B,

center). This increase of tone-evoked cortical processing was long-lasting and persisted over

approximately 30 min after the cessation of laser stimulation (Fig. 2 B, right).

Laminar specificity of transient and long-lasting effects of VTA stimulation

In order to relate the observed changes of in current flow to distinct cortical layers, we

separately analyzed CSD-traces from respective cortical layers (I/II, III/IV, Va, Vb/VIa and

VIb; see Materials and Methods). Prominent sink components of cortical layers III/IV, Va, and

Vb/VIa were related to early time windows of 0-50 ms after tone onset, while cortical layers

I/II and VIb felt into later time windows of >50 ms determined by sink onset (cf. Fig. 2-1).

Described effects during and after VTA stimulation might impact on sensory processing in a

less stimulus-locked manner and influence trial-by-trial signal-to-noise ratio (Kroener et al.,

2009). In order to capture such variability, we have utilized a single-trial analysis using LME

models (Chang et al., 2018) on peak amplitudes of each layer as a function of time before and

after VTA stimulation separately for BF, near- ($\pm$ 1-2 octaves) and non-BF ($\pm$ 3-4 octaves)

stimulation for all groups (Fig. 3). Within each frequency bin, data was normalized to pre-

measurements in order to characterize relative changes over time and compared both groups

with laser treatment (C1V1, YFP) against the control group. Within the control group, we

observed a variation of tone-evoked responses of up to $\pm$ 10% over the entire recording time.

Therefore, to obtain main effects of the laser treatment and account for general variance, we

quantified and statistically tested only data that crossed this $\pm$ 10% criterion of the normalized

pre-measurements (see dashed lines in Fig. 3). In the C1V1 group, VTA stimulation led to a

significant increase of peak amplitudes in infragranular layers Vb/VIa most prominently after

BF-stimulation for up to 45 minutes. The main current sink in thalamocortical input layers

III/IV, in contrast, showed only minor modulation after VTA stimulation. Only 30 minutes after

VTA stimulation off-BF evoked activity was increased. Activity in layer Va was unaffected by

VTA stimulation. Late onset sink components in layers I/II and VIb showed opposing temporal

effects after VTA stimulation. While supragranular activity increased gradually and long-

lasting mainly for BF/near-BF stimulation, infragranular layer VIb activity was most strongly

amplified for non-BF only during combined tone-laser stimulation. In the YFP-group peak

amplitudes of all sink components showed a general trend to be reduced during laser-

stimulation and recovered over time within the >45min of post recordings. These effects were

potentially due to light-related heating of the stimulation site (Stujenske et al., 2015; Arias-Gil

et al., 2016), which might shift the balance of excitation and inhibition in the VTA (Johnson

and North, 1992; Paladini and Tepper, 1999; Tan et al., 2012; Van Zessen et al., 2012). Data

from control animals displayed a mainly stable amplitude range within the $\pm$ 10% range of pre-

measurements.

Based on the most prominent changes of evoked activity in the C1V1-group during VTA

stimulation (layer VIb) and over the longer period of >30 min after stimulation (layers I/II and

Vb/VIa), we compared slopes of evoked peak amplitudes as a function of stimulation frequency

(Fig. 4). All data were normalized to BF-evoked amplitudes of pre-measurements within each

layer separately. Slopes across the BF/near-BF/non-BF range therefore allow us to characterize

changes in tuning sharpness. Frequency-independent gain increase, in contrast, would not affect

the slope, but rather affect the intercept. In the pre-measurement, the most prominent frequency

tuning was observed in thalamocortical input layers III/IV. In the pre-condition amplitudes in

the non-BF bin were reduced about 60% compared to BF-evoked responses. Across frequency

bins, this led to a slope of 0.28 indicating a significant dependence of amplitude and stimulation

frequency (Fig. 4, blue line, $p < 0.001$). During laser stimulation and 30 minutes after, slopes did

not change (change of slope > 0.01; $p > 0.05$) indicative of a stable frequency tuning in layers

III/IV. In layers Vb/VIa, a slope of 0.13 indicated a less prominent frequency tuning in the pre-

condition. During VTA stimulation the frequency tuning was unaffected (change of slope

< 0.01; $p < 0.05$) but showed a significant increase of sharpness 30 minutes later (change of slope

pre:30 min of +0.05; $p < 0.001$). Comparably, in supragranular layers, I/II slopes of the pre-

condition (0.09; $p < 0.001$) did not significantly increase during VTA stimulation (change of

slope +0.04; $p = 0.09$), but significantly increased >30 min later (change of slope pre:30 min

+0.08, $p < 0.001$). Hence, cortical layers Vb/VIa and I/II showed no change of tuning bandwidth

during the paired tone/VTA stimulation, but a significant increase in spectral integration

prolonged for up to 30 minutes afterwards. Activity in layer VIb showed a prominent increase

during VTA stimulation selectively for the off-BF bins (cf. Fig. 3; bottom right), and, in

accordance, we found slopes to be significantly decreased during laser stimulation (change of

slope pre:laser –0.06; $p < 0.001$), which recovered after 30 minutes (change of slope Pre:30 min

–0.02, $p = 0.10$). Layer Va similarly showed a decrease in slope during laser-stimulation (change

of slope pre:laser –0.04; p<0.001), and also recovered 30 minutes later (p>0.05). VTA

stimulation yielded a transient increase in response bandwidth in cortical layers Va and VIb.

Based on the layer-specific modulation of early tone-evoked inputs in thalamocortical-recipient

layers Vb/VIa and subsequent synaptic inputs in supragranular layers, we speculate that

stimulation of the VTA affects early and late sensory processing on a local and corticocortical

level, respectively (cf. Happel et al., 2014).

Impact on local intracolumnar and corticocortical synaptic inputs

In a next step, we therefore aimed to disentangle early activity associated with local

thalamocortical input, and subsequent activity associated with corticocortical processing. We

compared changes in the tone-evoked average rectified CSD (AVREC) and the residuals of the

CSD (ResidualCSD) during early (0-50 ms) and late (80-300 ms) time windows (Fig. 5 A).

This rationale is based on previous findings demonstrating that the residual CSD reflects the

spatiotemporal ratio of unbalanced sinks and sources providing a quantitative measure of

horizontal intercolumnar inputs to a given cortical site (Happel et al., 2010; Happel and Ohl,

2017). The AVREC, on the other hand, reflects the overall intracolumnar current flow at a given

recording site (Givre et al., 1994; Schroeder et al., 1998). Fig. 5 A (left) shows changes of both

the grand average of the AVREC (top) and relative CSD (bottom) waveforms before, during,

and, in certain time windows, after VTA stimulation in the C1V1 group. Comparable changes

were not observed in neither the YFP nor control group (Fig. 3-1). To further investigate

temporal changes of BF-evoked responses in more detail, time course of binned root mean

square (RMS) values for early and late averaged and normalized AVREC and ResidualCSD

windows of each group were plotted as a function of time (normalized to pre-measurements).

RMS values of the AVREC during the early phase (0-50 ms) displayed a significant increase

in the C1V1 group from 15 min until 52 min after VTA stimulation. Significant effects were

validated with a LME model (Fig. 5 B; p<0.001-0.025). During the late phase (80-300 ms) after

tone onset, RMS of the averaged AVREC did not show any significant changes between each

group (Fig. 5 B). During laser stimulation, a significant increase has only been observed for the

averaged early ResidualCSD indicative of amplified lateral contributions to early tone-evoked

columnar activity in the C1V1 group (p=0.04). Furthermore, early and late ResidualCSD were

both significantly enhanced during later time points after the VTA stimulation (early

ResidualCSD: 22.5-37.5 min; p=0.014-0.039; late ResidualCSD: 37.5-45 min, p=0.008-0.033).

Averaged RMS tuning curves of the AVREC and ResidualCSD of the C1V1 group revealed an

increase of tone-evoked activity during and after laser stimulation across the entire range of BF,

near-BF (± 1-2 octaves) and non-BF (± 3-4 octaves) stimulation (Fig. 5 C).

Integration of local and corticocortical synaptic inputs in dependence of the stimulation

frequency

All single-trial AVREC and ResidualCSD RMS values were normalized to pre measurements.

Normalized values were analyzed as a function of time before and after VTA stimulation binned

for BF, near-, and non-BF stimulation for all groups (Fig.6). Fig. 6 A (top) shows a gradual

increase of the early AVREC compared to the control- and YFP-group independent of

stimulation frequency. For BF- and near-BF-stimulation, this yielded a significant increase at

>30 min after VTA stimulation. The ResidualCSD also showed a strong increase almost

independent of the stimulation frequency already during the combined tone-laser presentation

and all following post measurements. Consistently with the layer-specific activity, there was a

less pronounced decrease of the near/non-BF-evoked AVREC in early and late time windows

in the YFP-group during laser stimulation, which started to recover over time (Fig. 6 A,

bottom).

In the later time window of 80-300 ms after tone-onset, only the ResidualCSD showed a significant increase during and up to 30 minutes after VTA stimulation (Fig. 6 B). Altogether, VTA stimulation led to a gradual increase of the overall early current flow most pronounced for BF- and near-BF stimulation. This effect was long-lasting and peaked around 30 minutes after stimulation. We further found an immediate and long-lasting increase of imbalance between sinks and sources based on the early and late ResidualCSD in all frequency bins indicative of amplified corticocortical activity.

Discussion

The VTA is a key structure to convey information about stimulus salience and valence to target areas distributed throughout the brain (Bromberg-Martin et al., 2010). Receiving both sensory as well as VTA inputs, the primary sensory cortex is ideally positioned to integrate such bottom-up and top-down information to shape behavior through reinforcement learning. We here

demonstrate for the first time that activation of VTA-reward circuits amplifies sensory input

via a yet largely overlooked thalamocortical projection targeting infragranular layers Vb/VIa of

the auditory cortex (Szymanski et al., 2009; Intskirveli et al., 2016). In contrast, granular layer

III/IV input, a well-known substrate for sensory gain modulation (Bruno and Sakmann, 2006;

Liu et al., 2007; Happel et al., 2010; Intskirveli et al., 2016), was unaffected by VTA

stimulation. The impact of the observed infragranular gain modulation on supragranular layers

manifests as a translaminar long-lasting enhancement of frequency tuning (see Fig. 7).

Furthermore, we could demonstrate with a residual CSD analysis, that the frequency-specific

gain in deeper layers emerges on a local columnar level and yields an increased corticocortical

integration displayed in lateral input most likely terminating in upper layers. Thereby, the VTA

impact on the sensory cortex transcends the time scales commonly associated with reward-

prediction error coding. This highlights the diverse, and yet elusive functions of cortical layers

in the sensory cortex (Adesnik and Naka, 2018).

VTA-activation enhances infragranular thalamocortical inputs in the auditory cortex

Optogenetic stimulation of VTA-projection neurons allowed us to investigate effects of reward-

related cross-regional circuits (Fig. 1) (Brocka et al., 2018). Roughly 8% of non-sensory inputs

of gerbil ACx arise from direct projections of the ipsilateral VTA, which terminate across the

entire cortical column (Budinger et al., 2008). Therefore, direct stimulation of the VTA

projection neurons is the most effective method to activate this circuit. In contrast, the success

of optogenetic approaches to directly stimulate cortical terminals have been challenged in recent

studies and particularly do not allow for an homogenous activation across all cortical layers

(Yizhar et al., 2011; Stujenske et al., 2015; Yona et al., 2016). VTA stimulation led to increased

early tone-evoked (0-50 ms) local columnar current flow and increased long-range

corticocortical activity. This effect persisted for up to 50 minutes after VTA stimulation and

was most prominent for the respective BF of the given recording site (Fig. 5). Hence, VTA

stimulation promotes the local gain of columnar input over a prolonged period and lead to

substantial corticocortical integration of spectral information. Notably, the gain amplification

of sensory input appeared exclusively in thalamocortical-recipient layers Vb/VIa but was absent

in the main thalamocortical-recipient granular layers III/IV (Fig. 3). This enhancement of

infragranular activity was locally relayed to supragranular layers I/II. This mechanism may

provide a way for intratelencephalic neurons in infragranular layers (Harris and Mrsic-Flogel,

2013) to provide a translaminar gain control of sensory representation depending on the

experience of reward (Castejon et al., 2016; Jarvis et al., 2018). Although direct projections of

the ipsilateral VTA provide a significant contribution to non-sensory inputs of the gerbil ACx

(Budinger et al., 2008), our approach may recruit other relay stations that could contribute to

the observed effects.

Previously, reported effects of dopamine on tuning sharpness in the ACx (Bao et al., 2001;

Deliano et al., 2018) raised the question on which specific cortical circuit motifs reward

information modulates spectral information. Our data showed that stimulating the VTA

projections only affected early auditory tuning sharpness in deep cortical layers, but not in the

main thalamocortical input layer III/IV (Sugimoto et al., 1997; Intskirveli et al., 2016). Sharper

frequency representation was relayed to superficial layers I/II and persisted over time after VTA

stimulation. Our observations are thereby consistent with recent findings from layer-specific

high-field fMRI in human auditory cortex showing sharper tuning in upper layers depending

on attention and task engagement (De Martino et al., 2015).

As a result, reward-circuit activation promotes a long-lasting gain enhancement of salient

auditory stimuli via strengthened infragranular thalamocortical inputs and increased sensory

integration in supragranular circuits. Our finding now provides an explanatory role of the

anatomically and physiologically well-described early infragranular inputs in layers Vb/VIa of

the sensory cortex (Szymanski et al., 2009; Happel et al., 2010; Constantinople and Bruno,

2013; Intskirveli et al., 2016) as a circuit for integrating sensory and task-related information.

Persistent cortical modulation might underlie dopamine-dependent influences on learning

Recent reports have highlighted that sensory-evoked responses in deep layers are modulated by

behavioral states (Pais-Vieira et al., 2015; Zhao et al., 2016) and undergo substantial learning-

related plasticity (Biane et al., 2016) not described for middle layer thalamocortical input

(Froemke et al., 2007). Our data now assigns a functional impact of the VTA on deep cortical

layers—particularly under the control of dopaminergic neuromodulation (Fig. 7).

Optogenetic VTA stimulation preferentially recruits the main output population of

dopaminergic projection neurons making up roughly 55-65% of all VTA neurons (Nair-Roberts

et al., 2008; Sesack and Grace, 2010; Liu et al., 2012). Nevertheless, it has been demonstrated

that the small population of non-dopaminergic projection neurons also convey reward

information (Gorelova et al., 2012; Wang et al., 2015; Yoo et al., 2016) and particularly

promote the activation of brain-wide regions including the auditory cortex (Brocka et al., 2018).

Furthermore, there is overlap between dopaminergic and glutamatergic cells through

transmitter co-release (Stuber et al., 2010; El Mestikawy et al., 2011; Hnasko et al., 2012).

Nevertheless, the long duration of responses modulation in our data set strongly argues for a

significant dopaminergic component, as glutamatergic effects are generally more transient

(Mylius et al., 2015). Further evidence for a dopaminergic origin of the long-lasting effects is

provided by the distribution of dopamine receptors. In the sensory cortex, they are concentrated

mainly in infragranular and supragranular layers where we see modulatory effects (Compare

Fig. 1 B) (Lewis et al., 1987; Phillipson et al., 1987; Lidow et al., 1991; Williams and Goldman-

Rakic, 1993; Campbell et al., 1996; Schicknick et al., 2008).

Thereby, the current study pursues the idea of dopamine-modulated columnar local gain of

behaviorally relevant stimuli on infragranular thalamocortical terminals via presumably

presynaptic dopamine receptors (Izhikevich, 2007; Blundon and Zakharenko, 2013; Happel et

al., 2014). Previously described accumulative effects of dopamine on auditory learning and

long-term memory consolidation (constant increase of cortical dopamine within 30 min after

training onset; cf. Stark and Scheich, 1997) could then be the result of the long-lasting action

of dopamine at the presynaptic terminal (Stark and Scheich, 1997; Schicknick et al., 2008).

Thereby, the sensory gain amplification described in this study potentially promotes a sustained

perceptual salience of incoming stimuli (Froemke et al., 2013; Kato et al., 2015; Sohoglu and

Chait, 2016). These previous observations are already implying functions of the VTA beyond

mere reward-prediction error signaling to the auditory cortex (Gardner et al., 2018). Our

findings now provide a circuit level explanation of the necessary integration of transient reward

information and persistent modulation of cortical activity. Thereby, this circuit might constitute

a substrate for credit assignment in reinforcement learning theory concerned with the temporal

delay between sensory input and its behavioral consequences (Hull, 1944).

References:

Adesnik H, Naka A (2018) Cracking the Function of Layers in the Sensory Cortex. Neuron

100:1028–1043.

Arias-Gil G, Ohl FW, Takagaki K, Lippert MT (2016) Measurement, modeling, and

prediction of temperature rise due to optogenetic brain stimulation. Neurophotonics

3:045007.

Bao S, Chan VT, Merzenich MM (2001) Cortical remodelling induced by activity of ventral

tegmental dopamine neurons. Nature 412:79–83.

Biane JS, Takashima Y, Scanziani M, Conner JM, Tuszynski MH (2016) Thalamocortical

Projections onto Behaviorally Relevant Neurons Exhibit Plasticity during Adult Motor

Learning. Neuron 89:1173–1179.

Blundon JA, Zakharenko SS (2013) Presynaptic Gating of Postsynaptic Synaptic Plasticity at

thalamocortical synapses: a plasticity filter in the adult auditory cortex. The

Neuroscientist 19:465–478.

Brocka M, Helbing C, Vincenz D, Scherf T, Montag D, Goldschmidt J, Angenstein F, Lippert

M (2018) Contributions of dopaminergic and non-dopaminergic neurons to VTA-

stimulation induced neurovascular responses in brain reward circuits. NeuroImage

177:88–97.

Bromberg-Martin ES, Matsumoto M, Hikosaka O (2010) Dopamine in motivational control:

rewarding, aversive, and alerting. Neuron 68:815–834.

Bruno RM, Sakmann B (2006) Cortex Is Driven by Weak but Synchronously Active

Thalamocortical Synapses. Science 312:1622–1627.

Budinger E, Laszcz A, Lison H, Scheich H, Ohl FW (2008) Non-sensory cortical and

subcortical connections of the primary auditory cortex in Mongolian gerbils: bottom-up

and top-down processing of neuronal information via field AI. Brain research 1220:2–32.

Budinger E, Scheich H (2009) Anatomical connections suitable for the direct processing of

neuronal information of different modalities via the rodent primary auditory cortex.

Hearing research 258:16–27.

Campbell AD, Kohl RR, McBride WJ (1996) Serotonin-3 receptor and ethanol-stimulated

somatodendritic dopamine release. Alcohol.

Castejon C, Barros-Zulaica N, Nuñez A (2016) Control of Somatosensory Cortical Processing

by Thalamic Posterior Medial Nucleus: A New Role of Thalamus in Cortical Function.

PloS one 11:e0148169.

Chang A, Bosnyak DJ, Trainor LJ (2018) Beta oscillatory power modulation reflects the

predictability of pitch change. Cortex 106:248–260.

Constantinople CM, Bruno RM (2013) Deep cortical layers are activated directly by

thalamus. Science (New York, NY) 340:1591–1594.

De Martino F, Moerel M, Ugurbil K, Goebel R, Yacoub E, Formisano E (2015) Frequency

preference and attention effects across cortical depths in the human primary auditory

cortex. Proceedings of the National Academy of Sciences of the United States of

America 112:16036–16041.

Deliano M, Brunk MGK, El-Tabbal M, Zempeltzi MM, Happel MFK, Ohl FW (2018)

Dopaminergic neuromodulation of high gamma stimulus phase-locking in gerbil primary

auditory cortex mediated by D1/D5-receptors. European Journal of Neuroscience.

Dinopoulos A, Parnavelas JG (1991) The development of ventral tegmental area (VTA)

projections to the visual cortex of the rat. Neuroscience letters 134:12–16.

El Mestikawy S, Wallén-Mackenzie Å, Fortin GM, Descarries L, Trudeau L-E (2011) From

glutamate co-release to vesicular synergy: vesicular glutamate transporters. Nature

Reviews Neuroscience 12:204–216.

Froemke RC, Carcea I, Barker AJ, Yuan K, Seybold BA, Martins ARO, Zaika N, Bernstein

H, Wachs M, Levis PA, Polley DB, Merzenich MM, Schreiner CE (2013) Long-term

modification of cortical synapses improves sensory perception. Nature Neuroscience

16:79–88.

Froemke RC, Merzenich MM, Schreiner CE (2007) A synaptic memory trace for cortical

receptive field plasticity. Nature 450:425–429.

Gardner MPH, Schoenbaum G, Gershman SJ (2018) Rethinking dopamine as generalized

prediction error. Proceedings of the Royal Society B: Biological Sciences 285:20181645.

Givre SJJ, Schroeder CEE, Arezzo JCC (1994) Contribution of extrastriate area V4 to the

surface-recorded flash VEP in the awake macaque. Vision Research 34:415–428.

Gorelova N, Mulholland PJ, Chandler LJ, Seamans JK (2012) The glutamatergic component

of the mesocortical pathway emanating from different subregions of the ventral

midbrain. Cerebral cortex (New York, NY : 1991) 22:327–336.

Guo S, Chen S, Zhang Q, Wang Y, Xu K, Zheng X (2014) Optogenetic activation of the

excitatory neurons expressing CaMKIIα in the ventral tegmental area upregulates the

locomotor activity of free behaving rats. BioMed research international 2014:687469.

Happel MFK (2016) Dopaminergic impact on local and global cortical circuit processing

during learning. Behavioural Brain Research 299:32–41.

Happel MFK, Deliano M, Handschuh J, Ohl FW (2014) Dopamine-modulated recurrent

corticoefferent feedback in primary sensory cortex promotes detection of behaviorally

relevant stimuli. The Journal of neuroscience : the official journal of the Society for

Neuroscience 34:1234–1247.

Happel MFK, Jeschke M, Ohl FW (2010) Spectral Integration in Primary Auditory Cortex

Attributable to Temporally Precise Convergence of Thalamocortical and Intracortical

Input. Journal of Neuroscience 30:11114–11127.

Happel MFK, Ohl FW (2017) Compensating Level-Dependent Frequency Representation in

Auditory Cortex by Synaptic Integration of Corticocortical Input Malmierca MS, ed.

PLOS ONE 12:e0169461.

Harding GW (1992) The currents that flow in the somatosensory cortex during the direct

cortical response. Experimental Brain Research 90:29–39.

Harris KD, Mrsic-Flogel TD (2013) Cortical connectivity and sensory coding. Nature 503:51–
58.

Helbing C, Brocka M, Scherf T, Lippert MT, Angenstein F (2016) The role of the mesolimbic
dopamine system in the formation of blood-oxygen-level dependent responses in the
medial prefrontal/anterior cingulate cortex during high-frequency stimulation of the rat
perforant pathway. Journal of Cerebral Blood Flow & Metabolism 36:2177–2193.

Hnasko TS, Hjelmstad GO, Fields HL, Edwards RH (2012) Ventral Tegmental Area
Glutamate Neurons: Electrophysiological Properties and Projections. Journal of
Neuroscience 32:15076–15085.

Hull CL (1944) Principles of Behavior: An Introduction to Behavior Theory.

Intskirveli I, Joshi A, Vizcarra-Chacón BJ, Metherate R (2016) Spectral breadth and laminar
distribution of thalamocortical inputs to A1. Journal of Neurophysiology 115:2083–
2094.

Izhikevich EM (2007) Solving the distal reward problem through linkage of STDP and
dopamine signaling. Cerebral Cortex.

Jarvis S, Nikolic K, Schultz SR (2018) Neuronal gain modulability is determined by dendritic
morphology: A computational optogenetic study Gutkin BS, ed. PLOS Computational
Biology 14:e1006027.

Johnson SW, North RA (1992) Two types of neurone in the rat ventral tegmental area and
their synaptic inputs. The Journal of Physiology 450:455–468.

Kato HK, Gillet SN, Isaacson JS (2015) Flexible Sensory Representations in Auditory Cortex
Driven by Behavioral Relevance. Neuron 88:1027–1039.

King AJ, Teki S, Willmore BDBB (2018) Recent advances in understanding the auditory

cortex. F1000Research 7:1555.

Kroener S, Chandler LJ, Phillips PEM, Seamans JK (2009) Dopamine Modulates Persistent

Synaptic Activity and Enhances the Signal-to-Noise Ratio in the Prefrontal Cortex

Linden R, ed. PLoS ONE 4:e6507.

Lewis DA, Campbell MJ, Foote SL, Goldstein M, Morrison ' JH (1987) The Distribution of

Tyrosine Hydroxylase-lmmunoreactive Fibers in Primate Neocortex Is Widespread but

Regionally Specific. The Journal of Neuroscience 7:279–290.

Lidow MS, Goldman-Rakic PS, Gallager DW, Rakic P (1991) Distribution of dopaminergic

receptors in the primate cerebral cortex: Quantitative autoradiographic analysis using

[3H]raclopride, [3H]spiperone and [3H]SCH23390. Neuroscience 40:657–671.

Lippert MT, Takagaki K, Weidner T, Brocka M, Tegtmeier J, Ohl FW (2018) Optogenetic

Intracranial Self-Stimulation as a Method to Study the Plasticity-Inducing Effects of

Dopamine. In: Handbook of Behavioral Neuroscience, pp 311–326.

Liu BH, Wu GK, Arbuckle R, Tao HW, Zhang LI (2007) Defining cortical frequency tuning

with recurrent excitatory circuitry. Nature Neuroscience 10:1594–1600.

Liu CL, Gao M, Jin GZ, Zhen X (2012) GABA Neurons in the Ventral Tegmental Area

Responding to Peripheral Sensory Input. PLoS ONE 7:1–8.

Mitzdorf U (1985) Current source-density method and application in cat cerebral cortex:

investigation of evoked potentials and EEG phenomena. Physiological reviews 65:37–

100.

Mylius J, Happel MFK, Gorkin AG, Huang Y, Scheich H, Brosch M (2015) Fast transmission

from the dopaminergic ventral midbrain to the sensory cortex of awake primates. Brain

Structure and Function.

Nair-Roberts RG, Chatelain-Badie SD, Benson E, White-Cooper H, Bolam JP, Ungless MA

(2008) Stereological estimates of dopaminergic, GABAergic and glutamatergic neurons

in the ventral tegmental area, substantia nigra and retrorubral field in the rat.

Neuroscience 152:1024–1031.

Ohl FW (2014) Role of cortical neurodynamics for understanding the neural basis of

motivated behavior - lessons from auditory category learning. Current Opinion in

Neurobiology 31:88–94.

Ohl FW, Scheich H (2005) Learning-induced plasticity in animal and human auditory cortex.

Current Opinion in Neurobiology 15:470–477.

Pais-Vieira M, Chiuffa G, Lebedev M, Yadav A, Nicolelis MAL (2015) Building an organic

computing device with multiple interconnected brains. Scientific Reports 5:11869.

Paladini CA, Tepper JM (1999) GABAA and GABAB antagonists differentially affect the

firing pattern of substantia nigra dopaminergic neurons in vivo. Synapse 32:165–176.

Phillipson OT, Kilpatrick IC, Jones MW (1987) Dopaminergic innervation of the primary

visual cortex in the rat, and some correlations with human cortex. Brain Research

Bulletin 18:621–633.

Radtke-Schuller S, Schuller G, Angenstein F, Grosser OS, Goldschmidt J, Budinger E (2016)

Brain atlas of the Mongolian gerbil (Meriones unguiculatus) in CT/MRI-aided

stereotaxic coordinates. Brain structure & function:1–272.

Saldeitis K, Happel MFK, Ohl FW, Scheich H, Budinger E (2014) Anatomy of the auditory

thalamocortical system in the mongolian gerbil: Nuclear origins and cortical field-, layer-

, and frequency-specificities. Journal of Comparative Neurology 522:2397–2430.

Schicknick H, Reichenbach N, Smalla K-H, Scheich H, Gundelfinger ED, Tischmeyer W

(2012) Dopamine modulates memory consolidation of discrimination learning in the

auditory cortex. European Journal of Neuroscience 35:763–774.

Schicknick H, Schott BH, Budinger E, Smalla KH, Riedel A, Seidenbecher CI, Scheich H,

Gundelfinger ED, Tischmeyer W (2008) Dopaminergic modulation of auditory cortex-

dependent memory consolidation through mTOR. Cerebral Cortex 18:2646–2658.

Schroeder CE, Metha a D, Givre SJ (1998) A spatiotemporal profile of visual system

activation revealed by current source density analysis in the awake macaque. Cerebral

Cortex 8:575–592.

Schultz W (2015) Neuronal Reward and Decision Signals: From Theories to Data.

Physiological reviews 95:853–951.

Sesack SR, Grace AA (2010) Cortico-Basal Ganglia reward network: microcircuitry.

Neuropsychopharmacology : official publication of the American College of

Neuropsychopharmacology 35:27–47.

Sohoglu E, Chait M (2016) Detecting and representing predictable structure during auditory

scene analysis. eLife 5.

Stark H, Scheich H (1997) Dopaminergic and serotonergic neurotransmission systems are

differentially involved in auditory cortex learning: a long-term microdialysis study of

metabolites. Journal of neurochemistry 68:691–697.

Stuber GD, Hnasko TS, Britt JP, Edwards RH, Bonci A (2010) Dopaminergic Terminals in

the Nucleus Accumbens But Not the Dorsal Striatum Corelease Glutamate. Journal of

Neuroscience 30:8229–8233.

Stujenske JM, Spellman T, Gordon JA (2015) Modeling the Spatiotemporal Dynamics of

Light and Heat Propagation for In Vivo Optogenetics. Cell Reports 12:525–534.

Sugimoto S, Sakurada M, Horikawa J, Taniguchi I (1997) The columnar and layer-specific

response properties of neurons in the primary auditory cortex of Mongolian gerbils.

Hearing Research 112:175–185.

Szymanski FD, Garcia-Lazaro JA, Schnupp JWH (2009) Current Source Density Profiles of

Stimulus-Specific Adaptation in Rat Auditory Cortex. Journal of Neurophysiology

102:1483–1490.

Tan KR, Yvon C, Turiault M, Mirzabekov JJ, Doehner J, Labouèbe G, Deisseroth K, Tye

KM, Lüscher C (2012) GABA Neurons of the VTA Drive Conditioned Place Aversion.

Neuron 73:1173–1183.

Van Zessen R, Phillips JL, Budygin EA, Stuber GD (2012) Activation of VTA GABA

Neurons Disrupts Reward Consumption. Neuron 73:1184–1194.

Wang H-L, Qi J, Zhang S, Wang H, Morales M (2015) Rewarding Effects of Optical

Stimulation of Ventral Tegmental Area Glutamatergic Neurons. Journal of Neuroscience.

Williams SM, Goldman-Rakic PS (1993) Characterization of the dopaminergic innervation of

the primate frontal cortex using a dopamine-specific antibody. Cerebral Cortex.

Yizhar O, Fenno LE, Davidson TJ, Mogri M, Deisseroth K (2011) Optogenetics in neural

systems. Neuron 71:9–34.

Yona G, Meitav N, Kahn I, Shoham S (2016) Realistic Numerical and Analytical Modeling of

Light Scattering in Brain Tissue for Optogenetic Applications(1,2,3). eNeuro

3:ENEURO.0059-15.2015.

Yoo JH, Zell V, Gutierrez-Reed N, Wu J, Ressler R, Shenasa MA, Johnson AB, Fife KH,

Faget L, Hnasko TS (2016) Ventral tegmental area glutamate neurons co-release GABA

and promote positive reinforcement. Nature Communications 7:1–13.

Zhao WJ, Kremkow J, Poulet JFA (2016) Translaminar Cortical Membrane Potential

Synchrony in Behaving Mice. Cell Reports.

 Figure legends:

Figure 1 - Anatomical validation of fiber position relative to target side and optogenetic

intracranial self-stimulation

(A) Schematic overview of fiber implant position for optogenetic stimulation of the VTA, and

the ipsilateral AI recording side. Dashed boxes indicate representative regions of interest shown

in B) identified by yellow and red asterisks. (B) Top, YFP expression in the temporal cortex

covering the region of AI after virus transduction of the VTA indicates fiber terminals from the

VTA targeting mainly upper layers and collaterals in deeper layers (white arrows). YFP

expression was detectable in multiple regions of the neocortex (not shown). Bottom, co-

fluorescence of YFP (AAV-CamKIIα-C1V1 (E162T)-p2A-eYFP) and TH-immunostaining

(Alexa Fluor 546). Area of co-fluorescence is indicative of overlapping neuronal populations

expressing the virus and TH mainly found in the VTA, and not the substantia nigra (Lippert et

al., 2018). Scale bars indicate 250 and 200 µm, respectively. (C) Schematic representation of

optogenetic intracranial self-stimulation: While nose-poking the lever, animals induce a light

flash towards the VTA resulting in extensive self-stimulating behavior (Brocka et al., 2018).

(D) Gerbil skull with prominent landmarks (bregma, lambda and occipital crest) for anatomical

reference (modified from Radtke-Schuller et al., 2016). For experiments, bregma has been used

for reference of medio-lateral (ML) and anterior-posterior (AP) position of the VTA. (E)

Estimated actual AP/ML coordinates with mean coordinates (AP:-3.57 mm; ML: 584.14 µm)

relative to bregma of the C1V1group (n=7) as estimated with the gerbil brain atlas (Radtke-

Schuller et al., 2016). (F) Fiber distance (mean ± SEM) towards area of co-fluorescence (146.8

± 38.9 µm; n=7). Compare with bottom picture of B). (G) Averaged lever pressing rates (lever

presses/ min) of the C1V1 (n=12) and YFP (n=8) groups over 10 days. A representative video

of a C1V1 animal is accessible as Movie 1-1.

Figure 2 - Electrophysiological measurements and effects of VTA stimulation on tone-evoked CSD profiles in A1

(A) Schematic representation of the timeline of measurements. Pre baseline measurements (Tone stimulation: 0.125-32 kHz, tone duration: 200 ms, ISI: 0.6-0.8, 50 pseudorandomized repetitions, 65 dB SPL, 7.5 min) were performed until CSD patterns had stabilized (45-75 min). A single measurement of combined tone and laser stimulation (25 Hz, 473 nm, 10 mW) was carried out and followed by a series of post-measurements (up to 60 min). (B) Representative example of a CSD profile after BF- (top) and off-BF (bottom) stimulation before, during and 30 minutes after the combined tone/VTA-laser measurement from an animal of the C1V1 group. Before VTA stimulation, BF-evoked CSD profiles revealed a canonical feedforward pattern of stimulus-evoked current flow starting with initial sink components in cortical layers III/IV and Vb/VIa. Subsequent sink activity was found in layers Va, I/II and VIb . Activity after off-BF stimulation showed a less pronounced columnar activation in the pre-condition. During paired tone and VTA stimulation, both stimulation frequencies evoked stronger and prolonged translaminar current flow, which was still enhanced 30 minutes later. Cortical layers are indicated. Dashed vertical lines indicate tone onset and offset.

Figure 3 - Binned time courses of single-trial sink peak amplitude data of layer-specific CSD-traces

Temporal development of single-trial sink peak amplitudes for early sink activity (sink onset <50 ms) in layers III/IV, Va and Vb/VIa and late sinks (sink onset >50 ms) in layers I/II and VIb for BF, near- and non-BFs. Layer I/II peak amplitudes showed a significant increase for the C1V1 group during laser and in post-measurements, surpassing the +10% criterion within

the BF and near-BF bins. Significant differences from the control group are again revealed by

LME models and indicated by asterisks. Layer I/II peak amplitudes in the YFP and control

group did not show corresponding increases over the recording time but showed a decrease and

recovery for the YFP group. Layer III/IV activity did not reveal differences over time for BF-

evoked responses in all groups. With spectral distance from the BF the C1V1 group showed a

moderate increase yielding significance at the >30 min time bin compared to control animals.

Within the non-BF bin, peak amplitudes showed a significant decrease in the YFP group during

and after laser stimulation. Layer Va peak amplitudes were most stable across time and groups

only displaying a significant decrease for the YFP group in the non-BF bin at <30 min. Layer

Vb/VIa peak amplitudes in the C1V1 group were significantly increased in all post-

measurements after BF-stimulation. For stimulation with near-BF a similar trend was found.

Significant changes between the YFP and control group were again due to decreased peak

amplitudes in the YFP group. Layer VIb displayed a highly significant increase in the C1V1

group during laser stimulation, most prominent in the non-BF bin, and less prominent in near-

BF and BF bins. Laser-induced increase only persisted for the next time bin, but then recovered

to pre-condition. While the control group showed stable peak amplitudes over the time course,

YFP animals again showed a moderate decrease most prominent in off-BF bins.

Figure 4 - Layer-specific early and late changes of tuning sharpness of the C1V1 group

For data from the C1V1 group, peak amplitudes within each layer were normalized to BF-

responses before laser stimulation. We fitted a linear line across evoked responses across the

BF-, near-BF- and non-BF-bin and calculated the slope as indicator for spectral tuning

sharpness (for pre, laser, >30 min). Sharpest tuning properties of peak amplitudes were found

in layers III/IV (slope of 0.28). Tuning in layers III/IV was not affected by VTA stimulation.

Layers I/II showed less prominent tuning revealed by a shallower slope in the pre-condition.

While VTA stimulation did not yield an immediate significant difference, slopes indicated a

significantly sharper tuning >30 min after VTA stimulation ($p < 0.01$). This was due to mainly

an increase in BF-evoked peak amplitudes in accordance with Fig. 5. Peak amplitudes in

infragranular layers Vb/VIa showed a similar effect as layers I/II with a trend of sharper tuning

during VTA stimulation that yielded significance >30 min later. In contrast, only in late

infragranular layer VIb, peak amplitudes revealed a significantly shallower slope indicative of

a broader frequency tuning during laser stimulation (Pre:laser; $p < 0.01$), which recovered after

>30 min (Pre:<30 min; $p = 0.1$). This can be seen by the flat slope between non- and BF bins.

Layer Va only showed a minor decrease of the slope during VTA stimulation that recovered

partially >30 min later. Significances indicated are based on slope comparisons performed using

the lsmeans package in R (version 2.27-62). For statistical explanation, see text.

Figure 5 - Temporal development of early and late AVREC and ResidualCSD activity

(A) Grand average traces of the BF-evoked AVREC (top) and ResidualCSD (bottom) of the

C1V1 group (n=12) for binned pre, laser and post (<30, >30 &>45 min)-measurements. Right,

insets illustrate that AVREC activity is mainly associated with local columnar activity, while

the ResidualCSD quantifies contributions from lateral corticocortical input (cf. Happel et al.,

2014, 2010). (B) Temporal time courses of Pre-normalized early/late AVREC and early/late

ResidualCSD traces of C1V1 (red), control (blue) and YFP (green) groups (n=12, n=7, n=7,

respectively) shown for BF stimulation. Early AVREC signal displayed a significant increase

in post-measurements for the C1V1 group whereas time courses of the control and YFP groups

were stable. Late AVREC signal showed no significant differences between either group. RMS

ResidualCSD values of early and late time windows displayed a similar temporal development

as the early AVREC becoming significantly increased in the later post-measurements within

the C1V1 group. Control and YFP groups showed no significant changes over time. Note the

peaking in the early ResidualCSD RMS during laser stimulation, which is only present in the

C1V1 group. Asterisks mark significant changes according to the applied mixed linear models

of the normalized data towards the control group (see Materials and Methods). (C) Tuning

curves of the RMS value normalized to BF-evoked responses in the pre-condition for early/late

AVREC and early/late ResidualCSD for binned time points for C1V1 group. Note the general

increase of the overall tuning curve relative to pre measurements.

Figure 6 - Normalized single-trial early and late AVREC and ResidualCSD RMS values over time

(A) Temporal time course of averaged early single-trial RMS AVREC (top) and ResidualCSD (bottom) values normalized to pre-measurements. Data for BF, near-BF and non-BF stimulation is shown separately and was binned for pre-measurements, laser, and post-measurements (<30 min, >30, >45 min). Note the prominent increase of early RMS AVREC values for the C1V1 group whereas control and YFP group displayed a more stable or decreasing time course, respectively. Also, early RMS ResidualCSD values significantly increased during and after the Laser measurements for BF, near- and non-BF bins of the C1V1 group, whereas YFP and control display a more stable time course. (B) Temporal time course of late Pre-normalized RMS AVREC and ResidualCSD values of single-trial data for BF, near-BF and non-BF. Late RMS AVREC values for the C1V1showed a mild, yet not significant, increase that did not surpass the +10% criterion. The control group displayed a stable and slightly declining time course for the late RMS AVREC values, while the YFP group showed a moderate decrease that recovered over time. Late RMS ResidualCSD values increased during and after the Laser measurements for BF, near- and non-BF bins of the C1V1 group, whereas YFP and control groups both showed a more stable time course. Significances in A) and B) were calculated using mixed linear models on the normalized data. Asterisks indicate corresponding significances in comparison to the control group, in case values exceed the $\pm$ 10% pre-criterion. Single-trial CSD data in following figure is presented equally.

Figure 7 - Schematic illustration of convergent deep layer inputs from sensory thalamus

and VTA exhibiting a local frequency-specific gain amplification.

VTA projections mainly target upper and infragranular layers of the gerbil auditory cortex (see

also Fig. 1 B). Stimulating the VTA projections led to a frequency-specific gain amplification

of early sensory-evoked responses in thalamocortical recipient layers Vb/VIa. This gain

increase effectuated an enhanced spectral tuning representation in supragranular layers I/II. The

supragranular gain modulation might hence be inherited via intratelencephalic neurons

targeting upper layers and bypassing granular input circuits (Harris and Mrsic-Flogel, 2013),

may originate from a direct influence of dopamine released in upper layers or a combination of

both. MGB, medial geniculate body.

Supplementary Information:

Movie 1-1 - Optogenetic intracranial self-stimulation induced by lever pressing

Animal video clip of an intracranial self-stimulation training session. Animals were placed into

a 25x35x22cm³ Plexiglas box for 20 min once a day. Optogenetic implants were attached via

an optical fiber with a shutter-controlled laser (10 mW, 473 nm). By pressing an installed lever,

animals were able to trigger a laser pulse (25 Hz, 5 ms) for VTA stimulation. Highly addicted

animals ($\geq$ 850 lever presses per session), were kept for optogenetic experiments whereas non-

897 addicted animals were rejected from this study (Compare behavior of the animal in the

898 background).

Figure 2-1 - Assignment of layers to CSD channels and semi-automatic detection of sink-activity

Based on previous studies, earliest tone-evoked components are related to layers Vb/VIa and

III/IV (Szymanski et al., 2009; Happel et al., 2010). Subsequent synaptic activity is conveyed

to layer Va and VIb, and layers I/II, as indicated in the representative example. Channels

assigned to individual layers were averaged and activity attributed to sources (positive values)

were dismissed. For visual purposes, a corresponding average-sink trace from layer III/IV was

inverted. Based on the pre-stimulus baseline signals (200 ms), the baseline mean value + 1

standard deviation (STD) was calculated as response threshold for each layer-dependent CSD

trace. A deflection that surpassed a 1-fold STD was used to determine the onset and offset of

early and late sink activations (see Materials and Methods). Significant sink onset activity

within the first 50 ms after tone onset was assigned as early activity, while later onsets were

assigned as late activity.

Figure 3-1 - Grand average AVREC and ResidualCSD traces for YFP and control groups

(A) AVREC and ResidualCSD traces of the YFP group (n=7) for binned pre-measurements,

laser-measurement, post-measurements < 30 min, post-measurements > 30 min and post-

measurements > 45 min. Dashed lines indicate tonal on- and offset. Areas for early and late

RMS calculations are underlined with gray. Group sorting was performed according to the BF

of the granular sink peak amplitudes. (B) AVREC and ResidualCSD traces of the control group

(n=7) for binned pre-measurements, laser-measurement, post-measurements < 30 min, post-

measurements > 30 min and post-measurements > 45 min. Dashed lines indicate tonal on- and

offset. Areas for early and late RMS calculations are underlined with gray. Group sorting was

performed according to the BF of the granular sink peak amplitudes.

A Fiber implant · Shaft electrode · AI · VTA

B VI · V · III-IV · I-II · 250μm · Fiber implant · VTA · SNR · 200μm · *CamKIIα-C1V1(E162T)-Ts-p2A-eYFP*

C Optical VTA-self stimulation · PC · Laser · 473 nm · AI · VTA

D bregma · ML · AP · lambda · occipital crest

E Rel. distances to bregma · AP-Axis [mm] · ML-Axis [μm] · 584.14 μm, -3.57 mm

F Distance fiber tip to VTA [μm]

G C1V1, n= 12 · YFP, n= 8 · Lever presses per minute · Sessions [days]

ACx

I/II

III/IV

Va

Vb/VIa

VIb

VTA

auditory MGB