Optogenetic stimulation of the VTA modulates a frequency-specific gain of

thalamocortical inputs in infragranular layers of the auditory cortex

Authors: Michael G. K. Brunk*[1], Katrina E. Deane[1], Martin Kisse[1], Matthias Deliano[1], Silvia

Vieweg[1], Frank W. Ohl[1,2,3], Michael T. Lippert[1,2], Max F. K. Happel*[1,3]

1 Department of Systems Physiology of Learning, Leibniz Institute for Neurobiology, 39118

Magdeburg, Germany

2 Center for Behavioral Brain Sciences (CBBS), 39106 Magdeburg, Germany

3 Institute for Biology, Otto-von-Guericke-University, 39120 Magdeburg, Germany

* Corresponding authors

Correspondence should be addressed to:

Michael Brunk or Max Happel

Department of Systems Physiology of Learning

Leibniz Institute for Neurobiology

Brenneckestr. 6

D-39118 Magdeburg

Tel.: +49-(0)391-6263-94411

Fax: +49-(0)391-6263-95489

Mail:

mbrunk@lin-magdeburg.de

mhappel@lin-magdeburg.de

Abstract

Background: Reward associations during auditory learning induce cortical plasticity in the primary auditory cortex. A prominent source of such influence is the ventral tegmental area (VTA), which conveys a dopaminergic teaching signal to the primary auditory cortex. It is currently unknown, however, how the VTA circuitry thereby influences cortical frequency information processing and spectral integration. In this study, we therefore investigated the temporal effects of direct optogenetic stimulation of the VTA onto spectral integration in the auditory cortex on a synaptic circuit level by current-source-density analysis in anesthetized Mongolian gerbils.

Results: While auditory lemniscal input predominantly terminates in the granular input layers III/IV, we found that VTA-mediated modulation of spectral processing is relayed by a different circuit, namely enhanced thalamic inputs to the infragranular layers Vb/VIa. Activation of this circuit yields a frequency-specific gain amplification of local sensory input and enhances corticocortical information transfer, especially in supragranular layers I/II. This effects further persisted over more than 30 minutes after VTA stimulation.

Conclusions: Altogether, we demonstrate that the VTA exhibits a long-lasting influence on sensory cortical processing via infragranular layers transcending the signaling of a mere reward-prediction error. Our findings thereby demonstrate a cellular and circuit substrate for the influence of reinforcement-evaluating brain systems on sensory processing in the auditory cortex.

Keywords:

Dopamine, auditory cortex, Mongolian gerbil, current source density, ventral tegmental area, optogenetics

Background

The sensory cortex receives both bottom-up input relaying stimulus information from the sensory epithelia and top-down input from, for example, reinforcement-evaluating brain structures [1]. Among the latter, the ventral tegmental area (VTA) is a key structure associated with the coding of reward, reward prediction, and reward prediction error [2]. Especially in the framework of reward prediction error coding, projections of dopamine neurons in the VTA to the striatum and prefrontal cortex have been investigated in great detail [3]. In contrast, the anatomy of projections from the VTA towards sensory cortices [4–6], and consequently its direct impact on cortical processing, has remained rather elusive.

Dopamine released in the sensory cortex may complement bottom-up stimulus processing with a behaviorally relevant representation of stimulus value and salience to support adaptive behavior [2, 7, 8]. Consistently, for the case of the auditory cortex (ACx), intracortical dopamine levels [9] and their experimental manipulation [10, 11] were shown to affect behavioral measures in auditory learning scenarios. Thus, dopamine appears to be involved in several general behaviorally relevant functions, including auditory perceptual decision making, prediction, and learning, which are increasingly realized to be supported by the ACx [12]. In agreement, we have previously shown that the pharmacological stimulation of D1/D5 receptors influences sensory processing at the level of both local and wide-spread circuits in auditory cortex [13].

In order to determine the contribution of VTA projections to these cortical effects of dopamine, we optogenetically stimulated the projection neurons of the VTA in adult male Mongolian gerbils (*Meriones unguiculatus*) and measured the layer-specific processing in the auditory cortex by tone-evoked current-source density (CSD) analysis. We demonstrate that VTA stimulation effectuated a sensory gain amplification via thalamocortical inputs in the deep layers Vb/VIa, rather than via recurrent excitation in layer III/IV [14, 15]. Our results

demonstrate for the first time a functional diversification of the anatomically distinct

thalamocortical input systems in the sensory cortex. Reward-modulated sensory input in deep

layer neurons therefore might provide a cellular substrate for integrating sensory and task-

related information in the service of sensory-based decision-making and reinforcement

learning.

Results

Optogenetic stimulation of the VTA evokes self-stimulation behavior

To selectively target the excitatory projection neurons of the VTA, we expressed the red-shifted

opsin C1V1 under the CamKIIα-promotor [16] by a stereotactically guided microinjection (Fig.

1 A; 700 nl; AAV-CamKIIα-C1V1(E162T)-p2A-eYFP, 3e12 particles per ml, UNC Vector

Core). Opsin expression in the VTA overlapped significantly with tyrosine hydroxylase (TH)-

immunostaining, selective for dopaminergic cell populations. The connectivity of the

transduced neurons with the auditory cortex was confirmed by histology (Fig. 1 A,B) in

agreement with previous reports [4, 17]. After viral transduction, we waited 2-3 weeks for

sufficient transgene expression before subjects were trained over the course of 10 days in an

optogenetic intracranial self-stimulation task (Fig. 1 C; Additional file 1; 10 pulses, 25 Hz, 473

nm laser-light per nose-poke; 10 mW power at fiber implant) [18]. Animals that showed more

than 50 presses/minute were chosen as the C1V1 group for electrophysiological experiments

(Fig. 1 G) [19, 20]. Anatomical validation of the fiber position relative to Bregma (AP:-3.57

mm, ML: 584.14 μm; cf. Fig. 1 D and E), as well as the fiber distance towards the area of opsin

expression (Fig. 1 F, 146.8 ± 38.9 μm), was performed in a subset of animals (n=7) in the C1V1

group (see Materials and Methods). In contrast to the opsin group, animals of the YFP-control

group showed no self-stimulation behavior (Fig. 1 G).

Effect of VTA-activation on tone-evoked columnar activity in the auditory cortex

After evaluation of self-stimulation behavior in the C1V1 and YFP group, we performed acute laminar local field potential electrophysiological recordings from their auditory cortices under ketamine-xylazine anesthesia. This approach allows the investigation of circuitry effects by the stimulation of the VTA on sensory cortical processing. Additionally, we also recorded from a control group of naïve animals, which underwent no implantation or laser stimulation, to determine which effects were due to opsin activation or laser light. Pure tones were presented in iso-octave steps in order to cover the main tonotopic range of the gerbil (Stimulation frequencies ranged from 0.125 kHz – 32 kHz, duration: 200 ms, ISI 0.6-0.8s, 50 rep., 65 dB SPL). After insertion of the recording electrode (NeuroNexus A1x32-50-413), sets of 50 repetitions per stimulus (~7.5 minutes) were recorded over 45-75 min to guarantee stable recordings (Fig. 2 A; see Materials and Methods). We then paired the presentation of each pure tone with laser-stimulation of the VTA (473 nm; 25 Hz; 10 ms pulse length, 200 ms duration). Thereby, the VTA was stimulated 400 times within 7.5 minutes of the combined laser/tone presentation (Fig. 2). After this paired measurement, consecutive tone-only measurements (post) were performed over the course of 60 minutes. This procedure allowed us to observe phasic and long-lasting effects of optogenetic VTA stimulation on auditory cortex tone-evoked activity.

Fig. 2 B displays a representative example of pure-tone evoked CSD-profiles after stimulation with the best frequency (BF, top) and an off-BF (bottom, 2 octaves apart) before (left), during (center) and 30 minutes after (right) VTA- stimulation in a C1V1-transduced animal. Before VTA stimulation, tone-evoked CSD profiles show the canonical feed-forward pattern of early current sink activity in layers III/IV and Vb/VIa (Fig. 2 B, left) [15, 21, 22]. Subsequent later supra- and infragranular sink activities can be related to intracortical synaptic circuits in layers

I/II, Va and VIb (see semi-automatic detection of sink activity in Materials and Methods and

Fig. 2 B and Additional file 2). Responses to frequencies apart from the BF generally evoke

less prominent tone-evoked activity across all cortical layers. When pure tone stimulation is

paired with VTA stimulation, we observed prominent changes of amplitudes, but not

spatiotemporal flow of synaptic activity (Fig. 2 B, center). This increase of tone-evoked cortical

processing was long-lasting and persisted over approximately 30 min after the cessation of laser

stimulation (Fig. 2 B, right).

Laminar specificity of transient and long-lasting effects of VTA stimulation

In order to relate the observed changes of in current flow to distinct cortical layers, we

separately analyzed CSD-traces from respective cortical layers (I/II, III/IV, Va, Vb/VIa and

VIb; see Materials and Methods). Prominent sink components of cortical layers III/IV, Va, and

Vb/VIa were related to early time windows of 0-50 ms after tone onset, while cortical layers

I/II and VIb felt into later time windows of >50 ms determined by sink onset (cf. Additional

file 2).

Described effects during and after VTA stimulation might impact on sensory processing in a

less stimulus-locked manner and influence trial-by-trial signal-to-noise ratio [23]. In order to

capture such variability, we have utilized a single-trial analysis using LME models [24] on

peak amplitudes of each layer as a function of time before and after VTA stimulation separately

for BF, near- ($\pm$ 1-2 octaves) and non-BF ($\pm$ 3-4 octaves) stimulation for all groups (Fig. 3).

Within each frequency bin, data was normalized to pre-measurements in order to characterize

relative changes over time and compared both groups with laser treatment (C1V1, YFP) against

the control group. Within the control group, we observed a variation of tone-evoked responses

of up to $\pm$ 10% over the entire recording time. Therefore, to obtain main effects of the laser

treatment and account for general variance, we quantified and statistically tested only data that

crossed this ± 10% criterion of the normalized pre-measurements (see dashed lines in Fig. 3).

In the C1V1 group, VTA stimulation led to a significant increase of peak amplitudes in

infragranular layers Vb/VIa most prominently after BF-stimulation for up to 45 minutes. The

main current sink in thalamocortical input layers III/IV, in contrast, showed only minor

modulation after VTA stimulation. Only 30 minutes after VTA stimulation off-BF evoked

activity was increased. Activity in layer Va was unaffected by VTA stimulation. Late onset sink

components in layers I/II and VIb showed opposing temporal effects after VTA stimulation.

While supragranular activity increased gradually and long-lasting mainly for BF/near-BF

stimulation, infragranular layer VIb activity was most strongly amplified for non-BF only

during combined tone-laser stimulation. In the YFP-group peak amplitudes of all sink

components showed a general trend to be reduced during laser-stimulation and recovered over

time within the >45min of post recordings. These effects were potentially due to light-related

heating of the stimulation site [25, 26], which might shift the balance of excitation and

inhibition in the VTA [27–30]. Data from control animals displayed a mainly stable amplitude

range within the ± 10% range of pre-measurements.

Based on the most prominent changes of evoked activity in the C1V1-group during VTA

stimulation (layer VIb) and over the longer period of >30 min after stimulation (layers I/II and

Vb/VIa), we compared slopes of evoked peak amplitudes as a function of stimulation frequency

(Fig. 4). All data were normalized to BF-evoked amplitudes of pre-measurements within each

layer separately. Slopes across the BF/near-BF/non-BF range therefore allow us to characterize

changes in tuning sharpness. Frequency-independent gain increase, in contrast, would not affect

the slope, but rather affect the intercept. In the pre-measurement, the most prominent frequency

tuning was observed in thalamocortical input layers III/IV. In the pre-condition amplitudes in

the non-BF bin were reduced about 60% compared to BF-evoked responses. Across frequency

bins, this led to a slope of 0.28 indicating a significant dependence of amplitude and stimulation

frequency (Fig. 4, blue line, p<0.001). During laser stimulation and 30 minutes after, slopes did

not change (change of slope >0.01; p>0.05) indicative of a stable frequency tuning in layers

III/IV. In layers Vb/VIa, a slope of 0.13 indicated a less prominent frequency tuning in the pre-

condition. During VTA stimulation the frequency tuning was unaffected (change of slope

<0.01; p<0.05) but showed a significant increase of sharpness 30 minutes later (change of slope

pre:30 min of +0.05; p<0.001). Comparably, in supragranular layers, I/II slopes of the pre-

condition (0.09; p<0.001) did not significantly increase during VTA stimulation (change of

slope +0.04; p=0.09), but significantly increased >30 min later (change of slope pre:30 min

+0.08, p<0.001). Hence, cortical layers Vb/VIa and I/II showed no change of tuning bandwidth

during the paired tone/VTA stimulation, but a significant increase in spectral integration

prolonged for up to 30 minutes afterwards. Activity in layer VIb showed a prominent increase

during VTA stimulation selectively for the off-BF bins (cf. Fig. 3; bottom right), and, in

accordance, we found slopes to be significantly decreased during laser stimulation (change of

slope pre:laser –0.06; p<0.001), which recovered after 30 minutes (change of slope Pre:30 min

–0.02, p=0.10). Layer Va similarly showed a decrease in slope during laser-stimulation (change

of slope pre:laser –0.04; p<0.001), and also recovered 30 minutes later (p>0.05). VTA

stimulation yielded a transient increase in response bandwidth in cortical layers Va and VIb.

Based on the layer-specific modulation of early tone-evoked inputs in thalamocortical-recipient

layers Vb/VIa and subsequent synaptic inputs in supragranular layers, we speculate that

stimulation of the VTA affects early and late sensory processing on a local and corticocortical

level, respectively [cf. 13].

Impact on local intracolumnar and corticocortical synaptic inputs

In a next step, we therefore aimed to disentangle early activity associated with local

thalamocortical input, and subsequent activity associated with corticocortical processing. We

compared changes in the tone-evoked average rectified CSD (AVREC) and the residuals of the

CSD (ResidualCSD) during early (0-50 ms) and late (80-300 ms) time windows (Fig. 5 A).

This rationale is based on previous findings demonstrating that the residual CSD reflects the

spatiotemporal ratio of unbalanced sinks and sources providing a quantitative measure of

horizontal intercolumnar inputs to a given cortical site [15, 21]. The AVREC, on the other hand,

reflects the overall intracolumnar current flow at a given recording site [31, 32]. Fig. 5 A (left)

shows changes of both the grand average of the AVREC (top) and relative CSD (bottom)

waveforms before, during, and, in certain time windows, after VTA stimulation in the C1V1

group. Comparable changes were not observed in neither the YFP nor control group (Additional

file 3). To further investigate temporal changes of BF-evoked responses in more detail, time

course of binned root mean square (RMS) values for early and late averaged and normalized

AVREC and ResidualCSD windows of each group were plotted as a function of time

(normalized to pre-measurements). RMS values of the AVREC during the early phase (0-50

ms) displayed a significant increase in the C1V1 group from 15 min until 52 min after VTA

stimulation. Significant effects were validated with a LME model (Fig. 5 B; $p<0.001$-0.025).

During the late phase (80-300 ms) after tone onset, RMS of the averaged AVREC did not show

any significant changes between each group (Fig. 5 B). During laser stimulation, a significant

increase has only been observed for the averaged early ResidualCSD indicative of amplified

lateral contributions to early tone-evoked columnar activity in the C1V1 group ($p=0.04$).

Furthermore, early and late ResidualCSD were both significantly enhanced during later time

points after the VTA stimulation (early ResidualCSD: 22.5-37.5 min; $p=0.014$-0.039; late

ResidualCSD: 37.5-45 min, $p=0.008$-0.033). Averaged RMS tuning curves of the AVREC and

ResidualCSD of the C1V1 group revealed an increase of tone-evoked activity during and after

laser stimulation across the entire range of BF, near-BF ($\pm$ 1-2 octaves) and non-BF ($\pm$ 3-4

octaves) stimulation (Fig. 5 C).

Integration of local and corticocortical synaptic inputs in dependence of the stimulation frequency

All single-trial AVREC and ResidualCSD RMS values were normalized to pre measurements. Normalized values were analyzed as a function of time before and after VTA stimulation binned for BF, near-, and non-BF stimulation for all groups (Fig.6). Fig. 6 A (top) shows a gradual increase of the early AVREC compared to the control- and YFP-group independent of stimulation frequency. For BF- and near-BF-stimulation, this yielded a significant increase at >30 min after VTA stimulation. The ResidualCSD also showed a strong increase almost independent of the stimulation frequency already during the combined tone-laser presentation and all following post measurements. Consistently with the layer-specific activity, there was a less pronounced decrease of the near/non-BF-evoked AVREC in early and late time windows in the YFP-group during laser stimulation, which started to recover over time (Fig. 6 A, bottom).

In the later time window of 80-300 ms after tone-onset, only the ResidualCSD showed a significant increase during and up to 30 minutes after VTA stimulation (Fig. 6 B). Altogether, VTA stimulation led to a gradual increase of the overall early current flow most pronounced for BF- and near-BF stimulation. This effect was long-lasting and peaked around 30 minutes after stimulation. We further found an immediate and long-lasting increase of imbalance between sinks and sources based on the early and late ResidualCSD in all frequency bins indicative of amplified corticocortical activity.

Discussion

The VTA is a key structure to convey information about stimulus salience and valence to target areas distributed throughout the brain [2]. Receiving both sensory as well as VTA inputs, the

primary sensory cortex is ideally positioned to integrate such bottom-up and top-down

information to shape behavior through reinforcement learning. We here demonstrate for the

first time that activation of VTA-reward circuits amplifies sensory input via a yet largely

overlooked thalamocortical projection targeting infragranular layers Vb/VIa of the auditory

cortex [22, 33]. In contrast, granular layer III/IV input, a well-known substrate for sensory gain

modulation [14, 15, 33, 34], was unaffected by VTA stimulation. The impact of the observed

infragranular gain modulation on supragranular layers manifests as a translaminar long-lasting

enhancement of frequency tuning (see Fig. 7). Furthermore, we demonstrate that the frequency-

specific gain in deeper layers emerges on a local columnar level and yields an increased

corticocortical integration displayed in lateral input most likely terminating in upper layers.

This highlights the diverse, and yet elusive functions of cortical layers in the sensory cortex

[35]. The long-lasting modulation after VTA stimulation reveals that the influence on sensory

cortex transcends the time scales commonly associated with reward-prediction error coding.

VTA-activation enhances infragranular thalamocortical inputs in the auditory cortex

Optogenetic stimulation of VTA-projection neurons allowed us to investigate effects of reward-

related cross-regional circuits (Fig. 1) [19]. Roughly 8% of non-sensory inputs of gerbil ACx

arise from direct projections of the ipsilateral VTA, which terminate across the entire cortical

column [4]. Therefore, direct stimulation of the VTA projection neurons is the most effective

method to activate this circuit. In contrast, the success of optogenetic approaches to directly

stimulate cortical terminals have been challenged in recent studies and particularly do not allow

for an homogenous activation across all cortical layers [26]. VTA stimulation led to increased

early tone-evoked (0-50 ms) local columnar current flow and increased long-range

corticocortical activity. This effect persisted for up to 50 minutes after VTA stimulation and

was most prominent for the respective BF of the given recording site (Fig. 5). Hence, VTA

stimulation promotes the local gain of columnar input over a prolonged period and lead to

substantial corticocortical integration of spectral information. Notably, the gain amplification

of sensory input appeared exclusively in thalamocortical-recipient layers Vb/VIa but was absent

in the main thalamocortical-recipient granular layers III/IV (Fig. 3). This enhancement of

infragranular activity was locally relayed to supragranular layers I/II. This mechanism may

provide a way for intratelencephalic neurons in infragranular layers [36] to provide a

translaminar gain control of sensory representation depending on the experience of reward [37,

38]. Although direct projections of the ipsilateral VTA provide a significant contribution to

non-sensory inputs of the gerbil ACx [4], our approach may recruit other relay stations that

could contribute to the observed effects.

Previously, reported effects of dopamine on tuning sharpness in the ACx [39, 40] raised the

question on which specific cortical circuit motifs reward information modulates spectral

information. Our data showed that stimulating the VTA projections only affected early auditory

tuning sharpness in deep cortical layers, but not in the main thalamocortical input layer III/IV

[33, 41]. Sharper frequency representation was relayed to superficial layers I/II and persisted

over time after VTA stimulation. Our observations are thereby consistent with recent findings

from layer-specific high-field fMRI in human auditory cortex showing sharper tuning in upper

layers depending on attention and task engagement [42].

As a result, reward-circuit activation promotes a long-lasting gain enhancement of salient

auditory stimuli via strengthened infragranular thalamocortical inputs and increased sensory

integration in supragranular circuits. Our finding now provides an explanatory role of the

anatomically and physiologically well-described early infragranular inputs in layers Vb/VIa of

the sensory cortex [15, 22, 33, 43] as a circuit for integrating sensory and task-related

information.

 ### Persistent cortical modulation might underlie dopamine-dependent influences on

 ### learning

Recent reports have highlighted that sensory-evoked responses in deep layers are modulated by

behavioral states [44, 45] and undergo substantial learning-related plasticity [46] not described

for middle layer thalamocortical input [47]. Our data now assigns a functional impact of the

VTA on deep cortical layers—particularly under the control of dopaminergic neuromodulation

(Fig. 7).

Optogenetic VTA stimulation preferentially recruits the main output population of

dopaminergic projection neurons making up roughly 55-65% of all VTA neurons [48–50].

Nevertheless, it has been demonstrated that the small population of non-dopaminergic

projection neurons also convey reward information [51–53] and particularly promote the

activation of brain-wide regions including the auditory cortex [19]. Furthermore, there is

overlap between dopaminergic and glutamatergic cells through transmitter co-release [54–56].

Nevertheless, the long duration of responses modulation in our data set strongly argues for a

significant dopaminergic component, as glutamatergic effects are generally more transient [57].

Further evidence for a dopaminergic origin of the long-lasting effects is provided by the

distribution of dopamine receptors. In the sensory cortex, they are concentrated mainly in

infragranular and supragranular layers where we see modulatory effects (Fig. 1 B) [11, 58–62].

Conclusions

Phasic release of dopamine from the ventral tegmental area has long been associated with the

reward prediction error signal postulated by reinforcement learning theory. How dopamine

released by the VTA affects the plasticity that integrates sensory information and such learning-

related signals in sensory cortex, however, is unknown. Our findings therefore assign a specific

function to this rather understudied cortical circuit: VTA-modulated input to deep layer neurons

in the sensory cortex mediates the integration of sensory and task-related information even

when they are not precisely temporally contingent. Our study pursues the idea of a VTA-

mediated local gain of sensory input at infragranular thalamocortical terminals via presumably

presynaptic dopamine receptors [13, 63, 64]. Previous studies have described accumulative

levels of dopamine in the auditory cortex over 30 minutes after training onset impacting on

learning and long-term memory consolidation [9] suggesting a long-lasting action of dopamine

at the presynaptic terminals [11]. Thereby, the sensory gain amplification within deep layer

neurons in the sensory cortex described in this study potentially promotes a sustained perceptual

salience of incoming stimuli [65–67]. Our findings now provide a circuit level explanation of

the necessary integration of sensory and task-related information even when they are not

precisely temporally contingent. Thereby, this circuit might constitute a substrate for the credit

assignment in reinforcement learning theory concerned with the temporal delay between

sensory input and its behavioral consequences [68].

338   Methods

All experiments were carried out in adult male Mongolian gerbils (*Meriones unguiculatus*, age

4-8 months, body weight: 70-100 g; n= 26). The C1V1 group (n=12) consisted of animals that

were expressing the viral construct, allowing for optogenetic stimulation of the VTA whilst

recording. The control group (n=7) consisted of non-transfected animals to control for temporal

stability of the signal. In order to investigate effects due to laser-stimulation itself, an opsin-less

YFP group (n=7) was tested. Experiments were conducted in accordance with ethical animal

research standards defined by the German Law and approved by an ethics committee of the

State of Saxony-Anhalt.

Viral transfection of the VTA

Gerbils were anesthetized by intraperitoneal injection (0.06-0.08 ml/h) of 45% v/v ketamine (50 mg/ml, Ratiopharm GmbH), 5% v/v xylazine (Rompun, 2%, Bayer Vital GmbH) and 50% v/v of isotonic sodium-chloride solution (154 mmol/l, B. Braun AG). Animals were fixed in an automated stereotaxic injector (Neurostar) for virus transduction (700 nl; 100 nl/min; 5 min resting; C1V1: AAV5-CamKIIα-C1V1(E162T)-p2A-eYFP or YFP: AAV2-CamKIIα-eYFP 3E12 particles/ml, UNC Vector Core; target side: AP: -3.8-4.0; ML: 0.5; DV: -6.2 mm, bregma for reference) and fiber implantation with a planar setting of bregma and lambda. Custom made fiber implants (fiber implant length 8 mm, Ø 230 μm, NA 0.39) were implanted at target side (AP: - 3.8-4.0; ML: 0.5; DV: -5.8 mm). Fiber positions were determined by histological analysis and compared with the gerbil brain atlas (Fig. 1 D and E) [69]. Animals were kept for 3 weeks to allow recovery and enough protein expression.

Intracranial self-stimulation using optogenetic VTA activation

Successful activation of the VTA through optogenetic stimulation was confirmed in an intracranial self-stimulation paradigm. Animals were trained for 10 consecutive days (at 20 min per day). By activating a lever with their nose, they could elicit brief laser stimulation (10 pulses, 25 Hz, 473 nm, 10 mW) delivered to the VTA through the implanted fiber (Fig. 1 A, B and C). Only animals pressing the lever more than 50 presses/minute were kept for the C1V1 group (Fig. 1 G) [18–20].

Electrophysiological recordings

Surgical procedure for electrophysiological recordings have been described in detail elsewhere [40]. Briefly, animals have been anesthetized by intraperitoneal injection (0.06-0.08 ml/h) of 45% v/v ketamine (50 mg/ml, Ratiopharm GmbH), 5% v/v xylazine (Rompun, 2%, Bayer Vital GmbH) and 50% v/v of isotonic sodium-chloride solution (154 mmol/l, B. Braun AG). The right auditory cortex was exposed by trepanation and target location of AI was chosen based on vascular landmarks. Status of anesthesia was checked via paw withdrawal-reflex on a regular basis (10-20 min). Body temperature was kept stable at 34°C. Due to an epileptic seizure during the electrophysiological recordings we had to omit an animal from the YFP group.

Experiments have been carried out in a Faraday-shielded acoustic soundproof chamber. The animals head was fixed 1m away from the speaker (Tannoy arena satellite KI-8710-32) by screwing the head post to a custom-made head holder. Local field potentials (LFPs) have been recorded with a linear 32-channel shaft electrode (NeuroNexus A1x32-50-413), which was implanted perpendicular to the surface area of AI [15]. LFPs have been pre-amplified 500-fold and band-pass filtered (0.7-300 Hz) with a PBX2 preamplifier (Plexon Inc.). Data were digitized at a sampling frequency of 1 kHz with a multichannel recording system (Multichannel Acquisition Processor, Plexon Inc.).

Pure tones spanning 8 octaves (frequency range between 125 Hz-32 kHz, tone duration: 200 ms, ISI: 0.6-0.8, 50 pseudorandomized repetitions, 65 dB SPL, 7.5 min per measurement) were generated in MATLAB, converted into an analog signal by a data acquisition card (NI PCI-BNC2110, National Instruments, Germany), rooted through a controllable attenuator (gPAH, Guger, Technologies, Austria) and amplified by an audio amplifier (Thomas Tech Amp75). A measurement microphone and conditioning amplifier were used to calibrate acoustic stimuli (G.R.A.S. 26AM and B&K Nexus 2690-A, Bruel & Kjaer, Germany).

Electrophysiological measurements were performed according to the scheme of Fig. 2 A. After

implantation, laminar LFPs were measured for 45-75 min until evoked CSD profiles had been

stabilized (Pre-measurements). The last three recordings before laser treatment were taken as

the referencing Pre-measurements. For the C1V1and YFP groups, light fibers were additionally

connected to a laser for VTA stimulation. Laser stimulation (10 mW at the optical fiber of the

setup) was then coupled and synchronized with tone presentation for one measurement (25 Hz,

473 nm, 10 mW; 400 repetitions in total). This measurement was taken as the laser

measurement followed by a series of post-measurements (>60 min). After the experiment,

gerbils were perfused with 4% PFA as described previously [70] and brains were removed for

anatomical and histological validation.

Current source density analysis

Based on the tone-evoked LFPs for each presented stimulus we calculated the second spatial

derivative, indicating the laminar evoked current-source density (CSD) distribution in AI [71]:

$(1) -CSD \approx \frac{\delta^2\emptyset(z)}{\delta z^2} = \frac{\emptyset(z+n\Delta z)-2\emptyset(z)+\emptyset(z-n\Delta z)}{(n\Delta z)^2}$

The formula reflects the relation of the field potential ($\emptyset$), the spatial coordinate of the cortical

laminae (z), the inter-channel distance (Δz, 50 μm) and the differentiation grid (n). Prior to

CSD calculation, LFPs were smoothed using a weighted average of 9 channels (Hamming

window, spatial filter kernel size of 400 μm; linear extrapolation of 3 channels at boundaries)

[15].

CSD profiles revealed patterns of current influx (sinks) and efflux (sources). Based on CSD

profiles of the Pre-measurements, cortical layers were assigned to the respective channels of

sink components. Early sink components were used as indicators of layers III/IV, Va, and

Vb/VIa, whereas later sink activities were assigned to layers I/II, and VIb (Fig. 2; Additional

file 2).

CSD data was also transformed by rectifying and averaging the waveforms of each single

channel (*n*). This average rectified CSD (AVREC) provides a temporal waveform of the overall

local columnar transmembrane current flow [31, 32].

(2) $AVREC = \frac{\sum_{i=1}^{n} |CSD_i|(t)}{n}$

We further quantified the residue of the CSD defined as the sum of the non-rectified magnitudes

divided by the rectified magnitudes for each channel. Thereby, the ResidualCSD quantifies the

balance of the transmembrane charge transfer along the recording axis [72] and gives rise to the

lateral corticocortical contribution to stimulus related activity [15].

(3) $ResidualCSD = \frac{\sum_{i=1}^{n} CSD_i(t)}{n}$

In order to quantify the early (0-50 ms) and late (80-300 ms) contributions of the AVREC and

ResidualCSD, root mean square (RMS) amplitudes were calculated. Based on the previously

described layer indexing, averaged layer-dependent CSDs were calculated, which were used

for the semi-automatic sink detection (Additional file 2). Briefly, before detection of significant

tone-evoked activity, sources were omitted before calculating the mean of CSD traces within a

cortical layer. Then, the 1.5-fold standard deviation (SD) of the baseline (-200-0 ms before tone

presentation onset for all channels and stimuli) was used to calculate the intercepts of the

smoothed (10 ms gaussian sliding window) mean CSD trace. For early tone-evoked sink

activity in cortical layers III/IV, Va, and Vb/VIa, we used an onset of the first intercept between

0-50 ms after tone onset. For subsequent tone-evoked sink components in layers I/II and VIb

onsets were only taken >50 ms after tone-onset. Offsets of each sink were defined as the

crossings below the 1.5 SD baseline and for each cortical layer. Peak amplitudes of layer-

dependent CSD-traces were determined within time windows of on- and offsets from the

average of the corresponding CSD channels. For calculating the RMS value of AVREC and

ResidualCSD we used fixed time windows of 0-50 ms after tone-onset for early and 80-300 ms

for late activity.

Data sorting and analysis

Tuning curves of layer-specific CSD-traces, AVREC, and ResidualCSD were averaged

according to the best-frequency of each animal separately to plot an averaged group tuning

curve. For illustration, all non-BF frequencies above and below the BF were binned pairwise,

resulting into the grouping of near-BFs (± 1-2 octaves) and non-BFs (± 3-4 octaves) in 5

frequency bins [- non-BF, - near-BF, BF, + near-BF, + non-BF]. For further analysis of BF-,

near- and non-BF evoked responses, we observed data from time windows before, during, and

after laser stimulation. Evoked responses were normalized within each animal using the

corresponding mean values of the last three Pre-measurements before laser-stimulation (Fig. 2

A.

For statistical comparison, parameters of interest were analyzed on a single-trial level using

linear-mixed effect (LME) models (R Studio; R 3.5.1) [24] binned across measurements as

follows: pre 1-3 (-22.5 - -7.5 min), laser (0 min), post 1-3 (7.5-22.5 min), post 4-5 (30-37.5

min), post 6-7 (45-52.5 min). Peak amplitudes of single-trial data were determined within the

time windows from on- and offsets revealed on averaged CSD traces. LME models were

calculated and fitted using normalized data with the lmerTest-package (3.0-1) and a Wald-chi-

square-test was performed using the car-package (3.0-2). Animals were used as "by-subject"

random effects, to account for the repeated measurements during and across time points

whereas measurements and groups were used as fixed effects to explain physiological data. As

significance criterion for effects on single-trial level, the normalized signal needed to i)

exceeded a ± 10% criterion compared to Pre-measurement (based on the SEM of the average

jitter before laser treatment), and ii) show significant difference from control group data

revealed by significant fixed effects of the LME models. For slope estimations and comparison

between measurements, we used the R-package lsmeans (version 2.27-62) to calculate a least

square linear fit for non-, near- and BF data bins and check significant changes in tuning slope

properties between measurements.

Anatomical estimation of fiber position

PFA fixated brains of the C1V1 group (n =7) were cut in 50 μm thick slices around the region

of the VTA. Every second slice was used for Nissl staining to anatomically evaluate the fiber

position, whereas the remaining slices were used for immunostaining to determine the distance

from fiber tip to areas of tyrosine hydroxylase (TH) and YFP-co-fluorescence with FIJI ImageJ

(1.51g). Pictures were taken with a Leica microscope (Axioscop 2, Leica GmbH).

Immunostaining was performed as previously described [19] with a primary antibody against

tyrosine hydroxylase (1:1000 rabbit anti-TH, Milipore) and a secondary Alexa Fluor 546 goat-

anti-rabbit (1:400 anti-rabbit, Molecular probes) using an antibody diluent (Zytomed Systems).

Position of the fiber was estimated using the recently published gerbil brain atlas for reference

[69].

List of abbreviations

ACx   Auditory cortex

AI   Primary auditory cortex

AVREC   Average rectified CSD

BF   Best frequency

488	C1V1	Animal group expressing a functional chimeric opsin in the VTA
489	DA	Dopamine
490	CSD	Current source density
491	LFP	Local field potential
492	LME model	Linear mixed effect model
493	PFA	Paraformaldehyde
494	RMS	Root mean square
495	SEM	Standard error of mean
496	TH	Tyrosine hydroxylase
497	VTA	Ventral tegmental area
498	YFP	Animal group without a functional opsin only expressing a yellow fluorescent
499	protein in the VTA

Declarations

Author contributions:

FWO, MTL and MFKH designed research. MGKB and KED performed electrophysiology.
MGKB and SV performed histology and behavioral experiments. MGKB, KED, MK, MD and
MFKH analyzed data. MGKB, KED and MD contributed analytic tools. MGKB and MFKH
prepared the figures and wrote the initial draft. MGKB, KED, FWO, MTL and MFKH wrote
the manuscript. All authors reviewed and revised the manuscript.

Conflict of interest:

The authors declare no competing financial interests.

Acknowledgements:

We like to thank Janet Stallmann, Anja Gürke and Kathrin Ohl for assistance with surgeries,

histology- and immunostaining of brain slices. We would like to thank PD Dr. Eike Budinger

and Dr. Julia Henschke for their help and assistance in anatomical questions. Many thanks to

our interns for their assistance with anatomical reconstructions: Tarik Drewes, Katharina Eick,

Vincent Koop, Jan-Niklas Löbner, Julia Rieger and Ceylan Steinecke. This project and

corresponding research have been funded by the Leibniz Association WGL by the Leibniz

Postdoctoral Network, LPN (to MTL and MFKH) and, the Deutsche Forschungsgemeinschaft

DFG by the SFB779 (to MFKH and FWO) and SPP1665 (to MTL and FWO).

References:

1. Ohl FW. Role of cortical neurodynamics for understanding the neural basis of motivated

behavior - lessons from auditory category learning. Curr Opin Neurobiol. 2014;31:88–94.

doi:10.1016/j.conb.2014.08.014.

2. Bromberg-Martin ES, Matsumoto M, Hikosaka O. Dopamine in motivational control:

rewarding, aversive, and alerting. Neuron. 2010;68:815–34.

doi:10.1016/j.neuron.2010.11.022.

3. Schultz W. Neuronal Reward and Decision Signals: From Theories to Data. Physiol Rev.

2015;95:853–951. doi:10.1152/physrev.00023.2014.

4. Budinger E, Laszcz A, Lison H, Scheich H, Ohl FW. Non-sensory cortical and subcortical

connections of the primary auditory cortex in Mongolian gerbils: bottom-up and top-down

processing of neuronal information via field AI. Brain Res. 2008;1220:2–32.

doi:10.1016/j.brainres.2007.07.084.

5. Dinopoulos A, Parnavelas JG. The development of ventral tegmental area (VTA)

projections to the visual cortex of the rat. Neurosci Lett. 1991;134:12–6. doi:10.1016/0304-

3940(91)90497-H.

6. Gardner MPH, Schoenbaum G, Gershman SJ. Rethinking dopamine as generalized

prediction error. Proc R Soc B Biol Sci. 2018;285:20181645. doi:10.1098/rspb.2018.1645.

7. Ohl FW, Scheich H. Learning-induced plasticity in animal and human auditory cortex. Curr

Opin Neurobiol. 2005;15:470–7.

8. Happel MFK. Dopaminergic impact on local and global cortical circuit processing during

learning. Behav Brain Res. 2016;299:32–41. doi:10.1016/j.bbr.2015.11.016.

9. Stark H, Scheich H. Dopaminergic and serotonergic neurotransmission systems are

differentially involved in auditory cortex learning: a long-term microdialysis study of

metabolites. J Neurochem. 1997;68:691–7. http://www.ncbi.nlm.nih.gov/pubmed/9003057.

10. Schicknick H, Reichenbach N, Smalla K-H, Scheich H, Gundelfinger ED, Tischmeyer W.

Dopamine modulates memory consolidation of discrimination learning in the auditory cortex.

Eur J Neurosci. 2012;35:763–74. doi:10.1111/j.1460-9568.2012.07994.x.

11. Schicknick H, Schott BH, Budinger E, Smalla KH, Riedel A, Seidenbecher CI, et al.

Dopaminergic modulation of auditory cortex-dependent memory consolidation through

mTOR. Cereb Cortex. 2008;18:2646–58.

12. King AJ, Teki S, Willmore BDBB. Recent advances in understanding the auditory cortex.

F1000Research. 2018;7:1555. doi:10.12688/f1000research.15580.1.

13. Happel MFK, Deliano M, Handschuh J, Ohl FW. Dopamine-modulated recurrent

corticoefferent feedback in primary sensory cortex promotes detection of behaviorally

relevant stimuli. J Neurosci. 2014;34:1234–47. doi:10.1523/JNEUROSCI.1990-13.2014.

14. Liu BH, Wu GK, Arbuckle R, Tao HW, Zhang LI. Defining cortical frequency tuning

with recurrent excitatory circuitry. Nat Neurosci. 2007;10:1594–600.

15. Happel MFK, Jeschke M, Ohl FW. Spectral Integration in Primary Auditory Cortex

Attributable to Temporally Precise Convergence of Thalamocortical and Intracortical Input. J

Neurosci. 2010;30:11114–27. doi:10.1523/JNEUROSCI.0689-10.2010.

16. Guo S, Chen S, Zhang Q, Wang Y, Xu K, Zheng X. Optogenetic activation of the

excitatory neurons expressing CaMKIIα in the ventral tegmental area upregulates the

locomotor activity of free behaving rats. Biomed Res Int. 2014;2014:687469.

doi:10.1155/2014/687469.

17. Budinger E, Scheich H. Anatomical connections suitable for the direct processing of

neuronal information of different modalities via the rodent primary auditory cortex. Hear Res.

2009;258:16–27. doi:10.1016/j.heares.2009.04.021.

18. Lippert MT, Takagaki K, Weidner T, Brocka M, Tegtmeier J, Ohl FW. Optogenetic

Intracranial Self-Stimulation as a Method to Study the Plasticity-Inducing Effects of

Dopamine. In: Handbook of Behavioral Neuroscience. 2018. p. 311–26. doi:10.1016/B978-0-

12-812028-6.00017-3.

19. Brocka M, Helbing C, Vincenz D, Scherf T, Montag D, Goldschmidt J, et al.

Contributions of dopaminergic and non-dopaminergic neurons to VTA-stimulation induced

neurovascular responses in brain reward circuits. Neuroimage. 2018;177:88–97.

doi:10.1016/j.neuroimage.2018.04.059.

20. Helbing C, Brocka M, Scherf T, Lippert MT, Angenstein F. The role of the mesolimbic

dopamine system in the formation of blood-oxygen-level dependent responses in the medial

prefrontal/anterior cingulate cortex during high-frequency stimulation of the rat perforant

pathway. J Cereb Blood Flow Metab. 2016;36:2177–93. doi:10.1177/0271678X15615535.

21. Happel MFK, Ohl FW. Compensating Level-Dependent Frequency Representation in

Auditory Cortex by Synaptic Integration of Corticocortical Input. PLoS One.

2017;12:e0169461. doi:10.1371/journal.pone.0169461.

22. Szymanski FD, Garcia-Lazaro JA, Schnupp JWH. Current Source Density Profiles of

Stimulus-Specific Adaptation in Rat Auditory Cortex. J Neurophysiol. 2009;102:1483–90.

doi:10.1152/jn.00240.2009.

23. Kroener S, Chandler LJ, Phillips PEM, Seamans JK. Dopamine Modulates Persistent

Synaptic Activity and Enhances the Signal-to-Noise Ratio in the Prefrontal Cortex. PLoS

One. 2009;4:e6507. doi:10.1371/journal.pone.0006507.

24. Chang A, Bosnyak DJ, Trainor LJ. Beta oscillatory power modulation reflects the

predictability of pitch change. Cortex. 2018;106:248–60.

doi:10.1016/J.CORTEX.2018.06.008.

25. Arias-Gil G, Ohl FW, Takagaki K, Lippert MT. Measurement, modeling, and prediction

of temperature rise due to optogenetic brain stimulation. Neurophotonics. 2016;3:045007.

doi:10.1117/1.NPh.3.4.045007.

26. Stujenske JM, Spellman T, Gordon JA. Modeling the Spatiotemporal Dynamics of Light

and Heat Propagation for In Vivo Optogenetics. Cell Rep. 2015;12:525–34.

doi:10.1016/j.celrep.2015.06.036.

27. Johnson SW, North RA. Two types of neurone in the rat ventral tegmental area and their

synaptic inputs. J Physiol. 1992;450:455–68. doi:10.1113/jphysiol.1992.sp019136.

28. Paladini CA, Tepper JM. GABAA and GABAB antagonists differentially affect the firing

pattern of substantia nigra dopaminergic neurons in vivo. Synapse. 1999;32:165–76.

doi:10.1002/(SICI)1098-2396(19990601)32:3<165::AID-SYN3>3.0.CO;2-N.

29. Tan KR, Yvon C, Turiault M, Mirzabekov JJ, Doehner J, Labouèbe G, et al. GABA

Neurons of the VTA Drive Conditioned Place Aversion. Neuron. 2012;73:1173–83.

doi:10.1016/J.NEURON.2012.02.015.

30. Van Zessen R, Phillips JL, Budygin EA, Stuber GD. Activation of VTA GABA Neurons

Disrupts Reward Consumption. Neuron. 2012;73:1184–94.

doi:10.1016/j.neuron.2012.02.016.

31. Givre SJJ, Schroeder CEE, Arezzo JCC. Contribution of extrastriate area V4 to the

surface-recorded flash VEP in the awake macaque. Vision Res. 1994;34:415–28.

doi:10.1016/0042-6989(94)90156-2.

32. Schroeder CE, Metha a D, Givre SJ. A spatiotemporal profile of visual system activation

revealed by current source density analysis in the awake macaque. Cereb Cortex. 1998;8:575–

92. doi:10.1093/cercor/8.7.575.

33. Intskirveli I, Joshi A, Vizcarra-Chacón BJ, Metherate R. Spectral breadth and laminar

distribution of thalamocortical inputs to A1. J Neurophysiol. 2016;115:2083–94.

doi:10.1152/jn.00887.2015.

34. Bruno RM, Sakmann B. Cortex Is Driven by Weak but Synchronously Active

Thalamocortical Synapses. Science (80-). 2006;312:1622–7.

doi:10.1126/SCIENCE.1124593.

35. Adesnik H, Naka A. Cracking the Function of Layers in the Sensory Cortex. Neuron.

2018;100:1028–43. doi:10.1016/J.NEURON.2018.10.032.

36. Harris KD, Mrsic-Flogel TD. Cortical connectivity and sensory coding. Nature.

2013;503:51–8. doi:10.1038/nature12654.

37. Castejon C, Barros-Zulaica N, Nuñez A. Control of Somatosensory Cortical Processing by

Thalamic Posterior Medial Nucleus: A New Role of Thalamus in Cortical Function. PLoS

One. 2016;11:e0148169. doi:10.1371/journal.pone.0148169.

38. Jarvis S, Nikolic K, Schultz SR. Neuronal gain modulability is determined by dendritic

morphology: A computational optogenetic study. PLOS Comput Biol. 2018;14:e1006027.

doi:10.1371/journal.pcbi.1006027.

39. Bao S, Chan VT, Merzenich MM. Cortical remodelling induced by activity of ventral

tegmental dopamine neurons. Nature. 2001;412:79–83. doi:10.1038/35083586.

40. Deliano M, Brunk MGK, El-Tabbal M, Zempeltzi MM, Happel MFK, Ohl FW.

Dopaminergic neuromodulation of high gamma stimulus phase-locking in gerbil primary

auditory cortex mediated by D1/D5-receptors. Eur J Neurosci. 2018. doi:10.1111/ejn.13898.

41. Sugimoto S, Sakurada M, Horikawa J, Taniguchi I. The columnar and layer-specific

response properties of neurons in the primary auditory cortex of Mongolian gerbils. Hear Res.

1997;112:175–85.

42. De Martino F, Moerel M, Ugurbil K, Goebel R, Yacoub E, Formisano E. Frequency

preference and attention effects across cortical depths in the human primary auditory cortex.

Proc Natl Acad Sci U S A. 2015;112:16036–41. doi:10.1073/pnas.1507552112.

43. Constantinople CM, Bruno RM. Deep cortical layers are activated directly by thalamus.

Science. 2013;340:1591–4. doi:10.1126/science.1236425.

44. Zhao WJ, Kremkow J, Poulet JFA. Translaminar Cortical Membrane Potential Synchrony

in Behaving Mice. Cell Rep. 2016.

45. Pais-Vieira M, Chiuffa G, Lebedev M, Yadav A, Nicolelis MAL. Building an organic

computing device with multiple interconnected brains. Sci Rep. 2015;5:11869.

doi:10.1038/srep11869.

46. Biane JS, Takashima Y, Scanziani M, Conner JM, Tuszynski MH. Thalamocortical

Projections onto Behaviorally Relevant Neurons Exhibit Plasticity during Adult Motor

Learning. Neuron. 2016;89:1173–9. doi:10.1016/j.neuron.2016.02.001.

47. Froemke RC, Merzenich MM, Schreiner CE. A synaptic memory trace for cortical

receptive field plasticity. Nature. 2007;450:425–9. doi:10.1038/nature06289.

48. Liu CL, Gao M, Jin GZ, Zhen X. GABA Neurons in the Ventral Tegmental Area

Responding to Peripheral Sensory Input. PLoS One. 2012;7:1–8.

49. Nair-Roberts RG, Chatelain-Badie SD, Benson E, White-Cooper H, Bolam JP, Ungless

MA. Stereological estimates of dopaminergic, GABAergic and glutamatergic neurons in the

ventral tegmental area, substantia nigra and retrorubral field in the rat. Neuroscience.

2008;152:1024–31. doi:10.1016/j.neuroscience.2008.01.046.

50. Sesack SR, Grace AA. Cortico-Basal Ganglia reward network: microcircuitry.

Neuropsychopharmacology. 2010;35:27–47. doi:10.1038/npp.2009.93.

51. Wang H-L, Qi J, Zhang S, Wang H, Morales M. Rewarding Effects of Optical Stimulation

of Ventral Tegmental Area Glutamatergic Neurons. J Neurosci. 2015.

52. Gorelova N, Mulholland PJ, Chandler LJ, Seamans JK. The glutamatergic component of

the mesocortical pathway emanating from different subregions of the ventral midbrain. Cereb

Cortex. 2012;22:327–36. doi:10.1093/cercor/bhr107.

53. Yoo JH, Zell V, Gutierrez-Reed N, Wu J, Ressler R, Shenasa MA, et al. Ventral tegmental

area glutamate neurons co-release GABA and promote positive reinforcement. Nat Commun.

2016;7 May:1–13. doi:10.1038/ncomms13697.

54. Stuber GD, Hnasko TS, Britt JP, Edwards RH, Bonci A. Dopaminergic Terminals in the

Nucleus Accumbens But Not the Dorsal Striatum Corelease Glutamate. J Neurosci.

2010;30:8229–33. doi:10.1523/JNEUROSCI.1754-10.2010.

55. El Mestikawy S, Wallén-Mackenzie Å, Fortin GM, Descarries L, Trudeau L-E. From

glutamate co-release to vesicular synergy: vesicular glutamate transporters. Nat Rev Neurosci.

2011;12:204–16. doi:10.1038/nrn2969.

56. Hnasko TS, Hjelmstad GO, Fields HL, Edwards RH. Ventral Tegmental Area Glutamate

Neurons: Electrophysiological Properties and Projections. J Neurosci. 2012;32:15076–85.

doi:10.1523/JNEUROSCI.3128-12.2012.

57. Mylius J, Happel MFK, Gorkin AG, Huang Y, Scheich H, Brosch M. Fast transmission

from the dopaminergic ventral midbrain to the sensory cortex of awake primates. Brain Struct

Funct. 2015.

58. Phillipson OT, Kilpatrick IC, Jones MW. Dopaminergic innervation of the primary visual

cortex in the rat, and some correlations with human cortex. Brain Res Bull. 1987;18:621–33.

59. Lewis DA, Campbell MJ, Foote SL, Goldstein M, Morrison ' JH. The Distribution of

Tyrosine Hydroxylase-lmmunoreactive Fibers in Primate Neocortex Is Widespread but

Regionally Specific. J Neurosci. 1987;7 l:279–90.

60. Campbell AD, Kohl RR, McBride WJ. Serotonin-3 receptor and ethanol-stimulated

somatodendritic dopamine release. Alcohol. 1996.

61. Lidow MS, Goldman-Rakic PS, Gallager DW, Rakic P. Distribution of dopaminergic

receptors in the primate cerebral cortex: Quantitative autoradiographic analysis using

[3H]raclopride, [3H]spiperone and [3H]SCH23390. Neuroscience. 1991;40:657–71.

doi:10.1016/0306-4522(91)90003-7.

62. Williams SM, Goldman-Rakic PS. Characterization of the dopaminergic innervation of

the primate frontal cortex using a dopamine-specific antibody. Cereb Cortex. 1993.

63. Blundon JA, Zakharenko SS. Presynaptic Gating of Postsynaptic Synaptic Plasticity at

thalamocortical synapses: a plasticity filter in the adult auditory cortex. Neurosci.

2013;19:465–78. doi:10.1177/1073858413482983.

64. Izhikevich EM. Solving the distal reward problem through linkage of STDP and dopamine

signaling. Cereb Cortex. 2007.

65. Froemke RC, Carcea I, Barker AJ, Yuan K, Seybold BA, Martins ARO, et al. Long-term

modification of cortical synapses improves sensory perception. Nat Neurosci. 2013;16:79–88.

66. Sohoglu E, Chait M. Detecting and representing predictable structure during auditory

scene analysis. Elife. 2016;5. doi:10.7554/eLife.19113.

67. Kato HK, Gillet SN, Isaacson JS. Flexible Sensory Representations in Auditory Cortex

Driven by Behavioral Relevance. Neuron. 2015;88:1027–39.

doi:10.1016/j.neuron.2015.10.024.

68. Hull CL. Principles of Behavior: An Introduction to Behavior Theory. 1944.

69. Radtke-Schuller S, Schuller G, Angenstein F, Grosser OS, Goldschmidt J, Budinger E.

Brain atlas of the Mongolian gerbil (Meriones unguiculatus) in CT/MRI-aided stereotaxic

coordinates. Brain Struct Funct. 2016;:1–272. doi:10.1007/s00429-016-1259-0.

70. Saldeitis K, Happel MFK, Ohl FW, Scheich H, Budinger E. Anatomy of the auditory

thalamocortical system in the mongolian gerbil: Nuclear origins and cortical field-, layer-, and

frequency-specificities. J Comp Neurol. 2014;522:2397–430. doi:10.1002/cne.23540.

71. Mitzdorf U. Current source-density method and application in cat cerebral cortex:

investigation of evoked potentials and EEG phenomena. Physiol Rev. 1985;65:37–100.

http://www.ncbi.nlm.nih.gov/pubmed/3880898. Accessed 9 Feb 2016.

72. Harding GW. The currents that flow in the somatosensory cortex during the direct cortical

response. Exp Brain Res. 1992;90:29–39. doi:10.1007/BF00229253.

Figure 1 - Anatomical validation of fiber position relative to target side and optogenetic intracranial self-stimulation

(A) Schematic overview of fiber implant position for optogenetic stimulation of the VTA, and the ipsilateral AI recording side. Dashed boxes indicate representative regions of interest shown in B) identified by yellow and red asterisks. (B) Top, YFP expression in the temporal cortex covering the region of AI after virus transduction of the VTA indicates fiber terminals from the VTA targeting mainly upper layers and collaterals in deeper layers (white arrows). YFP expression was detectable in multiple regions of the neocortex (not shown). Bottom, co-fluorescence of YFP (AAV-CamKIIα-C1V1 (E162T)-p2A-eYFP) and TH-immunostaining (Alexa Fluor 546). Area of co-fluorescence is indicative of overlapping neuronal populations expressing the virus and TH mainly found in the VTA, and not the substantia nigra [18]. Scale bars indicate 250 and 200 μm, respectively. (C) Schematic representation of optogenetic

intracranial self-stimulation: While nose-poking the lever, animals induce a light flash towards

the VTA resulting in extensive self-stimulating behavior [19]. (D) Gerbil skull with prominent

landmarks (bregma, lambda and occipital crest) for anatomical reference [modified from 69].

For experiments, bregma has been used for reference of medio-lateral (ML) and anterior-

posterior (AP) position of the VTA. (E) Estimated actual AP/ML coordinates with mean

coordinates (AP:-3.57 mm; ML: 584.14 μm) relative to bregma of the C1V1group (n=7) as

estimated with the gerbil brain atlas [69]. (F) Fiber distance (mean ± SEM) towards area of co-

fluorescence (146.8 ± 38.9 μm; n=7). Compare with bottom picture of B). (G) Averaged lever

pressing rates (lever presses/ min) of the C1V1 (n=12) and YFP (n=8) groups over 10 days. A

representative video of a C1V1 animal is accessible as Movie (See Additional file 1).

Figure 2 - Electrophysiological measurements and effects of VTA stimulation on tone-evoked CSD profiles in A1

(A) Schematic representation of the timeline of measurements. Pre baseline measurements (Tone stimulation: 0.125-32 kHz, tone duration: 200 ms, ISI: 0.6-0.8, 50 pseudorandomized repetitions, 65 dB SPL, 7.5 min) were performed until CSD patterns had stabilized (45-75 min). A single measurement of combined tone and laser stimulation (25 Hz, 473 nm, 10 mW) was carried out and followed by a series of post-measurements (up to 60 min). (B) Representative example of a CSD profile after BF- (top) and off-BF (bottom) stimulation before, during and 30 minutes after the combined tone/VTA-laser measurement from an animal of the C1V1 group. Before VTA stimulation, BF-evoked CSD profiles revealed a canonical feedforward pattern of stimulus-evoked current flow starting with initial sink components in cortical layers III/IV and Vb/VIa. Subsequent sink activity was found in layers Va, I/II and VIb . Activity after off-BF stimulation showed a less pronounced columnar activation in the pre-condition. During paired tone and VTA stimulation, both stimulation frequencies evoked stronger and prolonged translaminar current flow, which was still enhanced 30 minutes later. Cortical layers are indicated. Dashed vertical lines indicate tone onset and offset.

Figure 3 - Binned time courses of single-trial sink peak amplitude data of layer-specific CSD-traces

Temporal development of single-trial sink peak amplitudes for early sink activity (sink onset <50 ms) in layers III/IV, Va and Vb/VIa and late sinks (sink onset >50 ms) in layers I/II and VIb for BF, near- and non-BFs. Layer I/II peak amplitudes showed a significant increase for the C1V1 group during laser and in post-measurements, surpassing the +10% criterion within

the BF and near-BF bins. Significant differences from the control group are again revealed by

LME models and indicated by asterisks. Layer I/II peak amplitudes in the YFP and control

group did not show corresponding increases over the recording time but showed a decrease and

recovery for the YFP group. Layer III/IV activity did not reveal differences over time for BF-

evoked responses in all groups. With spectral distance from the BF the C1V1 group showed a

moderate increase yielding significance at the >30 min time bin compared to control animals.

Within the non-BF bin, peak amplitudes showed a significant decrease in the YFP group during

and after laser stimulation. Layer Va peak amplitudes were most stable across time and groups

only displaying a significant decrease for the YFP group in the non-BF bin at <30 min. Layer

Vb/VIa peak amplitudes in the C1V1 group were significantly increased in all post-

measurements after BF-stimulation. For stimulation with near-BF a similar trend was found.

Significant changes between the YFP and control group were again due to decreased peak

amplitudes in the YFP group. Layer VIb displayed a highly significant increase in the C1V1

group during laser stimulation, most prominent in the non-BF bin, and less prominent in near-

BF and BF bins. Laser-induced increase only persisted for the next time bin, but then recovered

to pre-condition. While the control group showed stable peak amplitudes over the time course,

YFP animals again showed a moderate decrease most prominent in off-BF bins.

Figure 4 - Layer-specific early and late changes of tuning sharpness of the C1V1 group

For data from the C1V1 group, peak amplitudes within each layer were normalized to BF-responses before laser stimulation. We fitted a linear line across evoked responses across the BF-, near-BF- and non-BF-bin and calculated the slope as indicator for spectral tuning sharpness (for pre, laser, >30 min). Sharpest tuning properties of peak amplitudes were found in layers III/IV (slope of 0.28). Tuning in layers III/IV was not affected by VTA stimulation. Layers I/II showed less prominent tuning revealed by a shallower slope in the pre-condition. While VTA stimulation did not yield an immediate significant difference, slopes indicated a significantly sharper tuning >30 min after VTA stimulation (p< 0.01). This was due to mainly an increase in BF-evoked peak amplitudes in accordance with Fig. 5. Peak amplitudes in infragranular layers Vb/VIa showed a similar effect as layers I/II with a trend of sharper tuning during VTA stimulation that yielded significance >30 min later. In contrast, only in late infragranular layer VIb, peak amplitudes revealed a significantly shallower slope indicative of a broader frequency tuning during laser stimulation (Pre:laser; p<0.01), which recovered after >30 min (Pre:<30 min; p=0.1). This can be seen by the flat slope between non- and BF bins. Layer Va only showed a minor decrease of the slope during VTA stimulation that recovered partially >30 min later. Significances indicated are based on slope comparisons performed using the lsmeans package in R (version 2.27-62). For statistical explanation, see text.

Figure 5 - Temporal development of early and late AVREC and ResidualCSD activity

(A) Grand average traces of the BF-evoked AVREC (top) and ResidualCSD (bottom) of the C1V1 group (n=12) for binned pre, laser and post (<30, >30 &>45 min)-measurements. Right, insets illustrate that AVREC activity is mainly associated with local columnar activity, while the ResidualCSD quantifies contributions from lateral corticocortical input (cf. Happel et al., 2014, 2010). (B) Temporal time courses of Pre-normalized early/late AVREC and early/late ResidualCSD traces of C1V1 (red), control (blue) and YFP (green) groups (n=12, n=7, n=7, respectively) shown for BF stimulation. Early AVREC signal displayed a significant increase

in post-measurements for the C1V1 group whereas time courses of the control and YFP groups

were stable. Late AVREC signal showed no significant differences between either group. RMS

ResidualCSD values of early and late time windows displayed a similar temporal development

as the early AVREC becoming significantly increased in the later post-measurements within

the C1V1 group. Control and YFP groups showed no significant changes over time. Note the

peaking in the early ResidualCSD RMS during laser stimulation, which is only present in the

C1V1 group. Asterisks mark significant changes according to the applied mixed linear models

of the normalized data towards the control group (see Materials and Methods). (C) Tuning

curves of the RMS value normalized to BF-evoked responses in the pre-condition for early/late

AVREC and early/late ResidualCSD for binned time points for C1V1 group. Note the general

increase of the overall tuning curve relative to pre measurements.

Figure 6 - Normalized single-trial early and late AVREC and ResidualCSD RMS values over time

(A) Temporal time course of averaged early single-trial RMS AVREC (top) and ResidualCSD (bottom) values normalized to pre-measurements. Data for BF, near-BF and non-BF stimulation is shown separately and was binned for pre-measurements, laser, and post-measurements (<30 min, >30, >45 min). Note the prominent increase of early RMS AVREC values for the C1V1 group whereas control and YFP group displayed a more stable or decreasing time course,

respectively. Also, early RMS ResidualCSD values significantly increased during and after the

Laser measurements for BF, near- and non-BF bins of the C1V1 group, whereas YFP and

control display a more stable time course. (B) Temporal time course of late Pre-normalized

RMS AVREC and ResidualCSD values of single-trial data for BF, near-BF and non-BF. Late

RMS AVREC values for the C1V1showed a mild, yet not significant, increase that did not

surpass the +10% criterion. The control group displayed a stable and slightly declining time

course for the late RMS AVREC values, while the YFP group showed a moderate decrease that

recovered over time. Late RMS ResidualCSD values increased during and after the Laser

measurements for BF, near- and non-BF bins of the C1V1 group, whereas YFP and control

groups both showed a more stable time course. Significances in A) and B) were calculated using

mixed linear models on the normalized data. Asterisks indicate corresponding significances in

comparison to the control group, in case values exceed the ± 10% pre-criterion. Single-trial

CSD data in following figure is presented equally.

Figure 7 - Schematic illustration of convergent deep layer inputs from sensory thalamus

and VTA.

VTA projections mainly target upper and infragranular layers of the gerbil auditory cortex (see

also Fig. 1 B). Stimulating the VTA projections led to a frequency-specific gain amplification

of early sensory-evoked responses in thalamocortical recipient layers Vb/VIa. This gain

increase effectuated an enhanced spectral tuning representation in supragranular layers I/II. The

supragranular gain modulation might hence be inherited via intratelencephalic neurons

targeting upper layers and bypassing granular input circuits [36], may originate from a direct

influence of dopamine released in upper layers or a combination of both. MGB, medial

geniculate body.

Additional files:

Additional file 1- Optogenetic intracranial self-stimulation induced by lever pressing

Animal video clip of an intracranial self-stimulation training session. Animals were placed into a 25x35x22cm³ Plexiglas box for 20 min once a day. Optogenetic implants were attached via an optical fiber with a shutter-controlled laser (10 mW, 473 nm). By pressing an installed lever, animals were able to trigger a laser pulse (25 Hz, 5 ms) for VTA stimulation. Highly addicted animals (>= 850 lever presses per session), were kept for optogenetic experiments whereas non-addicted animals were rejected from this study (Compare behavior of the animal in the background).

Additional file 2- Assignment of layers to CSD channels and semi-automatic detection of

sink-activity

Based on previous studies, earliest tone-evoked components are related to layers Vb/VIa and

III/IV [15, 22]. Subsequent synaptic activity is conveyed to layer Va and VIb, and layers I/II,

as indicated in the representative example. Channels assigned to individual layers were

averaged and activity attributed to sources (positive values) were dismissed. For visual

purposes, a corresponding average-sink trace from layer III/IV was inverted. Based on the pre-

stimulus baseline signals (200 ms), the baseline mean value + 1 standard deviation (STD) was

calculated as response threshold for each layer-dependent CSD trace. A deflection that

surpassed a 1-fold STD was used to determine the onset and offset of early and late sink

activations (see Materials and Methods). Significant sink onset activity within the first 50 ms

after tone onset was assigned as early activity, while later onsets were assigned as late activity.

Additional file 3- Grand average AVREC and ResidualCSD traces for YFP and control groups

(A) AVREC and ResidualCSD traces of the YFP group (n=7) for binned pre-measurements, laser-measurement, post-measurements < 30 min, post-measurements > 30 min and post-measurements > 45 min. Dashed lines indicate tonal on- and offset. Areas for early and late RMS calculations are underlined with gray. Group sorting was performed according to the BF of the granular sink peak amplitudes. (B) AVREC and ResidualCSD traces of the control group

896   (n=7) for binned pre-measurements, laser-measurement, post-measurements < 30 min, post-

897   measurements > 30 min and post-measurements > 45 min. Dashed lines indicate tonal on- and

898   offset. Areas for early and late RMS calculations are underlined with gray. Group sorting was

899   performed according to the BF of the granular sink peak amplitudes.

900