A NOVEL PHOX/CD38/MCOLN1/TFEB AXIS IMPORTANT FOR MACROPHAGE ACTIVATION DURING BACTERIAL PHAGOCYTOSIS

Mehran Najibi, Joseph A. Moreau, Havisha H. Honwad, Javier E. Irazoqui*

Department of Microbiology and Physiological Systems and Program in Innate Immunity, University of Massachusetts Medical School, Worcester MA 01605

*Correspondence: javier.irazoqui@umassmed.edu

Word count: 9188 including Methods, 7623 without Methods

Keywords: TFEB, innate immunity, inflammation, infection, phagocyte, transcription, cytokines, IL-6, IL-1α, IL-1β

Abstract

Macrophages are a key and heterogenous class of phagocytic cells of the innate immune system, which act as sentinels in peripheral tissues and are mobilized during infection. Macrophage activation in the presence of bacterial cells and molecules entails specific and complex programs of gene expression. How such triggers elicit the gene expression programs is incompletely understood. We previously discovered that transcription factor TFEB is a key contributor to macrophage activation during bacterial phagocytosis. However, the mechanism linking phagocytosis of bacterial cells to TFEB activation remained unknown. In this article, we describe a previously unknown pathway that links phagocytosis with the activation of TFEB and related transcription factor TFE3 in macrophages. We find that phagocytosis of bacterial cells causes an NADPH oxidase (PHOX)-dependent oxidative burst, which activates enzyme CD38 and generates NAADP in the maturing phagosome. Phago-lysosome fusion brings Ca^{2+} channel TRPML1/MCOLN1 in contact with NAADP, causing Ca^{2+} efflux from the lysosome, calcineurin activation, and TFEB nuclear import. This drives TFEB-dependent expression of important pro-inflammatory cytokines, such as IL-1α, IL-1β, and IL-6. Thus, our findings reveal that TFEB activation is a key regulatory event for the activation of macrophages. These findings have important implications for infections, cancer, obesity, and atherosclerosis.

Introduction

Macrophages are phagocytic cells of the innate immune system, which act as

sentinels in peripheral tissues and are mobilized during infection [1]. Macrophages

are activated in the presence of bacteria, which entails specific programs of gene

expression. Depending on the stimulus and the microenvironment, particularly

the pathogen molecules and cytokines present, macrophages adopt phenotypes

(or "polarize") along a spectrum, from "classical" activation (a.k.a. M1 polariza-

tion), which is predominantly pro-inflammatory, to "alternative" activation (a.k.a.

M2 polarization), which is predominantly pro-resolution [2]. The activation state of

the macrophage has great functional consequences for health and disease.

Because transcriptional regulation contributes greatly to the macrophage phe-

notype, understanding the transcription factors that are involved is of paramount

importance. Similarly, it is imperative to fully elucidate the signaling pathways

that regulate such transcription factors under conditions of homeostasis and dis-

ease. Much work has characterized important signaling from the Toll-like recep-

tors (TLRs), which recognize molecules produced by pathogens such as bacteri-

al cell wall components and trigger the activation of transcription factors NF-κB,

IRF3, and AP1 [3]. All three examples of transcription factors are subject to multi-

ple regulatory layers, nuclear-cytoplasmic shuttling being a major mechanism of

regulation.

We recently identified TFEB as a transcription factor that is important for cyto-

kine and chemokine gene induction in macrophages following bacterial infection

[4]. We found that TFEB resides in the cytosol in resting murine macrophages, but

is imported to the nucleus after short infection with *Staphylococcus aureus* or

Salmonella Typhimurium [4,5]. Moreover, cells defective in TFEB expression exhibit

defective induction of important pro-inflammatory cytokines, including TNF-α, IL-

1β, and IL-6 [4]. Subsequent studies confirmed our initial observations, and ex-

tended them by showing that TFE3 is also activated by lipopolysaccharide (LPS),

a cell wall constituent of Gram-negative bacteria [6]. Moreover, an independent

study demonstrated how phagocytosis of opsonized particles triggers TFEB nu-

clear import and the subsequent activation of bactericidal mechanisms in murine

macrophages [7]. Thus, clearly activation of TFEB is an important regulatory event

in macrophage responses to bacterial pathogens. However, the regulatory

mechanisms upstream of TFEB during bacterial infection in macrophages re-

mained unclear.

We recently reported a novel pathway that is required for TFEB activation dur-

ing bacterial infection in macrophages [5]. In it, phosphatidyl choline-directed

phospholipase C (PC-PLC), which produces diacyl glycerol (DAG), activates pro-

tein kinase D1 (PrKD1) to promote TFEB nuclear import. While in the absence of

this pathway TFEB cannot be activated by infection, its induction is sufficient to

cause TFEB nuclear import. However, the molecular mechanism linking PrKD1

to TFEB remained unknown. Importantly, the link between phagocytosis of bacte-

ria and PC-PLC/PrKD1 activation remained unclear.

In the present article, we describe the discovery of a novel pathway that links

phagocytosis of bacterial cells to TFEB activation and downstream cytokine ex-

pression. Instead of the expected TLR signaling pathways, we found that signal

transduction for TFEB activation requires NADPH oxidase (PHOX) and the pro-

duction of reactive oxygen species (ROS). This event sets in motion the activa-

tion of CD38, which produces NAADP, an activating ligand for lysosomal Ca^{2+}

export channel TRPML1/MCOLN1. TRPML1/MCOLN1 activation, followed by

Ca^{2+} export, drives calcineurin, a Ca^{2+}-regulated protein phosphatase that

dephosphorylates TFEB on conserved S/T residues critical for TFEB cytosolic

sequestration [8], thus allowing TFEB nuclear import and downstream gene induc-

tion.

Because of the broad tissue expression of CD38 and other components of this

pathway [9], and because several stresses and diseases cause potentially TFEB-

activating ROS production from mitochondria [10–12], the findings reported here are

likely to have broad implications for the regulation of gene expression down-

stream of TFEB in multiple tissues and physiological conditions, including home-

ostasis and inflammatory diseases.

Results

Phagocytosis of Gram+ and Gram- bacteria activates TFE3 in macro-phages

We previously showed that *S. aureus* and *S.* Typhimurium could activate TFEB in murine macrophages [5]. Since then, several studies have suggested that TFE3 and TFEB share many features of upstream regulation [13–15]. In addition, previous work showed that extended incubation with LPS activates TFE3 in macrophages [6]. Therefore, we wondered if TFE3 might also respond to infection by bacteria. We found that infection with *S. aureus* or *S.* Typhimurium activated TFE3 with similar kinetics and amplitude as TFEB, measured by its nuclear accumulation (**Fig. 1A-H** and **S1A-C, F**). Moreover, peptidoglycan (PGN) from *S. aureus* alone was sufficient to induce TFE3 and TFEB nuclear accumulation (**Fig. 1C, F-H**), suggesting that phagocytosis of large particles was not a requirement. Both *S. aureus* and PGN also induced the formation of lysosomes, measured with Lysotracker staining (**Fig. 1I-R**). There was a striking correlation between nuclear accumulation of TFEB and lysosome induction, indicating that *S. aureus-* or PGN-induced TFEB activation was biologically relevant.

Salmonella infection produced a more nuanced response. As we previously showed, live and dead *Salmonella* differed in their ability to induce TFEB activation [5]. While live *Salmonella* induced rapid activation of TFEB and TFE3, dead *Salmonella* did so much more slowly (**Fig. S1A-D, F, G**). Interestingly, the kinet-

ics of TFEB and TFE3 activation by dead *Salmonella* resembled that of LPS,

suggesting that they might trigger TFEB/3 activation through a similar sensing

mechanism (**Fig. S1E, H**). In contrast to *S. aureus, Salmonella* did not induce the

accumulation of lysosomes, but rather decreased it below baseline (**Fig. S2A**).

LPS induced lysosomal accumulation, as PGN did (**Fig. S2C**). Thus, we suspect

that *Salmonella* possesses a mechanism to inhibit lysosomal biogenesis down-

stream of TFEB/3. This is consistent with previous results showing lysosomal in-

hibition by *Salmonella* [16,17].

The overall conclusion of these observations is that bacterial infection and

bacterial ligand stimulation of murine macrophages causes biologically significant

TFEB and TFE3 activation. However, the functional consequences of bacterially-

induced TFEB and TFE3 nuclear import were unclear.

TFEB and TFE3 are key contributors to the macrophage transcriptional

response to infection

To better understand the functional consequences of TFEB and TFE3 activa-

tion during bacterial infection, we performed RNAseq of a RAW264.7 murine

macrophage cell line harboring deletions in *Tfeb* and *Tfe3.* We detected induction

of 1,022 genes after 3 h of *S. aureus* infection in wild type macrophages (**Table**

S1, Fig. 2A). Compared to wild type cells, *Tfeb*[-/-] *Tfe3*[-/-] (double knockout, or

dKO) cells exhibited a drastically altered transcription profile (**Fig. 2A, C**). Over-

all, about two-thirds of *S. aureus*-induced genes in wild type macrophages were

not induced in dKO cells (**Fig. 2D**), indicating that a large majority of the tran-

scriptional response to *S. aureus* was TFEB/3-dependent. In addition, while wild

type macrophages induced *Ccl5, Nos2, Ptgs2,* and *Tnf,* indicating a classically-

activated state, dKO cells were completely defective in their induction (**Fig. 2B**).

This revealed that macrophage polarization to the classically activated state, or

M1, requires TFEB and/or TFE3.

A large group of innate immunity genes exhibited decreased expression in

dKO cells (**Fig. 2C**). Noteworthy examples included TLR genes *Tlr6, Tlr9,* and

Tlr13, NF-κB transcription factor genes *Nfkb2* and *Rel,* NF-κB inhibitor genes

Nfkbib and *Nfkbie,* and NLR genes *Nod1, Nod2, Naip2,* and *Naip5.* In addition,

several cytokine genes exhibited defective expression, including *Tnf, Il1b, Il6,*

Ifnb1, Tgfb1, Csf1, Csf2, and Csf3, and various interleukins (e.g. *Il23a, Il27,* and

Il33), as well as chemokines and chemokine receptors such as *Ccl3, Ccl6, Ccl7,*

Ccl12, Ccl17, Ccl22, Cxcl3, Cxcl10, Cxcl11, Ccrl2, Cx3cr1, and signaling compo-

nents such as *Ripk2, Jak3, Stat1, Stat2,* and *Stat5a.* Moreover, several autopha-

gy and lysosomal genes showed lower expression: Cathepsins (lysosomal prote-

ases *Ctsa, Ctsd, Ctse, Ctsz*), *Sqstm1, Mcoln1* and *Mcoln2, Atg7, Atg13, Irgm1*

and *Irgm2,* and *Map1lc3a* (LC3)*.* Finally, expression of ER unfolded protein re-

sponse transcription factor gene *Xbp1* and necroptosis effector gene *Mlkl* were

also reduced in dKO macrophages. These results strongly suggest that TFEB/3

are important for many key functions of macrophages, including pathogen recog-

nition, pro-inflammatory signaling, secretion of chemotactic signals, and autoph-

agy/lysosomal clearance.

More systematically, analysis of over-represented Reactome pathways [18] re-

vealed that the top affected categories were "Cytokine Signaling in Immune Sys-

tem", "Signaling by Interleukins", and "Innate Immune System". Other significant

categories affected were "Interferon signaling", "ER-Phagosome Pathway", and

various TLR pathways (**Fig. 2E**). Consistent with the transcriptome measure-

ments, we observed robust secretion of IL-6 in *S. aureus*-infected wild type mu-

rine macrophages, which was abrogated by deletion of *Tfeb* alone (**Fig. 2F, H**).

Wild type cells also secreted more IL-1β and TNF-α than *Tfeb*$^{\Delta LysM}$ macrophages,

even though in this case the secreted levels were noisier than IL-6 (**Fig. 2H, J**).

IL-1α exhibited a similar trend to IL-1β (**Fig. 2G**). Lactate dehydrogenase (LDH)

release assays ruled out that these differences were due to cytotoxicity (**Fig. 2K**).

Together, these observations provide strong evidence that TFEB/3 are essential

for the macrophage transcriptional response to *S. aureus* infection, for macro-

phage polarization and function, and for the production of pro-inflammatory cyto-

kines and chemokines. They also suggest that *Tfeb* deletion alone is sufficient to

confer a defect in cytokine production, consistent with previous results [6]. There-

fore, nuclear translocation of TFEB and TFE3 is a key regulatory event for the

macrophage response to infection. However, the signaling events upstream of

TFEB/3 activation remained undefined.

Bacterial infection activates TFEB and TFE3 independently of TLR signaling

Because TLRs constitute a major mechanism of innate recognition of bacterial ligands, we investigated the roles of TLR signaling in TFEB activation by bacterial cells and ligands. MyD88 is a protein adaptor that is important for signaling from TLR2/6 and TLR4, which detect PGN and LPS, respectively [19]. TRIF is an adaptor that is also important for signaling from endosomal TLR4 [20].Together, MyD88 and TRIF are essential for TLR signaling. Therefore, we examined TFEB activation in murine macrophages lacking both MyD88 and TRIF.

Much to our surprise, after infection with *S. aureus* and treatment with PGN *Myd88*[-/-] *Trif*[-/-] macrophages exhibited the same levels of TFEB activation as wild type cells (**Fig. 3**). Similarly, after infection with *Salmonella* or treatment with TLR4 agonists LPS or MPLA we observed no difference between *Myd88*[-/-] *Trif*[-/-] and wild type macrophages (**Fig. S2**). These results suggested that TLR signaling is dispensable for TFEB activation by bacterial cells and ligands, regardless of cell wall structure.

To test this conclusion further, we examined TFEB activation in macrophages deficient for TLR2 and TLR4 (**Fig. S3A, B**). After infection with *S. aureus* and PGN or Pam3Csk4 treatment, *Tlr2*[-/-] macrophages exhibited the same activation of TFEB as wild type cells (**Fig. S3C-L**). Likewise, after infection with live *Salmo-*

nella or treatment with LPS, TFEB activation in *Tlr4*[-/-] and wild type macrophages

was indistinguishable; MPLA still activated TFEB in *Tlr4*[-/-] cells, albeit more weak-

ly (**Fig. S3M-X**). Together, these results indicated that bacterial cells and ligands

activate TFEB by a TLR-independent pathway.

Bacterial activation of TFE3 requires the PC-PLC/PrKD1 pathway

Given this lack of relevance of TLR signaling, we sought alternative mecha-

nisms of TFE3 activation. We previously showed that TFEB activation in macro-

phages infected with *S. aureus* or *Salmonella* required the activity of a pathway

upstream of PrKD1 [5]. We found that prior inhibition of PC-PLC with D609 or of

PrKD with kb-NB142-70 completely abrogated TFE3 activation by *Salmonella,*

LPS, and *S. aureus,* similar to TFEB (**Fig. S4A-N**). Furthermore, we observed

strong TFE3 activation by the DAG analog PMA, which was prevented by PrKD

inhibition (**Fig. S4O, P**). Together, these data indicate that the PC-PLC/PrKD1

pathway is necessary and sufficient for TFE3 activation during infection in mac-

rophages, as we previously showed for TFEB. However, the connection between

PrKD1 and TFEB (or TFE3) nuclear import remained unclear. More importantly,

the mechanism connecting phagocytosis of bacteria to TFEB/3 activation re-

mained unknown. Therefore, we decided to examine regulatory events more

proximal to TFEB.

Bacterial activation of TFEB requires the TRPML1/MCOLN1 – Ca^{2+} –

calcineurin pathway

The Ca^{2+}-dependent protein phosphatase calcineurin was recently shown to be principally responsible for TFEB nuclear import induced by starvation in HeLa, HEK293, and human fibroblasts, as well as mouse muscle and MEFs. Such effect was mediated by its dephosphorylation of mTORC1 target sites in TFEB [21]. However, the role of calcineurin in TFEB control in infected macrophages has not been tested before. Therefore, we evaluated TFEB activation in infected macrophages lacking calcineurin function. Prior incubation with Ca^{2+}-chelating agent BAPTA prevented TFEB activation by *S. aureus* (**Fig. 4A-E**), suggesting that intracellular Ca^{2+} is important. Moreover, prior treatment with calcineurin inhibitor FK506 prevented *S. aureus*-triggered TFEB activation even more effectively (**Fig. 4D, E**), suggesting that calcineurin activity was required. Furthermore, siRNA-mediated silencing of genes *Ppp3cb,* which encodes calcineurin catalytic subunit β, or *Ppp3r1,* which encodes calcineurin regulatory subunit Bα, abrogated TFEB activation by *S. aureus* (**Fig. 4F-L**). These experiments indicate that calcineurin is required for TFEB activation during *S. aureus* phagocytosis. We found similar results during infection with *Salmonella* (**Fig. S5A-L**), indicating that calcineurin is a general requirement for TFEB activation during infection of macrophages. Furthermore, the Ca^{2+} ionophore ionomycin was sufficient to trigger TFEB nuclear import in uninfected macrophages (**Fig. 4M-O**). Thus, activation of calcineurin is necessary and sufficient for TFEB nuclear import in macrophages during infection with *S. aureus* or *Salmonella.*

Previous studies have identified TRPML1/MCOLN1 as a major Ca^{2+} export

channel in lysosomes [22]. Furthermore, TRPML1/MCOLN1 was shown to function

upstream of calcineurin for TFEB activation in several human and mouse cell

types during nutrient starvation, and TRPML1/MCOLN1 was required for TFEB

activation by FcγR during phagocytosis of opsonized beads in murine macro-

phages [7,21]. In our hands, silencing of *Mcoln1* completely prevented TFEB activa-

tion and lysosome induction by *S. aureus* (**Fig. 4P-T**) and *Salmonella* (**Fig. S5M-**

Q). Conversely, treatment of uninfected macrophages with TRPML1/MCOLN1

agonist ML-SA1 [23] resulted in strong TFEB nuclear import (**Fig. 4U-W**). Thus, in

macrophages TRPML1/MCOLN1 is necessary and sufficient for TFEB activation

during infection. Together, these results suggested that the TRPML1/MCOLN1-

calcineurin-TFEB pathway is functional in macrophages, and is necessary and

sufficient for TFEB nuclear import during infection. However, the mechanism of

TRPML1/MCOLN1 activation during infection remained unknown.

CD38 functions upstream of TRPML1/MCOLN1 for TFEB activation

Because of its relevance to lysosomal storage disorders, in particular

mucolipidosis Type IV, the regulation of TRPML1/MCOLN1 is of great interest [22].

Recently, it was reported that nicotinic acid adenine dinucleotide phosphate

(NAADP) can function as a TRPML1/MCOLN1 ligand to induce lysosomal Ca^{2+}

export in several cell types [24,25]. Therefore, we investigated whether NAADP

might function to induce TFEB activation downstream of MCOLN1 during infec-

tion in macrophages.

Prior treatment with the potent NAADP antagonist Ned-19 [26] rendered macro-

phages unable to translocate TFEB into the nucleus during infection with *S.*

aureus (**Fig. 5A-C, F**). Likewise, treatment with kuromanin or apigenin, which in-

hibit the NAADP-synthesizing enzyme CD38 [27,28], also prevented TFEB activa-

tion by *S. aureus* (**Fig. 5D-F**)*.* We obtained essentially the same results with

Salmonella (**Fig. S6A-F**). Moreover, silencing of *Cd38* prevented TFEB activation

by *S. aureus* (**Fig. 5G-J**). Together, these data showed that CD38 and NAADP

were required for TFEB activation during infection. Conversely, treatment of mac-

rophages with the NAADP analog NAADP-AM [29] resulted in strong induction of

TFEB nuclear import (**Fig. 5K, L, N**). Together, these observations showed that

CD38 and its product NAADP are necessary and sufficient for TFEB activation by

S. aureus in macrophages. In contrast, NAADP-AM-induced TFEB activation was

abrogated by silencing *Mcoln1* (**Fig. 5M, N**), suggesting that CD38 and NAADP

require TRPML1/MCOLN1 to induce TFEB translocation, consistent with their

proposed roles upstream of TRPML1/MCOLN1.

Immunofluorescence in untreated macrophages showed that CD38 localized

principally to the plasma membrane, as previously shown in several human lym-

phocyte cell types [30,31]. In contrast, *S. aureus*-infected macrophages exhibited

intracellular staining in what appeared to be cytosolic vesicles (**Fig. 5O, P**). Such

vesicles are presumably phagosomes [32]. We also noticed increased staining in

infected cells compared to baseline. To further examine this, we performed

immunoblots. In wild type macrophages, we observed a slight increase in CD38

expression, albeit of a smaller molecular weight form (45 kDa v 64 kDa), in in-

fected cells compared to control (**Fig. 5Q**). This increased expression occurred in

Tfeb[-/-] cells as well (**Fig. 5Q**). Together, these observations suggested that infec-

tion caused the relocalization of CD38 to endomembranes, its increased expres-

sion, and its presumed proteolytic conversion to a smaller form in a TFEB-

independent manner.

ROS function upstream of CD38 and of TFEB and TFE3

CD38 is a fascinating transmembrane enzyme with complex membrane topol-

ogy, subcellular localization, and regulation [31]. Upstream regulation of CD38 is

poorly understood. It is thought that CD38 can be activated by reactive oxygen

species (ROS) by an unknown mechanism [33,34]. Moreover, ROS activated

TRPML1/MCOLN1 in HEK293 cells, presumably through CD38 [35]. Therefore, we

investigated the role of ROS in TFEB activation during infection.

First, we verified the accumulation of ROS and H_2O_2 during infection with *S.*

aureus in wild type macrophages (**Fig. 6A, B**). To our surprise, prior treatment

with ROS-scavenging compounds N-acetyl cysteine (NAC) or N-acetyl cysteine

aldehyde (NACA) completely prevented TFEB nuclear translocation by *S. aureus*

(**Fig. 6C-G**) and *Salmonella* (**Fig. S6G-K**), suggesting that ROS were essential

for TFEB activation during infection. Conversely, treatment with CCCP, which

uncouples the mitochondrial electron transport chain and elicits ROS, caused

TFEB translocation in the absence of infection (**Fig. 6H, I, K**). Silencing of *Cd38*

completely abrogated TFEB translocation caused by CCCP (**Fig. 6J, K**). Togeth-

er, these data indicated that ROS, generated during infection with *S. aureus*, are

necessary and sufficient for TFEB activation by infection, in a mechanism that

requires CD38. However, the source of ROS during phagocytosis remained un-

clear.

NADPH oxidase functions upstream of the ROS-CD38-TRPML1/MCOLN1-

CN-TFEB axis

During phagocytosis, recruitment of NADPH oxidase (PHOX) to the maturing

phagosome is a key event for phagolysosome maturation and bactericidal killing

[36]. PHOX catalyzes the production of H_2O_2 from NADPH oxidation, thus initiating

the production of ROS in the phagolysosome [37]. Therefore, we examined PHOX

as a potential source of ROS for CD38 activation during bacterial phagocytosis.

First, we examined the oxidative burst in macrophages harboring a deletion in

Nox2, which encodes the gp91phox subunit. While wild type macrophages in-

fected with *S. aureus* induced high levels of ROS and superoxide, *Nox2*[-/-] cells

did not (**Fig. 7A, B**), consistent with PHOX being the main source of ROS in our

infection assays. Consistent with the requirement for ROS to activate TFEB dur-

ing infection, *S. aureus* did not induce TFEB nuclear translocation in *Nox2-*

deficient cells (**Fig. 7C-G**), suggesting that PHOX is required for TFEB activation

by infection. In support of this conclusion, chemical inhibition of PHOX activity

using apocynin completely prevented TFEB activation during *S. aureus* infection

(**Fig. H-K**). We observed similar results with *Salmonella* (**Fig. S6L-P**). Together,

these data suggested that PHOX functions upstream of TFEB, and that during

bacterial infection its activity is essential for TFEB activation.

Immunofluorescence analysis revealed that *S. aureus* infection caused NOX2

and CD38 colocalization to increase about 100% over baseline (**Fig. 7L-T**). We

obtained similar results with *Salmonella* (**Fig. S6Q-T**). Taken together, this series

of experiments supports a hypothetical model (**Fig. 7U**) in which phagocytosis of

bacterial cells may cause activated PHOX and CD38 to localize to the same

compartment, likely the phago-lysosome. ROS produced by PHOX may cause

CD38 to produce NAADP, which could activate TRPML1/MCOLN1 and induce

Ca^{2+} export from the phago-lysosome/lysosome compartment. Ca^{2+} thus re-

leased may activate calcineurin, which can dephosphorylate TFEB. Dephosphor-

ylated TFEB can translocate into the nucleus, where it drives the expression of

cytokines and chemokines, as previously demonstrated [4,6]. In addition, TFEB ac-

tivation results in enhanced autophagy and lysosomal biogenesis, thus amplify-

ing the phagocytic capacity of the cell via a positive feedback loop, and increases

the degradative capacity of the cell via a positive feedforward loop [7]. However,

this model did not account for the requirement of the PC-PLC/PrKD1 pathway for

TFEB activation.

Disruption of PC-PLC/PrKD1 signaling inhibits TRPML1/MCOLN1 locali-

zation to the lysosome

A previous study showed that PrKD1 is required for TRPML1/MCOLN1

transport from the Golgi apparatus to the lysosome in HeLa cells and human skin

fibroblasts [38]. We tested if this is also a functional mechanism in macrophages.

Using immunofluorescence in baseline murine macrophages, we observed low

expression and localization to LAMP1-positive vesicles of TRPML1/MCOLN1

(**Fig. S7A-C**). TRPML1/MCOLN1 expression was highly upregulated by treat-

ment with ML-SA1 and by infection with *S. aureus* (**Fig. SD-I, T**). Such

upregulation was expected, since *Mcoln1* is a direct TFEB target in macrophag-

es, and is upregulated by TFEB activation [6]. Under *S. aureus* infection and ML-

SA1 treatment, we also observed higher colocalization of TRPML1/MCOLN1 and

LAMP1 (**Fig. S7D-F, J-L, S**). *S. aureus* infection caused a 100% increase in

TRPML1/MCOLN1 – LAMP1 colocalization, which was partially abrogated by

prior treatment with PrKD1 inhibitor kb-NB142-70 (**Fig. S7M-O, S**). kb-NB142-70

also prevented the induced expression of TRPML1/MCOLN1 by *S. aureus* and

MLSA1 (**Fig. S7T**). Together, these observations support the notion that PrKD1

is necessary for TFEB activation because it is required for TRPML1/MCOLN1

transport to the phagolysosome/lysosome compartment, where it is required for

Ca^{2+} export downstream of NAADP.

Discussion

In this article, we show a novel axis for TFEB activation in macrophages during phagocytosis of bacterial cells, which is essential for macrophage activation. Our observations support a model for a novel mechanism of pathogen sensing that is TLR-independent and is based on the phagocytic pathway. This pathway connects productive phagocytosis, presumably sensed by the recruitment of PHOX to the maturing phagosome. PHOX recruitment thus places the source of ROS in the same compartment as CD38, which may be endocytosed during phagocytosis of the bacterial particle. ROS generated by PHOX may activate CD38, as reported for lymphokine-activated killer cells and coronary arterial myocytes [39,40], causing the accumulation of NAADP in the phagosome. After phago-lysosome fusion, TRPML1/MCOLN1 may come into contact with this reservoir of NAADP, thus triggering the release of lysosomal Ca^{2+} into the cytosol. In the cytosol, Ca^{2+} activates calcineurin, which dephosphorylates TFEB and enables its nuclear import. In the nucleus, TFEB drives the expression of a large majority of genes that are induced upon pathogen phagocytosis. Such gene induction may require direct and indirect TFEB-dependent mechanisms. As far as we can tell, TFE3 is regulated in a similar manner.

Likewise, we found that bacterial molecules activate TFE3 and TFEB through a TLR-independent mechanism. These observations and the hypothetical model they support provide an explanation for previous, intriguing observations that

phagocytosis is sufficient to trigger TFEB nuclear translocation, but only if the

phagocytic cup is able to close [7]. In this model, ROS serve dual functions as a

bactericidal agent in the phago-lysosome, and as a signal to activate CD38 and

initiate the signal cascade that results in TFEB/TFE3 activation. Since both

PHOX- and mitochondria-derived ROS activated TFEB and TFE3, this provides

redundant mechanisms of ROS production that report on phagocytosis and mito-

chondrial integrity. Thus, the model predicts that any condition that leads to

PHOX activation or mitochondrial disruption, increasing ROS, would result in

TFEB/3 activation (at least, in macrophages). This has important implications for

homeostasis and disease.

Since our discovery of the key role of TFEB in the transcriptional host re-

sponse to infection in macrophages [4], many studies have independently shown

that TFEB and/or TFE3 greatly contribute to innate immunity against a large

number of bacterial pathogens [13,15,41]. In addition, several signaling components

have been identified in the regulation of TFEB/3, including AGS3 during LPS

stimulation, the activation of TFEB by IFN-γ, and the contribution of

AMPK/mTORC1 regulation by several mechanisms [42,43]. Moreover, at least one

example of bacterial modulation of TFEB has been elucidated, particularly how

Mycobacterium tuberculosis represses TFEB through *mir-33* and *mir-33** [44]. The-

se discoveries underscore the central importance of TFEB/TFE3 as key contribu-

tors to the host response to infection, and provide a strong rationale for complete

understanding of the contributions of the novel PHOX/CD38/MCOLN1/TFEB

pathway described herein in distinct infectious diseases.

The events that occur downstream of TFEB/3 activation are of considerable in-

terest. Prior studies demonstrated that TFEB and TFE3 have redundant and non-

redundant functions important for the induction of key immune signaling mole-

cules, such as intracellular signaling pathway components and extracellular cyto-

kines and chemokines in LPS-stimulated macrophages [6]. Here we show that this

is also true in the case of bacterial phagocytosis. Thus, TFEB/3 are predicted to

be key contributors to the function of cellular networks in tissues that are under

pathogenic attack, both in the induction of antimicrobial responses and in the re-

cruitment of effector cells of the innate and adaptive immune systems.

Furthermore, we show that TFEB activation in macrophages induces

lysosomal biogenesis, including the induction of TRPML1/MCOLN1 expression.

Enhanced lysosomal biogenesis increases the availability of TRPML1/MCOLN1-

containing lysosomes for fusion with PHOX/CD38-positive maturing

phagosomes, thus enacting a positive phagocytosis feedback loop. In addition,

prior studies showed how TFEB activation in macrophages leads to increased

autophagy [45–47], consistent with results in many other cell types. Thus, activation

of the PHOX/CD38/TRPML1/CN/TFEB axis drives a positive phagocytosis

feedforward loop that enhances the compartment that is poised to receive the

mature phagolysosome for degradation, and enhances lysosomal exocytosis of

remaining bacterial debris.

In summary, our studies reveal a heretofore unknown mechanism of macro-

phage activation that is activated by phagocytosis of bacterial cells. Such mech-

anism is parallel to better-understood TLR-mediated pathogen detection mecha-

nisms. A key challenge for the future is to understand how diverse pathogen-

detection pathways interact within an integrated regulatory network to produce a

cellular response that is tailored to each particular pathogenic challenge. Addi-

tionally, understanding how pathogens have evolved to circumvent this new axis

may help understand microbial pathogenesis and reveal new mechanisms to

safely manipulate it for therapeutic purposes.

Materials and Methods

Isolation and differentiation of Bone Marrow Derived macrophages

(BMDMs)

Femurs and tibias from 8-12 weeks old mice were separated and cleaned. Bone marrow was flushed into 50 ml tubes under the sterile hood. Bone marrow was passed through the bone until the color of the bone turned white. After centrifugation at 1500 rpm for 5 minutes at 4 °C, cell pellets were resuspended and plated in BMDM media: DMEM (Fisher Scientific, MT10102CV), FBS 10% (Thermo Fisher, 16000069), AA 1% (Life Technologies, 15240-062), L-glutamine 1% (Life Technologies, 25030-081), MEN NEA AA 1% (Life Technologies, 11140-050), 2-Mercapto 0.1% (Life Technologies, 21985), IL-3 5 ng/ml (Peprotech: 213-13), MCSF 5 ng/ml (Peprotech: 315-02). Cell were used for experiments after 1 week of differentiation. To produce immortalized BMDMs, cells were transformed by CreJ2 virus [48].

Cell culture, transfection and Imaging

RAW264.7 and iBMDM cells were grown in Dulbecco's Modified Eagle's Medium (DMEM) - high glucose (Sigma-Aldrich, D6429-500ML) containing 10% endotoxin tested FBS (Thermo Fisher, 16000044), 1% Antibiotic-Antimycotic (Life Technologies 15240-062). Cells were passage 4 to 12 times. iBMDM GFP-TFEB stably transfected cells were created using pEGFP-N1-TFEB (pEGFP-N1-TFEB was a gift from Shawn Ferguson, Addgene plasmid # 38119), Lipofectamine

3000 (Thermo Fisher, L3000008) used according to manufacturer's instructions,

and G418 sulfate (Life Technologies, 10131) used for selection. 10 days after se-

lection, stable cells were separated by FACS. For Imaging we used 96-Well Op-

tical-Bottom Plates (Thermo Fisher, 165305). 6×10^4 cells were seeded in each

well. At the end of the experiments, cells were fixed using 4% paraformaldehyde

(Sigma-Aldrich, 158127) and incubated with Hoechst stain (Anaspec, AS-83218)

at room temperature for 20 minutes as nuclear staining. For lysotracker staining,

LysoTracker Red DND-99 (Thermo Fisher, L7528) was added to the media 30

minutes before fixing according to the manufacture instruction. Image acquisition

was automatically performed using a Cytation3 Imaging Plate Reader (Biotek).

The N/C ratio was measured using CellProfiler (Broad Institute), as described in

[5]. Colocalization analysis was performed using ImageJ software (NIH).

siRNA Knockdown

siRNA compounds were purchased from Dharmacon RNAi Technologies.

siGENOME NonTargeting siRNA #1 (D-001210-01-05). siGENOME Mouse Cd38

(12494) siRNA - SMARTpool (M-058632-01- 0005). siGENOME Mouse Ppp3cb

(19056) siRNA (M-063545-00-0005). siGENOME Mouse Ppp3r1 (19058) siRNA

(M-040744-01-0005). siGENOME Mouse Mcoln1 (94178) siRNA (M-044469-00-

0005). We used Lipofectamine RNAiMAX (Thermo Scientific, 13778030) for

transfection according to manufacturer's instructions.

Bacterial strains

Salmonella enterica serovar Typhimurium SL1344 wild type is a gift from Brian

Coombes (McMaster University, Canada), *Staphylococcus aureus* NCTC8325 is

a gift from Fred Ausubel, MGH Research Institute, USA.

In vitro infection

Bacteria were grown overnight at 37 °C in LB medium (Difco, BD) with 100

µg/ml streptomycin for Salmonella and Columbia medium (Difco, BD) with

10 µg/ml Nalidixic acid for *S. aureus*. The following day cultures were diluted 1:50

in the same medium and grown at 37 °C for 3 h to late-exponential phase,

washed twice in cold PBS, and cells were infected with MOI 10 for *S. aureus* and

MOI 10 for S*. enterica*, as in (Engelenburg and Palmer, 2010; Najibi et al., 2016;

Trieu et al., 2009; Visvikis et al., 2014). For gentamycin antibiotic (AB) - killed

bacteria, before addition to cells, gentamicin (100 µg/ml) was added to washed

bacteria in PBS for 2 hours and 100% killing of the bacteria was confirmed by

culture for 48 h on LB-streptomycin agar at 37°C**.** The appropriate amounts of

bacteria were resuspended in DMEM 10 % FBS without antibiotic.

Immunofluorescence

After treatment, cells were fixed with 4% paraformaldehyde (PFA) pH 7.4 at

room temperature for 10 min and washed 3 times in PBS (Gibco Life Technolo-

gies,10010) for 5 min each. PFA was neutralized with 50 mM NH_4Cl in PBS at

room temperature for 10 minutes with agitation. After 3 washes with PBS, cells

were permeabilized with 0.1% Triton X in PBS at room temperature on agitator

for 5 min and then blocked with 5% bovine serum albumin (Sigma-Aldrich,

A9647) in PBS for 1 h. After 3 washes with PBS cells were incubated with prima-

ry antibodies for 2 hours. Anti-TFEB Antibody (Cell Signaling, 4240). Anti-TFE3

antibody (Sigma-Aldrich, HPA023881-100UL). Anti-CD38 antibody (Abcam,

ab24978). Anti-NOX2/gp91phox antibody (Abcam, ab80508). Anti-LAMP1 (Cell

Signaling Technology, 9091). Anti-Mucolipin 1 (MCOLN1) (C-Term) antibody

(Antibodies-Online, ABIN571446). Cells were washed thrice in PBS and incubat-

ed with the fluorescent secondary antibody plus Hoechst stain (Anaspec, AS-

83218) at room temperature for 1 h. Image acquisition was performed using a

Cytation3 imaging plate reader (Biotek).

Immunoblotting

Cells were washed 3 times with PBS, harvested, and lysed with 1X SDS sam-

ple buffer Blue Loading Pack (Cell Signaling, 7722) at 100 µl per well of 6-well

plate. Lysates were heated at 100 °C for 5 min and then centrifuged for 5 min.

The supernatant was collected and sonicated, gel electrophoresis was performed

using Novex 4-20% Tris-Glycine Mini Gels (Invitrogen, XP04200BOX), and were

then transferred onto nitrocellulose (Life Technologies, LC2009). After wash with

TBS (Life Technologies, 28358) for 5 minutes, membranes were soaked in block-

ing buffer containing 1X TBS with 5% BSA for 1 hour at room temperature. After

3 washes with TBS-Tween (Life Technologies, 28360), membranes were incu-

bated overnight at 4 °C with primary antibodies and gentle agitation. Next mem-

branes were washed thrice with TBS-Tween and incubated with HRP-conjugated

secondary antibody (Cell Signaling, 7074 1:2000) for 1 h at room temperature

with gentle agitation. Membranes were then washed with TBS-Tween and incu-

bated with LumiGLO® (Cell signaling, 7003) for 1 min and exposed to x-ray film

(Denville Scientific, E3012). Primary antibodies and dilutions that were used are

as follows: β-actin antibody (Cell Signaling Technology, 4967, 1:1000), Anti-

CD38 Antibody (Cell Signaling Technology, 14637S 1:1000), Anti-TLR4 antibody

(). Anti-TLR2 antibody (). Anti-Mucolipin 1 (MCOLN1) (C-Term) antibody (Anti-

bodies-Online, ABIN571446 1:1000).

Drugs and reagents

Lipopolysaccharides (LPS) from *S. enterica* serotype Typhimurium (Sigma-

Aldrich, L6143-1MG, 10ug/ml): Peptidoglycan (PGN) from *S. aureus* (Invivogen,

tlrl-pgns2): Monophosphoryl Lipid A (MPLA-SM) (Invivogen, tlrl-mpla): ML-SA1

(Sigma Aldrich, SML0627) Ionomycin (Cayman Chemical Company, 10004974):

FK-506 (Cayman Chemical Company, 10007965): BAPTA AM (Cayman Chemi-

cal Company, 15551): Ned-19 (Cayman Chemical Company, 17527): Kuromanin

(Cayman Chemical Company, 16406): Apigenin (Cayman Chemical Company,

10010275): CCCP (Cayman Chemical Company, 25458): N-acetyl-L-Cysteine

(NAC) (Cayman Chemical Company, 20261): N-acetyl-L-Cysteine amide (NACA)

(Cayman Chemical Company, 25866): Apocynin (Cayman Chemical Company,

11976): D609 (Cayman Chemical, 13307, 50 μM): kb NB 142-70 (Cayman

Chemical Company, 18002).

ROS/Superoxide Detection Assay

Cellular ROS/Superoxide Detection Assay Kit (Abcam, ab139476) was used

according to the manufacture instruction. Mean fluorescence intensity was calcu-

lated by Biotek Gen5 Data Analysis Software.

RNA Sequencing and Differential Expression Analysis

RAW264.7 dKO were a one-time gift from Rosa Puertollano (NHLBI). Infection

was performed as described in *In Vitro Infection,* in three independent replicates.

RNA was extracted by TRIzol Plus RNA Purification Kit (Thermo Fisher,

12183555). Whole cell RNA was submitted to BGI Inc for ribosomal RNA deple-

tion, library construction, and sequencing on Illumina NextSeq machines. Reads

were trimmed and quality-filtered by BGI. Clean reads were mapped to the

GRCm38 mouse reference transcriptome using Salmon v0.13.1 and transcript-

quantified (Patro et al., 2017). Salmon quant outputs were analyzed using

tximport and DESeq2 v1.22.2 in R (Love et al., 2014). DESeq2 output counts

were used as input for interactive analysis and data plotting using DEBrowser

1.10.9 in R (Kucukural et al., 2019). Differentially expressed (fold change ≥ 2;

Padj ≤ 0.01; filter genes with fewer than 10 counts) gene sets were used as input

for g:Profiler v0.6.6 online tools (Raudvere et al., 2019).

ELISA Assay

Concentrations of IL-1α, IL-1β, IL-6 and TNF-α were measured by en-

zyme-linked immunosorbent assays (ELISAs) using kits purchased from R&D

Systems (DY400-05, DY401-05, DY406-05, DY8234-05, DY410-05, respective-

ly). All incubation periods occurred at room temperature, and plates were washed

with Tris buffered saline (TBS) containing 0.05% Tween 20 (Bethyl Laboratories,

E106) after each step. Briefly, clear microplates (R&D Systems, DY990) were

incubated overnight with capture antibody. Plates were blocked for one hour us-

ing TBS containing 1% BSA (Bethyl Laboratories, E104). Samples were added

undiluted (IL-1α, IL-1β, IL-6) or diluted 1:10 in TBS containing 1% BSA (TNF-α)

and plates were incubated for two hours. Detection antibody was added, and

plates were incubated for two hours. Streptavidin-Horseradish Peroxidase (Strep-

HRP) was added, and plates were incubated for 20 minutes. TMB One Compo-

nent HRP Microwell Substrate (Bethyl Laboratories, E102) was added, and reac-

tion was stopped after 20 minutes using ELISA Stop Solution (Bethyl Laborato-

ries, E115). Optical densities were measured at 450 nm and 540 nm using

SpectraMax Plus 384 Microplate Reader with SoftMax Pro software. Measure-

ments at 540 nm were subtracted from measurements at 450 nm to correct for

optical impurities of the plates, per manufacturer's recommendations. Concentra-

tions of samples were interpolated from each standard curve using Prism 8. Data

are represented as mean $\pm$ SEM.

LDH Assay

Lactate Dehydrogenase (LDH) activity was measured using an LDH assay kit

purchased from Abcam (ab102526). LDH assay was performed following manu-

facturer's guidelines. Briefly, samples were added to a clear microplate (R&D

Systems, DY990). Reaction mix was made using supplied LDH Assay Buffer and

LDH Substrate Mix. Reaction mix was then added to each sample. Optical densi-

ties were measured at 450 nm using SpectraMax Plus 384 Microplate Reader

with SoftMax Pro software. The amount of NADH in each sample was interpolat-

ed from the standard curve using Prism 8. LDH activity was quantified by dividing

the amount of NADH in each sample by the product of the reaction time in

minutes and the original volume of sample added into the reaction well in mL.

LDH activity was reported in mU/mL and plotted as mean $\pm$ SEM.

Figure Legends

Figure 1. Gram+ bacterial cells and ligands activate TFE3. A-H. GFP-TFEB immortalized BMDM (iBMDMs) were infected with *S. aureus* (MOI = 10) for 3 h. Incubation with 10 µg/ml PGN from *S. aureus* was 3 h. Cells were fixed and processed for anti-TFE3 immunofluorescence staining. Shown are representative images, and quantification of three biological replicates (n = 450). **A**. DMSO control, **B**. *S. aureus* infection, and **C**. *S. aureus* PGN (10 µg/ml). **D**. DMSO control, **E**. *S. aureus* infection, **F**. *S. aureus* PGN. **G.** GFP-TFEB intensity in the nucleus compared to the cytoplasm (N/C ratio) measured using CellProfiler. **** p ≤ 0.0001 (one-way ANOVA followed by Tukey's post-hoc test). (**H**) Anti-TFE3 fluorescence N/C ratio measured using CellProfiler. **** p ≤ 0.0001 (one-way ANOVA followed by Tukey's post-hoc test). **I-N**. GFP-TFEB iBMDMs were infected with *S. aureus* or incubated with *S. aureus* PGN as above, and stained with Lysotracker. Shown are representative images and quantification of three biological replicates (n = 60). **I**. DMSO control. **J**. *S. aureus* infection, and **K**. *S. aureus* PGN. **L-N**. Higher magnification micrographs for DMSO control (**L**), *S. aureus* (**M**), and PGN (**N**). **O**. Lysotracker intensity per cell (MFI/cell) measured with CellProfiler. **** p ≤ 0.0001 (one-way ANOVA followed by Tukey's post-hoc test). **P-R.** Comparison of Lysotracker intensities in cells with TFEB cytoplasmic localization (C) with that in cells with nuclear localization (N) following incubation with DMSO (**P**), *S. aureus* (**Q**), or PGN (**R**). **** p ≤ 0.0001 (two-sample two-tailed unpaired *t* test).

Figure 2. TFEB and TFE3 are required for the macrophage transcriptional response to *S. aureus.* A-E. Wild type (WT) and *Tfeb Tfe3* double knockout (dKO) RAW264.7 cells were infected as described in *Methods* for 3 h. N = 3 biological replicates. **A.** Heat map showing expression of genes that were differentially expressed in wild type cells. Shown are baseline (CO) and infected (SA) conditions, in both cell lines. Two groups of genes with defective induction in dKO cells are indicated with black boxes. **B.** Heat map showing expression of M1 and M2 marker genes. Colors indicate direction and magnitude of expression per row; circle diameter is proportional to absolute expression. Grey indicates no expression detected. **C.** Volcano plot showing differential gene expression in infected dKO cells relative to infected WT cells. Blue color denotes downregulation in dKO cells; red indicates upregulation in dKO cells. Green highlights noteworthy innate immunity genes, with corresponding adjacent labels. **D.** Pie chart showing percent of genes that are upregulated in infected v uninfected WT cells, and which are TFEB/3-dependent. **E.** Reactome pathway over-representation analysis of TFEB/3-dependent genes, plotting adjusted P values for each category (Padj). **F.** Verification of TFEB expression in *Tfeb*$^{flox/flox}$ (WT) and *Tfeb*$^{\Delta LysM}$ (*Tfeb*$^{-/-}$) iBMDMs by anti-TFEB immunoblot. **G-K.** Wild type (WT) and *Tfeb*$^{\Delta LysM}$ (*Tfeb*$^{-/-}$) iBMDMs were incubated with PBS (Ctrl) or infected with *S. aureus* (Sa, MOI = 10). Concentrations of IL-1α (**G**), IL-1β (**H**), IL-6 (**I**), TNF-α (**J**), and lactate dehydrogenase activity (LDH, cytolysis control, **K**) were measured in culture supernatants 13 h following infection, by ELISA and LDH assay, respectively. N = 5-6, *p

< 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001, one-way ANOVA followed by

Tukey's post-hoc test.

Figure 3. TLR signaling is dispensable for TFEB activation. A-N. Primary

BMDMs from WT and *Myd88 Trif* double knockout mice (*Myd88 Trif* dKO) were

infected, fixed, and processed for anti-TFEB immunofluorescence staining. TFEB

fluorescence N/C ratio per cell was measured using CellProfiler (n = 250). Shown

are representative images and quantification of three biological replicates. **A, B**.

DMSO control (t = 6 h); **C, D.** *Salmonella* Typhimurium (St; MOI = 1, t = 6 h). **E,**

F. Monophosphoryl lipid A (MPLA, 1 µg/ml, t = 6 h). **G, H.** *Salmonella*

Typhimurium lipopolysaccharide (LPS, 1 µg/ml, t = 6 h). **I, J**. *S. aureus* (Sa, MOI

= 10, t = 3 h). **K, L**. *S. aureus* peptidoglycan (PGN, 10 µg/ml, t = 3 h). **M, N.**

Quantification of TFEB N/C ratio per cell in WT (**M**) and *Myd88 Trif* dKO (**N**) cells.

*** p ≤ 0.001 (one-way ANOVA followed by Tukey's post-hoc test).

Figure 4. TFEB activation is mediated by TRPML1/MCOLN1, Ca^{2+}, and

calcineurin. A-E. GFP-TFEB iBMDMs were treated with DMSO without infection

(t = 3 h, **A**) or treated for 3 h and subsequently infected with *S. aureus* (**B**, MOI =

10, t = 3 h). In parallel, cells were treated with BAPTA (**C,** 10 µM, t = 3 h) or

FK506 (**D,** 5 µM, t = 6 h) and subsequently infected with *S. aureus* (MOI = 10, t =

3 h). **E.** Quantification of GFP-TFEB N/C Ratio by CellProfiler (3 biological repli-

cates, n = 350 cells). **** p ≤ 0.0001 (one-way ANOVA followed by Tukey's post-

hoc test). **F-K.** GFP-TFEB iBMDMs were treated with scrambled (Scr, **F**) or

siRNA against *Ppp3cb* (**G**), *Ppp3r1* (**H**) for 48 h. Cells were treated with Scr (**I**),

Ppp3cb (**J**), or *Ppp3r1* (**K**) siRNA for 48 h prior to infection with *S. aureus* (MOI =

10, t = 3 h). **L.** GFP-TFEB N/C Ratio by CellProfiler (3 biological replicates, n =

210 cells). *** $p \leq 0.001$ (one-way ANOVA followed by Tukey's post-hoc test). **M,**

N. GFP-TFEB iBMDMs were treated with DMSO (**M**, t = 6 h) or Ionomycin (**N**, 10

μM, t = 6 h). **O.** GFP-TFEB N/C Ratio by CellProfiler (3 biological replicates, n =

350 cells). **** $p \leq 0.0001$ (one-way ANOVA followed by Tukey's post-hoc test).

P-S. GFP-TFEB iBMDMs were treated with scrambled (Scr, **P**) or siRNA against

Mcoln1 (**Q**) for 48 h. Cells were treated with Scr (**R**) or *Mcoln1* (**S**) siRNA for 48 h

prior to infection with *S. aureus* (MOI = 10, t = 3 h). **T.** GFP-TFEB N/C Ratio by

CellProfiler (3 biological replicates, n = 300 cells). *** $p \leq 0.001$ (one-way ANOVA

followed by Tukey's post-hoc test). **U, V.** GFP-TFEB iBMDMs were treated with

DMSO (**U**, t = 3 h) or ML-SA1 (**V**, 10 μM, t = 3 h). **W.** GFP-TFEB N/C Ratio by

CellProfiler (3 biological replicates, n = 355 cells). **** $p \leq 0.0001$ (two-sample

two-sided *t* test).

Figure 5. CD38 activates TFEB through NAADP and TRPML1/MCOLN1. A-

E. GFP-TFEB iBMDMs were treated with DMSO without infection (DM., t = 6 h,

A) or treated for 6 h and subsequently infected with *S. aureus* (**B**, MOI = 10, t = 3

h). In parallel, cells were treated with Ned-19 (**C,** Ned., 10 μM, t = 4 h),

Kuromanin (**D,** Kuro., 100 μM, t = 6 h), or Apigenin (**E,** Api., 100 μM, t = 6 h) and

subsequently infected with *S. aureus* (MOI = 10, t = 3 h). **F.** Quantification of

GFP-TFEB N/C Ratio by CellProfiler (3 biological replicates, n = 200 cells). **** p

≤ 0.0001 (one-way ANOVA followed by Tukey's post-hoc test). **G-I.** GFP-TFEB

iBMDMs were treated with scrambled (Scr, **G**) or siRNA for 48 h. Cells were

treated with Scr (**H**) or *Cd38* (**I**) siRNA for 48 h prior to infection with *S. aureus*

(MOI = 10, t = 3 h). **J.** GFP-TFEB N/C Ratio by CellProfiler (3 biological repli-

cates, n = 350 cells). *** p ≤ 0.001 (one-way ANOVA followed by Tukey's post-

hoc test). **K-M.** GFP-TFEB iBMDMs were treated with DMSO (DM., t = 2 h, **K**) or

treated with scrambled siRNA (Scr, t = 48 h) and subsequently incubated with

NAADP-AM (100 nM, t = 2 h, **L**). In parallel, cells were treated with *Mcoln1*

siRNA (t = 48 h) and subsequently incubated with NAADP-AM (100 nM, t = 2 h,

M). **N.** Quantification of GFP-TFEB N/C Ratio by CellProfiler (3 biological repli-

cates, n = 300 cells). **** p ≤ 0.0001 (one-way ANOVA followed by Tukey's post-

hoc test). **O, P.** CD38 immunofluorescence in wild type primary BMDM. PBS con-

trol (**O**) and *S. aureus* infection (MOI = 10, t = 2 h, **P**). **Q.** CD38 immunoblot of

whole cell lysates from wild type and *Tfeb*$^{\Delta LysM}$ (*Tfeb*$^{-/-}$) iBMDMs incubated with

PBS or infected with *S. aureus* (MOI = 10, t = 3 h). β-actin is loading control.

Representative of three biological replicates.

Figure 6. ROS activate TFEB through CD38. ROS (**A**) and superoxide (**B**)

generated in wild type primary BMDM during infection with *S. aureus* (MOI = 10, t

= 1 h) shown as mean fluorescence intensity (MFI) per cell and measured with

Gen5 (n = 50 cells, 3 biological replicates). **** p ≤ 0.0001 (two-sided two-sample

t test). **C-F.** GFP-TFEB iBMDMs were treated with DMSO without infection (DM.,

t = 4 h, **C**) or treated for 4 h and subsequently infected with *S. aureus* (**D**, MOI =

10, t = 3 h). In parallel, cells were treated with NAC (**E,** 5 mM, t = 4 h) or NACA

(**F,** 1 mM, t = 4 h) and subsequently infected with *S. aureus* (MOI = 10, t = 3 h).

G. Quantification of GFP-TFEB N/C Ratio by CellProfiler (3 biological replicates,

n = 250 cells). *** p ≤ 0.001, **** p ≤ 0.0001 (one-way ANOVA followed by

Tukey's post-hoc test). **H-J.** GFP-TFEB iBMDMs were treated with DMSO (DM., t

= 3 h, **H**) or treated with scrambled siRNA (Scr, t = 48 h) and subsequently incu-

bated with CCCP (10 μM, t = 3 h, **I**). In parallel, cells were treated with *Cd38*

siRNA (t = 48 h) and subsequently incubated with CCCP (10 μM, t = 3 h, **J**). **K.**

Quantification of GFP-TFEB N/C Ratio by CellProfiler (3 biological replicates, n =

355 cells). **** p ≤ 0.0001 (one-way ANOVA followed by Tukey's post-hoc test).

Figure 7. Infection induced ROS and TFEB activation require NADPH ox-

idase. ROS (**A**) and superoxide (**B**) generated in wild type and *Nox2*$^{-/-}$ primary

BMDM during infection with *S. aureus* (MOI = 10, t = 1 h) shown as mean fluo-

rescence intensity (MFI) per cell and measured with Gen5 (n = 200 cells, 3 bio-

logical replicates). **** p ≤ 0.0001 (two-sided two-sample *t* test). **C-F.** TFEB im-

munofluorescence in wild type and *Nox2*$^{-/-}$ primary BMDM. PBS controls (**C, E**)

and infected cells (MOI = 10, t = 3 h, **D, F**). **G.** Quantification of TFEB N/C Ratio

by CellProfiler (3 biological replicates, n = 200 cells). *** p ≤ 0.001, ns: p =

0.4399 (one-way ANOVA followed by Tukey's post-hoc test). **H-J.** GFP-TFEB

iBMDMs were treated with DMSO without infection (DM., t = 4 h, **H**) or treated for

4 h and subsequently infected with *S. aureus* (**I**, MOI = 10, t = 3 h). In parallel,

cells were treated with Apocynin (**J,** 10 μM, t = 4 h) and subsequently infected

with *S. aureus* (MOI = 10, t = 3 h). **K.** Quantification of GFP-TFEB N/C Ratio by

CellProfiler (3 biological replicates, n = 250 cells). *** $p \le 0.001$ (one-way ANOVA

followed by Tukey's post-hoc test). **L-S.** Immunofluorescence detection of NOX2

and CD38 in wild type primary BMDM. Cells were incubated with PBS (**L-O**) or

infected with *S. aureus* (MOI = 10, t = 3 h, **P-S**). **T.** Colocalization analysis of

NOX2 and CD38. **** $p \le 0.0001$ (two-sided two-sample unpaired *t* test). **U**. Pro-

posed hypothetical model for the phagocytosis-triggered activation of TFEB.

Acknowledgments

The authors are grateful to members of the Irazoqui laboratory, the Fitzgerald laboratory, the Program in Innate Immunity, and the Department of Microbiology and Physiological Systems for helpful insights and discussions. The Fitzgerald laboratory provided several knockout primary and immortalized BMDMs, and the Sassetti laboratory provided *Nox2* mutant mice and CreJ2 virus. This work was partially funded by grant R01GM101056 (JEI) from the National Institutes of Health. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Competing Interests

The authors declare no competing interests.

Author Contributions

Conceptualization, JEI and MN; Methodology, JEI and MN; Investigation, MN, JAM, JEI, and HHH; Animals, JAM; Writing–original draft, JEI; Writing–review and editing, JEI, MN, JAM, and HH; Visualization, MN and JEI; Supervision, JEI; Funding acquisition, JEI.

References

1. Aderem, A. & Underhill, D. M. MECHANISMS OF PHAGOCYTOSIS IN MACROPHAGES. *Annu Rev Immunol* **17,** 593–623 (1999).

2. Lawrence, T. & Natoli, G. Transcriptional regulation of macrophage polarization: enabling diversity with identity. *Nat Rev Immunol* **11,** 750 (2011).

3. Medzhitov, R. & Horng, T. Transcriptional control of the inflammatory response. *Nat Rev Immunol* **9,** 692–703 (2009).

4. Visvikis, O. *et al.* Innate Host Defense Requires TFEB-Mediated Transcription of Cytoprotective and Antimicrobial Genes. *Immunity* **40,** 896–909 (2014).

5. Najibi, M., Labed, S., Visvikis, O. & Irazoqui, J. An Evolutionarily Conserved PLC-PKD-TFEB Pathway for Host Defense. *Cell Reports* **15,** 1728–1742 (2016).

6. Pastore, N. *et al.* TFEB and TFE3 cooperate in the regulation of the innate immune response in activated macrophages. *Autophagy* **12,** 1–19 (2016).

7. Gray, M. A. *et al.* Phagocytosis Enhances Lysosomal and Bactericidal Properties by Activating the Transcription Factor TFEB. *Curr Biology Cb* **26,** 1955–64 (2016).

8. Puertollano, R., Ferguson, S. M., Brugarolas, J. & Ballabio, A. The complex

relationship between TFEB transcription factor phosphorylation and subcellular

localization. *Embo J* **37,** e98804 (2018).

9. Malavasi, F. *et al.* Evolution and function of the ADP ribosyl cyclase/CD38

gene family in physiology and pathology. *Physiol Rev* **88,** 841 886 (2008).

10. Roberts, R. A. *et al.* Nitrative and Oxidative Stress in Toxicology and Dis-

ease. *Toxicol Sci* **112,** 4–16 (2009).

11. Lin, M. T. & Beal, F. M. Mitochondrial dysfunction and oxidative stress in neu-

rodegenerative diseases. *Nature* **443,** nature05292 (2006).

12. Halliwell, B. & Cross, C. Oxygen-derived species: their relation to human dis-

ease and environmental stress. *Environ Health Persp* **102,** 5–12 (1994).

13. Raben, N. & Puertollano, R. TFEB and TFE3: Linking Lysosomes to Cellular

Adaptation to Stress. *Annu Rev Cell Dev Bi* **32,** 1–24 (2015).

14. Martina, J. A. & Puertollano, R. Protein phosphatase 2A stimulates activation

of TFEB and TFE3 transcription factors in response to oxidative stress. *J Biol*

Chem **293,** 12525–12534 (2018).

15. El-Houjeiri, L. *et al.* The Transcription Factors TFEB and TFE3 Link the

FLCN-AMPK Signaling Axis to Innate Immune Response and Pathogen Re-

sistance. *Cell Reports* **26,** 3613-3628.e6 (2019).

16. Eswarappa, S., Negi, V., Chakraborty, S., Sagar, C. B. & Chakravortty, D.

Division of the Salmonella-containing vacuole and depletion of acidic lysosomes

in Salmonella-infected host cells are novel strategies of Salmonella enterica to

avoid lysosomes. *Infect Immun* **78,** 68 79 (2010).

17. McGourty, K. *et al. Salmonella* Inhibits Retrograde Trafficking of Mannose-6-

Phosphate Receptors and Lysosome Function. *Science* **338,** 963–967 (2012).

18. Fabregat, A. *et al.* The Reactome Pathway Knowledgebase. *Nucleic Acids*

Res gkx1132- (2017). doi:10.1093/nar/gkx1132

19. Takeda, K. & Akira, S. TLR signaling pathways. *Semin Immunol* **16,** 3–9

(2004).

20. Yamamoto, M. *et al.* Role of Adaptor TRIF in the MyD88-Independent Toll-

Like Receptor Signaling Pathway. *Science* **301,** 640–643 (2003).

21. Medina, D. L. *et al.* Lysosomal calcium signalling regulates autophagy

through calcineurin and TFEB. *Nat Cell Biol* **17,** 288–299 (2015).

22. Paola, S., Scotto-Rosato, A. & Medina, D. TRPML1: The Ca(2+)retaker of the

lysosome. *Cell Calcium* **69,** (2018).

23. Shen, D. *et al.* Lipid storage disorders block lysosomal trafficking by inhibiting

a TRP channel and lysosomal calcium release. *Nat Commun* **3,** 731 (2012).

24. Zhang, F. & Li, P.-L. Reconstitution and Characterization of a Nicotinic Acid

Adenine Dinucleotide Phosphate (NAADP)-sensitive Ca2+ Release Channel

from Liver Lysosomes of Rats. *J Biol Chem* **282,** 25259–25269 (2007).

25. Zhang, F., Jin, S., Yi, F. & Li, P. TRP□ML1 functions as a lysosomal

NAADP□sensitive Ca2+ release channel in coronary arterial myocytes. *J Cell*

Mol Med **13,** 3174–3185 (2009).

26. Naylor, E. *et al.* Identification of a chemical probe for NAADP by virtual

screening. *Nat Chem Biol* **5,** 220 226 (2009).

27. Kellenberger, E., Kuhn, I., Schuber, F. & Muller-Steffner, H. Flavonoids as

inhibitors of human CD38. *Bioorg Med Chem Lett* **21,** 3939–3942 (2011).

28. Escande, C. *et al.* Flavonoid Apigenin Is an Inhibitor of the NAD+ase CD38

Implications for Cellular NAD+ Metabolism, Protein Acetylation, and Treatment of

Metabolic Syndrome. *Diabetes* **62,** 1084–1093 (2013).

29. Parkesh, R. *et al.* Cell-permeant NAADP: A novel chemical tool enabling the

study of Ca2+ signalling in intact cells. *Cell Calcium* **43,** 531–538 (2008).

30. Funaro, A. *et al.* CD38 functions are regulated through an internalization

step. *J Immunol Baltim Md 1950* **160,** 2238–47 (1998).

31. ZOCCHI, E. *et al.* Ligand-induced internalization of CD38 results in intracellu-

lar Ca2+ mobilization: role of NAD+ transport across cell membranes. *Faseb J*

13, 273–283 (1999).

32. Kang, J. *et al.* The Role of CD38 in Fcγ Receptor (FcγR)-mediated Phagocy-

tosis in Murine Macrophages. *J Biol Chem* **287,** 14502–14514 (2012).

33. Wilson, H. L. *et al.* ADP-ribosyl Cyclase and Cyclic ADP-ribose Hydrolase Act

as a Redox Sensor A PRIMARY ROLE FOR CYCLIC ADP-RIBOSE IN

HYPOXIC PULMONARY VASOCONSTRICTION. *J Biol Chem* **276,** 11180–

11188 (2001).

34. Zhang, A. Y., Yi, F., Teggatz, E. G., Zou, A.-P. & Li, P.-L. Enhanced produc-

tion and action of cyclic ADP–ribose during oxidative stress in small bovine coro-

nary arterial smooth muscle. *Microvasc Res* **67,** 159–167 (2004).

35. Zhang, X. *et al.* MCOLN1 is a ROS sensor in lysosomes that regulates au-

tophagy. *Nat Commun* **7,** 12109 (2016).

36. DeLeo, F., Allen, L., Apicella, M. & Nauseef, W. NADPH oxidase activation

and assembly during phagocytosis. *J Immunol Baltim Md 1950* **163,** 6732–40

(1999).

37. Slauch, J. M. How does the oxidative burst of macrophages kill bacteria? Still

an open question. *Mol Microbiol* **80,** 580–583 (2011).

38. Marks, D. L. *et al.* Role of protein kinase d in Golgi exit and lysosomal target-

ing of the transmembrane protein, Mcoln1. *Traffic* **13,** 565 575 (2012).

39. Park, D.-R., Nam, T.-S., Kim, Y.-W., Bae, Y. & Kim, U.-H. Oxidative activation

of type III CD38 by NADPH oxidase–derived hydrogen peroxide in Ca2+ signal-

ing. *Faseb J* fj.201800235R (2018). doi:10.1096/fj.201800235r

40. Xu, M. *et al.* Contribution of NADPH Oxidase to Membrane CD38 Internaliza-

tion and Activation in Coronary Arterial Myocytes. *Plos One* **8,** e71212 (2013).

41. Brady, O. A., Martina, J. A. & Puertollano, R. Emerging roles for TFEB in the

immune response and inflammation. *Autophagy* **14,** 1–9 (2017).

42. Vural, A. *et al.* Activator of G-Protein Signaling 3–Induced Lysosomal Bio-

genesis Limits Macrophage Intracellular Bacterial Infection. *J Immunol* **196,** 846–

856 (2016).

43. Singh, N. *et al.* Antimycobacterial effect of IFNG (interferon gamma)-induced

autophagy depends on HMOX1 (heme oxygenase 1)-mediated increase in intra-

cellular calcium levels and modulation of PPP3/calcineurin-TFEB (transcription

factor EB) axis. *Autophagy* **14,** 1–57 (2018).

44. Ouimet, M. *et al.* Mycobacterium tuberculosis induces the miR-33 locus to

reprogram autophagy and host lipid metabolism. *Nat Immunol* **17,** 677–686

(2016).

45. Evans, T. D., Jeong, S.-J., Zhang, X., Sergin, I. & Razani, B. TFEB and tre-

halose drive the macrophage autophagy-lysosome system to protect against

atherosclerosis. *Autophagy* **14,** 1–7 (2018).

46. Sergin, I. *et al.* Exploiting macrophage autophagy-lysosomal biogenesis as a

therapy for atherosclerosis. *Nat Commun* **8,** ncomms15750 (2017).

47. Emanuel, R. *et al.* Induction of Lysosomal Biogenesis in Atherosclerotic Mac-

rophages Can Rescue Lipid-Induced Lysosomal Dysfunction and Downstream

Sequelae. *Arteriosclerosis Thrombosis Vasc Biology* **34,** 1942–1952 (2014).

48. Blasi, E., Radzioch, D., Merletti, L. & Varesio, L. Generation of macrophage

cell line from fresh bone marrow cells with a myc/raf recombinant retrovirus.

Cancer Biochem Bioph **10,** 303–17 (1989).