Low oxygen enhances trophoblast column growth by potentiating the extravillous lineage and promoting LOX activity

Jenna Treissman[1,2], Victor Yuan[,1,3], Jennet Baltayeva[1,2], Hoa T. Le[1,2], Barbara Castellana[1,2], Wendy P. Robinson[1,3], Alexander G. Beristain[1,2]

[1] The British Columbia Children's Hospital Research Institute, Vancouver, Canada.
[2] Department of Obstetrics & Gynecology, The University of British Columbia, Vancouver, Canada.
[3] Department of Medical Genetics, The University of British Columbia, Vancouver, Canada.

To whom correspondence should be addressed: Alexander G. Beristain, The British Columbia Children's Hospital Research Institute, The University of British Columbia, Vancouver, British Columbia, Canada. V5Z 4H4. Tel: (604) 875-3573; E-mail: aberista@mail.ubc.ca

Keywords: Hypoxia, placentation, trophoblast, extravillous trophoblast, villous cytotrophoblast, lysyl oxidase

Summary Statement: Low oxygen promotes extravillous trophoblast differentiation

Abbreviations

BAPN: β-aminopropionitrile

BrdU: Bromodeoxyuridine

CM: Conditioned media

CTB: Cytotrophoblast

DAP: 1,5-diaminopentane

DCT: Distal column trophoblast

DGE: Differential gene expression

ECM: Extracellular matrix

EVT: Extravillous trophoblast

FDR: False discovery rate

GO: Gene ontology

HIF1A: Hypoxia inducible factor 1 A

HLA-G: Human leukocyte antigen G

Hr: Hour

IF: Immunofluorescence

LOX: Lysyl oxidase

PCA: Principal component analysis

PCT: Proximal column trophoblast

PEG10: Paternally expressed gene 10

scRNA-seq: single cell RNA sequencing

SCT: Syncytiotrophoblast

UMAP: Uniform manifold approximation and projection

vCTB: Villous cytotrophoblast

ABSTRACT

Early placental development and the establishment of the invasive trophoblast lineage take place within a low oxygen environment. However, conflicting and inconsistent findings have obscured the role of oxygen in regulating invasive trophoblast differentiation. In this study, the effect of hypoxic, normoxic, and atmospheric oxygen on invasive extravillous pathway progression was examined using a human placental explant model. Here, we show that exposure to low oxygen enhances extravillous column outgrowth and promotes the expression of genes that align with extravillous trophoblast (EVT) lineage commitment. By contrast, super-physiological atmospheric levels of oxygen promote trophoblast proliferation while simultaneously stalling EVT progression. Low oxygen-induced EVT differentiation coincided with elevated transcriptomic levels of lysyl oxidase (*LOX*) in trophoblast anchoring columns, where functional experiments established a role for LOX activity in promoting EVT column outgrowth. The findings of this work support a role for low oxygen in potentiating the differentiation of trophoblasts along the extravillous pathway. Additionally, these findings generate insight into new molecular processes controlled by oxygen during early placental development.

INTRODUCTION

In mammalian development, the placenta forms the mechanical and physiological link between maternal and fetal circulations. In rodents and humans that have invasive haemochorial placentae, nutrient and oxygen transfer between mother and fetus is achieved through extensive uterine infiltration by placenta-derived cells of epithelial lineage called trophoblasts (Pijnenborg et al., 2011; Velicky et al., 2016). In humans, trophoblast differentiation into invasive cell subtypes, called extravillous trophoblast (EVT), is essential for optimal placental function (Tilburgs et al., 2015; Velicky et al., 2016). Molecular processes governing invasive EVT differentiation and specific EVT functions like uterine artery remodeling and immuno-modulation of maternal leukocytes are strictly controlled (Pollheimer et al., 2018; Wallace et al., 2012). Defects in trophoblast differentiation along the EVT pathway associate with impaired placental function and certain aberrant conditions of pregnancy that directly impact fetal and maternal health (Avagliano et al., 2012).

In early pregnancy, anchoring villi of the placental basal plate initiate cellular differentiation events leading to the formation of EVT. At specific villi-uterine attachment points, proliferative EVT progenitors establish multi-layered cellular structures called anchoring columns (Haider et al., 2016; Pollheimer et al., 2018). Trophoblasts residing within proximal regions of anchoring columns, termed proximal column trophoblasts (PCT), are highly proliferative and show evidence of initial molecular characteristics that are hallmarks of EVT (Haider et al., 2018; Turco et al., 2018). At distal regions within anchoring columns, column trophoblasts lose their proliferative phenotype and express many markers akin to invasive EVT, such as HLA-G, α5 and β1 integrins, NOTCH2, and ERBB2 (Fock et al., 2015a; Fock et al., 2015b; Haider et al., 2016; Kabir-Salmani et al., 2004; Zhou et al., 1997). The transition of PCT into distal column trophoblasts (DCT) represents a significant developmental step towards the formation of uterine-invading and tissue-remodeling EVT.

Anchoring columns of the placenta initially develop in the absence of maternal blood, and

subsequently, within a relatively low oxygen environment (~20 mmHg) (Jauniaux et al., 2000;

Jauniaux et al., 1999; Rodesch et al., 1992). By comparison, the partial pressure of arterial blood

oxygen is approximately 100 mmHg, while the partial pressure of oxygen within the placental bed by

14 weeks' gestation is estimated to be between 40-60 mmHg (Jauniaux et al., 2001). The role of

oxygen in controlling anchoring column formation and EVT differentiation has been the focus of many

studies (Caniggia et al., 2000; Genbacev et al., 1997; James et al., 2006; Lash et al., 2006; Wakeland et

al., 2017). Unfortunately, contradictory and inconsistent findings have obscured the role of oxygen

tension in controlling aspects of trophoblast biology, where low oxygen both promotes (Caniggia et al.,

2000; Genbacev et al., 1997) and restrains (James et al., 2006; Lash et al., 2006) column outgrowth,

and potentiates (Wakeland et al., 2017) and inhibits (Caniggia et al., 2000; Genbacev et al., 1997) EVT

differentiation. These reported differences on the effect of oxygen on anchoring column outgrowth and

EVT differentiation are likely attributed to differences in model platforms and methods used to isolate,

characterize, and culture trophoblasts. Nonetheless, the role of oxygen in controlling anchoring column

formation and column trophoblast differentiation has yet to be fully examined.

In this study, we examine how differing levels of oxygen affect first trimester placental column

outgrowth. Using placental villous tissue explants that recapitulate many of the morphological and

molecular events central to anchoring column formation and EVT differentiation *in vivo*, we show that

low levels of oxygen potentiate column outgrowth. We demonstrate that low oxygen drives hypoxia-

related gene programs and processes central to cell-extracellular matrix interaction, while exposure to

high oxygen promotes/maintains a strong proliferative phenotype. Moreover, we provide evidence that

supports a role for low levels of oxygen in promoting the differentiation of column trophoblasts along

the EVT pathway. We further identify *LOX* as a critical gene up-regulated in response to low oxygen

that supports column outgrowth, and provide important insight into novel molecular programs

impacted by oxygen that align with trophoblast differentiation along the EVT pathway.

RESULTS

Hypoxia potentiates trophoblast column outgrowth

To test the effect of exposure to differing levels of oxygen on trophoblast column establishment

and outgrowth, we utilized a Matrigel-imbedded placental explant model that closely reproduces

developmental processes of trophoblast column cell expansion and differentiation along the EVT

pathway (Bilban et al., 2009; Newby et al., 2005). Early column formation during human placental

development is characterized by the expansion of mitotically active Ki67^{+}/HLA-G^{lo} proximal column

trophoblasts (PCT) into HLA-G^{hi} non-proliferating distal column trophoblasts (DCT) and invasive

EVT (Figure 1A, 1B). To this end, Matrigel-imbedded placental explants recapitulate anchoring

column cell organization and EVT-lineage commitment, establishing the explant system as an

appropriate experimental tool to study the cellular and molecular underpinnings regulating human

trophoblast column formation and outgrowth (Figure 1C).

Chorionic villi harvested from early first trimester placentae (n=8; 5-8 weeks' gestation) were

imbedded into Matrigel matrix and allowed to establish for 24 hr at 5% oxygen (Figure 2A); this level

of oxygen represents a relative "normoxic" condition for early placental development (Jauniaux et al.,

2001). Characteristics (age, gestational age, BMI, smoking status) of each patient who donated their

placenta for explant culture are listed in Supplemental Table 1. Following this, explants derived from

the same placenta were transferred into one of three conditions for an additional 48 hr representing

either hypoxic (1%, ~10 mmHg O$_2$), normoxic (5%, ~35 mmHg O$_2$), or hyperoxic (20%, ~141 mmHg

O$_2$) environments relative to the first trimester of pregnancy (Figure 2A). Within all oxygen conditions,

explant outgrowth was observed (Figure 2B, 2C). However, column outgrowth was most pronounced

in 1% and 5% oxygen, where outgrowth area in both of these low oxygen conditions was significantly

greater than outgrowth observed in 20% oxygen (Figure 2C, 2D). While column outgrowth in 5%

oxygen was overall less variable and trended on producing smaller columns than explants cultured in

1% oxygen, there was no statistical difference between outgrowth in 1% and 5% oxygen (Figure 2D).

In summary, trophoblast column outgrowth was potentiated by low oxygen, whereas exposure to

atmospheric oxygen (20% oxygen) blunted column outgrowth.

Explant exposure to low or high levels of oxygen generate distinct transcriptomic signatures

To gain mechanistic insight into how hypoxic, physiologically normal, and high levels of

oxygen modulate trophoblast column outgrowth, global gene expression was analysed in placental

explant cultures using gene microarrays. For this experiment, placental explants from 5 unique

placentae (n=5; 5-7 weeks' gestation) were established as previously described except that cultures

were maintained in their respective oxygen condition (1%, 5%, or 20%) for 24 hr prior to RNA

isolation in order to capture molecular signatures central to column formation. Importantly, RNA was

extracted from only column trophoblasts and Matrigel-invading EVT; chorionic villi of explants were

carefully micro-dissected away from columns and EVT following an approach described within Bilban

et al (Bilban et al., 2009) (Figure 3A). Following standard probe filtering, and normalization

(Supplemental Figure 1), differential gene expression (DGE) analysis of explant trophoblasts was

performed. Using a false discover rate (FDR) < 0.05, we identified many differences in gene

expression between 1% versus 20% (293 genes up-regulated; 685 genes down-regulated) and 5%

versus 20% oxygen conditions (363 genes up-regulated; 406 genes down-regulated) (Figure 3B;

Supplemental Table 2). DGE analysis between 1% and 5% oxygen did not identify differentially

expressed genes (Figure 3B). Principal component analysis (PCA) shows clustering of explants

cultured in hypoxic, normoxic, and hyperoxic conditions (Figure 3C). PCA sample clustering showed

explant trophoblasts cultured in 1% and 5% oxygen generally clustered closer together, save for two

1% oxygen explant outliers; one clustered amongst 20% oxygen samples while the other clustered

separately to all other samples (Figure 3C). Despite these two explants being identified as significant

outliers using the Silhouette coefficient (Barghash and Arslan, 2016), we opted to retain them for the

remainder of our analyses as we could not confidently ascribe outlier classification due to technical or

batch-related artifacts.

Hierarchical clustering of the 15 column trophoblast samples segregated samples into two

statistically significant clusters (*sigclust*, p<0.05): a 20% oxygen dominated cluster and a cluster

comprised of column trophoblasts cultured in 1% and 5% oxygen (Figure 3D). A gene heat-map of the

top 40 differentially expressed genes (top 20 differentially expressed genes in 1%; top 20 differentially

expressed genes in 20%; FDR< 0.05, ranked by FC) highlights gene patterns across 1%, 5%, and 20%

oxygen cultures (Figure 3D). In both 1% and 5% oxygen conditions, the top hits identified by global

DGE analysis included genes associated with hypoxia (*EGLN3, RORA*), cell-matrix interaction/re-

structuring (*LOX, JAM2, EGLN3, PLAUR*), and gene transcription regulation (*MXI1, TSC22D3,*

RORA). In contrast, the most highly expressed genes in explants exposed to 20% oxygen were

exclusive to pro-mitotic/proliferative processes (*MKI67, KIF20A, KIF23, PEG10, CDK1, NCAPG,*

NCAPH, TOP2A, CDC7), indicating that column trophoblasts cultured in 20% oxygen possess a

proliferative phenotype.

While stringent DGE parameters did not identify differentially expressed genes between 1%

and 5% oxygen culture conditions, we did identify 123 and 53 unique genes to be up-regulated in 5%

and 1% oxygen cultures compared to samples cultured in 20% oxygen (Figure 3E). Further, 78 and 357

genes were shown to be uniquely down-regulated in 5% and 1% oxygen cultured explants compared to

20% cultures (Figure 3F). Gene ontology (GO) pathway analysis of the above 1% and 5% signatures

indicated that column trophoblasts exposed to 1% oxygen show enrichment for processes favoring

hypoxia-related- and oxidative stress-signaling (Supplemental Figure 2). Conversely, column

trophoblasts cultured in 5% oxygen showed enrichment of biological pathways linked to nucleotide

biosynthesis and metabolism, indicating that column trophoblasts in 1% and 5% oxygen do exhibit

underlying molecular differences potentially contributing to trophoblast column development

(Supplemental Figure 2).

Differing oxygen levels drive distinct molecular programs in trophoblast columns

To broadly examine how transcriptomic differences within column trophoblasts exposed to

hypoxic, normoxic, and hyperoxic conditions relate to differences in molecular pathways, gene

signatures determined by DGE analysis (FDR < 0.05; fold-change > 1.5; Supplemental Table 2) were

used to identify pathways enriched in explant column trophoblasts cultured in 1%, 5%, and 20%

oxygen. Unsurprisingly, pathways in explants cultured in 1% oxygen (293 genes; clusterProfiler)

showed enrichment for multiple pathways and molecular processes specific to hypoxia (Figure 4A).

Additionally, the 1% oxygen signature also showed enrichment for pathways related to extracellular

matrix (ECM) structure and organization, steroid hormone responses, and hydroxyproline metabolism

(Figure 4A). Similar to the 1% oxygen pathway readouts, the 5% oxygen signature (363 genes) showed

enrichment for genes specific to ECM composition, response to hypoxia, and response to steroid

hormones, but also showed enrichment of pathways related to bone development and viral entry into

cells (Supplemental Figure 3). By contrast, 20% oxygen (685 genes) showed enrichment of pathways

and cellular processes related to organelle fission, nuclear division, chromosome segregation, mitotic

nuclear division, and DNA packaging and replication, all of which link to heightened cell cycle activity

and proliferation (Figure 4B).

Using the Mitotic Nuclear Division, Chromosome Segregation, and DNA Replication curated

GO gene signatures enriched within 20% column trophoblasts (93 genes; Supplemental Table 3),

explant samples were subjected to hierarchical clustering and visualized by gene heat-map (Figure 4C).

Notably, samples segregated into two groups: One group defined by 1% oxygen samples (4/5 1%

oxygen samples), and the other group consisting of mostly of 5% and 20% oxygen samples (10/10

5%/20% samples) (Figure 4C). This later branch was further divided into two sub-branches, one

enriched by 5% oxygen samples also containing a 1% oxygen sample outlier, and the other sub-branch

made up entirely of 20% oxygen column trophoblasts (Figure 4C). Interestingly, gene heat-map

expression intensities suggest a step-wise increase in expression of pro-mitotic/proliferative genes in

column trophoblasts exposed to increasing levels of oxygen (Figure 4C). To verify if an increase in

exposure to oxygen tension translates into increased proliferation, a BrdU pulse-chase was performed

on a separate cohort of placental explants (n=3) cultured in 1%, 5%, and 20% oxygen. In support of the

gene array data, little/no evidence of cell proliferation within explant columns was observed in 1%

oxygen cultures following a 4 hr chase (Figure 4D, 4E). However, explants cultured in 5% oxygen

showed a significant increase in BrdU incorporation within column trophoblasts compared to 1%

oxygen columns, and an even greater level of BrdU positivity was measured within 20% oxygen

columns (20% versus 1%) (Figure 4D, 4E). Though a trend for greater proliferation in explant columns

cultured in 20% oxygen compared to 5% oxygen was observed, this difference was not significant

(Figure 4E). Overall, our findings suggest that explant column trophoblasts cultured in low oxygen up-

regulate molecular processes related to hypoxia/HIF1A signalling and ECM organization/remodeling,

while column trophoblasts exposed to hyper-physiological 20% oxygen adopt a predominantly pro-

proliferative phenotype.

Low oxygen promotes EVT differentiation

Differentiation of trophoblasts along the EVT pathway is in part defined by the exiting of PCT

located at the base of anchoring columns from the cell cycle (Velicky et al., 2018). Moreover, as

column trophoblasts located at distal portions of columns acquire pro-invasive EVT-like

characteristics, molecular pathways related to ECM-remodeling and protease functions are accordingly

up-regulated (Davies et al., 2016). Our observation that low oxygen (1% & 5%) promotes

transcriptional signatures linked to cell-ECM interaction and protease-ECM remodeling, while 20% oxygen promotes proliferation, suggests that exposure to low oxygen drives, while high oxygen restrains EVT differentiation.

To gain insight into how differing levels of oxygen affect column trophoblast differentiation, explant column trophoblasts were subjected to hierarchical cluster analysis using a signature of differentially expressed genes derived from a list of 47 trophoblast-related genes curated from recent high-dimensional data and differentiation studies focused on trophoblast biology (Supplemental Figure 4; Supplemental Table 4) (Bilban et al., 2009; Davies et al., 2016; Haider et al., 2018; Lee et al., 2018; Turco et al., 2018). This list includes genes associated with trophoblast lineage (*TFAP2A, KLF5, GATA3*), trophoblast pluripotency (*CDX2*), villous cytotrophoblast (CTB) state (*EGFR, SPINT1, ITGA6, PEG10, TEAD4, TP63*), EVT state (*HLA-G, HTRA1, LAIR2, FLT-1, ERBB2, ADAM12, AMAM19, MYC, ITGA5, TEAD2*), syncytiotrophoblast (SCT) state (*GCM1, CGA, ERVW1, ERVFRD-1, ENDOU*), and genes commonly used to identify proliferative CTB and PCT (*MK167, CCNA2, NOTCH1*). From this curated list, 14 genes were differentially expressed between 1%, 5%, and 20% oxygen cultured explants (Supplemental Figure 4A). Notably, this small signature was sufficient to segregate samples into two main groups: One group consisted almost entirely of column trophoblasts cultured in 20% oxygen (save for one 1% oxygen outlier), while the other group was made up of a mix of 1% and 5% oxygen cultured samples (Supplemental Figure 4A). Genes enriched within low-oxygen cultured explants included trophoblast lineage-related transcription factors (*GATA3*, *KLF5*, *TFAP2A*), genes related to ECM remodeling (*TIMP1*, *ADAM12, ADAM19*), genes related to CTB specification (*CDH1*, *EGFR*), and genes linked with the EVT sub-lineage (*ADAM12*, *FLT1*, *ITGA5, MYC*) (Supplemental Figure 4A). Samples that grouped primarily as 20% oxygen cultured explants showed enrichment for genes specific to proliferative CTB and PCT (*CCNA2*, *MKI67*). Notably, the imprinted paternally expressed gene, *PEG10,* was also highly expressed within 20% oxygen column trophoblasts

(Supplemental Figure 4A).

Examination of PEG10 localization within first trimester placental villi (n=3 placentae; 6-10

weeks' gestation) by immunofluorescence microscopy (IF) showed that PEG10 preferentially localizes

to CTB (Supplemental Figure 4B). IF localization of PEG10 within a new cohort of placental explants

(established from n=5 placentae) revealed that PEG10 signal, similar to the signal observed in placental

villi, is broadly localized to CTB in explants exposed to all three oxygen culture conditions

(Supplemental Figure 4C). However, within 20% oxygen explants, bright PEG10 signal was also

observed within multi-layered PCT; this column-specific signal was minimal/absent in 1% and 5%

oxygen cultures (Supplemental Figure 4C).

To more closely examine how global gene expression changes identified within low and high

oxygen exposed explants relate to trophoblast differentiation, we examined the expression of the top

fifteen up-regulated genes in column trophoblasts cultured in 1% and 20% oxygen (FDR < 0.05; fold-

change > 2) within a recently reported first trimester placenta single cell transcriptomic dataset (Vento-

Tormo et al., 2018). Using this dataset we focused exclusively on the 5 subtypes of trophoblasts that

were described (14,366 trophoblasts from 5 individual placentae): CTB, proliferative CTB, SCT,

proliferative EVT (likely PCT), and EVT (likely an admixture of DCT and invasive EVT) (Figure 5A)

(Vento-Tormo et al., 2018). Within UMAP-directed cell clusters, the specificity of each trophoblast

sub-lineage/type is shown by expression levels of *EGFR* (CTB), *ERVFRD-1* (syncytin-2; SCT), *HLA-G*

(column trophoblast & EVT), and *MKI67* (proliferating trophoblast) (Figure 5B). A heat-map

projection shows the pattern of gene expression of trophoblast lineage and subtype-specific genes (Pan

trophoblast: *KRT7, TFAP2A, GATA3, KLF5*; CTB: *EGFR, SPINT1, TP63*; proliferative CTB and PCT:

MKI67, CCNA2; SCT: *ENDOU, ERVFRD-1*; EVT: *HLA-G, ITGA5, ERBB2*), and the fifteen top up-

regulated genes in 20% (*OAS1, IFIT3, DLGAP5, GSTA3, CDK1, OXCT1, CENPF, KIF20A, PARP1,*

TOP2A, PLEKHH1, RFX5, NCPAH, KPNA2, NRP2) and 1% oxygen conditions (*TNFSF10, LOX,*

AKAP12, ACTC1, BIRC7, SPNS2, S100A4, JAM2, MXI1, EGLN3, RORA, PLAUR, AK4, TSC22D3,

NRN1) (Figure 5C). Interestingly, genes up-regulated within 1% oxygen-cultured columns showed

alignment with proliferative EVT, and this relationship was even greater with EVT (Figure 5C). In

contrast, the top genes identified within 20% oxygen columns aligned predominately with proliferative

CTB and proliferative EVT (Figure 5C).

Pseudotime trajectory analysis using a gene signature derived from the top 1000 variable genes

reproduced a lineage trajectory similar to that reported in Vento-Tormo *et al* (Vento-Tormo et al.,

2018), where a predicted cell origin state was identified as well as the two differentiation trajectories

aligning with the villous and extravillous pathways (Figure 5D). Correlating the expression of genes

specific to distinct states of trophoblast differentiation (i.e. *EGFR, HLA-G, ERVFRD-1*) with the

pseudotime trajectory confirmed the accuracy of the pseudotime modeling, where *EGFR* aligned to

cells committed to the villous pathway, *HLA-G* to a maturing EVT, and *ERVFRD-1* to a terminally

differentiated SCT (Figure 5E). Correlating trophoblast lineage trajectory with the top up-regulated 1%

and 20% oxygen genes showed that genes enriched within 20% oxygen aligned to both the villous and

extravillous pathway trajectories (Figure 5F). Notably, high-oxygen genes aligned closely with cell

states linked to proliferative CTB (*PARP1, RFX5, PLEKHH1, GSTA3, NRP2*) and proliferative PCT

(*OXCT1, KIF20A, NCAPH, CENPF, TOP2A, DLGAP5, CDK1, KPNA2*) (Figure 5F). By contrast, low

oxygen enriched genes primarily aligned with the tail-end of the extravillous pathway (i.e. *TNFSF10,*

LOX, SPNS2, S100A4, JAM2, PLAUR) (Figure 5F). Together, these data suggest that column

trophoblasts exposed to low oxygen adopt transcriptomic signatures that are reflective of EVT, while

column trophoblasts cultured in 20% oxygen express genes that align predominately with proliferating

CTB and PCT.

306 ***LOX* expression and activity is potentiated by low oxygen**

307   Our finding that low oxygen drives column outgrowth and potentiates the expression of genes

308 linked with the EVT phenotype suggests that genes highly expressed within low oxygen columns may

309 in part contribute to EVT differentiation and trophoblast column formation. Rank ordering of up-

310 regulated genes by fold-change in both 1% and 5% oxygen column trophoblasts identified multiple

311 conserved genes between the two oxygen conditions (Supplemental Table 2). Notably, *LOX*, the gene

312 encoding lysyl oxidase, a copper-dependent enzyme that catalyses collagen and elastin crosslinking,

313 was the number 2-ranked gene in both 1% and 5% oxygen cultured explants. Specifically, *LOX*

314 expression was 6.2- and 5.1-fold higher in 1% and 5% oxygen cultures compared to 20% oxygen

315 explants (Figure 6A; Supplemental Table 2). While previous work has identified a role for elevated

316 *LOX* expression in promoting tumor cell metastasis (Cox et al., 2015; Di Stefano et al., 2016), the role

317 of LOX in placental trophoblast column biology and trophoblast differentiation along the EVT pathway

318 has not been described.

319   As an initial step to examine the importance of LOX in anchoring column biology, *LOX* mRNA

320 *in situ* hybridization within first trimester placental villi (n=3; 6-8 weeks' gestation) was performed.

321 RNAscope *in situ* hybridization showed specific and intense *LOX* localization to cells within the

322 mesenchymal core of placental villi and to trophoblasts within anchoring columns (Figure 6B).

323 Little/no *LOX* signal was detected in CTB or SCT (Figure 6B). RNAscope analysis of *LOX* within

324 placental explants cultured in 1% and 20% oxygen supported the gene array finding that *LOX*

325 expression was elevated in column trophoblasts exposed to low oxygen (Figure 6C). While we were

326 unable to verify elevated LOX protein expression in low oxygen-cultured placental explants via IF

327 microscopy due to non-specific antibody signal, LOX enzymatic activity, measured in conditioned

328 media (CM) generated by placental explants cultured in 1% or 20% oxygen, showed that activity was

329 significantly higher in 1% cultures (Figure 6D). Use of the LOX inhibitor, β-aminopropionitrile

(BAPN), demonstrated LOX-substrate specificity, while recombinant active LOX served as a positive

control (Figure 6D). Taken together, these findings show that LOX expression and activity is elevated

in column trophoblasts cultured in low oxygen. Further, *LOX's* preferential *in vivo* expression within

the trophoblast anchoring column combined with its involvement in promoting tumor cell metastasis,

suggests that LOX may also play a role in controlling trophoblast column outgrowth and/or EVT

differentiation.

Impairment of LOX restrains column outgrowth

To test the function of LOX in column outgrowth, Matrigel-imbedded placental explants

cultured in 1% oxygen (n=3) were cultured in either control explant media or media containing the

competitive LOX inhibitor BAPN. The effectiveness of BAPN in inhibiting LOX activity was

measured as before by examining the ability of endogenous LOX in explant CM to oxidize substrate

(Figure 7A). Conditioned media harvested from control explants showed LOX activity levels slightly

below levels measured in recombinant LOX positive control reactions, but significantly higher than

explant media alone (Figure 7A). Importantly, treatment of explants with BAPN significantly blunted

LOX activity, though activity was not completely blocked (Figure 7A). Importantly, placental explant

treatment with BAPN led to a significant two-fold impairment in column outgrowth (Figure 7B, 7C).

Taken together, these results suggest that LOX expression in developing trophoblast columns promotes

column outgrowth and associates with an EVT phenotype.

DISCUSSION

Here we describe how exposure to different levels of oxygen differentially affect trophoblast column outgrowth and global gene expression. We provide evidence that exposure to low oxygen results in overall increases in column outgrowth accompanied by gene expression signatures that align with an EVT phenotype. By contrast, gene signatures in high oxygen-cultured column trophoblasts define a role for elevated oxygen in maintaining column growth through cell proliferation. In both hypoxic and normoxic conditions, we identify the gene *LOX*, as one of the most highly up-regulated genes within explant columns. We show that *LOX* expression associates with EVT lineage trajectory and demonstrate that impairment of LOX activity blunts column outgrowth. Together, this work supports a role for low oxygen in potentiating the EVT pathway. Moreover, this work also identifies novel oxygen-sensitive molecular processes that likely play roles in anchoring column formation during human placental development.

The role of oxygen in controlling column formation and the EVT pathway is controversial. Previous studies have shown that exposure of placental explants to low oxygen (i.e. 2% to 3% oxygen) promotes column expansion and outgrowth, where outgrowth is primarily attributed to HIF1A-directed cell proliferation (Caniggia et al., 2000; Genbacev et al., 1997). Consistent with this, evidence exists that CTB exposure to low oxygen restrains trophoblast progression along the EVT pathway (Caniggia et al., 2000; Genbacev et al., 1997; Lash et al., 2006). By contrast, rodents and rodent-derived trophoblast stem cells engineered to lack hypoxia-sensing machinery (i.e. *ARNT*, *HIF1A,* and/or *EPSA1* null) fail to differentiate into trophoblast lineages of the labyrinth zone and into trophoblast giant cells, trophoblast populations akin to invasive EVT in humans (Chakraborty et al., 2016; Chakraborty et al., 2011; Cowden Dahl et al., 2005; Gultice et al., 2009; Maltepe et al., 2005). In support of these observations, low oxygen was shown to potentiate an invasive phenotype in human primary trophoblasts with an accompaniment in the expression of EVT-associated genes HLA-G and α5

integrin, and the up-regulation of pro-migratory integrin-linked kinase signaling (Horii et al., 2016;

Robins et al., 2007; Wakeland et al., 2017). Our findings overall align with these later studies that

suggest low oxygen promotes differentiation along the EVT pathway. Notably, we show that low

oxygen drives expression of EVT-related genes and signatures that align with EVT lineage trajectory.

We show that low oxygen (1% and 5% oxygen) induces expression of hallmark EVT genes like

ITGA5, *ADAM12*, and *FLT1*, as well as the transcription factors *KLF5* and *GATA3* that are

preferentially expressed by EVT. Importantly, the use of cell lineage trajectory modeling using single

cell RNA sequencing data also provides evidence that low oxygen promotes a cell state consistent with

the EVT lineage. These later observations are in line with the association of low levels of oxygen

within the intervillous space and anchoring column formation and interstitial EVT infiltration into

decidual mucosa in early pregnancy. However, the relationship between relative placental hypoxia and

inadequate placentation in aberrant pregnancy conditions like preeclampsia (Farrell et al., 2019)

suggests that the impact of oxygen on trophoblast biology may differ according to stage of

development.

Our finding that high levels of oxygen promote column trophoblast proliferation is inconsistent

with previous studies investigating the role of hypoxia in anchoring column biology (Caniggia et al.,

2000; Genbacev et al., 1997). However, in agreement with our findings, James *et al* reported that

explant columns cultured in 8% oxygen have increased trophoblast cellularity compared with columns

exposed to 1.5% oxygen (James et al., 2006). Moreover, there seems to be consistent agreement that

exposure to low oxygen leads to greater column outgrowth, where oxygen tension levels between 1-5%

consistently generate larger columns than atmospheric oxygen conditions (Caniggia et al., 2000; James

et al., 2006). However, dissecting how enhanced column outgrowth is achieved, i.e. via greater cell

proliferation within the column or through increased trophoblast migration and invasion, is still not

completely resolved. Indeed, while our data suggests that low oxygen promotes an EVT phenotype and

enhances molecular pathways related to cell-ECM remodeling, we did not directly examine if low

oxygen affects EVT invasion. The stark differences on the effect of oxygen levels in regulating

trophoblast proliferation and EVT differentiation within explant columns between various studies is

difficult to explain, but may stem from differences in media composition, the matrix substratum used in

explant cultures, subtle variations in oxygen tension, and the gestational age of placental tissues/cells

used for establishing explant cultures. Future studies will need to specifically re-examine how differing

levels of oxygen impact column cell proliferation.

PEG10, a paternally expressed and maternally imprinted gene, was identified as a top up-

regulated gene within 20% oxygen column trophoblasts. Previous work has shown a critical role for

Peg10 in mouse placental development, where mice deficient in Peg10 show severe fetal growth

restriction and fetal demise by E10.5, in addition to defects in labyrinth and spongiotrophoblast

development (Ono et al., 2006). Intriguingly, a partial loss in *Peg10* imprinting leading to *Peg10* over-

expression also associates with impaired labyrinth formation (Koppes et al., 2015). These findings

suggest that a fine balance in Peg10 expression is required for proper placental development. Notably,

in humans, dysregulation of PEG10 associates with certain pregnancy disorders, including pre-

eclampsia, gestational hypertension, molar pregnancy, and spontaneous miscarriage (Dória et al., 2010;

Liang et al., 2014; Rahat et al., 2017). Our finding that *PEG10* expression and cell proliferation in

trophoblast columns associates with exposure to high oxygen is consistent with studies showing a role

for PEG10 in promoting tumor cell survival and growth (Bang et al., 2015; Ishii et al., 2017; Peng et

al., 2017). Recent findings also provide evidence that PEG10 promotes trophoblast cell line

proliferation (Abed et al., 2019). Given that PEG10 specifically localizes to CTB of floating villi and to

subsets of trophoblasts within proximal regions of anchoring columns suggests that PEG10 may play

roles in column establishment by promoting progenitor expansion. Further work is needed to dissect

the role of PEG10 in column formation and progression along the EVT pathway.

Our finding that *LOX* expression was consistently a gene highly up-regulated in both 1% and

5% column trophoblasts compared to those cultured in 20% oxygen indicates that LOX-related

processes are important in placental development and trophoblast column biology. While a role of

LOX in column formation has not been previously reported, the importance of LOX in tumorigenesis is

known, where elevated LOX associates with numerous types of cancers and LOX activity promotes

tumor cell metastasis (Cox et al., 2015; Di Stefano et al., 2016). Further, a recent report does provide

evidence that LOX expression promotes cell invasion of a trophoblast cell line (Xu et al., 2019). This

later finding is consistent with the association of LOX expression and the EVT phenotype. Aldehydes

produced by LOX-directed oxidation of lysine residues within collagen and elastin facilitate

collagen/elastin cross-linking and stability, which in turn provide a structural lattice for cell movement

(Kim et al., 2014). That *LOX* expression is greatest within both proximal and distal column

trophoblasts is interesting, as PCT are not considered to be migratory. Nonetheless, it is likely that

creating an appropriate substratum scaffold via LOX-directed collagen crosslinking may contribute to

column stability and provide a platform for EVT outgrowth. How or if LOX affects anchoring column

establishment or EVT differentiation is unknown, but future studies utilizing newly-derived trophoblast

organoids (Haider et al., 2018; Turco et al., 2018) may allow for deep mechanistic examination of LOX

and the EVT pathway.

In summary, the extravillous pathway is controlled by multiple intrinsic as well as extrinsic

factors, including level of oxygen. We provide evidence that supports a role for low oxygen levels in

promoting the differentiation of trophoblasts along the EVT pathway. This finding establishes insight

into critical developmental events during placentation that occur in early pregnancy. Further, these

findings may also provide a foundation for understanding cellular and molecular processes contributing

to conditions linked to aberrant placentation.

MATERIALS AND METHODS

Patient recruitment and tissue collection

Decidual and placental tissues were obtained with approval from the Research Ethics Board on the use

of human subjects, University of British Columbia (H13-00640). All samples were collected from

women (19 to 39 years of age) providing written informed consent undergoing elective terminations of

pregnancy at British Columbia's Women's Hospital, Vancouver, Canada. First trimester decidual

(N=1) and placental tissues (N=30) were collected from participating women (gestational ages ranging

from 5–12 weeks) having confirmed viable pregnancies by ultrasound-measured fetal heartbeat. The

decidual tissue sample was selected based on the presence of a smooth uterine epithelial layer and a

textured thick spongy underlayer. Patient clinical characteristics i.e. height and weight were

additionally obtained to calculate body mass index (BMI: kg/m^2) and all consenting women provided

self-reported information via questionnaire to having first hand exposure to cigarette smoke, and are

summarized in Supplemental Table 1.

Placental villous explant assay

 Ex vivo placental villous cultures were established as described in (Aghababaei et al., 2014; De

Luca et al., 2017; Perdu et al., 2016). Briefly, placental villi from 5-8 week old gestation placentas

(n=8) obtained from patients undergoing elective termination of pregnancy were dissected, washed in

cold PBS, and imbedded into Millicell cell culture inserts (0.4 μm pores, 12mm diameter. EMD

Millipore, Billerica, MA) containing 200 μl of growth-factor-reduced Phenol-red free Matrigel (BD

Biosciences, San Diego, CA). Explants, containing 400 μl DMEM/F12 1:1 (200 mM L-glutamine) in

the outer chamber, were allowed to establish overnight in a humidified 37 °C trigas incubator at 5%

oxygen, 5% CO_2. Following 24 hr of culture, explants were cultured in 200 μl DMEM/F12 1:1 media

and placed into 1%, 5%, or 20% oxygen incubators for either an additional 24 hr (gene expression

analyses) or 48 hr (explant outgrowth measurements). All explant media were supplemented with

penicillin/streptomycin and antimycotic solution (Thermofisher Scientific, Waltham, MA) Growing

explants were imaged at indicated times using a Nikon SMZ 7454T triocular dissecting microscope

(Minato, Japan) outfitted with a digital camera. EVT outgrowths were measured by ImageJ software.

Fold-change in outgrowth was determined by dividing the mean column area at 48 hr into the mean

area at 0 hr.

Immunofluorescence, RNAScope, and immunohistochemistry microscopy

Immunofluorescence: Placental villi (6-12 weeks gestation; n=5) or placental explants (derived

from n=11 placentae) were fixed in 2% paraformaldehyde overnight at 4 °C. Tissues were paraffin

embedded and sectioned at 6 μm onto glass slides. Immunofluorescence was performed as described

elsewhere (Aghababaei et al., 2015). Briefly, cells or placental tissues underwent antigen retrieval by

heating slides in a microwave for 5 X 2 minute intervals in a citrate buffer (pH 6.0). Sections were

incubated with sodium borohydride for 5 minutes, RT, followed by Triton X-100 permeabilization for

5 minutes, RT. Slides were blocked in 5% normal goat serum/0.1% saponin for 1h, RT, and incubated

with combinations of the indicated antibodies overnight, 4 °C: Anti-HLA-G (1:100, 4H84, Exbio,

Vestec, Czech Republic); anti-cytokeratin 7, mouse monoclonal IgG (1:50, RCK105, Santa Cruz

Biotechnology, Dallas, TX); anti-cytokeratin 7, rabbit monoclonal IgG (1:50, SP52, Ventana Medical

Systems, Oro Valley, AZ); anti-Ki67 (1:75, SP6, Thermo Fisher Scientific); anti-LOX (1:100, NB100-

2527, NovusBio, Centennial, CO); anti-PEG10 (1:100, 4C10A7, NovusBio); anti-BrdU (1:1000,

Bu20a, Cell Signalling Technology, Danvers, MA). Following overnight incubation, sections and

coverslips were washed with PBS and incubated with Alexa Fluor goat anti rabbit-488/-568 and goat

anti mouse-488/-568 conjugated secondary antibodies (Life Technologies, Carlsbad, CA) for 1 hr at RT

and washed in PBS, and mounted using ProLong Gold mounting media containing DAPI (Life

Technologies).

Slides were imaged with an AxioObserver inverted microscope (Car Zeiss, Jena, Germany)

using 20X Plan-Apochromat/0.80NA or 40X Plan-Apochromat oil/1.4NA objectives (Carl Zeiss). An

ApoTome .2 structured illumination device (Carl Zeiss) set at full Z-stack mode and 5 phase images

was used for image acquisition. For quantification of BrdU signal in column trophoblasts, 2-4 cell

columns per explant (n=9; obtained using a 20X objective) were used to calculate values; BrdU+ cell

proportions were calculated by BrdU/KRT7+ cells per column; BrdU fluorescence intensity thresholds

were used to calculate BrdU proportions within Matrigel explant columns. Images were obtained using

an Axiocam 506 monochrome digital camera and processed and analyzed using ZenPro software (Carl

Zeiss).

Immunohistochemistry: Decidua (10 weeks' gestation; n=1) from a first trimester pregnancy

was fixed in 2% paraformaldehyde for 24 hr at 4°C, paraffin imbedded, and serially sectioned at 6μm

onto glass slides. Heat-induced antigen retrieval was performed using sodium citrate (10mM, pH 6.0)

followed by quenching endogenous peroxidases with 3% hydrogen peroxide for 30 minutes at RT.

Sections were then permeabilized with 0.2% Triton-X-100 for 5 minutes at RT. Serum block was

performed with 5% BSA in tris-buffered saline with 0.05% Tween 20 (TBST). Sections were then

incubated with mouse monoclonal HLA-G (1:100, clone 4H84, ExBio) diluted in TBST overnight at

4°C. Following overnight incubation, sections were incubated with Envision+ Dual Link Mouse/Rabbit

HRP-linked secondary antibody (DAKO, Santa Clara, CA) for 1 hr at RT. IgG1 isotype controls and

secondary antibody-only negative controls were performed to confirm antibody specificity. Staining

was developed via 3,3'-diaminobenzidine (DAB) chromogen (DAB Substrate Kit, Thermo Scientific),

counterstained in Modified Harris Hematoxylin Solution (Sigma-Aldrich, St. Louis, MO) and

coverslips mounted with Entellan mounting medium (Electron Microscopy Sciences, Hatfield, PA).

RNAScope: RNA *in situ* hybridization was performed using RNAscope® 2.5 HD Assay-RED

(Advanced Cell Diagnostics, Newark, CA) following the manufacturer's instructions (Wang *et al.*, 2012). Briefly, placenta (n = 3; 6-8 weeks' gestation) and Matrigel-imbedded placental explants (n=15; 5 explants per oxygen condition) from first trimester pregnancies were fixed overnight at 4°C in 4% paraformaldehyde and paraffin-embedded. Tissue sections were serially sectioned at 6 μm, and following deparaffinization, antigen retrieval was performed using RNAscope® Target Retrieval Reagent (95°C for 15 minutes) and ACD Protease Plus Reagent (40°C for 30 minutes). RNAScope probes targeting LOX (Hs-LOX-C2; Ref. 415941-C2) and negative control (Probe-dapB; Ref. 310043) were incubated on sections for 2hr at 40°C, and following this, the RNAscope signal was amplified over 6 rounds of ACD AMP 1-6 incubation and application of RED-A and RED-B at a ratio of 1 volume of RED-B to 60 volumes of RED-A for 10 minutes at RT. Selected samples were counterstained with 50% Hematoxylin I for 2 minutes at RT and all samples were mounted using EcoMount (Biocare Medical, Pacheco, CA).

RNA purification

Total RNA was prepared from column and invasive EVT cells using TRIzol reagent (Life Technologies, Carlsbad, CA) followed by RNeasy MinElute Cleanup (Qiagen, Hilden, Germany) and DNase treatment (Life Technologies) according to the manufacturer's instructions. Care was taken so that explants were exposed to atmospheric oxygen no longer than 2 minutes during microscopic dissection and separation of placental villi away from columns prior to the addition of TRIzol. RNA purity was confirmed using a NanoDrop Spectrophotometer (Thermo Fisher Scientific) and by running RNA samples on an Agilent 2100 Bioanalyzer (Agilent Technologies, Santa Clara, CA). Only RNA samples having an RNA Integrity Number (RIN) > 8.0 were used.

Microarray hybridization, gene array data preprocessing, gene expression analysis

Total RNA samples extracted from explant columns were sent to Génome Québec Innovation

Centre (McGill University, Montréal, Canada) for RNA quantification. Briefly, RNA samples were

prepared for transcriptome profiling using the GeneChip ™ Pico Reagent Kit (Thermo Fisher

Scientific) as per manufacturer's protocol. Samples were run on the Clariom™ S Human Array to

measure gene expression at >20,000 genes in the human genome (Affymetrix, Santa Clara, CA). Raw

data generated from the arrays were read into R statistical software (version 3.5.1) with the

Bioconductor *oligo* package to convert raw Affymetrix CEL files into an expression matrix of intensity

values. The expression data was background corrected, Quantile normalized, and log-transformed. A

total of 13,787 control, duplicated, non-annotated or low intensity probes were filtered out of the data,

leaving 13,402 probes for further analysis. Pre-processing was monitored at each step by Principal

Component Analysis (PCA) and linear modelling. Principal component analysis was performed by the

`svd()` function from the *sva* package in R. Linear modelling was conducted using the R package

limma. Probe-wise variances were shrunk using empirical Bayes with the `eBayes` function, followed

by FDR adjustment for multiple testing (Benjamini and Hochberg, 1995). Differentially expressed

genes were defined based on an FDR < 0.05. Enrichment of pathways were identified and annotated

using the *clusterProfiler* package in Bioconductor (Carvalho and Irizarry, 2010). Volcano plots were

generate using the *ggplot2* package for RStudio. Cluster analysis of sample relations based on principal

components was generated using the `plotSampleRelation` function for the *Lumi* package. A

hierarchical cluster analysis was conducted using Euclidean distances on the top 40 differentially

expressed genes, selected by FDR <0.05 and ranked by fold-change > 1.5.

Single cell RNA-seq data analysis

565   Processed droplet-based and Smart-seq2 single cell RNA sequencing data was obtained from public

repositories (ArrayExpress experiment codes: E-MTAB-6701 and E-MTAB-6678) (Vento-Tormo et al., 2018). Cells belonging to clusters from trophoblast lineages (n=14,366) were merged from droplet-based and Smart-seq2 data, and corrected for batch-effects using canonical correlation analysis implemented in the R package *Seurat* (Butler et al., 2018). Pseudotime trajectory modelling was conducted using the *monocle 2* R package (Qiu et al., 2017; Trapnell et al., 2014) under the recommended unsupervised procedure called "`dpFeature`". Briefly, the first 10 principal components on log-normalized expression data were used to construct a TSNE projection, upon which density-peak clustering determined 13 number of clusters, using parameters rho = 2, delta = 10. The top 1000 differentially expressed genes between these clusters were then used for ordering the cells. Visualization of oxygen concentration -dependent genes were visualized along the inferred pesudotime-ordered branches using the R function `monocle::plot_genes_branched_heatmap` with the following settings: the number of clusters k = 5 for clustering the genes, and default parameters for all else.

LOX activity assay and LOX inhibition

Measurement of endogenous LOX activity in placental explant conditioned media was performed following a modified protocol described in Wiel et al (Wiel et al., 2013). Briefly, 600 μl of conditioned media from placental explants cultured in triplicate from either 1%, 5%, or 20% oxygen conditions was pooled, concentrated 12-fold using 7.5 kDa exclusion Amicon Millipore concentration columns (Millipore, Burlington, MA), and snap-frozen in liquid nitrogen. LOX activity in 15 μl of concentrated conditioned media was determined using the Amplex Red H_2O_2 detection kit following the manufacture's instructions (Life Technologies, Carlsbad, CA). This assay is based on the ability of endogenous LOX to oxidize 10 mM DAP (a LOX substrate) in the presence of 0.5 U/ml horseradish peroxidase; the reaction is incubated at 37 °C for 30 minutes. Oxidation of Amplex Red generates a

fluorescence signal measurable at 560nm/590nm excitation/emission wavelengths and was detected

using a fluorescence plate reader (BMG Labtech, Ortenberg, Germany) using a 96-well format. As a

positive control, 10 μg/ml recombinant active LOX (MyBioscource.com, San Diego, CA) was

separately incubated with DAP substrate. LOX activity/specificity was determined by co-incubating

reactions with 5 mM of BAPN. For endogenous inhibition of LOX within explant cultures, following

24 hr of explant establishment at 37 °C, 5% CO2, 5% O2 in a humidified tri-gas incubator, explant

media was replaced with media containing 500 μM of BAPN and explant cultures were placed into 1%

or 20% O_2 culture conditions for an additional 48 hr of culture prior to being imaged/measured.

BrdU incorporation assay

Pulse-chase labeling with BrdU (Sigma-Aldrich) was conducted on a verification cohort of

Matrigel embedded placental explants (n=3 distinct placentae, 4-11 columns per oxygen condition).

Explants were established in 5% oxygen for 24 hr followed by 24 hr of culture in either 1%, 5% or

20% oxygen. After 48 hr of culture, explants were exposed to a 4-hr pulse with culture media

containing 10μM of BrdU. Following 4 hr of labelling, explants were washed in PBS and fixed in 4%

PFA overnight. Explants were paraffin embedded and sectioned for immunofluorescence microscopy.

Immunofluorescent staining with anti-BrdU antibody (Bu20a, Sigma-Aldrich) with the addition of a

30-minute incubation in 2M hydrochloric acid between permeabilization and sodium borohydride

steps.

Statistical analysis

Data are reported as median values with standard deviations. All calculations were carried out

using GraphPad Prism software (San Diego, CA). For single comparisons, Mann-Whitney non-

parametric unpaired t-tests were performed. For multiple comparisons, one-way Kruskal-Wallis

ANOVA followed by Dunn's multiple comparison test was performed on explant outgrowth data as

outgrowth in 1% oxygen was not normally distributed. One-way ANOVA followed by Tukey post-test

were performed for all other multiple comparisons. The differences were accepted as significant at $P <$

0.05. For gene microarray and scRNA-seq statistical analyses, please refer to the Gene Array Data

Preprocessing and scRNA-seq analysis sections in methods.

DATA AVAILABILITY/ACCESSION NUMBER

The GEO accession number for the data reported in this paper is: GSE132421

AUTHOR CONTRIBUTIONS

FUNDING

This work was supported by a Natural Sciences and Engineering Research Council of Canada

(NSERC; RGPIN-2014-04466) Discovery Grant (to AGB), Canadian Institutes of Health Research

(CIHR; 201403MOP-325905; 201809PJT-407571-CIA-CAAA) operating grants to AGB, and a CIHR

Master's graduate studentship (to JT).

ACKNOWLEDGEMENTS

The authors extend their sincere gratitude to the hard work of staff at British Columbia's

Women's Hospital's CARE Program for recruiting participants to our study, and thank Dr. Megan K.

Barker, for her critical readings of the manuscript.

COMPETING INTERESTS

The authors declare that no competing interests exist.

REFERENCES

Abed, M., Verschueren, E., Budayeva, H., Liu, P., Kirkpatrick, D. S., Reja, R., Kummerfeld, S. K., Webster, J. D., Gierke, S., Reichelt, M., et al. (2019). The Gag protein PEG10 binds to RNA and regulates trophoblast stem cell lineage specification. *PLoS ONE* **14**, e0214110.

Aghababaei, M., Hogg, K., Perdu, S., Robinson, W. P. and Beristain, A. G. (2015). ADAM12-directed ectodomain shedding of E-cadherin potentiates trophoblast fusion. *Cell Death Differ.* **22**, 1970–1984.

Aghababaei, M., Perdu, S., Irvine, K. and Beristain, A. G. (2014). A disintegrin and metalloproteinase 12 (ADAM12) localizes to invasive trophoblast, promotes cell invasion and directs column outgrowth in early placental development. *Mol. Hum. Reprod.* **20**, 235–249.

Avagliano, L., Marconi, A. M., Romagnoli, S. and Bulfamante, G. P. (2012). Abnormal spiral arteries modification in stillbirths: the role of maternal prepregnancy body mass index. **25**, 2789–2792.

Bang, H., Ha, S. Y., Hwang, S. H. and Park, C.-K. (2015). Expression of PEG10 Is Associated with Poor Survival and Tumor Recurrence in Hepatocellular Carcinoma. *Cancer Res Treat* **47**, 844–852.

Becht, E., McInnes, L., Healy, J., Dutertre, C.-A., Kwok, I. W. H., Ng, L. G., Ginhoux, F. and Newell, E. W. (2018). Dimensionality reduction for visualizing single-cell data using UMAP. *Nat. Biotechnol.* **37**, 38–44.

Benjamini, Y. and Hochberg, Y. (1995). Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society: Series B (Methodological)* **57**, 289–300.

Bilban, M., Haslinger, P., Prast, J., Klinglmüller, F., Woelfel, T., Haider, S., Sachs, A., Otterbein, L. E., Desoye, G., Hiden, U., et al. (2009). Identification of novel trophoblast invasion-related genes: heme oxygenase-1 controls motility via peroxisome proliferator-activated receptor gamma. *Endocrinology* **150**, 1000–1013.

Butler, A., Hoffman, P., Smibert, P., Papalexi, E. and Satija, R. (2018). Integrating single-cell transcriptomic data across different conditions, technologies, and species. *Nat. Biotechnol.* **36**, 411–420.

Caniggia, I., Mostachfi, H., Winter, J., Gassmann, M., Lye, S. J., Kuliszewski, M. and Post, M. (2000). Hypoxia-inducible factor-1 mediates the biological effects of oxygen on human trophoblast differentiation through TGFbeta(3). *J. Clin. Invest.* **105**, 577–587.

Carvalho, B. S. and Irizarry, R. A. (2010). A framework for oligonucleotide microarray preprocessing. *Bioinformatics* **26**, 2363–2367.

Chakraborty, D., Cui, W., Rosario, G. X., Scott, R. L., Dhakal, P., Renaud, S. J., Tachibana, M., Rumi, M. A. K., Mason, C. W., Krieg, A. J., et al. (2016). HIF-KDM3A-MMP12 regulatory circuit ensures trophoblast plasticity and placental adaptations to hypoxia. *Proc. Natl. Acad. Sci. U.S.A.* **113**, E7212–E7221.

Chakraborty, D., Rumi, M. A. K., Konno, T. and Soares, M. J. (2011). Natural killer cells direct
hemochorial placentation by regulating hypoxia-inducible factor dependent trophoblast lineage
decisions. *Proc. Natl. Acad. Sci. U.S.A.* **108**, 16295–16300.

Cowden Dahl, K. D., Fryer, B. H., Mack, F. A., Compernolle, V., Maltepe, E., Adelman, D. M.,
Carmeliet, P. and Simon, M. C. (2005). Hypoxia-inducible factors 1alpha and 2alpha regulate
trophoblast differentiation. *Molecular and Cellular Biology* **25**, 10479–10491.

Davies, J. E., Pollheimer, J., Yong, H. E. J., Kokkinos, M. I., Kalionis, B., Murthi, P., Davies, J.
E., Yong, H. E. J., Kokkinos, M. I., Kalionis, B., et al. (2016). Epithelial-mesenchymal transition
during extravillous trophoblast differentiation. *Cell Adhesion & Migration* **10**, 310–321.

De Luca, L. C., Le, H. T., Mara, D. L. and Beristain, A. G. (2017). ADAM28 localizes to HLA-
G(+) trophoblasts and promotes column cell outgrowth. *Placenta* **55**, 71–80.

Dória, S., Sousa, M., Fernandes, S., Ramalho, C., Brandão, O., Matias, A., Barros, A. and
Carvalho, F. (2010). Gene expression pattern of IGF2, PHLDA2, PEG10 and CDKN1C imprinted
genes in spontaneous miscarriages or fetal deaths. *Epigenetics* **5**, 444–450.

Farrell, A., Alahari, S., Ermini, L., Tagliaferro, A., Litvack, M., Post, M. and Caniggia, I. (2019).
Faulty oxygen sensing disrupts angiomotin function in trophoblast cell migration and predisposes
to preeclampsia. *JCI Insight* **4**.

Fock, V., Plessl, K., Draxler, P., Otti, G. R., Fiala, C., Knöfler, M. and Pollheimer, J. (2015a).
Neuregulin-1-mediated ErbB2-ErbB3 signalling protects human trophoblasts against apoptosis to
preserve differentiation. *J. Cell. Sci.* **128**, 4306–4316.

Fock, V., Plessl, K., Fuchs, R., Dekan, S., Milla, S. K., Haider, S., Fiala, C., Knöfler, M. and
Pollheimer, J. (2015b). Trophoblast subtype-specific EGFR/ERBB4 expression correlates with
cell cycle progression and hyperplasia in complete hydatidiform moles. *Hum. Reprod.* **30**, 789–
799.

Genbacev, O., Zhou, Y., Ludlow, J. W. and Fisher, S. J. (1997). Regulation of human placental
development by oxygen tension. *Science* **277**, 1669–1672.

Gultice, A. D., Kulkarni-Datar, K. and Brown, T. L. (2009). Hypoxia-inducible factor 1alpha
(HIF1A) mediates distinct steps of rat trophoblast differentiation in gradient oxygen. *Biology of*
Reproduction **80**, 184–193.

Haider, S., Meinhardt, G., Saleh, L., Fiala, C., Pollheimer, J. and Knöfler, M. (2016). Notch1
controls development of the extravillous trophoblast lineage in the human placenta. *Proc. Natl.*
Acad. Sci. U.S.A. **113**, E7710–E7719.

Haider, S., Meinhardt, G., Saleh, L., Kunihs, V., Gamperl, M., Kaindl, U., Ellinger, A., Burkard,
T. R., Fiala, C., Pollheimer, J., et al. (2018). Self-Renewing Trophoblast Organoids Recapitulate
the Developmental Program of the Early Human Placenta. *Stem Cell Reports* **11**, 537–551.

Horii, M., Li, Y., Wakeland, A. K., Pizzo, D. P., Nelson, K. K., Sabatini, K., Laurent, L. C., Liu,
Y. and Parast, M. M. (2016). Human pluripotent stem cells as a model of trophoblast
differentiation in both normal development and disease. *Proc. Natl. Acad. Sci. U.S.A.* **113**, E3882–

91.

Ishii, S., Yamashita, K., Harada, H., Ushiku, H., Tanaka, T., Nishizawa, N., Yokoi, K., Washio,
M., Ema, A., Mieno, H., et al. (2017). The H19-PEG10/IGF2BP3 axis promotes gastric cancer
progression in patients with high lymph node ratios. *Oncotarget* **8**, 74567–74581.

James, J. L., Stone, P. R. and Chamley, L. W. (2006). The effects of oxygen concentration and
gestational age on extravillous trophoblast outgrowth in a human first trimester villous explant
model. *Human Reproduction* **21**, 2699–2705.

Jauniaux, E., Watson, A. and Burton, G. (2001). Evaluation of respiratory gases and acid-base
gradients in human fetal fluids and uteroplacental tissue between 7 and 16 weeks' gestation. *YMOB*
184, 998–1003.

Kabir-Salmani, M., Shiokawa, S., Akimoto, Y., Sakai, K. and Iwashita, M. (2004). The role of
alpha(5)beta(1)-integrin in the IGF-I-induced migration of extravillous trophoblast cells during the
process of implantation. *Molecular Human Reproduction* **10**, 91–97.

Koppes, E., Himes, K. P. and Chaillet, J. R. (2015). Partial Loss of Genomic Imprinting Reveals
Important Roles for Kcnq1 and Peg10 Imprinted Domains in Placental Development. *PLoS ONE*
10, e0135202.

Lash, G. E., Otun, H. A., Innes, B. A., Bulmer, J. N., Searle, R. F. and Robson, S. C. (2006). Low
oxygen concentrations inhibit trophoblast cell invasion from early gestation placental explants via
alterations in levels of the urokinase plasminogen activator system. *Biol. Reprod.* **74**, 403–409.

Lee, C. Q. E., Turco, M. Y., Gardner, L., Simons, B. D., Hemberger, M. and Moffett, A. (2018).
Integrin α2 marks a niche of trophoblast progenitor cells in first trimester human placenta.
Development **145**, dev162305.

Liang, X. Y., Chen, X., Jin, Y. Z., Chen, X. O. and Chen, Q. Z. (2014). Expression and significance
of the imprinted gene PEG10 in placenta of patients with preeclampsia. *Genet. Mol. Res.* **13**,
10607–10614.

Maltepe, E., Krampitz, G. W., Okazaki, K. M., Red-Horse, K., Mak, W., Simon, M. C. and
Fisher, S. J. (2005). Hypoxia-inducible factor-dependent histone deacetylase activity determines
stem cell fate in the placenta. *Development* **132**, 3393–3403.

Newby, D., Marks, L., Cousins, F., Duffie, E. and Lyall, F. (2005). Villous explant culture:
characterization and evaluation of a model to study trophoblast invasion. *Hypertens Pregnancy* **24**,
75–91.

Ono, R., Nakamura, K., Inoue, K., Naruse, M., Usami, T., Wakisaka-Saito, N., Hino, T., Suzuki-
Migishima, R., Ogonuki, N., Miki, H., et al. (2006). Deletion of Peg10, an imprinted gene
acquired from a retrotransposon, causes early embryonic lethality. *Nat. Genet.* **38**, 101–106.

Peng, Y.-P., Zhu, Y., Yin, L.-D., Zhang, J.-J., Wei, J.-S., Liu, X., Liu, X.-C., Gao, W.-T., Jiang,
K.-R. and Miao, Y. (2017). PEG10 overexpression induced by E2F-1 promotes cell proliferation,
migration, and invasion in pancreatic cancer. *J. Exp. Clin. Cancer Res.* **36**, 30.

Perdu, S., Castellana, B., Yooona, K., Chan, K., DeLuca, L. and Beristain, A. G. (2016). Maternal
obesity drives functional alterations in uterine NK cells. *JCI Insight* **1**.

Pijnenborg, R., Vercruysse, L. and Carter, A. M. (2011). Deep trophoblast invasion and spiral artery
remodelling in the placental bed of the chimpanzee. *Placenta* **32**, 400–408.

Pollheimer, J., Vondra, S., Baltayeva, J., Beristain, A. G. and Knöfler, M. (2018). Regulation of
Placental Extravillous Trophoblasts by the Maternal Uterine Environment. *Front Immunol* **9**, 297.

Qiu, X., Mao, Q., Tang, Y., Wang, L., Chawla, R., Pliner, H. A. and Trapnell, C. (2017). Reversed
graph embedding resolves complex single-cell trajectories. *Nat. Methods* **14**, 979–982.

Rahat, B., Mahajan, A., Bagga, R., Hamid, A. and Kaur, J. (2017). Epigenetic modifications at
DMRs of placental genes are subjected to variations in normal gestation, pathological conditions
and folate supplementation. *Sci Rep* **7**, 40774.

Robins, J. C., Heizer, A., Hardiman, A., Hubert, M. and Handwerger, S. (2007). Oxygen tension
directs the differentiation pathway of human cytotrophoblast cells. *Placenta* **28**, 1141–1146.

Tilburgs, T., Crespo, Â. C., van der Zwan, A., Rybalov, B., Raj, T., Stranger, B., Gardner, L.,
Moffett, A. and Strominger, J. L. (2015). Human HLA-G+ extravillous trophoblasts: Immune-
activating cells that interact with decidual leukocytes. *Proc. Natl. Acad. Sci. U.S.A.* **112**, 7219–
7224.

Trapnell, C., Cacchiarelli, D., Grimsby, J., Pokharel, P., Li, S., Morse, M., Lennon, N. J., Livak,
K. J., Mikkelsen, T. S. and Rinn, J. L. (2014). The dynamics and regulators of cell fate decisions
are revealed by pseudotemporal ordering of single cells. *Nat. Biotechnol.* **32**, 381–386.

Turco, M. Y., Gardner, L., Kay, R. G., Hamilton, R. S., Prater, M., Hollinshead, M. S.,
McWhinnie, A., Esposito, L., Fernando, R., Skelton, H., et al. (2018). Trophoblast organoids as
a model for maternal-fetal interactions during human placentation. *Nature* **213**, 1.

Velicky, P., Knöfler, M. and Pollheimer, J. (2016). Function and control of human invasive
trophoblast subtypes: Intrinsic vs. maternal control. *Cell Adhesion & Migration* **10**, 154–162.

Velicky, P., Meinhardt, G., Plessl, K., Vondra, S., Weiss, T., Haslinger, P., Lendl, T., Aumayr, K.,
Mairhofer, M., Zhu, X., et al. (2018). Genome amplification and cellular senescence are
hallmarks of human placenta development. *PLoS Genet.* **14**, e1007698.

Vento-Tormo, R., Efremova, M., Botting, R. A., Turco, M. Y., Vento-Tormo, M., Meyer, K. B.,
Park, J.-E., Stephenson, E., Polański, K., Goncalves, A., et al. (2018). Single-cell reconstruction
of the early maternal-fetal interface in humans. *Nature* **563**, 347–353.

Wakeland, A. K., Soncin, F., Moretto-Zita, M., Chang, C.-W., Horii, M., Pizzo, D., Nelson, K. K.,
Laurent, L. C. and Parast, M. M. (2017). Hypoxia Directs Human Extravillous Trophoblast
Differentiation in a Hypoxia-Inducible Factor–Dependent Manner. *The American Journal of*
Pathology **187**, 767–780.

Wallace, A. E., Fraser, R. and Cartwright, J. E. (2012). Extravillous trophoblast and decidual
natural killer cells: a remodelling partnership.   *Hum. Reprod. Update* **18**, 458–471.

Wang, F., Flanagan, J., Su, N., Wang, L.-C., Bui, S., Nielson, A., Wu, X., Vo, H.-T., Ma, X.-J. and
Luo, Y. (2012). RNAscope: a novel in situ RNA analysis platform for formalin-fixed, paraffin-
embedded tissues. *J Mol Diagn* **14**, 22–29.

Wiel, C., Augert, A., Vincent, D. F., Gitenay, D., Vindrieux, D., Le Calvé, B., Arfi, V., Lallet-
Daher, H., Reynaud, C., Treilleux, I., et al. (2013). Lysyl oxidase activity regulates oncogenic
stress response and tumorigenesis. *Cell Death Dis* **4**, e855–e855.

Xu, X.-H., Jia, Y., Zhou, X., Xie, D., Huang, X., Jia, L., Zhou, Q., Zheng, Q., Zhou, X., Wang, K.,
et al. (2019). Downregulation of lysyl oxidase and lysyl oxidase-like protein 2 suppressed the
migration and invasion of trophoblasts by activating the TGF-β/collagen pathway in preeclampsia.
Exp. Mol. Med. **51**, 20.

Zhou, Y., Fisher, S. J., Janatpour, M., Genbacev, O., Dejana, E., Wheelock, M. and Damsky, C.
H. (1997). Human cytotrophoblasts adopt a vascular phenotype as they differentiate. A strategy for
successful endovascular invasion? *J. Clin. Invest.* **99**, 2139–2151.

 TITLES AND LEGENDS TO FIGURES

Figure 1. Establishment of a human placental villous explant model. **(A)** Schematic illustration

showing trophoblast subtypes within placental villus, anchoring column, and maternal uterine tissue.

Depicted are villous cytotrophoblasts (CTB), syncytiotrophoblast (SCT), proximal and distal column

trophoblast (PCT & DCT), and invasive interstitial and endovascular EVT. **(B)** Immunofluorescence

and immunohistochemistry images showing Ki67 (red) and HLA-G (green, brown) expression in

human first trimester placental (8 weeks' gestation) and decidual (10 weeks' gestation) tissues. Shown

are villous cytotrophoblasts (CTB), proximal and distal column trophoblasts (PCT & DCT), interstitial

EVT (iEVT), and the mesenchymal core (MC). Bar = 100 μm. **(C)** Images showing gross villous

explant establishment and outgrowth as well as localization of KRT7 (magenta; white), HLA-G

(green), and Ki67 (magenta) to specific subtypes of trophoblasts within explant columns. Nuclei are

shown via DAPI staining (white). Bar = 100 μm.

Figure 2. Exposure to low oxygen promotes trophoblast column outgrowth. **(A)** Schematic illustration

depicting the experimental approach to establishing and culturing placental explants in 1%, 5%, and

20% oxygen. **(B)** Representative images showing how explant column outgrowth area is measured at 0

hr and 48 hr of culture. The direction of explant outgrowth/invasion is also indicated. **(C)**

Representative images of villous explants (n=8 distinct placentae; multiple explants per placenta)

cultured in 1%, 5%, and 20% oxygen. Shown within the inverted (invert) image is the area of column

outgrowth at 48 hr of culture. **(D)** Bar with scatter plots showing the fold-change of column outgrowth

between 48 hr and 0 hr of exposure to 1%, 5%, or 20% oxygen. Median values and standard deviations

are shown. Statistical analyses between groups were performed using ANOVA and two-tailed Dunn's

post-test; significance considered $p < 0.05$.

Figure 3. Comparison of global gene expression patterns between trophoblast columns exposed to 1%,

5%, and 20% oxygen. **(A)** Schematic illustration highlighting the removal of placental villi and

retention of column trophoblasts and invasive EVT for gene expression analysis. **(B)** Volcano plots

showing individual gene-targeting probes differentially expressed between 1% and 20%, 5% and 20%,

and 1% and 5% oxygen cultures. X-axis: coefficients from linear model in log2 scale; y axis: negative

log base 10 of the false discovery rate (FDR). Black circles indicate probes with an FDR > 0.05; blue

and orange circles indicate under-expressed and over-expressed probes with an FDR < 0.05. **(C)** PCA

of 1% (blue), 5% (purple), and 20% (red) oxygen cultured explants. **(D)** Dendrogram depicts

hierarchical clustering of 1%, 5%, and 20% oxygen cultured column trophoblasts using Z-scores from

the top 40 (DEGS from 1% vs 20% comparisons) genes; n=5 per oxygen group. Within the heatmap,

brown = low expression; white = mid-level expression; green = high expression. For each sample,

oxygen condition (1% = blue; 5% = purple; 20% = red) is indicated, as is the unique placental

sample/patient ID used to generate the explant (indicated by shade of grey). Venn diagram showing the

number of shared and unique genes (FDR < 0.05) **(E)** up-regulated and **(F)** down-regulated in 1% and

5% oxygen-cultured explants compared to 20% oxygen cultures.

Figure 4. Placental explant exposure to atmospheric oxygen promotes column trophoblast

proliferation. Gene ontology (GO) pathway analyses of DGE genes indicating the top 10 up-regulated

pathways in explants cultures in **(A)** 1% oxygen and **(B)** 20% oxygen. Adjusted p values (Bonferroni)

and the number of genes identified in each pathway category is represented beside each plot. **(C)**

Dendrogram depicts hierarchical clustering of 1%, 5%, and 20% oxygen cultured column trophoblasts

using a proliferation-specific GO signature (Nuclear Division; 93 genes). Within the heatmap, brown =

low expression; white = mid-level expression; green = high expression. For each sample, oxygen

condition (1% = blue 5% = purple; 20% = red) is indicated, as is the unique placental sample/patient ID

used to generate the explant (indicated by shade of grey). **(D)** Representative immunofluorescence

images of BrdU signal (green) within placental explant columns cultured in 1%, 5%, or 20% oxygen.

Trophoblasts are identified by KRT7 signal (magenta). Bar = 100 μm. **(E)** Bar/scatter plots show

quantification of BrdU incorporation into column trophoblasts cultured in 1%, 5%, or 20% oxygen.

Median values are shown and statistical analyses between groups were performed using ANOVA and

two-tailed Tukey post-test; significance $p < 0.05$.

Figure 5. Low oxygen drives column cell differentiation along the EVT pathway. **(A)** Previously

published scRNAseq data (Vento-Tormo et al., 2018) for 14,366 trophoblast cells from first trimester

placentae (n=5) were selected for investigating the cell-specificity of our list of identified oxygen-

associated gene expression changes. Uniform Manifold Approximation and Projection (UMAP) (Becht

et al., 2018) was used to visualize and cluster the cells after subsetting from other cell types in the

published dataset. Cells are labelled according to their previous characterization (Vento-Tormo et al.,

2018). **(B)** UMAP clusters cells by gene expression of canonical trophoblast marker genes. **(C)**

Heatmap of top 15 genes upregulated in 1% and 20% oxygen. Also shown are genes aligning with

general trophoblasts, villous cytotrophoblasts (CTB), proximal column trophoblast (PCT), and distal

column trophoblast (DCT)/extravillous trophoblast (EVT). Geneset expression patterns are compared

to averaged gene expression levels within single-cell informed cell types (CTB, proliferative CTB,

EVT, proliferative EVT, SCT). **(D)** Pseudotime analysis was applied using Monocle 2 (Qiu et al., 2017;

Trapnell et al., 2014) to visualize gene expression across trophoblast differentiation. Two lineage

trajectories were identified corresponding to the extravillous pathway and villous pathway. A cell state

of origin is also shown. **(E)** The inferred trajectory resulted into two distinct endpoints: one branch

leading to cells highly expressing SCT markers (e.g. ERVFRD-1), and another leading to cells highly

expressing EVT markers (HLA-G). **(F)** A heatmap was constructed using inferred pseudotime and the

top 15 upregulated genes in 1% and in 20% oxygen conditions. Pseudotime was ordered such that the

left and right ends represent the EVT and SCT endpoints. Hierarchical clustering was applied to the

genes (ordered along the rows) and separated into 5 clusters.

Figure 6. Low oxygen exposure promotes column-specific expression of *LOX*. **(A)** Gene array box-plot expression levels of *LOX* mRNA in placental explants cultured in 1%, 5%, and 20% oxygen. **(B)** Representative image showing *LOX* mRNA transcript *in situ* localization (dark pink signal) within a first trimester placental villus. The hashed box outlines the enlarged region to the right. Shown are annotations of trophoblast/placental subtypes; Mesenchymal core (MC), cytotrophoblast (CTB), syncytiotrophoblast (SCT), proximal column trophoblast (PCT), distal column trophoblast (DCT). Shown as an inset is an immunofluorescent image depicting nuclei (DAPI; blue), keratin-7 (K7; magenta), and HLA-G (orange) localization within a serial section of the same placental villus. Bars = 100 μm. **(C)** Representative images of *LOX* mRNA localization within placental explants cultured in 1% or 20% oxygen. Specific trophoblast subtypes are indicated as above, and the hashed box corresponds to the enlarged area shown below. Also shown are insets of immunofluorescent images depicting nuclei (DAPI; blue), keratin-7 (K7; magenta), and HLA-G (orange) localization within corresponding serial sectioned regions of the explant. Bars = 100 μm. **(D)** Bar/scatter plots show LOX activity levels within conditioned media (CM) of placental explants cultured in either 1% or 20% oxygen in the presence/absence of the LOX inhibitor BAPN (5 mM). Recombinant active lysyl oxidase (rLOX) served as a positive control whereas explant culture media alone served as a negative control. Activity corresponds to level of fluorescence intensity (590 nm). Median values are shown and statistical analyses between groups were performed using ANOVA and two-tailed Tukey post-test; significance $p < 0.05$.

Figure 7. Inhibition of lysyl oxidase (LOX) dampens column outgrowth. **(A)** Bar/scatter plots show

LOX activity levels within conditioned media (CM) of placental explants cultured in 1% oxygen in the

presence/absence of BAPN (500 μM). Recombinant active lysyl oxidase (rLOX) served as a positive

with explant culture media alone served as a negative control. Activity corresponds to level of

fluorescence intensity (590 nm). Median values are shown and statistical analyses between groups were

performed using ANOVA and two-tailed Tukey post-test. **(B)** Representative images of villous

explants (n=3 distinct placentae; multiple explants per placenta) cultured in 1% oxygen and in the

presence/absence of BAPN (500 μM). Column outgrowth is shown at 0hr and 48 hr of culture. **(C)** Box

plots showing the fold-change of column outgrowth between 48 hr and 0 hr following treatment with

BAPN. Median values are shown and statistical analyses between groups were performed using

ANOVA and two-tailed Tukey post-test; significance *p* <0.05.

Supplemental Figure Legends

Supplemental Figure 1. Normalization and PCA analysis of placental explant gene microarray data.

Average expression values for each explant sample **(A)** before and **(B)** after quantile normalization and

filtering/removal of probes. **(C and D)** Principal component analysis (PCA) of raw data, normalized

data, and filtered data.

Supplemental Figure 2. Pathway analyses of unique gene signatures of 1% and 5% oxygen-cultured

explants. **(A)** Top 20 gene pathways upregulated exclusively in 1% oxygen when compared to 20%

oxygen. **(B)** Top 20 gene pathways exclusively upregulated in 5% oxygen when compared to 20%

oxygen. The size of the dot represents the number of significant genes in the pathway and the colour

represents adjusted *P*-values as indicated in the legend.

Supplemental Figure 3. Top gene pathways identified in explant columns cultured in 5% oxygen. Top

20 pathways upregulated in 5% oxygen when compared to 20% oxygen. The size of the dot represents

the number of significant genes in the pathway and the colour represents adjusted *P*-values as indicated

in the legend.

Supplemental Figure 4. Low oxygen culture promotes trophoblastic and EVT gene signatures in

placental explants. **(A)** Dendrogram depicts hierarchical clustering of 1%, 5%, and 20% oxygen

cultured column trophoblasts using the respective 14-gene signature. The heatmap displays expression

profiles of trophoblast genes identified as being significantly different between 1%, 5%, and 20%

cultures. Within the heatmap, brown = low expression; white = mid-level expression; green = high

expression. For each sample, oxygen condition (1% = blue 5% = purple; 20% = red) is indicated, as is

the unique placental sample/patient ID used to generate the explant (indicated by shade of grey).

Representative immunofluorescence images showing expression of PEG10 (magenta) within **(B)** first

trimester placental villi and **(C)** Matrigel-imbedded placental explants cultured in either 1%, 5%, or

20% oxygen. Trophoblast are identified by keratin-7 staining (KRT7; green) and nuclei are stained

with DAPI (white). Shown are specific cell types: Syncytiotrophoblast (SCT), villous cytotrophoblasts

(CTB), proximal column trophoblast (PCT), distal column trophoblast (DCT), and the mesenchymal

core (MC). Bar = 100 μm**.**

Figure 1

A

PCT
Anchoring trophoblast column
DCT
Interstitial EVT
Endovascular EVT
Spiral artery
CTB
SCT
Uterine epithelium
Uterine stroma

B

DAPI Ki67 HLA-G
DCT
PCT
MC
CTB
8wk placenta

HLA-G
iEVT
DCT
PCT
MC
Anchoring column
10wk decidua

C

Villous Explant
Explant outgrowth
Placental villus

DAPI KRT7 HLA-G
iEVT
DCT
MC
PCT

Ki67 KRT7
iEVT
DCT
MC
PCT

Figure 2

A First trimester placenta → Established for 24 hr → 5%

Explants cultured in 1%, 5% or 20% oxygen for 48 hr. → Measure column outgrowth.

5% 1% 20%

B
Placental villi
Area
0 hr

Column
Outgrowth
48 hr

C

Oxygen Condition	0 hr	48 hr	48 hr (invert)
1%			
5%			
20%			

D
Change in column area (48 hr/0 hr)

p < 0.0001
p = 0.12
p = 0.0003

Oxygen Condition
1%   5%   20%

Figure 3

Figure 4

A GO Pathways enriched in 1% O$_2$

B GO Pathways enriched in 20% O$_2$

C

Proliferation gene signature

D

E Column cell proliferation

Figure 5

Figure 6

A

LOX Expression

Oxygen Condition

B LOX

CTB
MC   SCT
DAPI K7 HLA-G
PCT
PCT
DCT
DCT

LOX

C LOX   DAPI K7 HLA-G   LOX   DAPI K7 HLA-G

MC   MC   PCT   MC   MC
PCT   DCT   PCT   PCT   DCT
DCT

1% Oxygen

PCT
DCT

20% Oxygen

MC
PCT

DCT

D

Fluorescence (590 nm)

80000
60000
40000
20000

P< 0.0001
P= 0.0002
P= 0.002

BAPN
5 mM   -   +   -   +   -   +   -   +
         rLOX   (-)   1% O₂ CM 20% O₂ CM
         4 μg/mL

Figure 7

A

Fluorescence (590 nm)

$P< 0.0001$
$P= 0.001$
$P= 0.04$
$P= 0.03$

rLOX (-) (-) BAPN 500µM

1% O$_2$ Explant CM

B

0h   48h

Control

BAPN

1% O$_2$

C

Outgrowth Area (1% O$_2$)

$P= 0.0003$

Change in column area (48 hr/0 hr)

Control (n=3)   BAPN (n=3)