Metabolic pathway rerouting in *Paraburkholderia rhizoxinica* evolved long-overlooked derivatives

of coenzyme F$_{420}$.

Daniel Braga[a,d], Daniel Last[a], Mahmudul Hasan[a,d], Huijuan Guo[b], Daniel Leichnitz[b], Zerrin Uzum[c],

Ingrid Richter[c], Felix Schalk[b], Christine Beemelmanns[b], Christian Hertweck[c,d], and Gerald Lackner[a,d] *

Author affiliations

[a]Junior Research Group Synthetic Microbiology
Leibniz Institute for Natural Product Research and Infection Biology – Hans Knöll Institute
Beutenbergstr 11a, 07745 Jena, Germany
10
11   [b]Junior Research Group Chemical Biology of Microbe-Host Interactions
12   Leibniz Institute for Natural Product Research and Infection Biology – Hans Knöll Institute
13   Beutenbergstr 11a, 07745 Jena, Germany
14
15   [c]Department of Biomolecular Chemistry
16   Leibniz Institute for Natural Product Research and Infection Biology – Hans Knöll Institute
17   Beutenbergstr 11a, 07745 Jena, Germany
18
19   [d]Friedrich Schiller University, Jena
20
21   *Corresponding author:
22   Dr. Gerald Lackner
23   E-Mail: gerald.lackner@leibniz-hki.de
24   Phone: +49 3631 532 1104
25

Author contributions and conflict of interest

Daniel Braga performed research, analyzed data (molecular biology, malachite green assay, mass spectrometry)

and contributed to writing the manuscript, Daniel Last performed research and analyzed data (structure

elucidation, Fno assay, biogas plant studies), Mahmudul Hasan performed research (CofC/D enzyme assays),

Huijuan Guo performed research and analyzed data (structure elucidation), Daniel Leichnitz performed research

(chemical synthesis), Zerrin Uzum performed research (microscopy), Ingrid Richter performed research

(microscopy), Felix Schalk performed research (*cofE* constructs), Christine Beemelmanns designed research,

acquired funding, analyzed data (structure elucidation, synthesis) and edited the manuscript, Christian Hertweck

designed research, acquired funding and edited the manuscript, Gerald Lackner designed the study, acquired

funding and wrote the original manuscript. The authors declare no conflict of interest.

Keywords: Cofactors, natural products, biocatalysis, symbiosis, deazaflavin

1 Abstract

Coenzyme F_{420} is a specialized redox cofactor with a highly negative redox potential. It supports
biochemical processes like methanogenesis, degradation of xenobiotics or the biosynthesis of
antibiotics. Although well-studied in methanogenic archaea and actinobacteria, not much is known
about F_{420} in Gram-negative bacteria. Genome sequencing revealed F_{420} biosynthetic genes in the
Gram-negative, endofungal bacterium *Paraburkholderia rhizoxinica,* a symbiont of phytopathogenic
fungi*.* Fluorescence microscopy, high-resolution LC-MS, and structure elucidation by NMR
demonstrated that the encoded pathway is active and yields unexpected derivatives of F_{420} (3PG-F_{420}).
Further analyses of a biogas-producing microbial community showed that these derivatives are more
widespread in nature. Genetic and biochemical studies of their biosynthesis established that a
specificity switch in the guanylyltransferase CofC re-programmed the pathway to start from 3-
phospho-D-glycerate, suggesting a rerouting event during the evolution of F_{420} biosynthesis.
Furthermore, the cofactor activity of 3PG-F_{420} was validated, thus opening up perspectives for its use
in biocatalysis. The 3PG-F_{420} biosynthetic gene cluster is fully functional in *Escherichia coli*, enabling
convenient production of the cofactor by fermentation.

16 Introduction

Cofactors are essential for the catalytic power of many enzymes and thus play a key role in virtually all
metabolic pathways. Knowledge of their catalytic functions and biosynthesis is highly important for
the understanding of biochemical reactions as well as their application in biocatalysis and
biotechnology. An important subclass comprises redox-cofactors that mediate electron transfer
between molecules. The deazaflavin coenzyme F_{420} (Figure 1) is a specialized redox cofactor with a
lower redox potential (–350 mV) than NAD (1). This feature makes F_{420} an ideal electron carrier
between H_2 and NAD(P) in methanogenesis and renders it a strong reducing agent for challenging
reactions in biocatalysis (2). For instance, enzymatic processes involving F_{420} facilitate the degradation
of pollutants like aromatic nitro compounds (3) or the carcinogen aflatoxin (4). Furthermore, F_{420}-
dependent enzymes are important for asymmetric ene reductions (5, 6). In actinomycetes, coenzyme
F_{420} is involved in the biosynthesis of antibiotics like oxytetracycline (7), pyrrolobenzodiazepines (8), or
thiopeptins (9). Additionally, F_{420} has attracted considerable interest as a fitness factor of the human
pathogen *Mycobacterium tuberculosis*, being involved in nitrosative stress response (10) or prodrug-
activation (11).

Figure 1. Biosynthesis of coenzyme F_{420} A) F_O synthase FbiC (in archaea: CofG/H) catalyzes formation of the deazaflavin ring from tyrosine and 5-amino-6-(ribitylamino)-uracil, an intermediate of riboflavin biosynthesis. B) Biosynthetic scheme of F_{420}-n proposed for archaea. CofC and CofD catalyze the activation of 2-PL and transfer of the 2-PL moiety, respectively. CofE performs (oligo-)γ-glutamylation. The number of glutamate residues (n) varies depending on the organism. The origin of 2-PL is unknown. C) Biosynthesis of F_{420}-n proposed for mycobacteria: CofC and CofD activate PEP resulting in DF_{420} formation. The C-terminal domain of FbiB reduces DF_{420} to F_{420}. EPPG: enolpyruvyl-diphosphoguanosine, LPPG: lactyl-diphosphoguanosine, 2-PL: 2-phospho-L-lactate.

A key step during the biosynthesis of F_{420} is the formation of the deazaflavin fluorophore F_O (Figure 1A), a stable metabolic precursor of F_{420} originating from tyrosine and an intermediate of riboflavin biosynthesis. This chemically challenging step is catalyzed by the radical SAM enzyme complex CofG/H in archaea or the homologous dual-domain protein FbiC in actinobacteria (12). F_O is then further processed by CofC (EC 2.7.7.68) and CofD (EC 2.7.8.28). This pair of enzymes is responsible for the biosynthesis of the 2-phospho-L-lactate (2-PL) moiety (Figure 1B). Previous studies have shown that CofC from *Methanocaldococcus jannaschii* directly activates 2-phospho-L-lactate (2-PL) by guanylation (13) resulting in the formation of the short-lived metabolite lactyl-diphosphoguanosine (LPPG). CofD then forms F_{420}-0 by transfer of the 2-PL moiety from LPPG to F_O (14). The enzymes producing 2-PL, however, have remained elusive in all F_{420} producers so far. Recently, Bashiri *et al.* proposed a revised biosynthetic pathway by demonstrating that phosphoenolpyruvate (PEP) instead of 2-PL can serve as a substrate of CofC in *Mycobacteria* (15). The resulting dehydro-F_{420} (DF_{420}) is reduced to F_{420} by a flavin-dependent reductase domain present in the FbiB protein (Figure 1C). In mycobacteria, FbiB is a dual-

domain protein consisting of a γ-glutamyl ligase domain (CofE-like, EC 6.3.2.31) and the C-terminal

DF_{420} reductase domain (16). The γ-glutamyl ligase CofE finally decorates F_{420}-0 with a varying number

of (oligo-)γ-glutamate residues (17).

F_{420} is not ubiquitous in prokaryotes, but it is associated with certain phyla (18, 19). First discovered in

methanogenic archaea (20, 21), it was extensively studied as a potential drug target of pathogenic

mycobacteria (10) or as a cofactor enabling antibiotics biosynthesis in streptomycetes (2). Genome

sequencing revealed that some Gram-negative bacteria have acquired F_{420} genes by horizontal transfer

(19, 22). However, virtually nothing is known about the biosynthesis and role of F_{420} in these organisms.

By genome mining, we found a biosynthetic gene cluster (BGC) homologous to those previously

implicated in the biosynthesis of F_{420} in the endofungal bacterium *Paraburkholderia rhizoxinica* HKI 454

(Figure 2A and *Supporting Information* Table S3). This organism is an intracellular endosymbiont of the

phytopathogenic fungus *Rhizopus microsporus* supplying its host with antimitotic toxins that act as

virulence factors during infection of rice plants (23-25). We hypothesized that genes from Gram-

negative bacteria related to F_{420} biosynthesis could facilitate F_{420} production in *E. coli* or could reveal

novel biosynthetic routes towards this valuable molecule. Therefore, we set out to investigate if the

BGC is active and if it can be refactored to produce F_{420} in *E. coli*.

Here, we show that *P. rhizoxinica* produces unexpected F_{420}-derivatives (3PG-F_{420}) both in symbiosis as

well as in axenic culture. Heterologous expression and large-scale production in *E. coli* allowed for

elucidation of their chemical structure. By comparative analyses, we discovered related metabolites in

a biogas-producing microbial community, thus indicating their broader abundance and relevance.

Enzyme assays showed that a switch in substrate specificity of CofC is responsible for the biosynthesis

of 3PG-F_{420} and proved that it can serve as a substitute for F_{420} in biochemical reactions.

Results and Discussion

Figure 2. Deazaflavin biosynthesis in *P. rhizoxinica.* A) BGC of 3PG-F$_{420}$. Core genes are shown in dark grey. B-I) Microscopy photographs depict fluorescence characteristic of deazaflavins in blue. B-D) axenic *M. smegmatis* (B), *P. rhizoxinica* (C), and *E. coli* / pDB045 (D). In *R. microsporus* ATCC 62417, deazaflavins are correlated to the presence of *P. rhizoxinica* symbionts (green, Syto9 staining) (E). No fluorescence was detected in cured ATCC 62417 mycelium (F) or in the naturally symbiont-free strain, CBS 344.29 (G). The same pattern was observed in spores of either wild-type ATCC 62417 (H) or CBS 344.29 (I). Scale bars represent 10 μm. J) Refactored versions of the BGC and corresponding plasmids for heterologous expression in *E. coli.* Asterisks mark genes from *M. jannaschii.*

To test whether *P. rhizoxinica* is capable of producing deazaflavins, we investigated axenic cultures of symbiont (*P. rhizoxinica*) and host (*R. microsporus*) as well as symbiotic cultures by fluorescence microscopy (Figures 2B-I and *Supporting Information* Figures S1-4). Indeed, symbiotic *P. rhizoxinica* and the cytosol of colonized mycelia emitted strong fluorescence characteristic of deazaflavins. Notably, even fluorescence of bacteria present inside of fungal spores was observed. Axenic fungi, however, showed no fluorescence, whereas only low signals were measured from axenic bacteria under the same conditions. To corroborate the results obtained by microscopy, we extracted metabolites from axenic *P. rhizoxinica*, axenic *R. microsporus* as well as fungal host containing

endosymbionts and analyzed the extracts by LC-MS/MS. To our surprise, only F_O could be detected in

axenic *P. rhizoxinica* and in the fungal host containing endosymbionts, but none of the expected F_{420}-

n species.

To further test the biosynthetic capacity of the full BGC, we refactored it to obtain a single operon

under the control of a T7 promoter for heterologous expression yielding *E. coli/*pDB045 (Figure 2J).

Examination of transformed bacteria by fluorescence microscopy revealed strong fluorescence as in

the endofungal bacteria (Figure 2D), but LC-MS analyses again yielded only mass traces of F_O (mainly

found in the culture supernatant), but not of F_{420}-n. Since the impeded production could be attributed

to nonfunctional proteins, we analyzed proteins by SDS-PAGE and found that CofD was poorly soluble

in *E. coli*. Replacement of the *cofD* gene with the corresponding *M. jannaschii* homolog (14) provided

soluble protein (*E. coli/*pDB060)*,* however, again no trace of F_{420} could be detected. Therefore, we

reexamined the metabolome of *E. coli/*pDB045 for characteristic MS/MS fragments derived from the

F_O moiety (*m/z* 230.06, 346.10, 364.11). Surprisingly, the analysis revealed spectra with a similar

fragmentation pattern, yet derived from precursor ions that had a mass shift of 15.995 compared to

F_{420}, indicating the presence of an additional oxygen atom (Figures 3A-B and Supporting Information

Figures 5-8). Further analyses revealed an (oligo-)γ-glutamate series of the oxygenated compound

suggesting these species are congeners. According to MS/MS fragmentation, the additional oxygen

was present in the "phospholactyl" moiety of F_{420}-n thus forming a "phosphoglyceryl" moiety.

Extensive 1D- and 2D-NMR experiments (Figure 3C and Supporting Information *Section 2.2*) and

comparison to classical F_{420} (20) corroborated that this moiety corresponds to 3-phosphoglycerate (3-

PG). Therefore, we named the molecules 3PG-F_{420}. This finding was unexpected because 2-

phosphoglycerate is structurally more similar to PEP and 2-PL than 3-PG. Chemical degradation

followed by chiral UHPLC-MS finally substantiated that the additional stereocenter of 3PG-F_{420} is *R*-

configured (Figure 3D and Supporting Information Figure S51). Large-scale cultivation of *E. coli/*pDB045

also revealed traces of dehydro-F_{420} (DF_{420}), but the yields were too low for NMR studies. The structure

and occurrence of 3PG-F_{420} has not been reported before. To date, the only known derivatives of F_{420}-

n are factor F_{390}-A and F_{390}-G, 8-OH-AMP and 8-OH-GMP esters of F_{420}, respectively (26). In

methanogens, they are formed reversibly, e.g., during oxygen exposure, acting as a reporter

compound for hydrogen starvation (27). In contrast, the modifications seen in 3PG-F_{420} are not

temporary. Rather, 3PG-F_{420} seems to replace F_{420} as a natural deazaflavin-cofactor in *P. rhizoxinica*. At

least in this organism, it does not coexist with classical F_{420}. This situation is reminiscent of mycothiol,

a specialized thiol cofactor that replaced glutathione in actinobacteria (28).

Figure 3. Chemical analysis of 3PG-F_{420}. A) Extracted ion chromatograms of 3PG-F_{420}-2 produced in *E. coli*. I: *E. coli* / pDB045,

II: *cofD* exchanged by *M. jannaschii* homolog (pDB060), III: *cofD* and *cofC* exchanged by *M. jannaschii* homologs (pDB070) IV:

empty vector (pETDuet). B) Excerpt of the MS/MS spectrum of 3PG-F_{420}-2. Grey bars highlight m/z used for fragment ion

search of F_{420} derivatives. C) ^1H NMR comparison of F_{420}-n (D$_2$O), 3PG-F_{420}-n (0.1% ND$_3$ in D$_2$O) and 3PG-F_{420}-0 (0.1% ND$_3$ in

D$_2$O) indicated the replacement of the lactyl moiety in F_{420} with a glyceryl moiety in 3PG-F_{420}. D) Proposed structures of 3PG-

F_{420}-0, 3PG-F_{420}-n, and DF$_{420}$-n.

To investigate if 3PG-F_{420} is produced by wild-type *P. rhizoxinica*, we reanalyzed LC-MS data for the

presence of corresponding mass signals. Indeed, 3PG-F_{420} species were found in samples containing

bacteria (axenic culture, symbiosis), but not in symbiont-free host mycelia (Supporting Information

Figures S9-14). In extracts of *P. rhizoxinica*, no traces of F_{420} and DF$_{420}$ were detected. The presence of

3PG-F_{420} was restricted to the cell pellet, whereas F_O was abundant in culture supernatants. We thus

conclude that the fluorescence observed in bacterial cells of *P. rhizoxinica* is derived from 3PG-F_{420} and

F_O. So far, there is only preliminary evidence for the occurrence of deazaflavin cofactors in a few Gram-negative bacteria, e.g., *Oligotropha carboxidivorans* and *Paracoccus dentrificans* (19) as well as in the uncultured, but biosynthetically highly prolific '*Candidatus* Entotheonella factor' (22). The exact structure and function of F_{420} in most Gram-negative bacteria that harbor corresponding biosynthetic genes, however, is unknown. The fact that *P. rhizoxinica* produced 3PG-F_{420} under symbiotic cultivation conditions allows for the conclusion that it provides a fitness benefit in its natural habitat. Notably, none of the well-characterized F_{420}-dependent enzyme families (4, 18) are encoded in the *P. rhizoxinica* genome according to BLAST and conserved domains searches, not even any of the widespread regeneration systems like Fno or F_{420}-dependent glucose-6-phosphate dehydrogenase. Therefore, future investigations of 3PG-F_{420}-producing organisms are likely to reveal novel enzymes, regeneration systems, and cellular pathways depending on this cofactor.

To assess if 3PG-F_{420} is restricted to fungal endosymbionts or if it might be more widespread in the environment, we examined *M. jannaschii* and *M. smegmatis* for the presence of any F_{420} congeners. Only F_{420}-n was found from extracts of *M. jannaschii,* while F_{420}-n and DF_{420}-n were detected in extracts of *M. smegmatis*. (Supporting Information Figures S15-16). None of the 3PG-F_{420}-derivatives were found in the reference organisms. Since methanogens are a common source of F_{420} in nature, we analyzed (two independent) sludge samples from a local biogas production plant. To our surprise, extraction of the microbial community present in the biogas-producing sludge followed by LC-MS/MS eluted, besides classical F_{420}, a compound with identical retention time, exact mass and MS/MS fragmentation pattern as 3PG-F_{420} (Supporting Information Figure S17). As fluorescence and UV-based detection does usually not resolve classical F_{420} and 3PG-F_{420}, these derivatives might have been misidentified as F_{420} in the past. Since neither *P. rhizoxinica* nor its host *R. microsporus* are able to grow under anaerobic and thermophilic (temperatures >42 °C) conditions, they can be excluded as the source of these cofactors. The high complexity of biogas-producing microbiomes (29) do not allow for an educated guess of the producer, although methanogens would be reasonable candidates.

In order to rationalize how the biosynthetic pathway was redirected to form 3PG-F_{420} instead of F_{420}, we examined key steps of the biosynthesis more closely. We observed that production of 3PG-F_{420} was not abolished by the exchange of c*ofD* from *P. rhizoxinica* by *cofD* from *M. jannaschii* (plasmid pDB060). Hence, the phospholactyl transferase CofD could not be held accountable for the switch towards 3PG-F_{420}. According to the existing biosynthetic model, the most plausible scenario was that CofC incorporated 3-phospho-D-glycerate (3-PG), an intermediate of glycolysis, instead of 2-PL to form 3PG-F_{420}-0 and, to a minor extent, PEP to form DF_{420}-0. To test this hypothesis, we exchanged *cofC* and *cofD* by the corresponding *M. jannaschii* homologs. The resulting strain *E. coli*/pDB070 (Figure 2J and Supporting Information Figure S18) produced neither 3PG-F_{420} nor F_{420} but traces of DF_{420}. To further investigate the substrate specificity of CofC, we performed an in-vitro assay using CofC and CofD (13).

Genes *cofC* of *P. rhizoxinica* as well as *cofC* and *cofD* from *M. jannaschii* were cloned, corresponding

proteins produced as hexahistidine fusions in *E. coli* and purified by metal affinity chromatography for

in-vitro assays. The physiologically relevant isomers 2-phospho-D-glycerate (2-PG) and 3-phospho-D-

glycerate (3-PG), as well as PEP and 2-PL, served as substrates. Reaction products were monitored by

LC-MS. Indeed, when CofC from *P. rhizoxinica* was tested, the mass of $3PG-F_{420}$-0 appeared after

reaction with D-3-PG eluting at the same retention time as the in-vivo product (Supporting Information

Section 2.4). In addition, the formation of DF_{420}-0 and F_{420}-0 was detected, when the enzymes were

incubated with PEP and 2-PL, respectively. In contrast, reaction with 2-PG yielded mass signals close

to the noise level. Controls lacking CofC did not generate any of these products. In a direct substrate

competition assay (Figure 4A), 3-PG was found to be the preferred substrate with a relative turnover

of ca. 73% (2-PL: 23% PEP: 4%). This finding is in agreement with the structure of $3PG-F_{420}$ and the

occurrence of DF_{420} as a minor biosynthetic product in *E. coli*. Note that CofC from *M. jannaschii*

displayed a strong turnover of 2-PL (96.5%), weak turnover of PEP (3.5%) and no turnover of 3-PG. This

finding supports the notion that the CofC of *P. rhizoxinica* has undergone a substrate specificity switch

during evolution.

A

B

Figure 4. Combined CofC/D in-vitro assay. A) Relative turnover of substrates estimated from a substrate competition assays

(D-3-PG, 2-PL, and PEP). CofC from *P. rhizoxinica* accepted 3-PG (72.7%), 2-PL (23.4%), and PEP (3.9%). CofC from *M. jannaschii*

preferred 2-PL (96.5%), and PEP (3.5%). 3-PG was not turned over. CofD from *M. jannaschii* was used in all assays. Error bars

represent the standard deviation (SD) of three independent biological replicates (N=3). B) Proposed model of $3PG-F_{420}$

biosynthesis. 3-GPPG: 3-(guanosine-5'-disphospho)-D-glycerate.

Recently, Bashiri *et al.* claimed that PEP is the substrate of CofC in prokaryotes (15). Our results confirm

the hypothesis that PEP is the physiological substrate in mycobacteria, since we observed turnover of

PEP by all CofC homologs tested. However, in contrast to Bashiri *et al.*, 2-PL was the best substrate of

M. jannaschii CofC in our assay. Since Graupner and White detected significant amounts of 2-PL in

methanogenic archaea and observed the conversion of lactate into 2-PL by isotope labeling (30), we

conclude that 2-PL might still be a relevant substrate in archaea. From a phylogenetic perspective, our

results suggest that multiple metabolic re-wiring events occurred in the evolution of F_{420} biosynthesis.

While actinobacteria evolved the DF_{420} reductase (C-terminal domain of FbiB), archaea accomplished

to produce the (unusual) metabolite 2-PL. Other organisms, as exemplified by *P. rhizoxinca,* rerouted

the biosynthesis to the ubiquitous metabolite 3-PG.

To address the question if the γ-glutamyl ligase CofE adapted its substrate specificity to $3PG-F_{420}$, we

individually co-expressed *cofE* genes from *P. rhizoxinica*, *M. jannaschii* and *M. smegmatis* (*fbiB*)

together with a minimal BGC consisting of *fbiC*, *cofC,* and *cofD* in a two-plasmid system. Extraction of

metabolites and LC-MS/MS revealed that all three CofE homologs elongated $3PG-F_{420}$-0 to oligo-

glutamate chain lengths up to n=6 (Supporting Information Figures S63-65). Thus, we conclude that

CofE does not act as an additional specificity filter during chain elongation of $3PG-F_{420}$.

Figure 5. Cofactor function of $3PG-F_{420}$. A) Michaelis-Menten kinetics of Fno for F_{420} (left) and $3PG-F_{420}$ as substrates (right).

Three biological replicates were used to determine parameters. K_M for F_{420} was 3.6 ± 0.7 µM (standard error). K_M for $3PG-F_{420}$

was 5.1 ± 1.0 µM. Error bars indicate standard deviation of replicates (N=3). B) In-vivo reduction of malachite green

(absorbance: 618 nm) by the F_{420}-dependent reductase MSMEG_5998. Fno was used to regenerate $3PG-F_{420}H_2$. Left panel:

Time course of the malachite green depletion assay. Right panel: Bar chart of residual malachite green after 20 h: wt: *E. coli* BL21(DE3), pDB061: *E. coli* producing MSMEG_5998, pDB071: *E. coli* producing 3PG-F_{420}-0 + Fno. Exact means ± SD of biological triplicates were 0.234 ± 0.017 (wt), 0.124 ± 0.003 (pDB061), 0.169 ± 0.011 (pDB071), and 0.082 ± 0.0139 (pDB061/pDB071). An asterisk indicates statistical significance (one-way ANOVA, p<0.05, N=3). C) Engineered *E. coli* combining 3PG-F_{420}, Fno and reductase MSMEG_5998 (*red*) for reduction of malachite green.

The successful isolation of 3PG-F_{420} and reconstitution of its biosynthesis in *E. coli* motivated us to address the question of whether 3PG-F_{420} could substitute F_{420} in biocatalysis. To this end, we cloned a gene encoding Fno (F_{420}:NADPH oxidoreductase), an enzyme that serves as a regeneration system for $F_{420}H_2$ using NADPH/H^+ as an electron donor (31). We first examined if Fno can accept 3PG-F_{420} as a substrate. Indeed, we observed an efficient reduction of 3PG-F_{420} by recombinant Fno as mirrored by a rapid decrease of characteristic UV absorption. An examination of kinetic parameters (Figure 5A) revealed that the apparent K_M of Fno for F_{420} was 3.6 ± 0.7 µM. This value is similar to the reported K_M of 10 µM (32). Under identical assay conditions, the K_M for 3PG-F_{420} was only slightly higher (5.1 ± 1.0 µM). The v_{max} values were in a similar range as well (F_{420}: 1.3 ± 0.2 µM min^{-1}; 3PG-F_{420}: 0.88 ± 0.07 µM min^{-1}) pointing towards only a minor reduction of maximal turnover. Encouraged by the finding that 3PG-F_{420} can substitute F_{420}, we aimed at an in-vivo application of the cofactor for malachite green reduction as a proof of principle. To this end, we combined the *fno* gene with a minimal BGC producing 3PG-F_{420}-0 (Figure 5C and Supporting Information Figure S62) on a single vector (pDB071). Additionally, the F_{420}-dependent malachite green reductase gene MSMEG_5998 (33) from *M. smegmatis* was cloned and expressed from a compatible vector backbone (pDB061). Finally, co-expression of all components in *E. coli* yielded a strain (pDB061/pDB071) that was able to decolorize malachite green significantly faster than control strains expressing the reductase or the cofactor alone (Figure 5B). Thus, we conclude that 3PG-F_{420} can substitute F_{420} as a redox cofactor in this case. The production of classical F_{420} and its use for biotransformations in *E. coli* has just recently been achieved in moderate yields using *Mycobacterium* genes including the DF_{420} reductase domain (15).

In summary, we discovered a derivative of the redox cofactor F_{420} that is produced by the Gram-negative endofungal bacterium *P. rhizoxinica*. We fully elucidated its chemical structure and show its potential cofactor function. Thus, our work is a solid basis to unveil unknown enzyme families and bioprocesses depending on 3PG-F_{420}. Intriguingly, its presence in a biogas-producing digester suggests that the cofactor is more widespread in nature than expected. Furthermore, we could demonstrate that the guanylyltransferase CofC is responsible for the biosynthetic switch leading to the production of 3PG-F_{420}. Our results thus significantly refine and extend the biosynthetic pathway models to deazaflavin cofactors in several phyla. Notably, the pathway discovered here, offered an alternative route to heterologous production and reconstitution of F_{420}-dependent bioprocesses in *E. coli*. In recent years, there has been increasing interest in F_{420}-dependent enzymes for biocatalysis (5, 6, 34,

1 35). Future applications will comprise for instance enantioselective biotransformations or the creation

2 of a universal expression host for the production of antibiotics and other high-value compounds.

Methods

5 Materials and methods are summarized in Supporting Information (Section 1).

Acknowledgments

We thank Prof. Dr. Dina Grohmann and Dr. Ghader Bashiri for kindly providing *M. jannaschii* and

M. smegmatis / fbiABC, respectively. We thank Biogas Jena GmbH and Co. KG for the kind donation of

biogas plant samples. We thank Heike Heinecke for conducting NMR experiments. G.L. thanks the

Deutsche Forschungsgemeinschaft (DFG Grant LA 4424/1-1) and the Carl Zeiss Foundation for funding.

Financial support by the DFG (CRC 1127 ChemBioSys) to C.H. and C.B., and, BE-4799/2-1 to C.B., and

Leibniz Award to C.H., by the ERC (MSCA-IF-EF-RI Project 794343, to I.R.) and the JSMC to Z.U. is

gratefully acknowledged.

References

1.   Jacobson F & Walsh C (1984) Properties of 7,8-didemethyl-8-hydroxy-5-deazaflavins relevant
to redox coenzyme function in methanogen metabolism. *Biochemistry* 23(5):979-988.
2.   Greening C*, et al.* (2016) Physiology, biochemistry, and applications of F_{420}- and F_o-dependent
redox reactions. *Microbiol Mol Biol Rev* 80(2):451-493.
3.   Heiss G*, et al.* (2002) npd gene functions of *Rhodococcus* (*opacus*) *erythropolis* HL PM-1 in
the initial steps of 2,4,6-trinitrophenol degradation. *Microbiology* 148(Pt 3):799-806.
4.   Taylor MC*, et al.* (2010) Identification and characterization of two families of $F_{420}H_2$-
dependent reductases from Mycobacteria that catalyse aflatoxin degradation. *Mol Microbiol*
78(3):561-575.
5.   Taylor M, Scott C, & Grogan G (2013) F_{420}-dependent enzymes - potential for applications in
biotechnology. *Trends Biotechnol* 31(2):63-64.
6.   Mathew S, Trajkovic M, Kumar H, Nguyen QT, & Fraaije MW (2018) Enantio- and
regioselective ene-reductions using $F_{420}H_2$-dependent enzymes. *Chem Commun*
54(79):11208-11211.
7.   Wang P, Bashiri G, Gao X, Sawaya MR, & Tang Y (2013) Uncovering the enzymes that catalyze
the final steps in oxytetracycline biosynthesis. *J Am Chem Soc* 135(19):7138-7141.
8.   Li W, Khullar A, Chou S, Sacramo A, & Gerratana B (2009) Biosynthesis of sibiromycin, a
potent antitumor antibiotic. *Appl Environ Microbiol* 75(9):2869-2878.
9.   Ichikawa H, Bashiri G, & Kelly WL (2018) Biosynthesis of the thiopeptins and identification of
an $F_{420}H_2$-dependent dehydropiperidine reductase. *J Am Chem Soc* 140(34):10749-10756.
10.   Purwantini E & Mukhopadhyay B (2009) Conversion of NO_2 to NO by reduced coenzyme F420
protects mycobacteria from nitrosative damage. *Proc Natl Acad Sci U S A* 106(15):6333-6338.
11.   Singh R*, et al.* (2008) PA-824 kills nonreplicating *Mycobacterium tuberculosis* by intracellular
NO release. *Science* 322(5906):1392-1395.
12.   Decamps L*, et al.* (2012) Biosynthesis of F_0, precursor of the F_{420} cofactor, requires a unique
two radical-SAM domain enzyme and tyrosine as substrate. *J Am Chem Soc* 134(44):18173-
18176.

13. Grochowski LL, Xu H, & White RH (2008) Identification and characterization of the 2-phospho-L-lactate guanylyltransferase involved in coenzyme F_{420} biosynthesis. *Biochemistry* 47(9):3033-3037.

14. Graupner M, Xu H, & White RH (2002) Characterization of the 2-phospho-L-lactate transferase enzyme involved in coenzyme F(420) biosynthesis in *Methanococcus jannaschii*. *Biochemistry* 41(11):3754-3761.

15. Bashiri G*, et al.* (2019) A revised biosynthetic pathway for the cofactor F_{420} in prokaryotes. *Nat Commun* 10(1):1558.

16. Bashiri G*, et al.* (2016) Elongation of the poly-gamma-glutamate tail of F_{420} requires both domains of the F420:gamma-glutamyl ligase (FbiB) of *Mycobacterium tuberculosis*. *J Biol Chem* 291(13):6882-6894.

17. Li H, Graupner M, Xu H, & White RH (2003) CofE catalyzes the addition of two glutamates to F_{420}-0 in F_{420} coenzyme biosynthesis in *Methanococcus jannaschii*. *Biochemistry* 42(32):9771-9778.

18. Selengut JD & Haft DH (2010) Unexpected abundance of coenzyme F(420)-dependent enzymes in *Mycobacterium tuberculosis* and other actinobacteria. *J Bacteriol* 192(21):5788-5798.

19. Ney B*, et al.* (2017) The methanogenic redox cofactor F_{420} is widely synthesized by aerobic soil bacteria. *ISME J* 11(1):125-137.

20. Eirich LD, Vogels GD, & Wolfe RS (1978) Proposed structure for coenzyme F_{420} from *Methanobacterium*. *Biochemistry* 17(22):4583-4593.

21. Cheeseman P, Toms-Wood A, & Wolfe RS (1972) Isolation and properties of a fluorescent compound, factor 420 , from *Methanobacterium* strain M.o.H. *J Bacteriol* 112(1):527-531.

22. Lackner G, Peters EE, Helfrich EJ, & Piel J (2017) Insights into the lifestyle of uncultured bacterial natural product factories associated with marine sponges. *Proc Natl Acad Sci U S A* 114(3):E347-E356.

23. Lackner G, Moebius N, Partida-Martinez LP, Boland S, & Hertweck C (2011) Evolution of an endofungal lifestyle: deductions from the *Burkholderia rhizoxinica* genome. *BMC Genomics* 12:210.

24. Lackner G & Hertweck C (2011) Impact of endofungal bacteria on infection biology, food safety, and drug development. *PLoS Pathog* 7(6):e1002096.

25. Scherlach K, Busch B, Lackner G, Paszkowski U, & Hertweck C (2012) Symbiotic cooperation in the biosynthesis of a phytotoxin. *Angew Chem Int Ed Engl* 51(38):9615-9618.

26. Kiener A, Orme-Johnson WH, & Walsh CT (1988) Reversible conversion of coenzyme F_{420} to the 8-OH-AMP and 8-OH-GMP esters, F_{390}-A and F_{390}-G, on oxygen exposure and reestablishment of anaerobiosis in *Methanobacterium thermoautotrophicum*. *Arch Microbiol* 150(3):249-253.

27. Vermeij P, Pennings JL, Maassen SM, Keltjens JT, & Vogels GD (1997) Cellular levels of factor 390 and methanogenic enzymes during growth of *Methanobacterium thermoautotrophicum* deltaH. *J Bacteriol* 179(21):6640-6648.

28. Newton GL, Buchmeier N, & Fahey RC (2008) Biosynthesis and functions of mycothiol, the unique protective thiol of Actinobacteria. *Microbiol Mol Biol Rev* 72(3):471-494.

29. Zhang L, Loh KC, Lim JW, & Zhang JX (2019) Bioinformatics analysis of metagenomics data of biogas-producing microbial communities in anaerobic digesters: A review. *Renew Sust Energ Rev* 100:110-126.

30. Graupner M & White RH (2001) Biosynthesis of the phosphodiester bond in coenzyme F(420) in the methanoarchaea. *Biochemistry* 40(36):10859-10872.

31. Warkentin E*, et al.* (2001) Structures of $F_{420}H_2$: $NADP^+$ oxidoreductase with and without its substrates bound. *Embo J* 20(23):6561-6569.

32. Kunow J, Schwörer B, Stetter KO, & Thauer RK (1993) A F(420)-dependent NADP reductase in the extremely thermophilic sulfate-reducing *Archaeoglobus fulgidus*. *Arch Microbiol* 160(3):199-205.

33. Jirapanjawat T, *et al.* (2016) The redox cofactor F_{420} protects mycobacteria from diverse antimicrobial compounds and mediates a reductive detoxification system. *Appl Environ Microbiol* 82(23):6810-6818.

34. Mascotti ML, Kumar H, Nguyen QT, Ayub MJ, & Fraaije MW (2018) Reconstructing the evolutionary history of F-420-dependent dehydrogenases. *Sci Rep* 8.

35. Kumar H, Nguyen QT, Binda C, Mattevi A, & Fraaije MW (2017) Isolation and characterization of a thermostable F_{420}:NADPH oxidoreductase from *Thermobifida fusca*. *J Biol Chem* 292(24):10123-10130.