Metabolic pathway rerouting in *Paraburkholderia rhizoxinica* evolved long-overlooked derivatives

of coenzyme F$_{420}$.

Daniel Braga[a,d], Daniel Last[a], Mahmudul Hasan[a,d], Huijuan Guo[b], Daniel Leichnitz[b], Zerrin Uzum[c],

Ingrid Richter[c], Felix Schalk[b], Christine Beemelmanns[b], Christian Hertweck[c,d], and Gerald Lackner[a,d] [*]

Author affiliations

[a]Junior Research Group Synthetic Microbiology
Leibniz Institute for Natural Product Research and Infection Biology – Hans Knöll Institute
Beutenbergstr 11a, 07745 Jena, Germany
10
11 [b]Junior Research Group Chemical Biology of Microbe-Host Interactions
12 Leibniz Institute for Natural Product Research and Infection Biology – Hans Knöll Institute
13 Beutenbergstr 11a, 07745 Jena, Germany
14
15 [c]Department of Biomolecular Chemistry
16 Leibniz Institute for Natural Product Research and Infection Biology – Hans Knöll Institute
17 Beutenbergstr 11a, 07745 Jena, Germany
18
19 [d]Friedrich Schiller University, Jena
20
21 [*]**Corresponding author:**
22 Dr. Gerald Lackner
23 E-Mail: gerald.lackner@leibniz-hki.de
24 Phone: +49 3631 532 1104
25

Author contributions and conflict of interest

Daniel Braga performed research, analyzed data (molecular biology, malachite green assay, mass spectrometry)

and contributed to writing the manuscript, Daniel Last performed research and analyzed data (structure

elucidation, Fno assay, biogas plant studies), Mahmudul Hasan performed research (CofC/D enzyme assays),

Huijuan Guo performed research and analyzed data (structure elucidation), Daniel Leichnitz performed research

(chemical synthesis), Zerrin Uzum performed research (microscopy), Ingrid Richter performed research

(microscopy), Felix Schalk performed research (*cofE* constructs), Christine Beemelmanns designed research,

acquired funding, analyzed data (structure elucidation, synthesis) and edited the manuscript, Christian Hertweck

designed research, acquired funding and edited the manuscript, Gerald Lackner designed the study, acquired

funding and wrote the original manuscript. The authors declare no conflict of interest.

Keywords: Cofactors, natural products, biocatalysis, symbiosis, deazaflavin

Abstract

Coenzyme F_{420} is a specialized redox cofactor with a highly negative redox potential. It supports biochemical processes like methanogenesis, degradation of xenobiotics or the biosynthesis of antibiotics. Although well-studied in methanogenic archaea and actinobacteria, not much is known about F_{420} in Gram-negative bacteria. Genome sequencing revealed F_{420} biosynthetic genes in the Gram-negative, endofungal bacterium *Paraburkholderia rhizoxinica,* a symbiont of phytopathogenic fungi. Fluorescence microscopy, high-resolution LC-MS, and structure elucidation by NMR demonstrated that the encoded pathway is active and yields unexpected derivatives of F_{420} (3PG-F_{420}). Further analyses of a biogas-producing microbial community showed that these derivatives are more widespread in nature. Genetic and biochemical studies of their biosynthesis established that a specificity switch in the guanylyl transferase CofC re-programmed the pathway to start from 3-phospho-D-glycerate, suggesting a rerouting event during the evolution of F_{420} biosynthesis. Furthermore, the cofactor activity of 3PG-F_{420} was validated, thus opening up perspectives for its use in biocatalysis. The 3PG-F_{420} biosynthetic gene cluster is fully functional in *Escherichia coli*, enabling convenient production of the cofactor by fermentation.

Significance

Cofactors are essential molecules for many enzymatic reactions. Coenzyme F_{420} is an unusual redox cofactor found in prokaryotes involved in valuable enzymatic processes like biogas production, bioremediation, and antibiotics biosynthesis. Thus, it has attracted increasing interest in the field of biocatalysis and metabolic engineering. We discovered derivatives of F_{420} from bacteria living within fungal mycelia. Surprisingly, these molecules have been long overlooked in biogas-producing organisms. Our studies on their biosynthesis make an important contribution to validating and refining the just recently revised biochemical pathway of F_{420} formation. The novel cofactor is highly active and easily accessible, thus representing an ideal system for biotechnological applications. Its discovery will guide projects to unearth bioprocesses and enzymes involving F_{420}-derivatives in the future.

Introduction

Cofactors are essential for the catalytic power of many enzymes and thus play a key role in virtually all metabolic pathways. Knowledge of their catalytic functions and biosynthesis is highly important for the understanding of biochemical reactions as well as their application in biocatalysis and biotechnology. An important subclass comprises redox-cofactors that mediate electron transfer between molecules. The deazaflavin coenzyme F_{420} (Fig. 1) is a specialized redox cofactor with a lower redox potential (−350 mV) than NAD (1). This feature makes F_{420} an ideal electron carrier between H_2

and NAD(P) in methanogenesis and renders it a strong reducing agent for challenging reactions in biocatalysis (2). For instance, enzymatic processes involving F_{420} facilitate the degradation of pollutants like aromatic nitro compounds (3) or the carcinogen aflatoxin (4). Furthermore, F_{420}-dependent enzymes are important for asymmetric ene reductions (5, 6). In actinomycetes, coenzyme F_{420} is involved in the biosynthesis of antibiotics like oxytetracycline (7), pyrrolobenzodiazepines (8), or thiopeptins (9). Additionally, F_{420} has attracted considerable interest as a fitness factor of the human pathogen *Mycobacterium tuberculosis*, being involved in nitrosative stress response (10) or prodrug-activation (11).

A key step during the biosynthesis of F_{420} is the formation of the deazaflavin fluorophore F_O (Fig. 1A), a stable metabolic precursor of F_{420} originating from tyrosine and an intermediate of riboflavin biosynthesis. This chemically challenging step is catalyzed by the radical SAM enzyme complex CofG/H in archaea or the homologous dual-domain protein FbiC in actinobacteria (12). F_O is then further processed by CofC and CofD. This pair of enzymes is responsible for the biosynthesis of the 2-phospho-L-lactate (2-PL) moiety (Fig. 1B). Previous studies have shown that CofC from *Methanocaldococcus jannaschii* directly activates 2-phospho-L-lactate (2-PL) by guanylation (13) resulting in the formation of the short-lived metabolite lactyl-diphosphoguanosine (LPPG). CofD then forms F_{420}-0 by transfer of the 2-PL moiety from LPPG to F_O (14). The enzymes producing 2-PL, however, have remained elusive in all F_{420} producers so far. Recently, Bashiri *et al.* proposed a revised biosynthetic pathway by demonstrating that phosphoenolpyruvate (PEP) instead of 2-PL can serve as a substrate of CofC in *Mycobacteria* (15)*.* The resulting dehydro-F_{420} (DF$_{420}$) is reduced to F_{420} by a flavin-dependent reductase domain present in the FbiB protein (Fig. 1C). In mycobacteria, FbiB is a dual-domain protein consisting of a γ-glutamyl ligase domain (CofE-like) and the C-terminal DF$_{420}$ reductase domain (16). The γ-glutamyl ligase CofE finally decorates F_{420}-0 with a varying number of (oligo-)γ-glutamate residues (17). F_{420} is not ubiquitous in prokaryotes, but it is associated with certain phyla (18, 19). First discovered in methanogenic archaea (20, 21), it was extensively studied as a potential drug target of pathogenic mycobacteria (10) or as a cofactor enabling antibiotics biosynthesis in streptomycetes (2). Genome sequencing revealed that some Gram-negative bacteria have acquired F_{420} genes by horizontal transfer (19, 22). However, virtually nothing is known about the biosynthesis and role of F_{420} in these organisms. By genome mining, we found a biosynthetic gene cluster (BGC) homologous to those previously implicated in the biosynthesis of F_{420} in the endofungal bacterium *Paraburkholderia rhizoxinica* HKI 454 (Fig. 2A and *SI Appendix*, Table S3). This organism is an intracellular endosymbiont of the phytopathogenic fungus *Rhizopus microsporus* supplying its host with antimitotic toxins that act as virulence factors during infection of rice plants (23-25). We hypothesized that genes from Gram-negative bacteria related to F_{420} biosynthesis could facilitate F_{420} production in *E. coli* or could reveal

novel biosynthetic routes towards this valuable molecule. Therefore, we set out to investigate if the

BGC is active and if it can be refactored to produce F_{420} in *E. coli*.

Here, we show that *P. rhizoxinica* produces unexpected F_{420}-derivatives (3PG-F_{420}) both in symbiosis as

well as in axenic culture. Heterologous expression and large-scale production in *E. coli* allowed for

elucidation of their chemical structure. By comparative analyses, we discovered related metabolites in

a biogas-producing microbial community, thus indicating their broader abundance and relevance.

Enzyme assays showed that a switch in substrate specificity of CofC is responsible for the biosynthesis

of 3PG-F_{420} and proved that it can serve as a substitute for F_{420} in biochemical reactions.

Results

Discovery and structure elucidation of 3PG-F_{420}

To test whether *P. rhizoxinica* is capable of producing deazaflavins, we investigated axenic cultures of

symbiont (*P. rhizoxinica*) and host (*R. microsporus*) as well as symbiotic cultures by fluorescence

microscopy (Figs. 2B-I and *SI Appendix,* Figs. S1-4). Indeed, symbiotic *P. rhizoxinica* and the cytosol of

colonized mycelia emitted strong fluorescence characteristic of deazaflavins. Notably, even

fluorescence of bacteria present inside of fungal spores was observed. Axenic fungi, however, showed

no fluorescence, whereas only low signals were measured from axenic bacteria under the same

conditions. To corroborate the results obtained by microscopy, we extracted metabolites from axenic

P. rhizoxinica, axenic *R. microsporus* as well as fungal host containing endosymbionts and analyzed the

extracts by LC-MS/MS. To our surprise, only F_O could be detected in axenic *P. rhizoxinica* and in the

fungal host containing endosymbionts, but none of the expected F_{420}-n species.

To further test the biosynthetic capacity of the full BGC, we refactored it to obtain a single operon

under the control of a T7 promoter for heterologous expression yielding *E. coli/*pDB045 (Fig. 2J).

Examination of transformed bacteria by fluorescence microscopy revealed strong fluorescence as in

the endofungal bacteria (Fig. 2D), but LC-MS analyses again yielded only mass traces of F_O (mainly

found in the culture supernatant), but not of F_{420}-n. Since the impeded production could be attributed

to nonfunctional proteins, we analyzed proteins by SDS-PAGE and found that CofD was poorly soluble

in *E. coli*. Replacement of the *cofD* gene with the corresponding *M. jannaschii* homolog (14) provided

soluble protein (*E. coli/*pDB060)*,* however, again no trace of F_{420} could be detected. Therefore, we

reexamined the metabolome of *E. coli/*pDB045 for characteristic MS/MS fragments derived from the

F_O moiety (*m/z* 230.06, 346.10, 364.11). Surprisingly, the analysis revealed spectra with a similar

fragmentation pattern, yet derived from precursor ions that had a mass shift of 15.995 compared to

F_{420}, indicating the presence of an additional oxygen atom (Figs. 3A-B and *SI Appendix*, Figs. 5-8).

Further analyses revealed an (oligo-)γ-glutamate series of the oxygenated compound suggesting these

species are congeners. According to MS/MS fragmentation, the additional oxygen was present in the

"phospholactyl" moiety of F_{420}-n thus forming a "phosphoglyceryl" moiety. Extensive 1D- and 2D-NMR

experiments (Fig. 3C and *SI Appendix, Section 2.2*) and comparison to classical F_{420} (20) corroborated

that this moiety corresponds to 3-phosphoglycerate (3-PG). Therefore, we named the molecules 3PG-

F_{420}. This finding was unexpected because 2-phosphoglycerate is structurally more similar to PEP and

2-PL than 3-PG. Chemical degradation followed by chiral UHPLC-MS finally substantiated that the

additional stereocenter of 3PG-F_{420} is *R*-configured (Fig. 3D and *SI Appendix*, Fig. S51). Large-scale

cultivation of *E. coli*/pDB045 also revealed traces of dehydro-F_{420} (DF_{420}), but the yields were too low

for NMR studies. To investigate if 3PG-F_{420} is produced by wild-type *P. rhizoxinica*, we reanalyzed LC-

MS data for the presence of corresponding mass signals. Indeed, 3PG-F_{420} species were found in

samples containing bacteria (axenic culture, symbiosis), but not in symbiont-free host mycelia (*SI

Appendix*, Figs. S9-14). In extracts of *P. rhizoxinica*, no traces of F_{420} and DF_{420} were detected. The

presence of 3PG-F_{420} was restricted to the cell pellet, whereas F_O was abundant in culture supernatants.

We thus conclude that the fluorescence observed in bacterial cells of *P. rhizoxinica* is derived from

3PG-F_{420} and F_O.

Biosynthesis of 3PG-F_{420}

In order to rationalize how the biosynthetic pathway was redirected to form 3PG-F_{420} instead of F_{420},

we examined key steps of the biosynthesis more closely. We observed that production of 3PG-F_{420} was

not abolished by the exchange of c*ofD* from *P. rhizoxinica* by *cofD* from *M. jannaschii* (plasmid

pDB060). Hence, the phospholactyl transferase CofD could not be held accountable for the switch

towards 3PG-F_{420}. According to the existing biosynthetic model, the most plausible scenario was that

CofC incorporated 3-phospho-D-glycerate (3-PG), an intermediate of glycolysis, instead of 2-PL to form

3PG-F_{420}-0 and, to a minor extent, PEP to form DF_{420}-0. To test this hypothesis, we exchanged *cofC* and

cofD by the corresponding *M. jannaschii* homologs. The resulting strain *E. coli*/pDB070 (Fig. 2J and *SI

Appendix*, Fig. S18) produced neither 3PG-F_{420} nor F_{420} but traces of DF_{420}. To further investigate the

substrate specificity of CofC, we performed an in-vitro assay using CofC and CofD (13). Genes *cofC* of

P. rhizoxinica as well as *cofC* and *cofD* from *M. jannaschii* were cloned, corresponding proteins

produced as hexahistidine fusions in *E. coli* and purified by metal affinity chromatography for in-vitro

assays. The physiologically relevant isomers 2-phospho-D-glycerate (2-PG) and 3-phospho-D-glycerate

(3-PG), as well as PEP and 2-PL, served as substrates. Reaction products were monitored by LC-MS.

Indeed, when CofC from *P. rhizoxinica* was tested, the mass of 3PG-F_{420}-0 appeared after reaction with

D-3-PG eluting at the same retention time as the in-vivo product (Fig. 4A and *SI Appendix*, Section 2.4).

In addition, the formation of DF_{420}-0 and F_{420}-0 was detected, when the enzymes were incubated with

PEP and 2-PL, respectively. In contrast, reaction with 2-PG yielded mass signals close to the noise level.

Controls lacking CofC did not generate any of these products. In a direct substrate competition assay,

3-PG was found to be the preferred substrate with a relative turnover of ca. 72% (2-PL: 23% PEP: 4%).

This finding is in agreement with the structure of 3PG-F_{420} and the occurrence of DF_{420} as a minor

biosynthetic product in bacteria. Note that CofC from *M. jannaschii* displayed a strong turnover of 2-

PL (96.5%), weak turnover of PEP (3.5%) and no turnover of 3-PG. This finding supports the notion that

the CofC of *P. rhizoxinica* has undergone a substrate specificity switch during evolution.

To address the question if the γ-glutamyl ligase CofE represents another specificity layer that directs

the biosynthesis of F_{420} towards 3PG-F_{420}, we individually co-expressed *cofE* genes from *P. rhizoxinica*,

M. jannaschii and *Mycolicibacterium smegmatis* (*fbiB*) together with a minimal BGC consisting of *fbiC*,

cofC, and *cofD* in a two-plasmid system. Extraction of metabolites and LC-MS/MS revealed that all

three CofE homologs elongated 3PG-F_{420}-0 to oligo-glutamate chain lengths up to n=6 (*SI Appendix*,

Figs. S63-65). Thus, we conclude that CofE does not act as an additional specificity filter during chain

elongation of 3PG-F_{420}.

Cofactor role of 3PG-F_{420}

The successful isolation of 3PG-F_{420} and reconstitution of its biosynthesis in *E. coli* motivated us to

address the question of whether 3PG-F_{420} could substitute F_{420} in biocatalysis. To this end, we cloned

a gene encoding Fno (F_{420}:NADPH oxidoreductase), an enzyme that serves as a regeneration system

for $F_{420}H_2$ using NADPH/H^+ as an electron donor (26). We first examined if Fno can accept 3PG-F_{420} as

a substrate. Indeed, we observed an efficient reduction of 3PG-F_{420} by recombinant Fno as mirrored by

a rapid decrease of characteristic UV absorption. An examination of kinetic parameters (Fig. 5A)

revealed that the apparent K_M of Fno for F_{420} was 3.6 ± 0.7 μM. This value is similar to the reported K_M

of 10 μM (27). Under identical assay conditions, the K_M for 3PG-F_{420} was only slightly higher (5.1 ± 1.0

μM). The v_{max} values were in a similar range as well (F_{420}: 1.3 ± 0.2 μmol min^{-1}; 3PG-F_{420}: 0.88 ± 0.07

μmol min^{-1}) pointing towards only a minor reduction of maximal turnover. Encouraged by the finding

that 3PG-F_{420} can substitute F_{420}, we aimed at an in-vivo application of the cofactor for malachite green

reduction as a proof of principle. To this end, we combined the *fno* gene with a minimal BGC producing

3PG-F_{420}-0 (Fig. 5C and *SI Appendix,* Fig. S62) on a single vector (pDB071). Additionally, the F_{420}-

dependent malachite green reductase gene MSMEG_5998 (28) from *M. smegmatis* was cloned and

expressed from a compatible vector backbone (pDB061)*.* Finally, co-expression of all components in *E.*

coli yielded a strain (pDB061/pDB071) that was able to decolorize malachite green significantly faster

than control strains expressing the reductase or the cofactor alone (Fig. 5B). We, therefore, conclude

that 3PG-F_{420} can substitute F_{420} as a redox cofactor in this case.

Occurrence of 3PG-F_{420} in nature

Finally, we set out to test if 3PG-F_{420} is restricted to fungal endosymbionts or if it might be more

widespread in the environment. We examined *M. jannaschii* and *M. smegmatis* for the presence of

any F_{420} congeners*.* Only F_{420}-n was found from extracts of *M. jannaschii,* while F_{420}-n and DF$_{420}$-n were

detected in extracts of *M. smegmatis*. (*SI Appendix*, Figs. S15-16). None of the 3PG-F_{420}-derivatives

were found in the reference organisms. Next, we analyzed (two independent) sludge samples from a

local biogas production plant. To our surprise, extraction of the microbial community present in the

biogas-producing sludge followed by LC-MS/MS eluted, besides classical F_{420}, a compound with

identical retention time, exact mass and MS/MS fragmentation pattern as 3PG-F_{420} (*SI Appendix*, Fig.

S17). Therefore, we conclude that microbes present in the biogas-producing microbial community

produce 3PG-F_{420}-derivatives or highly similar congeners.

Discussion

Unexpected biological sources of F_{420}-derivatives

In the course of this study, we discovered a BGC consisting of genes related to F_{420}-biosynthesis in the

Gram-negative bacterium *P. rhizoxinica*. We could show that the BGC is highly active and mediates

production of 3PG-F_{420}, a derivative of coenzyme F_{420}, thus revealing an untapped terrain of

deazaflavin-based biochemistry in Gram-negative bacteria. So far, there is only preliminary evidence

for the occurrence of deazaflavin cofactors in a few Gram-negative bacteria, e.g., *Oligotropha*

carboxidivorans and *Paracoccus dentrificans* (19) as well as in the uncultured, but biosynthetically

highly prolific '*Candidatus* Entotheonella factor' (22). The exact structure and function of F_{420} in most

Gram-negative bacteria that harbor corresponding biosynthetic genes, however, is unknown. The fact

that *P. rhizoxinica* produced 3PG-F_{420} under symbiotic cultivation conditions allows for the conclusion

that it provides a fitness benefit in its natural habitat. Notably, none of the well-characterized F_{420}-

dependent enzyme families (4, 18) are encoded in the *P. rhizoxinica* genome according to BLAST and

conserved domains searches, not even any of the widespread regeneration systems like Fno or F_{420}-

dependent glucose-6-phosphate dehydrogenase. Therefore, future investigations of 3PG-F_{420}-

producing organisms are likely to reveal novel enzymes, regeneration systems, and cellular pathways

depending on this cofactor. Intriguingly, substantial amounts of phosphoglycerate-derived

deazaflavins were also extracted from a microbial community of a biogas-production plant. Since

fluorescence and UV-based detection does usually not resolve classical F_{420} and 3PG-F_{420} these

derivatives might have been misidentified as F_{420} in the past. Since neither *P. rhizoxinica* nor its host *R.*

*microsporu*s are able to grow under anaerobic and thermophilic (temperatures >42 °C) conditions,

they can be excluded as the source of these cofactors. The high complexity and variability of biogas-

producing microbiomes (29) do not allow for an educated guess of the producer, although

methanogens would be reasonable candidates. In any case, phosphoglycerate-derived deazaflavins

are more abundant in nature than anticipated.

Structural diversification of deazaflavin cofactors

The structure and occurrence of 3PG-F_{420} has not been reported before. To date, the only known

derivatives of F_{420}-n are factor F_{390}-A and F_{390}-G, 8-OH-AMP and 8-OH-GMP esters of F_{420}, respectively

(30). In methanogens, they are formed reversibly, e.g., during oxygen exposure, acting as a reporter

compound for hydrogen starvation (31). In contrast, the modifications seen in 3PG-F_{420} are not

temporary. Rather, 3PG-F_{420} seems to replace F_{420} as a natural deazaflavin-cofactor in *P. rhizoxinica*. At

least in this organism, it does not coexist with classical F_{420}. This situation is reminiscent of mycothiol,

a specialized thiol cofactor that replaced glutathione in actinobacteria (32). Notably, the redox-active

deazaflavin moiety of 3PG-F_{420} remains unchanged and, therefore, redox activity of 3PG-F_{420} was

retained. The phospholactate and glutamate-derived tail moiety of F_{420} appears to have two functions.

Firstly, the tail avoids loss of the cofactor, since F_O, in contrast to F_{420}, is crossing the bacterial cell

membranes and is found mainly in culture supernatants. This function is also fulfilled by the altered

tail moiety of 3PG-F_{420}. Secondly, the tail of F_{420} provides additional contacts for non-covalent

interactions (33). Our first results show that the apparent K_M of Fno for 3PG-F_{420} is in a similar low

micromolar range as the K_M for F_{420}, thus indicating comparable binding affinity. Importantly, 3PG-F_{420}

can be used as a cofactor in *E. coli* to reduce malachite green. A similar result was just recently obtained

using classical F_{420} and a different malachite green reductase (15).

Rerouting of F_{420} biosynthetic pathways

In addition to the question if 3PG-F_{420} can act as a cofactor, we clarified how a F_{420}-derivative can be

produced in a metabolic background lacking DF_{420} reductase and 2-PL. Obviously, a substrate specificity

switch in CofC directed the biosynthesis towards 3-PG. Recently, Bashiri *et al.* claimed that PEP is the

substrate of CofC in prokaryotes (15). Our results partially confirm their hypothesis, since we observed

the turnover of PEP as a potential substrate of both CofC homologs we tested. However, in contrast to

Bashiri *et al.*, 2-PL was the best substrate of *M. jannaschii* CofC in our assay. No DF_{420} was measurable

in cell extracts of *M. jannaschii*, while it was readily detected in *M. smegmatis* cells. Since Graupner

and White detected significant amounts of 2-PL in methanogenic archaea and observed the conversion

of lactate into 2-PL by isotope labeling (34), we conclude that 2-PL might still be the relevant substrate

in archaea. From an evolutionary perspective, our results suggest that multiple metabolic re-wiring

events occurred in the evolution of F_{420} biosynthesis. While actinobacteria evolved the DF_{420} reductase

(C-terminal domain of FbiB), archaea accomplished to produce the (unusual) metabolite 2-PL. Other

organisms, as exemplified by *P. rhizoxinca,* rerouted the biosynthesis to the ubiquitous metabolite 3-

PG. Notably, the altered pathway offered another route to heterologous production of F_{420}-derivatives

in *E. coli*. The production of classical F_{420} in *E. coli* has just recently been achieved in moderate yields

using *Mycobacterium* genes including the DF_{420} reductase domain (15).

Prospects for biotechnology

In recent years, there has been increasing interest in F_{420}-dependent enzymes for biocatalytic

applications (5, 6, 35, 36). Therefore, the convenient production in *E. coli* along with the anticipation

to discover novel enzymes and regeneration systems specialized for 3PG-F_{420} opens intriguing

prospects for biocatalysis and metabolic engineering. Here, as a proof of concept, 3PG-F_{420} has been

shown to be easily reduced by Fno and to substitute F_{420} for malachite green reduction in vivo. Besides

future in-vitro applications, for instance enantioselective biotransformations, metabolically engineered microbes can be envisioned to use 3PG-F_{420} to promote challenging reactions of industrial appeal, to help in bioremediation or to produce high-value compounds like antibiotics in a sustainable manner.

Materials and methods

Materials and methods are summarized in *SI Appendix,* Section 1.

Acknowledgments

We thank Prof. Dr. Dina Grohmann and Dr. Ghader Bashiri for kindly providing *M. jannaschii* and *M. smegmatis / fbiABC,* respectively. We thank Biogas Jena GmbH and Co. KG for the kind donation of biogas plant samples. We thank Heike Heinecke for conducting NMR experiments. G.L. thanks the Deutsche Forschungsgemeinschaft (DFG Grant LA 4424/1-1) and the Carl Zeiss Foundation for funding. Financial support by the DFG (CRC 1127 ChemBioSys) to C.H. and C.B., and, BE-4799/2-1 to C.B., and Leibniz Award to C.H., by the ERC (MSCA-IF-EF-RI Project 794343, to I.R.) and the JSMC to Z.U. is gratefully acknowledged.

References

1. Jacobson F & Walsh C (1984) Properties of 7,8-didemethyl-8-hydroxy-5-deazaflavins relevant to redox coenzyme function in methanogen metabolism. *Biochemistry* 23(5):979-988.
2. Greening C*, et al.* (2016) Physiology, biochemistry, and applications of F_{420}- and F_o-dependent redox reactions. *Microbiol Mol Biol Rev* 80(2):451-493.
3. Heiss G*, et al.* (2002) npd gene functions of *Rhodococcus* (*opacus*) *erythropolis* HL PM-1 in the initial steps of 2,4,6-trinitrophenol degradation. *Microbiology* 148(Pt 3):799-806.
4. Taylor MC*, et al.* (2010) Identification and characterization of two families of F_{420} H_2-dependent reductases from Mycobacteria that catalyse aflatoxin degradation. *Mol Microbiol* 78(3):561-575.
5. Taylor M, Scott C, & Grogan G (2013) F_{420}-dependent enzymes - potential for applications in biotechnology. *Trends Biotechnol* 31(2):63-64.
6. Mathew S, Trajkovic M, Kumar H, Nguyen QT, & Fraaije MW (2018) Enantio- and regioselective ene-reductions using $F_{420}H_2$-dependent enzymes. *Chem Commun* 54(79):11208-11211.
7. Wang P, Bashiri G, Gao X, Sawaya MR, & Tang Y (2013) Uncovering the enzymes that catalyze the final steps in oxytetracycline biosynthesis. *J Am Chem Soc* 135(19):7138-7141.
8. Li W, Khullar A, Chou S, Sacramo A, & Gerratana B (2009) Biosynthesis of sibiromycin, a potent antitumor antibiotic. *Appl Environ Microbiol* 75(9):2869-2878.
9. Ichikawa H, Bashiri G, & Kelly WL (2018) Biosynthesis of the Thiopeptins and Identification of an $F_{420}H_2$-Dependent Dehydropiperidine Reductase. *J Am Chem Soc* 140(34):10749-10756.
10. Purwantini E & Mukhopadhyay B (2009) Conversion of NO_2 to NO by reduced coenzyme F420 protects mycobacteria from nitrosative damage. *Proc Natl Acad Sci U S A* 106(15):6333-6338.
11. Singh R*, et al.* (2008) PA-824 kills nonreplicating *Mycobacterium tuberculosis* by intracellular NO release. *Science* 322(5906):1392-1395.

12. Decamps L, *et al.* (2012) Biosynthesis of F_0, precursor of the F_{420} cofactor, requires a unique two radical-SAM domain enzyme and tyrosine as substrate. *J Am Chem Soc* 134(44):18173-18176.

13. Grochowski LL, Xu H, & White RH (2008) Identification and characterization of the 2-phospho-L-lactate guanylyltransferase involved in coenzyme F_{420} biosynthesis. *Biochemistry* 47(9):3033-3037.

14. Graupner M, Xu H, & White RH (2002) Characterization of the 2-phospho-L-lactate transferase enzyme involved in coenzyme F(420) biosynthesis in *Methanococcus jannaschii*. *Biochemistry* 41(11):3754-3761.

15. Bashiri G, *et al.* (2019) A revised biosynthetic pathway for the cofactor F_{420} in prokaryotes. *Nat Commun* 10(1):1558.

16. Bashiri G, *et al.* (2016) Elongation of the poly-gamma-glutamate Tail of F_{420} requires both domains of the F420:gamma-glutamyl ligase (FbiB) of *Mycobacterium tuberculosis*. *J Biol Chem* 291(13):6882-6894.

17. Li H, Graupner M, Xu H, & White RH (2003) CofE catalyzes the addition of two glutamates to F_{420}-0 in F_{420} coenzyme biosynthesis in *Methanococcus jannaschii*. *Biochemistry* 42(32):9771-9778.

18. Selengut JD & Haft DH (2010) Unexpected abundance of coenzyme F(420)-dependent enzymes in *Mycobacterium tuberculosis* and other actinobacteria. *J Bacteriol* 192(21):5788-5798.

19. Ney B, *et al.* (2017) The methanogenic redox cofactor F_{420} is widely synthesized by aerobic soil bacteria. *ISME J* 11(1):125-137.

20. Eirich LD, Vogels GD, & Wolfe RS (1978) Proposed structure for coenzyme F_{420} from *Methanobacterium*. *Biochemistry* 17(22):4583-4593.

21. Cheeseman P, Toms-Wood A, & Wolfe RS (1972) Isolation and properties of a fluorescent compound, factor 420 , from *Methanobacterium* strain M.o.H. *J Bacteriol* 112(1):527-531.

22. Lackner G, Peters EE, Helfrich EJ, & Piel J (2017) Insights into the lifestyle of uncultured bacterial natural product factories associated with marine sponges. *Proc Natl Acad Sci U S A* 114(3):E347-E356.

23. Lackner G, Moebius N, Partida-Martinez LP, Boland S, & Hertweck C (2011) Evolution of an endofungal lifestyle: deductions from the *Burkholderia rhizoxinica* genome. *BMC Genomics* 12:210.

24. Lackner G & Hertweck C (2011) Impact of endofungal bacteria on infection biology, food safety, and drug development. *PLoS Pathog* 7(6):e1002096.

25. Scherlach K, Busch B, Lackner G, Paszkowski U, & Hertweck C (2012) Symbiotic cooperation in the biosynthesis of a phytotoxin. *Angew Chem Int Ed Engl* 51(38):9615-9618.

26. Warkentin E, *et al.* (2001) Structures of $F_{420}H_2$: $NADP^+$ oxidoreductase with and without its substrates bound. *Embo J* 20(23):6561-6569.

27. Kunow J, Schwörer B, Stetter KO, & Thauer RK (1993) A F(420)-dependent NADP reductase in the extremely thermophilic sulfate-reducing *Archaeoglobus fulgidus*. *Arch Microbiol* 160(3):199-205.

28. Jirapanjawat T, *et al.* (2016) The Redox Cofactor F_{420} Protects Mycobacteria from Diverse Antimicrobial Compounds and Mediates a Reductive Detoxification System. *Appl Environ Microbiol* 82(23):6810-6818.

29. Zhang L, Loh KC, Lim JW, & Zhang JX (2019) Bioinformatics analysis of metagenomics data of biogas-producing microbial communities in anaerobic digesters: A review. *Renew Sust Energ Rev* 100:110-126.

30. Kiener A, Orme-Johnson WH, & Walsh CT (1988) Reversible conversion of coenzyme F_{420} to the 8-OH-AMP and 8-OH-GMP esters, F_{390}-A and F_{390}-G, on oxygen exposure and reestablishment of anaerobiosis in *Methanobacterium thermoautotrophicum*. *Arch Microbiol* 150(3):249-253.

31. Vermeij P, Pennings JL, Maassen SM, Keltjens JT, & Vogels GD (1997) Cellular levels of factor 390 and methanogenic enzymes during growth of *Methanobacterium thermoautotrophicum* deltaH. *J Bacteriol* 179(21):6640-6648.

32. Newton GL, Buchmeier N, & Fahey RC (2008) Biosynthesis and functions of mycothiol, the unique protective thiol of Actinobacteria. *Microbiol Mol Biol Rev* 72(3):471-494.

33. Ney B, *et al.* (2017) Cofactor tail length modulates catalysis of bacterial F-420-dependent oxidoreductases. *Front Microbiol* 8:1902.

34. Graupner M & White RH (2001) Biosynthesis of the phosphodiester bond in coenzyme F(420) in the methanoarchaea. *Biochemistry* 40(36):10859-10872.

35. Mascotti ML, Kumar H, Nguyen QT, Ayub MJ, & Fraaije MW (2018) Reconstructing the evolutionary history of F-420-dependent dehydrogenases. *Sci Rep* 8.

36. Kumar H, Nguyen QT, Binda C, Mattevi A, & Fraaije MW (2017) Isolation and characterization of a thermostable $F_{420:}$NADPH oxidoreductase from *Thermobifida fusca*. *J Biol Chem* 292(24):10123-10130.

Figures

Fig. 1. Biosynthesis of coenzyme F_{420} A) F_O synthase FbiC (in archaea: CofG/H) catalyzes formation of the deazaflavin ring from tyrosine and 5-amino-6-(ribitylamino)-uracil, an intermediate of riboflavin biosynthesis. B) Biosynthetic scheme of F_{420}-n proposed for archaea. CofC and CofD catalyze the activation of 2-PL and transfer of the 2-PL moiety, respectively. CofE performs (oligo-)γ-glutamylation. The number of glutamate residues (n) varies depending on the organism. The origin of 2-PL is unknown. C) Biosynthesis of F_{420}-n proposed for mycobacteria: CofC and CofD activate PEP resulting in DF_{420} formation. The C-terminal domain of FbiB reduces DF_{420} to F_{420}. EPPG: enolpyruvyl-diphosphoguanosine, LPPG: lactyl-diphosphoguanosine, 2-PL: 2-phospho-L-lactate.

Fig. 2. Deazaflavin biosynthesis in *P. rhizoxinica.* A) BGC of 3PG-F$_{420}$. Core genes are shown in dark grey. B-I) Microscopy photographs depict fluorescence characteristic of deazaflavins in blue. B-D) axenic *M. smegmatis* (B), *P. rhizoxinica* (C), and *E. coli* / pDB045 (D). In *R. microsporus* ATCC 62417, deazaflavins are correlated to the presence of *P. rhizoxinica* symbionts (green, Syto9 staining) (E). No fluorescence was detected in cured ATCC 62417 mycelium (F) or in the naturally symbiont-free strain, CBS 344.29 (G). The same pattern was observed in spores of either wild-type ATCC 62417 (H) or CBS 344.29 (I). Scale bars represent 10 µm. J) Refactored versions of the BGC and corresponding plasmids for heterologous expression in *E. coli.* Asterisks mark genes from *M. jannaschii.*

Fig. 3. Chemical analysis of 3PG-F$_{420}$. A) Extracted ion chromatograms of 3PG-F$_{420}$-2 produced in *E. coli*.

I: *E. coli* / pDB045, II: *cofD* exchanged by *M. jannaschii* homolog (pDB060), III: *cofD* and *cofC* exchanged

by *M. jannaschii* homologs (pDB070) IV: empty vector (pETDuet). B) Excerpt of the MS/MS spectrum

of 3PG-F$_{420}$-2. Grey bars highlight m/z used for fragment ion search of F$_{420}$ derivatives. C) ^{1}H NMR

comparison of F$_{420}$-n (D$_2$O), 3PG-F$_{420}$-n (0.1% ND$_3$ in D$_2$O) and 3PG-F$_{420}$-0 (0.1% ND$_3$ in D$_2$O) indicated

the replacement of the lactyl moiety in F$_{420}$ with a glyceryl moiety in 3PG-F$_{420}$. D) Proposed structures

of 3PG-F$_{420}$-0, 3PG-F$_{420}$-n, and DF$_{420}$-n.

A

B

3-D-PG   3-GPPG   3PG-F$_{420}$-0

Fig. 4. Combined CofC/D in-vitro assay. A) Relative turnover of substrates estimated from a substrate competition assays (D-3-PG, 2-PL, and PEP). CofC from *P. rhizoxinica* accepted 3-PG (72.7%), 2-PL (23.4%), and PEP (3.9%). CofC from *M. jannaschii* preferred 2-PL (96.5%), and PEP (3.5%). 3-PG was not turned over. CofD from *M. jannaschii* was used in all assays. Error bars represent the standard deviation (SD) of three independent biological replicates (N=3). B) Proposed model of 3PG-F$_{420}$ biosynthesis. 3-GPPG: 3-(guanosine-5'-disphospho)-D-glycerate.

Fig. 5. Cofactor function of 3PG-F_{420}. A) Michaelis-Menten kinetics of Fno for F_{420} (left) and 3PG-F_{420} as

substrates (right). Three biological replicates were used to determine parameters. K_M for F_{420} was 3.6

± 0.7 µM (standard error). K_M for 3PG-F_{420} was 5.1 ± 1.0 µM. Error bars indicate standard deviation of

replicates (N=3). B) In-vivo reduction of malachite green (absorbance: 618 nm) by the F_{420}-dependent

reductase MSMEG_5998. Fno was used to regenerate 3PG-$F_{420}H_2$. Left panel: Time course of the

malachite green depletion assay. Right panel: Bar chart of residual malachite green after 20 h: wt: *E.*

coli BL21(DE3), pDB061: *E. coli* producing MSMEG_5998, pDB071: *E. coli* producing 3PG-F_{420}-0 + Fno.

Exact means ± SD of biological triplicates were 0.234 ± 0.017 (wt), 0.124 ± 0.003 (pDB061), 0.169 ±

0.011 (pDB071), and 0.082 ± 0.0139 (pDB061/pDB071). An asterisk indicates statistical significance

(one-way ANOVA, p<0.05, N=3). C) Engineered *E. coli* combining 3PG-F_{420}, Fno and reductase

MSMEG_5998 (*red*) for reduction of malachite green.

16