1

The turning and barrier course reveals gait parameters for detecting freezing of gait and

measuring the efficacy of deep brain stimulation

Johanna J. O'Day[1,2] , Judy Syrkin-Nikolau[3], Chioma M. Anidi[4], Lukasz Kidzinski[1],

Scott L. Delp[1], Helen M. Bronte-Stewart[2,5*]

[1] Department of Bioengineering, Stanford University, Stanford, California, United States of

America

[2] Stanford University Department of Neurology and Neurological Sciences, Stanford University,

Stanford, California, United States of America

[3] Cala Health, Burlingame, California, United States of America

[4] University of Michigan Medical School, Ann Arbor, Michigan, United States of America

[5] Department of Neurosurgery, Stanford University, Stanford, California, United States of

America

* Corresponding author

E-mail: hbs@stanford.edu (HBS)

Abstract

Freezing of gait (FOG) is a devastating motor symptom of Parkinson's disease that leads to falls, reduced mobility, and decreased quality of life. Reliably eliciting FOG has been difficult in the clinical setting, which has limited discovery of pathophysiology and/or documentation of the efficacy of treatments, such as different frequencies of subthalamic deep brain stimulation (STN DBS). In this study we validated an instrumented gait task, the turning and barrier course (TBC), with the international standard FOG questionnaire item 3 (FOG-Q3, r = 0.74, $p < 0.001$). The TBC is easily assembled and mimics real-life environments that elicit FOG. People with Parkinson's disease who experience FOG (freezers) spent more time freezing during the TBC compared to during forward walking ($p = 0.007$). Freezers also exhibited greater arrhythmicity during non-freezing gait when performing the TBC compared to forward walking ($p = 0.006$); this difference in gait arrhythmicity between tasks was not detected in non-freezers or controls. Freezers' non-freezing gait was more arrhythmic than that of non-freezers or controls during all walking tasks ($p < 0.05$). A logistic regression model determined that a combination of gait arrhythmicity, stride time, shank angular range, and asymmetry had the greatest probability of classifying a step as FOG (area under receiver operating characteristic curve = 0.754). Freezers' percent time freezing and non-freezing gait arrhythmicity decreased, and their shank angular velocity increased in the TBC during both 60 Hz and 140 Hz STN DBS ($p < 0.05$) to non-freezer values. The TBC is a standardized tool for eliciting FOG and demonstrating the efficacy of 60 Hz and 140 Hz STN DBS for gait impairment and FOG. The TBC revealed gait parameters that differentiated freezers from non-freezers and best predicted FOG; these may serve as relevant control variables for closed loop neurostimulation for FOG in Parkinson's disease.

Introduction

Gait impairment and freezing of gait (FOG) are common in Parkinson's disease, and lead to falls, [1–3] resulting in injury, loss of independence, institutionalization, and even death [4,5]. Over 10 million people are affected by Parkinson's disease (PD) worldwide, and over 80% of people with moderate to advanced PD develop FOG [6]. Gait impairment is characterized by the loss of regular rhythmic alternating stepping associated with normal locomotion. FOG is an intermittent, involuntary inability to perform alternating stepping and usually occurs when patients attempt to initiate walking, turn, or navigate obstacles.

Understanding and treating gait impairment and FOG are paramount unmet needs and were given the highest priority at the National Institute of Neurological Disorders and Stroke 2014 PD conference [7]. Both gait impairment and FOG have unpredictable responses to dopaminergic medication and continuous high frequency open loop subthalamic deep brain stimulation (DBS) [8,9]. Although gait impairment and FOG may improve on continuous lower frequency (60 Hz) DBS, Parkinsonian tremor may worsen, and many patients do not tolerate 60 Hz DBS for long periods of time [10–12]. A closed loop, adaptive system that can adjust stimulation appropriately may be able to improve therapy for FOG and impaired gait. Before this goal can be attained, however, it is important to determine which gait parameters are associated with freezing behavior, which predict freezing events, and the effect of different DBS frequencies on gait impairment and FOG.

Several studies have employed wearable inertial sensors to monitor, detect, and predict FOG using a variety of different gait parameters. The most popular approach has been to use a frequency-based analysis of leg accelerations to capture the "trembling of knees" associated with FOG, and many variations on this approach have been described including the "freeze index"

[13] and "Frequency Ratio" [14]. These studies have employed a variety of different FOG-

eliciting tasks, such as turning 360 degrees in place for two minutes, walking around cones, or

walking during dual tasking [14–22]. These tasks have improved the detection of FOG but are

not representative of real-world environments, or cannot objectively measure gait impairment,

such as arrhythmicity, which has been correlated with FOG [23–27]. Objective gait measures

and standardized gait tasks that reliably induce FOG are needed to study the progression of gait

impairment and FOG in PD, and the efficacy of therapeutic interventions.

The goals of this study were to (1) validate a standardized gait task, the turning and

barrier course (TBC), which mimics real-life environments and elicits FOG, (2) discover relevant

gait parameters for detecting FOG in Parkinson's disease in the TBC, and (3) evaluate the effects

of 60 Hz and 140 Hz subthalamic deep brain stimulation (DBS) on quantitative measures of non-

freezing gait and FOG.

Materials and methods

Human subjects

Twenty-three subjects with PD (8 female), and 12 age-matched healthy controls (11

female), participated in the study. Subjects were recruited from the Stanford Movement

Disorders Center and were not pre-selected based on a history of FOG. Subjects were excluded if

they had peripheral neuropathy, hip or knee prostheses, structural brain disorders, or any visual

or anatomical abnormalities that affected their walking. For all PD subjects, long-acting

dopaminergic medication was withdrawn over 24h (72h for extended-release dopamine

agonists), and short-acting medication was withdrawn over 12h before all study visits. A

certified rater performed the Unified Parkinson's Disease Rating Scale (UPDRS III) motor

disability scale [28], and the Freezing of Gait Questionnaire (FOG-Q, Giladi et al., 2009) on all

subjects. Four subjects had FOG-Q scores taken from a prior research visit within the last 4

months. Subjects were classified as a freezer or non-freezer based on the FOG-Q question 3

(FOG-Q3): *Do you feel that your feet get glued to the floor while walking, turning or when trying*

to initiate walking? The scores were as follows: *0 – never, 1 – about once a month, 2 – about*

once a week, 3 – about once a day, 4 – whenever walking. A freezer was defined as a subject

who reported a FOG-Q3 $\geq$ 2 or if the subject exhibited a freezing event during the tasks. Control

subjects were excluded if they reported neurological deficits or interfering pathology that

affected their gait. All subjects gave their written informed consent to participate in the study,

which was approved by the FDA and the Stanford IRB.

Experimental protocol

All experiments were performed off therapy (medication and/or DBS). Subjects

performed two gait tasks: Forward Walking (FW), which is a standard clinical test of

Parkinson's gait, and the TBC, in a randomized order at their self-selected pace. Both tasks

started with 20s of quiet standing. For the FW task, subjects walked in a straight line for 10m,

turned around and returned, and repeated this for a total of 40 m. We only analyzed data from the

straight walking parts of FW. The FW task was conducted in a hallway at least 1.7 m wide

formed by a wall and room dividers (Bretford Mobile Screens, Pivot Interiors Inc., Pleasanton,

CA). The room dividers were 1.98 m high and a maximum of 1.14 m wide. In the TBC, subjects

walked around and through a narrow opening formed by room dividers [25], Fig 1A.

Fig 1. **Turning and Barrier Course (TBC) dimensions and specifications.**

(A) The individual barrier and course dimensions. Tall barriers limited vision around turns and narrow passageways to simulate a real-world environment. (B) Front view with patient walking in the TBC and (C) aerial diagram of the TBC with barriers (dark grey bars) and wall (light grey bar). Subjects walked in two ellipses and then two figures of eight around the barriers; this task was repeated starting on both the left and right side, for a total of four ellipses and four figures of eight.

The TBC was enclosed by a row of dividers on one side and a wall on the other, Fig 1B, which limited the subjects' visual field; the aisles of the TBC were the same width as a standard minimum hallway (0.91 meters) in the U.S., and the narrow opening between dividers was the same width as a standard doorway (0.69 meters), Fig 1A. After the initial standing rest period, the subject was instructed to sit on the chair. At the 'Go' command, the subject was instructed to stand up, walk around the dividers twice in an ellipse, and then walk in a 'figure eight' (i.e., around and through the opening between the dividers), twice, before sitting down again, Fig 1C. The subject was then instructed to repeat the task in the opposite direction, for a total of four ellipses and four figures of eight. The direction order was randomized.

Data acquisition and analysis

Shank angular velocity was measured during the gait tasks using wearable inertial

measurement units (IMUs, APDM, Inc., Portland, OR), which were positioned in a standardized

manner on the lateral aspect of both shanks. We aligned the IMU on the shank so that the

positive Z-axis was directed laterally and measured the angular velocity of the shank in the

sagittal plane. Signals from the IMUs' triaxial gyroscope and accelerometer and magnetometer

were sampled at 128 Hz. The data were filtered using a zero phase 8[th] order low pass

Butterworth filter with a 9 Hz cut-off frequency, and principal component analysis was used to

align the angular velocity with the sagittal plane. Using the sagittal plane angular velocity, the

beginning of the swing phase (positive slope zero crossing), end of swing phase (subsequent

negative slope zero crossing), and peak shank angular velocities (first positive peak following the

beginning of swing phase) were identified, Fig 2.

Fig 2. Gait parameters extracted from inertial measurement units (IMU).

(Top) Schematic of one gait cycle with IMU on the shank used to define gait parameters

including stride time, forward swing time, swing angular range and peak angular velocities (peak

AV). (Bottom) Gait parameters extracted from shank sagittal angular velocity data for the left

(blue) and right (red) legs during periods of non-freezing walking, and freezing of gait (orange).

Forward swing times (time between subsequent zero crossings of the same leg) and stride times

(time between consecutive peak angular velocities) were calculated from these data, Fig 2.

Swing angular range was calculated by integrating the sagittal angular velocity curve during the

swing time. Swing times and stride times were used to calculate asymmetry and arrhythmicity

respectively, during periods when the subject was not freezing. Asymmetry was defined as:

asymmetry $= 100 \times |\ln(SSWT/LSWT)|$, where SSWT and LSWT correspond to the leg with the

shortest and longest mean swing time over the trials, respectively and arrhythmicity was defined

as: arrhythmicity = the mean stride time coefficient of variation of both legs [23,26,30]. A large

stride time coefficient of variation is indicative of a less rhythmic gait. We developed a "forward

freeze index" inspired by the "Freeze Index" [13], and used antero-posterior accelerations

instead of vertical accelerations, making it similar to the "Frequency Ratio" [14]. We used a

window of 2s rather than 4s because 2s was closer to the mean stride time, and therefore

consistent with our other stride-by-stride metrics. The forward freeze index was calculated as the

square of the total power in the freeze band (3-8 Hz) over a 2s window, divided by the square of

the total power in the locomotor band (0.5-3 Hz) over the same 2s window. External videos of all

tasks were acquired on an encrypted clinical iPad (Apple Inc., Sunnyvale, CA) and synchronized

with the APDM data capture system through the Videography application (Appologics Inc.,

Germany).

A logistic regression model of freezing of gait

We developed a logistic regression model to calculate the probability that a given stride

was part of a freezing episode. The model was trained using 8 gait parameters (peak shank

angular velocity, stride time, swing angular range, arrhythmicity, asymmetry, forward freeze

index, peak shank angular velocity of the previous step, stride time of the previous step) and

ground truth binary labels (FOG = 1, no FOG = 0), from an experienced neurologist's (HBS)

video-determined ratings of freezing behavior, defined as periods where patient's normal gait

pattern changed (usually prior to a freezing episode) and where such behavior ended. VCode

software (Hagedorn, Hailpern, & Karahalios, 2008), was used to mark periods of freezing

behavior in each video with an accuracy of 10ms. Individual strides were identified using the

shank angular velocity trace as described above, and gait parameters were extracted for each

stride. The following gait parameters were calculated for each leg independently: peak shank

angular velocity, stride time, swing time, and swing angular range. The stride time and peak

shank angular velocity were normalized to averages from the subject's FW trial so that subjects

could be combined and compared to one another in the model. A step is likely to be a freeze if

the step before it has characteristics of a freeze, so the peak shank angular velocity for the

previous stride was included as a model input [15]. The swing and stride times for both legs were

concatenated to calculate arrhythmicity and asymmetry over the past 6 strides.

Analysis of gait parameters was performed in MATLAB (version 9.2, The MathWorks

Inc. Natick, MA, USA), and the logistic regression model was constructed using R (R Core

Team (2017)). We used a logistic regression model with a sparse set of features determined by

L1 regularization (LASSO) to predict whether a step was freezing or not. To evaluate model

performance, we used leave-one-out cross validation (LOOCV), which we refer to as external

LOOCV, where we left out a single subject as the test set. We then used the remaining subjects

as a training set, and used internal LOOCV, leaving out another subject as an internal test set

with which we used L1 regularization (LASSO) to determine a sparse set of features for the

model. Regularization minimizes the coefficients of different gait parameters, and the severity to

which it does this is determined by the regularization parameter. We found the best

regularization parameter (0.01) from the internal training set. This was repeated so that all

subjects were left out. We found that the variables selected by the internal LOOCV were

consistent across all runs, giving the combination of variables that best identified FOG. In both

LOOCVs, we kept subjects, who had multiple visits' worth of data together. For example, if

Subject X had two different visits, then data from both visits were either in the training set *or* in

the test set.

Investigating effects of DBS frequency in a subset of the PD cohort

A subset of the cohort, twelve PD subjects (7 freezers and 5 non-freezers), had been

treated with at least 21 months of optimized, continuous high frequency subthalamic DBS using

an implanted, investigative, concurrent sensing, and stimulating, neurostimulator (Activa® PC +

S, FDA-IDE approved; model 3389 leads, Medtronic, Inc.). Kinematic recordings were obtained,

off medication, during randomized presentations of no, 60 Hz, and 140 Hz subthalamic DBS

while subjects performed the TBC. The voltage was the same at both frequencies for each

subject's subthalamic nucleus. At least five minutes was allotted between experiments to allow

the subjects to rest.

Statistics

A two-way repeated-measures multivariate analysis of variance (MANOVA) test was

conducted to assess the effect of Group (Control, Non-Freezer, Freezer) or Task (Forward

Walking, TBC Ellipse, TBC Figure Eight), on average peak shank angular velocity, stride time,

asymmetry, and arrhythmicity for the three groups during non-freezing walking while OFF DBS.

If a main effect was found in the MANOVA, follow up univariate ANOVAs were used to

evaluate significant parameters. Post-hoc pairwise effects were examined using a Bonferroni

correction. A three-way repeated measures ANOVA was used to compare the effect of DBS

frequency (OFF, 60 Hz, 140 Hz), Group (Non-Freezer, Freezer), or Task (TBC Ellipse, TBC

Figure Eight) during non-freezing walking in the TBC. Post hoc analyses were conducted to

compare between stimulation conditions. A students t-test was used to compare freezers' percent

time spent freezing in the TBC ellipses versus figures of eight. Students t-tests were used for the

comparison of demographics between the freezer, non-freezer and control groups. Paired t-tests

were used to compare UPDRS III scores between visits for subjects with repeated visits. The

relationship between percent time freezing and FOG-Q3 response was investigated using a

Spearman correlation analysis. The relationship between percent time freezing and average peak

shank angular velocity, stride time, asymmetry, and arrhythmicity during non-freezing walking

was investigated using a Pearson correlation analysis to compare freezers' non-freezing walking

with the severity of their freezing behavior. All statistical testing was performed in SPSS Version

21, or SigmaPlot (Systat Software, San Jose, CA) using two-tailed tests with significance levels

of $p < .05$.

Results

Human subjects

Among the 23 PD subjects, there were 8 freezers, 13 non-freezers, and 2 subjects who

converted from the definition of a non-freezer to a freezer between two visits. Non-freezers and

controls were of similar ages, while freezers were younger (65.9 ± 7.5, 66.9 ± 8.9 years, $57.9 \pm$

6.14, respectively, $p < 0.05$). Disease duration was similar between the freezer and non-freezer

groups (9.3 ± 2.8, 8.9 ± 4.2 years, respectively). Freezers had a higher off medication UPDRS III

score than non-freezers (39.8 ± 9.2, 24.1 ± 13.6 respectively, $p < 0.01$), and all PD patients had

higher UPDRS III scores than controls ($p < 0.001$). All subjects completed all walking tasks,

except two freezers who could not complete the TBC, and one non-freezer whose sensor data

was unusable; these three subjects were excluded from the analysis. Three healthy control

subjects were excluded due to arthritis (N=2) or essential tremor (N=1), which affected their

walking. The average total durations of FW and the TBC were 33.1 ± 8.7 and 157.4 ± 88.9

seconds, respectively.

Nine subjects had repeat visits. The length between repeated visits was 430 ± 112 days

(mean $\pm$ SD) and the repeated visit group's mean UPDRS III score trended higher at the second

visit (32.4 ± 12.0, 35.7 ± 14.8, respectively, $p = 0.05$), so the repeated patient visits were treated

independently. Data from 40 visits (9 from controls, 13 from freezers, and 18 from non-freezers)

were used to examine how the three different cohorts completed the gait tasks while OFF

stimulation. In assessing the effects of lower and high frequency subthalamic DBS on subjects in

the TBC, there were no repeat visits.

The TBC reliably elicits more FOG than forward walking and is

validated with the FOG-Q3

During the TBC, all freezers experienced a freezing episode. In total, 217 freezing

episodes were identified. Freezers spent more time freezing in the TBC figures of eight than the

TBC ellipses (38.23 ± 29.0 %, $23.6 \, 0 \pm 19.3$ %, respectively, $p < 0.01$). During FW only one

freezer experienced a freezing episode. Freezers spent an average of 33.0 ± 24.2 % of the time

freezing in the TBC compared to the one freezer who spent 2% of the time freezing during

forward walking and was a moderate to severe freezer who spent 59% of the TBC task freezing

(as determined by the blinded neurologist). There was a strong correlation between the time

spent freezing in the TBC and a subjects' report of freezing severity from the FOG-Q3 (r = 0.74,

$p < 0.001$), which validates the TBC as a tool for measuring FOG in Parkinson's disease. There

was no significant correlation between the time spent freezing during FW and a subjects' report

of freezing severity from the FOG-Q3 (r = 0.28, $p = 0.075$).

Arrhythmicity during non-freezing gait differentiates freezers from non-freezers

MANOVA results indicated a main effect of Group (freezer, non-freezer, control, $p <$

0.001) and Task (FW, TBC Ellipse, TBC Figures of Eight, $p < 0.001$), demonstrating that the

three groups were distinguishable regardless of task, and the tasks were distinguishable

regardless of group. All four of the gait parameters showed significant univariate effects of

Group, and all gait parameters except asymmetry showed significant univariate effects of Task.

There was an interaction effect of Task*Group ($p = 0.011$), with a univariate effect only in

arrhythmicity. Post-hoc pairwise comparisons showed that freezers' non-freezing gait was more

arrhythmic than that of non-freezers or controls during all tasks ($p < 0.05$ for all), Fig 3A.

Fig 3. Gait parameters during non-freezing walking over different tasks and groups.

(A) Arrhythmicity differentiates freezers from non-freezers and controls. (B) Average peak

shank angular velocity (C) Asymmetry (D) Stride time. Error bars represent standard deviation.

* denotes $p < 0.05$, ** denotes $p < 0.01$, *** denotes $p < 0.001$, ^ denotes significant difference

between FW in the same group ($p < 0.05$), ~ denotes significant difference between ellipses in

the same group ($p < 0.05$).

Post-hoc pairwise comparisons showed that freezers' non-freezing gait during both the ellipses

and figures of eight of the TBC demonstrated greater arrhythmicity compared to their non-

freezing gait during FW ($p = 0.001$, $p < 0.001$, respectively), and their arrhythmicity was greater

in the figures of eight than in the ellipses ($p < 0.001$), Fig 3A. No pairwise effect was detected

for non-freezers' or controls' gait arrhythmicity between TBC and FW, though non-freezers

were significantly more arrhythmic while walking in the TBC figures of eight versus walking in

the TBC ellipses ($p = 0.02$). There was no Task*Group interaction observed for shank angular

velocity, stride time or asymmetry, though the observed power for these variables was low.

Gait features in logistic regression model detect freezing on a step-by-step basis

A logistic regression model with only four parameters had an AUC of 0.754, third row Fig 4A. In this model, the gait parameter with the largest coefficient and thereby the strongest predictor of whether a step was part of a freeze, was arrhythmicity over the last six steps (coefficient of 2.034), followed by stride time (coefficient of 0.0931), swing angular range (coefficient of -0.0615), and finally asymmetry over the last six steps (coefficient of 0.0003), with a model intercept of 0.941. The logistic regression models with single parameters all had coefficients significantly different from zero ($p < 0.001$) but most were only moderately better than chance (AUC = 0.5), first row Fig 4A. A logistic regression model with all gait parameters, second row in Fig 4A, outperformed any single-parameter model but had an AUC (0.750) less than that of the four-parameter-model.

Fig 4. Logistic regression model performance for different gait parameters.

(A) Overall model performance: Area under the receiver operating characteristic curve (AUC)

values for different model iterations using leave-one-out cross validation on the freezer group.

First row: AUC values for models using individual gait parameters; second row: AUC value for

model using all gait parameters; third row: AUC value for model with sparse parameter set

chosen from regularization. (Peak Shank AV = Peak Shank Angular Velocity; some metrics are

calculated over a window of steps in time, so "t-3:t" represents a window from "t-3" or 3 steps

earlier, to and including the current step "t"). (B) Sparse parameter model-identified freezing of

gait (pink shading) was often shorter and contained within neurologist-identified freezing

behavior (orange shading) as seen in shank sagittal angular velocity traces and freezing labels for

a representative patient.

Since the AUC is a threshold-independent assessment of the model, we calculated the accuracy

of the model at a threshold of 0.50 (e.g. if the probability that the step was a freeze was over 50%

then it was determined to be a freeze). At this threshold, the accuracy of the model to correctly

identify a step as freezing or not freezing ((true positives + true negatives)/total number of steps),

was 90%. We found that the model often detected a freezing event within the interval defined as

freezing behavior by the neurologist, Fig 4B. In this case, the model overlapped with the

neurologist-identified freezing behavior, though it did not detect some of preceding or

succeeding freezing behavior identified by the neurologist. We defined such a case as correct

model-identification of a freezing event, and overall, the model correctly identified 77% of the

neurologist-identified freezing behavior events, overlapping with neurologist markings within a

2-stride window.

The time spent freezing in the TBC for all subjects, identified by the logistic regression

model, correlated with the subject's score on FOG-Q3 ($r = 0.68$, $p < 0.001$). The percent time

freezing predicted by the model for the control subjects and non-freezers was less than 1% for

each subject, except for one subject who had one step erroneously classified as freezing resulting

in 2.5% time spent freezing in the TBC.

Percent time spent freezing correlated with freezers' gait

parameters during non-freezing gait in the TBC

Freezers' gait arrhythmicity, during non-freezing gait in both ellipses and figures of eight

but not during FW, strongly correlated with their percent time freezing in the TBC, as

determined by the model ($r = 0.94$, $r = 0.92$ respectively, $p < 0.001$ for both), Fig 5.

Fig 5. Relationship between freezers' non-freezing gait arrhythmicity in different walking

tasks and freezing severity in the turning and barrier course (TBC). Freezers' arrhythmicity

during non-freezing walking in the TBC ellipses and figure eights, but not forward walking

(FW), correlates with percent time spent freezing (FOG) in the TBC. Regression line (black line)

and confidence interval of the correlation coefficient at 95% (shaded grey), and subjects (colored

dots).

Freezers' peak shank angular velocity during non-freezing gait in figures of eight, but not in

TBC ellipses or FW, also correlated with their percent time spent freezing in the TBC (r = -0.71,

$p < 0.01$). There was no correlation between gait asymmetry or stride time during non-freezing

walking in TBC, or between any gait parameter during FW, with percent time freezing in the

TBC. These results demonstrated that increased gait arrhythmicity and decreased peak shank

angular velocity of non-freezing gait during the TBC were strong markers of FOG severity in PD

freezers.

Sixty Hz and 140 Hz subthalamic DBS improved non-freezing gait

impairment and FOG in freezers during the TBC

Gait impairment and FOG improved during both 60 Hz and 140 Hz subthalamic DBS:

the percent time spent freezing in the TBC was lower during either 60 Hz or 140 Hz DBS

compared to when OFF DBS in freezers ($5 \pm 7\%$, $9 \pm 10\%$, $35 \pm 23\%$, respectively, $p < 0.05$)

and was not different from that of non-freezers (whose percent time spent freezing was zero). S1

Video highlights the decrease in percent time freezing seen during 60 Hz or 140 Hz DBS versus

OFF DBS in a representative patient walking in one ellipse and passing through the narrow

passageway in the TBC. This patient went from 53% task freezing while OFF DBS, to 6% task

freezing on both 60 Hz and 140 Hz DBS. There was a statistically significant effect of DBS

frequency (OFF, 60 Hz, 140 Hz) on shank angular velocity and arrhythmicity ($p < 0.01$, $p <$

0.05), as well as a statistically significant effect of Task (TBC Ellipse, TBC Figure Eight) on

shank angular velocity ($p < 0.001$) as determined by three-way repeated measures ANOVAs.

Freezers' gait arrhythmicity during the TBC decreased to values not statistically different from

those of non-freezers during both 60 Hz and 140 Hz DBS ($p > 0.05$), Fig 6A, despite freezers'

arrhythmicity being significantly higher than that of non-freezers OFF DBS ($p < 0.01$). Freezers'

shank angular velocity increased during either frequency of DBS ($p < 0.01$), Fig 6B, despite

being significantly less than that of non-freezers OFF DBS ($p = 0.036$).

Fig 6. Gait parameters during different presentations of deep brain stimulation (DBS).

Changes in (A) arrhythmicity and (B) average peak shank angular velocity in freezers (red) and

non-freezers (blue) during walking in the TBC while OFF and on 60 Hz and 140 Hz DBS.

During DBS, freezers' arrhythmicity and average peak shank angular velocity improve to values

characteristic of non-freezers. Healthy control averages shown (green line) with standard

deviations (shaded green). Error bars represent standard deviation. * denotes $p < 0.05$, ** denotes

$p < 0.01$.

When OFF DBS, we did not detect a difference in non-freezers' arrhythmicity, asymmetry or

stride time from that of controls ($p > 0.05$ for all), though non-freezers' shank angular velocity

was significantly less than that of controls ($p = 0.031$). DBS had no detectable effect on any of

the non-freezers' gait parameters, Fig 6. Freezers had significantly lower shank angular velocity,

higher arrhythmicity and asymmetry than controls OFF DBS ($p < 0.05$ for all), but a similar

402 stride time to controls OFF DBS ($p > 0.05$). There were no statistically significant differences

403 between freezers' or non-freezers' group means between the DBS versus OFF DBS conditions

404 for stride time or asymmetry as determined by three-way repeated measures ANOVAs,

405 demonstrating no detectable effect of DBS on these gait parameters.

Discussion

This study has validated the objective measurement of FOG from an instrumented gait

task, the turning and barrier course (TBC), with the international standard FOG questionnaire

(FOG-Q). The TBC mimicked real-life scenarios that trigger FOG in PD and was superior at

eliciting more arrhythmic non-freezing gait, and freezing episodes in freezers compared to 40

meters of forward walking. Freezers' non-freezing gait was more arrhythmic than that of non-

freezers or controls irrespective of task.

A logistic regression model demonstrated that a combination of stride time, swing

angular range, arrhythmicity, and asymmetry of the past six steps best predicted FOG during the

TBC (AUC = 0.754). Freezers' gait arrhythmicity was not only the strongest feature for

predicting FOG, but also the non-freezing gait parameter most highly correlated with freezing

severity (the percent time freezing).

Freezers' percent time freezing decreased during either 60 Hz or 140 Hz STN DBS and

their non-freezing gait arrhythmicity and shank angular velocity was restored to similar values as

those of non-freezers.

The TBC is a robust, standardized task for assessing impaired gait in PD

It has been difficult to develop an objective measure of FOG since it is challenging to elicit FOG in the clinic or laboratory where there are few obstacles, tight corners, or narrow door openings [31]. Tasks that have been shown to provoke FOG include rapid clockwise and counterclockwise 360 degree turns in place [32], in combination with walking through doorways [33], walking with dual tasking [14,34–36], and recently forward walking tasks including straight walking or turning around cones [37]. We have previously validated freezing behavior during a stepping in place task on dual force plates with the FOG-Q3 [23]. In designing the TBC, we desired a forward walking task that included standardized situational triggers for FOG that were representative of real-world scenarios [25]. The TBC elicited more arrhythmic gait and FOG events in freezers than FW, indicating that the TBC is able to exacerbate gait impairment specifically in this group. Within all groups, we expected to see increased arrhythmicity while walking in the TBC compared to straight walking in the FW task; however, we only saw increased arrhythmicity in the gait of the PD freezers when comparing their walking in the TBC to their walking in FW. Non-freezing gait arrhythmicity in the TBC differentiated freezers from non-freezers and controls, and correlated with the percent time spent freezing, which was not seen during FW. This result aligns with previous studies that have shown that freezers exhibit greater arrhythmicity than non-freezers during non-freezing walking or stepping [24,25,30,38,39], though this is the first study to demonstrate this during non-freezing walking and turning to the best of our knowledge. This confirms that the arrhythmicity of non-freezing gait elicited during the TBC is a useful metric to predict the severity of FOG freezers may

445 experience in the real world, and is a robust measure of freezing behavior even during non-

446 freezing gait.

A logistic regression model identified freezing events using gait parameters from the TBC

A logistic regression model identified gait arrhythmicity, swing angular range, stride

time, and asymmetry as the most important gait parameters for classifying freezing events during

the TBC. The model had an AUC of 0.754 and identified the freezing events within the

neurologist identified periods of freezing behavior with 77% accuracy. It was interesting that

both the neurologist and the model behaved as they were 'trained.' The model's definition of a

freezing event was within the neurologist's period, Fig 4B, as the latter identified gait behavior

leading up to and after an actual freezing episode, which encompassed complete halts in walking

often seen in freezing of gait, but also included gait shuffling, festination, trembling, and shorter

strides that often precede and succeed the complete gait arrest. This highlights another variable

in the definition of FOG, some definitions only include 'motor blocks' or events when forward

motion stops, whereas others include abnormal freezing behavior in the definition of FOG.

These variable definitions may have contributed to the variation in the accuracy of other

IMU-based FOG detection algorithms, which have reported sensitivities and specificities ranging

from 73-99% [13–16,18–20,22,40]. Some of these algorithms detected FOG based on high

frequency components of leg linear acceleration corresponding to leg trembling-FOG, with lower

sensitivity to non-trembling FOG, despite high specificity. The "forward freeze index", which

measures the relative component of high to low frequency gait components, has been shown to

be a useful predictor of FOG in a 360-degree turning task [14]; however this had a lower AUC

value in our model compared to other gait parameters, Fig 4A. Explanations for this may include

that the TBC task did not include 360 degree turning, which may specifically induce more leg

trembling high frequency components of freezing behavior. This supports the clinical experience

that FOG manifests with different types of gait impairment depending on what gait task the PD

person is trying to accomplish.

474   FOG and gait impairment in freezers improved during STN DBS

475   We demonstrated that both FOG and predictors of FOG during non-freezing gait

improved during 60 Hz and 140 Hz STN DBS while subjects walked in the TBC that mimicked

real-life environments that elicit FOG. During the TBC, freezers spent less time freezing when

on either frequency of DBS compared to OFF DBS, which is similar to our reports of the effect

of DBS on the stepping in place and forward walking tasks [24]. Freezers' gait arrhythmicity

also improved on both 60 Hz and 140 Hz DBS, to levels that were not different from that of non-

freezers'. Three out of four of non-freezers' gait parameters OFF DBS were not different from

those exhibited by healthy controls and all were left unchanged on either frequency of DBS. This

'if it isn't broken, it doesn't need fixing' effect of DBS has been observed in gait [24,41] and in

aspects of postural instability [41–43].

485   Sixty Hz DBS has been shown to be effective in improving axial symptoms in patients

with FOG [10,11], though it is not obvious whether 60 Hz versus 140 Hz is better for FOG in

real-world walking tasks. Using the clinical assessment of FOG from the MDS-UPDRS III,

Ramdhani et al. reported that lower frequency (60 Hz) DBS reduced FOG when high frequency

(130 Hz) DBS did not, even shortly after DBS was initiated [44]. Our previous investigations of

the effect of 60 Hz and 140 Hz DBS on repetitive stepping in place and on progressive

bradykinesia demonstrated that 60 Hz DBS promoted more regularity in ongoing movement,

[24,45]

In this study percent time freezing and gait arrhythmicity improved during either 60 Hz

or 140 Hz STN DBS, and to a similar degree. This aligns with a previous report that gait and

postural performances with low and high frequency stimulations were largely similar [41], and

another demonstrating that 140 Hz STN DBS increased stride length and foot clearing [46],

underscoring the increased shank angular velocities demonstrated during STN DBS in this study.

Altogether this is valuable assurance for people with PD and clinicians that STN DBS can

improve gait and FOG, and that both 60 Hz and 140 Hz improve FOG in real-world walking

tasks.

502 **Limitations**

Our logistic regression model utilized data from only one IMU from a small cohort of

freezers. Although this resulted in interpretable gait features and an accuracy within that of

several other FOG models, it could be improved. Multiple IMUs on different parts of the body

may add sensitivity. The model, being a binary classifier, attempted to capture all of the

variability in freezing behavior with just two labels: "FOG" or "not FOG". A different model

might use multiple classes, where the classifier discriminates between unimpaired walking, a

completely halted gait freeze event, shuffling, and a start hesitation. In addition, only freezers

were used to train and test the logistic regression model, so that the incidence of freezing events

was sufficient. Future models might include bootstrapping methods, evaluate the data from

multiple IMUs, or more data to increase the sizes of the training and test sets. Another limitation

is that natural walking speeds vary among PD and healthy subjects. We attempted to overcome

this variability by normalizing each individual's stride times and shank angular velocities in the

TBC to their averages from FW. This normalization procedure allowed comparison among

individuals. In future model versions using absolute measures could be tested.

518 Conclusions

Tools and tasks such as the instrumented TBC are necessary for designing and assessing

personalized interventions and therapies for gait impairment and FOG in PD. We have

demonstrated the utility of the instrumented TBC for eliciting FOG, for revealing gait parameters

that identify freezers and predict FOG during non-freezing gait, and for measuring the efficacy

of different frequencies of STN DBS. From the TBC experimental data and a logistic regression

model, we have identified the gait parameters that are most likely to predict freezing events and

which may be useful in closed loop DBS for gait impairment and FOG.

527 Acknowledgements

528   We thank Matthew Petrucci, Tom Prieto, Jordan Parker, Varsha Prabhakar, Raumin

529 Neuville, Ross Anderson, and Amaris Martinez for their support during the experiments and

530 helpful comments. We would also like to thank our dedicated patient population who contributed

531 their time to participating in our study (ClinicalTrials.gov Identifier: NCT02304848).

534 References

1.   Giladi N, McMahon D, Przedborski S, Flaster E, Guillory S, Kostic V, et al. Motor blocks

in Parkinson's disease. Neurology. 1992 Feb;42(2):333–9.

2.   Morgan D, Funk M, Crossley M, Basran J, Kirk A, Dal Bello-Haas V. The potential of

gait analysis to contribute to differential diagnosis of early stage dementia: current

research and future directions. Can J Aging. 2007;

3.   Lippa CF, Duda JE, Grossman M, Hurtig HI, Aarsland D, Boeve BF, et al. DLB and PDD

boundary issues: Diagnosis, treatment, molecular pathology, and biomarkers. Neurology.

2007;

4.   Brozova H, Stochl J, Roth J, Ruzicka E. Fear of falling has greater influence than other

aspects of gait disorders on quality of life in patients with Parkinson's disease. Neuro

Endocrinol Lett [Internet]. 2009;30(4):453–7. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/20010494

5.   Bloem BR, Hausdorff JM, Visser JE, Giladi N. Falls and freezing of gait in Parkinson's

disease: A review of two interconnected, episodic phenomena. Mov Disord [Internet].

2004 Aug;19(8):871–84. Available from: http://doi.wiley.com/10.1002/mds.20115

6.   Amboni M, Stocchi F, Abbruzzese G, Morgante L, Onofrj M, Ruggieri S, et al.

Prevalence and associated features of self-reported freezing of gait in Parkinson disease:

The DEEP FOG study. Park Relat Disord [Internet]. 2015;21(6):644–9. Available from:

http://dx.doi.org/10.1016/j.parkreldis.2015.03.028

7.   Sieber B-AA, Landis S, Koroshetz W, Bateman R, Siderowf A, Galpern WR, et al.

Prioritized research recommendations from the National Institute of Neurological

Disorders and Stroke Parkinson's Disease 2014 Conference. Ann Neurol [Internet]. 2014

Oct;76(4):469–72. Available from: http://doi.wiley.com/10.1002/ana.24261

8.   Merola A, Zibetti M, Angrisano S, Rizzi L, Ricchi V, Artusi CA, et al. Parkinson's

disease progression at 30 years: a study of subthalamic deep brain-stimulated patients.

Brain [Internet]. 2011 Jul 1;134(7):2074–84. Available from:

https://academic.oup.com/brain/article-lookup/doi/10.1093/brain/awr121

9.   Schlenstedt C, Shalash A, Muthuraman M, Falk D, Witt K, Deuschl G. Effect of high-

563   frequency subthalamic neurostimulation on gait and freezing of gait in Parkinson's

564   disease: a systematic review and meta-analysis. Eur J Neurol. 2017;24(1):18–26.

565   10.   Xie T, Vigil J, MacCracken E, Gasparaitis A, Young J, Kang W, et al. Low-frequency

566   stimulation of STN-DBS reduces aspiration and freezing of gait in patients with PD.

567   Neurology [Internet]. 2015 Jan 27;84(4):415–20. Available from:

568   http://www.neurology.org/cgi/doi/10.1212/WNL.0000000000001184

569   11.   Xie T, Padmanaban M, Bloom L, MacCracken E, Bertacchi B, Dachman A, et al. Effect

570   of low versus high frequency stimulation on freezing of gait and other axial symptoms in

571   Parkinson patients with bilateral STN DBS: a mini-review. Transl Neurodegener

572   [Internet]. 2017;6:13. Available from:

573   http://ovidsp.ovid.com/ovidweb.cgi?T=JS&PAGE=reference&D=prem&NEWS=N&AN=

574   28529730

575   12.   Moreau C, Defebvre L, Destee A, Bleuse S, Clement F, Blatt JL, et al. STN-DBS

576   frequency effects on freezing of gait in advanced Parkinson disease. Neurology [Internet].

577   2008 Jul 8;71(2):80–4. Available from:

578   http://www.neurology.org/cgi/doi/10.1212/01.wnl.0000303972.16279.46

579   13.   Moore ST, MacDougall HG, Ondo WG. Ambulatory monitoring of freezing of gait in

580   Parkinson's disease. J Neurosci Methods. 2008;167(2):340–8.

14.   Mancini M, Priest KC, Nutt JG, Horak FB. Quantifying freezing of gait in Parkinson's

disease during the instrumented timed up and go test. 2012 Annu Int Conf IEEE Eng Med

Biol Soc [Internet]. 2012;1198–201. Available from:

http://ieeexplore.ieee.org/document/6346151/

15. Palmerini L, Rocchi L, Mazilu S, Gazit E, Hausdorff JM, Chiari L. Identification of

characteristic motor patterns preceding freezing of gait in Parkinson's disease using

wearable sensors. Front Neurol. 2017;8(AUG):1–12.

16. Silva de Lima AL, Evers LJW, Hahn T, Bataille L, Hamilton JL, Little MA, et al.

Freezing of gait and fall detection in Parkinson's disease using wearable sensors: a

systematic review. J Neurol [Internet]. 2017;264(8):1642–54. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/28251357%0Ahttp://www.pubmedcentral.nih.gov/a

rticlerender.fcgi?artid=PMC5533840

17. Kwon Y, Park SH, Kim J-W, Ho Y, Jeon H-M, Bang M-J, et al. A practical method for

the detection of freezing of gait in patients with Parkinson's disease. Clin Interv Aging

[Internet]. 2014;9:1709–19. Available from:

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4199977&tool=pmcentrez&re

ndertype=abstract

18. Rezvanian S, Lockhart T. Towards Real-Time Detection of Freezing of Gait Using

Wavelet Transform on Wireless Accelerometer Data. Sensors [Internet]. 2016 Apr

2;16(4):475. Available from: http://www.mdpi.com/1424-8220/16/4/475

19. Coste C, Sijobert B, Pissard-Gibollet R, Pasquier M, Espiau B, Geny C. Detection of

Freezing of Gait in Parkinson Disease: Preliminary Results. Sensors [Internet]. 2014 Apr

15;14(4):6819–27. Available from: http://www.mdpi.com/1424-8220/14/4/6819/

20. Zach H, Janssen AM, Snijders AH, Delval A, Ferraye MU, Auff E, et al. Identifying

freezing of gait in Parkinson's disease during freezing provoking tasks using waist-

mounted accelerometry. Parkinsonism Relat Disord. 2015 Nov;21(11):1362–6.

21.   Khemani P, Dewey RB, BH Q, MP B, A B, RG C, et al. Deep Brain Stimulation of the

Subthalamic Nucleus. JAMA Neurol [Internet]. 2015;72(5):499. Available from:

http://archneur.jamanetwork.com/article.aspx?doi=10.1001/jamaneurol.2015.36

22.   Kim H, Lee HJ, Lee W, Kwon S, Kim SK, Jeon HS, et al. Unconstrained detection of

freezing of Gait in Parkinson's disease patients using smartphone. In: 2015 37th Annual

International Conference of the IEEE Engineering in Medicine and Biology Society

(EMBC). 2015. p. 3751–4.

23.   Nantel J, de Solages C, Bronte-Stewart H. Repetitive stepping in place identifies and

measures freezing episodes in subjects with Parkinson's disease. Gait Posture [Internet].

2011;34(3):329–33. Available from: http://dx.doi.org/10.1016/j.gaitpost.2011.05.020

24.   Anidi C, O'Day JJ, Anderson RW, Afzal MF, Syrkin-Nikolau J, Velisar A, et al.

Neuromodulation targets pathological not physiological beta bursts during gait in

Parkinson's disease. Neurobiol Dis. 2018;120.

25.   Syrkin-Nikolau J, Koop MM, Prieto T, Anidi C, Afzal MF, Velisar A, et al. Subthalamic

neural entropy is a feature of freezing of gait in freely moving people with Parkinson's

disease. Neurobiol Dis [Internet]. 2017;108(June):288–97. Available from:

http://dx.doi.org/10.1016/j.nbd.2017.09.002

26.   Plotnik M, Hausdorff JM. The role of gait rhythmicity and bilateral coordination of

stepping in the pathophysiology of freezing of gait in Parkinson's disease. Mov Disord.

2008;23(SUPPL. 2):444–50.

27.   Hausdorff JM. Gait dynamics in Parkinson's disease: common and distinct behavior

among stride length, gait variability, and fractal-like scaling. Chaos. 2009

Jun;19(2):26113.

28.   Goetz CG, Tilley BC, Shaftman SR, Stebbins GT, Fahn S, Martinez-Martin P, et al.

Movement Disorder Society-Sponsored Revision of the Unified Parkinson's Disease

Rating Scale (MDS-UPDRS): Scale presentation and clinimetric testing results. Mov

Disord. 2008;23(15):2129–70.

29.   Giladi N, Tal J, Azulay T, Rascol O, Brooks DJ, Melamed E, et al. Validation of the

Freezing of Gait Questionnaire in patients with Parkinson's disease. Mov Disord.

2009;24(5):655–61.

30.   Plotnik M, Giladi N, Balash Y, Peretz C, Hausdorff JM. Is freezing of gait in Parkinson's

disease related to asymmetric motor function? Ann Neurol. 2005;57(5):656–63.

31.   Nieuwboer A, Giladi N. The challenge of evaluating freezing of gait in patients with

Parkinson's disease. Br J Neurosurg [Internet]. 2008 Jan 1;22(sup1):S16–8. Available

from: https://doi.org/10.1080/02688690802448376

32.   Mancini M, Smulders K, Cohen RG, Horak FB, Giladi N, Nutt JG. The clinical

significance of freezing while turning in Parkinson's disease. Neuroscience [Internet].

2017;343:222–8. Available from: http://dx.doi.org/10.1016/j.neuroscience.2016.11.045

33.   Ziegler K, Schroeteler F, Ceballos-Baumann AO, Fietzek UM. A new rating instrument to

assess festination and freezing gait in Parkinsonian patients. Mov Disord.

2010;25(8):1012–8.

34.   Snijders AH, Haaxma CA, Hagen YJ, Munneke M, Bloem BR. Freezer or non-freezer:

clinical assessment of freezing of gait. Parkinsonism Relat Disord. 2012 Feb;18(2):149–

54.

35.   Spildooren J, Vercruysse S, Desloovere K, Wim Vandenberghe MD P, Kerckhofs PhD E,

Nieuwboer A. Freezing of Gait in Parkinson's Disease: The Impact of Dual-Tasking and

Turning. Mov Disord. 2010 Nov 15;25:2563–70.

36.   Rochester L, Nieuwboer A, Baker K, Hetherington V, Willems AM, Kwakkel G, et al.

Walking speed during single and dual tasks in Parkinson's disease: Which characteristics

are important? Mov Disord. 2008;23(16):2312–8.

37.   Mitchell T, Conradsson D, Paquette C. Gait and trunk kinematics during prolonged

turning in Parkinson's disease with freezing of gait. Park Relat Disord [Internet].

2019;(April):0–1. Available from: https://doi.org/10.1016/j.parkreldis.2019.04.011

38.   Plotnik M, Hausdorff JM. The role of gait rhythmicity and bilateral coordination of

stepping in the pathophysiology of freezing of gait in Parkinson's disease. Mov Disord.

2008;23 Suppl 2:S444-50.

39.   Hausdorff JM, Schaafsma JD, Balash Y, Bartels AL, Gurevich T, Giladi N. Impaired

regulation of stride variability in Parkinson's disease subjects with freezing of gait. Exp

Brain Res. 2003;149(2):187–94.

40.   Moore ST, Yungher DA, Morris TR, Dilda V, MacDougall HG, Shine JM, et al.

Autonomous identification of freezing of gait in Parkinson's disease from lower-body

segmental accelerometry. J Neuroeng Rehabil [Internet]. 2013;10(1):19. Available from:

http://jneuroengrehab.biomedcentral.com/articles/10.1186/1743-0003-10-19

41.   Vallabhajosula S, Haq IU, Hwynn N, Oyama G, Okun M, Tillman MD, et al. Low-

frequency versus high-frequency subthalamic nucleus deep brain stimulation on postural

control and gait in Parkinson's disease: a quantitative study. Brain Stimul. 2015;8(1):64–

75.

42.   Bronte-Stewart HM. Postural instability in idiopathic Parkinson's disease: the role of

medication and unilateral pallidotomy. Brain [Internet]. 2002;125(9):2100–14. Available

from: http://brain.oxfordjournals.org/content/125/9/2100.abstract

43.   Shivitz N, Koop MM, Fahimi J, Heit G, Bronte-Stewart HM. Bilateral subthalamic

nucleus deep brain stimulation improves certain aspects of postural control in Parkinson's

disease, whereas medication does not. Mov Disord. 2006;21(8):1088–97.

44.   Ramdhani RA, Patel A, Swope D, Kopell BH. Early Use of 60 Hz Frequency Subthalamic

Stimulation in Parkinson's Disease: A Case Series and Review. Neuromodulation.

2015;18(8):664–9.

45.   Blumenfeld Z, Koop MM, Prieto TE, Shreve LA, Velisar A, Quinn EJ, et al. Sixty-hertz

stimulation improves bradykinesia and amplifies subthalamic low-frequency oscillations.

Mov Disord. 2017;32(1):80–8.

46.   Hell F, Plate A, Mehrkens JH, Bötzel K. Subthalamic oscillatory activity and connectivity

during gait in Parkinson's disease. NeuroImage Clin [Internet]. 2018;19(May):396–405.

Available from: https://doi.org/10.1016/j.nicl.2018.05.001