A young age of subspecific divergence in the desert locust, *Schistocerca gregaria*

Marie-Pierre Chapuis[1,2], Louis Raynal[3,4], Christophe Plantamp[5], Christine N. Meynard[6], Laurence

Blondin[7], Jean-Michel Marin[3,4] and Arnaud Estoup[4,6]

[1] CIRAD, CBGP, Montpellier, France.

[2] CBGP, CIRAD, INRA, IRD, Montpellier SupAgro, Univ Montpellier, Montpellier, France.

[3] IMAG, Univ de Montpellier, CNRS, Montpellier, France.

[4] Institut de Biologie Computationnelle (IBC), Montpellier, France.

[5] ANSES, Laboratoire de Lyon, France.

[6] CBGP, INRA, CIRAD, IRD, Montpellier SupAgro, Univ Montpellier, Montpellier, France.

[7] CIRAD, BGPI, Montpellier, France.

Corresponding author: Marie-Pierre Chapuis, marie-pierre.chapuis@cirad.fr

Keywords: Approximate Bayesian Computation; colonization; desert locust; divergence time; Holocene;

Orthoptera; paleo-vegetation; Random Forest; *Schistocerca gregaria*

Short title: Evolutionary history of *Schistocerca gregaria*

Abstract

Dating population divergence within species from molecular data and relating such dating to climatic and biogeographic changes is not trivial. Yet it can help formulating evolutionary hypotheses regarding local adaptation and future responses to changing environments. Key issues include statistical selection of a demographic and historical scenario among a set of possible scenarios, and estimation of the parameter(s) of interest under the chosen scenario. Such inferences greatly benefit from new statistical approaches including Approximate Bayesian Computation - Random Forest (ABC-RF), the later providing reliable inference at a low computational cost, with the possibility to take into account prior knowledge on both biogeographical history and genetic markers. Here, we used ABC-RF, including independent information on evolutionary rate and pattern at microsatellite markers, to decipher the evolutionary history of the African arid-adapted pest locust, *Schistocerca gregaria*. We found that the evolutionary processes that have shaped the present geographical distribution of the species in two disjoint northern and southern regions of Africa were recent, dating back 2.6 Ky (90% CI: 0.9 – 6.6 Ky). ABC-RF inferences also supported a southern colonization of Africa from a low number of founders of northern origin. The inferred divergence history is better explained by the peculiar biology of *S. gregaria*, which involves a density-dependent swarming phase with some exceptional spectacular migrations, rather than a continuous colonization resulting from the continental expansion of open vegetation habitats during the past Quaternary glacial episodes.

Introduction

As in other regions of the world, Africa has gone through several major episodes of climate change since the early Pleistocene (deMenocal 1995 and 2004). During glaciation periods, the prevalent climate was colder and drier than nowadays, and became more humid during warmer interglacial periods. These climatic phases resulted in shifts of vegetation (Vivo and Carmignotto 2004) and are most likely at the origin of the current isolation between northern and southern distributions of arid-adapted species (Monod 1971). In Africa, at least fifty-six plant species show disjoint geographical distributions in southern and northern arid areas (Monod 1971; Jurgens 1997; Lebrun 2001). Similarly, a number of animal vertebrate species show meridian disjoint distributions on this continent, including eight mammals and 29 birds (Monod 1971; de Vivo and Carmignotto 2004; Lorenzen *et al.* 2012). The desert locust, *Schistocerca gregaria*, is among the few examples of insect species distributed in two distinct regions along the north-south axis of Africa. Other known disjunctions in insects are interspecific and concern species of the families Charilaidae (Orthoptera) and Mythicomyiidae (Diptera), and of the genus *Fidelia* (Hymenoptera) (Le Gall *et al.* 2010). Similarities in extant distributions of African arid-adapted species across divergent taxonomic groups point to a common climatic history and an important role of environmental factors. Yet, to our knowledge, studies relating evolutionary history and climatic history have rarely been carried out in this continent (but see mitochondrial studies by Miller *et al.* 2011 on the ostrich, Atickem *et al.* 2018 on the black-backed jackal, and Moodley *et al.* 2018 on the white rhinoceros).

Relating evolutionary and climatic histories often requires dating population differentiation events so that species or subspecies divergence can be understood in a broad biogeographic context. However, finding a reliable calibration to convert measures of genetic divergence into units of absolute time is challenging, especially so for recent evolutionary events (Ho *et al.* 2008). Extra-specific fossil calibration may lead to considerable overestimates of divergence times, but internal fossil records are often lacking (Ho *et al.* 2008). A sensible approach is to use an evolutionary rate estimated from sequence data of a related species for which internal calibration is available (Ho *et*

al. 2008). Unfortunately, on the African continent, fossils, such as radiocarbon-dated ancient

samples, remain relatively rare and are often not representative of modern lineages (*e.g.,* Le Gall

2010 for insects). The lack of paleontological and archaeological records is partly due to their

fragility under the aridity conditions of the Sahara. The end-result is that the options to relate

population divergence to biogeographic events in this region are very limited. Finding a dating

strategy that does not rely on fossils would therefore constitute a key advance in understanding the

region's biogeography.

In this context, the use of versatile molecular markers, such as microsatellite loci, for which

evolutionary rates can be obtained from direct observation of germline mutations in the species of

interest, represents a useful alternative. Microsatellite mutation rates exceed by several orders of

magnitude that of point mutation in DNA sequences, ranging from 10^{-6} to 10^{-2} events per locus and

per generation (Ellegren 2000). Such rates allow one to both observe mutation events in parent-

offspring segregation data of realistic sample size and to reconstruct the recent history of related

populations. However, the use of microsatellite loci to estimate divergence times at recent

evolutionary time-scales still needs to overcome significant challenges. Since microsatellite allele

sizes result from the insertion or deletion of single or multiple repeat units and are tightly

constrained, these markers can be characterized by high levels of homoplasy that can obscure

inferences about gene history (*e.g.*, Estoup *et al.* 2002). In particular, at large time scales (*i.e.*, for

distantly related populations), genetic distance values no longer follow a linear relationship with

time. Rather, they reach a plateau and therefore provide biased unreliable estimation of divergence

time over a certain time threshold (Takezaki and Nei 1996; Feldman *et al.* 1997; Pollock 1998).

Microsatellites remain informative with respect to divergence time only if the population split

occurs within the period of linearity with time (Feldman *et al.* 1997; Pollock 1998). The exact value

of the differentiation threshold above which microsatellite markers would no longer accurately

reflect divergence times will depend on constraints on allele sizes and population-scaled mutation

rates (Feldman *et al.* 1997; Pollock 1998). In this context, the Approximate Bayesian Computation -

random forest (ABC-RF) approach recently proposed by Raynal *et al.* (2019) is a singular statistical advance, allowing both to take into account prior knowledge on genetic markers and to compute accuracy at a local (*i.e.*, posterior) scale. Using this method, one can consider to evaluate the divergence time threshold above which posterior estimates would become biased, and in this way thoroughly evaluate the robustness of inferences.

The desert locust, *S. gregaria*, is a generalist herbivore that can be found in arid grasslands and deserts in both northern and southern Africa (Figure 1a). In its northern range, the desert locust is one of the most widespread and harmful agricultural pest species, with a very large potential outbreak area, spanning from West Africa to Southwest Asia. The desert locust is also present in the southwestern arid zone (SWA) of Africa, which includes South-Africa, Namibia, Botswana and south-western Angola. The southern populations of the desert locust are termed *S. g. flaviventris* and are geographically separated by nearly 2,500 km from populations of the nominal subspecies from northern Africa, *S. g. gregaria* (Uvarov 1977). The isolation of *S. g. flaviventris* and *S. g. gregaria* lineages was recently supported by highlighting distinctive mitochondrial DNA haplotypes and male genitalia morphologies (Chapuis *et al.* 2016). Yet, the precise history of divergence remains elusive.

The main objective of the present study is to unravel the historical and evolutionary processes that have shaped the present disjoint geographical distribution of the desert locust and the genetic variation observed both within and between populations of its two subspecies. To this aim, we applied an ABC-RF approach and show its full potential to help discriminate between alternative biogeographic scenarios. We first start by identifying a set of evolutionary alternatives relevant to the species from African paleo-vegetation maps that reflect potential past distributions of the desert locust. We then used molecular data obtained from microsatellite markers for which we could obtain independent information on evolutionary rates and allele size constraints in the species of interest from direct observation of germline mutations (Chapuis *et al.* 2015). We applied recently available algorithms of the ABC-RF on our microsatellite population genetic data to compare a set

of thoroughly formalized and justified evolutionary scenarios and estimate the divergence time

between *S. g. gregaria* and *S. g. flaviventris* under the most likely of our scenarios. Finally, we

interpret our results in light of past vegetation cover and desert locust biology.

Results

Table 1 shows the values of the summary statistics obtained from the observed population dataset

consisting in two unstructured pooled samples of the subspecies *S. g. gregaria* and *S. g. flaviventris*,

A total of 170 individuals (*i.e.*, 80 and 90 individuals for *S. g. gregaria* and *S. g. flaviventris*,

respectively) were genotyped at 23 microsatellite markers derived from either genomic DNA (14

loci) or messenger RNA (9 loci) resources (hereafter referred to as untranscribed and transcribed

microsatellite markers, respectively). The level of differentiation between the two subspecies (as

measured by the parameter F_{ST}) was 0.04 and 0.12 for untranscribed and transcribed microsatellite

markers, respectively). The level of genetic diversity was higher within *S. g. gregaria* (+7 and 14%

for the mean number of alleles and expected heterozygosity, respectively).

Formalization and choice of evolutionary scenarios

Using a rich corpus of vegetation data, we reconstructed the present time (Fig. 1c) and past time

(Figs. 1d-f) distribution ranges of *S. gregaria* in Africa, going back to the Last Glacial Maximum

period (LGM, 26 to 14.8 Ky ago). Maps of vegetation cover for glacial arid maximums (Figs. 1e

and 1f) showed an expansion of open vegetation habitats sufficient to make the potential range of

the species continuous from the Horn of Africa in the north-west to the Cape of Good Hope in the

south. Maps of vegetation cover for interglacial humid maximums (Fig. 1d) showed a severe

contraction of deserts. These maps helped us formalize twelve competing evolutionary scenarios

(Figure 2), as well as bounds of prior distributions for various parameters (see the section *Prior*

setting for divergence parameters in Materials and methods). The twelve competing scenarios were

generated by different combinations of presence *vs.* absence of three key evolutionary events that

we identified as having potentially played a role in setting up the observed disjoint distribution of

the two locust subspecies: (i) a long population size contraction in the ancestral population, due to

the reduction of open vegetation habitats during the interglacial periods, (ii) a bottleneck in the

southern subspecies *S. g. flaviventris* right after divergence, associated to a single long-distance

migration event of a small fraction of the ancestral population, and (iii) a discrete genetic admixture

event either unidirectional from *S. g. gregaria* into *S. g. flaviventris*, or bidirectional, in order to

consider the many climatic transitions of the last Quaternary.

ABC-RF analyses supported the same group of scenarios or the same best individual

scenario for all ten replicate analyses (Table 2). The classification votes and posterior probabilities

estimated for the observed microsatellite dataset were the highest for the groups of scenarios in

which (i) *S. g. flaviventris* experienced a bottleneck event at the time of the split (scenarios

SB+SCB+SBA+SCBA; average of 2829 votes out of 3,000 RF-trees; posterior probability = 0.926),

(ii) the ancestral population experienced a population size contraction (scenarios

SC+SCB+SCA+SCBA; 2035 of 3,000 RF-trees; posterior probability = 0.682), and (iii) no

admixture event occurred between populations after the split (scenarios S+SC+SB+SCB; 2013 of

3,000 RF-trees; posterior probability = 0.700). When considering the twelve scenarios separately,

the highest classification vote was for scenario SCB (1521 of 3,000 RF-trees), which congruently

excludes a genetic admixture event and includes a population size contraction in the ancestral

population as well as a bottleneck event at the time of divergence in the *S. g. flaviventris* subspecies.

The posterior probability of scenario SCB averaged 0.564 over the ten replicate analyses (Table 2).

Table S1.1 (Supplementary Material S1) shows that only two other scenarios (SB and $SCBA_U$)

obtained at least 5% of the votes. The scenario SB included only a single bottleneck event in *S. g.*

flaviventris (mean of 15.7% of votes) and the scenario $SCBA_U$ included a bottleneck event in *S. g.*

flaviventris, a population size contraction in the ancestral population and an unidirectional genetic

admixture event from *S. g. gregaria* into *S. g. flaviventris* (mean of 10.8% of votes). All other

scenarios obtained less than 5% of the votes and were hence even more weakly supported.

We found that the posterior error rates (*i.e.*, 1 minus the posterior probabilities) were lower

than the prior error rates for the analyses considering either groups of scenarios based on the

presence (or not) of a bottleneck in *S. g. flaviventris* (*i.e.*, 7.4% versus 19.0%) or the scenarios

separately (*i.e.*, 43.6% versus 58.4%). For other groups of scenarios, the discrimination power was

similar at both the global (prior error rates) and local (posterior error rates) scales, with values

ranging from 26.1% to 32.1% (Table 2). Altogether, these results indicate that the observed dataset

belongs to a region of the data space where the power to discriminate among scenarios is higher

than the global power computed over the whole prior data space, and that the presence or absence of

a bottleneck in *S. g. flaviventris* is the demographic event with the most robust prediction in our

ABC-RF treatments. These results can be visually illustrated by the projection of the reference table

datasets and the observed one on a single (when analyzing pairwise groups of scenarios) or on the

first two linear discriminant analysis (LDA) axes (when analyzing the twelve scenarios considered

separately) (Figure S1.1, Supplementary Material S1).

Figure S1.2, Supplementary Material S1, illustrates how RFs automatically rank the

summary statistics according to their level of information. It shows that the number and set of most

informative statistics is different depending on the comparisons (groups of scenarios or individual

scenarios). Two sample statistics that measure the amount of genetic variation shared between

populations (F_{ST}, DM2 and LIK) were among the most informative when discriminating among

groups of scenarios including or not an admixture event. For groups of scenarios differing by

population size variation events, statistics summarizing variation between the two subspecies

samples (F_{ST} and DM2 for the bottleneck event in *S. g. flaviventris*; DAS and LIK for the

population size contraction in the ancestral population) and statistics summarizing genetic variation

within subspecies samples (mean expected heterozygosity and mean number of alleles for both

population size variation events) were among the most discriminative ones. Only fifteen single

sample statistics were not informative (according to their position relatively to the noise statistics

added to our treatments) when considering the twelve individual scenarios separately. Most of those

non informative statistics were associated to the set of transcribed microsatellites (Figure S1.3,

Supplementary Material S1).

Parameter estimation under the best evolutionary scenario

Figure 3a shows point estimates with 90% credibility intervals of the posterior distribution of the

divergence time between the two subspecies under the best supported scenario SCB. Our

estimations point to a young age of subspecies divergence, with a median divergence time of 2.6 Ky

and a 90% credibility interval of 0.9 to 6.6 Ky, when assuming an average of three generations per

year (Roffey and Magor 2003; Table 3 and Table S1.2, Supplementary Material S1). The accuracy

of divergence time estimation was similar at both the global and local scales (*i.e.*, normalized mean

absolute errors of 0.369 and 0.359, respectively; Table 3). Constraints on allele sizes in conjunction

with high population-scaled mutation rates potentially strongly affect the linearity of the

relationship between mutation accumulation and time of divergence estimated from microsatellite

data. We thus evaluated the accuracy of ABC-RF estimation of the population divergence time as a

function of the time scale, under scenario SCB. Analyses of pseudo-observed datasets showed that

the ABC-RF median estimate of divergence time reached a plateau for time scales $\geq$ 100,000

generations (Figure 4). Thus, the divergence time between *S. g. flaviventris* and *S. g. gregaria*

estimated on our real microsatellite dataset (~10,000 generations) is positioned within the period of

linearity with time, well before reaching a plateau reflecting a saturation of genetic information at

microsatellite markers. It is hence expected to represent a sensible estimation of the actual

divergence time.

Using the median as a point estimate, we estimated that the population size contraction in

the ancestor could have occurred at a time approximately three times older than the divergence time

between the subspecies (Table 3). Estimations of the ratio of stable effective sizes of the *S. g.*

gregaria and *S. g. flaviventris* populations (*i.e., N_f / N_g*) showed large 90% credibility intervals and

include the rate value of 1 (Table 3). Accuracy analysis indicates that our genetic data withhold

little information on this composite parameter (Table 3). The bottleneck intensity during the colonization of south-western Africa (*i.e.*, db_f / Nb_f) shows the highest accuracy of estimation (Table 3). The median of 1 and the 90% credibility interval of 0.5 to 2.4 exclude severe and mild bottlenecks and support a strong to moderate event (Table 3).

The most informative summary statistics were different depending on the parameter of interest (results not shown). For the time since divergence between the two subspecies, the most informative statistics corresponded to the expected heterozygosity computed within the *S. g. flaviventris* sample and the mean index of classification from *S .g. flaviventris* to *S. g. gregaria* (Figure S1.4, Supplementary Material S1). The addition of noise variables in our treatments showed that most statistics characterizing genetic variation within the *S. g. gregaria* sample were not informative.

Discussion

A young age of subspecific divergence

With a 90% credibility interval of the posterior density distribution of the divergence time at 0.9 to 6.6 Ky, our ABC-RF analyses clearly point to a divergence of the two desert locust subspecies occurring during the present Holocene geological epoch (0 to 11.7 Ky ago; Figure 3a). The posterior median estimate (2.6 Ky) and interquartile range (1.8 to 3.7 Ky) postdated the middle-late Holocene boundary (4.2 Ky). The latter past time boundary corresponds to the last transition from humid to arid conditions in the African continent (Figure 3b). This increasing aridity was shown to be a progressive change, with a concomitant maximum in northern and southern Africa at around 4 to 4.2 Ky ago, where aridity caused a contraction of the forest at its northern and southern peripheries without affecting its core region (Guo *et al.* 2000; Maley *et al.* 2018). Interestingly, the earliest archeological records of the desert locust found in Tin Hanakaten (Algeria) and Saqqara (Egypt) archaeological sites date back to this period (Figure 3b; Meinzingen 1993; Malek 1997;

Aumassip 2002). Pollen records also showed that during this period the plant community was

dominated by the desert and semi-desert taxa found today, including some species of prime

importance for the current ecology of the desert locust (Kröpelin *et al.* 2008, Shi *et al.* 1998,

Duranton *et al.* 2012). Then, the past 4 Ky are thought to have been under environmental stability

and as dry as at present. One can therefore reasonably assume that, at the inferred divergence time

between the two locust subspecies, the connectivity between the two African hemispheres was still

limited by the moist equator, in particular at the west, and by the savannahs and woodlands of the

eastern coast (Figure 1c). Consequently, contrary to most phylogeographic studies on other African

arid-adapted species (Atickem *et al.* 2018, Moodley *et al.* 2018), it is unlikely that the rather ancient

Quaternary climatic history explained the Southern range extension of the desert locust.

Recent geological and palynological research has shown that a brief fragmentation of the

African primary forest occurred during the Holocene interglacial from 2.5 Ky to 2.0 Ky ago

(reviewed in Maley *et al.* 2018). This forest fragmentation period is characterized by relatively

warm temperatures and a lengthening of the dry season rather than an arid climate. Although this

period does not correspond to a phase of general expansion of savannas and grasslands, it led to the

opening of the Sangha River Interval (SRI) in the core of the tropical forest in Central Africa (see

Fig. 1 in Maley *et al.* 2018). The SRI corresponds to a 400 km wide (14–18° E) open strip

composed of savannas and grasslands dividing the rainforest in a north-south direction. The SRI

corridor is thought to have facilitated the southern migration of Bantu-speaking pastoralists, along

with cultivation of the semi-arid sub-Saharan cereal, pearl millet, *Pennisetum glaucum* (Schwartz

1992; Bostoen *et al*. 2015). The Bantu expansion took place between approximately 5 and 1.5 Ky

ago and reached the southern range of the desert locust, including northern Namibia for the Western

Bantu branch and southern Botswana and eastern South Africa for the Eastern Bantu branch

(Vansina 1995). We cannot exclude that the recent subspecific distribution of the desert locust has

been mediated by this recent ecological disturbance, which included a north-south corridor of open

vegetation habitats and the diffusion of agricultural landscapes through the Bantu expansion. The

progressive reappearance of forest vegetation 2 Ky ago would have then led to the present-day

isolation and subsequent genetic differentiation of the new southern populations from northern

parental populations. However, the level of climatic and habitat favorability of the SRI

environmental disturbance to the species, and thereby the likelihood of a south-eastern colonization

through the SRI corridor, remains to be evaluated in line with further data on this period.

On the role of dispersal on subspecific divergence

Our ABC-RF results indicate that a demographic bottleneck (*i.e.*, a strong transitory reduction of

effective population size) occurred in the nascent southern subspecies of the desert locust. The high

posterior probability value (92.6%) shows that this evolutionary event could be inferred with strong

confidence. This result is compatible with the abovementioned colonization hypothesis if the

proportion of suitable habitats for the desert locust in the SRI corridor was low, strongly limiting

the carrying capacity during the time for range expansion. This scenario reduces, at the same time,

the likelihood of a successful colonization through the SRI corridor. A more plausible alternative

for a bottleneck event in *S. g. flaviventris* is a southern colonization of Africa through an

exceptional long-distance migration event. In winged migratory species, movements are assisted by

high velocity winds and may ascend to high altitudes (2000 m in the desert locust; Uvarov 1977)

(Pedgley *et al.* 1995). Although largest insects exert some control over their direction of migration

by flying actively, or at least gliding within (Pedgley *et al.* 1995), accidental displacements in wind

directions markedly different from that of seasonal dominant winds that assist migrations are likely.

Such accidental displacements at long-distance (*i.e.*, a few to several thousand kms) were recorded

in a number of migratory species, such as cicadas, butterflies, moths and locusts (Pedgley *et al.*

1995; Lorenz, 2009).

To survive in its erratic arid and semi-arid habitat, the desert locust migrates downwind to

reach areas where rain has recently fallen and exploit newly available resources. The dynamics of

prevailing winds and pressure over Africa predicts the likelihood of a southeastern transport of

locusts (Nicholson 1996, Waloff and Pedgley 1986). In northern Africa, at least since 2.7Ky, the strong northeast trade winds bring desert locust swarms equatorward in the moist intertropical convergence zone (Figures 1a-b; Kröpelin *et al.* 1998). Most winds are westerlies (Figures 1a-b), nevertheless easterly winds flow parallel to the eastern coast of Africa in northern winter (*e.g.*, January; Figure 1b). In southern Africa, winds blow mostly from the north-east toward the extant south-western distribution of the desert locust in southern winter (*e.g.*, August; Figure 1a). In agreement with this, southward movements of desert locust have been documented along the eastern coast of Africa, in southern Tanzania during the plagues of 1926-1934, 1940-1948 and 1949-1963 (Waloff 1976), and even in Mozambique in January 1945 during the peak of the major plague of 1940-1948 (Waloff 1966).

Furthermore, most travels off the range listed in the history of the desert locust were associated with plague events (Waloff 1976), with other records including Portugal, the British Isles and the famous trans-Atlantic crossing observed in October 1988, where large numbers of locusts landed in the Caribbean islands (Richardson and Nemeth, 1991). Plagues typically culminate several years of above average rainfall, resulting in abundant vegetation that supports both cumulative locust population growth and full development of gregarious characteristics (Richardson and Nemeth, 1991). The gigantic numbers (over a billion) of swarming locusts may facilitate the success of long-distance migrations, in spite of high mortality. In addition, flight capacity and endurance of the gregarious phase are remarkable, with swarms of winged adults regularly traveling up to hundreds km in a day, in case of sudden lack of resources (Roffey and Magor 2003). Therefore, out-of-range displacements of the desert locust may be explained by the synergistic effects of exceptional environmental conditions (*e.g.*, unusual winds in strength or direction, rain favorable to plague) and peculiar biology of the gregarious phase of this species (*e.g.*, swarming behavior, huge numbers of dispersers).

In conclusion, dynamics of dominant winds in subtropical desert, historical records of exceptional migrations in the desert locust, and the very robust prediction for the presence of a

bottleneck in *S. g. flaviventris* right after divergence by our ABC-RF treatments altogether support

that a single or a few swarm(s) from the Central region of the desert locust range sourced during a

plague the colonization of south-western Africa, suggesting for a role of dispersal in the disjoint

distribution and divergence of desert locust populations. Interestingly, an exceptional trans-Atlantic

flight of the desert locust from Africa to South America was revealed to give rise to the radiation of

some 50 *Schistocerca* species in the western hemisphere (Lovejoy *et al.*, 2006).

On the influence of climatic cycles

It may appear surprising, at least at first sight, that the southern colonization of the desert locust did

not occur during one of the major glacial episodes of the last Quaternary cycle, since these periods

are characterized by a more continuous range of the desert locust (see paleo-vegetation maps in

Figures 1e-f). In particular, during the last glacial maximum (LGM, -14.8 Ky to -26 Ky), the Sahara

desert extended hundreds of km further South than at present and annual precipitation were lower

(i.e. ~200–1,000 mm/year). Several hypotheses explain why our evolutionary scenario choice

procedure provided low support to the possibility of a birth of the locust subspecies *S. g. flaviventris*

at older periods. First, we cannot exclude that our microsatellite genetic data allow making

inferences about the last colonization event only. Error rates at both local and global scales for the

choice of scenario groups including or not a genetic admixture event after the split indicated that

our discrimination power to infer this specific evolutionary event was poor (*i.e.*, 32.1% and 30.0%

respectively, Table 2). The recent north-to-south colonization event selected by our ABC-RF

treatment may hence have blurred traces of older colonization events.

Second, while there is large evidence that much of Africa was drier during the last glacial

phase, this remains debated for southwestern Africa (see the gray coloration in Figure 3b). Some

climate models show that at least some parts of this region, such as the Kalahari Desert, may have

experienced higher rainfall than at present (Cockcroft 1987; Ganopolski *et al.* 1998; Chase and

Meadows 2007). Such regional responses to glacial cycles may have prolonged until the middle

Holocene. In particular, the northern Younger Dryas (*i.e.*, -12.9 to -11.7 Ky) can be correlated only

partly with an arid period in the southern hemisphere (*i.e.*, -14.4 to -12.5 Ky). Such older climate

episodes in antiphase between hemispheres (see the sandy brown coloration in Figure 3b) may have

prevented from either a successful north-to-south migration event or a successful establishment and

spread in the new southern range.

Third, although semi-desert and desert biomes were more expanded than at present during

the LGM, extreme aridity and lowered temperatures may have actually been unfavorable to the

species. For example, mean temperatures lowered by 5 to 6°C in both southern-western Africa

(Stute and Talma 1997) and Central Sahara (Edmunds *et al.* 1999).The maintenance of desert locust

populations depends on the proximity of areas with rainfalls at different seasons or with the

capacity to capture and release water. For instance, in the African northern range, breeding success

of locust populations relies on seasonal movements between the Sahel-Saharan zones of inter-

tropical convergence, where the incidence of rain is high in summer, and the Mediterranean-

Saharan transition zone, with a winter rainfall regime (Rainey and Waloff 1951). In addition, adult

migration and nymphal growth of the desert locust are dependent upon high temperature (Roffey

and Magor 2003). It is hence possible then that the conjunction of hyper-aridity with intense cold

could not easily support populations of the desert locust, despite the high extent of their migrations.

Interestingly, climatic reconstructions during the Last Glacial Maximum (LGM) indeed showed

dramatic decreases of mean temperature throughout all of Africa and of precipitation in the Sahelian

region (Figs. S2.3 and S2.4, Supplementary Material S2). Accordingly, species distribution

modelling showed a LGM distribution for the desert locust very similar to its current distribution

(Figs. S2.5, S2.6 and S2.7, Supplementary Material S2), excluding the hypothesis of a more

continuous range of the species at this time.

While ABC-RF analyses did not support that the Quaternary climatic history explained the

subspecific divergence in the desert locust, they provided evidence for the occurrence of a large

contraction of the size of the ancestral population preceding the divergence. Using the median as a

point estimate, we estimated that the population size contraction in the ancestor could have occurred

at a time about three fold older than the divergence time between the subspecies. This corresponds

to the African humid period in the early and middle stages of the Holocene, though the large

credibility interval also included the last interglacial period of the Pleistocene (Figure 3b). Such

population size contraction was likely induced by the severe(s) contraction(s) of deserts that

prevailed prior the estimated divergence between the two subspecies. Interestingly, these humid

periods were more intense and prolonged in northern Africa, which corresponded to the presumed

center of origin of the most recent common ancestor (Scott 1993; Partridge 1997; Shi *et al.* 1998).

Statistical advances by means of ABC Random Forest

To our knowledge, the present study is the first one using recently developed ABC-RF algorithms

(Raynal *et al.* 2019) to carry out inferences about parameter estimation on a real multi-locus

microsatellite dataset (for scenario choice see for example Rougemont *et al.* (2016) and Fraimout *et*

al. (2017)). When compared with various ABC solutions, this new RF method offers several

advantages: a significant gain in terms of robustness to the choice of the summary statistics,

independence from any type of tolerance level, and a good trade-off in terms of quality of point

estimator precision of parameters and credible interval estimations for a given computing time

(Raynal *et al.* 2019). The RF approach that we used here includes three novelties in statistical

analyses that were particularly useful for reconstructing the evolutionary history of the divergence

between *S. g. gregaria* and *S. g. flaviventris* subspecies. First, our ABC-RF statistical treatments

benefited from the incorporation of previous estimations of mutation rates and allele size constraints

for the microsatellite loci used in this study (for details see the Materials and methods section

Microsatellite dataset, mutation rate and mutational model). Microsatellite mutation rate and

pattern of most eukaryotes remains to a large extent unknown, and, to our knowledge, the present

study is a rare one where independent information on mutational features was incorporated into the

microsatellite prior distributions. Second, given the low RF computing time, we could simulate

large pseudo-observed datasets to compute error measures conditionally to a subset of fixed

divergence time values chosen to cover the entire prior interval. In this way, we showed that

posterior distributions for estimation of divergence time between the two subspecies accurately

reflected true divergence time values. Third, because error levels may differ depending on the

location of an observed dataset in the prior data space (e.g. Pudlo *et al.* 2016), prior-based indicators

are poorly relevant, aside from their use to select the best classification method and set of

predictors, here our summary statistics. Therefore, in addition to global prior errors, we computed

local posterior errors, conditionally to the observed dataset. The latter errors measure prediction

quality exactly at the position of the observed dataset. For parameter estimation accuracy, we

propose an innovative way to approximate local posterior errors, relying partly on out-of-bag

predictions and implemented in a new version of the R library abcrf (version 1.8) available on R

CRAN (see the Supplementary Material S3 for a detailed description). Such statistical advances are

of general interest and will be useful for any statistical treatments of massive simulation data,

including for inferences using single nucleotide polymorphisms (*i.e,*. SNPs) obtained from new

generation sequencing technologies.

Materials & Methods

Formalization of evolutionary scenarios

To help formalize the evolutionary scenarios to be compared, we relied on maps of vegetation

cover in Africa from the Quaternary Environment Network Atlas (Adams and Faure 1997). We

considered the periods representative of arid maximums (LGM and YD; Fig.1e-f, humid maximums

(HCO; Fig.1d), and present-day arid conditions (Fig.1c), for which reliable vegetation

reconstructions have been published. Desert and xeric shrubland cover fits well with the present-day

species range during remission periods. Tropical and Mediterranean grasslands were added

separately since the species inhabits such environments during outbreak periods only. The

congruence between present maps of species distribution (Fig.1a) and of open vegetation habitats

(Fig.1c) suggests that vegetation maps for more ancient periods could be considered as good

approximations of the potential range of the desert locust in the past (but see section '*On the*

influence of climatic cycles' in the Discussion). Maps of vegetation cover during ice ages (Figs. 1e

and 1f) show an expansion of open vegetation habitats (*i.e.*, grasslands in the tropics and deserts in

both the North and South of Africa) sufficient to make the potential range of the species continuous

from the Horn of Africa in North-West to the Cape of Good Hope in the South. Notice that we also

explored species distribution modelling for the HCO and LGM periods as an alternative to using

vegetation as a surrogate for the locust range (Supplementary Materials S2). However, distribution

modelling provided a narrower set of alternative hypotheses than the vegetation-based scenarios

mentioned above and are therefore not discussed any further.

Based on the above paleo-vegetation map reconstructions, we considered a set of alternative

biogeographic hypotheses formulated into different types of evolutionary scenarios. First, we

considered scenarios involving a more or less continuous colonization of southern Africa by the

ancestral population from a northern origin. In this type of scenario, effective population sizes were

allowed to change after the divergence event, without requiring any bottleneck event (*i.e.*, without

any abrupt and strong reduction of population size) right after divergence. Second, we considered

the situation where the colonization of Southern Africa occurred through a single (or a few) long-

distance migration event(s) of a small fraction of the ancestral population. This situation was

formalized through scenarios that differed from a continuous colonization scenario by the

occurrence of a bottleneck event in the newly founded population. The bottleneck event occurred

into *S. g. flaviventris* right after divergence and was modelled through a limited number of founders

during a short period.

Because the last Quaternary cycle includes several arid climatic periods, including the

intense punctuation of the Younger Dryas (YD) and the last glacial maximum (LGM), we also

considered scenarios that incorporated the possibility of a discrete genetic admixture event, either

bidirectional or unidirectional from *S. g. gregaria* into *S. g. flaviventris*. Since previous tests based

on simulated data showed a poor power to discriminate between a single versus several admixture

events (results not shown), we considered only scenarios including a single admixture event.

Finally, at interglacial humid maximums, the map of vegetation cover showed a severe

contraction of deserts, which were nearly completely vegetated with annual grasses and shrubs and

supported numerous perennial lakes (Fig.1d; deMenocal *et al.* 2000). We thus considered the

possibility that vegetation-induced contractions of population sizes have pre-dated the separation of

the two subspecies. Hence, whereas so far scenarios involved a constant effective population size in

the ancestral population, we formalized alternative scenarios in which we assumed that a long

population size contraction event occurred into the ancestral population at a time t_C, with an

effective population size N_C for a duration d_C.

Combining the presence or absence of the three above-mentioned key evolutionary events (a

bottleneck in *S. g. flaviventris,* a bidirectional or an unidirectional genetic admixture from *S. g.*

gregaria into *S. g. flaviventris*, and a population size contraction in the ancestral population)

allowed defining a total of twelve scenarios, that we compared using ABC-RF. The twelve

scenarios with their historical and demographic parameters are graphically depicted in Figure 2. All

scenarios assumed a northern origin for the common ancestor of the two subspecies and a

subsequent southern colonization of Africa. This assumption is supported by recent mitochondrial

DNA data showing that *S. g. gregaria* have higher levels of genetic diversity and diagnostic bases

shared with outgroup and congeneric species, whereas *S. g. flaviventris* clade was placed at the

apical tip within the species tree (Chapuis *et al.* 2016). In agreement with this assumption,

preliminary analyses based on observed data showed a low support for a southern origin for the

common ancestor of the two subspecies and a subsequent northern colonization of Africa (results

not shown).

All scenarios considered three populations of current effective population sizes N_f for *S. g.*

flaviventris, N_g for *S. g. gregaria*, and N_a for the ancestral population, with *S. g. flaviventris* and *S.*

g. gregaria diverging t_S generations ago from the ancestral population. The bottleneck event which

potentially occurred into *S. g. flaviventris* was modelled through a limited number of founders N_B

during a short period d_B. The potential unidirectional genetic admixture into *S. g. flaviventris*

occurred at a time t_A, with a proportion r_{AU} of genes of *S. g. gregaria* origin. (or r_{AB}). In the case of

a bidirectional genetic admixture, still occurring at a time t_A, the proportion of *S. g. gregaria* genes

entering into the *S. g. flaviventris* population was r_{AB} and the proportion of *S. g. flaviventris* genes

entering into the *S. g. gregaria* population was 1- r_{AB}. The potential population size contraction

event occurred into the ancestral population at a time t_C, with an effective population size N_C during

a duration d_C.

Prior setting for historical and demographical parameters

Prior values for time periods between sampling and admixture, divergence and/or ancestral

population size contraction events (t_C, t_S and t_A, respectively) were drawn from log-uniform

distributions bounded between 100 and 500,000 generations, with $t_C > t_S > t_A$. Assuming an average

of three generations per year (Roffey and Magor 2003), this prior setting corresponds to a time

period that goes back to the second-to-latest glacial maximum (150 Ky ago) (de Vivo and

Carmignotto 2004, deMenocal *et al.* 2000). Preliminary analyses showed that assuming a uniform

prior shape for all time periods (instead of log-uniform distributions) do not change scenario choice

results, with posterior probabilities only moderately affected, and this despite a substantial increase

of out-of-bag prior error rates (*e.g.*, + 50% when considering the twelve scenarios separately; Table

S4.1, Supplementary Material S4). Analyses of simulated pseudo-observed datasets (pods) showed

that assuming a uniform prior rather than a log-uniform prior for time period parameters would

have also biased positively the median estimate of the divergence time and substantially increased

its 90% credibility interval (Figure S4.1 and Table S4.2, Supplementary Material S4). Using a log-

uniform distribution remains a sensible choice for parameters with ranges of values covering

several if not many log-intervals, as doing so allows assigning equal probabilities to each of the log-

intervals.

We used uniform prior distributions bounded between 1×10^4 and 1×10^6 diploid individuals

for the different stable effective population sizes N_f, N_g and N_a (Chapuis *et al.* 2014). The admixture

rate (*i.e.,* the proportion of *S. g. gregaria* genes entering into the *S. g. flaviventris* population), was

drawn from a uniform prior distribution bounded between 0.05 and 0.50 for a unidirectional event

(r_{AU}) and between 0.05 and 0.95 for a bidirectional event (r_{AB}). We used uniform prior distributions

bounded between 2 and 100 for both the numbers of founders (in diploid individuals) and durations

of bottleneck events (in number of generations). For the contraction event, we used uniform prior

distributions bounded between 100 and 10,000 for both the population size N_C (in diploid

individuals) and duration d_C (in number of generations). Assuming an average of three generations

per year (Roffey and Magor 2003), such prior choice allowed a reduction in population size for a

short to a relatively long period, similar for instance to the whole duration of the HCO (from 9 to

5.5 Ky ago) which was characterized by a severe contraction of deserts.

Microsatellite dataset, mutation rate and mutational model

We carried out our statistical inference on the microsatellite dataset previously published in Chapuis

et al. (2016). The 23 microsatellite loci genotyped in that dataset were derived from either genomic

DNA (14 loci) or messenger RNA (9 loci) resources, and were hereafter referred to as untranscribed

and transcribed microsatellite markers (following Blondin *et al.* 2013). These microsatellites were

shown to be genetically independent, free of null alleles and at selective neutrality (Chapuis *et al.*

2016). Previous levels of F_{ST} (Weir 1996) and Bayesian clustering analyses (Pritchard *et al.* 2000)

among populations showed a weak genetic structuring within each subspecies (Chapuis *et al.* 2014,

2017). For each subspecies, we selected and pooled three population samples in order to ensure

both a large sample size (*i.e.*, 80 and 90 individuals for *S. g. gregaria* and *S. g. flaviventris*,

respectively), while ensuring a non-significant genetic structure within each subspecies pooled

sample, as indicated by non-significant Fisher's exact tests of genotypic differentiation among the

three initial population samples within subspecies and exact tests of Hardy-Weinberg equilibrium

for each subspecies pooled sample (*i.e.*, *P*-value > 0.05; Genepop 4.0; Rousset 2008). More

precisely, the *S. g. gregaria* sample consisted in pooling the population samples 8, 15 and 22 of

Chapuis *et al.* (2014) and the *S. g. flaviventris* sample included the population samples 1, 2 and 6 of

Chapuis *et al.* (2017).

Mutations occurring in the repeat region of each microsatellite locus were assumed to follow

a symmetric generalized stepwise mutation model (GSM; Zhivotovsky *et al.* 1997; Estoup *et al.*

2002). Prior values for any mutation model settings were drawn independently for untranscribed

and transcribed microsatellites in specific distributions. Because allele size constraints exist at

microsatellite markers, we informed for each microsatellite locus their lower and upper allele size

bounds using values estimated in Chapuis *et al.* (2015), following the approach of Pollock *et al.*

(1998) and microsatellite data from several species closely related to *S. gregaria* (Blondin *et al.*

(2013). Prior values for the mean mutation rates ($\overline{\mu_R}$) were set to the empirical estimates inferred

from observation of germline mutations in Chapuis *et al.* (2015), *i.e.,* 2.8×10^{-4} and 9.1×10^{-5} for

untranscribed and transcribed microsatellites, respectively. The parameters for individual

microsatellites were then drawn from a Gamma distribution with mean = $\overline{\mu_R}$ and shape = 0.7

(Estoup *et al.* 2001) for both types of microsatellites. We ensured that the chosen value of shape

parameter generated the same inter-loci variance as estimated in Sun *et al.* (2012) from direct

observations of thousands of human microsatellites. Prior values for the mean parameters of the

geometric distributions of the length in number of repeats of mutation events ($\overline{P}$) were set to the

proportions of multistep germline mutations observed in Chapuis *et al.* (2015), *i.e.*, 0.14 and 0.67

for untranscribed and transcribed microsatellites, respectively. The *P* parameters for individual loci

were then standardly drawn from a Gamma distribution (mean = $\overline{P}$ and shape = 2). We also

considered mutations that insert or delete a single nucleotide to the microsatellite sequence. To

model this mutational feature, we used the DIYABC default setting values (*i.e.*, a uniform

distribution bounded between [10^{-8}, 10^{-5}] for the mean parameter $\overline{\mu_{SNI}}$ and a Gamma distribution

(mean = $\overline{\mu_{SNI}}$ and shape = 2) for individual loci parameters; Cornuet *et al.* 2010; see also DIYABC

user manual p. 13, http://www1.montpellier.inra.fr/CBGP/diyabc/).

Analyses using ABC Random Forest

We used the software DIYABC v.2.1.0 (Cornuet *et al.* 2014) to simulate datasets constituting the

so-called reference tables (*i.e.*, records of a given number of datasets simulated using the scenario

ID and the evolutionary parameter values sampled from prior distributions and summarized with a

pool of statistics). Random-forest computations were then performed using a new version of the R

library ABCRF (version 1.8) available on the CRAN. This version includes all ABC-RF algorithms

detailed in Pudlo *et al.* (2016), Raynal *et al.* (2019) and Estoup *et al.* (2018a) for scenario choice

and parameter estimation, as well as several statistical novelties allowing to compute error rates in

scenario choice and accuracy measures for parameter estimation (see details below). An overview

of the ABC-RF methods used in the present paper is provided in Supplementary Material S3.

Readers can also consult Pudlo *et al.* (2016), Rougemont *et al.* (2016), Fraimout *et al.* (2017),

Estoup *et al.* (2018a,b) and Marin *et al.* (2018) for scenario choice, and Raynal *et al.* (2019) for

parameter estimation to access to further detailed statistical descriptions, testing and applications of

ABC-RF algorithms.

For scenario choice, the outcome of the first step of the ABC-RF statistical treatment applied

to a given target dataset is a classification vote for each scenario which represents the number of

times a scenario is selected in a forest of *n* trees. The scenario with the highest classification vote

corresponds to the scenario best suited to the target dataset among the set of compared scenarios.

This step also provides an error rate relevant to the entire prior sampling space, the global prior

error. See the section *Global prior errors* in Supplementary Material S2 for details. The second RF

analytical step provides a reliable estimation of the posterior probability of the best supported

scenario. One minus such posterior probability yields the local posterior error associated to the

observed dataset (see the section *Local posterior errors* in Supplementary Material S3). In practice,

ABC-RF analyses were processed by drawing parameter values into the prior distributions

described in the two previous sections and by summarizing microsatellite data using a set of 32

statistics (see Table 1 for details about such summary statistics) and the one LDA axis or eleven

LDA axes (i.e. number of scenarios minus 1; Pudlo *et al.* 2016) computed when considering

pairwise groups of scenarios or individual scenarios, respectively. We processed ABC-RF

treatments on reference tables including 150,000 simulated datasets (*i.e.,* 12,500 per scenario).

Following Pudlo et al. (2016), we checked that 150,000 datasets was sufficient by evaluating the

stability of prior error rates and posterior probabilities estimations of the best scenario on 50,000,

80,000 and 100,000 and 130,000 simulated datasets (Table S1.3, Supplementary Material S1). The

number of trees in the constructed random forests was fixed to $n = 3,000$, as this number turned out

to be large enough to ensure a stable estimation of the prior error rate (Figure S1.5, Supplementary

Material S1). We predicted the best scenario and estimated its posterior probability and prior error

rate over ten replicate analyses based on ten different reference tables.

In order to assess the power of our approach to infer each specific evolutionary event of

interest, we first processed ABC-RF analyses by grouping scenarios based on the presence or

absence of each type of evolutionary event that we identified as having potentially played a role in

setting up the disjoint distribution of the two locust subspecies (*e.g.*, Roux *et al.* 2016, Leroy *et al*.

2017 and Estoup *et al.* 2018a). We conducted ABC-RF treatments on three pairwise groups of

scenario (with four scenarios per group): groups of scenarios with *vs.* without a bottleneck in *S. g.*

flaviventris, groups with *vs.* without a population size contraction in the ancestral population, and

groups with *vs.* without an asymmetrical genetic admixture event from *S. g. gregaria* into *S. g.*

flaviventris. We then conducted ABC-RF treatments on the twelve scenarios considered separately.

For parameter estimation, we constructed ten independent replicate RF treatments based on

ten different reference tables for each parameter of interest (Raynal *et al.* 2019): the time since

divergence, the ratio of the time of the contraction event into the ancestral population on the time

since divergence, the intensity of the bottleneck event in the sampled *S. g. flaviventris* population

(defined as the ratio of the bottleneck event of duration db_f on the effective population size Nb_f) and

the ratio of the stable effective population size of the two sampled populations. For each RF

treatment, we simulated a total of 100,000 datasets for the selected scenario (drawing parameter

values into the prior distributions described in the two previous sections and using the same 32

summary statistics). Following Raynal *et al.* (2019), we checked that 100,000 datasets was

sufficient by evaluating the stability of the measure of accuracy on divergence time estimation using

50,000, 80,000 and 90,000 simulated datasets (Table S1.4, Supplementary Material S1). The

number of trees in the constructed random forests was fixed to $n = 2,000$, as such number turned out

to be large enough to ensure a stable estimation of the measure of divergence time estimation

accuracy (Figure S1.6, Supplementary Material S1). For each RF treatment, we estimated the

median value and the 5% and 95% quantiles of the posterior distributions. It is worth noting that we

considered median values as the later provided more accurate estimations (according to out-of-bag

predictions) than when considering mean values (results not shown). Accuracy of parameter

estimation was measured using out-of-bag predictions and the normalized mean absolute error

(NMAE). NMAE corresponds to the mean of the absolute difference between the point estimate

(here the median) and the (true) simulated value divided by the simulated value (formula detailed in

Supplementary Material S3).

Finally, because microsatellite markers tend to underestimate divergence time for large time

scales due to allele size constraints, we evaluated how the accuracy of ABC-RF estimation of the

time of divergence between the two subspecies was sensitive to the time scale. To this aim, we used

DIYABC to produce pseudo-observed datasets assuming fixed divergence time values chosen to

cover the prior interval (100 ; 250 ; 500; 1,000 ; 2,500; 5,000 ; 10,000 ; 25,000 ; 50,000; 100,000;

250,000 generations) and using the best scenario. We simulated 5,000 of such test datasets for each

of the eleven divergence time values. Each of these test dataset was treated using ABC-RF in the

same way as the above target observed dataset.

Acknowledgements

This work was supported by research funds from the French Agricultural Research Centre for International Development (CIRAD), the project ANR-16-CE02-0015-01 (SWING), the INRA scientific department SPE (AAP-SPE 2016), and the Labex NUMEV (NUMEV, ANR10-LABX-20). The data used in this work were partly produced through the technical facilities of the Centre Méditerranéen Environnement Biodiversité, Montpellier. We are grateful to Pierre-Emmanuel Gay for assistance with maps, and Antoine Foucart, Gauthier Dobigny and Jean-Yves Rasplus for fruitful discussions. We thank Michael D. Greenfield and two anonymous reviewers for constructive and insightful comments on an earlier version of the manuscript.

Conflict of interest disclosure

The authors of this preprint declare that they have no financial conflict of interest with the content of this article. Arnaud Estoup is one of the *PCI Evolutionary Biology* recommenders.

Data accessibility

Microsatellite genotypic data and R scripts for ABC-RF analyses are available at https://dataverse.cirad.fr/.

References

Adams JM, Faure H (1997) (eds), QEN members. Review and Atlas of Palaeovegetation: Preliminary land ecosystem maps of the world since the Last Glacial Maximum. Oak Ridge National Laboratory, TN, USA.

Atickem A, Stenseth NC, Drouilly M, Bock S, Roos C, Zinner D (2018) Deep divergence among mitochondrial lineages in African jackals. *Zool Scripta*, **47**, 1-8.

Aumassip G (2002) L'algérie des premiers hommes. Ibis Press, 230 p.

Beaumont MA, Zhang W, Balding DJ (2002) Approximate Bayesian Computation in Population Genetics. *Genetics*, **162**, 2025–2035.

Blondin L, Badisco L, Pagès C, Foucart A, Risterucci AM, Bazelet CS, Vanden Broeck J, Song H, Ould Ely S, Chapuis M-P (2013) Characterization and comparison of microsatellite markers derived from genomic and expressed libraries for the desert locust. *J Appl Entomol*, **137**, 673–683.

Bond G *et al.* (1997) A pervasive millenial-scale cycle in North Atlantic Holocene and Glacial climates. Science, **278**, 1257-1265.

Bostoen K, Clist B, Doumenge C, Grollemund R, Hombert JM, Muluwa JK, Maley J (2015) Middle to Late Holocene paleoclimatic change and early Bantu expansion in the rain forests of Western Central Africa. *Curr Anthropol*, **56**, 354–384.

Breiman, L. (2001) Random Forests. *Machine Learning*, **45**(1), 5–32.

Chapuis M-P, Bazelet CS, Blondin L, Foucart A, Vitalis R, Samways MJ. (2016) Subspecific taxonomy of the desert locust, *Schistocerca gregaria* (Orthoptera: Acrididae), based on molecular and morphological characters. *Syst Entomol*, **41**, 516-530.

Chapuis M-P, Foucart A, Plantamp P, Blondin L, Leménager N, Benoit L, Gay P-E, Bazelet CS (2017) Genetic and morphological variation in non-polyphenic southern African populations of the desert locust. *Afr Entomol*, **25**, 13-23.

Chapuis M.-P, Plantamp P, Blondin B, Pagès C, Vassal J.-M., Lecoq M (2014) Demographic
processes shaping genetic variation of the solitarious phase of the desert locust. *Mol Ecol,*
23, 1749–1763.

Chapuis M-P, Plantamp C, Streiff R, Blondin L, Piou C (2015) Microsatellite evolutionary rate and
pattern in *Schistocerca gregaria* inferred from direct observation of germline mutations. *Mol
Ecol*, **24**, 6107-6119.

Chase BM, Meadows ME (2007) Late Quaternary dynamics of southern Africa's winter rainfall
zone. *Earth Sci Rev*, **84**, 103-138.

Chase BM *et al.* (2009) A record of rapid Holocene climate change preserved in hyrax middens
from SW Africa. *Geology* **37**, 703-706.

Cockcroft MJ, Wilkinson MJ, Tyson PD (1987) The application of a present-day climatic model to
the Late Quaternary in southern Africa. *Climate Change*, **10**, 161-181.

Cornuet J-M, Ravigne V, Estoup A (2010) Inference on population history and model checking
using DNA sequence and microsatellite data with the software DIYABC (v1.0). *BMC
Bioinformatics*, **11**, 401.

Cornuet J-M, Pudlo P, Veyssier J, Dehne-Garcia A, Gautier M, Leblois R, Marin J-M, Estoup A
(2014) DIYABC v2.0: a software to make Approximate Bayesian Computation inferences
about population history using Single Nucleotide Polymorphism, DNA sequence and
microsatellite data. *Bioinformatics*, **30**, 1187–1189.

deMenocal PB (1995) Plio-Pleistocene African climate, *Science*, 270, 53-59.

deMenocal PB, Ortiz J, Guilderson T, Adkins J, Sarnthein M, Baker L, Yarusinsky M (2000)
Abrupt onset and termination of the African Humid Period: Rapid climate response to
gradual insolation forcing. *Quat Sci Rev*, **19**, 347-361.

deMenocal PB (2004) African climate change and faunal evolution during the Pliocene-Pleistocene.
Earth and Planetary Science Letters, **220**, 3–24.

Dib C, Fauré S, Fizames C, Samson D, Drouot N, Vignal A, Millasseau P, Marc S, Hazan J, Seboun

E, *et al.* (1996) A comprehensive genetic map of the human genome based on 5264

microsatellites. *Nature*, 380, 152–154

Dupont LM (2011) Orbital scale vegetation change in Africa *Quat Sci Rev*, **30**, 3589-3602.

Duranton JF, Foucart A, Gay P-E (2012) Florule des biotopes du criquet pèlerin en Afrique de

l'Ouest et du Nord-Ouest à l'usage des prospecteurs de la lutte antiacridienne. Rome : FAO,

487 p.

Edmunds WM, Fellman E, Goni IB (1999) Lakes, groundwater and palaeohydrology in the Sahel of

NE Nigeria: evidence from hydrogeochemistry. *J Geol Soc Lond*, **156**, 345–355.

Ellegren H (2000) Microsatellite mutations in the germline: implications for evolutionary inference.

Trends Genet, **16**, 551–558.

Estoup A, Jarne P, Cornuet J-M (2002) Homoplasy and mutation model at microsatellite loci and

their consequences for population genetics analysis. *Mol Ecol*, **11**, 1591-1604.

Estoup A, Raynal L, Verdu P, Marin J-M (2018a) Model choice using Approximate Bayesian

Computation and Random Forests: analyzes based on model grouping to make inferences

about the genetic history of Pygmy human populations. *J Soc Fr Statistique*, **159**, 167-190.

Estoup A, Verdu P, Marin J-M, Robert C, Dehne-Garcia A, Cornuet J-M, Pudlo P (2018b)

Application of approximate Bayesian computation to infer the genetic history of Pygmy

hunter-gatherers populations from Western Central Africa. *Handbook of Approximate*

Bayesian Computation. Chapman and Hall/CRC.

Estoup A, Wilson IJ, Sullivan C, Cornuet J-M, Moritz C (2001) Inferring population history from

microsatellite et enzyme data in serially introduced cane toads, *Bufo marinus*. *Genetics*, **159**,

1671-1687.

Feldman MW, Bergman A, Pollock DD, Goldstein DB (1997) Microsatellite genetic distances with

range constraints: analytic description and problems of estimation. *Genetics*, **145**, 207–216.

Fraimout A, Debat V, Fellous S, Hufbauer RA, Foucaud J, Pudlo P, Marin J-M, Price DK, Cattel J
et al. (2017). Deciphering the routes of invasion of *Drosophila suzukii* by means of ABC
random forest. *Mol Biol Evol*, **34**(4), 980–996.

Ganopolski A, Rahmstorf S, Petoukohov V, Claussen M (1998) Simulation of modern and glacial
climates with a coupled global model of intermediate complexity. *Nature*, **391**, 351-356.

Gasse F (2000) Hydrological changes in the African tropics since the Last Glacial Maximum. *Quat
Sci Rev*, **19**, 189–211.

Goldstein DB, Linares AR, Cavalli-Sforza LL, Feldman MW (1995) An evaluation of genetic
distances for use with microsatellite loci. *Genetics*, **139**, 463-471.

Guo Z, Petit-Maire N, Kroepelin S (2000) Holocene non-orbital climatic events in present-day arid
areas of northern Africa and China. *Global Planet Change* v.26 p.97-103.

Hijmans RJ, Cameron SE, Parra JL, Jones PG, Jarvis A (2005) Very high resolution interpolated
climate surfaces for global land areas. *Int J Climatol*, **25**, 1965-1978.

Ho SY, Saarma U, Barnett R, Haile J, Shapiro B (2008) The Effect of Inappropriate Calibration:
Three Case Studies in Molecular Ecology. *PLoS One*, 3, e1615.

Jürgens N (1997) Floristic biodiversity and history of African arid regions. *Biodiv Conserv*, **6**, 495-
514.

Kröpelin S, Verschuren D, Lézine A-M, Eggermont H, Cocquyt C, Francus P, Cazet JP, Fagot M,
Rumes B, Russel JM, *et al.* (2008). Climate-driven ecosystem succession in the Sahara: The
past 6000 years. *Science*, **320**, 765–8.

Latinne A, Meynard CN, Herbreteau V, Waengsothorn S, Morand S, Michaux JR (2015) Influence
of past and future climate changes on the distribution of three Southeast Asian murine
rodents. *J Biogeogr*, **42**, 1714-1726.

Lebrun JP (2001) Introduction à la flore d'Afrique. 156 pp. CIRAD and Ibis Press.

Le Gall P, Silvain JF, Nel A, Lachaise D (2010) Les insectes actuels témoins des passés de

l'Afrique : essai sur l'origine et la singularité de l'entomofaune de la région afrotropicale.

Ann Soc Entomol Fr, **46**, 297-343.

Leroy T, Roux C, Villate L, Bodénès C, Romiguier J, Paiva JAP, Dossat C, Aury J-M, Plomion C,

Kremer A (2017) Extensive recent secondary contacts between four European white oak

species. *New Phytol* 214:865–878.

Lorenz MW (2009) Migration and trans-Atlantic flight of locusts. Quaternary International, **196**, 4-

12.

Lorenzen ED, Heller R, Siegismund HR (2012) Comparative phylogeography of African savannah

ungulates. *Mol Ecol*, **21**, 3656–3670.

Malek J (1997) The locusts on the daggers of Ahmose. In: E. Goring, N. Reeves and J. Riffle (eds.),

Chief of Seers: Egyptian Studies in Memory of Cyril Aldred, London, 207-219.

Maley J, Doumenge C, Giresse P, Mahé G, Philippon N, Hubau W, Lokonda MO, Tshibamba JM,

Chepstow-Lusty A (2018) Late Holocene forest contraction and fragmentation in central

Africa. *Quat Res*, **89**, 43–59.

Marin JM, Pudlo P, Estoup A, Robert CP (2018) Likelihood-free Model Choice. *Handbook of*

Approximate Bayesian Computation. Chapman and Hall/CRC.

Meinzingen WF (1993) A guide to migrant pest management in Africa. FAO, Rome, Italy.

Meynard CN, Gay PE, Lecoq M, Foucart A, Piou C, Chapuis M-P (2017) Climate-driven

geographic distribution of the desert locust during recession periods: subspecies' niche

differentiation and relative risks under scenarios of climate change. *Glob Change Biol,*

23(11), 4739-4749.

Michel AP, Sim S, Powell THQ, Taylor MS, Nosil P, Feder JL. (2010) Widespread genomic

divergence during sympatric speciation. *Proc Natl Acad Sci USA*, **107**, 9724–9729.

Miller JM, Hallager S, Monfort S, Newby J, Bishop K, Tidmus S, Black P, Houston B, Matthee C,

Robinson J, Fleischer RC (2011). Phylogeographic analysis of nuclear and mtDNA supports

subspecies designations in the Ostrich (*Struthio camelus*). *Conserv Genet*, **12**, 423–431.

Monod T (1971) Remarques sur les symetries floristiques des zones sèches nord et sud en Afrique.

Mitteilungen der Botanischen Staatssammlung München, **10**, 375-423.

Moodley Y, Russo I-RM, Robovsky J, Dalton D, Kotzé A, Smith S, Stejskal J, Ryder OA, Hermes

R, Walzer C, Bruford MW (2018) Contrasting evolutionary history, anthropogenic declines

and genetic contact in the northern and southern white rhinoceros (*Ceratotherium simum*).

Proc R Soc B, **285**, 1-9.

Nicholson SE (1996) A review of climate dynamics and climate variability in Eastern Africa. In:

Johnson TC, Odada EO (Eds.), The Limnology, Climatology and Paleoclimatology of the

East African Lakes. Gordon and Breach, Amsterdam, pp. 25-56.

Partridge TC (1997) Cainozoic environmental change in southern Africa, with special emphasis on

the last 200 000 years. *Progr Phys Geog*, **21**, 3-22.

Pedgley DE, Reynolds DR, Tatchell GM (1995) Long-range migration in relation to climate and

weather: Africa and Europe. In: Drake VA, Gatehouse AG (Eds.), Insect Migration.

Cambridge University Press, Cambridge, pp. 3–29.

Pollock DD, Bergman A, Feldman MW, Goldstein DB (1998) Microsatellite behaviour with range

constraints: parameter estimation and improved distances for use in phylogenetic

reconstruction. *Theor Popul Biol*, **53**, 256–271.

Pritchard JK, Stephens M, Donnelly P (2000) Inference of population structure using multilocus

genotype data. *Genetics*, 155, 945–959.

Pudlo P, Marin J-M, Estoup A, Cornuet J-M, Gautier M, Robert CP (2016) Reliable ABC model

choice via random forests. *Bioinformatics,* **32**, 859–866*.*

Rainey RC, Waloff Z (1951) Flying locusts and convection currents. *Anti-Locust Bull*, **9**, 51-70.

Raynal L, Marin J-M, Pudlo P, Ribatet M, Robert CP, Estoup A (2019) ABC random forests for

Bayesian parameter inference. *Bioinformatics*, **35**, 1720–1728.

Richardson, CH, Nemeth, DJ (1991) Hurricane-borne African locusts (*Schistocerca gregaria*) on

the Windward Islands. *GeoJournal*, **23**, 349–357.

Robert CP, Cornuet J-M, Marin J-M, Pillai NS (2011) Lack of confidence in approximate Bayesian

computation model choice. *P Natl Acad Sci USA*, **108**, 15112–15117.

Roberts N, Taieb M, Baker P, Damnati B, Icole M, Williamson D (1993) Timing of the Younger

Dryas event in East Africa from lake-level changes. *Nature*, **366**,146–8.

Roffey J, Magor JI (2003) Desert Locust population parameters. Desert Locust Field Research

Stations, Technical Series, 30, 29 p. FAO, Rome, Italy.

Rougemont Q, Roux C, Neuenschwander S, Goudet J, Launey S, Evanno G (2016) Reconstructing

the demographic history of divergence between European river and brook lampreys using

approximate Bayesian computations. *PeerJ*, **4**, e1910.

Rousset F (2008) GenePop'007: a complete re-implementation of the GenePop software for

Windows and Linux. *Mol Ecol Resour*, **8**, 103–106.

Roux C, Fraïsse C, Romiguier J, Anciaux Y, Galtier N, Bierne N (2016) Shedding light on the grey

zone of speciation along a continuum of genomic divergence. *PLOS Biol*, 14(12), e2000234.

Schwartz D (1992) Assèchement climatique vers 3000 B.P. et expansion Bantu en Afrique centrale

atlantique: quelques réflexions. *B Soc Geol Fr*, **163**, 353–61.

Scott L (1993) Palynological evidence for late Quaternary warming episodes in Southern Africa.

Palaeogeogr Palaeocl, **101**, 229-235.

Shi N, Dupont LM, Beug H-J, Schneider R (1998) Vegetation and climate changes during the last

21 000 years in S.W. Africa based on a marine pollen record. *Veg Hist Archaeobot*, **7**, 127-

140.

Stokes S, Thomas DSG, Washington R (1997) Multiple episodes of aridity in southern Africa since

the last interglacial period. *Nature*, **388**, 154–158.

Stute M, Talma AS (1997) Glacial temperatures and moisture transport regimes reconstructed from

noble gases and ^{18}O, Stampriet aquifer, Namibia. Proceedings of International Symposium

on Isotope Techniques in the Study of Past and Current Environmental Changes in the

Hydrosphere and the Atmosphere, Vienna, International Atomic Energy Agency.

Sun JX, Helgason A, Masson G *et al.* (2012) A direct characterization of human mutation based on

microsatellites. *Nat Genet*, **44**, 1161–1165.

Sunnåker M, Busetto AG, Numminen E, Corander J, Foll M, Dessimoz C (2013) Approximate

Bayesian computation. *PLoS Comput Biol*, **9**, e1002803.

Sword GA, Lecoq M, Simpson SJ (2010). Phase polyphenism and preventative locust management.

J Insect Physiol, **56**, 949–957.

Takezaki N, Nei M (1996) Genetic distances and reconstruction of phylogenetic trees from

microsatellite DNA. *Genetics*, **144**, 389–399.

Talma AS, Vogel JC (1992) Late Quaternary paleotemperatures derived from a speleothem from

Cango Caves. Cape Province, South Africa. *Quat. Res.*, **37**, 203-13.

Uvarov BP (1977) Grasshoppers and Locusts, vol. 2. Centre for Overseas Pest Research, London,

UK.

Van Andel TH, Tzedakis PC (1996). Palaeolithic landscapes of Europe and environs: 150,000-

25,000 years ago: an overview. *Quat Sci Rev*, **15**, 481-500.

Vansina J (1995) New Linguistic Evidence and the Bantu expansion. J Afr Hist, **36**, 173–195.

Vivo M, Carmignotto AP (2004) Holocene vegetation change and the mammal faunas of South

America and Africa. *J Biogeogr*, **31**, 943–957.

Walker MJC, Berkelhammer M, Björck S, Cwynar LC, Fisher DA, Long AJ, Lowe JJ, Newnham

RM, Rasmussen SO, Weiss H (2012) Formal subdivision of the Holocene Series/Epoch: a

Discussion Paper by a Working Group of INTIMATE (Integration of ice-core, marine and

terrestrial records) and the Subcommission on Quaternary Stratigraphy (International

Commission on Stratigraphy). *J Quat Sci*, **27**, 649–659

Waloff Z (1966) The upsurges and recessions of the desert locust plague: an historical survey. *Anti-*

Locust Memoir, **8**, 111 p.

Waloff Z (1976) Some temporal characteristics of desert locust plagues. *Anti-Locust Memoir*, **13**, 1–

36.

Waloff Z, Pedgley DE (1986) Comparative biogeography and biology of the South American

locust, *Schistocerca cancellata* (Serville), and the South African desert locust, *S. gregaria*

flaviventris (Burmeister) (Orthoptera: Acrididae): a review. *Bull Entomol Res*, **76,** 1-20.

Weir BS (1996) Genetic Data Analysis II. Sinauer Associates, Sunderland, Massachusetts.

Zhivotovsky LA, Feldman MW, Grishechkin SA (1997) Biased mutations and microsatellite

variation. *Mol Biol Evol*, **14**, 926–933.

 Supporting Material

 Additional supporting information may be found in the online version of this article.

Figure legends

Figure 1. **Present time distribution range of *Schistocerca gregaria* in Africa under remission**

periods with winds in August a) and January b), and vegetation habitats suitable for the

species during the present period b), the Holocene Climatic Optimum (HCO, 9 to 6 Ky ago)

d), the Younger Dryas (YD, 12.9 to 11.7 Ky ago) e) and the Last Glacial Maximum (LGM, 26

to 14.8 Ky ago) f).

(a-b) Distribution range and winds are adapted from Sword *et al.* (2010) and Nicholson (1996),

respectively. (c-f) Vegetation habitats are adapted from Adams and Faure (1997). Open vegetation

habitats suitable for the desert locust correspond to deserts (dark orange), xeric shrublands (light

orange) and tropical - Mediterranean grasslands (pink). Other unsuitable habitat classes (white) are

forests, woodlands and temperate shrublands and savannas.

Figure 2. Evolutionary scenarios compared using ABC-RF.

The subscripts *g*, *f* and *a* refer to the subspecies *S. g. gregaria*, *S. g. flaviventris* and their unsampled

common ancestor, respectively. Twelve scenarios are considered and identified by an acronym.

Such scenarios differ by the presence or absence of three evolutionary events: a bottleneck in *S. g.*

flaviventris (B) right after divergence between the two subspecies, a population size contraction in

the ancestral population (C) and a discrete genetic admixture event either unidirectional from *S. g.*

gregaria into *S. g. flaviventris* (A_U) or bidirectional (A_B) . For convenience, parameters associated

with an evolutionary event are represented graphically only in the first shown scenario that include

this event. Looking forward in time, time periods are t_C, the time of ancestral population size

contraction, t_S, the time of split between the two subspecies, and t_A, the time of the genetic

admixture between subspecies (with $t_C > t_S > t_A$). r_{AU} is the unidirectional admixture rate, *i.e.* the

proportion of genes from the *S. g. gregaria* lineage entering the *S. g. flaviventris* population at time

t_A. r_{AB} is the bidirectional admixture rate, *i.e.* the proportion of genes exchanged between the *S. g.*

gregaria lineage and the *S. g. flaviventris* lineage at time t_A. N_g, N_f and N_a are the stable effective

population sizes of *S. g. gregaria*, *S. g. flaviventris* and the ancestor, respectively. N_C is the effective

population size during the contraction event of duration d_C in the ancestor. N_B is the effective

population size during the bottleneck event of duration d_B.

Figure 3. Divergence time between *S. g. gregaria* and *S. g. flaviventris* inferred under the best

supported scenario (scenario SCB) a) in relation to bioclimatic changes in Northern b) and

Southern Africa c).

a) Dashed and solid lines represent the formal subdivision of the Holocene and Pleistocene epochs

(Walker at al. 2012). Dotted lines with labels on the right side are the median value and 90%

credibility interval of the posterior density distributions of the divergence time (assuming an

average of three generations per year; Roffey and Magor 2003). Asterisks refer to earliest

archeological records of the desert locust. In the Algerian Sahara, remains of locusts were found in

a special oven dating back to about 6Ky ago, in the rock shelter of Tin Hanakaten (Aumassip 2002).

In Egypt, locusts were depicted on daggers of the pharaoh Ahmose, founder of the Eighteenth

Dynasty (about 3.5 Ky ago) (Malek 1997) and, at Saqqara, on tombs of the Sixth Dynasty (about

4.2 to 4.4 Ky ago) that is thought to have felt with the impact of severe droughts (Meinzingen

1993). b-c) Climatic episodes include major cycles and additional transitions of aridity (sandy

brown) and humidity (steel blue). The grey coloration means that there is debate on the climatic

status of the period (arid vs. humid). HCO: Holocene Climatic Optimum; YD: Younger Dryas;

LGM: Last Glacial Maximum; LIG: Last Inter Glacial. Delimitations of climatic periods were based

on published paleoclimatic inferences from geological sediment sequences (*e.g.,* eolian deposition,

oxygen isotope data) and biological records (*e.g.*, pollen or insect fossils assemblages) from marine

cores or terrestrial lakes. References are Bond *et al.* (1997), Guo *et al.*(2000), Kröpelin *et al.*

(2008), Roberts *et al.* (1993) and van Andel and Tzedakis (1996) for northern Africa, and Talma

and Vogel (1992), Stokes *et al.* (1997), and Shi *et al.* (1998) for southern Africa. See also Gasse

(2000) for a review.

Figure 4. Median and 90% credibility interval a) and local accuracy b) of ABC-RF posterior

distributions of the divergence time under the best supported scenario (scenario SCB) as a

function of the time scale.

Simulated pseudo-observed datasets (5,000 per divergence time) were generated for fixed

divergence time values of 100 ; 250 ; 500; 1,000 ; 2,500; 5,000 ; 10,000 ; 25,000 ; 50,000; 100,000;

and 250,000 generations (cf. x-axis with a log-scale). a) The estimated median (plain lines) and

90% credibility interval (90% CI; dashed lines), averaged over the 5,000 datasets, are represented

(y-axis). b) The local accuracy is measured using out-of-bag predictions and the normalized mean

absolute error (NMAE). See Supplementary Material S3 for details.

Tables

Table 1. Summary statistics provided by DIYABC with corresponding values computed from the observed microsatellite dataset.

	Summary statistics	Observed values at 14 untranscribed markers	Observed values at 9 transcribed markers
S. g. gregaria **one-sample statistics**	NAL	28.8	15.8
	HET	0.92	0.79
	VAR	36.1	13.3
	MGW	0.92	0.86
S. g. flaviventris **one-sample statistics**	NAL	23.4	14.4
	HET	0.86	0.69
	VAR	33.4	16.7
	MGW	0.96	0.95
Two-samples statistics	FST	0.04	0.12
	DAS	0.07	0.16
	$LIK_{Sgg \to Sgf}$	3.61	2.82
	$LIK_{Sgf \to Sgg}$	3.20	2.55
	DM2	22.7	12.4
	N2P	35.0	21.1
	H2P	0.91	0.79
	V2P	40.2	18.2

The names of summary statistics are those given in the program DIYABC (Cornuet *et al.* 2014). *NAL*: mean number of alleles; *HET*: mean expected heterozygosity; *VAR*: variance of allele sizes in base pairs; *MGW*: M index of Garza and Williamson (2001); *FST*: pairwise differentiation estimator of Weir and Cockerham (1984); *DAS*: shared allele distance (Chakraborty and Jin 1993); *LIK*: the mean index of classification (Rannala and Moutain, 1997; Pascual *et al.* 2007); *DM2*: distance of Golstein *et al.* (1995). *N2P*, *H2P* and *V2P*: *NAL*, *HET* and *VAR* statistics computed after pooling the two population samples. Note that five "noise variables", randomly drawn into uniform distributions bounded between 0 and 1, and denoted NOISE1 to NOISE5 in the concerned illustrations, were added to the set of summary statistics processed by RF, in order

to evaluate which summary statistics of our genetic datasets were informative in our different inferential ABC-RF settings, when conducting scenario choice or parameter estimation. Such noise variables do not alter ABC-RF inferences (see Marin *et al.* 2018; Raynal *et al.* 2019).

Table 2. Scenario choice when analyzing groups of scenarios or scenarios separately.

	Analyses of groups of scenarios			Analysis of scenarios separately
	Group 1 = no B *vs.* group 2 = B	Group 1= no C *vs.* group 2 = C	Group 1 = no A *vs.* group 2 = A$_S$ vs. group 3 = A$_U$	
Prior error rate	19.0%	26.1%	32.1%	58.4%
Posterior probability (selected group or individual scenario)	0.926 (B)	0.682 (C)	0.700 (no A)	0.564 (scenario SCB)

Scenarios were grouped based on the presence or not of a bottleneck in *S. g. flaviventris* (B or no B), a population size contraction in the ancestor (C or no C) and a symmetrical (i.e., bidirectional) or an asymmetrical genetic admixture from *S. g. gregaria* into *S. g. flaviventris* (A$_B$ or A$_U$ or no A). We reported values for prior error rates and posterior probabilities of the selected group of scenarios or individual scenario, averaged over ten replicate analyses. The local posterior error rate (corresponding to a confidence measure of the selected scenario given the observation.) can be computed as 1 minus the posterior probability (see Supplementary material S3 for details). The number of records for each reference datasets simulated from DIYABC was set to 150,000 and the number of RF- trees was 3,000.

Table 3. Parameter estimation under the best supported scenario (scenario SCB).

	Posterior values			Prior values		
	Median	90% CI	NMAE	Median	90% CI	NMAE
t_S	7,723	2,785 – 19,708	0.369	1,212	124 – 73,795	0.359
t_C / t_S	2.75	1.11 – 35.47	0.596	12.17	1.24 – 762.26	1.077
N_f / N_g	5.43	0.52 – 25.56	1.332	1.00	0.12 – 8.11	1.726
d_B / N_B	1.06	0.49 – 2.41	0.323	1.00	0.13 – 7.57	0.299

t_S: time of divergence between the two desert locust subspecies (in number of generations); t_C: time of ancestral population size contraction; N_f : stable effective population size of *S. g. flaviventris*; N_g: stable effective population size of *S. g. gregaria;* d_B : duration of the bottleneck event; N_B: effective population size during the bottleneck event. For each evolutionary parameter, we reported posterior point estimates averaged over ten replicate analyses. CI: credibility interval. The number of records for each reference datasets simulated from DIYABC was set to 100,000 and the number of RF-trees was 2,000. Accuracy has been measured with the normalized mean absolute error (NMAE) which corresponds to the mean of the absolute difference between the point estimate of the parameter (here the median) and the (true) simulated value divided by the (true) simulated value. NMAE measures were computed using out-of-bag predictions either on the whole data space defined by the prior distributions (prior NMAE) or conditionally to the observed dataset (posterior NMAE); see Supplementary material S3 for details.

Fig. 1

a) Present distribution
(winds in Aug)

b) Present distribution
(winds in Jan)

c) Present habitats

Schistocerca gregaria gregaria

Schistocerca gregaria flaviventris

d) HCO habitats

e) YD habitats

f) LGM habitats

Legend: Desert   Xeric shrubland   Tropical and Mediterranean grassland

Fig. 2

Fig. 3

Fig. 4

Supplementary material online for the manuscript: « A young age of subspecific divergence in the desert locust *Schistocerca gregaria*"

Marie-Pierre Chapuis[1,2], Louis Raynal[3,4], Christophe Plantamp[5], Christine N. Meynard[6], Laurence Blondin[7], Jean-Michel Marin[3,4] and Arnaud Estoup[4,6]

[1] CIRAD, CBGP, Montpellier, France.

[2] CBGP, CIRAD, INRA, IRD, Montpellier SupAgro, Univ Montpellier, Montpellier, France.

[3] IMAG, Univ de Montpellier, CNRS, Montpellier, France.

[4] Institut de Biologie Computationnelle (IBC), Montpellier, France.

[5] ANSES, Laboratoire de Lyon, France.

[6] CBGP, INRA, CIRAD, IRD, Montpellier SupAgro, Univ Montpellier, Montpellier, France.

[7] CIRAD, BGPI, Montpellier, France.

<u>**Outline**</u>:

Supplementary material S1. Details on results from ABC Random Forest (ABC-RF) treatments.

Supplementary material S2. Hindcasting the desert locust potential distribution into the mid-Holocene (HCO) and the last glacial maximum (LGM) using distribution modelling.

Supplementary material S3. Overview of the used ABC Random Forest (ABC-RF) methods.

Supplementary material S4. Details on results from ABC-RF treatments when assuming uniform priors for the three time period parameters of the studied scenarios.

Supplementary material S1. Details on results from ABC-RF treatments.

Table S1.1. Scenario choice for each of the ten replicate analyses.

Reference table	Best scenario	Votes S	Votes SC	Votes SB	Votes SCB	Votes SA_U	Votes SCA_U	Votes SBA_U	Votes $SCBA_U$	Votes SA_B	Votes SCA_B	Votes SBA_B	Votes $SCBA_B$	Prior error rate	Posterior probability (best scenario)
1	scenario SCB	0.004	0.033	0.171	0.513	0.011	0.022	0.040	0.094	0.017	0.041	0.024	0.031	0.583	0.572
2	scenario SCB	0.003	0.026	0.178	0.515	0.006	0.018	0.023	0.093	0.023	0.022	0.051	0.042	0.583	0.549
3	scenario SCB	0.003	0.024	0.196	0.497	0.010	0.019	0.030	0.101	0.025	0.033	0.036	0.024	0.583	0.545
4	scenario SCB	0.004	0.026	0.144	0.483	0.006	0.018	0.043	0.126	0.041	0.032	0.050	0.026	0.586	0.570
5	scenario SCB	0.004	0.025	0.126	0.537	0.007	0.017	0.030	0.117	0.017	0.021	0.063	0.036	0.582	0.607
6	scenario SCB	0.004	0.039	0.161	0.501	0.004	0.023	0.036	0.128	0.020	0.026	0.037	0.021	0.585	0.568
7	scenario SCB	0.014	0.021	0.136	0.536	0.005	0.022	0.029	0.099	0.022	0.044	0.037	0.035	0.585	0.588
8	scenario SCB	0.005	0.016	0.149	0.491	0.005	0.019	0.026	0.124	0.042	0.030	0.034	0.058	0.586	0.565
9	scenario SCB	0.003	0.036	0.151	0.506	0.009	0.019	0.017	0.113	0.049	0.024	0.034	0.040	0.584	0.551
10	scenario SCB	0.006	0.031	0.178	0.491	0.002	0.029	0.028	0.082	0.041	0.032	0.033	0.047	0.585	0.528
Mean	**scenario SCB**	**0.005**	**0.028**	**0.159**	**0.507**	**0.007**	**0.021**	**0.030**	**0.108**	**0.030**	**0.030**	**0.040**	**0.036**	**0.584**	**0.564**

We report values for the proportion of votes, prior error rates and posterior probabilities of the best scenario on ten replicate analyses based on ten different reference tables. Scenarios are depicted in Figure 2. For each reference table, the number of datasets simulated using DIYABC was set to 150,000 and the number of RF-trees was 3,000. The

scenario SCB was the best supported for all replicate analyses: it involves a bottleneck event in *S. g. flaviventris* right after divergence, a population size contraction in the ancestral population and not any genetic admixture event.

Table S1.2. Estimation of the divergence time between *S. g. gregaria* and *S. g. flaviventris* for the ten replicate analyses processed under the best supported scenario (scenario SCB).

t_{div} (*G*)	Median	q5%	q95%
reference table 1	7440.0	2485.0	19380.0
reference table 2	8257.1	2668.0	21086.0
reference table 3	7930.3	2771.0	20310.9
reference table 4	7301.0	2888.0	19639.5
reference table 5	7598.6	2376.0	18260.6
reference table 6	7426.0	2975.7	19704.7
reference table 7	7776.0	3190.0	19290.2
reference table 8	7960.0	2812.0	19664.2
reference table 9	7552.3	2717.0	20685.0
reference table 10	7991.0	2966.9	19060.7
Mean	**7723.2**	**2785.0**	**19708.2**
SD	**307.3**	**240.5**	**817.8**

Replicate analyses have been processed on different reference tables. For each reference table, the number of datasets simulated using DIYABC was set to 100,000 and the number of RF-trees was 2,000. Divergence times are given in number of generations (G). SD stands for standard deviations computed from the ten values of median, 5% quantile and 95% quantile estimated from the ten replicate analyses.

Table S1.3 Effect of the number of simulated datasets in the reference table on scenario choice.

n_{ref}	50,000		80,000		100,000		130,000		150,000	
	Mean	**SD**	**Mean**	**SD**	**Mean**	**SD**	**Mean**	**SD**	**Mean**	**SD**
Votes S	0.006	0.002	0.005	0.002	0.006	0.003	0.006	0.004	0.005	0.003
Votes SC	0.030	0.015	0.031	0.009	0.029	0.006	0.028	0.006	0.028	0.007
Votes SB	0.155	0.025	0.155	0.028	0.165	0.033	0.164	0.036	0.159	0.022
Votes SCB	0.496	0.035	0.506	0.036	0.500	0.034	0.502	0.025	0.507	0.019
Votes SA$_U$	0.006	0.003	0.007	0.004	0.006	0.004	0.006	0.004	0.007	0.003
Votes SCA$_U$	0.025	0.005	0.023	0.004	0.020	0.005	0.022	0.006	0.021	0.008
Votes SBA$_U$	0.033	0.009	0.031	0.007	0.033	0.008	0.030	0.009	0.030	0.016
Votes SCBA$_U$	0.117	0.027	0.113	0.019	0.108	0.017	0.109	0.017	0.108	0.012
Votes SA$_B$	0.031	0.009	0.030	0.011	0.027	0.010	0.029	0.010	0.030	0.008
Votes SCA$_B$	0.032	0.012	0.029	0.007	0.029	0.009	0.031	0.012	0.030	0.011
Votes SBA$_B$	0.039	0.014	0.039	0.013	0.043	0.014	0.042	0.013	0.040	0.011
Votes SCBA$_B$	0.030	0.013	0.031	0.009	0.032	0.009	0.032	0.009	0.036	0.011
Prior error rate	**0.594**	**0.002**	**0.589**	**0.002**	**0.587**	**0.001**	**0.585**	**0.001**	**0.584**	**0.001**
Posterior probability (best model)	**0.567**	**0.037**	**0.562**	**0.034**	**0.555**	**0.034**	**0.563**	**0.028**	**0.564**	**0.023**

Scenarios are depicted in Figure 2. The number of records in the reference datasets (n_{ref}) simulated from DIYABC varied from 50,000 to 150,000. We report mean and standard deviation values for the proportion of votes for each scenario, and for prior error rates and posterior probabilities of the best scenario for ten replicate analyses. Replicate analyses have been processed on different reference tables. The number of RF-trees was 3,000. The scenario SCB was the best supported for all replicate analyses: it involves a bottleneck event in *S. g. flaviventris* right after divergence, a population size contraction in the ancestral population and not any genetic admixture event.

Table S1.4. Effect of the number of simulated datasets in the reference table on posterior point estimation values (A) and estimation accuracy (B) of the divergence time between *S. g. gregaria* and *S. g. f laviventris* under the best supported scenario (scenario SCB).

(A)

n_{ref}		50,000		80,000		90,000		100,000	
		Mean	**SD**	**Mean**	**SD**	**Mean**	**SD**	**Mean**	**SD**
Posterior estimations	Median	7731.8	448.9	7691.8	357.5	7724.8	318.0	7723.2	307.3
	q5%	2697.0	129.7	2706.8	235.2	2764.4	172.9	2785.0	240.5
	q95%	20295.5	1763.6	19711.6	1451.3	19508.7	1264.1	19708.2	817.8

(B)

n_{ref}		50,000	80,000	90,000	100,000
Accuracy measures	Prior NMAE	0.378	0.365	0.362	0.359
	Posterior NMAE	0.375	0.370	0.365	0.369

The number of records in the reference datasets (n_{ref}) simulated from DIYABC varied from 50,000 to 100,000. The number of RF-trees was set to 2,000. (a) Replicate analyses have been processed on ten reference tables. (b). The normalized mean absolute error (NMAE) is the absolute difference between the point estimate (here the median) and the (true) simulated value divided by the (true) simulated value.

Figure S1.1. Projection on a single (when analyzing pairwise groups of scenarios) or on the first two LDA axes (when analyzing the twelve scenarios separately) of the observed dataset and the simulated datasets recorded in the reference table. Colors correspond to group of scenarios or individual scenarios. The location of the desert locust observed dataset is indicated by a vertical black line or a star. Scenarios were grouped based on the presence or not of a bottleneck in *S. g. flaviventris* (B or no B), a population size contraction in ancestor (C or no C) and a genetic admixture either unidirectional from *S. g. gregaria* into *S. g. flaviventris* or bidirectional (A_U or A_B or no A). When considering the whole set of twelve scenarios separately (d), the projected points substantially overlapped for at least some of the scenarios. This suggests an overall low power to discriminate among scenarios considered. Conversely, considering pairwise groups of scenarios, one can observe a weaker overlap of projected points (at least for (a) and (b)) suggesting a stronger power to discriminate among groups of scenarios of interest than when considering all scenarios separately. One can note that the location of the observed dataset (indicated by a vertical line) suggests an association with the scenario group with a bottleneck event in *S. g. flaviventris* and with the scenario group with a population size contraction in the ancestral population.

(a) Group g1= no B *vs.* group g2 = B

(b) Group g1 = no C *vs.* group g2 =C

(c) Scenario group 1 = no A vs. group 2 = A_U vs. group 3 = A_B **(d)** All twelve scenarios considered separately

Figure S1.2. Contributions of ABC-RF summary statistics when choosing between groups of scenarios. The contribution of each 32 summary statistics and one LDA axis (or two LDA axes) is evaluated as the total amount of decrease in the Gini criterion (variable importance on the x-axis). The higher the contribution of the statistics, the more informative it is in the inferential process. The microsatellite set and subspecies sample are indicated at the end of each statistics by indices k_i for single population statistics and $k\_i.j$ for two population statistics, with $k=1$ for the set of untranscribed microsatellites or $k=2$ for the set of transcribed microsatellites, and $i(j)=1$ for the *S. g. flaviventris* subspecies or and $i(j)=2$ for the *S. g. gregaria* subspecies. See Table S1 for details on the summary statistics abbreviations. Five noise variables, randomly drawn into uniform distributions bounded between 0 and 1, and denoted NOISE1 to NOISE5 were added to the set of summary statistics processed by RF, in order to evaluate from which amount of decrease in the Gini criterion the summary statistics computed from our genetic datasets were not informative anymore (indicated by a red star).

(a) Group g1= no B *vs.* group g2 = B   (b) Group g1 = no C *vs.* group g2 =C   (c) Group 1 = no A vs. group 2 = AU vs. group 3 = AB

Figure S1.3. Contributions of ABC-RF summary statistics when choosing among the twelve individual scenarios. The contribution of each 32 summary statistics and eleven LDA axes is evaluated as the total amount of decrease in the Gini criterion (variable importance on the x-axis). The higher the contribution of the statistics, the more informative it is in the inferential process. The microsatellite set and subspecies sample are indicated at the end of each statistics by indices k_i for single population statistics and $k\_i.j$ for two population statistics, with $k=1$ for the set of untranscribed microsatellites or $k=2$ for the set of transcribed microsatellites, and $i(j)=1$ for the *S. g. flaviventris* subspecies or and $i(j)=2$ for the *S. g. gregaria* subspecies. See Table 1 for details on the summary statistics abbreviations. Five noise variables, randomly drawn into uniform distributions bounded between 0 and 1, and denoted NOISE1 to NOISE5 were added to the set of summary statistics processed by RF, in order to evaluate from which amount of decrease in the Gini criterion the summary statistics computed from our genetic datasets were not informative anymore (indicated by a red star).

Figure S1.4. Contributions of ABC-RF summary statistics when estimating the divergence time between the two desert locust subspecies under the best supported scenario (scenario SCB). The contribution of each 32 summary statistics is evaluated as the total amount of decrease of the residual sum of squares, divided by the number of trees, (variable importance on the x-axis). The higher the contribution of the statistics, the more informative it is in the inferential process. The microsatellite set and subspecies sample are indicated at the end of each statistics by indices k_i for single population statistics and $k\_i.j$ for two population statistics, with $k=1$ for the set of untranscribed microsatellites or $k=2$ for the set of transcribed microsatellites, and $i(j)=1$ for the *S. g. flaviventris* subspecies or and $i(j)=2$ for the *S. g. gregaria* subspecies. See Table S6.1 for details on the summary statistics abbreviations. Five noise variables, randomly drawn into uniform distributions bounded between 0 and 1, and denoted NOISE1 to NOISE5 were added to the set of summary statistics processed by RF, in order to evaluate from which amount of decrease in the variable importance criterion the summary statistics computed from our genetic datasets were not informative anymore (indicated by a red star).

Figure S1.5. Effect of the number of RF-trees for scenario choice. We here illustrate the effect of the number of trees in the forest on the prior error rate using an informed mutational prior setting and considering the twelve compared scenarios separately. The number of datasets in the reference table simulated using DIYABC was 150,000. The shape of the curve shows that the prior error rate stabilizes for a number of RF-trees > 2,000.

Figure S1.6. Effect of the number of RF-trees for parameter estimation. We here illustrate the effect of the number of trees in the forest on the RF mean square error of the divergence time between *S. g. gregaria* and *S. g. flaviventris* under the selected scenario SCB and using an informed mutational prior setting. The number of datasets in the reference table simulated using DIYABC was 100,000. The shape of the curve shows that the prior error rate stabilizes for a number of RF-trees > 1,500.

Supplementary material S2. Hindcasting the desert locust potential distribution into the mid-Holocene (HCO) and the last glacial maximum (LGM) using distribution modelling.

To complement distribution projections based on vegetation during the mid-Holocene (HCO; ~6,000 years ago) and Last Glacial Maximum (LGM; ~120,000 years ago) periods, we used species distribution modelling along with climatic reconstructions of the past. However, results from this modelling effort do not fully match vegetation reconstructions presented in the main text. Given that most Global Climate Models (GCMs) have been calibrated and tested largely with northern hemisphere data, uncertainties related to current and past extrapolations of climate over large areas of Africa are often unknown (e.g. Rowell et al. 2016). In this appendix, we present the results of the distribution modelling effort, while in the main text we base our formalization of evolutionary scenarios on paleo-vegetation reconstructions. Notice that the difference between the two approaches does not affect our major conclusions. Indeed, since paleo-vegetation maps show more severe changes of open vegetation distributions during glacial and interglacial periods, their use make us consider a wider spectrum of key biogeographic events than we would have identified based on climatic modelling (e.g. continuous colonization of southern Africa, population size contraction), and thus a larger set of possible scenarios. In addition, our inference of a most likely scenario including a long-distance migration event, and our estimation of a short time scale of divergence between the two clades, do not conflict with the results of any of these approaches (see also the section '*On the influence of climatic cycles*' in Discussion of the main text).

Methods

In a previous study (Meynard et al. 2017), we used Climond data (Kriticos et al. 2012) to characterize the current distribution of the two desert locust subspecies, and project their potential fate under different climate change scenarios into 2070. Here, we could not use the same models because the climatic variables used to calibrate those models, in particular radiation and wetness indices, are not available into the mid-Holocene (HCO) and Last Glacial Maximum (LGM) periods. However, we apply here a similar modelling procedure with a different subset of environmental predictors. Instead of Climond, here we downloaded Worldclim v1.4 data (Hijmans et al. 2005) at a 5 min resolution (approximately 9 km at the equator).

For hindcasting, we first need to calibrate a model under current conditions, which can then be used to draw maps of potential ranges under HCO and LGM conditions. To do so, we used the occurrence data published in Meynard et al. (2017), which included decades of monitoring of *S. g. gregaria* in northern Africa, the middle east and southern Asia during remission periods, as well as field records and literature records of *S. g. flaviventris* in southern Africa. However, rather than modelling each clade independently as we did before, here we decided to group presence records for the two clades as a single unit for the following reasons: (1) here the emphasis is in paleo-climates, at a time scale where the two clades were not yet supposed to be differentiated; (2) a niche analysis in Meynard et al. (2017) did not show evidence of niche differentiation between *S. g. gregaria* and *S. g. flaviventris*, suggesting that

the environment occupied by the southern clade is a subset of that occupied by the northern one; and (3) the biogeographic hypothesis supported by the molecular analysis presented here involves the colonization of the southern tip of south Africa by an isolated population of the northern sub-species, which is more generalist in terms of its environmental niche. We therefore considered that grouping occurrence records regardless of the subspecies would provide for a more accurate hindcasting scenario. However, preliminary analysis using separate modelling for each subspecies showed qualitatively similar projections when using *S. g. gregaria* alone (results not shown).

Finally, since here the emphasis is on Africa, we included only occurrence records in this continent, and used 10,000 pseudo-absences drawn at random from outside the combined current ranges of the two clades by using the distribution maps proposed in Meynard et al. (2017) to mask out areas available for pseudo-absence selection. We also assumed here that most of the African continent was accessible to the species during this geological time-scale, given the long known dispersal distances during outbreak periods, the wide distribution of the northern clade, and the fact that the species is present in both extremes of the continent. Consequently, here we do not delimit an accessible area, but we limit our modelling to continental Africa.

To model the species distribution we used four climatic variables to represent mean and variability in temperature and precipitation, while minimizing correlation between them. The four variables selected using these criteria were annual mean temperature (BIO1), annual precipitation (BIO12), annual temperature range (BIO7) and precipitation seasonality (BIO15). We also tested the same forward variable selection procedure presented in Meynard et al. (2017), which provided very similar predictions but, for simplicity, we will not present those results here. Using these four climatic predictors, we applied three statistical models in Rv3.6.1 to draw a consensus map for all predictions: Generalized Additive Model (GAM) using library *mgcv*, and limiting the maximum model complexity to k=4 (Wood 2006); boosted regression trees (BRT) using the library *gbm*, with 2000 trees and no interactions (Elith et al. 2008); and MAXENT, using the library *dismo* and all default parameters (Elith et al. 2011). As mentioned above, background data was sampled at random in Africa from outside the species current distribution. Pseudo-absences received a combined weight of 50% when using GAM or BRT, in order to balance the large number of pseudo-absences with respect to presence records in the modelling process while representing the range of values of the environmental predictors in the study area (Barbet-Massin et al. 2012). The consensus was drawn as the median predicted probability between the three models, and four different threshold values were calculated from this consensus: the MST threshold (the threshold that maximizes the sum of sensitivity + specificity), often recommended as a good strategy to optimize presence and absence classification rates (Liu et al. 2005), was calculated to delimit the predicted range of the species and calculate classification rate statistics; and 3 different thresholds that represent the highest predicted values that include a fixed percentage of presences (50%, 75% and 90%). These thresholds were used to visualize on the maps the core (50% threshold) versus the more marginal (between 75% and 90% thresholds) habitats within the range. In this context, the 50% threshold represents the core distribution, i.e. the highest predicted values that contain more than half of the occurrences, the 90% threshold represents the area that includes most of the occurrences, and the area outside the 90% threshold represents marginal or no habitat, where the species is unlikely to occur. The area between the

75% and 90% threshold can be considered as habitat that is rather marginal but that is still part of the predicted range.

Classification success rates (AUC, TSS, sensitivity and specificity) were calculated for the consensus on all the dataset, using an MST threshold when needed. The thresholds calculated using current climate and occurrence data were then projected into past scenarios (HCO and LGM) using 3 different global circulation models at the same resolution: CCSM4, MIROC-ESM and MPI-ESM-P. We chose these three GCMs because their projections were available for both time steps of interest. Although the three GCMs are based on climatological principles, each one works in slightly different ways, and therefore produces different results. Consensus between GCMs presented below represent median values of predictions between the three GCMs.

Results

The current distribution of *S. gregaria*, as modelled using both subclades as a single unit and limited to continental Africa, is quite similar to the one already published in Meynard et al. (2017) (see Figure S2.1). There are, however, differences in the outline of the distributions, but the overall predicted area is largely overlapping and classification rates were excellent for this model under current conditions (% correctly classified = 87%; AUC = 0.960 ± 0.003; sensitivity = 0.911 ± 0.014; specificity = 0.870 ± 0.002; TSS = 0.788 ± 0.016). As expected, the species is more likely to occur in regions with mid to high temperatures, wide annual temperature range, low precipitation and high precipitation seasonality (Figure S2.2).

Figure S2.1: Current range as predicted by a model combining both subspecies' occurrences and using four climatic predictors. The range predicted by this model (black contour) is compared to the one published in Meynard et al. (2017) using a different set of predictors and separating occurrences by subspecies (blue contour).

Figure S2.2: Response curves of S. gregaria against the four environmental predictors used in the modelling process, as derived from the GAM model. Notice that temperature is expressed as the real temperature x 10.

To better understand the results in terms of distributional changes, Figure S2.3 shows a comparison of the climatic variables used in the modelling process in the study region during the present, the HCO and the LGM periods across Africa. Although only one GCM scenario is shown (CCSM4), climatic conditions for the other two GCMs present similar trends. Notice that, although temperatures are thought to have been warmer than present during the HCO globally, those changes are not uniform across regions and are not reflected in these HCO scenarios for Africa. In all the scenarios in Figure S2.3, mean temperature showed a slight decrease across Africa, while precipitation increased slightly.

Figure S2.3: Comparison of climates across times (current climate, mid-Holocene, Last Glacial Maximum=LGM). Variables shown are those used in the modelling process. BIO1 = Mean Annual Temperature; BIO7 = Annual Temperature Range; BIO12 = Annual Precipitation; BIO15 = Precipitation Seasonality (Coefficient of Variation). Vertical color lines indicate mean values for each period: blue = current mean; red = HCO mean; green = LGM mean.

However, when looking at the distribution of those changes (Figure S2.4) some regions showed a decrease and others an increase in temperatures during the same period. Most of Africa shows changes that represent less than 10% of their current mean temperature values (Figure S2.4, upper left). Overall, conditions under HCO in Africa were therefore similar to current climatic conditions in terms of temperature but were moister in many regions that are relevant to the potential distribution of the desert locust (Figure S2.4, upper row).

Figure S2.4: Relative changes in annual mean temperature (BIO1) and annual precipitation (BIO12) during the HCO and LGM, expressed as a percent of current values per grid cell. Positive values therefore indicate an increase whereas negative values represent a decrease with respect to current climatic conditions.

During the LGM (Figure S2.3, right column), mean temperature decreased more dramatically throughout all of Africa, while precipitation increased in some regions but decreased in others (Figure S2.4, lower row). While the entire continent experienced a decrease in temperature, changes in precipitation were more heterogeneous (Figure S2.4, lower right). Given that these changes are heterogeneous across regions, changes in the potential distribution of *S. gregaria* are less dramatic than one would expect given global simplifications of these changes.

Indeed, the predicted distribution during the HCO and the LGM are very similar to the current distribution (Figure S2.5). There is an overall shrinkage of the core habitat (in red in all maps), but the contour of the distribution (black contour) remains large and mostly overlapping with the current distribution.

Figure S2.5: Comparison of current versus past potential distribution of S. gregaria during the HCO and LGM periods.

Although each GCM resulted in differences in terms of the projected distribution maps of *S. gregaria*, in the consensus distribution is projected to be very similar to the current distribution during the HCO, and slightly reduced during the LGM, with a shift of the core favorable climatic conditions towards a much reduced area, especially during the LGM (Figure S2.5).

Although climatic projections for each GCM result in different projected distributions during the HCO (Figure S2.6) and the LGM (Figure S2.7), overall the tendencies are the same: a similar distribution when comparing present and HCO conditions, and a more restricted potential distribution when considering LGM conditions.

Figure S2.6: Projected potential distribution of S. gregaria during the mid-Holocene (~6000 years BP) using three different Global Circulation Models (GCMs): CCRM4, MIROC-ESM and MPI-ESM-P.

Figure S2.7: Projected potential distribution of S. gregaria during the LGM (~20,000 years BP) using three different Global Circulation Models (GCMs): CCRM4, MIROC-ESM and MPI-ESM-P.

References

Barbet-Massin M, Jiguet F, Albert CH, Thuiller W. 2012. Selecting pseudo-absences for species distribution models: how, where and how many? Methods Ecol. Evol. DOI: 10.1111/j.2041-210X.2011.00172.x.

Elith J, Leathwick JR, Hastie T. 2008. A working guide to boosted regression trees. J. Anim. Ecol. 77:802–813.

Elith J, Phillips SJ, Hastie T, Dudik M, Chee YE, Yates CJ. 2011. A statistical explanation of MaxEnt for ecologists. Divers. Distrib. 17:43–57.

Hijmans RJ, Cameron SE, Parra JL, Jones PG, Jarvis A. 2005. Very High Resolution interpolated climate surfaces for global land areas. Int. J. Climatol. 25:1965–1978.

Kriticos DJ, Webber BL, Leriche A, Ota N, Macadam I, Bathols J, Scott JK. 2012. CliMond: global high-resolution historical and future scenario climate surfaces for bioclimatic modelling. Methods Ecol. Evol. 3:53–64.

Liu C, Berry PM, Dawson TP, Pearson RG. 2005. Selecting thresholds of occurrence in the prediction of species distributions. Ecography 28:385–393.

Meynard CN, Gay P-E, Lecoq M, Foucart A, Piou C, Chapuis M-P. 2017. Climate- driven geographic distribution of the desert locust during recession periods: Subspecies' niche differentiation and relative risks under scenarios of climate change. Glob. Change Biol. 23:4739–4749.

Rowell DP, Senior CA, Vellinga M, Graham RJ. 2016. Can climate projection uncertainty be constrained over Africa using metrics of contemporary performance? Clim. Change 134:621–633.

Wood SN. 2006. Generalized Additive Models: an introduction with R. Boca Raton, Florida: Chapman & Hall/CRC

Supplementary Material S3: Overview of the used ABC Random Forest (ABC-RF) methods

In this supplementary material, we provide readers with an overview of the Approximate Bayesian Computation Random Forest (hereafter ABC-RF) methods used in the present paper. We invite the reader to consult Pudlo et al. (2016), Estoup et al. (2018), and Raynal et al. (2018) for more in-depth explanations.

ABC framework

Let $\mathbf{y}$ denote the observed data and $\boldsymbol{\theta}$ a vector of parameters associated to a statistical model whose likelihood is $f(. \mid \boldsymbol{\theta})$. Under the Bayesian parametric paradigm the posterior distribution

$$\pi(\boldsymbol{\theta} \mid \mathbf{y}) \propto f(\mathbf{y} \mid \boldsymbol{\theta})\pi(\boldsymbol{\theta})$$

is of prime interest. It characterizes the distribution of $\boldsymbol{\theta}$ given the observation $\mathbf{y}$ and can be interpreted as an update of the prior distribution $\pi(\boldsymbol{\theta})$ by the likelihood of $\mathbf{y}$. The likelihood is hence pivotal, but unfortunately intractable in the evolutionary scenarios (models) we consider in the present study, as well as in many other evolutionary studies. As a matter of fact, the underlying Kingman's coalescent process (Kingman, 1982) does not allow a close expression for the likelihood because all the possible genealogies and mutational process yielding $\mathbf{y}$ should be considered. To solve this issue, some likelihood-free methods have been developed using the fact that, even though the likelihood is not available, generating artificial (i.e. simulated) data for a given value of $\boldsymbol{\theta}$ is much easier if not feasible (e.g. Beaumont (2010). Approximate Bayesian computation (ABC) is one of them (Beaumont et al., 2002).

In a nutshell, ABC consists in generating parameters $\boldsymbol{\theta}'$ and associated pseudo-data $\mathbf{z}$ from the scenario, and accepting $\boldsymbol{\theta}'$ as a realization from an *approximated* posterior if $\mathbf{z}$ is similar to $\mathbf{y}$. In standard ABC treatments, the notion of similarity is defined through the use of a distance ρ to compare $\eta(\mathbf{z})$ and $\eta(\mathbf{y})$, where $\eta(.)$ is a projection of the data in a lower dimensional space of summary statistics. Only pseudo-data providing distance lower than a threshold ϵ are retained. The choice of ρ, $\eta(.)$ and ϵ is a major issue in ABC (Beaumont, 2010).

ABC-RF is a recently derived ABC approach based on the supervised machine learning tool named Random Forest (Breiman, 2001), which has as major advantage to avoid the three abovementioned difficulties. Initially introduced in Pudlo et al. (2016) for model choice and then extended to parameter inference in Raynal et al. (2018), ABC-RF relies on the use of random forests on a set of simulated pseudo-data according to the generative Bayesian models under consideration. Let consider M Bayesian parametric models. For a given model index $m \in \{1, \ldots, M\}$, a prior probability $\mathbb{P}(\mathcal{M} = m)$ is defined, with $\boldsymbol{\theta}_m$ its associated parameters and $f_m(\mathbf{y} \mid \boldsymbol{\theta}_m)$ its likelihood. The generation process of a reference table made of H elements is described in Algorithm 1.

Algorithm 1: Generation of a reference table with H elements

---
1 for $j \leftarrow 1$ **to** H **do**
2   Generate $m^{(j)}$ from the prior $\mathbb{P}(\mathcal{M} = m)$
3   Generate $\boldsymbol{\theta}_{m^{(j)}}$ from the prior $\pi_{m^{(j)}}(.)$
4   Generate $\mathbf{z}^{(j)}$ from the model $f_{m^{(j)}}(. \mid \boldsymbol{\theta}_{m^{(j)}})$
5   Compute $\eta(\mathbf{z}^{(j)}) = \left(\eta_1(\mathbf{z}^{(j)}), \ldots, \eta_d(\mathbf{z}^{(j)})\right)$
6 end

The output takes the form of a matrix containing simulated model indexes, parameters and summary statistics, as described below

$$
\begin{bmatrix}
m^{(1)} & \theta_{m^{(1)}} & \eta_1(\mathbf{z}^{(1)}) & \eta_2(\mathbf{z}^{(1)}) & \ldots & \eta_d(\mathbf{z}^{(1)}) \\
m^{(2)} & \theta_{m^{(2)}} & \eta_1(\mathbf{z}^{(2)}) & \eta_2(\mathbf{z}^{(2)}) & \ldots & \eta_d(\mathbf{z}^{(2)}) \\
\vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
m^{(H)} & \theta_{m^{(H)}} & \eta_1(\mathbf{z}^{(H)}) & \eta_2(\mathbf{z}^{(H)}) & \ldots & \eta_d(\mathbf{z}^{(H)})
\end{bmatrix}.
$$

ABC-RF for model choice

The ABC-RF strategy for model choice is described in Algorithm 2. The output is the affectation of $\mathbf{y}$ to a model (scenario), this decision being made based on the majority class of the RF tree votes.

Algorithm 2: ABC-RF for model choice

Input : a reference table used as learning set, made of H elements, each one composed of a model index $m^{(H)}$ and d summary statistics. A possibly large collection of summary statistics can be used, including some obtained by machine-learning techniques, but also by scientific theory and knowledge

Learning : construct a classification random forest $\hat{m}(\cdot)$ to infer model indexes

Output : apply the random forest classifier to the observed data $\eta(\mathbf{y})$ to obtain $\hat{m}(\eta(\mathbf{y}))$

The selected scenario is the one with the highest number of votes in his favor. In addition to this majority vote, the posterior probability of the selected scenario can be computed as described in Algorithm 3.

Algorithm 3: ABC-RF computation of the posterior probability of the selected scenario

---

Input : the values of $\mathbb{I}\left\{m^{(h)} \neq \hat{m}(\eta(\mathbf{z}^{(h)}))\right\}$ for the trained random forest and corresponding summary statistics of the reference table, using the out-of-bag classifiers

Learning : construct a regression random forest $\hat{\mathbb{E}}(.)$ to infer $\mathbb{E}\left(\mathbb{I}\left\{m \neq \hat{m}(\eta(\mathbf{y}))\right\} \mid \eta(\mathbf{y})\right)$

Output : an estimate of the posterior probability of the selected model $\hat{m}(\eta(\mathbf{y}))$

$$\hat{\mathbb{P}}\left(m = \hat{m}(\eta(\mathbf{y})) \mid \eta(\mathbf{y})\right) = 1 - \hat{\mathbb{E}}\left(\mathbb{I}\left\{m \neq \hat{m}(\eta(\mathbf{y}))\right\} \mid \eta(\mathbf{y})\right)$$

Such posterior probability provides a confidence measure of the previous prediction at the point of interest $\eta(\mathbf{y})$. It relies on the building of a regression random forest designed to explain the model prediction error. More specifically, and as a first step, posterior probability computation makes use of out-of-bag predictions of the training dataset. Because each tree of the random forest is built on a bootstrap sampling of the H elements of the reference table (i.e. the training dataset), there is about one third of the reference table that remains unused per tree, and this ensemble of left aside datasets corresponds to the "out-of-bag". Thus, for each pseudo-data of the reference

table, one can obtain an out-of-bag prediction by aggregating all the classification trees in which the pseudo-data was out-of-bag. In a second step, the out-of-bag predictions $\hat{m}(\eta(\mathbf{z}^{(h)}))$ are used to compute the indicators $\mathbb{I}\left\{m^{(h)} \neq \hat{m}(\eta(\mathbf{z}^{(h)}))\right\}$. These 0 - 1 values are used as response variables for the regression random forest training, for which the explanatory variables are the summary statistics of the reference table. Predicting the observed data thanks to this forest allows the derivation of the posterior probability of the selected model (Algorithm 3). Note that using the out-of-bag procedure prevents over-fitting issues and is computationally parsimonious as it avoids the generation of a second reference table for the regression random forest training.

Model grouping   A recent useful add-on to ABC-RF has been the model-grouping approach developed in Estoup et al. (2018), where pre-defined groups of scenarios are analysed using Algorithm 2 and 3. The model indexes used in the training reference table are modified in a preliminary step to match the corresponding groups, which are then used during learning phase. When appropriate, unused scenarios are discarded from the reference table. This improvement is particularly useful when a high number of individual scenarios are considered and have been formalized through the absence or presence of some key evolutionary events (e.g. admixture, bottleneck, ...). Such key evolutionary events allow defining and further considering groups of scenarios including or not such events. This grouping approach allows to evaluate the power of ABC-RF to make inferences about evolutionary event(s) of interest over the entire prior space and assess (and quantify) whether or not a particular evolutionary event is of prime importance to explain the observed dataset (see Estoup et al. (2018) for details and illustrations).

ABC-RF for parameter estimation

Once the selected (i.e. best) scenario has been identified, the next step is the estimation of its parameters of interest under this scenario. The ABC-RF parameter estimation strategy is described in Algorithm 4 and takes a similar structure to Algorithm 2. The idea is to use a regression random forest for each dimension of the parameter space (i.e. for each parameter). For a given parameter of interest, the output of the algorithm is a vector of weights $\mathbf{w_y}$ that can be used to compute posterior quantities of interest such as expectation, variance and quantiles. $\mathbf{w_y}$ provides an empirical posterior distribution for $\theta_{m,k}$; see Raynal et al. (2018) for more details.

Algorithm 4: ABC-RF for parameter estimation

Input : a vector of $\theta_{m^{(h)},k}$ values (i.e. the k-th component of $\theta_{m^{(h)}}$) and d summary statistics

Learning : construct a regression random forest to infer parameter values

Output : apply the random forest to the observed data $\eta(\mathbf{y})$, to deduce a vector of weights
$\mathbf{w_y} = \{w_{\mathbf{y}}^{(1)}, \ldots, w_{\mathbf{y}}^{(H)}\}$, which provides an empirical posterior distribution for $\theta_{j,k}$
$\mathbf{w_y}$ is used to compute the estimators of the mean, the variance and the quantiles of the parameter of interest

$$\hat{\mathbb{E}}(\theta_{m,k} \mid \eta(\mathbf{y})), \quad \hat{\mathbb{V}}(\theta_{m,k} \mid \eta(\mathbf{y})), \quad \hat{\mathbb{Q}}_\alpha(\theta_{m,k} \mid \eta(\mathbf{y}))$$

Global prior errors

In both contexts, model choice or parameter estimation, a global quality of the predictor can be computed, which does not take the observed dataset (about which one wants to make inferences) into account. Random forests make it possible the computation of errors on the training reference table, using the out-of-bag predictions previously described in the section "ABC-RF for model choice".

For model choice, this type of error is called the prior error rate, which is the mis-classification error rate computed over the entire multidimensional prior space. It can be computed as

$$\frac{1}{H} \sum_{h=1}^{H} \mathbb{I} \left\{ m^{(h)} \neq \hat{m}(\eta(\mathbf{z}^{(h)})) \right\}.$$

For parameter estimation, the equivalent is the prior mean squared error (MSE) or the normalised mean absolute error (NMAE), the latter being less sensitive to extreme values. These errors are computed as

$$\text{MSE} = \frac{1}{H} \sum_{h=1}^{H} \left(\theta_{m^{(h)},k} - \hat{\theta}_{m^{(h)},k} \right)^2,$$

$$\text{NMAE} = \frac{1}{H} \sum_{h=1}^{H} \left| \frac{\theta_{m^{(h)},k} - \hat{\theta}_{m^{(h)},k}}{\theta_{m^{(h)},k}} \right|.$$

They can be perceived as Monte Carlo approximation of expectations with respect to the prior distribution.

Local posterior errors

In the present paper, we propose some posterior versions of errors, which target the quality of prediction with respect to the posterior distribution. As such errors take the observed dataset $\eta(\mathbf{y})$ into account, we mention them as local posterior errors.

For model choice, the posterior probability provided by Algorithm 3 is a confidence measure of the selected scenario given the observation. Therefore

$$1 - \hat{\mathbb{P}} \left(m = \hat{m}(\eta(\mathbf{y})) \mid \eta(\mathbf{y}) \right)$$

directly yields the posterior error associated to $\eta(\mathbf{y})$: $\hat{\mathbb{P}} \left(m \neq \hat{m}(\eta(\mathbf{y})) \mid \eta(\mathbf{y}) \right)$.

For parameter estimation, when trying to infer on $\theta_{m,k}$, a point-wise analogous measure of a local error can be computed as the posterior expectations

$$\mathbb{E} \left(\left(\theta_{m,k} - \hat{\theta}_{m,k} \right)^2 \mid \eta(\mathbf{y}) \right) \quad \text{and} \quad \mathbb{E} \left(\left| \frac{\theta_{m,k} - \hat{\theta}_{m,k}}{\theta_{m,k}} \right| \mid \eta(\mathbf{y}) \right). \tag{1}$$

We approximate these expectations by

$$\sum_{i=1}^{H} w_{\mathbf{y}}^{(h)} \left(\theta_{m^{(h)},k} - \hat{\theta}_{m^{(h)},k} \right)^2 \quad \text{and} \quad \sum_{i=1}^{H} w_{\mathbf{y}}^{(h)} \left| \frac{\theta_{m^{(h)},k} - \hat{\theta}_{m^{(h)},k}}{\theta_{m^{(h)},k}} \right|.$$

We again uses the out-of-bag information to compute $\hat{\theta}_{m^{(h)},k}$, hence avoiding the (time consuming) production of a second reference table, and assume that the weights $\mathbf{w_y}$ from the regression random forest are good enough to approximate any posterior expectations of functions of $\theta_{m,k}$: $\mathbb{E}(g(\theta_{m,k}) \mid \eta(\mathbf{y}))$.

Another more expensive strategy to evaluate the posterior expectations (1) is to construct new regression random forests using the out-of-bag vector of values

$$\left(\theta_{m^{(h)},k} - \hat{\theta}_{m^{(h)},k}\right)^2 \qquad \text{or} \qquad \left|\frac{\theta_{m^{(h)},k} - \hat{\theta}_{m^{(h)},k}}{\theta_{m^{(h)},k}}\right|,$$

depending on the targeted error. The observation $\eta(\mathbf{y})$ is then given to the forests, targeting the expectations (1).

Note that the values $\hat{\theta}_{m^{(h)},k}$ in the previous formulas can be replaced by either the approximated posterior expectations $\hat{\mathbb{E}}(\theta_{m^{(h)},k} \mid \eta(\mathbf{y}))$ or the posterior medians $\hat{\mathbb{Q}}_{50\%}(\theta_{m^{(h)},k} \mid \eta(\mathbf{y}))$, again using the out-of-bag information, to provide the local posterior errors. We found that both in the present paper (see main text, Materials and Methods section) and for various tests that we carried out on different inferential setups and datasets (results not shown), the posterior median provides a better accuracy of parameter estimation than the posterior expectation (aka posterior mean). This trends also holds for global prior errors that can be computed using either the mean or the median as point estimates.

As final comment, it is worth noting that so far a common practice consisted in evaluating the quality of prediction (for model choice or parameter estimation) in the neighborhood of the observed dataset, that is around $\eta(\mathbf{y})$ and not exactly for $\eta(\mathbf{y})$. For model choice, Estoup et al. (2018) use the so called posterior predictive error rate which is an error of this type. In this case, some simulated datasets of the reference table close to the observation are selected thanks to an Euclidean distance, then new pseudo-observed datasets are simulated using similar parameters, on which is computed the error (see also Lippens et al., 2017, for a similar approach in a standard ABC framework). However, the main problem of processing this way is the difficulty to specify the size of the area around the observation, especially when the number of summary statistics is large. We therefore do not recommend the use of such a "neighborhood" error anymore, but rather to compute the local posterior errors detailed above as the latter measured prediction quality exactly at the position of interest $\eta(\mathbf{y})$.

References

Beaumont, M. A. (2010). Approximate Bayesian Computation in Evolution and Ecology. *Annual Review of Ecology, Evolution, and Systematics*, 41:379–406.

Beaumont, M. A., Zhang, W., and Balding, D. (2002). Approximate Bayesian Computation in Population Genetics. *Genetics*, 162(4):2025–2035.

Estoup, A., Raynal, L., Verdu, P., and Marin, J.-M. (2018). Model choice using Approximate Bayesian Computation and Random Forests: analyses based on model grouping to make inferences about the genetic history of Pygmy human populations. *Journal de la Société Française de Statistique*, 159(3).

Kingman, J. F. C. (1982). On the Genealogy of Large Populations. *Journal of Applied Probability*, 19(A):27–43.

Lippens, C., Estoup, A., Hima, M. K., Loiseau, A., Tatard, C., Dalecky, A., Bâ, K., Kane, M., Diallo, M., Sow, A., Niang, Y., Piry, S., Berthier, K., Leblois, R., Duplantier, J. M., and Brouat, C. (2017). Genetic structure and invasion history of the house mouse (Mus musculus domesticus) in Senegal, West Africa: a legacy of colonial and contemporary times. *Heredity*, 119(2):64–75.

Pudlo, P., Marin, J.-M., Estoup, A., Cornuet, J.-M., Gautier, M., and Robert, C. P. (2016). Reliable ABC model choice via random forests. *Bioinformatics*, 32(6):859–866.

Raynal, L., Marin, J.-M., Pudlo, P., Ribatet, M., Robert, C. P., and Estoup, A. (2018). ABC random forests for Bayesian parameter inference. *Bioinformatics*. to appear.

Supplementary material S4. Details on results from ABC-RF treatments when assuming uniform priors for the three time period parameters of the studied scenarios.

Table S4.1. Scenario choice for each of the ten replicate analyses using uniform priors for the three time period parameters of the studied scenarios.

Reference table	Best scenario	Votes S	Votes SC	Votes SB	Votes SCB	Votes SA_U	Votes SCA_U	Votes SBA_U	Votes $SCBA_U$	Votes SA_B	Votes SCA_B	Votes SBA_B	Votes $SCBA_B$	Prior error rate	Posterior probability (best scenario)
1	scenario SCB	0.013	0.045	0.147	0.462	0.007	0.022	0.020	0.040	0.041	0.099	0.034	0.069	0.778	0.596
2	scenario SCB	0.018	0.027	0.140	0.365	0.012	0.026	0.033	0.096	0.044	0.075	0.072	0.092	0.776	0.398
3	scenario SCB	0.012	0.077	0.182	0.326	0.006	0.030	0.019	0.069	0.038	0.102	0.064	0.074	0.775	0.426
4	scenario SCB	0.014	0.035	0.175	0.405	0.007	0.022	0.022	0.061	0.023	0.091	0.040	0.104	0.777	0.563
5	scenario SCB	0.019	0.048	0.101	0.416	0.010	0.024	0.026	0.063	0.039	0.092	0.055	0.106	0.778	0.547
6	scenario SCB	0.018	0.030	0.130	0.295	0.011	0.030	0.013	0.124	0.075	0.093	0.098	0.082	0.778	0.398
7	scenario SCB	0.016	0.057	0.094	0.308	0.011	0.018	0.015	0.050	0.028	0.113	0.058	0.231	0.775	0.511
8	scenario SCB	0.019	0.037	0.109	0.359	0.007	0.018	0.015	0.049	0.057	0.080	0.105	0.145	0.777	0.474
9	scenario SCB	0.010	0.052	0.114	0.286	0.017	0.023	0.014	0.070	0.069	0.136	0.098	0.113	0.777	0.460
10	scenario SCB	0.013	0.057	0.124	0.249	0.009	0.016	0.012	0.067	0.061	0.133	0.091	0.168	0.778	0.442
Mean	**scenario SCB**	**0.015**	**0.047**	**0.132**	**0.347**	**0.010**	**0.023**	**0.019**	**0.069**	**0.048**	**0.101**	**0.072**	**0.118**	**0.777**	**0.481**

We empirically evaluated the influence of shape of prior distributions for the time periods on our inferences by conducting all ABC-RF analyses assuming a set of uniform priors bounded between 100 and 500,000 generations. In the main document, prior values for time periods were drawn from log-uniform distributions bounded between 100 and 500,000 generations. We report values for the proportion of votes, prior error rates and posterior probabilities of the best scenario on ten replicate analyses based on ten different reference tables. Scenarios are depicted in Figure 2. For each reference table, the number of datasets simulated using DIYABC was set to 150,000 and the number of RF-trees was 3,000. The scenario SCB was the best supported for all replicate analyses: it involves a bottleneck event in *S. g. flaviventris* right after divergence, a population size contraction in the ancestral population and not any secondary contact with asymmetrical genetic admixture from *S. g. gregaria* into *S. g. flaviventris*.

Table S4.2. Estimation of the divergence time between *S. g. gregaria* **and** *S. g. flaviventris* **for ten replicate analyses using uniform prior distributions for the three time period parameters under the best supported scenario (scenario SCB).**

t_{div} (G)	Median	q5%	q95%
reference table 1	10236.6	3678.5	25457.0
reference table 2	7603.0	2311.2	23020.3
reference table 3	8500.0	2788.8	19401.0
reference table 4	10598.5	3564.4	24444.5
reference table 5	9226.0	3147.9	21286.6
reference table 6	9665.0	3381.5	24445.2
reference table 7	8675.5	2572.3	26375.8
reference table 8	10281.6	2919.8	27349.0
reference table 9	8845.0	3271.9	24094.7
reference table 10	7909.7	2040.1	21333.4
Mean	**9154.1**	**2967.6**	**23720.8**
SD	**1027.5**	**541.6**	**2476.4**

We empirically evaluated the influence of shape of prior distributions for the time periods on our inferences by conducting all ABC-RF analyses assuming a set of uniform priors bounded between 100 and 500,000 generations. Median value and 90% CI for priors are 146,936 and 13,195 – 498,867, respectively. Replicate analyses have been processed on different reference tables. For each reference table, the number of datasets simulated using DIYABC was set to 100,000 and the number of RF- trees was 2,000. Divergence times are given in number of generations. SD stands for standard deviations computed from the ten values of median, 5% quantile and 95% quantile estimated from the ten replicate analyses.

Figure S4.1. Estimation of the time since divergence between the two desert locust subspecies as a function of time scales using uniform prior distributions for the three time period parameters under the best supported scenario (scenario SCB).
Simulated datasets (5,000 par divergence time) were generated for fixed divergence time values of 100 ; 250 ; 500; 1,000 ; 2,500; 5,000 ; 10,000 ; 25,000 ; 50,000; 100,000; and 250,000 generations. The median (plain lines) and 90% credibility interval (90% CI; dashed lines), averaged over the 5,000 datasets, are represented. Divergence time values are in number of generations.