Title

Accurate, ultra-low coverage genome reconstruction and association studies in Hybrid Swarm mapping populations

Cory A. Weller[1,2], Alan O. Bergland[1]

[1] Department of Biology, University of Virginia, Charlottesville, Virginia 22904

[2] E-mail: caw5cv@virginia.edu

Abstract

Genetic association mapping studies seek to uncover the link between genotype and phenotype, and often utilize inbred reference panels as a replicable source of genetic variation. However, inbred reference panels can differ substantially from wild populations in their genotypic distribution, and patterns of linkage-disequilibrium and nucleotide diversity. As a result, associations discovered using inbred reference panels may not reflect the genetic basis of phenotypic variation in natural populations. To address this problem, we evaluated a mapping population design where dozens to hundreds of inbred lines are outbred for few (e.g. five) generations, which we call the Hybrid Swarm. The Hybrid Swarm approach has likely remained underutilized relative to pre-sequenced inbred lines due to the costs of genome-wide genotyping. To reduce sequencing costs and make the Hybrid Swarm approach feasible, we developed a computational pipeline that reconstructs accurate whole genomes from ultra-low-coverage (0.05X) sequence data in Hybrid Swarm populations derived from ancestors with phased haplotypes. We compared the power and precision of GWAS using the Hybrid Swarm, inbred lines, recombinant inbred lines, and highly outbred populations across a range of allele frequencies and effect sizes, modeling genetic variation from the Drosophila Genetic Reference Panel as well as variation from neutral simulations. While inbred populations tended to perform best due to the intrinsic power benefits conferred by the lack of heterozygotes, association mapping with the Hybrid Swarm performed comparably to highly outbred (F_{50}) populations and has higher precision than mapping with inbred lines. Taken together, our results demonstrate the feasibility of the Hybrid Swarm as a cost-effective method of fine-scale genetic mapping.

Introduction

Genetic mapping studies seek to describe the link between genotype and phenotype.

For experimental crosses, mapping was traditionally conducted by scoring the

phenotypes of recombinant offspring descended from a limited number of parental lines.

While such QTL mapping studies can have high power to detect associations, they offer

minimal mapping resolution (Cheng *et al.* 2010), often detecting broad regions of

phenotypic association (Bergland *et al.* 2012). If linkage disequilibrium is lowered,

spurious associations become rarer (Li *et al.* 2005) and associations can be resolved at

the gene or nucleotide level, as in GWAS of large outbred populations (Nikpay *et al.*

2015; Wu *et al.* 2017; Monir and Zhu 2017). However, GWAS suffer from reduced

power to detect associations, necessitating a large sample size relative to QTL mapping

(Spencer *et al.* 2009).

To generate higher resolution mapping populations than the traditional biparental F2

design, Multiparent Populations (MPPs) are commonly used. By crossing together

multiple inbred lines, researchers can produce genetically diverse mapping populations

without sampling wild individuals. MPPs are commonly used for the dissection of

complex traits in model organisms (Chesler *et al.* 2008; Kover *et al.* 2009; King *et al.*

2012b) and agriculturally important crops (Huang *et al.* 2012; Singh *et al.* 2013; Krämer

et al. 2014). The mapping resolution of MPPs depends on the extent of linkage

disequilibrium, and resolution is improved by allowing for more recombination between

haplotypes, or by incorporating a greater number of genetically diverse haplotypes (Mott

et al. 2000; Chia *et al.* 2005).

One alternative approach for generating a high-resolution mapping population is to

substitute extensive recombination for increased haplotype diversity. By crossing

dozens to hundreds of inbred lines for a limited number of generations, heterozygous

mapping populations can be generated quickly with sufficiently reduced LD to detect

associations with high resolution. Unfortunately, the many-haplotypes few-generations

method is not without its drawbacks. First, including many haplotypes decreases the

frequency of the rarest alleles, reducing power to detect associations. Second, such an

outbred population would require recurring genotyping efforts (Yang *et al.* 2018) unlike

pre-sequenced homozygous lines. The net requirement of genotyping a large sample

size may explain the widespread use of pre-genotyped inbred reference panels for

genetic association experiments in model systems (Huang *et al.* 2011; King *et al.*

2012b; MacKay *et al.* 2012; Srivastava *et al.* 2017).

Here, we describe computational methods that allow for cost-effective association

mapping with a large outbred population. The Hybrid Swarm is founded by dozens to

hundreds of inbred lines, crossed for a limited number of generations. To reduce

genotyping costs of the Hybrid Swarm, we developed and evaluated a pipeline to

reconstruct whole genomes using ultra-low coverage sequencing data. We developed

and tested our pipeline by reconstructing whole genomes for thousands of simulated

Hybrid Swarm individuals. Our simulated genomes draw from natural variation in the

Drosophila melanogaster Genetic Reference Panel (DGRP), as well as from variation

generated from coalescent models representing a broad range of genetic diversity

parameters for common model systems. We show that the Hybrid Swarm approach

allows for highly accurate genotyping (average 99.9% genotypic accuracy) from ultra-

low-coverage (0.005-0.05X) whole-genome individual-based sequencing. We then

perform simulated GWAS to describe the power and precision of association mapping in

the Hybrid Swarm compared to inbred lines, recombinant inbred lines, and a highly

outbred (F_{50}) population. Our computational tools are capable of efficiently simulating

low-coverage reconstruction and GWAS power analysis of any model system. Together,

our results the feasibility of cost-effective high-resolution association mapping in a large

outbred population.

91   **Methods**

Generating and preparing simulated reference panels.

In order to evaluate low-coverage reconstruction for various degrees of genetic

diversity, we generated reference panels using haplotypes produced by coalescent

models across a range of genetic diversity levels. Haplotypes were generated using the

R (R Core Team 2016) package *scrm* (Paul R. Staab et al. 2015) and subsequently

restructured into VCF file format (Danecek et al. 2011). We generated ten independent

panels for each of all 18 combinations of population size ($N_e = 10^4, 10^5, 10^6$), mutation

rate ($\mu = 10^{-9}, 5 \times 10^{-9}, 10^{-8}$), and number of haplotypes (32, 128). The value for θ for

each simulation was defined as $4N_e\mu$. We simulated a chromosome-length locus of 25

Mb with a recombination rate of 1.5 cM/Mb. SNP positions output by *scrm* (a decimal

within the range of 0 to 1) were converted to base pair positions by multiplying the

decimal by chromosome length (25×10^6 base pairs for our simulations) and rounding

down to the nearest integer. Any sites with more than two alleles were converted to a

biallelic site by discarding tertiary or quaternary alleles. Genotype values were re-coded

as polarized signed integers: +1 for reference and -1 for alternate alleles. For every

position, reference and alternate alleles were defined by randomly selecting one of the

twelve non-repeating pairs of nucleotides. Reference genome FASTA files were

created with a custom python script that generated a 25 million length string of

nucleotide characters with weighted probability to achieve 45% GC-content, followed by

replacing variable positions with their respective reference alleles.

Preparing DGRP haplotype data

As a case study of low-coverage genome reconstruction in a model system, we

incorporated wild fruit fly genetic diversity from the Drosophila Genetic Reference Panel

(MacKay *et al.* 2012) DGRP freeze 2 as available from the *Drosophila* Genome Nexus

(Lack *et al.* 2015). To minimize missing data, we included the 129 lines (out of 205)

which exhibited aligned whole genome FASTA files with less than 50% of nucleotides

indicated by the ambiguity character N. We excluded insertions, deletions, fixed sites,

and sites with more than two alleles. Any heterozygous genotype calls were masked as

missing data. Diploid genotypes were re-coded as a single signed integer value, with +1

for homozygous reference, -1 for homozygous alternate, and 0 for missing data. This

resulted in a polarized VCF file containing only biallelic SNPs and only homozygous (or

missing) genotype calls.

Simulating Mapping Populations

To generate simulated populations, we developed a forward-simulator in R that stores

ancestral haplotype block maps instead of genotypes. Our analyses necessitated a

method of storing genotype information for thousands of individuals across thousands of

simulations. To do so, we leveraged information redundancy that exists between related

individuals in recombinant populations, generating haplotype block files. We achieved

between three and four orders of magnitude of compression relative to a VCF file. For

example, for a population containing 5000 diploid genotypes at nearly four million sites,

a compressed VCF file is approximately 6.5 GB, compared to approximately 3.5 MB for

a haplotype block file. This reduced file size is what allowed us to generate and store

28,000 total independent GWAS simulations (500 each for 56 parameter combinations).

When haplotype block ancestry is known and recorded, as is possible with simulations,

genotypes must only be recorded once (for the ancestral founders). Recombinant

individual genotypes can then be reconstituted by extracting ancestral genotypes from

ancestor and base pair position indices.

We simulated Hybrid Swarms through random mating over five non-overlapping

generations at a population size of 10,000. Simulations proceeded in the following

manner: first, a subset of either 32 or 128 founders was selected. Then, of that founder

subset, 10,000 individuals were sampled with replacement. All possible founders were

chosen with equal probability and assigned male or female sex with a 1:1 ratio, where

sex was determined by the presence of a designated sex chromosome. Sexual

reproduction was simulated by random sampling of recombinant gametes from male-

female pairs. Once 10,000 recombinant progeny were generated, the parental

generation was discarded. Reproduction continued until the F_5 population was

achieved. Recombination frequency was modeled as a Poisson process with an

expected value $\lambda = \Sigma(Morgans)$ per chromosome. For simulations of Drosophila

populations based on DGRP chromosomes, recombination occurred only in females,

with recombination frequency and position based on values from Comeron et al (2012).

For populations founded by simulated haplotypes, recombination occurred in both

sexes, with recombination occurring uniformly across each chromosome (Supplemental

Figure S1).

Simulating and Mapping Sequencing Data

We used *wgsim* (Li 2011) to generate simulated reads. To achieve a desired level of

sequencing coverage $C = 0.05$ or 0.005, we generated $N = (C \times S)/(2 \times L)$ reads per

chromosome, with read length $L = 100$ bp and chromosome length S bp. We specified a

base error rate of 0.001 and an indel fraction of 0. Remaining *wgsim* parameters were

left as default.

We assembled paired end reads using *PEAR* (Zhang *et al.* 2014) and separately

aligned the assembled and unassembled groups to a reference genome with *bwa*

0.7.14 using the BWA-MEM algorithm (Li 2013). Reads from DGRP-derived populations

were mapped to the *D. melanogaster* reference genome v5.39, and reads from

coalescent-derived populations were mapped to their respective simulated reference

genomes. After converting mapped reads to compressed BAM format with *samtools*

1.3.1 (Li *et al.* 2009), we removed PCR duplicates with *Picard tools* 2.0.1 (Broad

Institute 2015a).

Most Likely Ancestors Selection

To make chromosome reconstructions in the hybrid swarm computationally tractable

(Figure 1), we developed a method of accurately selecting a subset of most likely

ancestors for any single chromosome. We then used that ancestor subset to reconstruct

haplotype blocks using the *RABBIT* package (Zheng *et al.* 2015) in Mathematica.

RABBIT operates as a Hidden Markov Model (HMM) using the Viterbi algorithm to

return the most likely series of parental combinations (hidden states) across the

genome (SNP positions) given the observations (sequenced alleles). For every position

in the genome, the Viterbi algorithm evaluates relative likelihoods of transitioning to any

possible hidden state. Because the hidden states in our case are ancestor

combinations, there will be $(N^2 + N)/2$ combinations of N haplotypes to evaluate at

every site. This number of evaluations is tractable at smaller values of N but grows at a

quadratic rate. For example, increasing the number of founding haplotypes from 8 to

128 is a 16-fold increase in haplotypes, but it would incur orders of magnitude increases

in computational effort (Figure 1). Thus, in order to make reconstructions in *RABBIT*

computationally tractable for hybrid swarm individuals, it is necessary to identify a

subset of founders that accurately includes the true ancestors contributing to any given

chromosome.

We used the software package *HARP* (Kessner *et al.* 2013) to rank the population

founding lines based on likelihood of being a true ancestor of a chromosome to be

reconstructed. *HARP* was originally developed to estimate haplotype frequencies from

pooled sequence data, and we co-opted it to assess relative likelihood that any founder

contributed to a genomic window. We ran *HARP* with non-overlapping 100 kb windows

with a minimum frequency cutoff 0.0001, producing output which can be visualized as a

heat map of ancestor likelihood across the chromosome. A custom *R* script analyzed

this *HARP* output and ranked all possible founders in terms of likelihood of contribution

for a given chromosome. Briefly, a chromosome-wide significance threshold was

calculated, e.g. the 95% or 99% quantile of all likelihoods across all founders and all

chromosome windows. Then, every potential ancestor for each 100 kb window was

classified as falling above or below this threshold. Founding lines were then ranked in

descending order of the number of windows passing the threshold. We examined two

measures of effectiveness for this method across a range of quantile threshold values

(90%, 95%, 99%, and 99.9%) when selecting up to a maximum number of most likely

ancestral haplotypes. The first measure is the number of true ancestral founders

excluded; the second measure is the fraction of the chromosome derived from

ancestors missing from the selected subset.

Chromosome Reconstruction with RABBIT

We used the MAGIC reconstruct method of the Mathematica package RABBIT (Zheng

et al. 2015) to perform chromosome reconstructions, which has been shown to be

accurate for genotype estimation at sequencing coverage at 0.05X for a variety of

multiparent populations (Zheng *et al.* 2018). RABBIT requires three inputs: observed

genotypes in the individual being reconstructed; map distance (in cM units) of the same

loci; and genotypes for the potential ancestors at those same loci. For DGRP-derived

simulated populations, we specified map distance based on values reported by

Comeron et al. (2012) by performing linear interpolation of cumulative map units (cM) as

a function of base pair position. For populations derived from simulated haplotyopes, we

used a linear function of 37.5 cM over each 25 Mb chromosome. To specify genotype

information, we first counted reference and alternate reads using the Genome Analysis

Toolkit *ASEReadCounter* (Broad Institute 2015b). Because it is not possible to make

confident homozygote genotype calls from low coverage sequencing data where most

sites are observed only once and or twice, we only included diploid genotype

observations for sites where both reference and alternate alleles were observed. As

RABBIT allows for an ambiguous allele character, for all sites where only reference or

alternate reads were observed (but not both), we included one ambiguous allele.

To minimize memory and runtime requirements, we included at most 5,000 SNPs per

chromosome, selected for maximum ancestor-discerning information content. If an

observed (sequenced) allele is common, it will only slightly narrow down the possibility

of ancestors. If a sequenced allele is rare—at the most extreme, unique to one

individual—it provides greater information from which founder that site is derived. Thus,

we designate information-rich sites as those where the frequency of the sequenced

allele is the lowest with respect to the pool of most likely ancestors. In order to sample

sites with high information content spread throughout the chromosome, we used an

iterative approach. First, we included all heterozygous sites (i.e. where reference and

alternate alleles are both observed). Then, 10% of all SNPs were randomly sampled,

and we retained up to the top 0.2% most informative sites, repeating the random

sampling and retention until we designated 5,000 SNPs.

We ran RABBIT independently for each chromosome using the Viterbi decoding

function under the joint model, with all other RABBIT parameters left at default. RABBIT

output was converted to a phased chromosome haplotype map, which we then used to

extract and concatenate genotype information from a VCF file containing founder

genotypes. To calculate genotype reconstruction accuracy, we first imported true

(simulated) and estimated (reconstructed) genotypes using a custom *R* script. We

measured the fraction of all remaining sites where the estimated diploid genotype is

identical to the originally simulated diploid genotype, excluding fixed sites with respect

to the founding haplotypes, and excluding any sites with missing genotype information.

Because male individuals do not possess two copies of the sex chromosome, we only

evaluated accuracy for autosomes.

To measure accuracy of estimated frequency of recombination events, true and

estimated recombination counts were first summed over both copies of each

chromosome in a simulated individual. This removed the possibility of introducing error

by comparing the wrong copies of chromosomes. Only detectable recombination events

were considered, i.e. those that did not occur between homologous haplotypes. We

then used the *epi.ccc* function of the *R* package *epiR* (Stevenson 2018) to calculate

Lin's concordance correlation coefficient (ρ) between the true and estimated

recombination counts.

Modeling Computational Complexity of Chromosome Reconstruction

To estimate the rate at which computational requirements grows with data input, we

performed chromosome reconstructions with varying numbers of potential founders and

markers (SNPs). This allows us to extrapolate the runtime and memory for performing

the most resource intensive chromosome reconstructions (i.e. those with > 40 founding

lines). To generate runtime and memory usage data, we performed 900 reconstructions

using varying sizes of RABBIT input for a single example individual 2L chromosome.

Reconstructions included the four true ancestors of the simulated individual, plus 0 to 32

additional haplotypes (for a total of between 4 and 36 founders, in steps of 4) and a

random selection of marker SNPs (between 500 and 5000 in steps of 500). Ten

replicates, each with a unique random set of SNPs, was conducted for each

combination of N founding lines and S SNPs using a single core on the University of

Virginia computing cluster, with total runtime and peak memory usage as reported from

the SLURM workload manager (CPUtime and MaxRSS, respectively). We then

modeled the mean runtime and memory usage (averaged across 10 replicates per

parameter combination) as a function of number of founding lines and number of SNPs

fed into RABBIT. For runtime, simulations involving 8 or fewer founding lines were

omitted from the regression model because they ran too quickly to resolve non-zero

runtime. Memory was modeled as $Memory(GB) = 7.367 \times 10^{-9} \times SN^4 + 0.0316$, while

runtime was modeled as $Runtime(Minutes) = [1.189 \times 10^{-3} \times N^2 + 1.038 \times 10^{-6} \times$

$SN^2 + 2.649 \times 10^{-4} \times S]^2$.

Simulated GWAS

We performed GWAS on mapping populations produced by random sampling and

permutation of the previously-described forward-simulated populations. Although the

forward simulator we developed is efficient, it would not have been computationally

feasible to simulate 500 fully independent mapping populations (per parameter

combination) in a reasonable amount of time. Instead, we generated ten independent

forward-simulated populations, and for each of those, generated fifty randomly

permuted subsets (Figure 2). For a single simulated mapping population, we began by

sampling (with replacement) a random subset of 5,000 individuals, out of 10,000 total

individuals generated by forward-simulation. Then, we performed a permutation of

haplotype ancestry with a new, randomly-ordered (equally sized) subset of founders.

The permutation of ancestry was one-to-one, e.g. all haplotype blocks that were

previously derived from founder X would be translated to founder Y, and blocks

previously derived from Y would in turn be mapped to founder Z.

In addition to Hybrid Swarm populations, which we ran through the simulated

sequencing and mapping pipeline, we generated four additional types of mapping

populations for comparing GWAS performance: Highly outbred (F_{50}) populations, similar

to sampling wild individuals; Inbred Lines (ILs), similar to mapping with the DGRP); and

Recombinant Inbred Lines (RILs), similar to mapping with the DSPR.

The F_{50} populations were generated in same manner as the Hybrid Swarm, except for

fifty non-overlapping generations of recombination instead of five generations. The ten

resulting forward-simulated populations were resampled and permuted as we did with

the Hybrid Swarms.

We simulated ten initial sets of 800 RILs using the same forward-simulator as previously

described, each initialized with a random subset of eight DGRP haplotypes. Populations

randomly recombined at a population size of 10,000 for fifty non-overlapping

generations, after which 800 random male-female pairs of individuals were isogenzied

through 25 generations of full-sibling mating. This scenario roughly corresponds to the

Drosophila Synthetic Population Resource (King *et al.* 2012a). For computational

simplicity, after the 25 generations of isogenization we removed any remaining residual

heterozygosity by forcing the identity of a second chromosome copy to be identical to

the first copy. We then sampled 5,000 draws (with replacement) of the 800 RILs

followed by ancestry permutation as described above.

To simulate GWAS on Inbred Lines, no forward-simulation was necessary. For a single

simulated population, we first randomly selected 128 DGRP lines with high coverage

and low levels of heterozygosity as the set of founders. Then, those 128 lines were

randomly sampled with replacement 5,000 times. As with hybrid swarm and RILs, for

any parameter combination we generated a total of 500 mapping populations.

Phenotypes were modeled as probabilistic assignment to a case or control group

dependent on allele dosage at a purely additive single SNP. We designated a causal

locus as a random autosomal biallelic SNP segregating within 0.5% of a desired minor

allele frequency (50%, 25%, and 12.5%). We modeled SNPs at 5% and 10% percent

variation explained (PVE), where reference allele homozygotes were assigned to the

case group with probability $50\% - PVE/2$, and alternate allele homozygotes were

assigned to the case group with probability $50\% + PVE/2$. Heterozygotes were equally

likely to be assigned case and control.

To perform many replicates of GWAS for many parameter combinations, we performed

a simple χ^2 test of independence for reference and alternate allele counts between case

and control groups. To do so most efficiently, we developed a method of aggregating

allele counts that uses a haplotype map table in conjunction with a single table of

founder genotypes (Figure 2). Briefly, haplotype table breakpoints across all individuals

were sorted in ascending order. When iterating through ascending unique start and stop

positions, between any pair of breakpoints, all SNPs will be comprised of the same

number of each founding haplotype. Haplotype IDs could then be counted and sorted in

the same column position order as the table containing polarized allele status (-1 for

alternate, +1 for reference). Multiplying the genotype table by the haplotype count vector

results in final allele counts, polarized negative for alternate alleles and positive for

reference alleles. For inbred mapping populations, we corrected for non-independent

allele draws by dividing the χ^2 value by two.

To describe the accuracy of simulated GWAS, we measured the likelihood of including

a locus that is near the causal site when considering a set of the top N most significant

SNPs. Here, 'near' is defined as either exact-SNP resolution, or within 1, 10, or 100 Kb.

In the case of 1 kb precision, we first consider the set of SNPs +/- 1 kb from the most

significant locus (greatest chi-square statistic). Then, we consider the second set of

SNPs as those within +/- 1 kb of the most significant locus outside of the window

already accounted for. This selection of significant clusters was repeated iteratively for

the top 25 regions, for window sizes of 0 (exact SNP resolution), 1 kb, 10 kb, and 100

kb.

We calculated genomic inflation factor (GIF, λ_{1000}) as the value of $\chi^2_{observed}/\chi^2_{expected}$

with two degrees of freedom. Because GIF increases with sample size, we performed a

correction to report the level of GIF expected with a sample size of 1,000 case and

1,000 control individuals (Freedman *et al.* 2004).

Assessing counts of variable sites at appreciable frequency in the DGRP

There is a reduction in power to detect associations with alleles segregating at low

minor allele frequencies. When a population is founded by N lines, any SNP will be

segregating at a relative frequency of at least 1/N, given that the SNP is not fixed within

the population and haplotypes are equally represented. We counted the number of sites

on a given chromosome arm segregating above a minor allele frequency threshold of

MAF=(0.05, 0.125, and 0.25) for random draws (without replacement) when sampling

N=(2, 4, 8, 16, 32, 64, 128) haplotypes of the 129 included DGRP lines We performed

this sampling 20 times for each chromosome arm.

Data Availability Statement

The code used to generate, process, and plot our data is available on GitHub:

https://github.com/cory-weller/low-coverage-genome-reconstruction

384   **Results**

Computational Complexity of Chromosome Reconstruction

To determine reasonable limits for numbers of SNPs and haplotypes used for

chromosome reconstruction with RABBIT, we modeled peak memory usage and

runtime across a range of input sizes. Peak memory grew linearly with number of SNPs

used, and at a greater-than-linear rate with haplotypes (Figure 1A, $Memory =$

$7.367 \times 10^{-9} \times SN^4 + 0.0316), \ F = 3.534 \times 10^6, \ df = 1 \ \& \ 88, \ R^2 = 1$). The runtime of

RABBIT increased at a greater-than-linear rate for both number of SNPs and number of

haplotypes, though the *N* parameter dominates (Figure 1B, $Runtime = [1.189 \times 10^{-3} \times$

$N^2 + 1.038 \times 10^{-6} \times SN^2 + 2.649 \times 10^{-4} \times S], \ F = 4.316 \times 10^4, \ df = 3 \ \& \ 67, \ R^2 =$

0.9995). These models allowed us to estimate resource requirements at greater

numbers of haplotypes (Figure 1, C & D) which would be unfeasible to measure

empirically.

Most-Likely-Ancestor Selection

To reduce computational requirements of haplotype reconstructions with RABBIT, we

developed and evaluated an algorithm for selecting a minimum representative set of

Most-Likely-Ancestors (MLAs) for chromosome reconstruction. We found a HARP

threshold of 0.99 (see methods) discerned a minimal subset of founding lines that

tended to include a given chromosome's true ancestors (Figure 3). At this threshold,

outcomes became asymptotic at the computationally tractable cap of 16 founding lines

(Figure 1). Thus, we performed chromosome reconstruction using up to 16 most-likely-

ancestors as inferred with a HARP threshold of 0.99.

In all cases, decreasing the HARP threshold from 0.95 to 0.90 further reduced

chromosome representation while increasing the number of extraneous founding lines

selected for reconstruction. While a higher HARP threshold of 0.999 yielded the

smallest and most computationally tractable set sizes of MLAs ($\overline{N}$=2.4-3.5), the strict

threshold excluded true ancestors, resulting in a set that is least representative of

chromosomes to be reconstructed. For 128-founder populations, a threshold of 0.999

failed to identify founders constituting an average of 3.33% and 13.9% of chromosomes

for DGRP- and Coalescent-founded populations, respectively. In 32-founder

populations, the 0.999 threshold missed founders representing an average of 15.5%

and 29.7% of chromosomes from DGRP- and Coalescent-founded populations,

respectively.

Populations simulated with genetic variation derived from coalescent models described

above included the parameters $N_e = 10^6$ and $\mu = 5 \times 10^{-9}$. The effectiveness of most-

likely-ancestor selection for populations modeled across extended values of N_e and μ is

shown in supplemental Figures S2 and S3, respectively. Similarly, the number of most-

likely-ancestors chosen for reconstruction in RABBIT are shown in Figures S5 and S6.

Selected MLA set size is described in Figure S5 for 32-founder populations and Figure

S6 for 128-founder populations. Ancestor selection effectiveness for DGRP-derived

populations at two levels of sequencing coverage (0.005X and 0.05X) is shown in

supplemental Figure S4, and the corresponding number of most-likely-ancestors

chosen for reconstruction are shown in supplemental Figure S7.

Reconstruction Accuracy

Chromosome reconstruction of simulated F_5 Hybrid Swarm genomes at 0.05X

sequencing coverage yielded highly accurate genotype estimates (Figure 4). The

median percent of sites with correctly estimated genotypes was greater than 99.9%

whether the population was founded by 32 or 128 founding lines for either DGRP or

coalescent ($N_e = 10^6$ and $\mu = 5 \times 10^{-9}$) haplotypes. We additionally report

reconstruction accuracy in coalescent-derived populations across a range of N_e and μ

values in supplemental Figure S8.

For simulations founded by DGRP lines, 80.5% of reconstructed chromosomes from 32-

founder populations exhibited > 99.9% accuracy, with the remaining 19.5% of

reconstructions contributing to a long tail with a minimum of 84.37%. Increasing the

number of founding lines to 128 resulted in genotype accuracy above 99% for all cases

(minimum: 99.4%), with 83% of reconstructed chromosomes achieving greater than

99.9% accuracy.

Although median accuracy for coalescent-derived populations was equivalent to that of

DGRP-derived populations (99.9%), coalescent-derived populations with 32 founders

exhibited a greater number of low-accuracy reconstructions. While 82.5% of simulations

with 32 coalescent haplotypes were at least 99% accurate, the remaining 17.5% of

reconstructions contributing to a long tail with a minimum accuracy of 59.7%. Increasing

the number of founding lines to 128 resulted in 96.3% of simulations being greater than

99% accurate, with a minimum accuracy of 89.6%.

The number of recombination events estimated from chromosome reconstruction was

most accurate for populations founded by 128 lines (Table 1). Reconstructions of

DGRP- and Coalescent-derived chromosomes yielded recombination count estimates

that were 98.6% and 95.6% concordant with their respective true recombination counts

(Lin's concordance correlation coefficient, ρ). When populations were founded by 32

lines, recombination count estimates were more inaccurate, with DGRP- and

Coalescent-derived reconstructions achieving 50.2% and 75.9% concordance with their

respective true recombination counts. For 32-founder populations, DGRP-derived

reconstructions tended to slightly overestimate recombination counts, while the same

counts were underestimated for coalescent-derived populations.

Simulations that inferred an unlikely high number of recombination events tended to

exhibit reduced accuracy (Figure 4). All DGRP-derived simulated individuals (of 1600

total) exhibited ≤ 8 recombination events, and all but three coalescent-derived simulated

individuals (7197 of 7200 total) exhibited ≤ 9 recombination events. Accordingly, we

considered any reconstructions to be 'hyper-recombinant estimates' if the inferred

recombination count is greater than 8 for DGRP-derived populations or greater than 9

for coalescent-derived populations.

At 0.05X sequencing coverage, hyper-recombinant estimates did not occur for 128-

founder populations, and only rarely resulted from 32-founder populations. Within

DGRP-derived 32-founder populations, reconstructions with hyper-recombinant

estimates were below the sixth percentile of genotype accuracy (N=6/400 simulations,

genotype accuracy range=92.8%-98.8%). For coalescent-derived 32-founder

populations, reconstructions estimated as hyper-recombinant fell in the bottom 9% of

genotype accuracy (N=3/400 simulations, genotype accuracy range = 92.1%-95.6%).

Although hyper-recombinant estimates always fell in the bottom 10% of accuracy, the

least accurate reconstructions were not hyper-recombinant. For coalescent-derived 32-

founder populations, 4.25% (17/400) of reconstructions without hyper-recombinant

estimates exhibited lower genotype accuracy than the least accurate hyper-recombinant

simulation (range = 59.7%-92.1%). Similarly, for DGRP-derived 32-founder populations,

2.5% (10/400) of reconstructions without hyper-recombinant estimates exhibited lower

genotype accuracy than the least accurate hyper-recombinant simulation (range =

82.3%-92.8%).

Reducing sequencing coverage by an order of magnitude from 0.05X to 0.005X resulted

in more frequent hyper-recombinant reconstruction estimates, though overall median

genotype accuracy remained above 99% (Figure S9). Hyper-recombinant reconstructed

chromosomes exhibited genotype estimates with accuracy below 99%, while the

remaining simulations (with lower recombinant counts) achieved above 99% genotype

accuracy. For populations founded by 32 DGRP lines and sequenced at 0.005X

coverage, 14% of simulations produced hyper-recombinant estimates (N=56/400), of

which only 26.8% (N=15/56) surpassed 99% genotype accuracy (median=98.5%). The

remaining 86% of simulations (N=344/400) that were not hyper-recombinant retained

greater accuracy, with 89% of simulations resulting in at least 99% genotype accuracy

(median=99.5). Increasing the number of founding DGRP lines from 32 to 128 at 0.005X

coverage failed to eliminate hyper-recombinant estimates. With 128 founding lines,

14.5% of simulations were hyper-recombinant (N=58/400), of which 24.1% (N=14/58)

surpassed 99% genotype accuracy (median=98.6%). The 85.5% of simulations that

were not hyper-recombinant (N=342/400), exhibited accurate genotype estimates, with

86.5% (296/342) of simulations achieving over 99% genotype accuracy (median=99.6).

509   GWAS Simulation Accuracy

To report the power of a GWAS, we must first define a "true positive" result. Consider a

putative SNP identified by GWAS that is 50 kb from the causal SNP. Such a result

would be considered a false positive if the aim is to identify the exact responsible

nucleotide, but may be a true positive with respect to to identify an associated gene. To

cover both use cases, we describe a true positive in terms of both SNP-level resolution

(requiring an exact base-pair match), or region-level resolution (allowing for tolerance

up to 100kb between putative hits and the causal SNP). Additionally, it is unrealistic to

simply evaluate the single top result of a GWAS. Rather, a set of candidate loci may be

chosen for follow-up evaluation in confirmatory studies, and the probability of including

the causal SNP will increase as a greater number of putative SNPs are evaluated. Most

distinct changes in GWAS power occurred when including between most significant to

top 10 most significant candidate loci, after which power increased at a reduced rate, if

an asymptote was not already reached.

The estimated power of GWAS using a specific type of mapping population, i.e. the

fraction of simulations with a true positive, is shown in Figure 5. For simplicity, we focus

on GWAS power when including the top 10 most significant candidate loci—a

reasonable number of putative sites that may be investigated in follow-up studies.

$_5$ Hybrid Swarms founded by either 32 or 128 founding lines exhibited nearly

equivalent power compared to $_{50}$ outbred populations across all parameter

combinations. For common alleles, i.e. those segregating at 50% frequency, all outbred

populations achieved approximately 50% power to identify a causal variant with SNP-

level precision, and 70% power at the gene-level. Both inbred populations were highly

effective at detecting associations at the gene-level (99% and 99.8% for ILs and RILs,

respectively). SNP-level power was one fourth lower than gene-level power for RILs

(75.4%), but only marginally reduced for inbred lines (97.4%).

Power to detect associations is reduced when the causal allele is rare (segregating at

12.5% frequency). For such rare alleles, the gene-resolving power of ILs drops by

nearly half (to 54.8%), while RILs maintained high power (81.8%). All outbred

populations exhibited approximately 20% power to detect rare alleles at the gene-level.

Inbred lines were the sole frontrunner for identifying low frequency alleles with SNP-

level resolution (37.2%), followed by 128-founder Hybrid Swarm (10.8%), F50 outbred

(8.2%), 32-founder Hybrid Swarm (6%), and RILs (3.2%).

GWAS Genomic Inflation Factor

If individuals are assigned to case and control groups with equal probability, then the

resulting χ^2 statistics should follow the expected distribution. If individuals are not sorted

into groups randomly, i.e. allele state at a causal SNP dictates nonrandom group

assignment, then χ^2 values for that SNP should be inflated to some extent. Nonrandom

associations between a causal SNP and other loci can inflated test statistics across a

chromosome, or across a whole genome. The genome-wide inflation factor (λ) can be

expressed as the ratio of observed and expected median χ^2 values (Figure 6). Because

our simulations model a single causal SNP, λ is a reflection of greater-than-chance

associations arising due to linkage with the causal SNP being modeled, which can

serve as a proxy for false positive rate.

Because the median expected χ^2 statistic increases with sample size, we report λ_{1000}, a

sample-size-corrected value that is comparable across studies (Freedman *et al.* 2004).

We calculated λ as aggregated across three groups: linked, including only the

autosome arm containing the causal SNP; unlinked, including the unlinked autosome

that doesn't contain the causal SNP; and autosomal, for all sites across both autosomes

two and three

Inflation factor measured across autosomes two and three was greatest for ILs, followed

by 32-founder Hybrid Swarm, RILs, 128-founder Hybrid Swarm, and $_{50}$ Outbred

populations. This order was observed whether the causal allele was common or rare,

though with reduced values of λ at the lower allele frequency (Figure 7).

Only inbred populations displayed inflation on unlinked autosomes. When the causal

allele is common (50% frequency), inflation on unlinked sites was greater for Inbred

lines (median $\lambda = 1.17$, interquartile range or $IQR = 0.11$) than for RILs ($\lambda = 1.02, IQR =$

0.07). There was no inflation for unlinked chromosome in outbred populations, where

$\lambda = 1.0$ with varying degrees of dispersion ($IQR = 0.10, 0.06$ and 0.03, respectively, for

32-founder HS, 128-founder HS, and F50 outbred populations). Unlinked sites remained

inflated for ILs even when the causal allele was rare ($\lambda = 1.07, IQR = 0.09$). Distributions

for λ across an extended range of autosome groups, PVE and allele frequencies are

shown in Figure S11.

When we dissociated phenotype from genotype with purely random case-control

assignment (i.e. PVE was set to 0% in our simulations), λ was centered at 1 (Figure

S12). F_{50} outbred populations exhibited the lowest dispersion ($IQR = 0.02$), followed by

128-founder Hybrid Swarms ($IQR = 0.04$), RILs ($IQR = 0.06$), and 32-founder Hybrid

Swarms or ILs ($IQR = 0.07$ each).

Frequency of sites segregating at appreciable frequency

The number of SNPs segregating amongst DGRP haplotypes with at least a given MAF

strongly depends on the haplotype subset count for a given population (Figure 8). If only

considering SNPs segregating at or above a frequency of 12.5% on chromosome arm

2L, a population founded by 8 lines will yield approximately twice as many SNPs

compared to a population founded by 128 lines (N=8 lines yields a median of 140K

SNPs; N=128 lines yields a median of 71k SNPs). If the minimum MAF threshold is

instead set to 5%, then populations with a greater number of lines exhibit a greater

number of SNPs—with a maximum number of segregating sites with N=16 lines

(median of 231.6k SNPs), nearly as many for 128 lines (median of 194k SNPs), and

fewer for N=8 lines (median of 133k SNPs).

598   **Discussion**

Herein, we examined the feasibility and statistical properties of genome-wide

association mapping using the Hybrid Swarm, an outbred population derived from

limited and random outcrossing of an arbitrary number of founding strains. We show

that it is possible to accurately reconstruct whole genomes from Hybrid Swarm

populations using ultra-low coverage sequencing data (Figure 5). Genome-wide

association mapping using the Hybrid Swarm approach performs as well as mapping in

highly outbred F_{50} populations in a case-control GWAS framework (Figures 6,

Supplemental Figure S10). While mapping using the Hybrid Swarm approach generally

has reduced power compared to mapping using inbred lines (as would any outbred

population in general) a limited number of generations of recombination reduces false

positives arising from long-distance linkage disequilibrium present in founding strains

(Figure 7, Supplemental Figure S11). Together, our results demonstrate the feasibility

and potential of using the Hybrid Swarm approach for generating and genotyping

outbred mapping populations in a cost-effective and computationally-efficient (Figure 1)

manner.

Benefits of the Hybrid swarm Approach

The Hybrid Swarm approach is applicable to a wide variety of organisms and

experimental designs, conferring potential benefits over inbred reference panels. These

benefits are realized in three primary ways by: (1) allowing researchers to address

questions that require heterozygotes; 2) reducing labor and the influence of cage-effects

with random mating in a common environment; and 3) breaking down population

structure when incorporating individuals from divergent populations. These benefits are

possible due to the ability to reconstruct genomes accurately and in a cost-effective

manner for a large number of individuals.

Note that the Hybrid Swarm method is not limited to populations founded by inbred

lines, as the technique can be applied to populations where phased genomes are

available for all outbred founders. Research systems without inbred reference panels

can thus make an up-front investment of fully phasing founder genomes to realize

downstream savings of reconstructing progeny from low-coverage sequencing data.

Due to the relative ease of generating phased genomes from a variety of long-read

sequencing technologies (Pollard *et al.* 2018), the Hybrid Swarm method may enable

association mapping in a wide variety of organisms.

Representation of heterozygotes

One clear difference between inbred and outbred mapping populations is the presence

of heterozygotes. On the one hand, the presence of heterozygotes in outbred

populations decreases power to detect association relative to inbred lines for an (semi-)

additive allele with a given effect size (Figure 6, Supplemental Figure S10). However,

the reduced statistical power of association mapping in outbred populations may be

ameliorated by reduced inbreeding depression and by the ability to assess the

heterozygous effects of alleles.

The ability to assess heterozygous effects of alleles will provide valuable insights into

several interesting aspects of biology, such as the nature of dominance and the identity

of regulatory polymorphisms. An increased understanding of dominance relationships

and regulatory polymorphisms is important for advancing our understanding of

quantitative trait variation and evolution. For instance, several theoretical models have

shown that context dependent dominance of quantitative fitness traits can underlie the

stable maintenance of polymorphisms subject to seasonally variable (Wittmann *et al.*

2017) or sexually antagonistic (Connallon and Chenoweth 2019) selection. The ability

to efficiently map loci with context dependent dominance relationships will aid in the

understanding of the stability and abundance of polymorphisms maintained by these

forms of balancing selection.

Regulatory polymorphisms are known to underlie genetic variation in expression (Brem

et al. 2002; Cavet *et al.* 2003; Rockman and Kruglyak 2006) and this expression

variation can potentially be resolved to exact nucleotide differences (Grosveld *et al.*

1987; Rave-Harel *et al.* 1997; Bosma *et al.* 2002). The resulting differences in

expression can manifest as phenotypic changes to drive local adaptation (Kudaravalli

et al. 2009; Fraser *et al.* 2010; Fraser 2011, 2013). Allele-specific expression (ASE)

arising from cis-acting regulatory factors is a common mechanism to produce heritable

differences in expression (Yan *et al.* 2002; Cowles *et al.* 2002; Lo *et al.* 2003; Doss

2005). Because allelic expression biases are only produced (and detectable) in

heterozygotes, Hybrid Swarm populations facilitate the study of regulatory genetic

variation (i.e. ASE) as a driver of local adaptation in a variety of organisms.

Undirected outbreeding in a common environment

The Hybrid Swarm approach involves propagation of a single large outbred population

via undirected crossing. This design confers benefits over alternatives of either rearing

inbred lines separately or performing controlled crosses. First, a single population

reduces the influence of random block effects associated with rearing families or closely

related individuals in separate enclosures or defined areas. Second, random

outbreeding of a single population requires less labor compared to performing controlled

crosses or serial propagation of inbred lines. One drawback of the randomly outbred

method is susceptibility to loss of haplotypes due to genetic drift. The distribution of

haplotypes can also be skewed by line-specific differences in fitness or fecundity, with

such differences being observed for DGRP lines (Horváth and Kalinka 2016). To

attenuate haplotype dropout, it may be prudent to seed a Hybrid Swarm with a large

population of F1 hybrids produced by round-robin crosses. The F1 population would

then be followed by a limited number of generations (e.g., 4-5) of random outbreeding.

Hybrid Swarm breaks down population structure and linkage disequilibrium

Recombination between lines in the Hybrid Swarm approach allows for greater

dissection of functional polymorphisms segregating between genetically structured

populations. If an association study incorporates haplotypes from multiple distinct

source populations, causal variants would segregate along with other linked variants.

Thus, to identify genetic mechanisms of local adaptation and trait variation in general, it

is necessary to minimize false positives from linked non-causal loci. Corrections due to

relatedness can reduce the type I error rate to some degree (Yu *et al.* 2006; Price *et al.*

2010; Yang *et al.* 2014), and can be further reduced by a greater extent of

recombination. Within mapping populations with many haplotypes such as the DGRP,

long-distance linkage disequilibrium results from correlated occurrence of rare variants

(Huang *et al.* 2014), potentially contributing to false positives in GWAS. This is reflected

in our simulations by genome-wide inflation of λ, even across physically unlinked

chromosomes, whereas five generations of recombination were sufficient to reduce this

inflation (Figure 7).

Most notably, F_5 Hybrid Swarm populations performed equivalently to F_{50} outbred

population in a case-control GWAS framework. This is likely owed to the large number

of unique haplotypes within the Hybrid Swarm population, reducing the influence of long

distance LD, and in turn reducing false positive GWAS hits. One interpretation is that

only slightly recombinant populations comprised of a modest number of haplotypes are

sufficient representations of highly outbred (or wild) populations in a GWAS framework.

Inbred populations did exhibit greater power than outbred populations for identifying a

causal locus, although this result is to be expected. Because we simulated a purely

additive trait for which heterozygotes are equally likely to be assigned to either case or

control group, heterozygotes contribute no statistical signal of association. Accordingly,

for a causal allele segregating at 50% frequency, sample sizes for any outbred

populations will be effectively half that of an inbred population.

The Hybrid Swarm method is similar but distinct from advanced intercross populations

(AIPs), where AIPs result from crossing few lines (e.g., 8) for many generations

(Chesler 2014; Mackay and Huang 2018) and the Hybrid Swarm from crossing dozens
to hundreds of lines for few generations. The choice to use an AIP or hybrid swarm
population will influence the number of SNPs segregating at or above a desired minor
allele frequency (Figure 8). For an association test to detect a causal variant with single-
nucleotide precision, that variant must be segregating above a minor allele frequency
required to detect phenotypic association at a given effect size and sample size. If
sample size precludes sites segregating at a minor allele frequency below 1/8, then a
population founded by 8 haplotypes would yield the greatest number of variants. If
power is sufficient to detect association with alleles segregating above a frequency of
5%, then populations founded by 16+ lines would yield a greater number of variants
(Figure 8). In cases where only few founding haplotypes are available, an AIP may be
necessary, as the breakup of linkage disequilibrium can only be accomplished with
many generations of crosses instead of leveraging greater haplotype diversity.
726

Computational Considerations
The simulations conducted for this analysis were made feasible by three primary
innovations. First, the haplotype block file format allowed us to leverage information
redundancy between related individuals and store highly compressed, lossless
genotype information. With nearly 1/2000th the file size of a compressed VCF file,
haplotype block files greatly reduced both the disk storage footprint and time required
for disk write operations. Second, instead of performing forward-time simulations for
every single iteration of GWAS, permuted subsets of simulated populations allowed for
more rapid GWAS simulations. The format of haplotype block files facilitated
permutations of the ancestry contained within a population's mosaic haplotypes,
generating novel population genetic structure while preserving the forward-simulator's
influence of drift and meiotic recombination (Figure 2). Third, instead of extracting site-
specific genotypes for every individual, we decreased the number of computational
operations by performing aggregate counts across all sites between adjacent
recombination events in the population (Figure 3).
742

Importantly, selecting a subset of most-likely-ancestors results in maximum

computational complexity that remains constant with increasing number of founding

lines, instead of complexity increasing at a greater-than-linear rate. This means that the

larger the pool of unique haplotypes that an individual descends from, the greater

speedup of our pipeline relative to other methods. Although computational speed has

been shown to be reduced by haplotype pre-phasing (Howie *et al.* 2011), to our

knowledge, pre-phasing has not been demonstrated with ultra-low coverage sequencing

on the order of 0.005-0.05X. As a result, pre-phasing would likely require greater

sequencing effort, negating the benefit of low coverage reconstruction. Computational

search space can also be reduced if an individual's pedigree is known with certainty,

however controlled crosses can be laborious, and may lead to cage-specific effects.

Applying the Hybrid Swarm approach

At minimum, the Hybrid Swarm approach requires a sequenced set of individuals for

founding a recombinant population. Although our simulations presented here were

conducted with inbred founding lines, genome reconstructions can similarly be

performed with any phased genomes. For example, 16 phased outbred founders could

be treated as 32 independent haplotypes. Phased genomes are becoming increasingly

accessible with the advent of long-read sequencing platforms and phasing software

(Chin *et al.* 2016; Mostovoy *et al.* 2016; Seo *et al.* 2016), allowing this technique to be

applied to even more systems. Optionally, a recombination rate map for the population

can be provided, otherwise recombination is assumed to occur with equal user-defined

probability across any chromosome.

As a first step, power analyses using our rapid association test simulation pipeline will

inform choices of sample size and mapping population design (Figure 3). After

determining a feasible sample size for a given SNP of minimum percent variation

explained, researchers can evaluate the accuracy of low-coverage chromosome

reconstructions for a simulated proposed mapping population. Note that while we

performed association tests in a case/control framework, the relative power of the

Hybrid Swarm is expected to be the same for quantitative traits, which could garner

additional power from sampling individuals from phenotypic extremes (D. Li, Lewinger,

Gauderman, Murcray, & Conti, 2011).

For our simulations, we parameterized chromosome reconstructions using a maximum

of $N = 16$ most-likely ancestors (MLAs) and $S = 5000$ SNPs, which required less than 3

GB of memory and completed in under 5 minutes on a single core. However, these

values may not be ideal for all systems. It may be necessary to select greater number of

MLAs prior to reconstruction if haplotypes are difficult to differentiate due to being less

divergent (i.e. exhibiting lower θ_π) than those simulated here. For example,

reconstruction accuracy was low for coalescent-derived mapping populations modeled

with $\theta = 4 \times 10^{-5}$ (Supplemental Figure S8), which may reflect those of *C. elegans*

(Barriere and Félix 2005). Further, 5000 SNPs may be an over- or under-estimate of

those required in other systems. Because recombination between haplotypes can only

be inferred at sampled variable sites, SNP density directly influences how close inferred

breakpoints will be resolved with respect to their actual position. The models described

in Figure 1 can be used to estimate the memory and runtime required for a given

number of input ancestor haplotypes.

To evaluate whether low coverage sequencing data will yield accurate genotype

estimates for a given proposed mapping population, researchers can test reconstruction

accuracy *in silico*. We provide a convenient forward-simulation R script for this purpose

that generates output in the haplotype map format (Figure 2). Simulated individuals can

then be ran through the simulated sequencing and mapping pipeline at a desired level

of coverage. After generating simulated mapped individuals, researchers can optimize

the number of MLAs and HARP threshold that provide most effective MLA selection for

their mapping population (Figure 4). This step may reveal haplotypes that are

consistently problematic or inaccurately chosen, which can be excluded from further

simulations (and when generating the true mapping population). Researchers can then

perform chromosome reconstruction using the optimized MLA selection parameters and

evaluate whether accuracy is acceptable (Figure 5).

After performing chromosome reconstructions, a quality control step may be applied

whereby troublesome regions are masked. For example, a reconstructed chromosome

with a sequence of short recombination blocks could be masked prior to evaluating

genotyping accuracy or performing association testing. In our simulations, it was

surprisingly difficult to diagnose exact factors contributing to the least accurate

reconstructions. However, these highly recombinant reconstructions still achieved 90-

99% accuracy, suggesting that accuracy may be achieved even for anomalous hyper-

recombinant individuals (Figure 5). Optimized parameters can then be applied to a

genuine mapping population akin to the simulated one.

Conclusions

An outbred high-resolution mapping population that can be generated in little time is an

attractive option for researchers, but such mapping populations have been prohibited by

genotyping costs or computational requirements to impute genotypes from ultra-low

sequencing data. Our work demonstrates the feasibility of the Hybrid Swarm as a cost-

effective method of fine-scale genetic mapping in an outbred population and provides a

computationally efficient framework for GWAS power analysis.

References

Barrière, A. and Félix, M.-A. Natural variation and population genetics of
Caenorhabditis elegans (December 26, 2005), *WormBook*, ed. The *C. elegans*
Research Community, WormBook, doi/10.1895/wormbook.1.43.1,
http://www.wormbook.org.

Bergland A. O., H. Chae, Y.-J. Kim, and M. Tatar, 2012 Fine-Scale Mapping of Natural
Variation in Fly Fecundity Identifies Neuronal Domain of Expression and Function
of an Aquaporin. PLoS Genet. 8: e1002631.

Bosma P. J., J. R. Chowdhury, C. Bakker, S. Gantla, A. de Boer, *et al.*, 2002 The
Genetic Basis of the Reduced Expression of Bilirubin UDP-Glucuronosyltransferase
1 in Gilbert's Syndrome. N. Engl. J. Med. 333: 1171–1175.

Brem R. B., G. Yvert, R. Clinton, and L. Kruglyak, 2002 Genetic dissection of
transcriptional regulation in budding yeast. Science 296: 752–5.

Broad Institute, 2015a The Picard toolkit. https://broadinstitute.github.io/picard/.

Broad Institute, 2015b Genome Analysis Toolkit: Variant Discovery in High-Throughput
Sequencing Data. https://software.broadinstitute.org/gatk/.

Cavet G., P. S. Linsley, M. Mao, R. B. Stoughton, S. H. Friend, *et al.*, 2003 Genetics of
gene expression surveyed in maize, mouse and man. Nature 422: 297–302.

Cheng R., J. E. Lim, K. E. Samocha, G. Sokoloff, M. Abney, *et al.*, 2010 Genome-wide
association studies and the problem of relatedness among advanced intercross
lines and other highly recombinant populations. Genetics 185: 1033–1044.

Chesler E. J., D. R. Miller, L. R. Branstetter, L. D. Galloway, B. L. Jackson, *et al.*, 2008
The Collaborative Cross at Oak Ridge National Laboratory: Developing a powerful
resource for systems genetics. Mamm. Genome 19: 382–389.

Chesler E. J., 2014 Out of the bottleneck: the Diversity Outcross and Collaborative
Cross mouse populations in behavioral genetics research. Mamm. Genome 25: 3–
11.

Chia R., F. Achilli, M. F. W. Festing, and E. M. C. Fisher, 2005 The origins and uses of
mouse outbred stocks. Nat. Genet. 37: 1181–1186.

Chin C.-S., P. Peluso, F. J. Sedlazeck, M. Nattestad, G. T. Concepcion, *et al.*, 2016
Phased diploid genome assembly with single-molecule real-time sequencing. Nat.
Methods 13: 1050–1054.

Comeron J. M., R. Ratnappan, and S. Bailin, 2012 The Many Landscapes of
Recombination in Drosophila melanogaster. PLoS Genet. 8: 33–35.

Connallon T., and S. F. Chenoweth, 2019 Dominance reversals and the maintenance of
genetic variation for fitness. PLoS Biol. 17: 1–11.

Cowles C. R., J. N. Hirschhorn, D. Altshuler, and E. S. Lander, 2002 Detection of
regulatory variation in mouse genes. Nat. Genet. 32: 432–437.

Danecek P., A. Auton, G. Abecasis, C. A. Albers, E. Banks, *et al.*, 2011 The variant call
format and VCFtools. Bioinformatics 27: 2156–2158.

Doss S., 2005 Cis-acting expression quantitative trait loci in mice. Genome Res. 15:
681–691.

Fraser H. B., A. M. Moses, and E. E. Schadt, 2010 Evidence for widespread adaptive
evolution of gene expression in budding yeast. Proc. Natl. Acad. Sci. 107: 2977–
2982.

Fraser H. B., 2011 Genome-wide approaches to the study of adaptive gene expression
evolution. BioEssays 33: 469–477.

Fraser H. B., 2013 Gene expression drives local adaptation in humans. Genome Res.

23: 1089–1096.

Freedman M. L., D. Reich, K. L. Penney, G. J. McDonald, A. A. Mignault, *et al.*, 2004
Assessing the impact of population stratification on genetic association studies.
Nat. Genet. 36: 388–393.

Grosveld F., G. B. van Assendelft, D. R. Greaves, and G. Kollias, 1987 Position-
independent, high-level expression of the human beta-globin gene in transgenic
mice. Cell 51: 975–85.

Horváth B., and A. T. Kalinka, 2016 Effects of larval crowding on quantitative variation
for development time and viability in Drosophila melanogaster. Ecol. Evol. 6: 8460–
8473.

Howie B., J. Marchini, and M. Stephens, 2011 Genotype imputation with thousands of
genomes. G3 1: 457–70.

Huang X., M.-J. Paulo, M. Boer, S. Effgen, P. Keizer, *et al.*, 2011 Analysis of natural
allelic variation in Arabidopsis using a multiparent recombinant inbred line
population. Proc. Natl. Acad. Sci. U. S. A. 108: 4488–93.

Huang B. E., A. W. George, K. L. Forrest, A. Kilian, M. J. Hayden, *et al.*, 2012 A
multiparent advanced generation inter-cross population for genetic analysis in
wheat. Plant Biotechnol. J. 10: 826–839.

Huang W., A. Massouras, Y. Inoue, J. Peiffer, M. Ramia, *et al.*, 2014 Natural variation in
genome architecture among 205 Drosophila melanogaster Genetic Reference
Panel lines. Genome Res. 24: 1193–1208.

Kessner D., T. L. Turner, and J. Novembre, 2013 Maximum likelihood estimation of
frequencies of known haplotypes from pooled sequence data. Mol. Biol. Evol. 30:
1145–58.

King E. G., S. J. Macdonald, and A. D. Long, 2012a Properties and power of the
Drosophila synthetic population resource for the routine dissection of complex
traits. Genetics 191: 935–949.

King E. G., C. M. Merkes, C. L. McNeil, S. R. Hoofer, S. Sen, *et al.*, 2012b Genetic
dissection of a model complex trait using the Drosophila Synthetic Population
Resource. Genome Res. 22: 1558–1566.

Kover P. X., W. Valdar, J. Trakalo, N. Scarcelli, I. M. Ehrenreich, *et al.*, 2009 A
Multiparent Advanced Generation Inter-Cross to Fine-Map Quantitative Traits in
Arabidopsis thaliana, (R. Mauricio, Ed.). PLoS Genet. 5: e1000551.

Krämer N., N. Ranc, N. Meyer, M. Ouzunova, C. Lehermeier, *et al.*, 2014 Usefulness of
Multiparental Populations of Maize (Zea mays L.) for Genome-Based Prediction .
Genetics 198: 3–16.

Kudaravalli S., J. B. Veyrieras, B. E. Stranger, E. T. Dermitzakis, and J. K. Pritchard,
2009 Gene expression levels are a target of recent natural selection in the human
genome. Mol. Biol. Evol. 26: 649–658.

Lack J. B., C. M. Cardeno, M. W. Crepeau, W. Taylor, R. B. Corbett-Detig, *et al.*, 2015
The drosophila genome nexus: A population genomic resource of 623 Drosophila
melanogaster genomes, including 197 from a single ancestral range population.
Genetics 199: 1229–1241.

Li R., M. A. Lyons, H. Wittenburg, B. Paigen, and G. A. Churchill, 2005 Combining data
from multiple inbred line crosses improves the power and resolution of quantitative
trait loci mapping. Genetics 169: 1699–709.

Li H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan, *et al.*, 2009 The Sequence
Alignment/Map format and SAMtools. Bioinformatics 25: 2078–2079.

Li H., 2011 wgsim (short read simulator). https://github.com/lh3/wgsim.

Li H., 2013 Aligning sequence reads, clone sequences and assembly contigs with BWA-
MEM. arXiv:1303.3997.

Lo H. S., Z. Wang, Y. Hu, H. H. Yang, S. Gere, *et al.*, 2003 Allelic variation in gene
expression is common in the human genome. Genome Res. 13: 1855–62.

Mackay T. F. C., and W. Huang, 2018 Charting the genotype–phenotype map: lessons
from the Drosophila melanogaster Genetic Reference Panel. Wiley Interdiscip. Rev.
Dev. Biol. 7: e289.

MacKay T. F. C. C., S. Richards, E. A. Stone, A. Barbadilla, J. F. Ayroles, *et al.*, 2012
The Drosophila melanogaster Genetic Reference Panel. Nature 482: 173–178.

Monir M. M., and J. Zhu, 2017 Comparing GWAS Results of Complex Traits Using Full
Genetic Model and Additive Models for Revealing Genetic Architecture. Sci. Rep. 7:
38600.

Mostovoy Y., M. Levy-Sakin, J. Lam, E. T. Lam, A. R. Hastie, *et al.*, 2016 A hybrid
approach for de novo human genome sequence assembly and phasing. Nat.
Methods 13: 587–590.

Mott R., C. J. Talbot, M. G. Turri, A. C. Collins, and J. Flint, 2000 A method for fine
mapping quantitative trait loci in outbred animal stocks. Proc. Natl. Acad. Sci. 97:
12649–54.

Nikpay M., A. Goel, H. H. Won, L. M. Hall, C. Willenborg, *et al.*, 2015 A comprehensive
1000 Genomes-based genome-wide association meta-analysis of coronary artery
disease. Nat. Genet. 47: 1121–1130.

Paul R. Staab, Sha Zhu, Dirk Metzler, and Gerton Lunter, 2015 {scrm}: efficiently

simulating long sequences using the approximated coalescent with recombination.
Bioinformatics 31: 1680–1682.

Pollard M. O., D. Gurdasani, A. J. Mentzer, T. Porter, and M. S. Sandhu, 2018 Long
reads: their purpose and place. Hum. Mol. Genet. 27: R234–R241.

Price A. L., N. A. Zaitlen, D. Reich, and N. Patterson, 2010 New approaches to
population stratification in genome-wide association studies. Nat. Rev. Genet. 11:
459–463.

R Core Team, 2016 R: A Language and Environment for Statistical Computing

Rave-Harel N., E. Kerem, M. Nissim-Rafinia, I. Madjar, R. Goshen, *et al.*, 1997 The
molecular basis of partial penetrance of splicing mutations in cystic fibrosis. Am. J.
Hum. Genet. 60: 87–94.

Rockman M. V., and L. Kruglyak, 2006 Genetics of global gene expression. Nat. Rev.
Genet. 7: 862–872.

Seo J. S., A. Rhie, J. Kim, S. Lee, M. H. Sohn, *et al.*, 2016 De novo assembly and
phasing of a Korean human genome. Nature 538: 243–247.

Singh R., I. T. Lobina, M. Thomson, S. McCouch, C. Dilla-Ermita, *et al.*, 2013 Multi-
parent advanced generation inter-cross (MAGIC) populations in rice: progress and
potential for genetics research and breeding. Rice 6: 11.

Spencer C. C. A., Z. Su, P. Donnelly, and J. Marchini, 2009 Designing genome-wide
association studies: Sample size, power, imputation, and the choice of genotyping
chip. PLoS Genet. 5.

Srivastava A., A. P. Morgan, M. L. Najarian, V. K. Sarsani, J. S. Sigmon, *et al.*, 2017
Genomes of the Mouse Collaborative Cross. Genetics 206: 537–556.

Stevenson M., 2018 epiR: Tools for the Analysis of Epidemiological Data.
https://rdrr.io/cran/epiR/.

Wittmann M. J., A. O. Bergland, M. W. Feldman, P. S. Schmidt, and D. A. Petrov, 2017
Seasonally fluctuating selection can maintain polymorphism at many loci via
segregation lift. Proc. Natl. Acad. Sci. 114: E9932–E9941.

Wu Y., Z. Zheng, P. M. Visscher, and J. Yang, 2017 Quantifying the mapping precision
of genome-wide association studies using whole-genome sequencing data.
Genome Biol. 18: 1–10.

Yan H., W. Yuan, V. E. Velculescu, B. Vogelstein, and K. W. Kinzler, 2002 Allelic
variation in human gene expression. Science 297: 1143.

Yang J., N. A. Zaitlen, M. E. Goddard, P. M. Visscher, and A. L. Price, 2014 Advantages

and pitfalls in the application of mixed-model association methods. Nat. Genet. 46:
100–106.

Yang C., Y. Wang, W. Xu, Z. Liu, S. Zhou, *et al.*, 2018 Genome-wide association study
using diversity outcross mice identified candidate genes of pancreatic cancer.
Genomics 0–1.

Yu J., G. Pressoir, W. H. Briggs, I. V. Bi, M. Yamasaki, *et al.*, 2006 A unified mixed-
model method for association mapping that accounts for multiple levels of
relatedness. Nat. Genet. 38: 203–208.

Zhang J., K. Kobert, T. Flouri, and A. Stamatakis, 2014 PEAR: A fast and accurate
Illumina Paired-End reAd mergeR. Bioinformatics 30: 614–620.

Zheng C., M. P. Boer, and F. A. van Eeuwijk, 2015 Reconstruction of genome ancestry
blocks in multiparental populations. Genetics 200: 1073–1087.

Zheng C., M. P. Boer, and F. A. van Eeuwijk, 2018 Accurate genotype imputation in
multiparental populations from low-coverage sequence. Genetics 210: 71–82.

Figures

993
Figure 1. Resource usage of RABBIT during haplotype reconstruction.
All reconstructions involve the same simulated 2L chromosome arm comprised of four
haplotypes. Simulations included varied numbers of founding haplotypes (*N*) and a
randomly selected set of markers (number of SNPs, *S*, incremented in steps of 500). All
simulations included, at minimum, the four true haplotypes for the simulated individual.
In **A** and **B**, points depict the mean of empirical values (over 10 replicates) and gray
lines depict the defined regression models. Predicted peak memory usage and runtime
are displayed on a log scale over a greater range for number of founding haplotypes in
C and **D**, respectively.

1003
Figure 2. Basic structure of the forward simulator pipeline.
Inbred founding lines (**A**) are randomly intercrossed to produce a recombinant
population (**B**). Rapid generation of independent mapping populations is achieved by
random down-sampling (**C**) and permutation of ancestry (**D**). Population genetic data is
encoded in a highly compressed format **(E)** that references the positions of haplotype
blocks instead of genotypes at every site, enabling us to generate 500 mapping
populations for a given parameter combination. Individuals are probabilistically assigned
to case or control groups based on genotype at a randomly chosen causal SNP
segregating at a specified frequency.

Figure 3. Schematic for rapid association testing with the haplotype block files. For a given population represented by a haplotype map file (**A**), all SNPs between sorted breakpoints (indicated by dashed lines) will share identical aggregated haplotype frequencies (**B**). Haplotype frequencies are multiplied by a founder genotype matrix (**C**) where alleles are coded reference (black cells) and alternate (white cells). Conditional row sums of the resulting matrix (**D**) yields reference and alternate frequencies at each locus (**E**), to be used for χ^2 tests of independence.

1021
Figure 4. Optimization curves for Most-Likely-Ancestor (MLA) selection.
Increasing the upper limit for the number of MLAs chosen reduces the number of true
ancestors missed, similarly reducing the fraction of a given chromosome that is not
represented within the selected set of MLAs. Ancestors that fail to pass the HARP
threshold across all genomic windows are not selected, resulting in realized sets of
MLAs (Number of Ancestors Chosen) below the upper-limit allowed (x-axis). Data
shown reports means across 400 replicates made up of 100 simulated individuals (4
autosomes each for coalescent simulations, 4 autosome arms each for DGRP
simulations) per parameter combination. Coalescent-derived populations described
here were simulated with $N_e = 10^6$ and $\mu = 5 \times 10^{-9}$.

Figure 5. Accuracy of genome reconstruction pipeline for simulated F$_5$ Hybrid Swarm individuals.

Reconstructions were performed for populations simulated as being founded by either 32 or 128 inbred lines at 0.05X sequencing coverage with up to 16 MLAs as determined with a HARP threshold of 0.99. Accuracy, calculated as the per-chromosome fraction of variable sites with a correct diploid genotype estimate, is shown on logit-transformed scale. Values are coded depending on the number of estimated recombination events, with highly recombinant estimates ($\geq$10 recombination events) displayed as an $\times$. Each parameter combination includes 400 reconstructed autosomes (individual circles) for 100 simulated individuals. The coalescent-derived individuals displayed here were simulated with an effective population size of $N_e = 1 \times 10^6$ and mutation rate $\mu = 5 \times 10^{-9}$.

1045
Figure 6. Accuracy of simulated GWAS for various mapping populations.
Plots display the cumulative probability of including a causal SNP when selecting the
top N most significant SNPs, or 100kb windows around those SNPs, out of 500
simulated GWAS (each comprised of 5000 individuals phenotypically assigned in a
case-control framework). Homozygotes for the reference allele were assigned to the
case group with 45% probability, while homozygotes for the alternate allele were
assigned to the case group with 55% probability (a difference of 10%), and
heterozygotes are assigned to case and control groups with equal probability. **ILs**:
inbred lines. **RILs**: recombinant inbred lines. **HS**: Hybrid Swarm populations founded by
32 or 128 lines. **Outbred**: An F_{50} population founded by 128 lines.

1056

All autosomes ▤ Autosome arm with causal SNP ▢ Whole autosome without causal SNP

1057

Figure 7. Genomic Inflation Factor (GIF, λ_{1000}) for simulated GWAS with a causal allele
segregating at a specified frequency.
GIF is calculated genome-wide (across all autosomes); on the autosome arm containing
the causal allele (linked); and for sites on the autosome physically unlinked to the
causal allele. λ is calculated as the ratio of observed to expected χ^2 values, and a
correction is performed to produce the null expectation given the sample size had
actually been 1000 individuals (see Materials and Methods for details). Data are
averaged over 500 simulated GWAS (each comprised of 5000 individuals
phenotypically assigned in a case-control framework). Homozygotes for the reference
allele were assigned to the case group with 45% probability, while homozygotes for the
alternate allele were assigned to the case group with 55% probability (a difference of
10%), and heterozygotes are assigned to case and control groups with equal
probability. Boxes represent the median and interquartile range; whiskers extending to
the lower and upper bounds of the 95% quantiles. **ILs**: 128 Inbred Lines. **RILs**: 800
Recombinant Inbred Lines. **HS**: Hybrid Swarm with 32 or 128 founding lines. **Outbred**:
F_{50} population founded by 128 inbred lines.

1074

1075
Figure 8. Counts of variable sites depending on number of founding DGRP haplotypes
Each point represents the number of sites segregating at or above a given minor allele
frequency threshold when drawing *N* haplotypes, with 20 replicates per parameter
combination. With a minimum minor allele frequency (MAF) of 12.5%, a population
founded by eight haplotypes exhibits approximately double the number of variable sites
compared to a population founded by 128 haplotypes. With a minimum MAF of 5%,
populations with eight founding haplotypes present with fewer SNPs compared to
populations founded by 16 or more haplotypes.

Population	N Founders	ρ	$\overline{\Delta}$	σ_Δ
DGRP	128	0.986	-0.015	0.25
DGRP	32	0.502	0.17	2.15
Coalescent	128	0.956	-0.17	0.44
Coalescent	32	0.759	-0.31	1.26

1084

Table 1. Accuracy of estimated number of recombination events following chromosome
reconstruction.
A high concordance correlation coefficient (Lin's ρ) indicates agreement between
estimated and true recombination counts for 400 reconstructed chromosomes
(coalescent-derived populations) or chromosome arms (DGRP-derived populations).
Coalescent-derived populations are described across a range of values for effective
population size N_e and mutation rate μ. $\overline{\Delta}$ and σ_Δ denote mean and standard deviation,
respectively, of difference between estimated and true recombination counts.
Reconstructions were performed with a maximum of 16 most-likely-ancestors with a
HARP threshold of 0.99 (see methods for more details).

1095
Figure S1. Recombination probability functions used for simulated individuals.
Recombination is modeled as a Poisson process, with position sampled from linear
interpolation of recombination rates measured in *Drosophila melanogaster* by Comeron
et al. (2012). The frequency of recombination samples cumulative map distance (inset,
e.g. a 99 cM chromosome is modeled as a Poisson variable with an expected value of
$\lambda = 0.99$). For DGRP-derived individuals, recombination was simulated for full
chromosomes two and three, and reconstructions were then conducted independently
for arms 2L, 2R, 3L, and 3R.

1104
Figure S2. Optimization curves for Most-Likely-Ancestor inclusion, by count, in SCRM-
derived F_5 hybrid swarm individuals
The number of missed true ancestors is shown as a function of the number of
ancestors chosen across a range of HARP threshold values (0.9 to 0.999), effective
population sizes (N_e) and mutation rates (μ). Values are averaged across 400
reconstructed autosomes (from 100 individuals) per parameter combination.

Figure S3. Optimization curves for Most-Likely-Ancestor inclusion, by chromosome representation, in simulated SCRM-derived F_5 hybrid swarm individuals. The proportion of the chromosome not covered by the chosen ancestors is shown as a function of the number of ancestors chosen for populations founded by either 32 or 128 inbred founding lines across a range of HARP threshold values (0.9 to 0.999), effective population sizes (N_e) and mutation rates (μ). Each line summarizes the arithmetic mean fraction of sites where the true ancestor is not included within the inferred set of Most-Likely-Ancestors. Values are averaged across 400 reconstructed autosomes (from 100 individuals) per parameter combination.

1121
Figure S4. Optimization curves for Most-Likely-Ancestor (MLA) selection for DGRP-
derived F_5 hybrid swarm individuals.
Effectiveness is shown for populations founded by either 32 or 128 inbred founding lines
across a range of HARP threshold values (0.9 to 0.999), for two levels of sequencing
coverage. Values are averaged across 400 reconstructed autosomes (from 100
individuals) per parameter combination.

Figure S5. Distribution of Most-Likely-Ancestor counts for simulated, Coalescent-derived, 32-founder F_5 hybrid swarm individuals. The mean value ± 1 standard deviation is shown by the solid line and ribbon, respectively, across a range of HARP threshold values (0.9 to 0.999), effective population sizes (N_e) and mutation rates (μ). The number of Most-Likely-Ancestors dictates the computational complexity (runtime and memory requirements) of chromosome reconstruction. Each parameter combination includes 400 chromosomes (from 100 simulated individuals) simulated with 0.05X sequencing coverage.

1137
Figure S6. Distribution of Most-Likely-Ancestor counts for simulated, Coalescent-
derived, 128-founder F_5 hybrid swarm individuals.
The mean value ± 1 standard deviation is shown by the solid line and ribbon,
respectively, across a range of HARP threshold values (0.9 to 0.999), effective
population sizes (N_e) and mutation rates (μ). The number of Most-Likely-Ancestors
dictates the computational complexity (runtime and memory requirements) of
chromosome reconstruction. Each parameter combination includes 400 chromosomes
(from 100 simulated individuals) simulated with 0.05X sequencing coverage.

1146
Figure S7. Distribution of Most-Likely-Ancestor counts for simulated, DGRP-derived F_5
hybrid swarm individuals.
The mean value ± 1 standard deviation is shown by the solid line and ribbon,
respectively, for populations founded by either 32 or 128 inbred lines, across a range of
HARP threshold values (0.9 to 0.999) and two levels of sequencing coverage. Each
parameter combination includes 400 chromosomes (from 100 simulated individuals)
simulated with 0.05X sequencing coverage.
1154

1155
Figure S8. Accuracy of genome reconstruction for simulated, coalescent-derived F_5
hybrid swarm individuals.
Reconstructions were performed for populations simulated as being founded by either
32 or 128 inbred lines for various effective population sizes (N_e) and mutation rates (μ).
Accuracy is represented on a logit scale, as most points occur above 90%.
Reconstructed chromosomes that are predicted to exhibit $\geq$10 recombination events are
denoted by an X. Each parameter combination includes 400 reconstructed
chromosomes (from 100 simulated individuals).

1164

Figure S9 Accuracy of genome reconstruction for simulated, DGRP-derived F_5 hybrid
swarm individuals.
Reconstructions were performed for populations simulated as being founded by either
32 or 128 inbred lines for two levels of ultra-low sequencing coverage. Accuracy is
represented on a logit scale, as most points occur above 90%. Accuracy values are
marked depending on the number of estimated recombination events, with highly
recombinant estimates ($\geq$10 recombination events) displayed as an $\times$. Each parameter
combination includes 400 reconstructed chromosomes (from 100 simulated individuals).
1173

1174
1175

1176
Figure S10. Probability of selecting a causal SNP (or a nearby neighbor) in simulated
GWAS.
Each line represents the fraction of GWAS simulations (out of 500 total GWAS, each
comprised of 5000 individuals in a case-control framework) where the causal SNP is
selected within N most significant regions. Case-control status was assigned based on
reference allele dosage at a randomly selected causal SNP segregating with frequency
of 50%, 25%, or 12.5%. For 10% PVE, homozygotes for the reference allele were
assigned to the case group with 45% probability, while homozygotes for the alternate
allele were assigned to the case group with 55% probability (a difference of 10%). For
5% PVE, homozygotes for the reference allele were assigned to the case group with
47.5% probability, while homozygotes for the alternate allele were assigned to the case
group with 52.5% probability (a difference of 5%). All heterozygotes (and any individuals
modeled with 0% PVE, irrespective of genotype) were equally likely to be assigned to
case or control group (a difference of 0%).
1191

1192
1193 **Figure S11**. Genomic Inflation Factor (GIF, λ_{1000}) in simulated DGRP GWAS as a
1194 function of minor allele frequency and percent variation explained.
1195 GIF is stronger with greater PVE, and is elevated on the arms which contain (and are
1196 directly and most strongly linked to) the SNP associated with case and control status.
1197

1198

1199
1200 **Figure S12**. Genomic Inflation Factor (GIF, λ_{1000}) in simulated DGRP GWAS with no
1201 genotype-phenotype link.
1202 As the percent variation explained is 0% (and thus, case and control status is randomly
1203 assigned), GIF centers around 1.0 as expected.
1204
1205

Population	Coverage	N Founders	N_e	μ	ρ	$\overline{\Delta}$	σ_{Δ}
DGRP	0.005X	128	-	-	0.201	3.06	3.11
DGRP	0.005X	32	-	-	0.201	3.15	2.81
DGRP	0.05X	128	-	-	0.986	-0.015	0.25
DGRP	0.05X	32	-	-	0.502	0.17	2.15
Coalescent	0.05X	128	10^4	1×10^{-9}	0.029	-2.72	1.60
Coalescent	0.05X	32	10^4	1×10^{-9}	0.219	-1.84	1.50
Coalescent	0.05X	128	10^5	1×10^{-9}	0.288	-1.57	1.64
Coalescent	0.05X	32	10^5	1×10^{-9}	0.761	-0.53	0.99
Coalescent	0.05X	128	10^6	1×10^{-9}	0.742	-0.46	1.09
Coalescent	0.05X	32	10^6	1×10^{-9}	0.754	-0.42	1.22
Coalescent	0.05X	128	10^4	5×10^{-9}	0.203	-2.03	1.73
Coalescent	0.05X	32	10^4	5×10^{-9}	0.622	-0.89	1.17
Coalescent	0.05X	128	10^5	5×10^{-9}	0.652	-0.64	1.41
Coalescent	0.05X	32	10^5	5×10^{-9}	0.782	-0.26	1.16
Coalescent	0.05X	128	10^6	5×10^{-9}	0.956	-0.17	0.44
Coalescent	0.05X	32	10^6	5×10^{-9}	0.759	-0.31	1.26
Coalescent	0.05X	128	10^4	1×10^{-8}	0.238	-1.65	1.86
Coalescent	0.05X	32	10^4	1×10^{-8}	0.745	-0.66	1.05
Coalescent	0.05X	128	10^5	1×10^{-8}	0.776	-0.34	1.12
Coalescent	0.05X	32	10^5	1×10^{-8}	0.846	-0.25	0.93
Coalescent	0.05X	128	10^6	1×10^{-8}	0.937	-0.22	0.56
Coalescent	0.05X	32	10^6	1×10^{-8}	0.833	-0.32	0.95

Table S1. Accuracy of estimated number of recombination events following chromosome reconstruction. A high concordance correlation coefficient (Lin's ρ) indicates agreement between estimated and true recombination counts for 400 reconstructed chromosomes (coalescent-derived populations) or chromosome arms (DGRP-derived populations). Coalescent-derived populations are described across a range of values for effective population size N_e and mutation rate μ. $\overline{\Delta}$ and σ_{Δ} denote mean and standard deviation, respectively, for difference between estimated and true recombination count. Reconstructions were performed with a maximum of 16 most-likely-ancestors with a HARP threshold of 0.99 (see methods for more details).