Accurate, ultra-low coverage genome reconstruction and association studies in Hybrid Swarm mapping populations

Cory A. Weller[1,2*], Susanne Tilk[3], Subhash Rajpurohit[4], Alan O. Bergland[1]

[1] Department of Biology, University of Virginia, Charlottesville, Virginia 22904

[2] National Human Genome Research Institute, National Institutes of Health, Bethesda, MD 20892

[3] Department of Biology, Stanford University, Stanford CA 94305

[4] Department of Biological and Life Sciences, Ahmedabad University, Ahmedabad, India 380009

* To whom correspondence should be addressed: cory.weller@nih.gov

Abstract

Genetic association studies seek to uncover the link between genotype and phenotype, and often utilize inbred reference panels as a replicable source of genetic variation.
However, inbred reference panels can differ substantially from wild populations in their genotypic distribution, patterns of linkage-disequilibrium, and nucleotide diversity. As a
result, associations discovered using inbred reference panels may not reflect the genetic basis of phenotypic variation in natural populations. To address this problem,
we evaluated a mapping population design where dozens to hundreds of inbred lines are outbred for few generations, which we call the Hybrid Swarm. The Hybrid Swarm
approach has likely remained underutilized relative to pre-sequenced inbred lines due to the costs of genome-wide genotyping. To reduce sequencing costs and make the
Hybrid Swarm approach feasible, we developed a computational pipeline that reconstructs accurate whole genomes from ultra-low-coverage (0.05X) sequence data
in Hybrid Swarm populations derived from ancestors with phased haplotypes. We evaluate reconstructions using genetic variation from the Drosophila Genetic Reference
Panel as well as variation from neutral simulations. We compared the power and precision of GWAS using the Hybrid Swarm, inbred lines, recombinant inbred lines, and
highly outbred populations across a range of allele frequencies, effect sizes, and genetic architectures. Our simulations show that these different mapping panels vary in
their power and precision, largely depending on the architecture of the trait. The Hybrid Swam and RILs outperform inbred lines for quantitative traits, but not for monogenic
ones. Taken together, our results demonstrate the feasibility of the Hybrid Swarm as a cost-effective method of fine-scale genetic mapping.

Introduction

Genetic mapping studies seek to describe the link between genotype and phenotype. For experimental crosses, mapping was traditionally conducted by scoring the

phenotypes of recombinant offspring descended from a limited number of parental lines (Lander and Botstein 1989). While such QTL mapping studies can have high power to

detect association, they offer minimal mapping resolution (Cheng *et al.* 2010), often detecting broad regions of phenotypic association. If linkage disequilibrium is lowered,

regions of association can be resolved at the gene or nucleotide level (Li *et al.* 2005; Rockman and Kruglyak 2008), as in GWAS of large outbred populations (Nikpay *et al.*

2015; Wu *et al.* 2017; Monir and Zhu 2017). However, GWAS suffer from reduced power to detect associations (Long and Langley 1999), necessitating a large sample

size relative to QTL mapping (Spencer *et al.* 2009).

To generate higher resolution mapping populations than the traditional biparental crosses, Multiparental Populations (MPPs) are becoming increasingly used. By crossing

together multiple inbred lines, one can produce genetically diverse mapping populations without sampling wild individuals. MPPs are commonly used for the dissection of

complex traits in model organisms (Chesler *et al.* 2008; Kover *et al.* 2009; King *et al.* 2012) and agriculturally important crops (Huang *et al.* 2012a; Singh *et al.* 2013; Krämer

et al. 2014). Although MPPs are statistically powerful (Valdar *et al.* 2006; Svenson *et al.* 2012), they may have limited utility in addressing basic questions about the evolutionary

forces affecting causal variants in natural populations (Long *et al.* 2014) and are often restricted to existing mapping panels in a limited number of taxa.

66
The mapping resolution of MPPs depends on the extent of linkage disequilibrium.

Mapping resolution is improved by allowing for more recombination between haplotypes or by substituting extensive recombination for increased haplotype diversity (Mott *et al.*

2000; Chia *et al.* 2005). In the latter approach, by crossing dozens to hundreds of inbred lines for a limited number (~5) of generations and subsequently phenotyping and

genotyping outbred individuals, heterozygous mapping populations can be generated quickly with sufficiently reduced LD to potentially detect associations with high

resolution. We refer to such an outbred mapping population as a Hybrid Swarm. Whether Hybrid Swarm mapping populations can be generated efficiently, and whether

association mapping using such a population is useful, remain open questions.

To address these questions, we evaluate a method to reconstruct phased whole genomes from large Hybrid Swarm populations using ultra-shallow sequencing (<0.05X)

and assessed the power and precision of association mapping using a Hybrid Swarm relative to common alternatives [8-way MPP – DSPR (Long *et al.* 2014); inbred lines –

DGRP (MacKay *et al.* 2012)]. First, we developed and tested our genome reconstruction method by generating whole genomes for thousands of simulated Hybrid

Swarm individuals. Our simulated genomes draw from natural variation in the *Drosophila melanogaster* Genetic Reference Panel (DGRP), or variation generated from

coalescent models representing a broad range of genetic diversity parameters for common model systems. We show that the Hybrid Swarm approach allows for highly

accurate genotyping (average 99.9% genotypic accuracy) from ultra-low-coverage

whole-genome individual-based sequencing. We confirm the accuracy of genome
90 reconstructions using experimental animals from a Hybrid Swarm constructed in the lab
from 128 DGRP founder lines. We then simulated a range of genetic architectures to
92 discern the accuracy and precision of association mapping in the Hybrid Swarm
compared to inbred lines, recombinant inbred lines, and a highly outbred population.
94 GWA simulations confirm that inbred MPPs have the highest power to detect
associations, outbred Hybrid Swarm populations have intermediate power, and inbred
96 reference panels have the lowest power, particularly for quantitative traits. In addition,
we show that outbred Hybrid Swarms eliminate spurious signals of association that
98 arise using inbred lines. Together, our results show the feasibility of cost-effective
association mapping in a large outbred multi-parental population and provides tools for
100 genome-reconstruction and simulation.

102   **Methods**

Simulating a Hybrid Swarm. As a case study of low-coverage genome reconstruction
104 in a model system, we simulated a Hybrid Swarm using sequenced inbred lines from
the Drosophila Genetic Reference Panel (MacKay *et al.* 2012) as available from the
106 *Drosophila* Genome Nexus (Lack *et al.* 2015). We only included the 128 lines (out of
205) with the least amount of missing genotype data. We removed insertions, deletions,
108 and sites with more than two alleles. Any heterozygous genotype calls were masked as
missing data.
110
To generate simulated populations, we developed a forward-simulator in R that stores
112 ancestral haplotype block maps instead of genotypes. See Figure 1 for depictions of
mapping populations simulated. We simulated Hybrid Swarms through random mating
114 over five non-overlapping generations at a population size of 10,000. Sexual
reproduction was simulated by random sampling of recombinant gametes from male-
116 female pairs. Recombination frequency was modeled as a Poisson process with an
expected value $\lambda = \Sigma$(Morgans) per chromosome. For simulations of *Drosophila*
118 populations based on DGRP chromosomes, recombination occurred only in females,
with recombination frequency and position based on values from Comeron et al (2012).
120 See extended methods and Supplemental Figure S1 for additional detail.

122 We also simulated Hybrid Swarms using haplotypes derived from whole-genome
coalescent simulation using *scrm* (Staab *et al.* 2015). We generated simulations across
124 a range of diversity and recombination levels, encompassing the values for many
common model systems. For populations founded by simulated haplotypes,
126 recombination occurred in both sexes, with recombination occurring uniformly across
each chromosome.
128
We used *wgsim* (Li 2011) to simulate 100 bp reads at two average read depths (0.005X
130 and 0.05X). We specified a base error rate of 0.001 and an indel fraction of 0.
Remaining *wgsim* parameters were left as default. We assembled paired end reads
132 using *PEAR* (Zhang *et al.* 2014) and separately aligned the assembled and
unassembled groups to a reference genome with *bwa* 0.7.14 using the BWA-MEM

134 algorithm (Li 2013). Simulated *Drosophila* reads were mapped to the *D. melanogaster* reference genome v5.39. After converting mapped reads to compressed BAM format
136 with *samtools* 1.3.1 (Li *et al.* 2009), we removed PCR duplicates with *Picard tools* 2.0.1 ("Picard toolkit" 2019).
138

Genome reconstruction for a Hybrid Swarm. Genome reconstruction for Hybrid
Swarm individuals can take advantage of Hidden Markov Models developed for standard MPPs (Zheng *et al.* 2015b). In these models, the probability of each diplotype
– a unique diploid combination of founding haplotypes - is calculated. Calculating these probabilities is feasible with smaller numbers of founding lines (< 8, typically) but grows
at a quadratic rate. Therefore, increasing the number of founding haplotypes from 8 to 128 incurs orders of magnitude more computational effort (Supplemental Figure S2)
and is not feasible. To make chromosome reconstructions in the Hybrid Swarm computationally tractable, we developed a method of accurately selecting a subset of
most likely ancestors for any single chromosome. It is important to note that the computational requirement of selecting a subset of founding chromosomes requires that
a Hybrid Swarm only be propagated for a limited number of generations, especially if the number of founding lines is large.
152

We used the software package *HARP* (Kessner *et al.* 2013) to identify the most likely
154 founding lines based on the number of genomic windows with high probabilities of ancestry. *HARP* was originally developed to estimate haplotype frequencies from
156 pooled sequence data, and we co-opted it to assess relative likelihood that any founder contributed to a genomic window in a focal Hybrid Swarm chromosome. We ran *HARP*
158 with non-overlapping 100 kb windows with a minimum frequency cutoff 0.0001. Ideally, any given window would have four (or fewer) founders with high proportions of ancestry
160 (~25%) and unambiguous haplotype paths (Supplemental Figure S3). However, empirical evaluation of Harp output for sample chromosomes showed that many had
162 more than four founders with high proportion of ancestry and difficult-to-resolve haplotypes (Supplemental Figure S4).
164

We developed a heuristic method to identify the set of most likely ancestors. We ranked
all possible founders by their contribution to the focal chromosome. To do this, we counted the number of windows with extreme likelihoods of ancestry (i.e., greater than a
specified quantile) for each potential founding haplotype. Potential founding lines were then ranked by the number of windows passing this threshold. We examined two
measures of effectiveness for this method across a range of quantile threshold values (90%, 95%, 99%, and 99.9%) when selecting up to a maximum number of most likely
ancestral founders. The first measure is the number of true ancestral founders excluded; the second measure is the fraction of the chromosome derived from
ancestors missing from the selected subset.

We used the Mathematica package RABBIT (Zheng *et al.* 2015a) to perform chromosome reconstructions. This package has been shown to be accurate for
genotype estimation at sequencing coverage at 0.05X for a variety of multiparent populations (Zheng *et al.* 2018) but, the best of our knowledge, only for MPPs with

twelve or fewer potential founders. To input the observed genotype of each recombinant
       individual, we counted reference and alternate reads at variable sites (i.e. polymorphic
182   among the list of potential founders) using the Genome Analysis Toolkit
       ASEReadCounter tool (Broad Institute 2015). Because it is not possible to make
184   confident homozygote genotype calls from low coverage sequencing data where most
       sites are observed only once and or twice, we encoded individuals with only reference
186   or alternate alleles as "1N" or "2N"; observed heterozygotes were encoded as "12". We
       sampled 5000 highly informative SNPs per individual per chromosome for genome
188   reconstruction. Informative SNPs were identified as those at low frequency amongst
       potential founders identified in the HARP step. We ran RABBIT independently for each
190   chromosome using the Viterbi decoding function under the "joint model" with all other
       RABBIT parameters left at default. RABBIT output was converted to a phased
192   chromosome haplotype map, which we then used to extract and concatenate genotype
       information from a VCF file containing founder genotypes.
194
       We evaluated the accuracy of reconstruction by calculating genotype accuracy and the
196   number of inferred recombination events. To calculate genotype accuracy, we
       measured the fraction of sites where the estimated diploid genotype is identical to the
198   originally simulated diploid genotype. We only examined accuracy on the autosomes.
       To measure accuracy of estimated frequency of recombination events, true and
200   estimated recombination counts were first summed over both copies of each
       chromosome in a simulated individual. We then calculated Lin's concordance
202   correlation coefficient ρ between the true and estimated recombination counts using the
       epi.ccc function of the *R* package *epiR* (Stevenson 2018). An overview of our
204   reconstruction pipeline is shown in Supplemental Figure S5.

206   **Chromosome reconstruction with STITCH.** We estimated genotypes with alternative
       approach, STITCH, which is capable of imputing genotypes without reference panels
208   (Davies *et al.* 2016). To assess genotype estimation accuracy using STITCH, we
       generated and mapped simulated reads for chromosome 2L at 0.05X coverage for 5000
210   individuals in a F_5 Hybrid Swarm population (N=10000 individuals per generation).
       Because STITCH draws inference from haplotypes between related individuals,
212   genotype estimate accuracy is expected to increase with greater numbers of sequenced
       individuals. To capture the effect of sample size on STITCH accuracy, we ran STITCH
214   for sample sizes from 100 to 5000 sequenced individuals. We calculated genotype
       estimate accuracy as the fraction of variable sites with a correct diploid genotype
216   estimate using a custom R script. Because STITCH memory requirements increase with
       greater values of *k* (founding haplotypes), it was not possible to evaluate *k*=128
218   founders, or to use the *diploid* estimation model. As a result, we limited our evaluations
       to 32-founder populations, using the *pseudoHaploid* model.
220
       Generating a real Hybrid Swarm population. We generated a hybrid swarm mapping
222   population through undirected mating of all Drosophila Genetic Reference Panel
       (DGRP) inbred lines (Mackay et al. 2012) in large population cages (6' x 6' x 6').
224   Populations were seeded with males and females from each line and were expanded
       for four generations on cornmeal-molasses media. Cages reached an approximate final

population size of ~500,000 as determined by a volumetric assay of dead adults. Generations were discrete: we removed egg-laden media from the cages and replaced

egg-laden media in emptied cages after removal of the previous generation.

Library preparation and read mapping. We homogenized individual flies in 350 µL Lysis Buffer RLT Plus using 2-3 1 mm beads in a bead shaker for 2 minutes. DNA and

RNA were extracted using the AllPrep DNA/RNA Micro Kit (Qiagen product number 80284). Following DNA extraction, we purified samples with Ampure XP Beads

(Beckman Coulter product number A63880). We prepared 1 µL of DNA at ~2.5 ng/µL sequencing using a modified Nextera protocol developed by Baym *et al.* (2015),

indexing samples with custom primers, and selecting fragments between 450-500 bp in length using a SizeSelect e-Gel. As a final step to amplify the prepared DNA

sequencing libraries, we ran all size selected samples through additional 5 rounds of PCR. Each PCR reaction used 5 µL template DNA, 0.6 µL of 100 mM forward and

reverse primers (custom synthesized by IDT), 10µL of KAPA HiFi Ready Mix (KAPA Biosystem product number KK2611/2612), and 3.8 µL nuclease free water. Our PCR

protocol included 5 minutes of initial denaturation at 95°C followed by 4 rounds of 20 seconds denaturation (98°C), 20 seconds annealing (62°C), 30 seconds elongation

(72°C), followed by a final elongation at 72°C for 2 minutes. Following PCR amplification, we purified DNA libraries using Ampure XP beads and quantified

concentrations on a Life Technologies Qubit spectrophotometer and Agilent Bioanalyzer. We sequenced the same library preparations at low (~0.05X) and high (~8)

coverage on separate Illumina sequencing runs. Reads were mapped to the reference genome using the same methods as our simulations.

     Empirical evaluation of genotype accuracy in the Hybrid Swarm. We tested if

RABBIT genotype estimates accurately reflected genotypes from higher coverage data for six individuals. First, we called genotypes in the high coverage samples using a

heuristic based on reference and alternate allele counts produced by GATK's ASEReadCounter to call genotypes. Sites with at least three reference and three

alternate allele bearing short-reads were called as heterozygous; any site with at least six reference reads and zero alternate reads (or the converse) was called as

homozygous. All other sites with fewer six reads were excluded. We calculated accuracy as the proportion of sites where the low-coverage genotype derived from

reconstruction matched our higher-coverage genotype call. We note that our measurement of genotype accuracy from the empirical data is conservative as we

expect a moderate fraction of true heterozygous sites to be called as homozygous.

Simulating mapping populations for GWAS. We performed GWAS on simulations of various types of Hybrid Swarm designs and contrasted the signal of association using

these designs to that from a simulated 8-way MPP (akin to the DSPR), and the DGRP. Simulation of GWAS proceeded in two steps.

     For every type of genetic architecture or mapping population explored, we simulated

270   500 mapping populations comprised of 5000 individuals. Full genomes for individuals were generated using the forward simulation framework described above. We used the

272   DGRP as the founder chromosomes. One important note is that the genotypes we use
      for GWAS are the "true" genotypes and are not affected by genotyping error. This
274   decision was made to reduce computation time. Because genotyping error is randomly
      distributed throughout the genome (and rare on a per-site basis, see Figure 2), the use
276   of perfect genotype data should not bias the results of our simulations.

278   Next, we assigned phenotypes to each simulated individual. We assign case or control
      phenotypes to individuals based on a liability model, where an observable binary
280   phenotype is produced from an underlying continuous trait, known as its liability (Xu and
      Atchley 1996). Under this model, phenotypic variance will increase with additional
282   causal loci. We contrasted mapping results for a monogenic trait and a polygenic trait,
      using loci with large and small effects, assigning individuals to case or control groups
284   based on their genotype at causal loci. We randomly selected a causal locus (or
      multiple loci for the polygenic model) from all variable sites in the mapping population.
286   We assigned the effect size of this locus to be proportional to its frequency: low
      frequency variants had a large effect and common variants had a small effect
288   (Supplemental Figure S6). To achieve a population size of 5000, it is necessary to
      sample from the 128 inbred lines or 800 recombinant inbred lines with replacement.
290

      To test for association between phenotype and each SNP, we performed a χ^2 test of
292   independence for reference and alternate allele counts between case and control
      groups. For inbred mapping populations, we corrected for non-independent allele draws
294   by dividing the χ^2 value by two.

296   **Assessing GWAS accuracy.** To measure GWAS accuracy, we generated Receiver
      Operator Characteristic (ROC) curves and calculated Area Under the Curve (AUC).
298   Briefly, we calculated the true positive rate (sensitivity) as a function of false positive
      rate (1-specificty) using the *R* package *pROC* (Turck *et al.* 2011). In our simulations,
300   only the causal SNP may be considered a true positive. We summarize GWAS
      accuracy by averaging ROC curves from all replicate simulations, generating a single
302   representative ROC curve following Marigorta *et al.* (2018). We tested for statistical
      differences in the ROC curves between different mapping designs using a Wilcoxon test
304   on the distribution of simulated AUC values. We performed multiple-testing correction
      for the tests comparing AUC distributions using the Bonferroni method.
306

      Genomic inflation factor. We calculated the genomic inflation factor (GIF) to evaluate
308   the role of mapping design and genetic architecture on the overall signal of association.
      We evaluated GIF on chromosomes linked and unlinked to causal loci. We calculated
310   the GIF by correcting for sample size following Freedman et al (2004).

312   **Data availability statement.** The raw sequence data for our six reconstructed real-life
      hybrid swarm individuals is available at SRA BioProject accession PRJNA691163. The
314   most up-to-date code associated with this project is available on GitHub:
      https://github.com/cory-weller/HS-reconstruction-gwas and an archived version of our
316   repository is available at Zenodo (DOI: 10.5281/zenodo.4472955). This repository
      includes a containerized version of the reconstruction pipeline along with a test data-set.

Results

Genome reconstruction for a Hybrid Swarm. First, we evaluated our most-likely
322 ancestor selection algorithm, which selects a minimum representative set of Most-
Likely-Ancestors (MLAs). This step required optimizing a discrimination threshold for
324 likelihood values calculated with HARP (Kessner *et al.* 2013, see methods). We found
that classifying any potential founder as an MLA by counting the number of
326 chromosomal windows with observed likelihoods in the top 5% (threshold of 0.95)
identified all potential founders with high accuracy (Supplemental Figure S7). The
328 threshold of 0.99 performed slightly better when 128 founding lines were used to found
a population, while the threshold of 0.95 performed slightly better with only 32 founding
330 lines. HARP thresholds of 0.95 and 0.99 perform similarly well across a range of
simulated population parameters, suggesting this is a reasonable starting point for
332 optimizing the pipeline for other populations (Supplemental Figure S8A). Reducing
sequencing coverage by a factor of ten (from 0.05X to 0.005X) resulted in similar levels
334 of ancestor selection accuracy (Supplemental Figure S8B), demonstrating that
additional coverage may not necessarily improve reconstruction accuracy. For
336 simplicity, we next conducted all genome reconstructions with the 0.99 threshold at a
simulated sequencing coverage of 0.05X.

Next, we evaluated the accuracy of genotype calls following genome reconstruction
340 based on the most-likely ancestors identified for any individual chromosome. In general,
genotype reconstruction from simulated F_5 Hybrid Swarm individuals was very accurate:
342 the median percent of sites with correctly estimated genotypes was greater than 99.9%
whether the population was founded by 32 or 128 founding lines or made use of DGRP
344 or simulated haplotypes (Figure 2, Supplemental Figure S10). For simulations founded
by DGRP lines, 80.5% of reconstructed chromosomes from 32-founder populations
346 exhibited > 99.9% accuracy, with the remaining 19.5% of reconstructions contributing to
a long tail with a minimum of 84.5%. Increasing the number of founding lines to 128
348 resulted in genotype accuracy above 99% for all cases (minimum: 99.4%), with 83% of
reconstructed chromosomes achieving greater than 99.9% accuracy. Reducing
350 sequencing coverage by an order of magnitude from 0.05X to 0.005X resulted in more
frequent over-estimation of recombination, though overall median genotype accuracy
352 remained above 99% (Supplemental Figure S9). Using simulated haplotypes, we also
show that reconstruction accuracy improves with genetic diversity (Supplemental Figure
354 S10).

356 We also examined the accuracy of our estimates of recombination number, per
individual. The number of recombination events estimated from was generally high but
358 differed between Hybrid Swarm populations with different numbers of founders.
Reconstructions of individuals from Hybrid Swarm populations with more founders
360 generated more accurate estimates of recombination count (Lin's concordance
correlation coefficient: 98% and 50% for DGRP 128- and 32-way Hybrid Swarms,
respectively; see Table 1). For 32-founder populations, DGRP-derived reconstructions

tended to over-estimate recombination counts. Results on recombination count accuracy from simulated haplotypes are reported in Supplemental Table 1.

Although genome reconstruction using RABBIT accurately calls genotypes, and generally estimates the number of recombination events well, we observe some
reconstructions that are hyper-recombinant (i.e. > 10 recombination events for our simulations of an F5, Figures 2 and 3). These hyper-recombinant individuals had among
the lowest genotyping accuracy, and this was the case for both DGRP-derived and coalescent-derived hybrid swarm individuals. The cause of these hyper-recombinant
reconstructions is not always clear, but frequently appears due to RABBIT's estimation switching between closely-related haplotypes. Because the number of inferred
recombination events is determined by the number of generations of recombination, hyper-recombinant individuals (or regions of a genome) can be easily identified and
removed as part of quality control (see Erickson et al 2020).

We contrasted the accuracy of genotype calls made with RABBIT to those made with an alternative tool, STITCH (Davies *et al.* 2016). STITCH does not rely on priors of parental
haplotypes, but instead performs imputation and phasing based on patterns of linkage among the entire set of sequenced Hybrid Swarm recombinants. As expected, STITCH
accuracy improved with greater sample size of N sequenced individuals (Figure S11), providing 68.5% genotype accuracy with N=100, increasingly approximately linearly
until 94.5% accuracy at N=3000, and providing 99% accuracy with N=4000.

Genome reconstruction and genotype calls of real Hybrid Swarm individuals sequenced at ultra-shallow (~0.05X) and high (~8) coverage were highly concordant (~95%; Figure
3). Accuracy is particularly high for reconstructed chromosomes that predict fewer than 10 recombination events.

GWAS accuracy. We evaluated GWAS accuracy for several configurations of Hybrid
392 Swarms and compared these to simulations of GWAS using an 8-way RILs (modeled after the DSPR), and an inbred reference panel (the DGRP). Examples of
394 representative Manhattan plots for a single causal locus of large effect are shown in Figure 4. RILs tend to result in a strong and wide association peak near the causal
396 allele, while other populations result in only a small number of sites (typically on or neighboring the causal locus) with p-values below 10^{-5}.

To assess overall performance of these mapping populations across hundreds of
400 simulations, we generated Receiver Operator Characteristic (ROC) curves and calculated Area Under the Curve (AUC; Figure 5). In general, recombinant inbred lines
402 outperformed other mapping designed. RILs were only significantly outperformed for a single-locus trait of large effect mapped in a 128-founder inbred reference panel, though
404 both populations approach a median AUC of 1.0 (Figure 5B, median RIL AUC = 0.997, median 32-founder Inbred Line AUC = 0.999, p=1.101×10^{-6}). Inbred reference panels
406 consistently had the lowest AUC for multi-locus traits (Figure 5C-F). We suspect that the decreased accuracy of association mapping using inbred lines is due to linkage to
408 causal sites, as well as spurious long-distance linkage (see below, *Genomic Inflation*

Factor). See Supplemental Table S2 for comparisons between all mapping population
410   types.

412   Hybrid Swarm populations, composed of either 32- or 128-founders intermated for 5
       generations had intermediate performance (Figure 5), and generally resembled fully
414   outbred (F50) populations composed of 128 founders. We also evaluated the GWAS
       performance when crossing 32 or 128 founders to generate F1 and F2 mapping
416   populations (Supplemental Figure S12). For single-locus simulations, the number of
       generations of recombination did not influence accuracy, although 32-founder
418   populations performed better than 128-founders. For multi-locus simulations, GWAS
       accuracy improved with more generations of recombination, and this effect was more
420   pronounced with 10-locus traits compared to 5-locus traits. Statistical tests comparing
       different models can be found in Supplemental Table S2.
422
       Genomic Inflation Factor. An alternative way to assess the quality of GWAS using
424   different mapping designs is to compare the Genomic Inflation Factor. This metric
       describes how much the genome-wide distribution of test statistics differs from a null
426   expectation. Values greater than 1 can indicate an excess of low p-values and can
       reflect population structure (Reich and Goldstein 2000) or polygenicity (Yang *et al.*
428   2011). We calculated the genomic inflation factor for any given mapping population and
       genetic architecture genome-wide (across all autosomes), on the autosome arm
430   containing the causal allele (linked), and for sites on the autosome physically unlinked
       to the causal allele.
432
       The relative ranking of GIF remained the same between a mono- and polygenic
434   architecture with Inbred Lines producing the greatest GIF, and outbred F50 populations
       producing the lowest GIF (Figure 6). Hybrid Swarm populations showed intermediate
436   GIF. Interestingly, Inbred Lines display elevated GIF even for sites physically unlinked
       to causal loci, suggesting spurious long-distance linkage disequilibrium. This inflation on
438   unlinked chromosomes for inbred lines persists when filtering out sites below 5%
       frequency (Supplemental Figure S13), suggesting the pattern cannot be attributed to
440   low-frequency alleles alone. Elevated GIF on unlinked chromosomes when using inbred
       panels such as the DGRP suggests caution in interpreting GWAS using this type of
442   mapping population, and could contribute to the observed differences in mapping
       results of the same trait studied in both inbred lines and an advanced Hybrid Swarms
444   derived from the DGRP (Huang *et al.* 2012b, 2020). The Hybrid Swarm, propagated for
       at least two generations, was able to decrease GIF at unlinked loci and by five
446   generations the problem of elevated GIF on unliked chromosomes was ameliorated.

448   **Discussion**

       Here, we show that the Hybrid Swarm is a viable method for association mapping using
450   experimental outbred populations. We first demonstrate that ultra-low coverage,
       individual-based whole genome sequencing can be used to reconstruct accurate diploid
452   genomes for species across a range of diversity levels (Figure 2, Supplemental Figure
       S10), and we validate the quality of genome reconstruction using real Hybrid Swarm
454   individuals sequenced at both high and low coverages (Figure 3). Genome

reconstruction with a reduced subset of most-likely-ancestors is computationally
456 efficient (Supplemental Figure S2), and not dependent upon the number of individuals
genotyped unlike STITCH (Supplemental Figure S11), enabling a wide range of
458 experimental designs. Next, we demonstrate that association mapping using the Hybrid
Swarm can match or outperform various types of inbred lines and completely outbred
460 populations (Figures 4, 5, Supplemental Figure S12, Supplemental Table S2). Lastly,
we have described a set of computational tools to simulate various mapping populations
462 (including genome reconstructions, and GWAS), enabling this method to easily be
applied to a variety of organisms.
464

Considerations of the Hybrid Swarm approach. The Hybrid Swarm approach is
466 applicable to a wide variety of organisms and experimental designs, conferring potential
benefits over inbred reference panels. These benefits are realized in three primary ways
468 by: 1) allowing researchers to address questions that require heterozygotes; 2) reducing
labor and the influence of block-effects; and 3) breaking down linkage-disequilibrium
470 and population structure that arises from the sampling of individuals from the wild.
These benefits are possible due to the ability to accurately reconstruct genomes from
472 ultra-low-coverage sequencing data, i.e. from cost-effective diluted DNA library
preparation for a large number of individuals (Baym *et al.* 2015).
474

We evaluated simulated Hybrid Swarms founded by 32 or 128 founders to determine
476 the effect of founding haplotype count on reconstruction accuracy and GWAS
performance. One might expect 32-founder populations to yield more accurate
478 reconstructions than 128-founder populations, because it is simpler to make a correct
inference out of a smaller pool of ancestors. Yet, 32-founder populations were less
480 accurate in our DGRP simulations (Figure 2), and for some coalescent-based
simulations (Supplemental Figure S10). The difference in reconstruction accuracy is
482 perhaps due to the inverse relationship between SNP frequency and information
content. In a 128-way cross, rare information-rich SNPs are only likely to exist and be
484 sampled during low-coverage sequencing if nucleotide diversity is high. The DGRP
haplotypes happen to exist in a parameter space with higher nucleotide diversity,
486 resulting in 128-way reconstructions being more accurate. For GWAS, 32-founder
Hybrid Swarm populations tended to perform better than 128-founder populations if the
488 simulated trait was controlled by a single locus (Figure 5). Taking these two factors into
consideration, broadly speaking, a Hybrid Swarm with fewer founding haplotypes may
490 be more appropriate for reconstructing genomes when genetic diversity in the founding
lines is low, or if the trait of interest is presumably controlled by one or few loci.
492 Conversely, a greater number of founders may be desired when genetic diversity is
high, or the trait of interest is likely to be complex.
494

The Hybrid Swarm method is not limited to populations founded by inbred lines, as the
496 technique can be applied to populations where phased genomes are available for all
outbred founders. Phasing the outbred founders could be accomplished in two basic
498 ways. For instance, phased genomes could be generated using a variety of long-read
sequencing technologies (Pollard *et al.* 2018). Alternatively, trio- and quartet-phasing
methods (Patterson *et al.* 2015), coupled with high coverage sequence data from the

outbred parents plus a limited (≤2) number of F1 offspring, would enable accurate phasing of parents to be used for reconstruction of downstream recombinant genomes.

One experimental consideration is the effects of drift and selection that operates during the generation of the Hybrid Swarm (e.g., Thépot *et al.* (2014)). However, it is not clear
that such processes will affect the ability to reconstruct genomes nor the specific outcome of GWAS, unless the trait of interest was under strong selection itself. What
may be more likely is the reduction of haplotypic diversity due to drift or selection will slightly decrease power to detect associations, as we see in the differences between
32- and 128-way crosses (Supplemental Figure S10). The distribution of haplotypes can also be skewed by line-specific differences in fitness or fecundity, with such differences
being observed for DGRP lines (Horváth and Kalinka 2016). To attenuate haplotype dropout, it may be prudent to seed a Hybrid Swarm with a large population of directed
F1 hybrids produced by round-robin crosses. The F1 population would then be followed by a limited number of generations (e.g., 4-5) of random outbreeding. This approach
was used by Erickson *et al* (2020) and resulted in a relatively even distribution of founding haplotypes after 5 generations.
518
We compared genotype calls from our low-coverage reconstruction pipeline to genotype
520 calls from higher coverage data for six experimental Hybrid Swarm individuals (Figure 3). Our results show high concordance between reconstructed genomes and high
522 coverage genotypes, particularly if reconstructions with 10 or more inferred recombination events are excluded. Researchers wishing to minimize the impact of
524 inaccurate genotype estimates could implement a stringent limit on inferred recombination counts.
526
Representation of heterozygotes. One clear difference between inbred and outbred
528 mapping populations is the presence of heterozygotes. On the one hand, the presence of heterozygotes in outbred populations decreases power to detect association relative
530 to inbred lines for an (semi-) additive allele with a given effect size (Figure 5). This basic statistical effect may also be influenced by differences in realized genetic variance
532 between inbred and outbred populations (Genissel *et al.* 2004). However, the reduced statistical power of association mapping in outbred populations may be ameliorated by
534 reduced inbreeding depression (Lee *et al.* 2017) and by the ability to assess the heterozygous effects of alleles.
536
The ability to assess heterozygous effects of alleles will provide valuable insights into
several interesting aspects of biology, such as the nature of dominance and the identity of regulatory polymorphisms governing allele-specific expression in a variety of
heterogenous genomes. An increased understanding of dominance relationships and regulatory polymorphisms is important for advancing our understanding of quantitative
trait variation and evolution. For instance, several theoretical models have shown that context dependent dominance of quantitative fitness traits can underlie the stable
maintenance of polymorphisms subject to seasonally variable (Wittmann *et al.* 2017) or sexually antagonistic (Connallon and Chenoweth 2019) selection. The ability to
efficiently map loci with context dependent dominance relationships will aid in the

understanding of the stability and abundance of polymorphisms maintained by these forms of balancing selection.

Regulatory polymorphisms are known to underlie genetic variation in expression (Brem *et al.* 2002; Cavet *et al.* 2003; Rockman and Kruglyak 2006) and this expression
variation can potentially be resolved to exact nucleotide differences (Grosveld *et al.* 1987; Rave-Harel *et al.* 1997; Bosma *et al.* 2002). The resulting differences in
expression can manifest as phenotypic changes to drive local adaptation (Kudaravalli *et al.* 2009; Fraser *et al.* 2010; Fraser 2011, 2013). Allele-specific expression (ASE)
arising from cis-acting regulatory factors is a common mechanism to produce heritable differences in expression (Yan *et al.* 2002; Cowles *et al.* 2002; Lo *et al.* 2003; Doss
2005). Because allelic expression biases are only produced (and detectable) in heterozygotes, Hybrid Swarm populations facilitate the study of regulatory genetic
variation (i.e. ASE) as a driver of local adaptation in a variety of organisms.

Undirected outbreeding in a common environment. The Hybrid Swarm approach involves propagation of a single large outbred population via undirected crossing. This
design confers benefits over alternatives of either rearing inbred lines separately or performing controlled crosses. The relative value of these benefits may vary across
organisms and experimental designs.

Individuals within a Hybrid Swarm are reared in a common environment. This reduces the influence of random block effects associated with rearing families or closely related
individuals in separate enclosures or defined areas. Therefore, studies examining genotype-by-environment interactions can dramatically expand the number of
environmental factors used because all individuals were previously reared in a common environment, enabling a Hybrid Swarm to be randomly scatted across environmental
treatments. This feature is in contrast to rearing each inbred family across all environments, a practice which generally limits the number of environmental treatments
used. Rearing individuals in a common environment also likely reduces genotype-environment correlation that could be exacerbated by vial effects.
578

Random outbreeding of a single population can require less labor compared to
580 performing controlled crosses or serial propagation of inbred lines. The relative value of these benefits is likely to be species specific. For instance, for larger animals (e.g.,
582 mice), controlled crosses may be logistically easier than a randomly mating swarm. For many other species, however, the Hybrid Swarm would be logistically advantageous.
584 The cost of maintaining inbred reference panels also varies among taxa, most notably between those species which can be kept as seeds or in a cryogenically preserved
586 state. For other species, such as flies, the monetary and environmental cost of maintaining hundreds to thousands of inbred lines may be prohibitive.
588

One drawback draw-back of the Hybrid Swarm is the lack of genetic replicates for
590 phenotyping. Inbred panels allow for a single genotype to be phenotyped multiple times, reducing the effects of error associated with phenotyping and micro-environmental
592 variation (Mackay and Huang 2018). Because this benefit is typically not possible in the

Hybrid Swarm (unless clones can be propagated), the Hybrid Swarm method
594 represents a tradeoff, reducing the influence of block effects while increasing error associated with phenotyping.
596

Hybrid Swarm breaks down population structure and linkage disequilibrium.
Recombination between lines in the Hybrid Swarm approach allows for greater dissection of functional polymorphisms segregating between genetically structured
populations. If an association study incorporates haplotypes from multiple distinct source populations, causal variants would segregate along with other linked variants.
Thus, to identify genetic mechanisms of local adaptation and trait variation in general, it is necessary to minimize false positives from linked non-causal loci. Corrections due to
relatedness can reduce the type I error rate to some degree (Yu *et al.* 2006; Price *et al.* 2010; Yang *et al.* 2014), and can be further reduced by a greater extent of
recombination. Therefore, the Hybrid Swarm approach allows researchers to tailor the mapping population to the traits of interest, by perhaps selecting founders from diverse
origins to test specific hypotheses about the distribution of functional variation in the wild.
610
A limited number of rounds of recombination during the propagation of the Hybrid
612 Swarm also reduces long-distance linkage disequilibrium within a panel of inbred lines derived from a single locality. Long-distance LD results from correlated occurrence of
614 rare variants (Huang *et al.* 2014), potentially contributing to false positives in GWAS (Figure 6). Our simulations showed a genome-wide inflation of p-values, even across
616 physically unlinked chromosomes, for inbred panels and this pattern persisted even when rare variants were excluded (Supplemental Figure S13). Two or more generations
618 of recombination were sufficient to reduce this inflation (Figure 6), at least when using association statistics that do not explicitly account for genetic structure in the sample.
620 Whether such long-distance linkage disequilibrium substantially affects the results of GWAS in practice, remains an open question.
622
Conclusions. Here, we demonstrate the feasibility of genome-reconstruction in Hybrid
624 Swarm populations derived from many founding haplotypes and evaluate the power of this approach for association mapping. Our results suggest that the Hybrid Swarm
626 approach can be a useful alternative to other MPP breeding designs and is applicable to model and non-model organisms.
628

630 **Author Contributions**. **CW**: conceptualization, data curation, methodology, formal analysis, software, validation, visualization, writing the manuscript; **ST**: investigation,
632 review of manuscript; **SR**: investigation; **AOB**: conceptualization, funding acquisition, project administration, resources, supervision, review and editing of the manuscript.
634
<div align="center">

References

</div>
636

Baym M., S. Kryazhimskiy, T. D. Lieberman, H. Chung, M. M. Desai, *et al.*, 2015
638 Inexpensive multiplexed library preparation for megabase-sized genomes. PLoS

One 10: 1–15.

640 Bosma P. J., J. R. Chowdhury, C. Bakker, S. Gantla, A. de Boer, *et al.*, 2002 The
Genetic Basis of the Reduced Expression of Bilirubin UDP-Glucuronosyltransferase

642 1 in Gilbert's Syndrome. N. Engl. J. Med. 333: 1171–1175.

Brem R. B., G. Yvert, R. Clinton, and L. Kruglyak, 2002 Genetic dissection of

644 transcriptional regulation in budding yeast. Science 296: 752–5.

Broad Institute, 2015 Genome Analysis Toolkit: Variant Discovery in High-Throughput

646 Sequencing Data

Cavet G., P. S. Linsley, M. Mao, R. B. Stoughton, S. H. Friend, *et al.*, 2003 Genetics of

648 gene expression surveyed in maize, mouse and man. Nature 422: 297–302.

Cheng R., J. E. Lim, K. E. Samocha, G. Sokoloff, M. Abney, *et al.*, 2010 Genome-wide

650 association studies and the problem of relatedness among advanced intercross
lines and other highly recombinant populations. Genetics 185: 1033–1044.

652 Chesler E. J., D. R. Miller, L. R. Branstetter, L. D. Galloway, B. L. Jackson, *et al.*, 2008
The Collaborative Cross at Oak Ridge National Laboratory: Developing a powerful

654 resource for systems genetics. Mamm. Genome 19: 382–389.

Chia R., F. Achilli, M. F. W. Festing, and E. M. C. Fisher, 2005 The origins and uses of

656 mouse outbred stocks. Nat. Genet. 37: 1181–1186.

Comeron J. M., R. Ratnappan, and S. Bailin, 2012 The Many Landscapes of

658 Recombination in Drosophila melanogaster. PLoS Genet. 8: 33–35.

Connallon T., and S. F. Chenoweth, 2019 Dominance reversals and the maintenance of

660 genetic variation for fitness. PLoS Biol. 17: 1–11.

Cowles C. R., J. N. Hirschhorn, D. Altshuler, and E. S. Lander, 2002 Detection of

662 regulatory variation in mouse genes. Nat. Genet. 32: 432–437.

Danecek P., A. Auton, G. Abecasis, C. A. Albers, E. Banks, *et al.*, 2011 The variant call

664 format and VCFtools. Bioinformatics 27: 2156–2158.

Davies R. W., J. Flint, S. Myers, and R. Mott, 2016 Rapid genotype imputation from

666 sequence without reference panels. Nat. Genet. 48: 965–969.

Doss S., 2005 Cis-acting expression quantitative trait loci in mice. Genome Res. 15:

668 681–691.

Fraser H. B., A. M. Moses, and E. E. Schadt, 2010 Evidence for widespread adaptive

670 evolution of gene expression in budding yeast. Proc. Natl. Acad. Sci. 107: 2977–
2982.

672 Fraser H. B., 2011 Genome-wide approaches to the study of adaptive gene expression
evolution. BioEssays 33: 469–477.

674 Fraser H. B., 2013 Gene expression drives local adaptation in humans. Genome Res.
23: 1089–1096.

676 Genissel A., T. Pastinen, A. Dowell, T. F. C. Mackay, and A. D. Long, 2004 No
Evidence for an Association between Common Nonsynonymous Polymorphisms in

678 Delta and Bristle Number Variation in Natural and Laboratory Populations of
Drosophila melanogaster. Genetics 166: 291–306.

680 Grosveld F., G. B. van Assendelft, D. R. Greaves, and G. Kollias, 1987 Position-
independent, high-level expression of the human beta-globin gene in transgenic

682 mice. Cell 51: 975–85.

Horváth B., and A. T. Kalinka, 2016 Effects of larval crowding on quantitative variation

684 for development time and viability in Drosophila melanogaster. Ecol. Evol. 6: 8460–

8473.

686   Huang B. E., A. W. George, K. L. Forrest, A. Kilian, M. J. Hayden, *et al.*, 2012a A multiparent advanced generation inter-cross population for genetic analysis in
688   wheat. Plant Biotechnol. J. 10: 826–839.

Huang W., S. Richards, M. A. Carbone, D. Zhu, R. R. H. Anholt, *et al.*, 2012b Epistasis
690   dominates the genetic architecture of Drosophila quantitative traits. Proc. Natl. Acad. Sci. 109: 15553–15559.

692   Huang W., A. Massouras, Y. Inoue, J. Peiffer, M. Ramia, *et al.*, 2014 Natural variation in genome architecture among 205 Drosophila melanogaster Genetic Reference
694   Panel lines. Genome Res. 24: 1193–1208.

Huang W., T. Campbell, M. A. Carbone, W. E. Jones, D. Unselt, *et al.*, 2020 Context-
696   dependent genetic architecture of Drosophila life span. PLoS Biol. 18: e3000645.

Kessner D., T. L. Turner, and J. Novembre, 2013 Maximum likelihood estimation of
698   frequencies of known haplotypes from pooled sequence data. Mol. Biol. Evol. 30: 1145–58.

700   King E. G., C. M. Merkes, C. L. McNeil, S. R. Hoofer, S. Sen, *et al.*, 2012 Genetic dissection of a model complex trait using the Drosophila Synthetic Population
702   Resource. Genome Res. 22: 1558–1566.

Kover P. X., W. Valdar, J. Trakalo, N. Scarcelli, I. M. Ehrenreich, *et al.*, 2009 A
704   Multiparent Advanced Generation Inter-Cross to Fine-Map Quantitative Traits in Arabidopsis thaliana, (R. Mauricio, Ed.). PLoS Genet. 5: e1000551.

706   Krämer N., N. Ranc, N. Meyer, M. Ouzunova, C. Lehermeier, *et al.*, 2014 Usefulness of Multiparental Populations of Maize (Zea mays L.) for Genome-Based Prediction .
708   Genetics 198: 3–16.

Kudaravalli S., J. B. Veyrieras, B. E. Stranger, E. T. Dermitzakis, and J. K. Pritchard,
2009 Gene expression levels are a target of recent natural selection in the human genome. Mol. Biol. Evol. 26: 649–658.

Lack J. B., C. M. Cardeno, M. W. Crepeau, W. Taylor, R. B. Corbett-Detig, *et al.*, 2015 The drosophila genome nexus: A population genomic resource of 623 Drosophila
melanogaster genomes, including 197 from a single ancestral range population. Genetics 199: 1229–1241.

Lander E. S., and S. Botstein, 1989 Mapping mendelian factors underlying quantitative traits using RFLP linkage maps. Genetics 121: 185.

Lee Y. C. G., Q. Yang, W. Chi, S. A. Turkson, W. A. Du, *et al.*, 2017 Genetic architecture of natural variation underlying adult foraging behavior that is essential
for survival of Drosophila melanogaster. Genome Biol. Evol. 9: 1357–1369.

Li R., M. A. Lyons, H. Wittenburg, B. Paigen, and G. A. Churchill, 2005 Combining data
722   from multiple inbred line crosses improves the power and resolution of quantitative trait loci mapping. Genetics 169: 1699–709.

724   Li H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan, *et al.*, 2009 The Sequence Alignment/Map format and SAMtools. Bioinformatics 25: 2078–2079.

726   Li H., 2011 wgsim (short read simulator). GitHub repository: https://github.com/lh3/wgsim

728   Li H., 2013 Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM

730   Lo H. S., Z. Wang, Y. Hu, H. H. Yang, S. Gere, *et al.*, 2003 Allelic variation in gene

expression is common in the human genome. Genome Res. 13: 1855–62.

732 Long A. D., and C. H. Langley, 1999 The power of association studies to detect the contribution of candidate genetic loci to variation in complex traits. Genome Res. 9:
734 720–731.

Long A. D., S. J. Macdonald, and E. G. King, 2014 Dissecting complex traits using the
736 Drosophila Synthetic Population Resource. Trends Genet. 30: 488–495.

Mackay T. F. C., and W. Huang, 2018 Charting the genotype-phenotype map: lessons
738 from the Drosophila melanogaster Genetic Reference Panel. Wiley Interdiscip. Rev. Dev. Biol. 7: e289.

740 MacKay T. F. C. C., S. Richards, E. A. Stone, A. Barbadilla, J. F. Ayroles, *et al.*, 2012 The Drosophila melanogaster Genetic Reference Panel. Nature 482: 173–178.

742 Marigorta U. M., J. A. Rodríguez, G. Gibson, and A. Navarro, 2018 Replicability and Prediction: Lessons and Challenges from GWAS. Trends Genet. 34: 504–517.

744 Monir M. M., and J. Zhu, 2017 Comparing GWAS Results of Complex Traits Using Full Genetic Model and Additive Models for Revealing Genetic Architecture. Sci. Rep. 7:
746 38600.

Mott R., C. J. Talbot, M. G. Turri, A. C. Collins, and J. Flint, 2000 A method for fine
748 mapping quantitative trait loci in outbred animal stocks. Proc. Natl. Acad. Sci. 97: 12649–54.

750 Nikpay M., A. Goel, H. H. Won, L. M. Hall, C. Willenborg, *et al.*, 2015 A comprehensive 1000 Genomes-based genome-wide association meta-analysis of coronary artery
752 disease. Nat. Genet. 47: 1121–1130.

Patterson M., T. Marschall, N. Pisanti, L. van Iersel, L. Stougie, *et al.*, 2015 WhatsHap :
754 Weighted Haplotype Assembly for Future-Generation Sequencing Reads. J. Comput. Biol. 22: 498–509.

756 Picard toolkit, 2019 Broad Institute, GitHub Repository: https://github.com/broadinstitute/picard; Broad Institute.

758 Pollard M. O., D. Gurdasani, A. J. Mentzer, T. Porter, and M. S. Sandhu, 2018 Long reads: their purpose and place. Hum. Mol. Genet. 27: R234–R241.

760 Price A. L., N. A. Zaitlen, D. Reich, and N. Patterson, 2010 New approaches to population stratification in genome-wide association studies. Nat. Rev. Genet. 11:
762 459–463.

R Core Team, 2016 R: A Language and Environment for Statistical Computing
764 Rave-Harel N., E. Kerem, M. Nissim-Rafinia, I. Madjar, R. Goshen, *et al.*, 1997 The molecular basis of partial penetrance of splicing mutations in cystic fibrosis. Am. J.
766 Hum. Genet. 60: 87–94.

Reich D. E., and D. B. Goldstein, 2000 Detecting association in a case-control study
768 while correcting for population stratification. Genet. Epidemiol. 20: 4–16.

Rockman M. V., and L. Kruglyak, 2006 Genetics of global gene expression. Nat. Rev.
770 Genet. 7: 862–872.

Rockman M. V., and L. Kruglyak, 2008 Breeding designs for recombinant inbred
772 advanced intercross lines. Genetics 179: 1069–1078.

Singh R., I. T. Lobina, M. Thomson, S. McCouch, C. Dilla-Ermita, *et al.*, 2013 Multi-
774 parent advanced generation inter-cross (MAGIC) populations in rice: progress and potential for genetics research and breeding. Rice 6: 11.

776 Spencer C. C. A., Z. Su, P. Donnelly, and J. Marchini, 2009 Designing genome-wide

association studies: Sample size, power, imputation, and the choice of genotyping
778   chip. PLoS Genet. 5.

Staab P. R., S. Zhu, D. Metzler, and G. Lunter, 2015 scrm: efficiently simulating long
780   sequences using the approximated coalescent with recombination. Bioinformatics
        31: 1680–1682.

782 Stevenson M., 2018 epiR: Tools for the Analysis of Epidemiological Data

Svenson K. L., D. M. Gatti, W. Valdar, C. E. Welsh, R. Cheng, *et al.*, 2012 High-
784   resolution genetic mapping using the mouse Diversity Outbred population. Genetics
        190: 437–447.

786 Thépot S., G. Restoux, I. Goldringer, F. Hospital, D. Gouache, *et al.*, 2014 Efficiently
        tracking selection in a multiparental population: The case of earliness in wheat.
788   Genetics 199: 609–623.

Turck N., L. Vutskits, P. Sanchez-Pena, X. Robin, A. Hainard, *et al.*, 2011 pROC: an
790   open-source package for R and S+ to analyze and compare ROC curves. BMC
        Bioinformatics 8: 12–77.

792 Valdar W., J. Flint, and R. Mott, 2006 Simulating the collaborative cross: Power of
        quantitative trait loci detection and mapping resolution in large sets of recombinant
794   inbred strains of mice. Genetics 172: 1783–1797.

Wittmann M. J., A. O. Bergland, M. W. Feldman, P. S. Schmidt, and D. A. Petrov, 2017
796   Seasonally fluctuating selection can maintain polymorphism at many loci via
        segregation lift. Proc. Natl. Acad. Sci. 114: E9932–E9941.

798 Wu Y., Z. Zheng, P. M. Visscher, and J. Yang, 2017 Quantifying the mapping precision
        of genome-wide association studies using whole-genome sequencing data.
800   Genome Biol. 18: 1–10.

Xu S., and W. R. Atchley, 1996 Mapping quantitative trait loci for complex binary
802   diseases using line crosses. Genetics 143: 1417–1424.

Yan H., W. Yuan, V. E. Velculescu, B. Vogelstein, and K. W. Kinzler, 2002 Allelic
804   variation in human gene expression. Science 297: 1143.

Yang J., M. N. Weedon, S. Purcell, G. Lettre, K. Estrada, *et al.*, 2011 Genomic inflation
806   factors under polygenic inheritance. Eur. J. Hum. Genet. 19: 807–812.

Yang J., N. A. Zaitlen, M. E. Goddard, P. M. Visscher, and A. L. Price, 2014 Advantages
808   and pitfalls in the application of mixed-model association methods. Nat. Genet. 46:
        100–106.

810 Yu J., G. Pressoir, W. H. Briggs, I. V. Bi, M. Yamasaki, *et al.*, 2006 A unified mixed-
        model method for association mapping that accounts for multiple levels of
812   relatedness. Nat. Genet. 38: 203–208.

Zhang J., K. Kobert, T. Flouri, and A. Stamatakis, 2014 PEAR: A fast and accurate
814   Illumina Paired-End reAd mergeR. Bioinformatics 30: 614–620.

Zheng C., M. P. Boer, and F. A. van Eeuwijk, 2015a Reconstruction of genome ancestry
816   blocks in multiparental populations. Genetics 200: 1073–1087.

Zheng C., M. P. Boer, and F. A. van Eeuwijk, 2015b Reconstruction of genome ancestry
818   blocks in multiparental populations. Genetics 200: 1073–1087.

Zheng C., M. P. Boer, and F. A. van Eeuwijk, 2018 Accurate genotype imputation in
820   multiparental populations from low-coverage sequence. Genetics 210: 71–82.

Figures and Figure Legends

Figure 1. Depictions of various mapping populations explored. Each color represents a
826 unique haplotype.

Figure 2. Genotype estimation accuracy for Hybrid Swarm populations.
Reconstructions were performed for populations simulated using 32 or 128 inbred
founder . Accuracy, calculated as the per-chromosome fraction of variable sites with a
correct diploid genotype estimate, is shown on logit-transformed scale. Values are
coded depending on the number of estimated recombination events, with highly
recombinant estimates (>10 recombination events) displayed as an ✕. Each parameter
combination includes 400 reconstructed autosomes (individual circles) for 100 simulated
individuals. The coalescent-derived individuals displayed here were simulated with an
effective population size of $N_e=1×10^6$ and mutation rate $μ=5×10^{-9}$.

Figure 3. RABBIT chromosome reconstructions for real-life hybrid swarm individuals. A: Depictions of estimated haplotypes for six low-coverage F_5 *Drosophila melanogaster*, identified by sample number. B: Genotype estimate concordance between our reconstruction pipeline and higher-coverage genotype calls. Colors correspond to individuals 1-6 in panel A. Our low-coverage genotype estimates tend to be 97-99% concordant with high-coverage calls, except in cases where RABBIT estimates are highly recombinant. A dashed line at 10 recombination events marks our threshold for excluding assumedly inaccurate hyper-recombinant reconstructions.

Figure 4. Examples of Manhattan plots from our simulated GWAS. These simulations correspond to a single causal locus of large effect, with position indicated by the vertical red line. Recombinant Inbred Lines tend to display a wide association peak near the causal locus, while other populations show only a small number of sites tightly clustered on the causal locus.

Figure 5. Receiver Operator Characteristic (ROC) curves for GWAS simulations. ROC curves represent the aggregate of 500 GWAS simulations (each comprised of 5000 individuals) per parameter combination. To generate single representative ROC curves for each population, we stepped through all specificity values and calculated the mean sensitivity. Inset boxplots display the distribution of AUCs, with whiskers spanning the middle 95% of data. See methods for descriptions for generating each mapping population.

Figure 6. Genomic Inflation Factor (GIF, λ_{1000}) for simulated GWAS with monogenic trait
866   of with a large effect locus, or multi-locus trait of smaller individual effects.
GIF is calculated genome-wide (across all autosomes); on the autosome arm containing
868   the causal allele (linked); and for sites on the autosome physically unlinked to the
causal allele. λ is calculated as the ratio of observed to expected $\chi 2$ values, and a
870   correction is performed to produce the null expectation of a sample size equal to 1000
individuals (see Materials and Methods for details). Data are averaged over 500 GWAS
872   simulations (each comprised of 5000 individuals), with phenotypes assigned in a case-
control framework. Boxes represent the median and interquartile range; whiskers
874   extending to the lower and upper bounds of the 95% quantiles. Hybrid Swarm
populations here used 128 founders. Outbred: F50 population founded by 128 inbred
876   lines.

Population	N Founders	ρ	Mean Δ	σ_Δ
DGRP	128	0.986	-0.015	0.25
DGRP	32	0.502	0.17	2.15
Coalescent	128	0.956	-0.17	0.44
Coalescent	32	0.759	-0.31	1.26

Table 1. Accuracy of estimated number of recombination events following chromosome
880 reconstruction from 0.05X sequencing data. A high concordance correlation coefficient
(Lin's ρ) indicates agreement between estimated and true recombination counts for 400
882 reconstructed chromosomes (coalescent-derived populations) or chromosome arms
(DGRP-derived populations). Coalescent-derived populations are described across a
884 range of values for effective population size N_e and mutation rate μ. Δ represents the
difference between estimated and true recombination counts, and σ_Δ represents the
886 mean of 400 standard deviations of Δ. Reconstructions were performed with a
maximum of 16 most-likely-ancestors with a HARP threshold of 0.99 (see methods for
888 more details). The coalescent-derived populations described here were simulated with
an effective population size of $N_e=1\times10^6$ and mutation rate $\mu=5\times10^{-9}$.
890

892   **Supplemental Figure S1**. Basic structure of the forward simulator pipeline.
        Inbred founding lines (**A**) are randomly intercrossed to produce a recombinant
894   population (**B**). Rapid generation of independent mapping populations is achieved by
        random down-sampling (**C**) and permutation of ancestry (**D**). Population genetic data is
896   encoded in a highly compressed format **(E)** that references the positions of haplotype
        blocks instead of genotypes at every site, enabling us to generate 500 mapping
898   populations for a given parameter combination.

Supplemental Figure S2. Resource usage of RABBIT during haplotype reconstruction. All reconstructions involve the same simulated 2L chromosome arm comprised of four haplotypes. Simulations included varied numbers of founding haplotypes (*N*) and a randomly selected set of markers (number of SNPs, *S*, incremented in steps of 500). All simulations included, at minimum, the four true haplotypes for the simulated individual. In **A** and **B**, points depict the mean of empirical values (over 10 replicates) and gray lines depict the defined regression models. Regression models for memory usage and runtime are extrapolated over a greater range for number of founding haplotypes in **C** and **D**, respectively. Peak memory grew linearly with number of SNPs used, and at a greater-than-linear rate with haplotypes (A). The runtime of RABBIT increased at a greater-than-linear rate for both number of SNPs and number of haplotypes, with the N parameter contributing most to resource cost (B). These models allowed us to estimate resource requirements at greater numbers of haplotypes (C & D) which would be unfeasible to measure empirically.

916   **Supplemental Figure S3.** Example of unambiguous haplotype inference from HARP. For each chromosome, each haplotype is indicated by red and yellow dotted lines.

Supplemental Figure S4. Example of HARP freq output with ambiguous haplotype
920   inference. For each chromosome, each haplotype is indicated by red and yellow dotted
lines. For this individual's autosomes, at least one haplotype cannot be unambiguously
922   resolved.

Supplemental Figure S5. Workflow for reconstructing whole genomes from ultra-low
926   coverage sequencing data. Three required inputs are sequencing reads (.fastq), a
reference genome (.fasta), and homozygous (or phased) founder haplotypes (.vcf file).
928   A recombination map can optionally be provided for use with RABBIT to inform variation
in recombination rates across chromosomes. Flowchart designed in draw.io.
930

Supplemental Figure S6. Liability phenotype model for GWAS simulations. for
934   transforming additive risk scores to probability of assignment to the case population, i.e.
probability of developing a disease. Rare alleles increase risk score, which in turns
936   increases probability of being assigned a case phenotype. **A**: The relationship between
allele frequency and effect on liability risk score. Simulations were modeled as larger or
938   smaller effect with a user-defined effect size coefficient. Error bars describe the
standard deviation of a gaussian sample which constitutes random phenotypic noise. **B**:
940   Logistic transformation of the form $y=(1+e^{-x})^{-1}$ translates the liability score into probability
of being assigned to case population.

Supplemental Figure S7. Optimization curves for Most-Likely-Ancestor (MLA) selection.
To ensure reconstruction was both accurate and computationally feasible, we selected the smallest set of most-likely-ancestors that still represent the greatest fraction of to-be-reconstructed genomes. Increasing the upper limit for the number of MLAs chosen improves the fraction of genome represented with diminishing returns. The HARP threshold of 0.99 performed best for 128-founder populations, for both DGRP-derived and coalescent-derived simulations. Conversely, 32-founder populations performed best with a HARP threshold of 0.95. Data shown reports means across 400 replicates made up of 100 simulated individuals (4 autosomes each for coalescent simulations, 4 autosome arms each for DGRP simulations) per parameter combination. Coalescent-derived populations described here wer1e simulated with $N_e=1\times10^6$ and mutation rate $\mu=5\times10^{-9}$.

Supplemental Figure S8. Optimization curves for Most-Likely-Ancestor inclusion for
960 additional parameter combinations. A: SCRM (coalescent) derived populations across
different parameters; B: accuracy is similar for both 0.05X and 0.005X sequencing
962 coverage in populations simulated with DGRP genomes.

Supplemental Figure S9. Accuracy of genome reconstruction for simulated, DGRP-
966   derived F_5 hybrid swarm individuals. Reconstructions were performed for populations
simulated as being founded by either 32 or 128 inbred lines for two levels of ultra-low
968   sequencing coverage. Accuracy is represented on a logit scale, as most points occur
above 90%. Reconstructed chromosomes estimated to have $\geq$10 recombination events
970   are denoted by an ✕, offset from the bulk of the distribution. Each parameter
combination includes 400 reconstructed chromosomes (from 100 simulated individuals).

974 **Supplemental Figure S10.** Accuracy of genome reconstruction for simulated, coalescent-derived F$_5$ hybrid swarm individuals. Reconstructions were performed for
976 populations simulated as being founded by either 32 or 128 inbred lines for various effective population sizes (N$_e$) and mutation rates (μ). Accuracy is represented on a logit
978 scale, as most points occur above 90%. Reconstructed chromosomes estimated to have ≥10 recombination events are denoted by an ✕, offset from the bulk of the
980 distributions. Each parameter combination includes 400 reconstructed chromosomes (from 100 simulated individuals).
982

984   **Supplemental Figure S11**. Genotyping accuracy of STITCH for simulated 32-founder
       F5 hybrid swarm populations sequenced at 0.05X coverage. Genotype accuracy
986   improves with greater numbers of sequenced individuals, as STITCH infers missing
       genotypes from other haplotypes in the population. Boxes represent the median and
988   interquartile range; whiskers extending to the lower and upper bounds of the 95%
       quantiles. See methods for parameters used.

Supplemental Figure S12. Receiver Operator Characteristic (ROC) curves for additional GWAS simulations. To generate single representative ROC curves across 500 GWAS simulations per population, we stepped through all specificity values and calculated the mean sensitivity. Inset boxplots display the median and interquartile range of Area Under the Curve (AUC) distributions, with whiskers spanning the middle 95% of data. **ILs**: inbred lines. **HS**: Hybrid Swarm populations founded by 32 or 128 lines.

1002   **Supplemental Figure S13.** Genomic Inflation Factor with and without low-frequency
filtering. Excluding alleles with a minor allele frequency below 5% does not reduce
1004   genomic inflation factor in inbred lines. Box depicts the middle 50% quantiles with
whiskers extending to the middle 95% of data points. Traits were modeled as a single
1006   locus of large effect.

1008

Population	Coverage	N Founders	N_e	μ	ρ	Mean Δ	σ_Δ
DGRP	0.005X	128	-	-	0.201	3.06	3.11
DGRP	0.005X	32	-	-	0.201	3.15	2.81
DGRP	0.05X	128	-	-	0.986	-0.015	0.25
DGRP	0.05X	32	-	-	0.502	0.17	2.15
Coalescent	0.05X	128	10^4	1×10^{-9}	0.029	-2.72	1.60
Coalescent	0.05X	32	10^4	1×10^{-9}	0.219	-1.84	1.50
Coalescent	0.05X	128	10^5	1×10^{-9}	0.288	-1.57	1.64
Coalescent	0.05X	32	10^5	1×10^{-9}	0.761	-0.53	0.99
Coalescent	0.05X	128	10^6	1×10^{-9}	0.742	-0.46	1.09
Coalescent	0.05X	32	10^6	1×10^{-9}	0.754	-0.42	1.22
Coalescent	0.05X	128	10^4	5×10^{-9}	0.203	-2.03	1.73
Coalescent	0.05X	32	10^4	5×10^{-9}	0.622	-0.89	1.17
Coalescent	0.05X	128	10^5	5×10^{-9}	0.652	-0.64	1.41
Coalescent	0.05X	32	10^5	5×10^{-9}	0.782	-0.26	1.16
Coalescent	0.05X	128	10^6	5×10^{-9}	0.956	-0.17	0.44
Coalescent	0.05X	32	10^6	5×10^{-9}	0.759	-0.31	1.26
Coalescent	0.05X	128	10^4	1×10^{-8}	0.238	-1.65	1.86
Coalescent	0.05X	32	10^4	1×10^{-8}	0.745	-0.66	1.05
Coalescent	0.05X	128	10^5	1×10^{-8}	0.776	-0.34	1.12
Coalescent	0.05X	32	10^5	1×10^{-8}	0.846	-0.25	0.93
Coalescent	0.05X	128	10^6	1×10^{-8}	0.937	-0.22	0.56
Coalescent	0.05X	32	10^6	1×10^{-8}	0.833	-0.32	0.95

Supplemental Table S1. Accuracy of estimated number of recombination events
1010 following chromosome reconstruction. A high concordance correlation coefficient (Lin's
ρ) indicates agreement between estimated and true recombination counts for 400
1012 reconstructed chromosomes (coalescent-derived populations) or chromosome arms
(DGRP-derived populations). Coalescent-derived populations are described across a
1014 range of values for effective population size N_e and mutation rate μ. Δ represents the
difference between estimated and true recombination counts, and σ_Δ represents the
1016 mean of 400 standard deviations of Δ. Reconstructions were performed with a
maximum of 16 most-likely-ancestors with a HARP threshold of 0.99 (see methods for
1018 more details).

1020   **Supplemental Table S2.** Test statistics and p-values for 270 pairwise comparisons of
        simulated GWAS AUC distributions. Our p-value significance threshold at α=0.05 is
1022 1.85×10^{-4}.

1024 **Supplemental Methods**

1026 **Generating simulated reference panels across a range of diversity levels**
To evaluate low-coverage reconstruction for various degrees of genetic diversity, we
1028 generated reference panels using haplotypes produced by coalescent models across a
range of genetic diversity levels. Haplotypes were generated using the R (R Core Team
1030 2016) package scrm (Staab *et al.* 2015) and subsequently restructured into VCF file
format (Danecek *et al.* 2011). We generated ten independent panels for each of all 18
1032 combinations of population size (Ne=104, 105, 106), mutation rate (µ=10-9, 5×10-9, 10-
8), and number of haplotypes (32, 128). The value θ for each simulation was defined as
1034 4Neµ. We simulated a chromosome-length locus of 25 Mb with a recombination rate of
1.5 cM/Mb. SNP positions output by scrm (a decimal within the range of 0 to 1) were
1036 converted to base pair positions by multiplying the decimal by chromosome length (25
×106 base pairs for our simulations) and rounding down to the nearest integer. Any
1038 sites with more than two alleles were converted to a biallelic site by discarding tertiary or
quaternary alleles. Genotype values were re-coded as polarized signed integers: +1 for
1040 reference and -1 for alternate alleles. For every position, reference and alternate alleles
were defined by randomly selecting one of the twelve non-repeating pairs of
1042 nucleotides. Reference genome FASTA files were created with a custom python script
that generated a 25 million length string of nucleotide characters with weighted
1044 probability to achieve 45% GC-content, followed by replacing variable positions with
their respective reference alleles.
1046

Extended details for simulating GWAS
1048 *Simulated haplotypes.* Although the forward simulator we developed is efficient, it would
not have been computationally feasible to simulate 500 fully independent mapping
1050 populations (per parameter combination) in a reasonable amount of time. Instead, we
generated ten independent forward-simulated populations, and for each of those,
1052 generated fifty randomly permuted subsets. For a single simulated mapping population,
we began by sampling (with replacement) a random subset of 5,000 individuals, out of
1054 10,000 total individuals generated by forward-simulation. Then, we performed a
permutation of haplotype ancestry with a new, randomly-ordered (equally sized) subset
1056 of founders. The permutation of ancestry was one-to-one, e.g. all haplotype blocks that
were previously derived from founder X would be translated to founder Y, and blocks
1058 previously derived from Y would in turn be mapped to founder Z.

1060 For all simulated GWAS, we began with a set of 129 DGRP haplotypes with the least
missing data, e.g. high coverage and low levels of heterozygosity. This allowed us to
1062 perform leave-one-out subsampling for the 128-founder populations. In addition to
Hybrid Swarm populations, which we ran through the simulated sequencing and
1064 mapping pipeline, we generated four additional types of mapping populations for
comparing GWAS performance: Highly outbred (F50) populations; Inbred Lines (ILs) to
1066 represent the DGRP; and Recombinant Inbred Lines (RILs), similar to the Drosophila
Synthetic Population Resource, or DSPR (King *et al.* 2012).
1068

The F50 populations were generated with 128 founders in same manner as the Hybrid
1070   Swarm, except that populations were simulated over fifty non-overlapping generations
of recombination instead of five generations. The ten resulting forward-simulated
1072   populations were resampled and permuted as we did with the Hybrid Swarms.

1074   We simulated ten initial sets of 800 RILs using the same forward-simulator as previously
described, each initialized with a random subset of eight DGRP haplotypes. Populations
1076   randomly recombined at a population size of 10,000 for fifty non-overlapping
generations, after which 800 random male-female pairs of individuals were isogenzied
1078   through 25 generations of full-sibling mating. This scenario roughly corresponds to the
DSPR. For computational simplicity, after the 25 generations of isogenization we
1080   removed any remaining residual heterozygosity by forcing the identity of a second
chromosome copy to be identical to the first copy. We then sampled 5,000 draws (with
1082   replacement) of the 800 RILs followed by ancestry permutation as described above.

1084   To simulate GWAS on Inbred Lines, no forward-simulation was necessary. For a single
simulated population, we first randomly selected 128 DGRP lines, then randomly
1086   sample with replacement 5,000 times. As with hybrid swarm and RILs, for any
parameter combination we generated a total of 500 mapping populations.
1088
Liability model. Using the liability framework, we assign case or control phenotypes to
1090   mapping populations derived from DGRP chromosome 2L haplotypes. First, a
genotype-dependent risk score is calculated for every individual, which is translated into
1092   probabilities for case or control group assignment. We parameterized our model such
that individuals begin with equivalent odds of being assigned to case or control groups,
1094   and that probability is modified depending on allele status at causal loci.

1096   For a given simulation, N causal loci are randomly selected with equal probability out of
all segregating sites in the mapping population. The effect of individual causal loci are
1098   simulated as a Gaussian variable drawn with a small extent of noise (sd=0.005) with an
expected (mean) value dependent on the allele frequency at that locus. Intermediate
1100   frequency loci (i.e. allele frequency = 0.5) are simulated to contribute on average no
effect on case/control assignment; alleles near fixation decrease risk; rare alleles
1102   increase risk. The linear center of the curve, where most individual risk scores exist
under the parameter combinations used in our simulations, approximates additive
1104   genetic architecture. The sigmoid tails allow for the continuous risk score to translate to
a phenotype necessarily bounded by zero and one. The relationship between allele
1106   frequency, risk score, and phenotypic assignment is diagrammed in Supplemental
Figure S9. Within the linear portion of the curve (which is the domain for the majority of
1108   simulations), a singleton minor allele will modify 'case' assignment probability by
approximately +5% (small effect) or +10% (large effect), beginning at an initial
1110   assignment probability of 50%. The major allele, conversely, would decrease
assignment by an opposite amount. An allele at 50% frequency will not modify
1112   case/control assignment (aside from a very small degree of random noise equal to
about +/- 0.06%). Note that we performed GWAS simulations assuming 100% genotype
accuracy.

We implemented a method of efficiently aggregating allele counts compatible with our haplotype map format. Briefly, haplotype map breakpoints across all individuals are
sorted in ascending order. When iterating through ascending unique start and stop positions, between any pair of breakpoints, all SNPs will be comprised of the same
number of each founding haplotype. Haplotype IDs could then be counted and sorted in the same column position order as the table containing polarized allele status (-1 for
alternate, +1 for reference). Multiplying the genotype table by the haplotype count vector results in final allele counts, polarized negative for alternate alleles and positive for
reference alleles.