NITROGEN FERTILISERS SHAPE THE COMPOSITION AND PREDICTED FUNCTIONS

OF THE MICROBIOTA OF FIELD-GROWN TOMATO PLANTS

Federica Caradonia[1,3*], Domenico Ronga[1,#], Marcello Catellani[1,‡], Cleber Vinícius Giaretta

Azevedo[1,2], Rodrigo Alegria Terrazas[3], Senga Robertson-Albertyn[3], Enrico Francia[1*] and

Davide Bulgarelli [3*]

[1]Department of Life Sciences, Centre BIOGEST-SITEIA, University of Modena and Reggio

Emilia, Italy; [2]Faculty of Agricultural and Veterinary Sciences, Sào Paulo State University,

Jaboticabal, Brasil; [3]Plant Sciences, School of Life Sciences, University of Dundee,

Dundee, United Kingdom.

Present addresses:

[#]CREA, Centre for Animal Production and Aquaculture (CREA-ZA), Lodi, Italy;

[‡]ENEA, Department for Sustainability, C.R. Trisaia, Rotondella (MT), Italy.

* Correspondence:

Federica Caradonia

federica.caradonia@unimore.it

Enrico Francia

enrico.francia@unimore.it

Davide Bulgarelli

d.bulgarelli@dundee.ac.uk

ABSTRACT

The microbial communities thriving at the root-soil interface have the potential to improve plant growth and sustainable crop production. Yet, how agricultural practices, such as the application of either mineral or organic nitrogen fertilisers, impact on the composition and functions of these communities remains to be fully elucidated. By deploying a two-pronged 16S rRNA gene sequencing and predictive metagenomics approach we demonstrated that the bacterial microbiota of field-grown tomato (*Solanum lycopersicum*) plants is the product of a selective process that progressively differentiates between rhizosphere and root microhabitats. This process initiates as early as plants are in a nursery stage and it is then more marked at late developmental stages, in particular at harvest. This selection acts on both the bacterial relative abundances and phylogenetic assignments, with a bias for the enrichment of members of the phylum Actinobacteria in the root compartment. Digestate-based and mineral-based nitrogen fertilisers trigger a distinct bacterial enrichment in both rhizosphere and root microhabitats. This compositional diversification mirrors a functional diversification of the root-inhabiting communities, manifested predominantly by the differential enrichment of genes associated to ABC transporters and the two-component system. Together, our data suggest the microbiota thriving at the tomato root-soil interface is modulated by and in responses to the type of nitrogen fertiliser applied to the field.

Additional keywords: *Solanum lycopersicum*, Rhizosphere, Root, Microbiota, Nitrogen, Fertilisers, Digestate.

INTRODUCTION

Limiting the negative impact of agricultural practices on the environment while preserving sustainable crop yield is one of the key challenges facing agriculture in the years to come.

As an essential element for plant nutrition, nitrogen represents a paradigmatic example of such a challenge. Moreover, due to the combined effect of elevated solubility and little retention in soils, the lack of this element is and will be one of the major yield-limiting factors worldwide (Tilman et al., 2011). At the same time, the application of synthetic nitrogen fertilisers is, in many agricultural systems, a low efficiency approach which has been linked with the degradation of natural resources (Elser and Bennett, 2011). One of the strategies adopted to limit the economic and environmental footprint of crop production while maintaining sustainable yield is the "recycling" of, mineral-rich, biodegradable products of the livestock and agricultural sectors.

One example of this approach is the application of the digestate, a by-product the anaerobic digestion of organic waste for the production of biogas (Möller and Müller, 2012) as renewable soil amendment for crop production. The digestate is a mixture of partially-degraded organic matter, microbial biomass and inorganic compounds (Alburquerque et al., 2012). We recently demonstrated how the digestate can be efficiently used as innovative fertiliser and plant growing media (Ronga et al., 2018b; Ronga et al., 2018a; Ronga et al., 2019), yet the impact of digestate applications on the agroecosystem remains to be fully elucidated.

For instance, the digestate can be a source of phytoavailable nitrogen, in particular ammonium, capable of impacting on organic matter mineralisation and emission of carbon dioxide from the soil profile (Grigatti et al., 2011). Therefore, it is legitimate to hypothesize that such treatments impact on the composition of the microbial communities thriving at the root-soil interface, collectively referred to as the rhizosphere and root microbiota, which

play a critical role in mobilisation of mineral elements for plant uptake (Alegria Terrazas et al., 2016). Congruently, several studies indicate that the application of biogas by-product enhances soil microbial activity (Möller, 2015) . However, the intertwined relationship among microbiota composition, soil characteristics and amendments as well as host plant species-specificity (Bulgarelli et al., 2013) makes it difficult to infer first principles.

This is particularly true for field-grown crops such as tomato (*Solanum lycopersicum L.*), one of the most cultivated horticultural crops globally with plantations occupying an area of 4.8 million of hectares with a production of 182 million tonnes in 2017 (http://www.fao.org/faostat/en/). Notably, this species is also an excellent experimental model for basic science investigations: tomato was one of the first crops whose genome was sequenced (Consortium, 2012) and provided a superb platform to test the significance of genome editing for evolutionary studies and plant breeding (Zsögön et al., 2018). Perhaps not surprisingly, tomato is gaining momentum as an experimental system to study host-microbiota interactions in crop plants. Recent investigations revealed novel insights into the assembly cues of the microbiota associated to this plant (Bergna et al., 2018; Toju et al., 2019) and the contribution of microbes thriving at the tomato root-soil interface to pathogen protection (Chialva et al., 2018; Kwak et al., 2018). However, the composition and functional potential of the tomato microbiota and their interdependency from nitrogen fertilisers remain to be elucidated.

Here we report the metagenomics characterisation of the microbiota thriving at the root-soil interface of field-grown tomato plants. We hypothesize that nitrogen treatments shape and modulate the contribution of the tomato microbiota for crop yield. To test this hypothesis, we focused on processing tomato exposed to different nitrogen fertilisers, either digestate-based or containing a mineral fraction. By using a 16S rRNA amplicon sequencing survey we deciphered how the microhabitat (i.e., either rhizosphere or root)

94   sculpts the tomato microbiota which, in turn, is fine-tuned by the type of fertiliser applied.

95   Finally, by using a predictive metagenomics survey, we inferred the functional

96   diversification imposed by the nature of the fertilisers on the root microbiota.

MATERIALS AND METHODS

Field site

A field trial was established in a tomato farm near the city of Ravenna (44°25'40.8"N 12°05'53.3"E), Emilia Romagna Region, Italy, during the 2017 growing season. During the period from transplant to harvest, the minimum and the maximum average temperatures recorded were 17.1°C and 32.8°C, respectively, and the rainfall was 101.7 mm. The soil had a silty loam texture (14% clay, 51% silt, 35% sand), a pH 8.3 (in H_2O), 1.1 g kg^{-1} total N (Kjeldahl method), 7 mg kg^{-1} available P (Olsen method), 129 mg kg^{-1} exchangeable K (Ammonium acetate), and 9 g kg^{-1} organic matter (Walkey-Black method). A schematic illustration of the field trial is depicted in Figure 1.

Plant material

We used the tomato cultivar 'Fokker', a processing-type genotype with blocky fruit, late fruit ripening and suitable for tomato puree, for the experimentation. Seedlings were provided by Bronte Soc. Coop. Agr. A.R.L. (Mira, Italy). Processing tomato seedlings were transplanted at the end of May when they were 6-week old corresponding to plants at the fourth true leaf stage. Plant density was 3 plants m^{-2}. Plants were transplanted into single row, with a spacing of 0.22 m between plants in each row and 1.50 m between rows.

Experimental design

We established a randomized complete design with three replicates and seven treatments: pelleted digestate (hereafter PE), liquid digestate (LD), slow-acting liquid digestate (SRLD), organo-mineral fertiliser based on digestate (SC), synthetic fertiliser (MF), slow-acting synthetic fertiliser (SRMF), and no fertilization treatment (NT). The composition of the treatments is summarised in Table 1. For each treatment, we applied a total amount of nitrogen in the ratio 150 N kg ha^{-1} on the basis of soil analysis, crop rotation and crop nutrients required. Nitrogen was supplied at transplanting time with the exception of the synthetic fertiliser treatment. For this latter treatment, the amount of total Nitrogen was

125 equally divided and applied in 3 times (transplanting, full flowering and fruit ripening) using

126 ammonium nitrate in the first treatment and calcium nitrate in the second and in the third

127 ones. During the trial, 600 m^3 ha^{-1} of irrigation water was distributed by drip irrigation to

128 each treatment. Weeds and pests were controlled according to the production rules of

129 Emilia Romagna Region, Italy.

Yield traits

131 At harvest we determined the marketable yield (t ha^{-1}), as a weight of fully ripe fruits, and

132 the solid soluble content (Brix t ha^{-1}) as a proxy for fruit quality.

Root, Rhizosphere and Bulk soil Sampling and DNA Extraction

At transplanting time (May 2017), 5 root specimens per treatment were collected. Upon

uprooting, soil particles loosely bound to roots were dislodged by hand shaking and root

segments of ~ 6 cm were placed in sterile 50 mL tubes. The samples were stored in a

portable cooler (~ 4°C), transported to the laboratory and immediately processed. Root

specimens were incubated in 30 mL of PBS (Phosphate buffered saline) and placed on a

shaker for 20 minutes in order to separate the soil tightly adhering to plant material, which

we operationally defined as "rhizosphere", from the roots. The first tubes were centrifuged

for 20 minutes at 4,000 x g and the rhizosphere pelleted was collected in liquid nitrogen

and stored at -80°C. The roots were moved to a new sterile tube containing 30 mL PBS

and sonicated by Ultrasonics Sonomatic Cleaner (Langford Ultrasonics, Birmingham, UK)

for 10 minutes (intervals of 30 seconds pulse and 30 seconds pause) at 150 W, as

previously reported (Schlaeppi et al., 2014) to enrich for endophytic microorganisms.

Roots were then washed in the same new buffer and dried on sterile filter paper. After few

minutes, the roots were moved to 50 mL tubes and frozen in liquid nitrogen for storage at -

80°C. Three independent soil samples were harvested from unplanted soil in different

points of the field, frozen in liquid nitrogen and stored at -80°C. At harvest time (September

2017) the whole plants were harvested, 5 roots per treatment and 3 bulk soil samples were

collected, prepared and stored like the previous samples. Frozen root samples were pulverized in a sterile mortar using liquid nitrogen prior DNA preparation. DNA was extracted from all the specimens (i.e., bulk soil, rhizosphere and pulverized roots) using the FastDNA® SPIN Kit for Soil (MP Biomedicals, Solon, USA) following the instruction manual provided by manufacturer. DNA samples were diluted using 50 µL DES water and quantified using the Nanodrop 1000 Spectrophotometer (Thermo Scientific, Wilmington, United States).

16S rRNA Gene Sequencing

The sequencing library was generated using primers specific (515F 5'-GTGCCAGCMGCCGCGGTAA-3' and 806R 5'-GGACTACHVGGGTWTCTAAT-3') for hypervariable V4 region of the 16S rRNA gene. The reverse primers included a 12-mer unique "barcode" sequences (Caporaso et al., 2012) to facilitate the multiplexing of the samples into a unique sequencing run. Individual PCR reactions were performed as previously reported (Robertson-Albertyn et al., 2017), with the exception of the concentration of the Bovine Serum Albumin, added at 10 µg/reaction, and the addition of a Peptide Nucleic Acid (PNA) blocker (PNA Bio, Newbury Park, United States) at a concentration of 0.5 µM/reaction to inhibit plastidial amplification. For each barcoded primers, three technical replicates and a no-template control (NTC) were organised and processed starting from a unique master mix. Five microliters of amplified samples and cognate NTCs were inspected on a 1% (w/v) agarose gel. Two independent sets of triplicated amplicons, displaying the expected amplicon size and lacking detectable contaminations, were combined in a barcode-wise manner and purified using the Agencourt AMPure XP kit (Beckman Coulter, Brea, United States) with a ratio of 0.7 mL AMPure XP beads per 1 mL of sample. Purified DNA samples were quantified using Picogreen (Thermo Fisher, United Kingdom) and combined in an equimolar ratio into an amplicon pool. This latter material was used for the preparation of a MiSeq run at the

Genome Technology facilities of the James Hutton Institute (Invergowrie, UK) as previously reported (Robertson-Albertyn et al., 2017).

OTU Table Generation and pre-processing

We used QIIME, version 1.9.0 (Caporaso et al., 2010) to process the sequencing output of the MiSeq machine. Briefly, the command join_paired_ends.py was used to decompress and merged (minimum overlap 5bp) forward and reverse read FASTQ files. Next, we removed in silico low-quality sequencing reads and sequencing reads without the barcode information. Then, the reads were assigned to individual samples. In these analyses, the command split_libraries_fastq.py was used imposing a minimum PHRED score of 20. The resulted high-quality reads were assembled into an Operational Taxonomic Unit (OTU) table at 97% sequence identity. We used a 'closed reference' approach for OTU-picking using the command pick_closed_reference_otus.py. We imposed the Green Genes database version 13_5 (DeSantis et al., 2006) as a references database to identify microbial OTUs and prune for chimeric sequences. We used SortMeRNA algorithm for OTU -picking and taxonomy assignment. Finally, OTUs whose representative sequences were classified as either chloroplast or mitochondria, as well as OTUs accruing only one sequencing read over the entire dataset (i.e., singletons), were depleted in silico using the function filter_otus_from_otu_table.py.

Data visualisation and statistical analyses

Agronomic traits were analysed by Analysis of variance (ANOVA) using GenStat 17th (VSN International, Hemel Hempstead, UK). Means were compared using Bonferroni's test at the 5% level.

The OTU table produced in QIIME was analysed in R using a custom script developed from Phyloseq (McMurdie and Holmes, 2013).

Initially, the data were filtered removing the samples with less than 1,000 reads and the OTUs with less than 10 reads in at least the 5% of the samples. For alpha-diversity calculation, sequencing reads were rarefied at an even sequencing deep of 18,467 reads per sample retaining 2,439 unique OTUs. The number of observed OTUs and Chao1 index was used as richness estimators, while the Shannon index was used for evaluating the evenness. Upon inspecting distribution of the data using a Shapiro-Wilk test, the means of rhizosphere and root samples at harvest time were compared using a non-parametric Wilcoxon rank sum test.

For beta-diversity calculation, the original counts (i.e., not rarefied) were transformed to relative abundances and we imposed an abundance threshold to target PCR-reproducible OTUs. The differences among microbial communities of the samples were computed using Bray Curtis index and weighted Unifrac index, with this latter index including phylogenetic information in the analysis (Lozupone and Knight, 2005). A principal coordinates analysis was generated to visualize similarities and dissimilarities of microhabitats and treatments. In order to assess the effects of microhabitats and the treatments on the bacterial community composition a Permutational Multivariate Analysis Of Variance (PERMANOVA) was performed using the distance matrices. Before performing the analysis, bulk and nursery samples were removed from dataset.

Finally, original counts data were used to perform a differential analysis to identify individual bacteria differentially enriched in the tested samples using DESeq2 (Love et al., 2014).

The phylogenetic tree was constructed using the representative sequences of the OTUs significantly enriched in rhizosphere and root specimens and annotated with iTOL (Letunic and Bork, 2006).

Functional predictions

Tax4Fun (Asshauer et al., 2015) package in R was used as a predictive tool to obtain a

functional profile based on 16S rRNA gene data. Metabolic capabilities are calculated by

linking the amplicon data phylogenetic and abundance profile to a set of pre-computed

metabolic reference profiles, based on the KEGG Ortholog (KO) database (Kanehisa et

al., 2008). The input for this analysis was an OTU table obtained with the representative

sequences of the OTU table previously generated (see above), reclassified using

SILVA_115 taxonomic database (Quast et al., 2013). Similar to a previously reported

operational protocol (Kavamura et al., 2018), we focused our analysis in prokaryotic

functional categories related to amino acid metabolism, carbohydrate metabolism, cell

motility, energy metabolism, membrane transport, metabolism of terpenoids and

polyketides and two-component system, trimming the rest of predicted functions from the

Tax4fun output. A statistical comparison between two groups using a Welch's t-test

(Bluman, 2009) filtered at a p-value < 0.01 with Storey's correction for false discovery rate

(Storey and Tibshirani, 2003) was performed in STAMP, Statistical Analysis of

Metagenomic Profiles (Parks et al., 2014).

Data and scripts availability

The 16S rRNA gene sequences presented in this study are available at the European

Nucleotide archive under the study accession number PRJEB32219. The scripts to

reproduce the statistical analysis and figures are available at

https://github.com/BulgarelliD-Lab/Tomato_nitrogen. Data frames required for scripts

reproducibility are included in Supplementary Database 1.

RESULTS

Fertiliser treatment impacts on yield and quality of processing tomato

At harvest time the two most important parameters such as marketable yield and fruit

quality were collected to evaluate the effect of 7 different fertiliser performances on

processing tomato (Table 2). The fertiliser treatments had a significant effect on fruit fresh biomass (ANOVA, Bonferroni's test, $P < 0.001$). Pelleted digestate registered the best performance followed by synthetic fertiliser and slow acting liquid digestate. In addition, the different fertilisers influenced significantly also the quality of processing tomato (Table 2) (ANOVA, Bonferroni's test, $P < 0.001$).

The assembly dynamics of the bacterial microbiota of field-grown processing tomato

To gain insights into the relationships between yield traits and microbiota composition in field-grown processing tomato plants, we generated 5,546,303 high quality 16S rRNA gene sequences for 86 samples generated in this study.

Upon in silico depletion of OTUs classified as Mitochondria we reduced the number of analysable sequences to 4,645,503 with a retaining proportion of 83.7% of the original sequences (mean per samples = 54,017.48 reads; max = 111,213 reads; min = 272 reads). The data were further filtered removing the samples with less than 1,000 reads as well as the OTUs with less than 10 reads in 5% of samples. This allowed us to retain 2,515 unique OTUs accounting for 4,308,580 high quality reads and 85 samples.

Then, we computed alpha-diversity calculations on a dataset rarefied at 18,467 reads per sample and alpha-diversity was investigated considering two microhabitats (root and rhizosphere) and the seven fertilisers treatments. OTUs richness was assessed by Chao1 index and observed OTUs while the OTUs evenness was assessed by Shannon index. This analysis revealed a significant effect of the microhabitat on the characteristics of the microbiota thriving at the tomato root-soil interface: regardless of the treatment, the root microhabitat emerged as less diverse and even compared to the rhizosphere one (Wilcoxon rank sum, $p < 0.01$, Figure 2). This observation suggests that root microhabitat represents a gated community compared to the surrounding soil environment.

Congruently, beta-diversity analysis computed on the non-rarefied dataset using both weighted Unifrac and Bray-Curtis indicated a microhabitat-dependent microbiota diversification. In particular, the weighted Unifrac matrix visualised using a Principal Coordinated Analysis revealed such a microhabitat effect on samples processed at harvest time along the axis accounting for the major variation. Interestingly, younger nursery samples displayed a similar degree of diversification, although their communities were separated from the harvest samples on the axis accounting for the second source of variation (Figure 3). These data were supported by a permutational analysis of variance which attributed a R^2 of 37% to the treatment, a R^2 of 30% to the microhabitat and a R^2 of 5% to their interactions (Adonis test, 5,000 permutations, p <0.01; "nursery" samples considered as a treatment). Remarkably, the Bray-Curtis matrix produced a congruent results, although the temporal effect (i.e., nursery vs. harvest time) explained slightly more variation (~ 29%; Supplementary Figure S1) than microhabitat diversification manifested along the second axis of variation (~ 26%; Supplementary Figure S1). Crucially, also in this case the observed diversification was supported by a permutational analysis of variance which attributed a R^2 of 39% to the treatment, a R^2 of 23% to the microhabitat and a R^2 of 7% to their interactions (Adonis test, 5,000 permutations, p <0.01; "nursery" samples considered as a treatment).

Differential bacterial enrichments define microhabitat and treatment "signatures" on the field grown tomato microbiota

To gain insights into individual members of the tomato microbiota responsible for the observed diversification we implemented a series of pair-wise comparison among microhabitats and treatments at harvest stage. We took a two-pronged approach. First, we identified bacterial underpinning the microhabitat effect i.e., the selective enrichment of bacteria in the roots and the rhizosphere microhabitats amended with no fertiliser. Next,

we assessed the effect of the fertiliser treatment on roots and rhizosphere bacterial composition by comparison with bacteria enriched in untreated samples.

This allowed us to identify 170 bacterial OTUs whose abundance was significantly enriched in and differentiated between rhizosphere specimens and unplanted soil samples (Wald test, $p < 0.01$, FDR corrected; Supplementary database 1). Similarly, we identified 374 bacterial OTUs whose abundance was significantly enriched in and differentiated between root specimens and unplanted soil samples (Wald test, $p < 0.01$, FDR corrected; Supplementary database 1). Of these differentially enriched bacteria, 96 OTUs represented a set of tomato-competent OTUs capable of colonising both the rhizosphere and root environments. When we then looked into the taxonomic affiliations of this tomato-competent microbiota, we discovered that the field-grown tomato microbiota is dominated by members of Actinobacteria, Bacteroidetes, Alpha-, Beta-, Gamma- and Deltaproteobacteria as well as Verrucomicrobia (Figure 4). Strikingly, the taxonomic investigation revealed a bias for Actinobacteria in the root compartment, possibly reflecting an adaptive advantage of member of this phylum in colonising the endophytic environment.

Interestingly, each fertiliser treatment had a distinct impact on these tomato-enriched microbiota. The pelleted digestate (PE) and the slow-acting synthetic fertiliser (SRMF) yielded the highest number of uniquely enriched OTUs regardless of the microhabitat investigated, albeit with a distinct pattern: the SRMF had a more pronounced effect on the rhizosphere communities while the PE impacted more on the bacteria thriving in association with root tissues. (Wald test, $p < 0.01$, FDR corrected; Figure 5; Supplementary database 1).

Interestingly, when we inspected the taxonomic composition of the bacteria differentially impacted by the fertiliser treatment we observed an increase of the number of OTUs

belonging to phylum of Actinobacteria. In particular, PE had 12 OTUs out of 80 and 14

OTUs out of 105, in root and rhizosphere, respectively, belonging to phylum

Actinobacteria. While, MF had 15 OTUs out of 38 and 22 OTUs out of 49 in root and

rhizosphere, respectively, belonging to phylum Actinobacteria (Supplementary database

S1). Within this phylum we observed the presence of OTUs classified as *Streptomyces*

spp., *Agromyces sp.*, *Microbispora sp.* and *Actinoplanes spp.*

Together these data suggested that the enrichment of specific bacteria underpins the

observed microhabitat effect whose magnitude is fine-tuned by the applied fertiliser.

Organic- and synthetic-based fertiliser trigger different metabolic capacities in the tomato
root microbiota

To investigate the ecological significance of the observed differential recruitments among

fertiliser treatment we employed a predictive metagenomics approach. Briefly, we inferred

in silico the functions encoded by the tomato microbiota at harvest stage (Material and

Methods) and we grouped the samples in digestate-based (i.e., PE, LD and SRLD;

hereafter 'organic') and treatments containing at least a synthetic component (i.e., SC, MF

and SRMF; hereafter 'mineral'). We observed that the functions putatively encoded by the

communities exposed to either organic and mineral fertilisers can discriminated between

treatments in both microhabitat (PERMANOVA: Rhizosphere samples R^2 = 14%, p value

<0.01, 5,000 permutations; Root samples R^2 = 16%, p value <0.01, 5,000 permutations)

Congruently, we identified a set of 14 functions differentially enriched between root

communities exposed to either group of treatments (Welch t-test, p <0.01, FDR corrected;

Figure 6). Interestingly, we observed a striking dichotomy between the two groups of

treatments: communities exposed to mineral fertilisers are predicted to enrich for genes

implicated in the ABC transporter machinery while bacteria exposed to the organic

treatments are predicted to enrich for genes implicated in the two-component system.

These two set of genes are dominant in communities exposed to both treatments and are

also associated to additional distinct enrichment patterns, most notably including nitrogen metabolism (organic communities) and tetracycline biosynthesis (mineral communities).

These results suggest that, within tomato roots, the observed taxonomic diversification underpins a functional specialisation of the microbiota which, in turn, may impact on plant growth development and health.

DISCUSSION

This study revealed that all nitrogen treatments led to an increase of tomato production in comparison with the no fertilization treatment (~ + 142.5 ± 30.5%) confirming that, in the tested conditions, nitrogen limits the yield potential of processing tomato crops as observed in previous studies (Ronga et al., 2015; Ronga et al., 2017). Yet, despite the same amount of nitrogen was applied in each treatment (i.e., 150 kg ha^{-1}), all the treatments were statistically different from each other. A prediction of this observation is that, under the tested conditions, the nature of the fertilisers, rather than the amount of nitrogen per se, affect the yield and the fruit quality of tomato plants. These observations and the contribution to plant nutrition of the microbial communities thriving at the root-soil interface (Alegria Terrazas et al., 2016), motivated us to investigate relationships between yield traits and the composition of the tomato rhizosphere and root microbiota under field conditions.

The tomato rhizosphere and root microbiota are gated communities

First, we characterised the rhizosphere and root microbiota of processing tomato with no treatment. Both alpha and beta diversity discriminated between the communities of seedlings and adult plants (Figures 2 and 3). Despite these differences, which could be attributed to both abiotic, e.g., time of residence in soil (Dombrowski et al., 2017), and biotic factors, e.g., developmental-conditioned rhizodeposits (Chaparro et al., 2014), it is striking to note how tomato plants displayed a rhizosphere and root compartmentalisation

regardless of the developmental stage. This is congruent with the observation that in rice,

the assembly and structural diversification of the microbiota is a rapid process which

reaches a steady-state level within a few weeks from germination (Edwards et al., 2015).

Closer inspection of the rhizosphere and root profiles at harvest stage indicates that these

plant-associated communities are phylogenetically related to those of unplanted soil,

suggesting that the initial tomato microbiota is further modulated by the growing

conditions.

Despite this apparent relatedness, the selective enrichment of individual bacterial

members of the microbiota discriminates between rhizosphere and root communities for

mature plants from unplanted soil profiles (Figure 4). These enrichment displayed a bias

for members of the phyla Actinobacteria, Bacteroidetes, Proteobacteria (alpha, beta, delta

and gamma) as well as Verrucomicrobia (Figure 4). Members of these taxa have been

routinely reported in studies focussing on plant-competent bacteria under both laboratory

and field conditions (Bulgarelli et al., 2013; Walters et al., 2018), suggesting that the

experimental approach followed in this study can be considered representative for field-

grown processing tomato.

However, we noticed a differential selective pressure on the bacteria thriving either in the

rhizosphere or in the root tissue: this latter environment produced more distinct profiles,

i.e. more differentially enriched bacteria compared to unplanted soil, than the ones

retrieved from the soil surrounding the roots (compare the number of OTUs in the outer

circles of Figure 4). This indicates that the diversification of the tomato-inhabiting microbial

communities from the surrounding soil biota initiates in the rhizosphere and progresses

through the root tissue, where it produces a more pronounced microbiota diversification

compared to unplanted specimens. This observation is reminiscent of the recruitment

patterns of other crops such as barley (Bulgarelli et al., 2015) but it is in striking contrast

with studies conducted with both model (Bulgarelli et al., 2012) and field-grown (Rathore et

al., 2017) Brassicaceae, whose 'rhizosphere effect' appear negligible.

We further noticed that the "root effect" on the microbiota was exerted also at phylogenetic

level with a bias for the enrichment Actinobacteria. This observation is in apparent contrast

with results gathered from the recent seed-to-seed characterisation of the tomato

microbiota which revealed that, albeit averaging 8% of the sequencing reads across

microhabitats, members of this phylum did not significantly discriminate root from

rhizosphere specimens (Bergna et al., 2018). However, it is worth mentioning that these

two studies differed in terms of both soil type and plant genotype used.

Together, our results suggest that both species- and soil-specific traits govern the

assembly of the tomato microbiota in field-grown crops.

Nitrogen source impacts on the structural and functional composition of the tomato
microbiota

Next, we investigated the impact of the type of nitrogen fertiliser on the tomato microbiota

and we demonstrated that each treatment produced "distinct signatures", represented by

specific selective enrichment, on both the rhizosphere and root communities (Figure 5).

Despite microhabitat-associated variation, the effect of the application of pelleted digestate

(PE) resulted in the most distinct microbial profile in the root compartment and the second

largest number of specifically enriched OTUs in the rhizosphere. Of note, the slow-acting

mineral fertiliser (SRMF) follow a "complementary" pattern: its application yielded the

greatest and the second greatest number of differentially enriched OTUs compared to

untreated samples in the rhizosphere and root profiles, respectively. Remarkably, these

two treatments had a discernible effect also on crop yield, with the PE treatment producing

the best performance among the various fertilisers (Table 1). Our data are congruent with

studies conducted on wheat which observed a structural diversification of the soil and

plant-associated communities exposed to either mineral or organic fertilisers (Kavamura et

al., 2018). Yet, the numerical shift in terms of OTUs differentially enriched per se cannot

explain the potential impact of these communities on crop yield.

We therefore focused our attention on the taxonomical composition of the rhizosphere and

root communities. In particular, we noticed that the proliferation of Actinobacteria in the

root compartment was retained in the various treatments. The enriched Actinobacteria

included *Streptomyces spp., Agromyces sp.*, *Microbispora sp.* and *Actinoplanes spp.*

(Supplementary Database 1). *Streptomyces spp.* are well-known bacteria able to produce

a wide diversity of bioactive compounds able to promote plant growth and health (de Jesus

Sousa and Olivares, 2016). For example, a recent study conducted on tomato revealed

that the exogenous application of *Streptomyces pactum* Act12 to tomato plants increased

host virus resistance and promoted the recruitment of beneficial microbes in the

rhizosphere (Li et al., 2019). On the other hand, members of the genus Streptomyces are

responsible of economically relevant plant diseases, most notably common scab of potato

caused by S*. scabies* (Loria et al., 2006). Remarkably, the molecular characterisation of S.

scabies *87-22* revealed that the pangenome of the species encodes for a tomatinase, a

gene previously identified in tomato pathogens (Seipke and Loria, 2008).

Thus, the taxonomic diversification triggered by both microhabitat and treatment may

underpin a functional diversification of the microbiota at the cross-road of mutualism and

inter-species competition.

This functional diversification of the root communities is manifested in the root

communities by the differential enrichments of ABC transporter genes (mineral) and the

two-component system (organic). ABC transporters have previously been identified as

genes underpinning rhizosphere competence in the microbiota of wheat and cucumber

(Ofek-Lalzar et al., 2014) while the two-component system is required for the rhizosphere

colonisation of the biocontrol agent *Pseudomonas fluorescens* WCS365 (De Weert et al.,

2006). Owing the role played by these classes of genes in uptake of organic compounds

(e.g., root exudates, cellular secretion) and stimulus-response mechanisms (e.g.,

chemotaxis), it is tempting to hypothesize that the different source of nitrogen define a

different metabolic status in and in the vicinity of tomato roots which, in turn, requires a

prompt adaptation of the root-inhabiting communities.

For instance, experimental data indicate that abundance of phytoavailable nitrogen, i.e.,

the scenario of mineral fertiliser treatments, tend to repress the proliferation and activity of

members of the microbiota (Ramirez et al., 2012; Terrazas et al., 2019), and this in turn

may be reflected in the metabolism of secondary compounds (terpenoid and polyketide

metabolism) and membrane transport (ABC transporters).

Congruently with previous observations (Möller, 2015), the digestate treatments appear to

sustain a higher microbial metabolism, reflected by the function associated to energy

metabolism, carbohydrate metabolism and two component systems. Considering that

phytohormons, such as gibberellins, indole acetic acid, auxin-like and auxin-active

molecules have previously been identified in digestate preparations (Scaglia et al., 2017),

it is legitimate to speculate that these preparations may act as both biostimulants and

prebiotics for the plants and their associated microbial communities, respectively.

We further hypothesize that this adaptation is modulated by mineral nitrogen availability,

as manifested by the differential enrichment of functions associated to nitrogen

metabolism *per se* and aminoacids. This observation is congruent with results gathered

from monocots wheat (Kavamura et al., 2018) and rice (Zhang et al., 2019) and suggests

a cross-species pattern whereby plant's adaptation to nitrogen forms and availability is

mediated, at least in part, by the associated microbiota.

Finally, it is interesting to note how the production of antibiotics, namely tetracyline, is also

among the functions differentially enriched between fertilisers. It is becoming increasingly

clear how plant-associated bacteria can act as a reservoir of antimicrobial genes (Cernava

et al., 2019) which can be deployed during inter-organismal competition in the plant

microbiota.

Our investigation suggests that the bacterial microbiota of field-grown processing tomato is

the product of a selective process that progressively differentiates between rhizosphere

and root microhabitats. This process initiates as early as plants are in a nursery stage and

it is then more marked when plants reached the harvest stage. This selection *a*) acts both

on the relative abundances and phylogenetic assignments of members of the tomato

microbiota, *b*) is modulated, at least in part, by the nitrogen fertiliser provided which, in

turn, *c*) triggers different microbial metabolic specialisations within tomato roots.

It is important to mention that the nitrogen fertiliser may also represent a microbial

inoculant *per se*, in particular in the case of organic-based amendments. Indeed, the initial

substrate subjected to biogas production is a determinant of the microbial communities

retrieved from digestate preparations (Studer et al., 2017).

Towards a lab-in-the-field approach to harness the potential of plant microbiota for climate-
smart agriculture
Our experiments represent an example of how cultivation-independent approaches can be

efficiently deployed to investigate the plant microbiota under field conditions. Although this

type of investigation is not novel *per se* in tomato (Toju et al., 2019), our results revealed

fundamentally novel insights into plant's adaptation to nitrogen fertilisers and the

implication for crop yield. Similar to what has recently been postulated for tomato pathogen

protection (Kwak et al., 2018), our results predicts that the use of field-derived, sequencing

data will allow scientists to identify "signatures" of the plant microbiota that can be targeted

to enhance plant performance. This approach, which we define as lab-in-the-field, will be

key towards the rationalisation of nitrogen (and other treatments) application in agriculture

and we anticipate will pave the way for the effective exploitation of the plant microbiota for

agricultural purposes (Schlaeppi and Bulgarelli, 2015; Toju et al., 2018).

507 AUTHORS CONTRIBUTION

508 DR conceived of and designed the field experiment. DR and FC harvested the field data

509 and samples. FC, RAT, EF and DB conceived of and designed the analysis of the

510 microbiota. FC, MC, CVGA, SR-A performed the microbiota experiments. FC, RAT and DB

511 analysed the sequencing data. FC, RAT and DB wrote the initial draft of the manuscript.

512 All the authors discussed the results and commented on the manuscript.

513 FUNDING

This research was partially founded by GENBACCA project (Regione Emilia Romagna,

POR-FESR 2014/2020 Initiative). The generation and analysis of the sequencing data was

supported by a Royal Society of Edinburgh/Scottish Government Personal Research

Fellowship co-funded by Marie Curie Actions awarded to DB. SR-A is supported by a

BBSRC iCASE studentship awarded to DB (BB/M016811/1) and partnered by the James

Hutton Limited (Invergowrie, United Kingdom). RAT and DB are supported by the H2020

Innovation Action 'Circles' (European Commission, Grant agreement 818290) awarded to

the University of Dundee.

522 ACKNOWLEDGEMENTS

We thank Dr Guido Bezzi (CIB - Italian Biogas Association, Italy) for providing us with the

field, plant material and the digestated treatments, Dr. Massimo Zaghi (CAT - COOP.

Agroenergetica Territoriale Correggio S.C.A., Italy) for providing the pelleted digestate and

Dr Stefano Tagliavini (SCAM Spa, Italy) for providing the organo-mineral fertiliser.

We thank Malcolm Macaulay, Jenny Morris and Dr Pete Hedley (The James Hutton

Institute, Invergowrie, UK) for their support in preparing the sequencing library and

generating the sequencing data.

We thank Prof Marco Candela (U Bologna, Italy) for the critical comments on the

manuscript.

CONFLICTS OF INTERESTS

533 The authors declare no conflicts of interest.

LITERATURE CITED

Alburquerque, J.A., de la Fuente, C., Ferrer-Costa, A., Carrasco, L., Cegarra, J., Abad, M.,

and Bernal, M.P. 2012. Assessment of the fertiliser potential of digestates from farm

and agroindustrial residues. Biomass and bioenergy 40:181-189.

Alegria Terrazas, R., Giles, C., Paterson, E., Robertson-Albertyn, S., Cesco, S., Mimmo,

T., Pii, Y., and Bulgarelli, D. 2016. Plant-Microbiota Interactions as a Driver of the

Mineral Turnover in the Rhizosphere. Adv Appl Microbiol 95:1-67.

Asshauer, K.P., Wemheuer, B., Daniel, R., and Meinicke, P. 2015. Tax4Fun: predicting

functional profiles from metagenomic 16S rRNA data. Bioinformatics 31:2882-2884.

Bergna, A., Cernava, T., Rändler, M., Grosch, R., Zachow, C., and Berg, G. 2018. Tomato

seeds preferably transmit plant beneficial endophytes. Phytobiomes Journal 2:183-

193.

Bluman, A.G. 2009. Elementary statistics: A step by step approach. McGraw-Hill Higher

Education New York.

Bulgarelli, D., Schlaeppi, K., Spaepen, S., van Themaat, E.V.L., and Schulze-Lefert, P.

2013. Structure and functions of the bacterial microbiota of plants. Annual review of

plant biology 64:807-838.

Bulgarelli, D., Garrido-Oter, R., Munch, P.C., Weiman, A., Droge, J., Pan, Y., McHardy,

A.C., and Schulze-Lefert, P. 2015. Structure and function of the bacterial root

microbiota in wild and domesticated barley. Cell host & microbe 17:392-403.

Bulgarelli, D., Rott, M., Schlaeppi, K., van Themaat, E.V.L., Ahmadinejad, N., Assenza, F.,

Rauf, P., Huettel, B., Reinhardt, R., and Schmelzer, E. 2012. Revealing structure

and assembly cues for Arabidopsis root-inhabiting bacterial microbiota. Nature

488:91.

Caporaso, J.G., Lauber, C.L., Walters, W.A., Berg-Lyons, D., Huntley, J., Fierer, N.,

Owens, S.M., Betley, J., Fraser, L., Bauer, M., Gormley, N., Gilbert, J.A., Smith, G.,

and Knight, R. 2012. Ultra-high-throughput microbial community analysis on the

Illumina HiSeq and MiSeq platforms. The ISME journal 6:1621-1624.

Caporaso, J.G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F.D., Costello, E.K.,

Fierer, N., Pena, A.G., Goodrich, J.K., Gordon, J.I., Huttley, G.A., Kelley, S.T.,

Knights, D., Koenig, J.E., Ley, R.E., Lozupone, C.A., McDonald, D., Muegge, B.D.,

Pirrung, M., Reeder, J., Sevinsky, J.R., Turnbaugh, P.J., Walters, W.A., Widmann,

J., Yatsunenko, T., Zaneveld, J., and Knight, R. 2010. QIIME allows analysis of

high-throughput community sequencing data. Nature methods 7:335-336.

Cernava, T., Erlacher, A., Soh, J., Sensen, C.W., Grube, M., and Berg, G. 2019.

Enterobacteriaceae dominate the core microbiome and contribute to the resistome

of arugula (Eruca sativa Mill.). Microbiome 7:13.

Chaparro, J.M., Badri, D.V., and Vivanco, J.M. 2014. Rhizosphere microbiome

assemblage is affected by plant development. Isme Journal 8:790-803.

Chialva, M., Salvioli di Fossalunga, A., Daghino, S., Ghignone, S., Bagnaresi, P.,

Chiapello, M., Novero, M., Spadaro, D., Perotto, S., and Bonfante, P. 2018. Native

soils with their microbiotas elicit a state of alert in tomato plants. New Phytologist

220:1296-1308.

Consortium, T.G. 2012. The tomato genome sequence provides insights into fleshy fruit
evolution. Nature 485:635.

de Jesus Sousa, J.A., and Olivares, F.L. 2016. Plant growth promotion by streptomycetes:
ecophysiology, mechanisms and applications. Chemical and Biological
Technologies in Agriculture 3:24.

De Weert, S., Dekkers, L.C., Bitter, W., Tuinman, S., Wijfjes, A.H., Van Boxtel, R., and
Lugtenberg, B.J. 2006. The two-component colR/S system of Pseudomonas
fluorescens WCS365 plays a role in rhizosphere competence through maintaining
the structure and function of the outer membrane. FEMS microbiology ecology
58:205-213.

DeSantis, T.Z., Hugenholtz, P., Larsen, N., Rojas, M., Brodie, E.L., Keller, K., Huber, T.,
Dalevi, D., Hu, P., and Andersen, G.L. 2006. Greengenes, a Chimera-Checked 16S
rRNA Gene Database and Workbench Compatible with ARB. Applied and
Environmental Microbiology 72:5069-5072.

Dombrowski, N., Schlaeppi, K., Agler, M.T., Hacquard, S., Kemen, E., Garrido-Oter, R.,
Wunder, J., Coupland, G., and Schulze-Lefert, P. 2017. Root microbiota dynamics
of perennial Arabis alpina are dependent on soil residence time but independent of
flowering time. Isme Journal 11:43-55.

Edwards, J., Johnson, C., Santos-Medellin, C., Lurie, E., Podishetty, N.K., Bhatnagar, S.,
Eisen, J.A., and Sundaresan, V. 2015. Structure, variation, and assembly of the
root-associated microbiomes of rice. Proceedings of the National Academy of
Sciences of the United States of America 112:E911-920.

Elser, J., and Bennett, E. 2011. A broken biogeochemical cycle. Nature 478:29.

Grigatti, M., Di Girolamo, G., Chincarini, R., Ciavatta, C., and Barbanti, L. 2011. Potential
nitrogen mineralization, plant utilization efficiency and soil CO_2 emissions following
the addition of anaerobic digested slurries. biomass and bioenergy 35:4619-4629.

Kanehisa, M., Araki, M., Goto, S., Hattori, M., Hirakawa, M., Itoh, M., Katayama, T.,

Kawashima, S., Okuda, S., Tokimatsu, T., and Yamanishi, Y. 2008. KEGG for

linking genomes to life and the environment. Nucleic acids research 36:D480-484.

Kavamura, V.N., Hayat, R., Clark, I.M., Rossmann, M., Mendes, R., Hirsch, P.R., and

Mauchline, T.H. 2018. Inorganic Nitrogen Application Affects Both Taxonomical and

Predicted Functional Structure of Wheat Rhizosphere Bacterial Communities.

Frontiers in Microbiology 9.

Kwak, M.-J., Kong, H.G., Choi, K., Kwon, S.-K., Song, J.Y., Lee, J., Lee, P.A., Choi, S.Y.,

Seo, M., and Lee, H.J. 2018. Rhizosphere microbiome structure alters to enable wilt

resistance in tomato. Nature biotechnology 36:1100.

Letunic, I., and Bork, P. 2006. Interactive Tree Of Life (iTOL): an online tool for

phylogenetic tree display and annotation. Bioinformatics 23:127-128.

Li, Y., Guo, Q., Li, Y., Sun, Y., Xue, Q., and Lai, H. 2019. Streptomyces pactum Act12

controls tomato yellow leaf curl virus disease and alters rhizosphere microbial

communities. Biol Fert Soils 55:149-169.

Loria, R., Kers, J., and Joshi, M. 2006. Evolution of plant pathogenicity in Streptomyces.

Annual Review of Phytopathology 44:469-487.

Love, M.I., Huber, W., and Anders, S. 2014. Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. Genome biology 15:550.

McMurdie, P.J., and Holmes, S. 2013. phyloseq: an R package for reproducible interactive

analysis and graphics of microbiome census data. PloS one 8:e61217.

Möller, K. 2015. Effects of anaerobic digestion on soil carbon and nitrogen turnover, N

emissions, and soil biological activity. A review. Agronomy for sustainable

development 35:1021-1041.

Möller, K., and Müller, T. 2012. Effects of anaerobic digestion on digestate nutrient

availability and crop growth: a review. Engineering in Life Sciences 12:242-257.

Ofek-Lalzar, M., Sela, N., Goldman-Voronov, M., Green, S.J., Hadar, Y., and Minz, D.

2014. Niche and host-associated functional signatures of the root surface

microbiome. Nature communications 5:4950.

Parks, D.H., Tyson, G.W., Hugenholtz, P., and Beiko, R.G. 2014. STAMP: statistical

analysis of taxonomic and functional profiles. Bioinformatics 30:3123-3124.

Pulvirenti, A., Ronga, D., Zaghi, M., Tomasselli, A.R., Mannella, L., and Pecchioni, N.

2015. Pelleting is a successful method to eliminate the presence of Clostridium spp.

from the digestate of biogas plants. Biomass and Bioenergy 81:479-482.

Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., and

Glockner, F.O. 2013. The SILVA ribosomal RNA gene database project: improved

data processing and web-based tools. Nucleic acids research 41:D590-596.

Ramirez, K.S., Craine, J.M., and Fierer, N. 2012. Consistent effects of nitrogen

amendments on soil microbial communities and processes across biomes. Global

Change Biology 18:1918-1927.

Rathore, R., Dowling, D.N., Forristal, P.D., Spink, J., Cotter, P.D., Bulgarelli, D., and

Germaine, K.J. 2017. Crop Establishment Practices Are a Driver of the Plant

Microbiota in Winter Oilseed Rape (Brassica napus). Frontiers in Microbiology 8.

Robertson-Albertyn, S., Alegria Terrazas, R., Balbirnie, K., Blank, M., Janiak, A., Szarejko,

I., Chmielewska, B., Karcz, J., Morris, J., Hedley, P.E., George, T.S., and Bulgarelli,

D. 2017. Root Hair Mutations Displace the Barley Rhizosphere Microbiota. Frontiers

in plant science 8:1094.

Ronga, D., Loyelli, S., Zaccardelli, M., Perrone, D., Ulrici, A., Francia, E., Milc, J., and

Pecchioni, N. 2015. Physiological responses of processing tomato in organic and

conventional Mediterranean cropping systems. Sci Hortic-Amsterdam 190:161-172.

Ronga, D., Zaccardelli, M., Lovelli, S., Perrone, D., Francia, E., Milc, J., Ulrici, A., and

Pecchioni, N. 2017. Biomass production and dry matter partitioning of processing

tomato under organic vs conventional cropping systems in a Mediterranean environment. Sci Hortic-Amsterdam 224:163-170.

Ronga, D., Pellati, F., Brighenti, V., Laudicella, K., Laviano, L., Fedailaine, M., Benvenuti, S., Pecchioni, N., and Francia, E. 2018a. Testing the influence of digestate from biogas on growth and volatile compounds of basil (Ocimum basilicum L.) and peppermint (Mentha x piperita L.) in hydroponics. Journal of Applied Research on Medicinal and Aromatic Plants 11:18-26.

Ronga, D., Setti, L., Salvarani, C., De Leo, R., Bedin, E., Pulvirenti, A., Milc, J., Pecchioni, N., and Francia, E. 2019. Effects of solid and liquid digestate for hydroponic baby leaf lettuce (Lactuca sativa L.) cultivation. Sci Hortic-Amsterdam 244:172-181.

Ronga, D., Caradonia, F., Setti, L., Hagassou, D., Giaretta Azevedo, C., Milc, J., Pedrazzi, S., Allesina, G., Arru, L., and Francia, E. (2018b). Effects of innovative biofertilizers on yield of processing tomato cultivated in organic cropping systems in northern Italy. In XV International Symposium on Processing Tomato 1233, pp. 129-136.

Scaglia, B., Pognani, M., and Adani, F. 2017. The anaerobic digestion process capability to produce biostimulant: the case study of the dissolved organic matter (DOM) vs. auxin-like property. Sci Total Environ 589:36-45.

Schlaeppi, K., and Bulgarelli, D. 2015. The plant microbiome at work. Molecular plant-microbe interactions : MPMI 28:212-217.

Schlaeppi, K., Dombrowski, N., Oter, R.G., van Themaat, E.V.L., and Schulze-Lefert, P. 2014. Quantitative divergence of the bacterial root microbiota in Arabidopsis thaliana relatives. Proceedings of the National Academy of Sciences of the United States of America 111:585-592.

Seipke, R.F., and Loria, R. 2008. Streptomyces scabies 87-22 possesses a functional tomatinase. J Bacteriol 190:7684-7692.

Storey, J.D., and Tibshirani, R. 2003. Statistical significance for genomewide studies.

Proceedings of the National Academy of Sciences of the United States of America

100:9440-9445.

Studer, I., Boeker, C., and Geist, J. 2017. Physicochemical and microbiological indicators

of surface water body contamination with different sources of digestate from biogas

plants. Ecological Indicators 77:314-322.

Terrazas, R.A., Pietrangelo, L., Corral, A.M., Torres-Cortés, G., Robertson-Albertyn, S.,

Balbirnie-Cumming, K., Morris, J., Hedley, P.E., Barret, M., and Paterson, E. 2019.

Nitrogen Availability Modulates the Host Control of the Barley Rhizosphere

Microbiota. bioRxiv:605204.

Tilman, D., Balzer, C., Hill, J., and Befort, B.L. 2011. Global food demand and the

sustainable intensification of agriculture. Proceedings of the National Academy of

Sciences of the United States of America 108:20260-20264.

Toju, H., Okayasu, K., and Notaguchi, M. 2019. Leaf-associated microbiomes of grafted

tomato plants. Scientific reports 9:1787.

Toju, H., Peay, K.G., Yamamichi, M., Narisawa, K., Hiruma, K., Naito, K., Fukuda, S.,

Ushio, M., Nakaoka, S., and Onoda, Y. 2018. Core microbiomes for sustainable

agroecosystems. Nat. Plants 4:247-257.

Walters, W.A., Jin, Z., Youngblut, N., Wallace, J.G., Sutter, J., Zhang, W., González-Peña,

A., Peiffer, J., Koren, O., Shi, Q., Knight, R., Glavina del Rio, T., Tringe, S.G.,

Buckler, E.S., Dangl, J.L., and Ley, R.E. 2018. Large-scale replicated field study of

maize rhizosphere identifies heritable microbes. Proceedings of the National

Academy of Sciences.

Zhang, J., Liu, Y.-X., Zhang, N., Hu, B., Jin, T., Xu, H., Qin, Y., Yan, P., Zhang, X., and

Guo, X. 2019. NRT1. 1B is associated with root microbiota composition and

nitrogen use in field-grown rice. Nature biotechnology:1.

Zsögön, A., Čermák, T., Naves, E.R., Notini, M.M., Edel, K.H., Weinl, S., Freschi, L.,

Voytas, D.F., Kudla, J., and Peres, L.E.P. 2018. De novo domestication of wild

tomato using genome editing. Nature biotechnology.

TABLES

Table 1. Composition and information on fertilisers used in this study.

TOC = Total organic carbon; N = Nitrogen; P = Phosphorus; K = Potassium; H2O = water content

Treatment	TOC%	(N)%	(P)%	(K)%	H_2O%	Additional information
Synthetic fertiliser (MF)		41				Ammonium nitrate (N 26%) and calcium nitrate (N 15%)
Pelleted digestate (PE)	39.70	1.50	2.50	2.00	7.80	(Pulvirenti et al., 2015)
Slow acting liquid digestate (SRLD)	3.74	0.34		0.95		Liquid digestate plus the nitrogen stabilizer Vizura® (BASF, 2 L ha^{-1}),
Liquid digestate (LD)	3.74	0.34		0.95		EC 1.07 dS m-1 and pH 8.3
Organo-mineral fertiliser (SC)	10.50	10.00	5.00	15.00	7.00	Produced by SCAM Spa (Modena, Italy), based on solid digestate for the organic fraction
Slow acting synthetic fertiliser (SRMF)		15.00	15.40	15.00		NPK Original Gold® (Compo Expert)

Table 2. Agronomic and qualitative parameter collected at harvest time.

Different letters mean statistical significant differences among treatments by ANOVA.

Means were compared using Bonferroni's test.

Treatment	Marketable yield t ha^{-1}		Brix t ha^{-1}	
No treatmet	61.9	g	2.7	d
MF	159.8	b	7.1	ab
PE	169.1	a	7.3	a
SRLD	153.0	c	7.1	ab
LD	112.0	f	4.9	c
SC	141.8	d	6.0	abc
SRMF	128.6	e	5.7	bc
	P <0.001		P <0.001	

FIGURE LEGENDS

Figure 1. The tomato field described in this study

Image of the filed at (**A**) plant's transplant and details of (**B**) an individual tomato plant and (**C**) the plantation at harvest stage.

Figure 2. The tomato root microbiota is a gated community

Average **(A)** number of observed OTUs, **(B)** Chao 1 index and **(C)** Shannon index computed on the indicated rhizosphere and root specimens. Abbreviations LD (Liquid Digestate), SRLD (Slow acting Liquid Digestate), PE (Pelleted Digestate); SC (Organo-mineral fertiliser); MF (Mineral Fertiliser); SRMF (Slow acting Mineral Fertiliser). Asterisks denote statistically significant differences between microhabitat by non-parametric Wilcoxon rank sum test ($P < 0.01$).

Figure 3. The tomato rhizosphere and root microbiota host compositionally different communities.

PCoA calculated using a weighted UniFrac matrix among the indicated microhabitat and treatments.

Figure 4. The enrichment of Actinobacteria is a distinctive feature of the tomato root microbiota.

Phylogenetic relationships of the OTUs enriched in rhzosphere and root compartment. Individual external nodes represent one of the OTUs enriched in either (or both) rhizosphere or root samples in no treatment conditions (Wald test, P value < 0.01, FDR corrected) whose colour reflects their taxonomic affiliation at Phylum level. A black bar in the outer rings depicts whether that given OTUs was identified in the rhizosphere- or root-enriched sub-communities, respectively. Phylogenetic tree constructed using OTUs 16S rRNA gene representative sequences.

Number of OTUs significantly enriched (Wald test, P value < 0.01, FDR corrected) in the indicated treatment versus untreated controls in (**A**) rhizosphere and (**B**) roots. In each panel, blue bars denote the total number of enriched OTUs for a given treatment, the black bars denote the magnitude of the enrichment in either the individual treatment or among two or more treatments highlighted by the interconnected dots underneath the panels. Abbreviations LD (Liquid Digestate), SRLD (Slow acting Liquid Digestate), PE (Pelleted Digestate); SC (Organo-mineral fertiliser); MF (Mineral Fertiliser); SRMF (Slow acting Mineral Fertiliser).

Prokaryotic functions discriminating between Digestate-based (indicated as 'R-organic': Liquid Digestate; Slow acting Liquid Digestate and Pelleted Digestate) and mineral-based fertilisers (indicated as 'R-mineral': Organo-mineral; Mineral Fertiliser and Slow acting Mineral Fertiliser) retrieved from Tax4Fun functional profiles (Storey's FDR corrected, $p<0.01$).

Environment/Microhabitat

Nursery   Production field; Rhizosphere   Production field; Roots

Microhabitat

● Bulk   ▲ Rhizosphere   ■ Root

Treatment

● Nursery   ● MF   ● LD

● No treatment   ● PE   ● SRLD

● SC   ● SRMF

Root_OTUs

Rhizosphere_OTUs

OTU taxonomy

🟥 Actinobacteria		🟩 Alphaproteobacteria	🟦 Bacteroidetes
🟧 Betaproteobacteria		🟪 Deltaproteobacteria	🟨 Gammaproteobacteria
🟫 Verrucomicrobia		⬜ Other classes	Tree scale: 0.1 ⊢———⊣

95% confidence intervals

ko02010; ABC transporters
ko02020; Two-component system
ko00500; Starch and sucrose metabolism
ko00520; Amino sugar and nucleotide sugar metabolism
ko00052; Galactose metabolism
ko00330; Arginine and proline metabolism
ko00630; Glyoxylate and dicarboxylate metabolism
"ko00260; Glycine, serine and threonine metabolism"
ko00910; Nitrogen metabolism
ko00720; Carbon fixation pathways in prokaryotes
ko00903; Limonene and pinene degradation
ko00562; Inositol phosphate metabolism
ko00902; Monoterpenoid biosynthesis
ko00253; Tetracycline biosynthesis

R_Organic
R_Mineral

Mean proportion (%)

0.0   15.7

Difference in mean proportions (%)

−1.5   −1.0   −0.5   0.0   0.5   1.0

q-value (corrected)

7.60e-3
7.06e-3
7.60e-3
7.06e-3
7.06e-3
8.89e-3
7.06e-3
7.06e-3
7.60e-3
7.06e-3
7.60e-3
7.06e-3
7.06e-3
7.06e-3