1
10 Virulence phenotypes result from an interaction between pathogen ploidy and genetic
11 background
12
13 Dorian J. Feistel, Rema Elmostafa and Meleah A. Hickman*
30 Running title: Pathogen ploidy and genetic background interact to impact fungal virulence
31
32 Department of Biology, Emory University
33 *corresponding author

Abstract

Studying fungal virulence is often challenging and frequently depends on many contexts, including host immune status and pathogen genetic background. However, ploidy has often been overlooked when studying virulence in eukaryotic pathogens. Since fungal pathogens, including the human opportunistic pathogen *Candida albicans*, can display extensive ploidy variation, assessing how ploidy impacts virulence has important clinical relevance. Here, we assessed how *C. albicans* ploidy and genetic background impact virulence phenotypes in both healthy and immunocompromised nematode hosts. In addition to reducing overall host survival, *Candida* negatively impacted host reproduction, which allowed us to survey lethal and non-lethal virulence phenotypes. While we did not detect any global differences in virulence between diploid and tetraploid pathogens, there were significant interactions between ploidy and *C. albicans* genetic background, regardless of host immune function.

Introduction

Virulence is measured by the reduction of host fitness resulting from a host-pathogen interaction[1-3]. Therefore, virulence is not solely the property of the pathogen, but rather the product of the interaction between a host and its pathogen[4,5]. While many biotic and abiotic factors contribute to virulence[6,7], the genotype-by-genotype interaction between hosts and pathogens is a primary determinant of whether a host gets infected and the resulting level of virulence[8,9]. One important, yet understudied, element of an organism's genotype is its ploidy level. Theoretical studies demonstrate that host-pathogens co-evolution should drive hosts towards diploidy and pathogens towards haploidy[10]. However, polyploidy may have an important role in host-pathogen dynamics. For example, polyploidy of some host species is associated with elevated immune response[11,12]. Furthermore, polyploidy and aneuploidy are well documented in fungal pathogens that infect plant and/or animal hosts[13-15]. However there have only been a limited number of studies that investigate whether pathogen ploidy impacts virulence phenotypes and the few that have, often result in contradictory findings[13,16-19].

One potential source of these contradictory results regarding ploidy is the genetic background or allelic composition of the pathogen. Allelic composition refers to not just the specific alleles present in a genome, but the amount of heterozygosity throughout the genome. Pathogen virulence depends on the pathogen's specific allelic combination[20] and phenotypic analysis of diverse clinical isolates within pathogenic species clearly demonstrate that pathogen genetic background contributes to its virulence[21,22]. Ploidy intrinsically impacts allelic composition – haploids contain a single set of gene alleles, whereas diploids and polyploids can

either be homozygous or heterozygous for any given locus, and dominance can mask recessive

alleles. Allelic composition in polyploids is further complicated by multiple allelic ratios, in which

one to four alleles may be present, depending on the mechanism and age of the polyploidization

event. Thus, a major challenge in determining the specific role ploidy has on pathogen virulence

is disentangling it from allelic composition.

The opportunistic fungal pathogen *Candida albicans*, while typically a highly

heterozygous diploid[23-25], shows tremendous ploidy variation, ranging from haploid to

polyploid[26-29] and is highly tolerant of aneuploidy[26,30-34]. *C. albicans* is a commensal in the

human gastrointestinal microbiota and various other niches[35]. Despite its commensal existence,

C. albicans causes a range of infection, including superficial mucosal infections to life

threatening systemic infections[36]. The severity of fungal infection is closely linked to the immune

status of the host, with superficial mucosal infections occurring in healthy individuals and

serious bloodstream infections occurring almost exclusively in immunocompromised hosts, the

latter of which has a mortality of up to 50%[36]. Extensive allelic and ploidy variation, including

loss-of-heterozygosity, aneuploidy and polyploidy, are well-documented in laboratory and

clinical *C. albicans* isolates[21-25,37,38]. However, these large-scale genomic perturbations are often

identified and studied in the context of antifungal drug resistance[22,31,32], and only limited studies

have specifically investigated how *C. albicans* ploidy impacts virulence phenotypes[16-18].

In this study we sought to identify how *C. albicans* ploidy impacts its virulence by using

four diploid-tetraploid pairs of strains, with each pair representing a distinct genetic background.

We assessed virulence by monitoring four measures of host fitness using healthy and

immunocompromised *C. elegans* hosts[39]. While we find almost no overall relationship between

ploidy and virulence, there are detectable differences among *C. albicans* genetic backgrounds

and clear interactions between *C. albicans* genetic background and its ploidy state on virulence

phenotypes. We also observe these interactions in immunocompromised hosts, however, in

some cases, the ploidy-specific pattern and/or the degree of virulence severity is different

between healthy and immunocompromised hosts. Taken together, these results emphasize the

importance of genotypic interactions on virulence phenotypes, in which genotype includes ploidy

state.

Results

Interactions between pathogen ploidy and genetic background impact virulence phenotypes in

healthy hosts

First, we wanted to investigate whether *C. albicans* genetic background differentially
impacted host fitness. To do this we infected *C. elegans* hosts with four genetically distinct
strain backgrounds of *C. albicans.* Two of these genetic backgrounds are laboratory strains: a
'laboratory heterozygous,' which consists of the SC5314 reference strain and a 'laboratory
homozygous,' a derivative of SC5314 in which the genome is completely homozygous. The
other two genetic backgrounds are clinical strains: a 'bloodstream,' isolated from a candidemia
infection of a male immunosuppressed patient and an 'oral/vaginal,' a pair of clinical strains
isolated from a single immune-competent female patient with vulvovaginal candidiasis. For each
C. albicans genetic background, we assessed four measures of host fitness: host survival (Fig.
1A), host lineage growth (Fig. 1B), host brood size (Fig. 1C), and host reproductive timing (Fig.
1D). For survival and lineage growth, all *C. albicans* genetic backgrounds were virulent, with
significant differences observed between uninfected and infected treatments, and the only
statistical difference between *C. albicans* genetic backgrounds was between the laboratory
heterozygous and oral/vaginal strains for host survival. For brood size, only the laboratory
homozygous genetic background was virulent and resulted in significantly smaller host brood
sizes compared to the two clinical *C. albicans* backgrounds. The impact on host reproductive
timing also depended on pathogen genetic background: the laboratory heterozygous and the
clinical bloodstream strains were virulent (i.e. reduced amount of normal reproductive timing)
whereas the other two *C. albicans* genetic backgrounds were not. The laboratory heterozygous
strain impacted host reproductive timing significantly more than the three other pathogen
genetic backgrounds. Together, these results suggest that while pathogen genetic background
may not obviously contribute to host survival phenotypes, it may be important for other
measures of fungal infection that are less lethal.
One factor that may dampen any differences in virulence among *C. albicans* genetic
backgrounds is ploidy. For each genetic background, we used a related diploid strain and a
tetraploid strain (Table S1). The tetraploids for both the 'laboratory' strains were produced via
mating in the laboratory. The 'bloodstream' pair consisted of a diploid strain isolated early in the
infection and its corresponding tetraploid strain was isolated mid- to late- infection following
antifungal treatment. The 'oral/vaginal' pair consisted of a diploid strain isolated from the oral
cavity and its corresponding tetraploid was isolated from the urogenital tract following antifungal
treatment for vulvovaginal candidiasis. There have been limited and conflicted reports on the
role of tetraploid in fungal virulence[16,17]. To assess how pathogen ploidy impacts virulence, we
compared all the diploid strains to the tetraploid strains for host survival (Fig. 1E), lineage
growth (Fig. 1F), host brood size (Fig. 1G), and host reproductive timing (Fig. 1H) in wildtype

hosts. We did not observe a significant difference between these two ploidy states for any of the

host fitness measures tested, suggesting that ploidy may not impact virulence phenotypes.

However, when we account for *C. albicans* strain background, differences between pathogen

ploidy emerge but depend on strain genetic background and thus, we detect a significant

interaction between genetic background and ploidy for host lineage growth ('interaction'

p=0.0354, two-way ANOVA; Fig. S1B), brood size ('interaction' p<0.0001, two-way ANOVA; Fig.

S1B), and reproductive timing ('interaction' p=0.0283, two-way ANOVA; Fig. S1B). While we

cannot directly test for an interaction using host survival curves, we do detect differences

between diploids and tetraploids for each *C. albicans* genetic background (Fig. S1 and Table

S2). Taken together, these results suggest that there is no global pattern in ploidy state and

virulence, but ploidy in combination with genetic background does significantly contribute to *C.*

albicans virulence phenotypes.

When we look at the specific diploid-tetraploid pairs, representing different *C. albicans*

genetic backgrounds, we see significant differences in virulence between the diploid and

tetraploid for at least one fitness measure, for every *C. albicans* genetic background (Fig. 2). For

two genetic backgrounds, the laboratory homozygous and clinical bloodstream strains, the

diploid strain was more virulent than its tetraploid counterpart. For the laboratory heterozygous

and clinical oral/vaginal strains, the tetraploid strain was more virulent than its diploid

counterpart, when differences between ploidy states were detected. Furthermore, these genetic

background specific ploidy patterns are generally consistent across host fitness measures. For

example, the laboratory heterozygous and clinical oral/vaginal tetraploids are also more virulent

than their diploid counterparts for host brood size (Fig 2C), and the clinical bloodstream diploid

was more virulent than its tetraploid counterpart for lineage growth and delayed host

reproduction (Fig. 2B and D). We performed every pairwise comparison between treatments for

host survival (Table S2), lineage growth (Table S3), brood size (Table S4), and reproductive

timing (Table S5) and find significant differences in virulence between different *C. albicans*

genetic backgrounds and ploidy states for most host fitness measures, except host lineage

growth, were very few differences between *C. albicans* strains were detected. Taken together,

these data support that *C. albicans* ploidy does contribute to its virulence phenotypes, but

whether it attenuates or enhances virulence depends on its genetic background.

Host immune status and pathogen genetic background contribute to virulence phenotypes

We have previously established that *C. albicans* and other non-*albicans Candida*

species cause more severe infections in an immunocompromised *C. elegans* hosts compared to

healthy *C. elegans* hosts [39]. Given that we see pathogen ploidy patterns depend on genetic

background for virulence phenotypes in wildtype healthy hosts, we next wanted to assess if we

could detect comparable patterns in immunocompromised hosts. For each *C. albicans* strain,

we assessed four measures of host fitness and compared *C. albicans* virulence in

immunocompromised *sek-1 C. elegans* hosts: host survival (Fig. 3A), host lineage growth (Fig.

3B), host brood size (Fig. 3C), and host reproductive timing (Fig. 3D). Nearly all the *C. albicans*

genetic backgrounds significantly reduced host fitness compared to uninfected controls, the only

exception was the laboratory homozygous strains did not significantly delay host reproduction.

The laboratory homozygous background is also less virulent than the clinical oral/vaginal

genetic background for host. Together, these results suggest that global differences in pathogen

genetic background are revealed in hosts with compromised immune function for both lethal and

non-lethal measures of host fitness.

To assess how pathogen ploidy impacts virulence in immunocompromised hosts, we

compared all the diploid strains to the tetraploid strains for host survival (Fig. 3E), host lineage

growth (Fig. 3F), host brood size (Fig. 3G), and host reproductive timing (Fig. 3H). Similar to

healthy wildtype hosts, we did not observe a significant difference between these two ploidy

states for and of the host fitness measures. However, when we account for *C. albicans* strain

background, differences between pathogen ploidy emerge but depends on strain genetic

background and a significant interaction between genetic background and ploidy is detected for

host lineage growth ('interaction' p=0.0012, two-way ANOVA; Fig. S2B), brood size ('interaction'

p=0.0090, two-way ANOVA; Fig. S2C), and reproductive timing ('interaction' p=0.0167, two-way

ANOVA; Fig. S2D). While we cannot directly test for an interaction using host survival curves,

we do detect differences between diploids and tetraploids for each *C. albicans* genetic

background (Fig. S2 and Table S2). Taken together, these results further support the

observation that there is no global correlation between ploidy state and virulence, but ploidy

does have an important contribution to virulence phenotypes within pathogen genetic

backgrounds regardless of host immune status.

When we look at specific diploid-tetraploid pairs, representing different *C. albicans*

genetic background, we see significant differences in virulence in immunocompromised hosts

between the diploid and tetraploid state for at least one fitness measure, for all *C. albicans*

genetic backgrounds except for the laboratory homozygous (Fig. 4). For the laboratory

heterozygous genetic background, the tetraploid counterpart was more virulent than its diploid

counterpart when differences between ploidy states were detected (lineage growth and brood

size), similar to the pattern observed in healthy hosts (Fig. 2). Furthermore, the clinical

bloodstream diploid was more virulent than its tetraploid counterpart for immunocompromised
host survival and brood size (Fig. 4A and C), similar to the pattern observed in healthy hosts.
However, the oral/vaginal strain had significant differences between its diploid and tetraploid
counterparts for host survival and reproductive timing (Fig. 4A and D), however the tetraploid
was more virulent in the former, while the diploid was more virulent in the latter. We also
performed every pairwise comparison between treatments for host survival (Table S2), lineage
growth (Table S3), brood size (Table S4), and reproductive timing (Table S5) and find significant
differences in virulence between different *C. albicans* genetic backgrounds and ploidy states for
most host fitness measures, except host reproductive timing, were very few differences between
C. albicans strains were detected. These results emphasize that while global differences
between *C. albicans* genetic backgrounds or ploidy states may be difficult to identify in
immunocompromised hosts (Fig. 3), ploidy-specific and genetic background differences in
virulence are detectable across multiple host fitness measures.
216
Ploidy-specific interactions between host and pathogen genotypes.
Finally, we were curious if host immune status impacted the virulence relationship
between each diploid-tetraploid pair of strains. Since there are significant differences for most
host fitness measures between healthy and immunocompromised hosts even when uninfected
(Table S6), we normalized each *C. albicans*-infected host fitness metric to that of the uninfected
control to directly compare the severity of *C. albicans* infection between host genotypes. This
analysis shows that immunocompromised hosts are more significantly more susceptible to *C.*
albicans infection and show larger reductions in survival, brood size and lineage growth
compared to those observed in healthy hosts, while the amount of delayed reproduction caused
by *C. albicans* infection is similar (Fig. 5A-D). We also detect significant interactions between *C.*
albicans strain and host immune status for lineage growth ('interaction' p=0.0004, two-way
ANOVA), brood size ('interaction' p=0.0042, two-way ANOVA; Fig. S2C), and reproductive
timing ('interaction' p=0.0001, two-way ANOVA).
We next examined whether the relationship between ploidy-specific virulence differences
changed depending on host immune status. To do this, we plotted the relative impact of the
diploid (solid lines) and tetraploid (dotted lines) in healthy (N2) and immunocompromised (*sek-*
1) host backgrounds for all four *C. albicans* genetic backgrounds (Fig 5E-H). In general, there
was a high degree of similarity in the relationship between diploid and tetraploids across for both
host genotypes, as indicated by a non-significant interaction term measured by two-way
ANOVA. However, there were a couple of notable exceptions, particularly in the bloodstream

pair of strains. In healthy hosts, we observed the diploid displaying more severe virulence

phenotypes than its tetraploid counterpart for lineage growth and reproductive timing, yet these

differences are diminished in immunocompromised hosts. Thus, we detect a significant

interactions between host immune status and *C. albicans* ploidy for the bloodstream genetic

background. Strikingly, we detect the reverse pattern for host survival with the *C. albicans*

bloodstream diploid and tetraploid strains, in which no detectable differences are observed in

healthy hosts, but the diploid is significantly more virulent than the tetraploid in

immunocompromised hosts, yet regardless of the host context or specific fitness measure, the

diploid is more virulent than its tetraploid counterpart. From these results, we conclude that

interactions between host immune status, pathogen genetic background and ploidy state

determine the severity of virulence phenotypes in *C. albicans*.

Discussion

Here, we sought to understand how the genetic background and ploidy of the human

fungal opportunistic pathogen *Candida albicans* contributes to virulence phenotypes in the

nematode host *C. elegans*. Overall, we found that most of the strains we tested were virulent for

at least one measure of host fitness, but cannot generalize any over-arching patterns in how

ploidy or genetic background contribute to virulence phenotypes. Rather, we have found that

there is a significant interaction between *C. albicans* ploidy and its genetic background. We

detect this interaction in both healthy and immunocompromised host contexts. However,

immunocompromised hosts display more severe fungal infections compared to healthy hosts,

regardless of *C. albicans* ploidy and/or genetic background.

C. albicans was considered an 'obligate' heterozygous diploid for many years [40-43], and

its diploid-tetraploid parasexual cycle only discovered and characterized in the last two

decades[15,44]. Much of the research regarding *C. albicans* ploidy states have focused on the

mechanisms involved with the parasexual cycle and how the ploidy reduction process that

tetraploids undergo promote cellular heterogeneity and genetic variation[26,45-49]. There are only a

small number of studies investigating if different ploidy states impact its virulence. These earlier

studies investigating whether *C. albicans* ploidy impacts virulence had contradictory results –

Hubbard et al demonstrated that derivative tetraploid had similar results in a murine tail vein

infection model as its diploid progenitor[17], whereas Ibrahim et al demonstrate that tetraploids are

less virulent than diploids[16]. Given that tetraploid *C. albicans* have been identified in clinical

settings[37] and that other fungal pathogens display extensive ploidy variation[14,15], assessing how

ploidy impacts virulence has newfound clinical relevance. We think our systematic approach of

comparing paired diploid and tetraploid strain across genetic backgrounds and host contexts

provides a more comprehensive analysis of ploidy and virulence. Furthermore, our results

support the contradictory findings reported earlier, as we observe that sometimes certain

tetraploids are less virulent than diploids, sometimes there are no differences between ploidy

states, and sometimes certain tetraploids are more virulent than diploids (Tables S2-S5).

Importantly, not only does our analysis account for differences in *C. albicans* genetic

background, but, by utilizing *C. elegans* as an infection system we can assess multiple

measures of host fitness beyond host survival. As an opportunistic pathogen, *C. albicans*

causes a wide-range of infections with the most severe infections occurring in

immunocompromised individuals and only a fraction of these severe infections result in patient

death. As such, we need to broaden our scope to investigate fungal virulence beyond a lethal

phenotype. We have previously demonstrated that *C. albicans* infection of *C. elegans* not only

results in reduced survival, but also has reduced fecundity and delays in reproduction[39]. Here,

we have leveraged this infection system to screen for differences in virulence between *C.*

albicans ploidy and genetic backgrounds. While we did not find any overall pattern in virulence

between diploids and tetraploids, we did identify ploidy-specific differences depending on

genetic background (Figs. 2 and 4). Importantly, these specific patterns were consistent across

multiple host fitness measures and host immune status (Fig. 5). Furthermore, we found that

immunocompromised hosts had significantly more severe infections than immune competent

hosts (Fig. 5), similar to what is observed clinically.

It is important to note that the clinical *C. albicans* genetic backgrounds, bloodstream and

oral/vaginal, are not completely isogenic between the diploid and tetraploid strains, and that

some allelic variation exists in addition to their differences in ploidy[37]. It is feasible that some of

the differences in virulence that we observe between the diploid and tetraploids isolates in these

two clinical *C. albicans* backgrounds (i.e. the bloodstream diploid is more virulent than its

tetraploid, whereas the oral/vaginal tetraploid is more virulent than its diploid) is due to these

allelic differences. By also measuring virulence phenotypes in laboratory-derived genomes that

only differ in the number of chromosome sets they contain, we can directly assess the impact

ploidy has on virulence. Here, we still observe different patterns of ploidy-specific virulence,

depending on genetic background. In the laboratory heterozygous genome, we consistently

observe the tetraploid as more virulent than its diploid in both healthy (Fig. 2) and

immunocompromised (Fig. 4) host contexts. However, in the laboratory homozygous genome,

we frequently failed to find any significant differences between diploid and tetraploid for any of

the host fitness measures, the only exception being healthy host survival, in which the tetraploid

was avirulent and the diploid was virulent (Fig. 2A). This result is consistent with previous

findings that *C. albicans* homozygous genomes do not show significant growth differences *in*

vitro or *in vivo* between ploidy states[27,46].

308   In this work we found that ploidy and genetic background interact to contribute to *C.*

albicans virulence (Figs. S1 and S2). We also observe significant interactions between *C.*

albicans strains and host immune status (Fig. 5). These results indicate that virulence is not

simply a binary 'avirulent/virulent' classification, but rather a complex trait and we need to start

dissecting fungal virulence from this perspective. Recently, genome analysis of clinical isolates

revealed genomic features that differ between clinical isolates and SC5314, the laboratory

reference strain of *C. albicans,* and/or identify genetic determinants of antifungal drug

resistance[22,38,50-53]. Only a small number have attempted to identify the genetic loci underpinning

virulence and only two genes were identified and validated, *EFG1*[21,51] and *PHO100*[46].

Importantly, these approaches may overlook other factors and contexts, such as ploidy or host

immune status, that contribute to virulence. While it is clear that there variation in virulence

occurs across *C. albicans* genetic backgrounds, there is still much work to do in elucidating the

drivers of virulence.

Material and Methods

Strains and media

324   For this study, the *C. albicans* strains used are described in Table S1. *C. elegans* N2

Bristol (WT) and *sek-1* mutant[54] were used to test host survival, fecundity and population

growth in healthy and immunocompromised hosts, respectively. *C. elegans* populations were

maintained at 20°C on lite nematode growth medium (NGM, US Biological) with *E. coli* OP50 as

a food source and were transferred to a newly seeded *E. coli* plate every 3-4 days. For survival,

fecundity and lineage growth assays, NGM was supplemented with 0.08 g/L uridine, 0.08 g/L

histidine, and 0.04 g/L adenine to facilitate growth of auxotrophic *C. albicans* strains and 0.2 g/L

streptomycin sulfate to inhibit *E. coli* overgrowth so *C. albicans* strains could proliferate.

Survival and Fecundity Assays

334   Seeding NGM plates and *C. elegans* egg preparation for survival and fecundity assays

were performed as previously described[39]. Briefly, *C. albicans* strains and *E. coli* OP50 were

inoculated in 3 mL of YPD or 5 mL of LB, respectively, and incubated at 30°C for 1-2 days. *C.*

albicans cultures were diluted to a final volume of 3.0 OD_{600} per mL and *E. coli* was pelleted and

washed twice with 1 mL of ddH_2O. The washed pellet was centrifuged for 60 sec at maximum

and any excess liquid removed. The pellet was weighed and suspended with ddH$_2$O to a final

volume of 200 mg/mL. For uninfected treatments, 6.25 μL *E. coli* was mixed with 43.75 μL

ddH$_2$O. For *C. albicans* treatments, 1.25 μL *C. albicans* was mixed with 6.25 μL *E. coli* and

brought to a final volume of 50 μL with ddH$_2$O. All 50 μL was spotted onto the center of a 35mm

supplemented-NGM Lite agar plate and incubated at room temperature overnight before

addition of nematode eggs or transferred nematode.

To synchronize host populations, *C. elegans* populations (~100 nematodes) were

washed off NGM plates with M9 buffer and transferred to a 15 mL conical tube and pelleted by

centrifugation for 2 min at 1200 rpm. The pellet was re-suspended in a 5.25% bleach solution,

transferred to a microcentrifuge tube and inverted for 90-120 sec and subsequently centrifuged

for 30 sec at 1500 rpm. The pellet was washed thrice with 1 mL M9 buffer and resuspended in

500 μL M9 buffer. ~100 eggs were added to uninfected or *C. albicans* treatment plates. 48 h

later, a single L4 nematode (x10 per treatment) was randomly selected and transferred to an

uninfected or *C. albicans* treatment plate and incubated at 20°C. Each founder nematode was

transferred to new treatment plates every 24 h for seven consecutive days. For survival

analysis, we documented whether the founder was alive, dead, or censored (i.e. crawled off the

plate or were otherwise unaccounted). For fecundity analysis, any eggs laid for each 24-hour

interval remained undisturbed on the plate and incubated at 20°C for an additional 24 h and the

number of viable progeny produced per day was scored. Total brood size is the sum of viable

progeny produced over seven days. Delayed reproduction is calculated by the number of

progeny produced on Day 4 or later divided by the total progeny produced for each founder

nematode. All experiments were performed in triplicate or more.

Lineage Growth Assays

Lineage growth assays were performed as previously described[39]. In brief, a single L4

founder nematode (x6 founders per treatment) was randomly selected from a synchronized

population and transferred to a 100mm treatment plate that contained a 300 μl seed of *C.*

albicans and/or *E. coli* and incubated at 20°C for five days in which the founder produces F1

and F2 progeny. The entire population derived from the single founder were washed with M9

buffer and transferred to 15mL conical tubes and brought to a final volume of 10mL. Tubes were

placed at 4°C for 1h to allow the nematodes to settle at the bottom. 20 μL samples were taken

from each population and counted six independent times to determine the final population size

for each founder nematode. All experiments were performed at least in duplicate.

Statistical Analyses

   All statistical analyses were performed with GraphPad Prism. Survival curves were tested for significant differences using log-rank (Mantel Cox) tests. Lineage growth, total brood size and delayed reproduction data sets were tested for normality using the D'Agostino & Pearson omnibus normality test. For comparisons across genetic backgrounds, Kruskal-Wallis and Dunn's multiple comparison tests were performed. For pairwise comparisons between ploidy or host context, Mann Whitney tests were performed. Two-way ANOVAs were performed to test for interactions between *C. albicans* genetic background and ploidy or between *C. albicans* ploidy and host context.

Author contributions

DFJ and MAH devised the experiments. DFJ and RE performed the experiments. DJF and MAH analyzed the data. MAH wrote the manuscript.

Competing interests

The author(s) declare no competing interests.

Literature Cited

1. Read, A. F. The evolution of virulence. *Trends Microbiol.* **2,** 73–76 (1994).
2. Lipsitch, M., Siller, S. & Nowak, M. A. THE EVOLUTION OF VIRULENCE IN PATHOGENS WITH VERTICAL AND HORIZONTAL TRANSMISSION. *Evolution* **50,** 1729–1741 (1996).
3. Cressler, C. E., McLEOD, D. V., Rozins, C., van den Hoogen, J. & Day, T. The adaptive evolution of virulence: a review of theoretical predictions and empirical tests. *Parasitology* **143,** 915–930 (2016).
4. Casadevall, A. & Pirofski, L.-A. Host-Pathogen Interactions: Redefining the Basic Concepts of Virulence and Pathogenicity. *Infect Immun* **67,** 3703–3713 (1999).
5. Jabra-Rizk, M. A. *et al.* Candida albicans Pathogenesis: Fitting within the Host-Microbe Damage Response Framework. *Infect Immun* **84,** 2724–2739 (2016).
6. Katsir, L., Schilmiller, A. L., Staswick, P. E., He, S. Y. & Howe, G. A. COI1 is a critical component of a receptor for jasmonate and the bacterial virulence factor coronatine. *Proc Natl Acad Sci U S A* **105,** 7100–7105 (2008).
7. Mauch-Mani, B. & Mauch, F. The role of abscisic acid in plant-pathogen interactions. *Curr. Opin. Plant Biol.* **8,** 409–414 (2005).
8. Schulte, R. D., Makus, C., Hasert, B., Michiels, N. K. & Schulenburg, H. Multiple reciprocal adaptations and rapid genetic change upon experimental coevolution of an animal host and its microbial parasite. *Proc Natl Acad Sci USA* **107,** 7359–7364 (2010).
9. Lambrechts, L., Scott, T. W. & Gubler, D. J. Consequences of the expanding global distribution of Aedes albopictus for dengue virus transmission. *PLoS Negl Trop Dis* **4,** e646 (2010).

10.   Nuismer, S. L. & Otto, S. P. Host-parasite interactions and the evolution of ploidy. *Proc Natl Acad Sci USA* **101,** 11036–11039 (2004).

11.   Osnas, E. E. & Lively, C. M. Host ploidy, parasitism and immune defence in a coevolutionary snail-trematode system. *Journal of evolutionary biology* **19,** 42–48 (2006).

12.   King, K. C., Seppälä, O. & Neiman, M. Is more better? Polyploidy and parasite resistance. *Biol Lett* **8,** 598–600 (2012).

13.   Lin, X., Nielsen, K., Patel, S. & Heitman, J. Impact of mating type, serotype, and ploidy on the virulence of Cryptococcus neoformans. *Infect Immun* **76,** 2923–2938 (2008).

14.   Zhu, Y. O., Sherlock, G. & Petrov, D. A. Whole Genome Analysis of 132 Clinical Saccharomyces cerevisiae Strains Reveals Extensive Ploidy Variation. *G3 (Bethesda)* **6,** 2421–2434 (2016).

15.   Morrow, C. A. & Fraser, J. A. Ploidy variation as an adaptive mechanism in human pathogenic fungi. *Seminars in Cell & Developmental Biology* **24,** 339–346 (2013).

16.   Ibrahim, A. S. *et al.* Effects of ploidy and mating type on virulence of Candida albicans. *Infect Immun* **73,** 7366–7374 (2005).

17.   Hubbard, M. J., Poulter, R. T., Sullivan, P. A. & Shepherd, M. G. Characterization of a tetraploid derivative of Candida albicans ATCC 10261. *J Bacteriol* **161,** 781–783 (1985).

18.   Suzuki, T., Kanbe, T., Kuroiwa, T. & Tanaka, K. Occurrence of ploidy shift in a strain of the imperfect yeast *Candida albicans*. *J. Gen. Microbiol.* **132,** 443–453 (1986).

19.   Purnell, D. M. & Martin, G. M. Heterozygous diploid strains of Aspergillus nidulans: enhanced virulence for mice in comparison to a prototrophic haploid strain. *Mycopathol Mycol Appl* **49,** 307–319 (1973).

20.   Manning, S. D. *et al.* Variation in virulence among clades of Escherichia coli O157:H7 associated with disease outbreaks. *Proc Natl Acad Sci U S A* **105,** 4868–4873 (2008).

21.   Hirakawa, M. P. *et al.* Genetic and phenotypic intra-species variation in Candida albicans. *Genome Res* **25,** 413–425 (2015).

22.   Ford, C. B. *et al.* The evolution of drug resistance in clinical isolates of Candida albicans. *eLife Sciences* **4,** 3178–27 (2015).

23.   Jones, T. *et al.* The diploid genome sequence of *Candida albicans*. *Proc Natl Acad Sci USA* **101,** 7329–7334 (2004).

24.   Muzzey, D., Schwartz, K., Weissman, J. S. & Sherlock, G. Assembly of a phased diploid Candida albicans genome facilitates allele-specific measurements and provides a simple model for repeat and indel structure. *Genome Biol* **14,** R97 (2013).

25.   Abbey, D., Hickman, M., Gresham, D. & Berman, J. High-resolution SNP/CGH microarrays reveal the accumulation of loss of heterozygosity in commonly used *Candida albicans* strains. *G3 (Bethesda)* **1,** 523–530 (2011).

26.   Hickman, M. A., Paulson, C., Dudley, A. & Berman, J. Parasexual Ploidy Reduction Drives Population Heterogeneity Through Random and Transient Aneuploidy in Candida albicans. *Genetics* **200,** 781–794 (2015).

27.   Hickman, M. A. *et al.* The 'obligate diploid' Candida albicans forms mating-competent haploids. *Nature* **494,** 55–59 (2013).

28.   Gerstein, A. C., Lim, H., Berman, J. & Hickman, M. A. Ploidy tug-of-war: Evolutionary and genetic environments influence the rate of ploidy drive in a human fungal pathogen. *Evolution* **71,** 1025–1038 (2017).

29.   Selmecki, A., Forche, A. & Berman, J. Genomic Plasticity of the Human Fungal Pathogen Candida albicans. *Eukaryotic Cell* **9,** 991–1008 (2010).

30.   Arbour, M. *et al.* Widespread occurrence of chromosomal aneuploidy following the routine production of *Candida albicans* mutants. *FEMS Yeast Res* **9,** 1070–1077 (2009).

31.   Selmecki, A., Bergmann, S. & Berman, J. Comparative genome hybridization reveals widespread aneuploidy in *Candida albicans* laboratory strains. *Mol Microbiol* **55,** 1553–1565 (2005).

32. Selmecki, A., Forche, A. & Berman, J. Aneuploidy and isochromosome formation in drug-
resistant *Candida albicans*. *Science* **313,** 367–370 (2006).

33. Lenardon, M. D. & Nantel, A. Rapid detection of aneuploidy following the generation of
mutants in Candida albicans. *Methods Mol Biol* **845,** 41–49 (2012).

34. Selmecki, A. M., Dulmage, K., Cowen, L. E., Anderson, J. B. & Berman, J. Acquisition of
aneuploidy provides increased fitness during the evolution of antifungal drug resistance.
5, e1000705 (2009).

35. Underhill, D. M. & Iliev, I. D. The mycobiota: interactions between commensal fungi and
the host immune system. *Nature reviews. Immunology* **14,** 405–416 (2014).

36. Brown, G. D. *et al.* Hidden killers: human fungal infections. *Sci Transl Med* **4,** 165rv13
(2012).

37. Abbey, D. A. *et al.* YMAP: a pipeline for visualization of copy number variation and loss of
heterozygosity in eukaryotic pathogens. *Genome Med* **6,** 100 (2014).

38. Ropars, J. *et al.* Gene flow contributes to diversification of the major fungal pathogen
Candida albicans. *Nat Commun* **9,** 1–10 (2018).

39. Feistel, D. J. *et al.* A Novel Virulence Phenotype Rapidly Assesses Candida Fungal
Pathogenesis in Healthy and Immunocompromised Caenorhabditis elegans Hosts.
mSphere **4,** 405 (2019).

40. Olaiya, A. F. & Sogin, S. J. Ploidy determination of *Candida albicans*. *J Bacteriol* **140,**
1043–1049 (1979).

41. Riggsby, W. S., Torres-Bauza, L. J., Wills, J. W. & Townes, T. M. DNA content, kinetic
complexity, and the ploidy question in *Candida albicans*. *Mol Cell Biol* **2,** 853–862 (1982).

42. Berman, J. & Sudbery, P. E. Candida albicans: A molecular revolution built on lessons
from budding yeast. *Nat Rev Genet* **3,** 918–931 (2002).

43. Noble, S. M. & Johnson, A. D. Genetics of Candida albicans, a Diploid Human Fungal
Pathogen. *Annu Rev Genet* **41,** 193–211 (2007).

44. Sherwood, R. K. & Bennett, R. J. Fungal meiosis and parasexual reproduction--lessons
from pathogenic yeast. *Curr Opin Microbiol* **12,** 599–607 (2009).

45. Bennett, R. J. & Johnson, A. D. Completion of a parasexual cycle in *Candida albicans* by
induced chromosome loss in tetraploid strains. *EMBO J* **22,** 2505–2515 (2003).

46. Gerstein, A. C., Lim, H., Berman, J. & Hickman, M. A. Ploidy tug-of-war: Evolutionary and
genetic environments influence the rate of ploidy drive in a human fungal pathogen.
Evolution **71,** 1025–1038 (2017).

47. Hirakawa, M. P., Chyou, D. E., Huang, D., Slan, A. R. & Bennett, R. J. Parasex
Generates Phenotypic Diversity de Novo and Impacts Drug Resistance and Virulence in
Candida albicans. *Genetics* **207,** 1195–1211 (2017).

48. Forche, A. *et al.* The Parasexual Cycle in Candida albicans Provides an Alternative
Pathway to Meiosis for the Formation of Recombinant Strains. *PLoS Biol* **6,** e110 (2008).

49. Berman, J. & Hadany, L. Does stress induce (para)sex? Implications for *Candida
albicans* evolution. *Trends Genet* **28,** 197–203 (2012).

50. Cavalieri, D. *et al.* Genomic and Phenotypic Variation in Morphogenetic Networks of Two
Candida albicans Isolates Subtends Their Different Pathogenic Potential. *Front. Immunol.*
8, 3166–18 (2018).

51. Castillo, L. *et al.* Property Differences among the Four Major Candida albicans Strain
Clades. *Eukaryotic Cell* **8,** 373–387 (2009).

52. Bensasson, D. *et al.* Diverse Lineages of Candida albicansLive on Old Oaks. *Genetics*
211, 277–288 (2019).

53. Wang, J. M., Bennett, R. J. & Anderson, M. Z. The Genome of the Human Pathogen
Candida albicansIs Shaped by Mutation and Cryptic Sexual Recombination. **9,** a019604–
16 (2018).

54.   Kim, D. H. *et al.* A conserved p38 MAP kinase pathway in Caenorhabditis elegans innate
515   immunity. *Science* **297,** 623–626 (2002).
516

Figure 1: Different *C. albicans* genetic backgrounds reduce multiple measures of wildtype *C. elegans* host fitness.

A) Survival curves for *C. elegans* populations that are either uninfected (exposed to just an E. coli food source, grey) or when infected with different *C. albicans* strains (indicated in legend). The number of worms analyzed (n) for each treatment is indicated in Table S1. Statistical significance was tested using pairwise comparisons of survival curves with Log-rank (Mantel-Cox) test. Astrisks denote statistical differences compared to the uninfected control (* indicates p <0.05, **** indicates p < 0.0001). *C. albicans* treatments that share letters are not significantly different, whereas treatments with differing letters are stastically different.
B) Box and whiskers plot of host lineage growth which represents the total population size (representing the number of F1 and F2 progeny) produced within 7 days from a single founder *C. elegans* host infected with *C. albicans*. Boxes indicate the 25-75th quartiles with median indicated. Error bars are the normalized range of the data and circles indicate outliers. The mean and 95% CI of the uninfected control treatment are indicated by the grey dashed line and shaded grey box. Statistical significance was tested using one-way ANOVA. Astrisks denote statistical differences compared to the uninfected control (* indicates p <0.05, *** indicates p < 0.001). C. albicans treatments that share letters are not significantly different, whereas treatments with differing letters are stastically different, post-hoc Dunn's multiple comparison test. **C)** Total brood size and **D)** Average percentage of host progeny produced during days 1-3 of adulthood (normal reproductive timing) of *C. elegans* infected with *C. albicans*. Data and statistical analysis are the same as (B).
E) % host survival on Day 7 for diploid (dip) and tetraploid (tet) *C. albicans* strains (colored symbols indicate specific *C. albicans* genetic background). Statistical significance was tested using Wilcoxon matched-pairs signed rank test and p values are indicated.
F) Lineage growth, **G)** Brood size, and **H)** Reproductive timing of hosts infected with *C. albicans* diploid and tetraploid strains. Data and statistical analysis are the same as for (E).

Figure 2: Ploidy-specific differences across *C. albicans* genetic backgrounds.
A) Survival curves for *C. elegans* populations that are either uninfected (exposed to just an E. coli food source, grey) or when infected with diploid or tetraploid *C. albicans* strains from laboratory homozygous (pink), laboratory heterozygous (green), bloodstream (orange), or oral/vaginal (blue) genetic backgrounds. Statistical significance was tested using pairwise comparisons of diploid and tetraploid survival curves with Log-rank (Mantel-Cox) test and p-values are indicated and significant differences are highlighted in bold text.
B) Box and whiskers plot of host lineage growth which represents the total population size (representing the number of F1 and F2 progeny) produced within 7 days from a single founder *C. elegans* host infected with *C. albicans*. Boxes indicate the 25-75th quartiles with median indicated. Error bars are the normalized range of the data and circles indicate outliers. The mean and 95% CI of the uninfected control treatment are indicated by the grey dashed line and shaded grey box. Statistical significance between diploid and tetraploid strains was determined using Mann Whitney test and p-values are indicated and significant differences are highlighted in bold text.
C) Total brood size and **D)** Average percentage of host progeny produced during days 1-3 of adulthood (normal reproductive timing) of *C. elegans* infected with diploid and tetraploid *C. albicans*. Data and statistical analysis are the same as (B).

Figure 3: Immunocompromised hosts are highly susceptible to *C. albicans* infection regardless of genetic backgrounds or ploidy.

A) Survival curves for *sek-1 C. elegans* populations that are either uninfected (exposed to just an *E. coli* food source, grey) or when infected with different *C. albicans* strains (indicated in legend). Statistical significance was tested using pairwise comparisons of survival curves with Log-rank (Mantel-Cox) test. Astrisks denote statistical differences compared to the uninfected control (* indicates p <0.05, **** indicates p < 0.0001). *C. albicans* treatments that share letters are not significantly different, whereas treatments with differing letters are stastically different.

B) Box and whiskers plot of host lineage growth which represents the total population size (representing the number of F1 and F2 progeny) produced within 7 days from a single founder *sek-1 C. elegans* host infected with *C. albicans*. Boxes indicate the 25-75th quartiles with median indicated. Error bars are the normalized range of the data and circles indicate outliers. The mean and 95% CI of the uninfected control treatment are indicated by the grey dashed line and shaded grey box. Statistical significance was tested using one-way ANOVA. Astrisks denote statistical differences compared to the uninfected control (* indicates p <0.05, *** indicates p < 0.001). *C. albicans* treatments that share letters are not significantly different, whereas treatments with differing letters are stastically different, post-hoc Dunn's multiple comparison test.

C) Total brood size and **D)** Average percentage of host progeny produced during days 1-3 of adulthood (normal reproductive timing) of *sek-1 C. elegans* infected with *C. albicans*. Data and statistical analysis are the same as (B).

E) % *sek-1* host survival on Day 7 for diploid (dip) and tetraploid (tet) *C. albicans* strains (colored symbols indicate specific *C. albicans* genetic background). Statistical significance between diploid and tetraploids was tested using Wilcoxon matched-pairs signed rank test and p values are indicated.

F) Lineage growth, **G)** Brood size, and **H)** Reproductive timing of *sek-1* hosts infected with *C. albicans* diploid and tetraploid strains. Data and statistical analysis are the same as for (E).

Figure 4: Ploidy-specific differences across *C. albicans* genetic backgrounds in immunocompromised hosts.
A) Survival curves for *sek-1 C. elegans* populations that are either uninfected (exposed to just an E. coli food source, grey) or when infected with diploid or tetraploid *C. albicans* strains from laboratory homozygous (pink), laboratory heterozygous (green), bloodstream (orange), or oral/vaginal (blue) genetic backgrounds. Statistical significance was tested using pairwise comparisons of diploid and tetraploid survival curves with Log-rank (Mantel-Cox) test and p-values are indicated and significant differences are highlighted in bold text.
B) Box and whiskers plot of host lineage growth which represents the total population size (representing the number of F1 and F2 progeny) produced within 7 days from a single founder *sek-1 C. elegans* host infected with *C. albicans*. Boxes indicate the 25-75th quartiles with median indicated. Error bars are the normalized range of the data and circles indicate outliers. The mean and 95% CI of the uninfected control treatment are indicated by the grey dashed line and shaded grey box. Statistical significance between diploid and tetraploid strains was determined using Mann Whitney test and p-values are indicated and significant differences are highlighted in bold text.
C) Total brood size and **D)** Average percentage of host progeny produced during days 1-3 of adulthood (normal reproductive timing) of *sek-1 C. elegans* infected with diploid and tetraploid *C. albicans*. Data and statistical analysis are the same as (B).

Figure 5: Ploidy-specific interactions between healthy and immunocompromised hosts.
A) Relative impact on host survival from Day 7 survival data (*C. albicans* D7 survival/uninfected D7 survival for each host type) for all *C. albicans* strains (colored symbols indicate specific *C. albicans* genetic background) in healhty (N2) and immunocompromised hosts (*sek-1*). The mean and 95% CI of the uninfected control treatment are indicated by the grey dashed line and shaded box. Statistical significance between host genotypes was tested using Wilcoxon matched-pairs signed rank test and p values are indicated.
B) Box and whiskers plot of relative host lineage growth, **C)** Brood size, and **D)** Reproductive timing between healthy. Data and statistical analysis are the same as (A).
E) Relative impact of *C. albicans* ploidy on host survival. Relative virulence of diploid (solid lines) and tetraploid (dotted lines) *C. albicans* lab homozygous (pink), lab heterozygous (green), bloodstream (orange), and oral/vaginal (blue) genetic backgrounds in healthy (N2) and immunocompromised hosts (sek-1). Y-axis scale bar is the same as in (A).
F) Relative host lineage growth, **G)** brood size, and **H)** reproductive timing between healthy and immunocompromised hosts across *C. albicans* genetic backgrounds. Symbols represent the mean value and error bars +/-SD. Y-axis scale bar is the same as in (B, C and D) respectively. Stastistical significance was tested by two-way ANOVA and 'interaction' p value indicated.