Demographic history of a remote ichthyofauna assemblage reveals common

processes driving colonization and persistence in endemic coral reef fishes

Erwan Delrieu-Trottin[1,2,3], Nicolas Hubert[2], Emily C. Giles[1],Pascaline Chifflet-Belle[4,5],

Arnaud Suwalski[4,5], Valentina Neglia[1], Cristian Rapu-Edmunds[6], Stefano Mona[4,5,†],

Pablo Saenz-Agudelo[1,†]

[1]Instituto de Ciencias Ambientales y Evolutivas, Universidad Austral de Chile, Valdivia,

Chile

[2]Institut de Recherche pour le Développement, UMR 5554 (UM-CNRS-IRD-EPHE),

ISEM, Montpellier, France.

[3]Museum für Naturkunde, Leibniz-Institut für Evolutions-und Biodiversitätsforschung

an der Humboldt-Universität zu Berlin, Berlin, Germany

[4] Institut de Systématique, Evolution, Biodiversité, ISYEB, Muséum

national d'Histoire naturelle, CNRS, Sorbonne Université, EPHE, Université des

Antilles, Paris, France

[5]EPHE, PSL Research University, Paris, France

[6]Mike Rapu Diving Center, Caleta Hanga Roa O'tai, Chile

[†]These authors jointly supervised this work.

Author for correspondence: Erwan Delrieu-Trottin, e-mail:

erwan.delrieu.trottin@gmail.com

Keywords: endemism, community assembly, colonization, ddRADs, Easter Island,

population genetics

Abstract

Elucidating demographic history during the settlement of ecological communities is

crucial to properly inferring the mechanisms that shape species diversity and their

persistence through time. We used genomic data to elucidate for the first time the

demographic dynamics associated with the settlement by endemic reef fish fauna of

one of the most isolated islands of the Pacific Ocean, Rapa Nui (Easter Island). We

compared the demographic history of nine endemic species including seven small-

range (restricted to Rapa Nui and Motu Motiro Hiva) and two large-range (present in

Rapa Nui and other southern subtropical islands of the Pacific) endemic species in

order to explore the onset of community settlement and associated demographic

history. We found that most Rapa Nui endemic species share a common demographic

history, with a demographic expansion initiated during the last interglacial period and

related to the last sea level high-stand. The commonality of this pattern suggests that

eustatic fluctuations associated with Milankovitch cycles have played a central role in

species demographic history and the final stage of contemporary community assembly

of Rapa Nui reef fishes. We discuss the potential role of seamounts in the colonization /

extinction / recolonization dynamics of populations of the Rapa Nui Archipelago.

1. Introduction

Local and regional processes jointly drive community assembly, *i.e.* the dynamics of species aggregations in ecological communities and the number of coexisting species at a given place. The importance and nature of local (e.g. environmental filtering, species interactions inside a community) and regional (e.g. dispersal, immigration of new species from outside of a community) processes in the origin and maintenance of biodiversity in communities is one of the oldest question in ecology [1,2]. Phylogenetic analysis of community assembly can give a better understanding of the relative importance of local and regional dynamics in shaping community assembly [3] as they provide a temporal framework [4]. However, the difficulty of reconstructing accurate divergence times between species due to the incompleteness of available phylogenies limit the power of such analyses. In the meantime, community assembly, especially in island systems, is known to be dynamic, with regular colonization / extinction / and recolonization of populations occurring through time [2,5,6], processes which cannot be assessed using species level approaches. Investigating shallower time scale, population genetics can provide insights into both assembly time and species demographic history during community assembly, hence helping understanding the mechanisms by which species persist through time. Thus by comparing the demographic histories of multiple species in a community and estimating the timing of the early stages of community assembly, it is possible to disentangle not only the relative contribution of historical (*e.g.* tectonic, eustatic changes) and ecological (*e.g.* competitive exclusion, habitat filtering) dynamics but also the scale of the processes (local vs. regional) in shaping the final stage of contemporary community assembly.

Rapa Nui (Easter Island, Chile, 166 km^2) is one of the most remote islands on Earth and hosts an exceptional reef fish community. The mechanisms at the origin of

this diversity and its persistence through time remain enigmatic. Rapa Nui (RN) is the

second hotspot of endemism of the Indo-Pacific Ocean, with 21.7% of fishes endemic

to these islands [7,8]. This reef fish community is known for being extremely species

poor (only 169 fish species; 139 shore fishes) compared to the species-rich islands of

the Indo-Australian Archipelago for instance. RN and the islet Motu Motiro Hiva (MMH)

(Salas y Gómez, 0.15 km^2), located 400 km further east, which shares the same reef

fish fauna as RN [9], constitute the only two emerged islands of the Easter Chain. RN

and MMH are relatively young; 2.5 and 1.7 My, respectively [10], but they are

embedded in a network of numerous seamounts. This seamount chain extends 2,232

km east to the Nazca seamount (23°360 S et 83°300 W) [10,11] and these mounts

have emerged to various degrees during periods of low sea level. In this context, an

"Ancient Archipelago" hypothesis has been formulated to account for the high level of

endemism observed in RN and MMH, young islands, hypothesizing that "most if not all

endemic species were acquired from elsewhere in the region" [12]. Seamounts, which

were once likely islands, could have provided potentially suitable habitat for at least the

past 29 My for endemics of the region to evolve and persist up to present times [12].

However, a recent analysis of the divergence times of endemic species from their

closest relatives has shown that small-range endemics are not older that the

emergence of RN and MMH; they are thus neoendemics [13], thereby questioning the

ancient origin of RN endemics. Yet, the seamounts nearby RN could have played a

major role in long-term species persistence through time, especially during the

Pleistocene. This era is characterized by notable Milankovitch cycles, historical

processes resulting in the alternation of glacial and inter-glacial time that resulted in

sea levels 150 m below present level. Such paleo-environmental perturbations might

be expected to have left a footprint of bottlenecks and expansions in the demographic

history of populations of many marine species [14–17]. These major sea level changes

altered shallow water habitats; exposing continental shelves during low sea levels but

also giving access to habitat otherwise too deep to colonize such as relatively shallow

seamounts. Thus, these features of the Pleistocene could have temporally expanded

the range distribution of endemic species and their population sizes[18].

Reef fish community assembly in remote islands is dependent on colonization /

extinctions / recolonization processes. The RN reef ichtyofauna is very dynamic;

several species that were reported as being abundant in 1969 in RN waters were then

reported as rare or even absent 16 years later. Along the same line, numerous species

are reported as present in RN waters but are actually vagrants, i.e. species that

colonize RN from time to time without being able to establish a population locally [7].

The endemic RN reef ichtyofauna is composed of two types of endemics, small-range

and large-range endemic species. It is possible to hypothesize that *local* processes, at

the scale of the RN and MMH community, influence the demographic history of small-

range endemics as these fishes are restricted to these two islands. On the contrary,

large-range endemics are present both in RN and in other southern subtropical islands

of the Pacific. As such, populations of these fishes are embedded in metapopulations

of larger geographic ranges than those of small-range endemics. Thus the population

dynamics of large-range endemics are likely to also be affected by *regional* processes

such as fragmentation and dispersal outside of the RN and MMH community.

Through genome-wide sequencing (ddRAD), we explore the demographic

history of nine endemic species of RN. The species studied here represent the two

types of endemics (small-range and large-range), seven major reef fish families, and

three different reproductive strategies endemics. We examine whether demographic

histories vary (1) according to the current range distribution of the species; (2) with life

history traits; doing so, we investigate the consequence of historical processes such as

sea level changes or ecological dynamics driven by reproductive strategy on the

genetic diversity. If historical processes are more important than ecological processes

in the maintenance of endemic species in RN, we expect to find similar population

dynamic for the two types of endemic species. In the same way, if local processes are

more important than regional processes in the maintenance of endemic species in RN,

we expect to find similar assembly time for the two types of endemic species.

Implications for the biogeography of the area are discussed.

2. Material and methods

(a) Sampling

A total of 143 reef fishes (nine species) were collected using polespears or an

anesthetic (clove oil) in Rapa Nui in October 2016 (Table 1). Species were classified as

either small-range or large-range endemics. Small-range endemics only present

around RN and MMH and have a maximum range of <500 km in linear distance, see

Delrieu-Trottin et al. [19]. Southern subtropical endemics have large-ranges (1,000–

8,000 km in linear distance, see Delrieu-Trottin et al. [19], hereafter large-range

endemic species) and are distributed from Southern Polynesia to RN (regional

endemics in Friedlander et al. [9]). The species analyzed here also possess different

reproductive strategies, with five species producing pelagic eggs, three species

producing demersal eggs, and one species brooding eggs in their mouths (Table 1).

(b) Library preparation and sequencing

Whole genomic DNA was extracted from fin or gill tissue preserved in 96% ethanol

using the GeneJet Genomic DNA purification kit according to the manufacturer's

protocols (Thermo Fisher Scientific). Double-digest restriction-associated DNA

(ddRAD) libraries were prepared following Peterson et al.' protocol [20]. The genomic

libraries obtained were sequenced in 3 lanes of a HiSeq 2500 Illumina sequencer

(single end, 125 pb). Illumina reads are available from the Sequence Read Archive

(SRA) at NCBI under the Accession nos. XXXXXX–XXXXXXX.

(c) De novo assembly

We used the 'process_radtags.pl' pipeline in STACKS version 2.0 [21,22] to

demultiplex and quality filter the sequences obtained. In the absence of reference

genomes for the species under study, RADSeq loci were assembled de novo using the

'denovo_map.pl' pipeline in STACKS. We used the parameter combination

recommended by Mastretta-Yanes et al. [23]; this included minimum read depth to

create a stack (m) = 3, number of mismatches allowed between loci within individuals

(M) = 3, number of mismatches allowed between loci within catalogue (n) = 3 and

required a locus to be present in all individuals of each species (r = 1). Following de

novo mapping, an initial data-filtering step was performed using the population

component of STACKS removing all loci with maximum observed heterozygosity higher

than 0.8. We first kept all single-nucleotide polymorphisms (SNP) per stack (*i.e.* locus)

and did not use any threshold regarding the minor allele frequencies as we obtained a

high coverage for each individual. We then removed all loci displaying more than three

SNPs to avoid potential paralogs. Summary statistics based on the resulting vcf file

(Supplementary material) such as nucleotide diversity (π on variable loci),

heterozygosity (*H*) and F_{IS}, were calculated for every SNP using

the *populations* program in STACKS. The folded Site (Allele) Frequency Spectrum

(SFS) was computed in R [24] with the package *pegas* [25].

(d) Demographic analyses

Variation in the effective population size (*Ne*) through time was investigated using the

composite likelihood approach implemented in the software *stairwayplot* [26]. The

stairwayplot is a non-parametric model where *Ne* is free to vary at each coalescent

interval. The composite likelihood is evaluated as the difference between the observed

SFS and its expectation under a specific demographic history. To confirm the results

obtained by the *stairwayplot*, we ran an additional approximate Bayesian computation

algorithm based on coalescent simulations following Maisano Delser et al [27]. Briefly,

we performed 1,000,000 coalescent simulations of a demographic model with three

instantaneous changes of *Ne*. The model is therefore defined by seven parameters:

four values of *Ne* and three instantaneous time changes (hereafter, *T*). We set the

same uniform distribution for the four *Ne* values and incremental uniform distribution for

the three *T* parameters, similarly to Maisano Delser et al [27]. Coalescent simulations

and SFS computation were performed with *fastsimcoal* [28]. We used the SFS and the

mean pairwise differences (computed with a custom R script) as summary statistics.

We retained the best 5,000 simulations to perform a local linear regression [29] and

reconstructed the abc skyline at user specified time points starting from the posterior

distribution as in Maisano Delser et al [27]. We used a mutation rate (μ) of 1.0×10^{-8} /

site / generation following [30–34] and generation times found in the literature (Table 1)

or inferred using maximum standard length.

As data did not meet assumptions of normality, we computed the Wilcoxon-

Mann-Whitney and Kruskal-Wallis non parametric tests to examine differences in the

Time to the Most Recent Common Ancestor (TMRCA) considering four factors: range

size, family, generation time, and reproductive strategies. All statistical analyses were

performed in R, using the package *vegan* [35] and *ggplot2* was used for graphical

representations [36].

3. Results

Raw sequence filtering, assembly, and SNP calling. A total of 31 543 967 reads of 121

bp each were obtained for the 143 individual samples from the nine species endemic to

RN. An average of 3 504 885 reads were found per species (min: 2 983 062, max: 4

430 571). The different filtering steps resulted in the building of an average of 41 470

loci (min: 25 331; max: 63 590) per species out of which an average of 23 693 (min: 19

246; max: 28 855) were variable (Table 2). The variable loci harbored an average of 37

107 SNPs (min: 28 239; max: 46 718). The depth of coverage per SNP ranged from

31x to 50x (43x averaged across species).

*Genetic diversity statistics***.** Different levels of genetic diversity were found for the nine

species (Table 2). *Cantherhines rapanui* displayed the highest values of heterozygosity

(max: 0.262) and nucleotide diversity (32.73 x 10^{-4}). *Chrysiptera rapanui* showed the

lowest values of heterozygosity (0.136) while *Ostorhinchus chalcius* displayed the

lowest values of nucleotide diversity (0.063).

Community assembly time. The mean TMRCA retrieved with the stairway plots for the

nine species was very recent 253,871 YBP (+/- 119,482 YBP) and ranged from

105,232 YBP (*Ostorhinchus chalcius*) to 398,058 YBP (*Sargocentron wilhelmi*) (Figure

1). The TMRCAs retrieved did not differ significantly among types of endemics (W = 5,

p = 0.6667), families (H_6 = 7.4667, *p* = 0.2798), or reproductive strategy (H_2 =2.56, *p* =

0.278). The only significant factor was generation time (H_2 = 6.1111, *p* = 0.0471).

Changes in historical effective population size. We recovered very similar demographic

patterns with the two methods (stairway plots and the abc skylines; Figure 1, S1).

Patterns of population expansions were recovered for seven out of nine endemic

species studied, constant population size through time was found for one species

(*Sargocentron wilhelmi*) and a decrease in population size was recovered for one

species (*Cantherhines rapanui*). Expansion times retrieved were highly similar between

the two methods for four out of the seven species, while abc skylines estimated older

expansion times than those inferred with the stairway plots for the three other cases. In

the same way, the bottleneck found for *Cantherhines rapanui* was dated slightly older

with the abc skyline than with the stairway plot analysis.

4. Discussion

Rapa Nui endemic species share a common history, with population expansions

dominating the demographic histories dated during the last interglacial period for both

small-range and large-range endemics. These expansions indicate that their current

effective population size established in RN during the last glacial maxima (25 000 - 115

000 YBP), a period characterized by climatic cooling and decreased sea levels [37].

These results agree with those found for expansions and bottlenecks of other marine

organisms suggesting that sea level oscillations during the last 800 000 years level

[38,39] has greatly influenced reef fish populations [40–49]. Rapa Nui endemics,

however, differ in their ecology compared to organisms of other reefs in the Pacific

region, as they are far less restricted in terms of depth range. Many of the RN endemic

species have the particularity to be found from shallow waters to the mesophotic zone

(100 - 150 m) in RN water and in seamounts nearby RN (Table 1). Considering the

network of seamounts surrounding RN, low sea levels during glacial maxima would

have produced periods of maximum reef habitat extension (Figure 2). As such,

population expansions during glacial maxima could reflect the colonization /

recolonization history of RN from MMH or other seamounts of the RN archipelago.

While the Ancient Archipelago hypothesis of Newman & Foster [12] has been

invalidated at the species level for this island system [13], this hypothesis could explain

what is observed at the population level for RN endemic species (i.e. a recolonization

from the seamounts rather than from elsewhere in the Pacific). Overall, historical

metapopulation dynamics within the RN Archipelago combined with sea level

fluctuations during glacial maxima could have driven the population dynamics and

hence community assembly of RN endemics.

The common demographic and temporal patterns found for large-range and

small-range endemics provide insights into the mechanisms that shaped the RN

ichthyofauna. Community assembly in remote islands is expected to rely on

colonization / extinction / recolonization processes. The concordant expansions

detected for all species except *Cantherines rapanui* and *Sargocentron wilhelmi* suggest

a concomitant establishment of current population sizes during the last glacial cycle.

This pattern does not seem to be influenced by the size of the species range

distribution as both small-range and large-range endemics display similar patterns.

Actually, populations of large-range endemics sampled around RN could potentially

operate at a regional scale, being also present in several other islands of the South

Pacific, but the similar expansion times observed here for both types of endemism

argue in favor of population demography driven by local historical processes.

Local processes rather than regional processes seem to have driven the final stage of

community assembly among endemic species around RN, as indicated by the similar

TMRCAs retrieved here for both types of endemism. This suggests that the

demography of these species has been poorly affected by neighbouring populations.

This is consistent with the isolation of RN, as supported by the lower species richness

observed there compared to in other Pacific islands. The low species diversity in RN

reefs might not only be due to the difficulty for larvae to colonize such remote islands,

but also due to the difficulty to establish a viable population / metapopulation that

functions at such local scales. This hypothesis is reinforced by the fact that the two

large range endemics studied here are among the most abundant species of RN [9],

yet they are rare throughout the rest of their distribution [50]. Assessing the amount of

gene flow between RN populations with those of Pitcairn or Austral Island for these

large range endemics could help to test this hypothesis.

The relatively small range of TMRCAs retrieved here could also reflect the

speciation itself for small-range endemics. This has been hypothesized for French

Polynesian small-range endemics which display similarly population expansions older

than the Last Glacial Maximum [17]. Speciation is not an instantaneous process, and

could takes at least 2 My to complete in various vertebrate clades [51]. Divergence

times, upper bound of the speciation time, are known to be quite recent for several

small-range species of this study (*Cantherhines rapanui* 0.54 Ma (0.07 - 1.3 95%

HPD), *Chrysiptera rapanui* 1.39 Ma (0.50 - 2.83 95% HPD), *Chromis randalli* 2.02 (0.58

- 4.3 95 % HPD), *Coris debueni*: 2.31 (0.72 - 4.11 95% HPD); see [13]). Only a study in

a genomic framework based on pairs of sister species could provide a better estimate

of the timing of speciation (**e.g** [33]).

Coalescence time does vary across loci and across sets of individuals. By

screening thousands of loci, the inferences made on the demographic history of the

populations should not be biased by the loci observed. However, uncertainty in both

mutation rates and generation times can potentially bias molecular dating. Mutation

rates for SNP data of fish range from 2.5×10^{-8} [52] to $3.5 \ 10^{-9}$ [53] and their selection

for estimating demographic histories is not often justified. As such, we chose the fish

SNP mutation rate most often used in the litterature. Finally, estimations of generation

times in the wild for fishes are scarce. Most of the generation times we used (six out of

nine) were selected from the literature from species that are phylogenetically close-

related to the species studied here (same genus). Additionally, we used maximal length

to infer generation times for three species as maximum standard length and generation

time are positively correlated [54].

Conclusion

We elucidate for the first time the community assembly time of the RN endemic

reef fish community using genomic data and demographic inferences. We show that

the RN reef fish endemic community shares a common history, with expansion

occurring during the last glacial cycle for the two different types of endemic species.

Local processes based on the seamounts system around RN should have played a

major role in the foundation and persistence of this endemic reef fish community

through dynamic of colonization / extinction / recolonization throughout the RN

Archipelago.

Ethics. All applicable institutional guidelines for the care and use of animals were

followed. Specimens were collected under permit No. 724, 8 March 2016 obtained from

the Chilean Subsecretary of Fishing. The Universidad Austral de Chile Ethical Care

Committee and Biosecurity Protocol approved our use and handling of animals.

Data accessibility. All Fastq sequence files are available from the GenBank at the

National Center for Biotechnology Information short-read archive database (accession

number: *forthcoming*).

Authors' contributions.

E.D.-T. and P.S.-A. conceived the study; E.D.-T., P.S.-A. and S.M. acquired the

funding; E.D.-T., E.C.G., V.N, C.R.E. and P.S.-A., collected the field data; EDT, P.C.-

B. and A.S. produced the data; E.D.-T.; E.D.-T., N.H., S.M, and P.S.-A. analyzed the

data; E.D.-T., N.H., E.C.G., S.M, and P.S.-A wrote the manuscript. All authors gave

final approval for publication.

Competing interests. We declare we have no competing interests.

Funding. This work was supported by the FONDECYT Postdoctorado fellowship

N°3160692 to E. Delrieu-Trottin and by the ATM grant 2016 from the Museum National

d'Histoire Naturelle to Stefano Mona.

Acknowledgements. We thank Rebeca Tepano, Nina, Taveke Olivares Rapu, Liza

Garrido Toleado (SERNAPESCA), Ludovic Tuki (Mesa del Mar, Te Mau O Te Vaikava

O Rapa Nui), and the people of the Rapa Nui Island for their kind and generous

support. We thank Luiz A. Rocha for kindly sharing photographs and information

related to maximum depth of several RN endemic species. We are grateful to the

Genotoul bioinformatics platform Toulouse Midi-Pyrenees (Bioinfo Genotoul;

http://bioinfo.genotoul.fr/) for providing computing resources.

References

1. Macarthur R, Levins R. 1967 The Limiting Similarity, Convergence, and

Divergence of Coexisting Species. *Am. Nat.* (doi:10.1086/282505)

2. MacArthur RH, Wilson EO. 1967 *The Theory of Island Biogeography*. Princeton

university press.

3. Emerson BC, Gillespie RG. 2008 Phylogenetic analysis of community assembly

and structure over space and time. *Trends Ecol. Evol.* **23**, 619–630.

(doi:10.1016/j.tree.2008.07.005)

4.   Graham CH, Fine PVA. 2008 Phylogenetic beta diversity: Linking ecological and

evolutionary processes across space in time. *Ecol. Lett.* **11**, 1265–1277.

(doi:10.1111/j.1461-0248.2008.01256.x)

5.   Whittaker RJ, Triantis KA, Ladle RJ. 2008 A general dynamic theory of oceanic

island biogeography. *J. Biogeogr.* (doi:10.1111/j.1365-2699.2008.01892.x)

6.   Ricklefs RE, Bermingham E. 2002 The concept of the taxon cycle in

biogeography. *Glob. Ecol. Biogeogr.* **11**, 353–361. (doi:10.1046/j.1466-

822x.2002.00300.x)

7.   Randall JE, Cea A. 2011 *Shore Fishes of Easter Island*. University of Hawai'i

Press. See

http://books.google.com/books?hl=en&lr=&id=iXtCTSRTcb8C&pgis=1.

8.   Cea A. 2016 *Ika Rapa Nui*. Rapa Nui Press.

9.   Friedlander AM, Ballesteros E, Beets J, Berkenpas E, Gaymer CF, Gorny M,

Sala E. 2013 Effects of isolation and fishing on the marine ecosystems of Easter

Island and Salas y Gómez, Chile. *Aquat. Conserv. Mar. Freshw. Ecosyst.* **23**,

515–531. (doi:10.1002/aqc.2333)

10.   Clouard V, Bonneville A. 2005 Ages of seamounts, islands, and plateaus on the

Pacific plate. *Spec. Pap. 388 Plates, plumes Paradig.* **388**, 71–90.

(doi:10.1130/0-8137-2388-4.71)

11.   Ray JS, Mahoney JJ, Duncan RA, Ray J, Wessel P, Naar DF. 2012 Chronology

and geochemistry of lavas from the Nazca Ridge and Easter Seamount Chain:

An ~30 myr hotspot record. *J. Petrol.* **53**, 1417–1448.

(doi:10.1093/petrology/egs021)

12.   Newman W, Foster B. 1983 The Rapanuian faunal district (Easter and Sala y

Gomez): in search of ancient archipelagos. *Bull. Mar. Sci.* **33**, 633–644.

13.   Delrieu-Trottin E, Brosseau-Acquaviva L, Mona S, Neglia V, Giles EC, Rapu-

Edmunds C, Saenz-Agudelo P. 2019 Understanding the origin of the most

isolated endemic reef fish fauna of the Indo-Pacific: Coral reef fishes of Rapa

Nui. *J. Biogeogr.* **46**, 723–733. (doi:10.1111/jbi.13531)

14.   Bard E, Hamelin B, Arnold M, Montaggioni L, Cabioch G, Faure G, Rougerie F.

1996 Deglacial sea-level record from Tahiti corals and the timing of global

meltwater discharge. *Nature* **382**, 241–244. (doi:10.1038/382241a0)

15.   Hewitt GM. 2004 The structure of biodiversity -- insights from molecular

phylogeography. *Front. Zool.* **1**, 4. (doi:10.1186/1742-9994-1-4)

16.   Woodroffe CD, Brooke BP, Linklater M, Kennedy DM, Jones BG, Buchanan C,

Mleczko R, Hua Q, Zhao JX. 2010 Response of coral reefs to climate change:

Expansion and demise of the southernmost pacific coral reef. *Geophys. Res.*

Lett. **37**, L15602. (doi:10.1029/2010GL044067)

17.   Delrieu-Trottin E, Mona S, Maynard J, Neglia V, Veuille M, Planes S. 2017

Population expansions dominate demographic histories of endemic and

widespread Pacific reef fishes. *Sci. Rep.* **7**, 40519. (doi:10.1038/srep40519)

18.   Pinheiro HT, Bernardi G, Simon T, Joyeux J-C, Macieira RM, Gasparini JL,

Rocha C, Rocha LA. 2017 Island biogeography of marine organisms. *Nature*

549, 82.

19.   Delrieu-Trottin E, Maynard J, Planes S. 2014 Endemic and widespread coral

reef fishes have similar mitochondrial genetic diversity. *Proc. R. Soc. B Biol. Sci.*

281. (doi:10.1098/rspb.2014.1068)

20.   Peterson BK, Weber JN, Kay EH, Fisher HS, Hoekstra HE. 2012 Double digest

412   RADseq: An inexpensive method for de novo SNP discovery and genotyping in

413   model and non-model species. *PLoS One* **7**.

414   (doi:10.1371/journal.pone.0037135)

415 21. Catchen JM, Amores A, Hohenlohe P, Cresko W, Postlethwait JH. 2011

416   Stacks⬚: Building and Genotyping Loci De Novo From Short-Read Sequences.

417   *G3 Genes|Genomes|Genetics* **1**, 171–182. (doi:10.1534/g3.111.000240)

418 22. Catchen J, Hohenlohe PA, Bassham S, Amores A, Cresko WA. 2013 Stacks: An

419   analysis tool set for population genomics. *Mol. Ecol.* **22**, 3124–3140.

420   (doi:10.1111/mec.12354)

421 23. Mastretta-Yanes A, Arrigo N, Alvarez N, Jorgensen TH, Piñero D, Emerson BC.

422   2015 Restriction site-associated DNA sequencing, genotyping error estimation

423   and de novo assembly optimization for population genetic inference. *Mol. Ecol.*

424   *Resour.* **15**, 28–41. (doi:10.1111/1755-0998.12291)

425 24. R Core Team. 2017 R: A Language and Environment for Statistical Computing.

426   *R Found. Stat. Comput. Vienna, Austria.* **0**, {ISBN} 3-900051-07-0.

427   (doi:http://www.R-project.org/)

428 25. Paradis E. 2010 pegas: an {R} package for population genetics with an

429   integrated--modular approach. *Bioinformatics* **26**, 419–420.

430 26. Liu X, Fu YX. 2015 Exploring population size changes using SNP frequency

431   spectra. *Nat. Genet.* **47**, 555–559. (doi:10.1038/ng.3254)

432 27. Maisano Delser P, Corrigan S, Hale M, Li C, Veuille M, Planes S, Naylor G,

433   Mona S. 2016 Population genomics of C. melanopterus using target gene

434   capture data: demographic inferences and conservation perspectives. *Sci. Rep.*

435   **6**, 33753. (doi:10.1038/srep33753)

28.   Excoffier L, Dupanloup I, Huerta-Sánchez E, Sousa VC, Foll M. 2013 Robust

Demographic Inference from Genomic and SNP Data. *PLoS Genet.* **9**.

(doi:10.1371/journal.pgen.1003905)

29.   Beaumont MA, Zhang W, Balding DJ. 2002 Approximate Bayesian computation

in population genetics. *Genetics*

30.   Jacobs A, Hughes M, Robinson P, Adams C, Elmer K. 2018 The Genetic

Architecture Underlying the Evolution of a Rare Piscivorous Life History Form in

Brown Trout after Secondary Contact and Strong Introgression. *Genes (Basel).*

9, 280. (doi:10.3390/genes9060280)

31.   Le Moan A, Gagnaire PA, Bonhomme F. 2016 Parallel genetic divergence

among coastal-marine ecotype pairs of European anchovy explained by

differential introgression after secondary contact. *Mol. Ecol.*

(doi:10.1111/mec.13627)

32.   Souissi A, Bonhomme F, Manchado M, Bahri-Sfar L, Gagnaire PA. 2018

Genomic and geographic footprints of differential introgression between two

divergent fish species (Solea spp.). *Heredity (Edinb).* (doi:10.1038/s41437-018-

0079-9)

33.   Rougeux C, Bernatchez L, Gagnaire PA. 2017 Modeling the multiple facets of

speciation-with-gene-flow toward inferring the divergence history of lake

whitefish species pairs (Coregonus clupeaformis). *Genome Biol. Evol.* **9**, 2057–

2074. (doi:10.1093/gbe/evx150)

34.   Tine M *et al.* 2014 European sea bass genome and its variation provide insights

into adaptation to euryhalinity and speciation. *Nat. Commun.* **5**, 5770.

(doi:10.1038/ncomms6770)

35.   Oksanen J *et al.* 2012 vegan: Community Ecology Package. R package version

2.1-20/r2309. *www.R-project.org.* , 1–255.

36. Wickham H. 2009 *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag
New York. See http://ggplot2.org.

37. Miller KG *et al.* 2005 The Phanerozoic Record of Global Sea-Level Change.
Science (80-.). **310**, 1293–1298. (doi:10.1126/science.1116412)

38. Augustin L *et al.* 2004 Eight glacial cycles from an Antarctic ice core. *Nature*
(doi:10.1038/nature02599)

39. McManus JF. 2004 A great grand-daddy of ice cores. *Nature*
(doi:10.1038/429611a)

40. Robertson DR. 2004 High genetic diversities and complex genetic structure in an
Indo-Paci c tropical reef sh (Chlorurus sordidus): evidence of an unstable
evolutionary past? *Mar. Biol.* , 757–767. (doi:10.1007/s00227-003-1224-3)

41. Craig MT, Eble JA, Bowen BW, Robertson DR. 2007 High genetic connectivity
across the Indian and Pacific Oceans in the reef fish Myripristis berndti
(Holocentridae). *Mar. Ecol. Prog. Ser.* **334**, 245–254. (doi:10.3354/meps334245)

42. Craig MT, Eble JA, Bowen BW. 2010 Origins, ages and population histories:
Comparative phylogeography of endemic Hawaiian butterflyfishes (genus
Chaetodon). *J. Biogeogr.* **37**, 2125–2136. (doi:10.1111/j.1365-
2699.2010.02358.x)

43. Crandall ED, Frey MA, Grosberg RK, Barber PH. 2008 Contrasting demographic
history and phylogeographical patterns in two Indo-Pacific gastropods. *Mol. Ecol.*
17, 611–626. (doi:10.1111/j.1365-294X.2007.03600.x)

44. Fauvelot C, Bernardi G, Planes S. 2003 Reductions in the mitochondrial DNA
diversity of coral reef fish provide evidence of population bottlenecks resulting

from Holocene sea-level change. *Evolution* **57**, 1571–1583. (doi:10.1111/j.0014-

3820.2003.tb00365.x)

45. Gaither MR, Toonen RJ, Robertson DR, Planes S, Bowen BW. 2010 Genetic

evaluation of marine biogeographical barriers: Perspectives from two

widespread Indo-Pacific snappers (Lutjanus kasmira and Lutjanus fulvus). *J.*

Biogeogr. **37**, 133–147. (doi:10.1111/j.1365-2699.2009.02188.x)

46. Hellberg ME. 2001 Climate-Driven Range Expansion and Morphological

Evolution in a Marine Gastropod. *Science (80-.).* **292**, 1707–1710.

(doi:10.1126/science.1060102)

47. Horne JB, van Herwerden L, Choat JH, Robertson DR. 2008 High population

connectivity across the Indo-Pacific: Congruent lack of phylogeographic

structure in three reef fish congeners. *Mol. Phylogenet. Evol.* **49**, 629–638.

(doi:10.1016/j.ympev.2008.08.023)

48. Klanten OS, Choat JH, Van Herwerden L. 2007 Extreme genetic diversity and

temporal rather than spatial partitioning in a widely distributed coral reef fish.

Mar. Biol. **150**, 659–670. (doi:10.1007/s00227-006-0372-7)

49. Winters KL, van Herwerden L, Choat JH, Robertson DR. 2010 Phylogeography

of the Indo-Pacific parrotfish Scarus psittacus: Isolation generates distinctive

peripheral populations in two oceans. *Mar. Biol.* **157**, 1679–1691.

(doi:10.1007/s00227-010-1442-4)

50. Randall JE. 2005 *Reef and Shore Fishes of the South Pacific : New Caledonia*

to Tahiti and the Pitcairn Islands. University of Hawai'i Press Honolulu.

51. Avise JC, Walker D, Johns GC. 1998 Speciation durations and Pleistocene

effects on vertebrate phylogeography. *Proc. R. Soc. B Biol. Sci.* **265**, 1707–

1712. (doi:10.1098/rspb.1998.0492)

52. Kavembe GD, Kautt AF, Machado-Schiaffino G, Meyer A. 2016 Eco-
morphological differentiation in Lake Magadi tilapia, an extremophile cichlid fish
living in hot, alkaline and hypersaline lakes in East Africa. *Mol. Ecol.* **25**, 1610–
1625. (doi:10.1111/mec.13461)

53. Malinsky M, Svardal H, Tyers AM, Miska EA, Genner MJ, Turner GF, Durbin R.
2018 Whole-genome sequences of Malawi cichlids reveal multiple radiations
interconnected by gene flow. *Nat. Ecol. Evol.* **2**, 1940–1955.
(doi:10.1038/s41559-018-0717-x)

54. Froese R, Binohlan C. 2000 Empirical relationships to estimate asymptotic
length, length at first maturity and length at maximum yield per recruit in fishes,
with a simple method to evaluate length frequency data. *J. Fish Biol.* **56**, 758–
773. (doi:10.1111/j.1095-8649.2000.tb00870.x)

55. Pante E, Simon-Bouhet B. 2013 marmap: A Package for Importing, Plotting and
Analyzing Bathymetric and Topographic Data in R. *PLoS One* **8**, 1–4.
(doi:10.1371/journal.pone.0073051)

56. Bowen BW, Muss A, Rocha LA, Grant WS. 2006 Shallow mtDNA coalescence in
Atlantic pygmy angelfishes (genus Centropyge) indicates a recent invasion from
the Indian Ocean. *J. Hered.* **97**, 1–12. (doi:10.1093/jhered/esj006)

57. Gajdzik L, Bernardi G, Lepoint G, Frédérich B. 2018 Genetic diversity mirrors
trophic ecology in coral reef fish feeding guilds. *Mol. Ecol.* **27**, 5004–5018.
(doi:10.1111/mec.14936)

58. Easton EE, Sellanes J, Berkenpas E, Gaymer CF, Morales N, Gorny M. 2017
Diversity of deep-sea fi shes of the Easter Island Ecoregion. *Deep. Res. Part II*
137, 78–88. (doi:10.1016/j.dsr2.2016.12.006)

Figure captions

Figure 1. Stairway plots for large-range (red) and small-range endemics (black) representing the variation of effective population size through time and sea level fluctuations for the past 400 000 years using data from [37]. Dark tones indicate glacial era characterized by rises of the sea level while greys tones indicate inter-glacial era with lower sea level.

Figure 2. Surface of potential habitat for Rapa Nui endemic reef fish species computed using the package marmap [55] in R [24]. Present conditions (red) and past conditions during the glacial maximums (orange).

Supplementary Material.

Figure S1. Median of effective population sizes (Ne) based on stairway plots and abc skylines analyses.

550 **Table 1 Ecological data on the species of interest of this study. Range size (S: small-range size; L: large-range size), Family,**

551 **Maximum Depth where the species have been observed (Max. depth) and reference associated ([1]: specimens photographed at**

552 **those depths by Luiz A. Rocha; [2]: personal observation from Cristian Rapu-Edmunds freediving); generation time used for this**

553 **study and references associated, maximum size of the species. Codes for reproductive strategy are: [p] pelagic eggs, [b]**

554 **demersal eggs and [m] mouth brooding.**

Range Size	Family	Species	Max. depth (m)	Generation time (years)	Max. size (TL, cm)
S	Monacanthidae [b]	*Cantherhines rapanui*	20 [7]	3	20
S	Apogonidae [m]	*Ostorhinchus chalcius*	25 [7]	2	16
L	Pomacanthidae [p]	*Centropyge hotumatua*	50 [7]	1 [56]	9
S	Pomacentridae [b]	*Chrysiptera rapanui*	72[1]	1 [57]	7.8
S	Labridae [p]	*Coris debueni*	70[1]	3	27
L	Holocentridae [p]	*Myripristis tiki*	80[2]	3 [41]	26.5
S	Pomacentridae [b]	*Chromis randalli*	105[1]	2 [57]	15
S	Chaetodontidae [p]	*Chaetodon litus*	105	2 [42]	15.5
S	Holocentridae [p]	*Sargocentron wilhlmi*	157 [58]	3 [41]	19.5

Species	No. of ind	No. of Sites TOTAL	No. of RADSeq loci	No. of variable RADSeq loci	No. SNP	π accross variable sites	Ho	He	F_{IS}	π accross all sites [1] (10^{-4})
Cantherhines rapanui	15	5 373 281	44 233	20 052	*28 239*	**0.271**	**0.262**	**0.260**	0.023	**32.73**
Centropyge hotumatua	15	3 563 265	29 195	21 684	38 816	*0.163*	0.156	0.158	0.032	25.21
Chaetodon litus	15	5 024 341	41 439	21 905	33 421	0.227	0.220	0.219	0.022	29.86
Chromis randalli	13	4 247 760	34 893	24 673	43 844	0.172	0.161	0.165	**0.043**	26.34
Chrysiptera rapanui	18	*3 104 187*	*25 331*	*19 246*	35 735	0.145	*0.137*	*0.141*	0.039	23.25
Coris debueni	14	5 817 876	47 949	28 594	45 870	0.216	0.208	0.209	0.029	29.94
Myripristis tiki	18	5 522 762	45 432	**28 855**	**46 718**	0.165	0.159	0.160	0.025	23.00
Ostorhinchus chalcius	19	**7 702 975**	**63 590**	24 019	32 224	0.166	0.165	0.162	*0.008*	*19.19*

| *Sargocentron wilhlmi* | 16 | 5 009 870 | 41 234 | 24 207 | 37 860 | 0.224 | 0.217 | 0.217 | 0.026 | 30.98 |