Exhaustive identification of conserved upstream open reading frames with potential translational

regulatory functions from animal genomes

Hiro Takahashi[1,2#]*, Shido Miyaki[2#], Hitoshi Onouchi[3#], Taichiro Motomura[1], Nobuo Idesako[2], Anna Takahashi[4],

Masataka Murase[1], Shuichi Fukuyoshi[5], Toshinori Endo[6], Kenji Satou[7], Satoshi Naito[3,8], and Motoyuki Itoh[9]*

[1]Graduate School of Medical Sciences, Kanazawa University, Kanazawa 920-1192, Japan

[2]Graduate School of Horticulture, Chiba University, Matsudo 271-8510, Japan

[3]Graduate School of Agriculture, Hokkaido University, Sapporo 060-8589, Japan

[4]Faculty of Information Technologies and Control, Belarusian State University of Informatics and Radio

Electronics, Minsk 220013, Belarus

[5]Institute of Medical, Pharmaceutical and Health Sciences, Kanazawa University, Kakuma-machi, Kanazawa,

Ishikawa 920-1192, Japan

[6]Graduate School of Information Science and Technology, Hokkaido University, Sapporo 060-0814, Japan

[7]Faculty of Biological Science and Technology, Institute of Science and Engineering, Kanazawa University,

Kanazawa 920-1192, Japan

[8]Graduate School of Life Science, Hokkaido University, Sapporo 060-0810, Japan

[9]Graduate School of Pharmaceutical Science, Chiba University, Chuo-ku, Chiba 260-8675, Japan

*Correspondence. Tel: +81-76-234-4484; Fax: +81-76-234-4484; Email: takahasi@p.kanazawa-u.ac.jp

Correspondence may also be addressed to Motoyuki Itoh. Email: mito@chiba-u.jp

[#]Joint first authors.

Key words: upstream open reading frame; translational regulation; bioinformatics; nascent peptide

Abstract

Background: Upstream open reading frames (uORFs) are present in the 5′-untranslated regions of many eukaryotic mRNAs, and some peptides encoded in these regions play important regulatory roles in controlling main ORF (mORF) translation. To comprehensively identify uORFs encoding functional peptides, genome-wide searches for uORFs with conserved peptide sequences (CPuORFs) have been conducted in various organisms using comparative genomic approaches. However, in animals, CPuORFs have been identified only by comparing uORF sequences between a limited number of closely related species, and how many previously identified CPuORFs encode regulatory peptides is unclear.

Results: Here, we conducted exhaustive genome-wide searches for animal CPuORFs conserved in various taxonomic ranges, using the ESUCA pipeline, which we recently developed for efficient comprehensive identification of CPuORFs. ESUCA can efficiently compare uORF sequences between an unlimited number of species using BLAST and automatically determine the taxonomic ranges of uORF sequence conservation. By applying ESUCA to human, chicken, zebrafish, and fruit fly genomes, 1,517 (1,425 novel and 92 known) CPuORFs were identified. We examined the effects of 17 human CPuORFs on mORF translation using transient expression assays. Through these analyses, we identified seven novel regulatory CPuORFs that repressed mORF translation in a sequence-dependent manner, including the one conserved only among Eutheria.

Conclusions: We discovered a much higher number of animal CPuORFs than previously identified. Since most human CPuORFs identified in this study are conserved across a wide range of Eutheria or a wider taxonomic range, many CPuORFs encoding regulatory peptides are expected to be found in the identified CPuORFs.

Background

The human genome contains many regions encoding potential functional small peptides outside of the well-annotated protein-coding regions (Ingolia, et al., 2014). Some upstream open reading frames (uORFs), which are located in the 5′-untranslated regions (5′-UTRs) of mRNAs, have been shown to encode such functional small peptides. Most uORF-encoded peptides play regulatory roles in controlling the translation of protein-coding main ORFs (mORFs) (Cruz-Vera, et al., 2011; Ito and Chiba, 2013; Morris and Geballe, 2000; Somers, et al., 2013). During the translation of these regulatory uORFs, nascent peptides interact inside the ribosomal exit tunnel to cause ribosome stalling (Bhushan, et al., 2010). Ribosome stalling on a uORF results in translational repression of the downstream mORF because stalled ribosomes block scanning of subsequent pre-initiation complexes and prevent them from reaching the start codon of the mORF (Wang and Sachs, 1997). In some genes, uORF peptides are involved in translational regulation in response to metabolites (Ito and Chiba, 2013).

To comprehensively identify uORFs encoding functional peptides, genome-wide searches for uORFs with conserved peptide sequences (CPuORFs) have been conducted using comparative genomic approaches in plants (Hayden and Jorgensen, 2007; Takahashi, et al., 2019; Takahashi, et al., 2012; Tran, et al., 2008; van der Horst, et al., 2018; Vaughn, et al., 2012). To date, 157 CPuORF families have been identified by comparing 5′-UTR sequences between plant species. Of these, 101 families were identified in our previous studies by applying our original methods, BAIUCAS (Takahashi, et al., 2012) and ESUCA (an advanced version of BAIUCAS) (Takahashi, et al., 2019) to genomes of *Arabidopsis*, rice, tomato, poplar, and grape.

ESUCA has many unique functions (Takahashi, et al., 2019), such as efficient comparison of uORF sequences between an unlimited number of species using BLAST, automatic determination of taxonomic ranges of CPuORF sequence conservation, systematic calculation of K_a/K_s ratios of CPuORF sequences, and wide compatibility with any eukaryotic genome whose sequence database is registered in ENSEMBL (Zerbino, et al., 2018). More importantly, to distinguish between 'spurious' CPuORFs conserved because they encode parts of mORF-encoded proteins and 'true' CPuORFs conserved because of the functions of their encoded small peptides, ESUCA assesses whether a transcript containing a fusion of a uORF and an mORF is a major or minor form among homologous transcripts (Takahashi, et al., 2019). By using these functions, ESUCA can efficiently

identify CPuORFs likely to encode functional small peptides. In fact, our recent study demonstrated that poplar

CPuORFs encoding regulatory peptides were efficiently identified by selecting ones conserved across diverse

eudicots using ESUCA (Takahashi, et al., 2019).

To date, only a few studies on genome-wide identification of animal CPuORFs have reported. In these

previous studies, uORF sequences were compared between a limited number of closely related species, such as

human and mouse or several species in dipteran, leading to identification of 204 and 198 CPuORFs in human

and mouse, respectively (Crowe, et al., 2006), and 44 CPuORFs in fruit fly (Hayden and Bosco, 2008).

Additionally, the relationships between taxonomic ranges of CPuORF conservation and the likelihood of having

a regulatory function have not been studied in animals.

Accordingly, in this study, we applied ESUCA to genomes of fruit fly, zebrafish, chiken, and human to

exhaustively identify animal CPuORFs and to determine the taxonomic range of their sequence conservation.

Using ESUCA, we identified 1,517 animal (1,425 novel and 92 known) CPuORFs belonging to 1,430 CPuORF

families. We examined the effects of 17 CPuORFs conserved in various taxonomic ranges on mORF translation,

using transient expression assays. Through this analysis, we identified seven novel regulatory CPuORFs that

repress mORF translation in a sequence-dependent manner.

Materials and methods

Extraction of CPuORFs using ESUCA

ESUCA was developed as an advanced version of BAIUCAS (Takahashi et al., 2012) in our previous study

(Takahashi et al., 2019). ESUCA consists of six steps, and some of these steps are divided into substeps, as

shown in Fig. 1A and 1B. To identify animal CPuORFs using ESUCA, the following eight-step procedures were

conducted, including the six ESUCA steps: 0) data preparation for ESUCA, 1) uORF extraction from the 5′-UTR,

2) calculation of uORF-mORF fusion ratios, 3) uORF-tBLASTn against transcript sequence databases, 4)

mORF-tBLASTn against downstream sequence datasets for each uORF, 5) calculation of K_a/K_s ratios, 6)

determination of the taxonomic range of uORF sequence conservation, and 7) manual validation after ESUCA.

See the Supplementary Materials and Methods for details.

Determination of the taxonomic range of uORF sequence conservation for animal CPuORFs

To apply ESUCA to animal genomes, we defined 20 animal taxonomic categories, as shown in Fig. 1C. See the

Supplementary Materials and Methods for details.

Plasmids and reporter assays

DNA fragments containing a CPuORF with the wild-type amino acid sequence or the frameshift mutant version

of the 17 selected genes were subcloned into pSV40:UTR:Fluc. Reporter assays were conducted using a

Dual-Luciferase Reporter Assay system (Promega, Madison, WI, USA). See the Supplementary Materials and

Methods for details.

Results

Genome-wide search for animal CPuORFs using ESUCA

Prior to ESUCA application (Fig. 1A and 1B), we counted the number of protein-coding genes for four species,

i.e., fruit fly, zebrafish, chiken, and human. As shown in Supplementary Table S1, 13,938, 25,206, 14,697, and

19,956 genes were extracted for fruit fly, zebrafish, chiken, and human, respectively. After step 1 of ESUCA, we

calculated the numbers of uORFs and protein-coding genes with any uORF for each species. As shown in

Supplementary Table S1, 17,035 (7,066), 39,616 (14,453), 8,929 (3,535), and 44,085 (12,321) uORFs (genes)

were extracted from fruit fly, zebrafish, chicken, and human genomes, respectively. In this analysis, when

multiple uORFs from a gene shared the same stop or start codon, they were counted as one. Potential candidate

CPuORFs were narrowed down by selection at step 2 of ESUCA in a step-by-step manner, as shown in

Supplementary Table S1. The numbers of BLAST hits (expressed sequence tag [EST], transcriptome shotgun

assembly [TSA], assembled EST/TSA, and RefSeq RNA sequences) extracted at step 3.2 are also shown in

Supplementary Table S1. After the final step of ESUCA, 49, 192, 261, and 1,495 candidate CPuORFs were

extracted from fruit fly, zebrafish, chiken, and human, respectively. We conducted manual validation for the

extracted candidate CPuORFs as described in our previous study (Takahashi, et al., 2019) and in the

Supplementary Materials and Methods. We selected CPuORFs conserved in at least two orders other than the

order to which the original species belongs. In total, 1,517 animal CPuORFs (37 for fruit fly, 156 for zebrafish,

230 for chicken, and 1,094 for human) were identified (Fig. 1D). Of these, 1,425 CPuORFs were newly

identified in the current study. Detailed information on the identified CPuORFs is shown in Supplementary Table

S2. The identified CPuORF-containing genes were classified into 1,363 ortholog groups on the basis of

similarities of mORF-encoded amino acid sequences, using OrthoFinder ver. 1.1.4 (Emms and Kelly, 2015).

CPuORFs with similar amino acid sequences from the same ortholog groups were categorized as the same

CPuORF families (homology groups [HGs]; Supplementary Materials and Methods). The identified 1,517

CPuORFs were classified into 1,430 HGs (Fig. 1D). We assigned HG numbers to 1,430 HGs in an order based

on numbers of orders in which any CPuORF belonging to each HG was extracted, the taxonomic range of the

sequence conservation of each HG, and gene ID numbers. When multiple CPuORF families were identified in

the same ortholog groups, the same HG number with a different subnumber was assigned to each of the families

(e.g., HG0004.1 and HG0004.2; Supplementary Table S2).

Sequence-dependent effects of CPuORFs on mORF translation

To address the relationship between taxonomic ranges of CPuORF conservation and likelihood of having

regulatory function, we selected 17 human CPuORFs conserved in various taxonomic ranges, including a

previously identified sequence-dependent regulatory CPuORF, the *PTP4A1* CPuORF (Hardy, et al., 2019), as a

positive control, and examined their sequence-dependent effects on the expression of the downstream reporter

gene using transient expression assays (Fig. 2). Other uORFs overlapping any of the selected CPuORFs were

eliminated by introducing mutations that changed the ATG codons of the overlapping uORFs to other codons but

did not alter the amino acid sequences of the CPuORFs (Supplementary Fig. S1). The resulting modified

CPuORFs were used as CPuORFs bearing the wild-type amino acid sequences (WT-aa CPuORFs) (Fig. 2B). To

assess the importance of amino acid sequences for the effects of these CPuORFs on mORF translation, frameshift

mutations were introduced into the WT-aa CPuORFs such that the amino acid sequences of their conserved

regions could be altered (see Supplementary Materials and Methods and Supplementary Fig. S1 for details). In

eight of the 17 CPuORFs, the introduced frameshift mutations significantly upregulated the expression of the

reporter gene, indicating that these CPuORFs repressed mORF translation in a sequence-dependent manner (Fig.

2C). One of the eight sequence-dependent regulatory CPuORFs, the *TMEM184C* CPuORF, is conserved only

among Eutheria (Fig. 2A). This result suggests that CPuORFs conserved only among Eutheria can have

seqence-dependent regulatory effects.

158 **Discussion**

In the current study, by applying ESUCA to four animal genomes, we identified 1,517 CPuORFs belonging to

1,430 HGs. Taxonomic ranges of sequence conservation of these CPuORFs largely vary, demonstrating that

ESUCA can identify CPuORFs conserved in various taxonomic ranges (Supplementary Table S3). We examined

the effects of 17 human CPuORFs conserved beyond Euarchontoglires on mORF translation, and identified

seven novel sequence-dependent regulatory CPuORFs (in the *MKKS*, *SLC6A8*, *FAM13B*, *MIEF1*, *KAT6A*,

LRRC8B, and *TMEM184C* genes). Of these, the *TMEM184* CPuORF is one of those conserved in the narrowest

taxonomic range among the tested CPuORFs. This suggests that human CPuORFs conserved beyond

Euarchontoglires are likely to be conserved because of functional constraints of their encoded peptides. Of the

1,094 CPuORFs extracted from the human genome, 1,082 are conserved beyond Euarchontoglires

(Supplementary Table S3). Therefore, many CPuORFs encoding regulatory peptides are expected to be found in

the human CPuORFs identified in this study.

Of the sequence-dependent regulatory CPuORFs identified here, the *MKKS* CPuORF has been

previously reported to be a translational regulator that represses the production of a protein involved in

McKusick-Kaufman syndrome (Akimoto, et al., 2013); however, the amino acid sequence dependence of the

CPuORF function was not reported. Interestingly, the *MIEF1* CPuORF-encoded peptide is a functional peptide

localized in the mitochondria (Samandi, et al., 2017). Thus, the *MIEF1* CPuORF may have dual functions.

Chemical screening recently identified a compound that causes nascent peptide-mediated ribosome

stalling in the mORF of the human *PCSK9* gene, resulting in specific translational inhibition of *PCSK9* and a

reduction in total plasma cholesterol levels (Lintner, et al., 2017). Nascent peptide-mediated ribosome stalling in

some of the previously identified regulatory CPuORFs is promoted by metabolites, such as polyamine, arginine,

and sucrose (Ito and Chiba, 2013; Yamashita, et al., 2017). Therefore, compounds that promote nascent

peptide-mediated ribosome stalling in CPuORFs could be identified by chemical screening through a method

similar to that used for the screening of the stall-inducing compound for *PCSK9*. The data from the current study

may be useful for selection of CPuORFs as potential targets for pharmaceutical drugs and for identification of

regulatory CPuORFs.

Funding: This work was supported by the Japan Society for the Promotion of Science (JSPS) KAKENHI (grant

nos. JP16K07387 to H.O., JP19K22892 to H.T., JP18H03330 to H.T. and H.O., JP18H02568 to M.I.); the

Ministry of Education, Culture, Sports, Science and Technology (MEXT) KAKENHI (grant nos. JP17H05658

to S.N., JP26114703 to H.T, JP17H05659 to H.T); and the Naito Foundation (to H.O.).

Competing interests: none declared.

References

Akimoto, C*., et al.* (2013) Translational repression of the McKusick-Kaufman syndrome transcript by unique upstream open reading frames encoding mitochondrial proteins with alternative polyadenylation sites, *Biochim Biophys Acta*, **1830**, 2728-2738.

Bhushan, S*., et al.* (2010) Structural basis for translational stalling by human cytomegalovirus and fungal arginine attenuator peptide, *Mol. Cell*, **40**, 138-146.

Crowe, M.L., Wang, X.Q. and Rothnagel, J.A. (2006) Evidence for conservation and selection of upstream open reading frames suggests probable encoding of bioactive peptides, *BMC Genomics*, **7**, 16.

Cruz-Vera, L.R*., et al.* (2011) Nascent polypeptide sequences that influence ribosome function, *Curr. Opin. Microbiol.*, **14**, 160-166.

Emms, D.M. and Kelly, S. (2015) OrthoFinder: solving fundamental biases in whole genome comparisons dramatically improves orthogroup inference accuracy, *Genome Biol.*, **16**, 157.

Hardy, S*., et al.* (2019) Magnesium-sensitive upstream ORF controls PRL phosphatase expression to mediate energy metabolism, *Proc Natl Acad Sci U S A*, **116**, 2925-2934.

Hayden, C.A. and Bosco, G. (2008) Comparative genomic analysis of novel conserved peptide upstream open reading frames in *Drosophila melanogaster* and other dipteran species, *BMC Genomics*, **9**, 61.

Hayden, C.A. and Jorgensen, R.A. (2007) Identification of novel conserved peptide uORF homology groups in *Arabidopsis* and rice reveals ancient eukaryotic origin of select groups and preferential association with transcription factor-encoding genes, *BMC Biol.*, **5**, 32.

Ingolia, N.T*., et al.* (2014) Ribosome profiling reveals pervasive translation outside of annotated protein-coding genes, *Cell Rep.*, **8**, 1365-1379.

Ito, K. and Chiba, S. (2013) Arrest peptides: cis-acting modulators of translation, *Annu. Rev. Biochem.*, **82**, 171-202.

Lintner, N.G*., et al.* (2017) Selective stalling of human translation through small-molecule engagement of the

ribosome nascent chain, *PLoS Biol*, **15**, e2001882.

Morris, D.R. and Geballe, A.P. (2000) Upstream open reading frames as regulators of mRNA translation, *Mol.*

Cell Biol., **20**, 8635-8642.

Samandi, S., *et al.* (2017) Deep transcriptome annotation enables the discovery and functional characterization of

cryptic small proteins, *Elife*, **6**.

Somers, J., Poyry, T. and Willis, A.E. (2013) A perspective on mammalian upstream open reading frame

function, *Int. J. Biochem. Cell Biol.*, **45**, 1690-1700.

Takahashi, H., *et al.* (2019) Comprehensive genome-wide identification of angiosperm upstream ORFs with

peptide sequences conserved in various taxonomic ranges using a novel pipeline, ESUCA, *bioRxiv*, 524090.

Takahashi, H., *et al.* (2012) BAIUCAS: a novel BLAST-based algorithm for the identification of upstream open

reading frames with conserved amino acid sequences and its application to the *Arabidopsis thaliana* genome,

Bioinformatics, **28**, 2231-2241.

Tran, M.K., Schultz, C.J. and Baumann, U. (2008) Conserved upstream open reading frames in higher plants,

BMC Genomics, **9**, 361.

van der Horst, S., *et al.* (2018) Novel pipeline identifies new upstream ORFs and non-AUG initiating main ORFs

with conserved amino acid sequences in the 5' leader of mRNAs in *Arabidopsis thaliana*, *RNA*, **25**,

292-304.

Vaughn, J.N., *et al.* (2012) Known and novel post-transcriptional regulatory sequences are conserved across plant

families, *RNA*, **18**, 368-384.

Wang, Z. and Sachs, M.S. (1997) Ribosome stalling is responsible for arginine-specific translational attenuation

in *Neurospora crassa*, *Mol. Cell Biol.*, **17**, 4904-4913.

Yamashita, Y., *et al.* (2017) Sucrose sensing through nascent peptide-meditated ribosome stalling at the stop

codon of Arabidopsis *bZIP11* uORF2, *FEBS Lett.*, **591**, 1266-1277.

Zerbino, D.R., *et al.* (2018) Ensembl 2018, *Nucleic Acids Res.*, **46**, D754-D761.

Figure Legends

Figure 1. Identification of animal CPuORFs using ESUCA. (A) Data preparation. (B) Outline of the ESUCA pipeline. Numbers with parenthesis indicate datasets labeled with the same numbers in A. (C) Defined animal taxonomic categories. (D) Numbers of identified CPuORFs.

Figure 2. Taxonomic conservation and experimental validation of 17 selected human CPuORFs. (A) Taxonomic ranges of conservation of CPuORFs examined in transient assays. Filled cells in each taxonomic category indicate the presence of uORF-tBLASTn and mORF-tBLASTn hits for CPuORFs of the indicated genes. (B) Reporter constructs used for transient assays. The hatched box in the frameshift (fs) mutant CPuORF indicates the frame-shifted region. Dotted boxes represent the first five nucleotides of the mORFs associated with the 17 human CPuORFs. (C) Relative luciferase activities of WT-aa (white) or frameshift (gray) CPuORF reporter plasmids. Means ± SDs of three biological replicates are shown. $*p < 0.05$.

Figure 1

D

Species	CPuORF
D. melanogaster	37
D. rerio	156
G. gallus	230
H. sapiens	1094

Number of animal CPuORFs: 1517
Number of animal HGs: 1430

Figure 2