Generation of comprehensive ecosystems-specific reference databases with species-level resolution by high-throughput full-length 16S rRNA gene sequencing and automated taxonomy assignment (AutoTax)

Authors: Morten Simonsen Dueholm*[§], Kasper Skytte Andersen[§], Simon Jon McIlroy**, Jannie Munk Kristensen, Erika Yashiro, Søren Michael Karst, Mads Albertsen, Per Halkjær Nielsen*

Affiliation:

Center for Microbial Communities, Department of Chemistry and Bioscience, Aalborg University, Aalborg, Denmark.

*Correspondence to: Morten Simonsen Dueholm and Per Halkjær Nielsen, Center for Microbial Communities, Department of Chemistry and Bioscience, Aalborg University, Fredrik Bajers Vej 7H, 9220 Aalborg, Denmark; Phone: (+45) 9940 8503; Fax: Not available; E-mail: md@bio.aau.dk , phn@bio.aau.dk

Present address: Centre for Microbiome Research, School of Biomedical Sciences, Queensland University of Technology (QUT), Translational Research Institute, Woolloongabba, Qld. 4072, Australia

§ contributed equally to the work

Running title:

Comprehensive ecosystems-specific 16S rRNA gene databases

Abstract

High-throughput 16S rRNA gene amplicon sequencing is an essential method for studying the diversity and dynamics of microbial communities. However, this method is presently hampered by the lack of high-identity reference sequences for many environmental microbes in the public 16S rRNA gene reference databases, and by the absence of a systematic and comprehensive taxonomy for the uncultured majority. Here we demonstrate how high-throughput synthetic long-read sequencing can be applied to create ecosystem-specific full-length 16S rRNA gene amplicon sequence variant (FL-ASV) reference databases that include high-identity references (>98.7% identity) for nearly all abundant bacteria (>0.01% relative abundance) using Danish wastewater treatment systems and anaerobic digesters as an example. In addition, we introduce a novel sequence identity-based approach for automated taxonomy assignment (AutoTax) that provides a complete seven-rank taxonomy for all reference sequences, using the SILVA taxonomy as a backbone, with stable placeholder names for unclassified taxa. The FL-ASVs are perfectly suited for the evaluation of taxonomic resolution and bias associated with primers commonly used for amplicon sequencing, allowing researchers to choose those that are ideal for their ecosystem. The AutoTax taxonomy greatly improves the classification of short-read 16S rRNA gene amplicon sequence variants (ASVs) at the genus- and species-level, compared to the commonly used universal reference databases. Importantly, the placeholder names provide a way to explore the unclassified environmental taxa at different taxonomic ranks, which in combination with *in situ* analyses can be used to uncover their ecological roles.

Introduction:

Microbial communities underpin key biochemical transformations in natural and engineered ecosystems. A deep understanding of these systems requires reliable identification of the microbes present, which can then be associated with their metabolic and ecological functions. Identification at the lowest taxonomic rank is preferred, as microbial traits vary in their degree of phylogenetic conservation, and many ecologically important traits are conserved only at the genus and species level (1).

Identification of microbes is commonly achieved by high-throughput 16S rRNA gene amplicon sequencing, where a segment of the 16S rRNA gene spanning one to three hypervariable regions (usually 200-500 bp long) is amplified by PCR and sequenced. The amplicons are then clustered into operational taxonomic units (OTUs) or used to infer exact amplicon sequence variants (ASVs) with denoising algorithms such as Deblur (2), DADA2 (3), or Unoise3 (4). The taxonomy of these amplicons is then assigned based on their comparison to a reference database (5–7).

ASVs are often preferred over OTUs because they provide single-nucleotide resolution and can be applied as consistent labels for microbial identification independently of a 16S rRNA gene reference database (8). This approach is used with short-read ASVs in several large-scale projects, including the Earth Microbiome Project (EMP) (9) and the American Gut project (10), to provide detailed insight into the factors that shape the overall microbial community diversity and dynamics. However, ASVs are not ideal as references for linking microbial identity with the physiology and ecology of key community members, which is crucial if we want to use microbial community structure to predict ecosystem functions or process performance in engineered systems. Firstly, without taxonomic assignment, it is not possible to compare results across studies that have used primers targeting different regions of the 16S rRNA gene. Secondly, short-read ASVs alone do not contain enough evolutionary information to resolve their phylogeny confidently (11, 12). This makes it impossible to report and infer how microbial traits are conserved at different phylogenetic scales. As such, functional properties for uncultured lineages predicted from the annotation of metagenome-assembled genomes (MAGs) with complete rRNA genes (Hiqh-quality MIMAGs standard) (11), or determined through *in situ* studies, cannot be confidently linked to the shorter ASV sequences (12, 13). A robust taxonomic assignment is therefore crucial for cross-study comparisons and the accumulation and dissemination of knowledge relating to uncultured lineages.

Taxonomic assignment to ASVs relies on a classifier, e.g., the SINTAX (7) or the RDP

classifier (6), which uses statistical algorithms to compare each ASV to a full-length 16S rRNA

gene reference database to propose the best estimate of their taxonomy. Confident classification at

the lowest taxonomic ranks (genus and species) requires high-identity reference sequences (≥98.7%

identity) and a complete seven-rank taxonomy for all references (13). Neither of these criteria is met

with the most commonly applied universal reference databases, e.g., Greengenes (14), SILVA (15),

and RDP (16), as they lack sequences or taxonomic assignment for many uncultivated

environmental taxa. Given the vast diversity predicted for microbial life on Earth (17), it will be

some time before reference sequences for all species are generated, and the manual curation of their

taxonomy will not be feasible.

A potential solution to the problems mentioned above is to create ecosystem-specific

reference databases. These can either be ecosystem-curated versions of universal databases or

smaller independent databases that only include sequences from the specific ecosystem. The

MiDAS 2 database for microbes in biological wastewater treatment systems (18) and the

Dictyopteran gut microbiota reference Database (DictDb) (19) are examples of ecosystem-curated

versions of the SILVA databases, where the taxonomies for the abundant and process-critical

microorganisms are manually curated and maintained. However, the mostly manual taxonomic

curation of central reference databases is time-consuming and subjective. The universal reference

databases are also clustered at 99% identity or below to keep the unique sequences to a manageable

number, which reduces the taxonomic resolution.

Examples of independent ecosystem-specific databases include the human intestinal tract 16S

taxonomic database (HITdb) (20), the human oral microbiome database (HOMD) (21), the

freshwater-specific FreshTrain database (22, 23), the honey bee gut microbiota database (24), and

the rumen and intestinal methanogen database (25). While such databases have been shown to

improve the rate of classifications for amplicons, they generally contain a relatively limited number

of sequences and are therefore associated with an inherent risk of over- or misclassification if the

sequence being classified is not represented in the database. To address this issue, Rohwer *et al*.

introduced the TaxAss algorithm that classifies amplicons using two reference databases: a

universal database and a small ecosystem-specific database (23). Amplicons are first mapped to the

ecosystem-specific database to determine the percent identity with the best hit, and those above a

user-defined threshold are classified using the ecosystem-specific database. The remaining

sequences are classified using the more extensive universal database. While higher rates of

classification were achieved, an issue with the approach is the potential for closely related sequences that fall on either side of the user-defined threshold, to receive very different taxonomies, especially if the ecosystem-specific database is not updated to reflect the evolving taxonomy of the universal reference database. As such, while current strategies to create ecosystem-specific databases have shown promise, there are critical issues that need to be resolved before their potential can be realized.

The recent development of methods for high-throughput full-length 16S rRNA gene sequencing, e.g., synthetic long-read sequencing on the Illumina platform (26, 27), along with PacBio (28) or Nanopore (29) consensus sequencing, now allows for the generation of millions of high-quality reference sequences within days. Importantly, these technologies now allow for the high-throughput generation of high-identity reference databases with broad coverage of the true diversity. However, improving sequence coverage alone will not solve the problem of poor taxonomic assignments for many uncultured taxa. We have therefore developed the AutoTax pipeline, which provides a simple and efficient strategy for the creation of comprehensive ecosystem-specific taxonomies that cover all seven taxonomic ranks. AutoTax uses the SILVA taxonomy as a backbone and provides stable placeholder names for unclassified taxa, based on *de novo* clustering of sequences according to statistically supported identity thresholds for each taxonomic rank (12). Importantly, AutoTax databases are easily updated with subsequent releases of the SILVA taxonomy – avoiding the divergence of generated ecosystem-specific taxonomies with the universal reference database. The strict computational nature of the taxonomy assignment means that it is objective and reproducible. The simplicity of the applied *de novo* clustering also ensures that the placeholder names are maintained even though the database is expanded with additional reference sequences.

We demonstrate the potential of the AutoTax method by sequencing almost a million full-length 16S rRNA gene sequences from Danish biological wastewater treatment and bioenergy systems. The sequences were denoised to resolve full-length exact amplicon sequence variants (FL-ASVs) with single-nucleotide resolution. Taxonomy was assigned to the FL-ASVs using the AutoTax pipeline to create an ecosystem-specific reference database. As evidence supporting the value of our approach, mapping of short-read amplicon data revealed that a substantially higher proportion of sequences were matched to high-identity references, and received species and genus level classification when the FL-ASV database was used compared to the much larger public universal reference databases.

Materials and methods:

General molecular methods

Concentration and quality of nucleic acids were determined using a Qubit 3.0 fluorometer (Thermo Fisher Scientific) and an Agilent 2200 Tapestation (Agilent Technologies), respectively. Agencourt RNAClean XP and AMPure XP beads were used as described by the manufacturer, except for the washing steps, where 80% ethanol was used. RiboLock RNase inhibitor (Thermo Fisher Scientific) was added to the purified RNA to minimize degradation. All commercial kits were used according to the protocols provided by the manufacturer unless otherwise stated. Oligonucleotides used in this study can be found in **Table S1**.

Samples and nucleic acid purification

Activated sludge and anaerobic digester biomass were obtained as frozen aliquots (-80°C) from the MiDAS collection (18). Sample metadata is provided in **Table S2**. Total nucleic acids were purified from 500 µL of sample thawed on ice using the PowerMicrobiome RNA isolation kit (MO BIO Laboratories) with the optional phenol-based lysis or with the RiboPure RNA purification kit for bacteria (Thermo Fisher Scientific). Purification was carried out according to the manufacturers' recommendations, except that cell lysis was performed in a FastPrep-24 instrument for 4x 40 s at 6.0 m/s to increase the yield of nucleic acids from bacteria with sturdy cell walls (30). The samples were incubated on ice for 2 min between each bead beating to minimize heating due to friction. DNA-free total RNA was obtained by treating 41 µL of the purified nucleic acid with the DNase Max kit (MO BIO Laboratories), followed by clean up using 1.0x RNAClean XP beads with elution into 25 µL nuclease-free water.

Primer-free full-length 16S rRNA library preparation and sequencing

Purified RNA obtained from biomass samples was pooled separately for each sample source type (activated sludge or anaerobic digester) to give equimolar amounts of 16S rRNA determined based on peak area in the TapeStation analysis software A.02.02 (SR1). Full-length SSU sequencing libraries were then prepared, as previously described (26). The SSU_rRNA_RT2 (activated sludge) and SSU_rRNA_RT3 (anaerobic digester) reverse transcription primers and the SSU_rRNA_l adaptor were used for the molecular tagging (**Table S1**), and approximately 1,000,000 tagged molecules from each pooled sample were used to create the clonal library. The final library was sequenced on a HiSeq2500 using on-board clustering and rapid run mode with a HiSeq PE Rapid

Cluster Kit v2 (Illumina) and HiSeq Rapid SBS Kit v2, 265 cycles (Illumina), as previously

described (26). Raw sequence reads were binned based on unique molecular tags, *de novo*

assembled into synthetic long-read sequences, and trimmed equivalent to *E. coli* position 8 and

1507 using the fSSU-pipeline-RNA_v1.2.sh scripts script

(https://github.com/KasperSkytte/AutoTax) (26).

Primer-based full-length 16S rRNA gene library preparation and sequencing

The purified nucleic acids obtained from the biomass samples were pooled separately for each

sample source type (activated sludge or anaerobic digester) with an equal weight of DNA from each

sample. Full-length 16S rRNA sequencing libraries were then prepared, as previously described

(26). The f16S_rDNA_pcr1_fw1 (activated sludge) or f16S_rDNA_pcr1_fw2 (anaerobic digester)

and the f16S_rDNA_pcr1_rv (**Table S1**) were used for the molecular tagging, and approximately

1,000,000 tagged molecules from each pooled sample were used to create the clonal library. The

final library was sequenced on a HiSeq2500 using on-board clustering and rapid run mode with a

HiSeq PE Rapid Cluster Kit v2 (Illumina) and HiSeq Rapid SBS Kit v2, 265 cycles (Illumina) as

previously described (26). Raw sequence reads were binned based on unique molecular tags, *de*

novo assembled into synthetic long-read sequences, and trimmed equivalent to *E. coli* position 28

and 1491 using the fSSU-pipeline-DNA_v1.2.sh scripts script

(https://github.com/KasperSkytte/AutoTax) (26).

Extraction of high-quality "full-length" 16S rRNA gene sequences from SILVA

High-quality bacterial and archaeal 16S rRNA gene sequences in the SILVA 138 SSURef NR99

ARB-database was selected using the query pintail_slv = 100 and tax_slv = Bacteria* or tax_slv =

Archaea*. Bacterial and Archaeal sequences were exported separately in the "fastawide" format

after terminal trimming. Bacterial sequences were trimmed between the 27F and 1391R (31) primer

binding sites equivalent to position 1,044 and 41,788 in the global SILVA alignment. Archaeal

sequences were trimmed between the 20F (32) and the SSU1000ArR (33) primer binding sites

equivalent to position 1,041 and 32,818 in the global SILVA alignment. A list of names for "full-

length" sequences spanning the positions above was created using the Extract_full-

length_16S_rRNA_names_from_SILVA.sh script, which takes advantage of the fact that ARB uses

the period to specify terminal gaps and therefore indicates truncated sequences in the exported

FASTA-files. The names were used to select and export the "full-length" bacterial or archaea

sequences without trimming from the SILVA ARB-database.

Generation of reference databases using AutoTax

AutoTax was created as a modular multi-step Linux BASH script that 1) generates FL-ASV and

full-length 16S rRNA gene operational taxonomic units clustered at 99% identity (FL-OTU)

reference sequences from high-quality, full-length 16S rRNA sequences. 2) Assigns a

comprehensive seven-rank taxonomy to all reference sequences based on the SILVA taxonomy

with the addition of placeholder names for unclassified taxa defined by *de novo* clustering of

sequences using specific identity thresholds for each taxonomic rank (12). 3) Produces formatted

reference databases, which can be directly used for classification using SINTAX or classifiers in the

QIIME 2 framework.

AutoTax combines several software tools (GNU parallel v. 20161222 (34), USEARCH v.

11.0.667 (35), SINA v.1.6.0 (36), and R v.3.5.0 with the following packages: biostrings (37),

doParallel (38), stringr (39), data.table (40), tidyr (41), and dplyr (42)) into a single BASH script

that otherwise only requires a single FASTA file with the user provided full-length 16S rRNA gene

sequences and the FASTA-formatted SILVA_138_SSURef_NR99_tax_silva reference database as

input. The script, as well as a docker container image with all required software (except

USEARCH, as the required 64 bit version it is non-free and must be purchased from

https://drive5.com/usearch), is available on the GitHub repository

https://github.com/KasperSkytte/AutoTax. The script is composed of separate, individual BASH

functions to both allow for customization of the script as well as unit testing using the BASH

Automated Testing System (https://github.com/bats-core/bats-core) where possible. Core functions

of AutoTax are briefly described below. Expanded descriptions can be found in the supplementary

information (**Text S1**).

Resolving full-length 16S rRNA amplicon sequence variant (FL-ASV)

Input sequences are oriented based on the SILVA 138 SSURef NR99 database using the *usearch -*

orient command, dereplicated using *usearch -fastx_uniques* with the *-sizeout, -strand plus*, and *-*

threads 1 options, and finally denoised to produce the FL-ASVs using the *usearch -unoise3*

command with the *-minsize 2* option.

Preparation of chimera-filtered full-length 16S rRNA OTUs (FL-OTUs)

Dereplicated sequences (before denoising) from above are clustered at 99% sequence identity using

usearch -cluster_smallmem command with the *-id 0.99*, *-maxrejects 0, -centroids*, and *-sortedby*

size options. Potential chimeras are identified and extracted using the *usearch -uchime2_ref*

command with the *-strand plus, -mode sensitive*, and *-chimeras* options with the FL-ASVs from

above as the reference database. The chimeras are finally removed to create the final FL-OTUs

using the *usearch -search_exact* command with the *-strand plus* and *-dbnotmatched* options.

Taxonomy assignment

A complete taxonomy from kingdom to species is automatically assigned to each FL-ASV. In brief,

the AutoTax script identifies the closest relative of each FL-ASV in the SILVA database using the

usearch -usearch_global command, obtains the taxonomy for this sequence, and discards

information at taxonomic ranks not supported by the sequence identity and the thresholds for

taxonomic ranks proposed by Yarza *et al*. (12). The identity thresholds used for each of the

taxonomic ranks are: Phylum (75.0%), Class (78.5%), Order (82.0%), Family (86.5%), Genus

(94.5%), and Species (98.7%). For the species-level classification, the script identifies all type

strains within the species-level threshold in the SILVA database and assigns a species-level

classification to the FL-ASV if only a single species fits within the threshold. In addition, FL-ASVs

are *de novo* clustered using the *usearch -cluster_smallmem* command using the thresholds for each

taxonomic rank. The *de novo* clusters are labeled according to the format denovo_x_y, where x is a

one-letter abbreviation for the taxonomic rank (k, p, c, o, f, g, and s), and y represents the FL-ASV

number of the cluster centroid for each taxon. These labels act as a placeholder taxonomy, where

the SILVA taxonomy does not provide any taxonomy information.

Amplicon sequencing and analyses

Bacterial community analyses were performed by amplicon sequencing of the V1-3 variable region

of the 16S rRNA gene as previously described (43) using the 27F (5'-

AGAGTTTGATCCTGGCTCAG-3') (31) and 534R (5'-ATTACCGCGGCTGCTGG-3') (44)

primers. Forward reads were processed using USEARCH v.11.0.667. Raw fastq files were filtered

for phiX sequences using *usearch -filter_phix*, trimmed to 250 bp using *usearch -fastx_truncate -*

trunclen 250, and quality filtered using *usearch -fastq_filter* with the *-fastq_maxee 1.0* option. The

sequences were dereplicated using *usearch -fastx_uniques* with the *-sizeout* option. Exact amplicon

sequence variants (ASVs) were resolved using *usearch -unoise3* (4). ASV-tables were created by

mapping the raw reads to the ASVs using *usearch -otutab* with the *-zotus* and *-strand both* options.

Taxonomy was assigned to ASVs using *usearch -sintax* with *-strand both* and *-sintax_cutoff 0.8*

(13).

Construction of phylogenetic trees and primer evaluation

FL-ASVs aligned to SILVA 138 NR99 ARB-database was obtained from the AutoTax output

(temp/FL-ASVs_SILVA_aligned.fa) and loaded into SILVA 138 NR99 ARB-database. All

bacterial FL-ASVs were selected and exported as a FASTA-file using the ssuref:bacteria positional

variability by parsimony filter. A rough tree was created from the alignment using FastTree v.

2.1.10 (45) with the *-nt -gtr -gamma* options and loaded into ARB. The specificity of commonly

used amplicon primers was determined for each FL-ASV using the *analyze_primers.py* script from

Primer Prospector v. 1.0.1 (46). The specificity of primer sets was defined based on the overall

weighted scores (OWS) for the primer with the highest score as follows: Perfect hit (OWS = 0),

partial hit (OWS > 0, and ≤ 1), poor hit (OWS > 1). A comma-separated table with the specificity of

each primer set for each FL-ASV was made in R, loaded into ARB, and used to color the tree.

Data analyses and visualization

USEARCH v.11.0.667 was used for mapping sequences to references with *-usearch_global -id 0 -*

maxrejects 0 -maxaccepts 0 -top_hit_only -strand plus, unless otherwise stated. Data were imported

into R v.3.6.3 (47) using RStudio IDE (48), aggregated using the tidyverse package v.1.2.1

(https://www.tidyverse.org/), and analyzed and visualized using ggplot2 v.3.1.0 (49) and Ampvis2

v.2.4.0 (50).

Data and code availability

Raw and assembled sequencing data is available at the European Nucleotide Archive

(https://www.ebi.ac.uk/ena) under the project number PRJEB26558. The AutoTax script is

available at https://github.com/KasperSkytte/AutoTax. AutoTax processed FL-ASV and FL-OTU

expanded reference database in SINTAX and QIIME format, is available at

https://figshare.com/articles/Data_used_in_AutoTax_paper/12377741/1. The AutoTax processed

SILVA 138 SSURef NR99 database in SINTAX, and QIIME format is available at

https://doi.org/10.6084/m9.figshare.12366626. R-markdown scripts used for data analyses and

figures are available at https://github.com/msdueholm/Publications/tree/master/Dueholm2019a.

Results and Discussion:

Sampling and high-throughput sequencing of full-length 16S rRNA sequences

To obtain 16S rRNA gene reference sequences for Danish wastewater treatment plants (WWTPs) and anaerobic digesters (ADs), we sampled biomass from 22 typical WWTPs and 16 ADs treating waste activated sludge located at Danish wastewater treatment facilities (**Table S2**). These facilities represent an important engineered ecosystem containing complex microbial communities of both bacteria and archaea, with the vast majority of microbes being uncultured and poorly characterized (51).

DNA and RNA were extracted and used for synthetic long-read 16S rRNA gene sequencing using both a primer-based and primer-free approach (26) (**Figure 1a**). A total of 926,507 full-length 16S rRNA gene sequences were obtained after quality filtering. These were denoised with UNOISE3 to generate a comprehensive reference database of 9,521 FL-ASVs with an error-rate below the detection limit according to theoretical calculations and analysis of 7,816 sequences previously obtained from the eight strain ZymoBIOMICS Microbial Community DNA Standard (26) (**Text S1**).

To estimate the number of FL-ASVs belonging to novel taxa, FL-ASVs were mapped to the SILVA 138 SSURef NR99 database (15) using global mapping with USEARCH, and the identity of their closest relatives was compared to the thresholds for taxonomic ranks proposed by Yarza *et al.* 2014) (**Table 1**). The majority of the FL-ASVs (94%) had references in the SILVA database above the genus-level threshold (identity >94.5%), but 26% lacked references above the species-level threshold (identity > 98.7%), which are crucial for confident taxonomy assignment to ASVs.

The FL-ASVs have better ecosystem coverage than universal reference databases

To evaluate if the FL-ASV database contained high-identity references for all bacteria in the ecosystem, we mapped V1-3 ASVs obtained from two sources: (1) the same samples used to create the FL-ASVs, and (2) samples from unrelated Danish WWTP and ADs. The ecosystem-specific FL-ASV database (9,521 seq.) included more high-identity references (>98.7% identity) for the abundant ASVs (relative abundance cutoff at 0.01%) in all samples analyzed, compared to the much larger universal databases: MiDAS 2.1 (548,447 seq.) (18), SILVA 138 SSURef NR99 (510,984 seq.) (15), SILVA 138 SSURef (2,225,272 seq.) (15), GreenGenes 16S v.13.5 (1,262,986 seq.) (14), and the full RDP v.11.5 (3,356,808 seq.) (16) (**Figure 1b and Figure S1**). ASVs were also mapped to the 16S rRNA database derived from the Genome Taxonomy Database (GTDB) release 89

(17,460 seq.) (52). However, this database lacked high-identity references for almost all ASVs. The
poor coverage likely relates to the fact that 16S rRNA genes often fail to assemble in MAGs
produced by short-read sequencing data (53). This problem will likely disappear in the future with
the introduction of more high-quality MAGs with complete rRNA genes into the GTDB as a result
of long-read sequencing technologies such as Nanopore and PacBio (54, 55).

When the rare biosphere was included in the analysis (relative abundance cutoff at 0.001%),
a decrease in the percentage of ASVs with high-identity reference sequences was observed (**Figure**
1b). This may be a problem in ecosystems with high diversity, such as soil and sediments (9),
where low-abundant microbes constitute a considerable fraction of the community, but also in
engineered systems where transient or low abundant bacteria, such as pathogens or bacteria
degrading micropollutants, may be important. To get a better representation of the rare biosphere,
we created an additional reference database, which besides the FL-ASVs, contained the chimera-
filtered, full-length 16S rRNA gene sequences clustered at 99% identity (FL-OTUs). This database
greatly increased the coverage for the rare biosphere because it includes FL-OTUs that represent
singletons. However, the improved coverage is achieved at the expense of taxonomic resolution
(see later).

Since only Danish WWTPs and ADs were used to establish the comprehensive high-identity
FL-ASV reference database, published amplicon data from non-Danish WWTPs (56, 57) was also
evaluated (**Figure 1c-d, and S2**). Compared to the analyzed universal reference databases, the
Danish reference FL-ASVs performed better or as well for most of the investigated non-Danish
WWTPs, which indicates that even with less than 10,000 sequences, the database covers many of
the microbes that are common to WWTP across the world, particularly for systems with nutrient
removal. We anticipate that our ongoing sampling of more than 1,000 WWTPs and AD systems
across all seven continents and different process-configurations (MiDAS Global,
https://www.midasfieldguide.org/global) for high-throughput full-length 16S rRNA gene
sequencing will provide references for the region-specific taxa in the near future, providing a
comprehensive database of reference sequences for this ecosystem.

A new comprehensive taxonomic framework
A major limitation in the classification of amplicon data from environmental samples is the lack of
genus and species names for many uncultivated bacteria in the universal reference databases. To
address this, we developed a simple taxonomic framework (AutoTax), which provides a consistent

taxonomic classification of all reference sequences to all seven taxonomic ranks using identity thresholds (**Figure 2**).

In the AutoTax pipeline, the FL-ASVs (or FL-OTUs) are first mapped to the SILVA 138 SSURef NR99 database, which provides the taxonomy and percent identity of the closest relative in the database. This taxonomy is assigned to the FL-ASV down to the taxonomic rank supported by the sequence identity thresholds proposed by Yarza *et al.* (2014) (**Table 1**). Because of the stringent mapping and the taxonomy trimming based on percent identity, we obtained an overall better taxonomy assignment compared to commonly used classifiers, as revealed by a leave-one-out classification test (**Figure S3**).

Since species-level classification is desired wherever possible and the official SILVA taxonomy for bacteria and archaea is not curated at the species level (58), FL-ASVs were also mapped to the 16S rRNA gene sequences from type strains extracted from the SILVA 138 SSURef NR99 database as these carry official species names. Species-level classifications were assigned to the FL-ASVs if they shared more than 98.7% identity with only one species. If the FL-ASVs matched more than one, they were not classified at the species level due to the high risk of misclassification.

Because the SILVA taxonomy does not provide a complete seven-rank taxonomy for all sequences, the missing classifications are covered with a *de novo* placeholder taxonomy. This taxonomy was created based on the clustering of the FL-ASVs at identity thresholds corresponding to each taxonomic rank (12). The clusters were labeled according to the format denovo_x_y, where x is a one-letter abbreviation for the taxonomic rank (k, p, c, o, f, g, and s), and y represents the number of the FL-ASV which is the cluster centroid of the particular taxon. Because the applied clustering algorithm processes the sequences sequentially in the order they appear in the input file, sequences are always clustered in the same way, even if additional FL-ASVs are later added to the database. This strategy may not always yield the most optimal clusters, but the reproducibility is critical if the clusters are to be used as a robust placeholder taxonomy.

Merging of the SILVA- and the *de novo*-based taxonomies resulted in a few conflicts, e.g., where different FL-ASVs from the same species associate with more than one genus. In such cases, the genus-level classification for the centroid FL-ASV is adapted for all FL-ASVs within that species. However, these types of conflicts only applied to lower rank taxa (species), which were located close to the taxonomic threshold of the higher rank taxa (genus), and it only affected the classification of a low number of FL-ASVs (approx. 1%).

As AutoTax is based on the SILVA taxonomy, the taxonomy generated will change if

another version of the SILVA SSURef NR99 reference database is used. Accordingly, users must

specify which version of the SILVA database has been used when they publish databases created

with AutoTax. We recommend that AutoTax generated databases are updated when there is a new

version of the SILVA database. This ensures that the taxonomy is in agreement with the current

central taxonomy.

Taxonomy assignment to FL-ASVs with AutoTax

AutoTax provided placeholder names for many previously undescribed taxa in our FL-ASV

database (**Table 2, Figure S4**). 95% of all species, 73% of all genera, 47% of all families, and 24%

of all orders obtained placeholder names from the *de novo* taxonomy and would otherwise have

remained unclassified. The novel taxa were affiliated with several phyla, especially the

Proteobacteria, Planctomycetota, Patescibacteria, Firmicutes, Chloroflexi, Bacteroidota,

Actinobacteriota, and Acidobacteriota (**Figure S4**). A prominent example is the Chloroflexi, where

5 out of 14 orders, 26 out of 34 families, and 142 out of 152 genera observed were assigned a *de*

novo placeholder taxonomy. We believe that this will have important implications for future studies

and the accumulation and transfer of knowledge about these taxa in WWTPs, given their high

diversity and abundance, and their association with serious operational problems related to the

settling of activated sludge (bulking) and foaming (59, 60). It should be noted that the placeholder

taxonomy does not provide the same degree of support as traditional phylogenetic analyses, and

should only be used until an official taxonomy is established.

Improved classification of ASVs from WWTP and anaerobic digesters

To benchmark the FL-ASV database, we classified V1-3 amplicon data obtained from activated

sludge and anaerobic digester samples (**Table S2**) using this database with the SINTAX-classifier

and compared the results to classifications obtained using the universal reference databases (**Figure**

3a). With the use of the FL-ASV database, many more of the abundant ASVs (relative

abundance >0.01%) were classified to the genus and species level (89.9±4.3% and 78.5±4.0%,

respectively), compared to the use of popular public reference databases, including SILVA 138

SSURef NR99 (30.4±3.5% and 0%), GreenGenes 16s v. 13.5 (24.5±4.4% and 1.4±0.4%), GTDB

r89 (22.1±2.8% and 11.6±1.2%), the RDP 16S v16 training set (20.3±4.0% and 0%), and even the

MiDAS 2.1 (59.7±4.5% and 0.4±0.3%) - which is a manually ecosystem-specific curated version of

the SILVA 123 SSURef NR99 database.

We have here shown that increased coverage of the rare biosphere can be achieved by

adding FL-OTUs clustered at 99% to the FL-ASV database. The FL-OTU expanded database

increased the classification rate at the genus level for the abundant bacteria (relative abundance >

0.01%) (94.2±1.4% vs. 89.9±4.3%), but reduced classification at the species level (67.7±2.6% vs.

78.5±4.0%) (**Figure 3a**). When ASVs from the rare biosphere (relative abundance > 0.001%) were

included in the analysis, the advantage of including FL-OTUs for genus-level classification was

even more pronounced (93.0±0.6% vs. 80.7±1.7%), and an improved classification rate at the

species level was also observed (69.9±1.5% vs. 66.8±2.4%) (**Figure S5**). We hypothesize that the

reduced classification rate at the species level for the abundant ASVs is caused by sequencing errors

as well as low-divergence chimera, which cannot be detected by chimera filtering (61). It should be

noted that FL-OTUs are problematic for databases that are to be maintained and updated with

additional references in the future. This is because the placeholder taxonomy will change if FL-

OTUs are removed, and sequencing errors and chimeras are propagated if they are kept. We

therefore recommend that FL-OTUs are only added for exploratory purposes.

To investigate the influence of AutoTax on taxonomic assignment, independent of our

ecosystem-specific database, we applied the pipeline directly to high-quality, full-length sequences

from the SILVA 138 SSURef_NR99 database. This increased the percentage of ASVs classified

with SILVA at the genus level from 30.4±3.5% to 63.7±4.8%, and at the species-level from 0% to

28.2±2.4%, suggesting that the large universal databases would also benefit from the use of

AutoTax. An advantage of the AutoTax processed SILVA database, which we have made publicly

available, is that the placeholder taxonomy is universally applicable, providing a unique opportunity

for studying the ecology of unclassified taxa across ecosystems.

Taxonomic resolution of ASVs in combination with a comprehensive reference database

Classification of amplicon sequences can be challenging due to the limited taxonomic information

in short-read sequences (12, 13). However, this may change with the access to reference databases

with perfect references for the majority of all ASVs and a complete seven-rank taxonomy for all

reference sequences. To determine the confidence of the amplicon classification in this scenario, we

extracted ASVs *in silico* from the bacterial FL-ASVs corresponding to commonly amplified 16S

rRNA regions, including full-length amplicons. These ASVs were classified against the FL-ASV

database. We then calculated the fraction of amplicons correctly classified to the same genus and

species as their corresponding FL-ASVs (**Figure 3b and Figure S6a**). Nearly all ASVs (95-99%)

were assigned to the correct genus and most (72-91%) to the right species, depending on the

taxonomic conservation of 16S rRNA region covered by the *in silico* amplicons. The primers

targeting the V1-3 variable region performed exceptionally well for species-level identification

(90.7% correct classifications), which is the same as for the full-length 16S rRNA gene amplicons.

The commonly used primers targeting the V4 variable region were the worst (72.5% correct

classifications). Very few of the sequences that did not receive the same taxonomic classification as

their source reference sequence where misclassified (<0.2% at genus-level and <0.8% at species-

level), with the majority not receiving any classification at the specific taxonomic rank (**Figure

S6a**).

Sequencing costs on the Illumina platforms can be reduced considerably if single reads are

used instead of merged reads. To evaluate the effect of reduced amplicon length, we compared the

classification of 200 bp forward reads and reverse reads to those of full-length amplicons (**Figure

3b**). The decrease in the percentage of correct classifications was highly dependent on the 16S

rRNA region targeted and from which direction the single reads were obtained. For the ecosystem

studied here, the V1-3 and V4-5 forward read provided almost the same specificity as the full-

length amplicons, whereas the reverse reads performed much worse for species-level classification.

For the V3-4, V5-7, and V5-8, the reverse reads performed better than the forward reads, revealing

the importance of choosing the right direction for single read amplicons.

To evaluate the effect of sequencing errors and low-divergence chimeras in the reference

database, we also classified the *in silico* ASVs against the reference databases expanded with the

FL-OTUs (**Figure S6b**). The result confirmed our prior observations that the inclusion of error

prone references had a negative impact on our ability to classify short-read amplicons correctly;

however, the effect was marginal at the genus-level. Full-length amplicons were less affected

(**Figure S6b**), highlighting a clear advantage of using longer amplicons in combination with

universal databases, which are likely to contain sequencing errors and low-divergence chimeras

despite chimera filtering (61).

Overall, the analysis demonstrated that confident classification of short-read ASV sequences

at the genus to species level is possible. However, it requires a reference database with a complete

seven-rank taxonomy and perfect references for the majority of all ASVs. The scripts made

available with this study can be used to confirm whether this is the case for samples from other

studies.

Ecosystem-specific evaluation of primer-bias

When choosing primers for amplicon sequence analyses, it is essential also to take primer-bias into

account as some primer sets may result in ecosystem-relevant species being severely

underestimated or absent from the analyses (30). The ecosystem-specific FL-ASV databases

provide a near-perfect reference to determine the theoretical coverage of different primer sets for

the given ecosystem so that an informed selection can be made (46, 62). It should be noted that

primers used to generate the full-length 16S rRNA sequences for FL-ASV databases may introduce

a bias, and we have here include primer-free (RNA-based) libraries to account for this. Evaluation

of different primer sets using our FL-ASV database revealed a clear taxonomic bias associated with

several primer sets (**Figure 3c**). The primers targeting the V4-5 and V5-8 regions had the best

coverage of the FL-ASVs. The V5-7 primers demonstrated a very poor coverage. Because the FL-

ASVs were trimmed after the forward priming site of the V1-3 primers, we were not able to

evaluate the coverage of this primer pair here. However, it has previously been shown that the

primers have a good overall agreement with metagenomic data for wastewater treatment systems

and capture most of the process-critical organisms (31).

Perspectives

High-throughput full-length 16S rRNA gene sequencing and automated taxonomy assignment

(AutoTax) now allow individual research groups to develop their own FL-ASV ecosystem-specific

reference databases for community profiling analyses. In addition, such databases can be used to

evaluate the ecosystem-specific coverage and specificity of amplicon primers and fluorescence *in*

situ hybridization (FISH) probes. The high quality of the FL-ASVs furthermore allows for the

design of new primers and probes with improved confidence of the coverage and taxonomic

resolution when applied within the target ecosystem. Collectively, the approach importantly allows

for the identification and subsequent characterization of novel numerically important taxa for the

specific environment, which would have otherwise been overlooked.

We acknowledge that the ability to generate new 16S rRNA gene reference databases quickly

poses a risk for the development of several competing divergent taxonomies. We, therefore,

recommend the custom databases are only used for exploratory purposes and combined with

traditional phylogenetic analyses of key taxa. Ecosystem-specific databases that are broadly applied, such as the MiDAS database, should be created as open-source community efforts, and universal reference databases processed with AutoTax should be published and maintained in agreement with current developers of such databases. An important benefit of this approach is that the placeholder taxonomy can be used as a common language within the field, or in the case of universal reference databases, across all fields. This has major implications, e.g., in wastewater treatment systems, as it allows for the identification of unclassified taxa that are process-critical and decisive for process performance. If the current universal reference databases are used, the majority of ASVs will not get a genus-level classification, making it impossible to compare their prevalence across studies. Given that hundreds of amplicon-based studies are carried out every year worldwide, a considerable amount of useful information is lost when the data generated across studies is not comparable.

We have chosen to use the SINTAX-classifier for our analyses because it applies a simple algorithm that does not require training, and we expect the results to be less biased. However, some classifiers which use Bayesian inference, e.g., q2-feature-classifier (5) may yield better classifications. Kaehler et al. (63) recently demonstrated what environment-specific taxonomic abundance information could be used as weights for such classifiers to improve the accuracy of the taxonomy assignment. This approach is interesting because the abundance of raw full-length 16S rRNA gene sequences used to generate the FL-ASV database may be used as phylogenetically informative weights for the specific ecosystem.

We used the SILVA SSURef NR99 as the backbone taxonomy for AutoTax in this study, as this is currently the most comprehensive database. However, we anticipate the GTDB may replace SILVA in a future release of AutoTax when more high-quality genomes and MAGs with 16S rRNA genes (MIMAG standard (11)) are added to the database as the result of advances in long-read sequencing technology (Nanopore and PacBio). This will importantly link the 16S rRNA gene taxonomy with that derived from the more robust phylogenomic-based analyses, creating a unified language across the field of microbiology.

Acknowledgments
We would like to thank the 22 wastewater treatment plants involved in the project for providing samples. This work was supported by the Danish Research Council [6111-00617A to P.H.N.]; and the Villum foundation [13351 to P.H.N.].

Conflict of Interest

The authors declare no conflict of interest.

References:

1. Martiny JBH, Jones SE, Lennon JT, Martiny AC. 2015. Microbiomes in light of traits: A phylogenetic perspective. Science 350: aac9823-18.

2. Amir A, Daniel M, Navas-Molina J, Kopylova E, Morton J, Xu ZZ, Eric K, Thompson L, Hyde E, Gonzalez A, Knight R. 2017. Deblur Rapidly Resolves Single- Nucleotide Community Sequence Patterns. Am Soc Microbiol 2:1–7.

3. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. 2016. DADA2: High-resolution sample inference from Illumina amplicon data. Nat Methods 13:581–583.

4. Edgar RC. 2016. UNOISE2: improved error-correction for Illumina 16S and ITS amplicon sequencing. bioRxiv 81257.

5. Bokulich NA, Kaehler BD, Rideout JR, Dillon M, Bolyen E, Knight R, Huttley GA, Gregory Caporaso J. 2018. Optimizing taxonomic classification of marker-gene amplicon sequences with QIIME 2's q2-feature-classifier plugin. Microbiome 6:90.

6. Wang Q, Garrity GM, Tiedje JM, Cole JR. 2007. Naive Bayesian classifier for rapid assignment of rRNA sequences into the new bacterial taxonomy. Appl Environ Microbiol 73:5261–5267.

7. Edgar R. 2016. SINTAX: a simple non-Bayesian taxonomy classifier for 16S and ITS sequences. bioRxiv 074161.

8. Callahan BJ, McMurdie PJ, Holmes SP. 2017. Exact sequence variants should replace operational taxonomic units in marker-gene data analysis. ISME J 11:2639–2643.

9. Thompson LR, Sanders JG, McDonald D, Amir A, Ladau J, Locey KJ, Prill RJ, Tripathi A, Gibbons SM, Ackermann G, Navas-Molina JA, Janssen S, Kopylova E, Vázquez-Baeza Y,

González A, Morton JT, Mirarab S, Xu ZZ, Jiang L, Haroon MF, Kanbar J, Zhu Q, Song SJ,

Kosciolek T, Bokulich NA, Lefler J, Brislawn CJ, Humphrey G, Owens SM, Hampton-

Marcell J, Berg-Lyons D, McKenzie V, Fierer N, Fuhrman JA, Clauset A, Stevens RL,

Shade A, Pollard KS, Goodwin KD, Jansson JK, Gilbert JA, Knight R, Agosto Rivera JL,

Al-Moosawi L, Alverdy J, Amato KR, Andras J, Angenent LT, Antonopoulos DA, Apprill

A, Armitage D, Ballantine K, Bárta J, Baum JK, Berry A, Bhatnagar A, Bhatnagar M, Biddle

JF, Bittner L, Boldgiv B, Bottos E, Boyer DM, Braun J, Brazelton W, Brearley FQ,

Campbell AH, Caporaso JG, Cardona C, Carroll JL, Cary SC, Casper BB, Charles TC, Chu

H, Claar DC, Clark RG, Clayton JB, Clemente JC, Cochran A, Coleman ML, Collins G,

Colwell RR, Contreras M, Crary BB, Creer S, Cristol DA, Crump BC, Cui D, Daly SE,

Davalos L, Dawson RD, Defazio J, Delsuc F, Dionisi HM, Dominguez-Bello MG, Dowell R,

Dubinsky EA, Dunn PO, Ercolini D, Espinoza RE, Ezenwa V, Fenner N, Findlay HS,

Fleming ID, Fogliano V, Forsman A, Freeman C, Friedman ES, Galindo G, Garcia L,

Garcia-Amado MA, Garshelis D, Gasser RB, Gerdts G, Gibson MK, Gifford I, Gill RT,

Giray T, Gittel A, Golyshin P, Gong D, Grossart HP, Guyton K, Haig SJ, Hale V, Hall RS,

Hallam SJ, Handley KM, Hasan NA, Haydon SR, Hickman JE, Hidalgo G, Hofmockel KS,

Hooker J, Hulth S, Hultman J, Hyde E, Ibáñez-Álamo JD, Jastrow JD, Jex AR, Johnson LS,

Johnston ER, Joseph S, Jurburg SD, Jurelevicius D, Karlsson A, Karlsson R, Kauppinen S,

Kellogg CTE, Kennedy SJ, Kerkhof LJ, King GM, Kling GW, Koehler A V., Krezalek M,

Kueneman J, Lamendella R, Landon EM, Lanede Graaf K, LaRoche J, Larsen P, Laverock

B, Lax S, Lentino M, Levin II, Liancourt P, Liang W, Linz AM, Lipson DA, Liu Y, Lladser

ME, Lozada M, Spirito CM, MacCormack WP, MacRae-Crerar A, Magris M, Martín-Platero

AM, Martín-Vivaldi M, Martínez LM, Martínez-Bueno M, Marzinelli EM, Mason OU,

Mayer GD, McDevitt-Irwin JM, McDonald JE, McGuire KL, McMahon KD, McMinds R,

Medina M, Mendelson JR, Metcalf JL, Meyer F, Michelangeli F, Miller K, Mills DA, Minich

J, Mocali S, Moitinho-Silva L, Moore A, Morgan-Kiss RM, Munroe P, Myrold D, Neufeld

JD, Ni Y, Nicol GW, Nielsen S, Nissimov JI, Niu K, Nolan MJ, Noyce K, O'Brien SL,

Okamoto N, Orlando L, Castellano YO, Osuolale O, Oswald W, Parnell J, Peralta-Sánchez

JM, Petraitis P, Pfister C, Pilon-Smits E, Piombino P, Pointing SB, Pollock FJ, Potter C,

Prithiviraj B, Quince C, Rani A, Ranjan R, Rao S, Rees AP, Richardson M, Riebesell U,

Robinson C, Rockne KJ, Rodriguezl SM, Rohwer F, Roundstone W, Safran RJ, Sangwan N,

Sanz V, Schrenk M, Schrenzel MD, Scott NM, Seger RL, Seguinorlando A, Seldin L, Seyler

LM, Shakhsheer B, Sheets GM, Shen C, Shi Y, Shin H, Shogan BD, Shutler D, Siegel J,

Simmons S, Sjöling S, Smith DP, Soler JJ, Sperling M, Steinberg PD, Stephens B, Stevens

MA, Taghavi S, Tai V, Tait K, Tan CL, Taş N, Taylor DL, Thomas T, Timling I, Turner BL,

Urich T, Ursell LK, Van Der Lelie D, Van Treuren W, Van Zwieten L, Vargas-Robles D,

Thurber RV, Vitaglione P, Walker DA, Walters WA, Wang S, Wang T, Weaver T, Webster

NS, Wehrle B, Weisenhorn P, Weiss S, Werner JJ, West K, Whitehead A, Whitehead SR,

Whittingham LA, Willerslev E, Williams AE, Wood SA, Woodhams DC, Yang Y, Zaneveld

J, Zarraonaindia I, Zhang Q, Zhao H. 2017. A communal catalogue reveals Earth's multiscale

microbial diversity. Nature 551:457–463.

10.   Mcdonald D, Hyde E, Debelius JW, Morton JT, Gonzalez A, Ackermann G, Aksenov AA,

Behsaz B, Brennan C, Chen Y, Goldasich D, Dorrestein PC, Dunn RR, Fahimipour AK,

Gaffney J, Gilbert JA, Gogul G, Green JL, Hugenholtz P, Humphrey G, Huttenhower C,

Jackson MA, Janssen S, Jeste D V, Jiang L, Kelley ST, Knights D, Kosciolek T, Ladau J,

Leach J, Marotz C, Meleshko D, Melnik A V, Metcalf JL, Mohimani H, Montassier E,

Rahnavard G, Robbins-pianka A, Sangwan N, Shorenstein J, Smarr L, Vázquez-baeza Y,

Vrbanac A, Wischmeyer P, Wolfe E, Zhu Q, Gut A, Jt M, Gonzalez A, Ackermann G,

Behsaz B, Brennan C, Chen Y, Goldasich L, Pc D, Rr D, Jl G, Hugenholtz P, Humphrey G,

Huttenhower C, Ma J, Janssen S, Jiang L, St K, Knights D, Kosciolek T. 2018. American

Gut : an Open Platform for Citizen Science. mSystems 3:1–28.

11.   Bowers RM, Kyrpides NC, Stepanauskas R, Harmon-Smith M, Doud D, Reddy TBK, Schulz

F, Jarett J, Rivers AR, Eloe-Fadrosh EA, Tringe SG, Ivanova NN, Copeland A, Clum A,

Becraft ED, Malmstrom RR, Birren B, Podar M, Bork P, Weinstock GM, Garrity GM,

Dodsworth JA, Yooseph S, Sutton G, Glöckner FO, Gilbert JA, Nelson WC, Hallam SJ,

Jungbluth SP, Ettema TJG, Tighe S, Konstantinidis KT, Liu WT, Baker BJ, Rattei T, Eisen

JA, Hedlund B, McMahon KD, Fierer N, Knight R, Finn R, Cochrane G, Karsch-Mizrachi I,

Tyson GW, Rinke C, Lapidus A, Meyer F, Yilmaz P, Parks DH, Eren AM, Schriml L,

Banfield JF, Hugenholtz P, Woyke T. 2017. Minimum information about a single amplified

genome (MISAG) and a metagenome-assembled genome (MIMAG) of bacteria and archaea.

Nat Biotechnol 35:725–731.

12.   Yarza P, Yilmaz P, Pruesse E, Glöckner FO, Ludwig W, Schleifer K-H, Whitman WB,

Euzéby J, Amann R, Rosselló-Móra R. 2014. Uniting the classification of cultured and

uncultured bacteria and archaea using 16S rRNA gene sequences. Nat Rev Microbiol

12:635–645.

13.   Edgar RC. 2018. Accuracy of taxonomy prediction for 16S rRNA and fungal ITS sequences.

PeerJ 6:e4652.

14.   Desantis TZ, Hugenholtz P, Larsen N, Rojas M, Brodie EL, Keller K, Huber T, Dalevi D, Hu

P, Andersen GL. 2006. Greengenes , a Chimera-Checked 16S rRNA Gene Database and

Workbench Compatible with ARB. Appl Environ Microbiol 72:5069–5072.

15.   Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glöckner FO. 2013.

The SILVA ribosomal RNA gene database project: Improved data processing and web-based

tools. Nucleic Acids Res 41:D590-6.

16.   Cole JR, Wang Q, Fish JA, Chai B, McGarrell DM, Sun Y, Brown CT, Porras-Alfaro A,

Kuske CR, Tiedje JM. 2014. Ribosomal Database Project: data and tools for high throughput

rRNA analysis. Nucleic Acids Res 42:D633-42.

17.   Louca S, Mazel F, Doebeli M, Parfrey LW. 2019. A census-based estimate of earth's

bacterial and archaeal diversity. PLoS Biol 17:1–30.

18.   McIlroy SJ, Kirkegaard RH, McIlroy B, Nierychlo M, Kristensen JM, Karst SM, Albertsen

M, Nielsen PH. 2017. MiDAS 2.0: An ecosystem-specific taxonomy and online database for

the organisms of wastewater treatment systems expanded for anaerobic digester groups.

Database 2017:1–9.

19.   Mikaelyan A, Köhler T, Lampert N, Rohland J, Boga H, Meuser K, Brune A. 2015.

Classifying the bacterial gut microbiota of termites and cockroaches: A curated phylogenetic

reference database (DictDb). Syst Appl Microbiol 38:472–482.

20.   Ritari J, Salojärvi J, Lahti L, de Vos WM. 2015. Improved taxonomic assignment of human

intestinal 16S rRNA sequences by a dedicated reference database. BMC Genomics 16:1056.

21.   Chen T, Yu W-H, Izard J, Baranova O V, Lakshmanan A, Dewhirst FE. 2010. The Human

Oral Microbiome Database: a web accessible resource for investigating oral microbe

taxonomic and genomic information. Database (Oxford) 2010:baq013.

22.   Newton RJ, Jones SE, Eiler A, McMahon KD, Bertilsson S. 2011. A Guide to the Natural
669   History of Freshwater Lake Bacteria. Microbiol and Mol Biol Rev 75:14-49.

670   23.   Rohwer RR, Hamilton JJ, Newton RJ, McMahon KD. 2018. TaxAss: Leveraging a Custom
671   Freshwater Database Achieves Fine-Scale Taxonomic Resolution. mSphere 3:1–14.

672   24.   Newton IL, Roeselers G. 2012. The effect of training set on the classification of honey bee
673   gut microbiota using the Naïve Bayesian Classifier. BMC Microbiol 12:221.

674   25.   Seedorf H, Kittelmann S, Henderson G, Janssen PH. 2014. RIM-DB: a taxonomic framework
675   for community structure analysis of methanogenic archaea from the rumen and other
676   intestinal environments. PeerJ 2:e494.

677   26.   Karst SM, Dueholm MS, McIlroy SJ, Kirkegaard RH, Nielsen PH, Albertsen M. 2018.
678   Retrieval of a million high-quality, full-length microbial 16S and 18S rRNA gene sequences
679   without primer bias. Nat Biotechnol 36:190–195.

680   27.   Burke CM, Darling AE. 2016. A method for high precision sequencing of near full-length
681   16S rRNA genes on an Illumina MiSeq. PeerJ 4:e2492.

682   28.   Callahan BJ, Wong J, Heiner C, Oh S, Theriot CM, Gulati AS, McGill SK, Dougherty MK.
683   2019. High-throughput amplicon sequencing of the full-length 16S rRNA gene with single-
684   nucleotide resolution. Nucleic Acids Res 47:e103.

685   29.   Karst SM, Ziels RM, Kirkegaard RH, Albertsen M. 2019. Enabling high-accuracy long-read
686   amplicon sequences using unique molecular identifiers and Nanopore sequencing. bioRxiv
687   645903.

688   30.   Albertsen M, Karst SM, Ziegler AS, Kirkegaard RH, Nielsen PH. 2015. Back to basics - the
689   influence of DNA extraction and primer choice on phylogenetic analysis of activated sludge
690   communities. PLoS One 10:e0132783.

691   31.   Lane DJ. 1991. 16S/23S rRNA sequencing, p. 115–175. *In* Stackebrandt, E, Goodfellow, M
692   (eds.), Nucleic Acid Techniques in Bacterial Systematics. John Wiley and Sons, Chichester,
693   United Kingdom.

694   32.   Massana R, Murray AE, Preston CM, DeLong EF. 1997. Vertical distribution and
695   phylogenetic characterization of marine planktonic Archaea in the Santa Barbara Channel.
696   Appl Environ Microbiol 63:50–56.

697   33.   Bahram M, Anslan S, Hildebrand F, Bork P, Tedersoo L. 2018. Newly designed 16S rRNA
698   metabarcoding primers amplify diverse and novel archaeal taxa from the environment.
699   Environ Microbiol Rep 11:487–494.

700   34.   Tange O. 2018. GNU Parallel 2018. Ole Tange.

701   35.   Edgar RC. 2010. Search and clustering orders of magnitude faster than BLAST.
702   Bioinformatics 26:2460–2461.

703   36.   Pruesse E, Peplies J, Glöckner FO. 2012. SINA: Accurate high-throughput multiple sequence
704   alignment of ribosomal RNA genes. Bioinformatics 28:1823–1829.

705   37.   Pagès H, Aboyoun P, Gentleman R, DebRoy S. 2019. Biostrings: Efficient manipulation of
706   biological strings.

707   38.   Corporation M, Weston S. 2019. doParallel: Foreach Parallel Adaptor for the "parallel"
708   Package.

709   39.   Wickham H. 2019. stringr: Simple, Consistent Wrappers for Common String Operations.

710   40.   Dowle M, Srinivasan A. 2019. data.table: Extension of `data.frame`.

711   41.   Wickham H, Henry L. 2019. tidyr: Easily Tidy Data with "spread()" and "gather()"
712   Functions.

713   42.   Wickham H, François R, Henry L, Müller K. 2019. dplyr: A Grammar of Data Manipulation.

714   43.   Kirkegaard RH, McIlroy SJ, Kristensen JM, Nierychlo M, Karst SM, Dueholm MS,
715   Albertsen M, Nielsen PH. 2017. The impact of immigration on microbial community
716   composition in full-scale anaerobic digesters. Sci Rep 7:9343

717   44.   Muyzer G, de Waal EC, Uitterlinden AG. 1993. Profiling of complex microbial populations
718   by denaturing gradient gel electrophoresis analysis of polymerase chain reaction-amplified

genes coding for 16S rRNA. AEM 59:695–700.

45.   Price MN, Dehal PS, Arkin AP. 2010. FastTree 2--approximately maximum-likelihood trees
for large alignments. PLoS One 5:e9490.

46.   Walters WA, Caporaso JG, Lauber CL, Berg-Lyons D, Fierer N, Knight R. 2011.
PrimerProspector: De novo design and taxonomic analysis of barcoded polymerase chain
reaction primers. Bioinformatics 27:1159–1161.

47.   R Core Team. 2016. R: A language and environment for statistical computing. R Foundation
for Statistical Computing, Vienna, Austria.

48.   RStudio Team. 2015. RStudio: Integrated Development Environment for R. Boston, MA.

49.   Wickham H. 2009. ggplot2 - Elegant Graphics for Data AnalysisSpringer. Springer Science
& Business Media.

50.   Andersen KS, Kirkegaard RH, Karst SM, Albertsen M. 2018. ampvis2: an R package to
analyse and visualise 16S rRNA amplicon data. bioRxiv 299537.

51.   Wu L, Ning D, Zhang B, Li Y, Zhang P, Shan X, Zhang Q, Brown M, Li Z, Van Nostrand
JD, Ling F, Xiao N, Zhang Y, Vierheilig J, Wells GF, Yang Y, Deng Y, Tu Q, Wang A,
Zhang T, He Z, Keller J, Nielsen PH, Alvarez PJJ, Criddle CS, Wagner M, Tiedje JM, He Q,
Curtis TP, Stahl DA, Alvarez-Cohen L, Rittmann BE, Wen X, Zhou J. 2019. Global diversity
and biogeography of bacterial communities in wastewater treatment plants. Nat Microbiol
4:1183-1195.

52.   Parks DH, Chuvochina M, Waite DW, Rinke C, Skarshewski A, Chaumeil P-A, Hugenholtz
P. 2018. A standardized bacterial taxonomy based on genome phylogeny substantially
revises the tree of life. Nat Biotechnol 36:996-1004.

53.   Parks DH, Rinke C, Chuvochina M, Chaumeil P-A, Woodcroft BJ, Evans PN, Hugenholtz P,
Tyson GW. 2017. Recovery of nearly 8,000 metagenome-assembled genomes substantially
expands the tree of life. Nat Microbiol 2:1533–1542.

54.   Moss EL, Maghini DG, Bhatt AS. 2020. Complete, closed bacterial genomes from

microbiomes using nanopore sequencing. Nat Biotechnol. https://doi.org/10.1038/s41587-
020-0422-6

55. Singleton CM, Petriglieri F, Kristensen JM, Kirkegaard RH, Michaelsen TY, Andersen MH,
Kondrotaite Z, Karst SM, Dueholm MS, Nielsen PH, Albertsen M. 2020. Connecting
structure to function with the recovery of over 1000 high-quality activated sludge
metagenome-assembled genomes encoding full-length rRNA genes using long-read
sequencing. bioRxiv 088096.

56. Isazadeh S, Jauffur S, Frigon D. 2016. Bacterial community assembly in activated sludge:
mapping beta diversity across environmental variables. MicrobiologyOpen 5:1050–1060.

57. Gonzalez-Martinez A, Rodriguez-Sanchez A, Lotti T, Garcia-Ruiz MJ, Osorio F, Gonzalez-
Lopez J, Van Loosdrecht MCM. 2016. Comparison of bacterial communities of conventional
and A-stage activated sludge systems. Sci Rep 6:18786.

58. Yilmaz P, Parfrey LW, Yarza P, Gerken J, Pruesse E, Quast C, Schweer T, Peplies J, Ludwig
W, Glöckner FO. 2014. The SILVA and "All-species Living Tree Project (LTP)" taxonomic
frameworks. Nucleic Acids Res 42:D643–D648.

59. McIlroy SJ, Karst SM, Nierychlo M, Dueholm MS, Albertsen M, Kirkegaard RH, Seviour
RJ, Nielsen PH. 2016. Genomic and in situ investigations of the novel uncultured Chloroflexi
associated with 0092 morphotype filamentous bulking in activated sludge. ISME J 10:2223-
2234.

60. Nierychlo M, Mcilroy SJ, Kucheryavskiy S, Jiang C, Ziegler AS, Kondrotaite Z, Stokholm-
Bjerregaard M, Nielsen PH. 2020. Candidatus Amarolinea and Candidatus Microthrix Are
Mainly Responsible for Filamentous Bulking in Municipal Danish Wastewater Treatment
Plants. Front Microbiol. doi: 10.3389/fmicb.2020.01214

61. Edgar RC. 2016. UCHIME2: improved chimera prediction for amplicon sequencing. bioRxiv
074252.

62. Klindworth A, Pruesse E, Schweer T, Peplies J, Quast C, Horn M, Glöckner FO. 2013.
Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-
generation sequencing-based diversity studies. Nucleic Acids Res 41:1–11.

63.   Kaehler BD, Bokulich NA, McDonald D, Knight R, Caporaso JG, Huttley GA. 2019.

774   Species abundance information improves sequence taxonomy classification accuracy. Nat

775   Commun 10:1–10.

776   64.   Peterson J, Bonazzi V, Starke-Reed P, Read J, Zakhari S, Baker CC, Garges S, Howcroft TK,

777   Wang L, Giovanni M, Grandison L, McEwen JE, Watson B, Lunsford RD, Akolkar B, Guyer

778   M, Giblin C, David H, Khalsa J, Humble M, Pontzer C, Mills M, Salomon R, Wright M,

779   Schloss JA, Begg L, Belachew T, Sayre MH, Karp RW, Davis C, Mullins C, Peavy H, Di

780   Francesco V, Wetterstrand KA, McInnes P, Wellington CR, Portnoy M, Little AR, Deal C.

781   2009. The NIH Human Microbiome Project. Genome Res 19:2317–2323.

782   65.   Apprill A, Mcnally S, Parsons R, Weber L. 2015. Minor revision to V4 region SSU rRNA

783   806R gene primer greatly increases detection of SAR11 bacterioplankton. Aquat Microb

784   Ecol 75:129–137.

785   66.   Parada AE, Needham DM, Fuhrman JA. 2016. Every base matters: Assessing small subunit

786   rRNA primers for marine microbiomes with mock communities, time series and global field

787   samples. Environ Microbiol 18:1403–1414.

788   67.   Bodenhausen N, Horton MW, Bergelson J. 2013. Bacterial Communities Associated with the

789   Leaves and the Roots of Arabidopsis thaliana. PLoS One 8:e56329.

790   68.   Chelius MK, Triplett EW. 2001. The diversity of archaea and bacteria in association with the

791   roots of Zea mays L. Microb Ecol 41:252–263.

792   69.   Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing

793   reads. EMBnet.journal 17:10–12.

Tables:

Table 1: Numbers and percentage of FL-ASVs estimated to belong to novel taxa. FL-ASVs were mapped to SILVA 138 SSURef NR99 to find the identity with the closest relative in the database. The novelty was determined based on the identity for each FL-ASV using the taxonomic thresholds proposed by Yarza et al. (2014).

	Sequences	Percentage
Phylum (<75.0%)	1	0.01%
Class (<78.5%)	1	0.01%
Order (<82.0%)	3	0.03%
Family (<86.5%)	16	0.17%
Genus (<94.5%)	548	5.76%
Species (<98.7%)	2449	25.7%

Table 2: Numbers and percentage of taxa that were assigned *de novo* names.

	De novo taxa	Percentage
Phylum	1	2.22%
Class	12	10.17%
Order	66	24.35%
Family	276	47.18%
Genus	1324	72.91%
Species	3973	95.28%

Table S1: Oligonucleotides used for full-length 16S rRNA gene library preparation. Unique

molecular tags, and sample barcodes are marked with blue and red, respectively. The GGGGGG

motif marked with bold typefaces may form strong secondary structures known as guanine

tetraplexes. This motif was replaced by GCGCGG in the DNA-based protocol, which increased the

robustness of library preparation. Oligoes marked with "*" are identical between the RNA- and

DNA-based protocols.

Name	Sequence	Protocol step
SSU_rRNA_RT2	A**GGGGGG**CAAAGATGAAGATNNNNNNNNNNNNNNNCTAGTACGACTTGCCTGTCGCTCTATCTTCTTTTTTTTTTTTTTTTTTTTVN	First strand cDNA synthesis
SSU_rRNA_RT3	A**GGGGGG**CAAAGATGAAGATNNNNNNNNNNNNNNNTTCTGCCTACTTGCCTGTCGCTCTATCTTCTTTTTTTTTTTTTTTTTTTTVN	
SSU_rRNA_s	5'Phos-GGGCAATATCAGCACCAACAGAAA-3'SpC3	Adaptor ligation
SSU_rRNA_l	CTCCACCCAGACTCATCCATNNNNNNNNNNNNNNNATAGAGGCTTTCTGTTGGTGCTGATATTGC	Second strand cDNA syntehsis
SSU_rRNA_pcr_fw*	CTCCACCCAGACTCATCCAT	Library and clonal amplification
SSU_rRNA_pcr_rv	A**GGGGGG**CAAAGATGAAGAT	
SSU_rRNA_readtag_fw*	CAAGCAGAAGACGGCATACGAGATGTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTCTCCACCCAGACTCATCCAT	Read-tag library preparation
SSU_rRNA_readtag_rv	CAAGCAGAAGACGGCATACGAGATGTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTA**GGGGGG**CAAAGATGAAGAT	
SSU_rRNA_linktag_fw	AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGACGCTCTTCCGATCTGAAGATAGAGCGACAGGCAAGT	Linked-tag library preparation
SSU_rRNA_linktag_rv	CAAGCAGAAGACGGCATACGAGATCGGTCTCGGCATTCCTGCTGAACCGCTCTTCCGATCTGCAATATCAGCACCAACAGAAA	
SSU_rRNA_read2_fw*	GCTCTTCCGATCTCTCCACCCAGACTCATCCAT	Illumina MiSeq/HiSeq sequencing
SSU_rRNA_read2_rv	GCTCTTCCGATCTA**GGGGGG**CAAAGATGAAGAT	
f16S_rDNA_pcr1_fw1	CTCCACCCAGACTCATCCATNNNNNNNNNNNNNNNTGCCTCTTAGAGTTTGATCMTGGCTCAG	Adaptor tagging
f16S_rDNA_pcr1_fw2	CTCCACCCAGACTCATCCATNNNNNNNNNNNNNNNTCCTCTACAGAGTTTGATCMTGGCTCAG	
f16S_rDNA_pcr1_rw-v2	A**GCGCGG**CAAAGATGAAGATNNNNNNNNNNNNNNNCGTACTAGTACGGYTACCTTGTTACGACTT	
f16S_rDNA_pcr2_fw*	CTCCACCCAGACTCATCCAT	Library and clonal amplification
f16S_rDNA_pcr2_rv-v2	A**GCGCGG**CAAAGATGAAGAT	
f16S_rDNA_readtag_fw*	CAAGCAGAAGACGGCATACGAGATGTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTCTCCACCCAGACTCATCCAT	Read-tag library preparation
f16S_rDNA_readtag_rv	CAAGCAGAAGACGGCATACGAGATGTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTA**GCGCGG**CAAAGATGAAGAT	
f16S_rDNA_linktag_fw	CAAGCAGAAGACGGCATACGAGATCGGTCTCGGCATTCCTGCTGAACCGCTCTTCCGATCTCTGAGCCAKGATCAAACTCT	Linked-tag library preparation
f16S_rDNA_linktag_rv	AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGACGCTCTTCCGATCTAAGTCGTAACAAGGTARCCGTA	
f16S_rDNA_read2_fw*	GCTCTTCCGATCTCTCCACCCAGACTCATCCAT	Illumina MiSeq/HiSeq sequencing
f16S_rDNA_read2_rv-v2	GCTCTTCCGATCTA**GCGCGG**CAAAGATGAAGAT	

811 **Table S2: Samples used in this study.**

Sample/SeqID	Environment	Location	Sampling_date	Latitude	Longitude	Design	PMID
MQ180605-111	WWTP	Fredericia	150824	55.552368	9.720404	EBPR	This study
MQ180605-112	WWTP	Randers	150831	56.453767	10.070249	EBPR	This study
MQ180605-113	WWTP	Avedoere	150902	55.607888	12.448949	EBPR	This study
MQ180605-114	WWTP	Damhusaaen	150831	55.641184	12.505800	EBPR	This study
MQ180605-115	WWTP	Viborg	150825	56.425251	9.455198	EBPR	This study
MQ180605-116	WWTP	Ribe	150824	55.329053	8.74336	EBPR	This study
MQ180605-117	WWTP	Skive	150825	56.565132	9.042158	EBPR	This study
MQ180605-118	WWTP	Ejby Moelle	150825	55.399534	10.414787	EBPR	This study
MQ180605-119	WWTP	Esbjerg W	150921	55.488097	8.430505	BNR	This study
MQ180605-120	WWTP	Aalborg E	150824	57.045161	10.045761	EBPR	This study
MQ180605-121	WWTP	Haderslev	150902	55.249786	9.508609	EBPR	This study
MQ180605-122	WWTP	Odense NE	150908	55.432604	10.458855	EBPR	This study
MQ180605-123	WWTP	Odense NW	150908	55.421534	10.366234	BNR	This study
MQ180605-124	WWTP	CP Kelco	150901	55.507272	12.145488	BNR	This study
MQ180605-125	WWTP	Aalborg W	150824	57.049513	9.864788	EBPR	This study
MQ180605-126	WWTP	Hjoerring	150831	57.421265	9.975411	EBPR	This study
MQ180605-127	WWTP	Egaa	150825	56.21314	10.242467	EBPR	This study
MQ180605-128	WWTP	Esbjerg E	150826	55.458597	8.489092	BNR	This study
MQ180605-129	WWTP	Bjergmarken	150825	55.649286	12.058568	EBPR	This study
MQ180605-130	WWTP	Boeslum	150921	56.198467	10.728742	EBPR	This study
MQ180605-131	WWTP	Mariagerfjord	150831	56.741769	10.115426	EBPR	This study
MQ180605-132	WWTP	Hirtshals	150914	57.577275	9.992971	EBPR	This study
16SAMP-4598	WWTP	Lundtofte	140623	55.802079	12.538763	EBPR	28496434
16SAMP-4601	WWTP	Ringkoebing	140625	56.089793	8.236342	EBPR	28496434
16SAMP-2340	WWTP	Aaby	100819	56.150696	10.176274	EBPR	28496434
16SAMP-2643	WWTP	Aars	100819	56.810414	9.547714	BNR	28496434
16SAMP-2589	WWTP	Fornaes	110819	56.437583	10.913985	BNR	28496434
MQ180605-133	Anaerobic digester	Bjergmarken	150826	55.6494	12.0589	Thermophilic	This study
MQ180605-134	Anaerobic digester	Ejby Moelle	150826	55.3981	10.415	Mesophilic	This study
MQ180605-135	Anaerobic digester	Fredericia	150826	55.5752	9.7476	THP + Mesophilic	This study
MQ180605-136	Anaerobic digester	Randers	150826	56.4697	10.0373	Mesophilic	This study
MQ180605-137	Anaerobic digester	Slagelse	150826	55.4163	11.3404	Mesophilic	This study
MQ180605-138	Anaerobic digester	Viborg	150826	56.4251	9.4543	Mesophilic	This study
MQ180605-139	Anaerobic digester	Aaby	150826	56.1512	10.1756	Thermophilic	This study
MQ180605-140	Anaerobic digester	Aalborg W	150826	57.0493	9.8635	Thermophilic	This study
MQ180605-141	Anaerobic digester	Esbjerg W	150826	55.4882	8.4307	Mesophilic	This study
MQ180605-142	Anaerobic digester	Damhusaaen	150902	55.6408	12.5059	Mesophilic	This study
MQ180605-143	Anaerobic digester	Fornaes	150902	56.4377	10.9145	Mesophilic	This study
MQ180605-144	Anaerobic digester	Hjoerring	150902	57.4196	9.9756	Mesophilic	This study
MQ180605-145	Anaerobic digester	Avedoere	150902	55.6087	12.4505	Mesophilic	This study
MQ180605-146	Anaerobic digester	Mariagerfjord	150902	56.741769	10.115189	Mesophilic	This study
MQ180605-147	Anaerobic digester	Naestved	150826	55.2111	11.7285	THP + Mesophilic	This study
MQ180605-148	Anaerobic digester	Soeholt	150826	56.1753	9.5826	Mesophilic	This study
16SAMP-3460	Anaerobic digester	Hobro	130828	56.6375	9.8085	Mesophilic	28839166
16SAMP-3463	Anaerobic digester	Lundtofte	130828	55.8024	12.5388	Mesophilic	28839166
16SAMP-4077	Anaerobic digester	Ringkoebing	130828	56.0882	8.2568	Mesophilic	28839166
16SAMP-4081	Anaerobic digester	Egaa	130501	56.21314	10.242467	Mesophilic	This study
16SAMP-3459	Anaerobic digester	Hobro	130821	56.1353	8.9731	NA	This study
16SAMP-4076	Anaerobic digester	Nysted	130821	54.699063	11.712259	NA	This study

* THP: thermal hydrolysis pretreatment; PMID: Pubmed ID reference

Table S3: Classification of raw assemblies, full-length 16S rRNA, and FL-ASVs.

Environment	Library type	Data set	nSeqs	A16S	A23S	B16S	B23S	E18S	E28S	Unclassified
Activated sludge	RNA based	Raw assemblies	274,240	2,730	306	119,947	129,851	6,139	13,689	1,540
Activated sludge	RNA based	Full-length 16S rRNA	45,907	396	0	45,496	0	15	0	0
Activated sludge	DNA based	Raw assemblies	506,519	0	0	506,518	0	0	0	1
Activated sludge	DNA based	Full-length 16S rRNA	468,426	0	0	468,425	0	0	0	1
Anaerobic digester	RNA based	Raw assemblies	361,091	73,583	8,990	79,057	196,916	0	206	2,339
Anaerobic digester	RNA based	Full-length 16S rRNA	38,834	16,494	0	22,340	0	0	0	0
Anaerobic digester	DNA based	Raw assemblies	402,232	0	0	402,232	0	0	0	0
Anaerobic digester	DNA based	Full-length 16S rRNA	373,340	0	0	373,340	0	0	0	0
Combined	Combined	FL-ASVs	9,521	75	0	9,446	0	0	0	0

Figures:

Figure 1:

Figure 1. Construction and evaluation of the FL-ASV and FL-OTU expanded reference

databases. a) Preparation of FL-ASVs and FL-OTUs. Samples were collected from WWTPs and

ADs, and DNA and RNA were extracted. Purified DNA or RNA was used for the preparation of

primer-based and "primer-free" full-length 16S rRNA libraries, respectively. These were sequenced

and processed bioinformatically to produce the FL-ASVs and FL-OTUs b) Mapping of V1-3

amplicon data to the FL-ASV reference database, the FL-OTU expanded database, and commonly

applied universal reference databases. ASVs were obtained from activated sludge and anaerobic

digester samples and filtered based on their relative abundance before the analyses to uncover how

well the databases cover the rare biosphere. The fraction of the microbial community represented by

the remaining ASVs after the filtering (coverage) is shown as the mean ± standard deviation across

plants. c) Mapping of V1-3 ASVs from Dutch WWTPs based on raw data from Gonzalez-Martinez

et al. (57). For details, see Figure S2a. d) Mapping of V3-5 ASVs from Canadian WWTPs, based

on raw data from Isazadeh et al. 2016 (56). For details, see Figure S2b.

Figure 2. The AutoTax taxonomic framework. (1) FL-ASVs (or FL-OTUs) are first mapped to

the SILVA 138 SSURef NR99 database to identify the closest relative and the shared percent

identity. (2) Taxonomy is adopted from this sequence after trimming based on percent identity and

the taxonomic thresholds proposed by Yarza *et al.* (12). To gain species-level information, FL-

ASVs are also mapped to sequences from type strains extracted from the SILVA database, and

species names are adopted if the identity is >98.7% and only a single species is within the threshold.

(3) FL-ASVs are also clustered at different percent identity, corresponding to the thresholds

proposed by Yarza *et al.* (12). The clustering is used to generate a stable *de novo* taxonomy. (4)

Finally, a comprehensive taxonomy is obtained by filling gaps in the SILVA-based taxonomy with

the *de novo*-taxonomy. Colored squares represent sources of taxonomic classifications of FL-ASVs:

orange SILVA SSURef NR99, yellow: SILVA type strains, blue: *de novo* names, grey: names

rejected during the AutoTax workflow.

Figure 3. Evaluation of amplicon classification and primer specificity. a) The fraction of V1-3

ASVs from activated sludge and anaerobic digester sample with a relative abundance higher than

0.01% that were classified to the genus and species level using the FL-ASV reference database, the

FL-OTU expanded database, and commonly applied universal reference databases, including the

Autotax-processed SILVA 138 SSURef NR99 database. b) Classification of *in silico* bacterial

ASVs, corresponding to amplicons produced using typical amplicon primers and the FL-ASV

database. Results are shown for the complete amplicons as well as for partial amplicons

corresponding to the first 200 bp from the forward or reverse read. c) Predicted primer bias

associated with commonly used amplicon primers. The trees show the bacterial FL-ASVs, and the

branch colors represent perfect hits (gray), partial hits (orange), and poor hits (red) (see materials

and methods for definitions). The following primers were used: V13 (Lane 1991) (31), V18, V19,

V34, and V58 (Klindworth et al. 2013) (62), V35 (Peterson et al. 2009) (64), V4 (Apprill et al.

2015) (65), V45 (Parada et al. 2016) (66), V57 (Chelius and Triplett 2001 and Bodenhausen et al.

2013) (67, 68).

Figure S1:

Figure S1: Mapping of activated sludge and anaerobic digester amplicon data to the FL-ASV
and FL-OTU expanded reference database and public reference databases. V1-3 ASVs were
obtained from the individual activated sludge (a) and anaerobic digester (b) samples used to create
the FL-ASV reference database (black labels) and from other WWTPs (red labels).

Figure S2: Mapping of amplicon data from non-Danish WWTP to the FL-ASV and FL-OTU

expanded reference database and public reference databases. a) ASVs were created based on

raw V1-3 amplicon reads obtained from Gonzarlez-Martinez *et al.* (57). Raw 454 genome

sequencer reads in sff format were converted to fastq using CLC genomic workbench v. 20.0.

Sequences were trimmed for the forward primer and truncated to 250 bp using *usearch -*

fastx_truncate -stripleft 19 -trunclen 250, and quality filtered using *usearch -fastq_filter -*

fastq_maxee 1. Processed reads were dereplicated using *usearch -fastx_uniques -sizeout*, denoised

using *usearch -unoise3* (4). An ASV-table was produced with *usearch -otutab -zotus -*

sample_delim . -strand both, and analyzed in R as described in the main article. All plants are based

on conventional activated sludge except Dokhaven, which is configured with an Adsorption-

Belebungsverfahren (AB) process. b) ASVs were created based on raw V3-5 amplicon reads

obtained from Isazadeh *et al.* (2016)(56). Read 1 in SRA data were first oriented based on the

SILVA 138 SSURef NR99 database using *usearch -orient*. Sequences were trimmed for the

forward primer using *cutadapt -g CTACGGRNGGCWGC* (69), truncated to 250bp using *usearch -*

fastx_truncate -trunclen 250, and quality filtered using *usearch -fastq_filter -fastq_maxee 1*.

Processed reads were dereplicated using *usearch -fastx_uniques -sizeout*, denoised using *usearch -*

unoise3 (4). An ASV-table was produced with *usearch -otutab -zotus -sample_delim . -strand both*,

and analyzed in R as described in the main article. The data originate from samples collected at

eight Canadian WWTPs, which among others, differs in influent composition and treatment

processes, for details see Isazadeh *et al.* (2016)(56). Marieville, Farnham, and Cowansville:

Oxidation ditch. LaPrairie, Granby, Pincount, Salaberry: Conventional aeration. Vaudreuil:

Sequencing batch reactor carrousel, which employs the principles of oxidation ditch.

Figure S3:

Figure S3. Evaluation of the taxonomy assignment to FL-ASVs. A leave-one-out validation test was performed using the AutoTax processed SILVA 138 SSURef NR99 database without the *de novo* taxonomy. This database only contains high-quality, full-length 16S rRNA sequences and improper-taxonomic entries (e.g., entries containing the words uncultured, unknown, incertae sedis, and possible) were removed. The database was divided into a 10,000 sequence test set, and the

remaining database which was used for classification. a) Top hit identity distribution of the leave-one-out test data and for the FL-ASVs from this study mapped to the full SILVA 138 SSURef NR99 database (real data). The distributions are in the same range, confirming that the test data reflect actual data processed by AutoTax. b) Comparison of classification performance metrics for the AutoTax taxonomy assignment and commonly applied classifiers. The metrics used were the same as described by Edgar (13), but they were here applied to each taxonomic rank. Accuracy=TP/(K+OC); Misclassification rate=MC/K; Over-classification rate=OC/L; True positive rate=TP/K; Under-classification rate: UC/K, where K is the number of test sequences with a known taxonomy, L the number of test sequences without a known taxonomy, TP the number of names correctly assigned, MC the number of misclassification errors, OC the number overclassification errors, and UC the number of underclassification errors. The following classifiers were used: QIIME 2 feature-classifier with classify-consensus-blast (q2b), classify-consensus-vsearch (q2v), and classify-sklearn (q2sk) (5), USEARCH v.11 with SINTAX, NBC (Reimplementation of RDP), and USEARCH top hit (default parameters) (7, 13).

Figure S4. Novel taxa classified in the FL-ASV reference database. The number of taxa, which obtained names from the *de novo* taxonomy in different phyla. Archaeal phyla are highlighted in red and *de novo* phyla in blue.

Figure S5:

921 **Figure S5. Effect of the rare biosphere on amplicon classification.** a) The fraction of V1-3 ASVs

922 from activated sludge and anaerobic digester sample with a relative abundance higher than 0.001%

923 which was classified at the genus and species level using the FL-ASV reference database, the FL-

924 OTU expanded database, and commonly applied universal reference databases, including the

925 Autotax-processed SILVA 138 SSURef NR99 database.

Figure S6:

Figure S6. The specificity of taxonomy assignment using SINTAX and the FL-ASV and FL-OTU expanded reference database. a) Percentage of *in silico* bacterial ASVs, corresponding to amplicons produced using typical amplicon primers, that was correctly, wrongly, or not classified based on the FL-ASV database. b) Determining the effect of chimeras in the reference database. *In silico* bacterial ASVs were classified using the FL-ASV database and the FL-OTU expanded reference database.

SUPPLEMENTARY INFORMATION FOR

Generation of comprehensive ecosystems-specific reference databases with

species-level resolution by high-throughput full-length 16S rRNA gene

sequencing and automated taxonomy assignment (AutoTax)

Morten Simonsen Dueholm, Kasper Skytte Andersen, Simon Jon McIlroy, Jannie Munk Kristensen,

Erika Yashiro, Søren Michael Karst, Mads Albertsen, and Per Halkjær Nielsen

Center for Microbial Communities, Department of Chemistry and Bioscience, Aalborg University,

Aalborg, Denmark.

Detailed description of the AutoTax procedure:

Full-length 16S rRNA exact amplicon sequence variant (FL-ASV) calling

Full-length 16S rRNA gene sequences are first oriented according to the SILVA 138 SSURef Nr99

database (1) using the *usearch -orient* command. The sequences are then dereplicated using *usearch*

-fastx_uniques with the *-sizeout, -strand plus*, and *-threads 1* options. The *-sizeout* option put a size

annotation (how many times a unique sequence was observed) in the FASTA header. The *-strand*

plus option ensures that only correctly oriented sequences are considered when identifying identical

sequences. The dereplicated sequences are sorted and numbered based on the size annotation. The -

threads 1 option ensures that sequences with the same size are always sorted and numbered the

same way in the output. The dereplicated sequences are finally denoised to produce the FL-ASVs

using the *usearch -unoise3* command with the *-minsize 2* option. The *-minsize* options specify the

minimum abundance required for detecting an ASV. The default value for short-read amplicons is

8; however as the synthetic long-read sequences are independently amplified (due to the unique

molecular identifiers (UMIs) added before the PCR steps) and have a very low error-rate we can

lower the threshold to 2 with little risk of introducing sequences with errors, see "Calculation of FL-

ASV error-rate" below.

Preparation of chimera-filtered full-length 16S rRNA OTUs (FL-OTUs)

Dereplicated sequences from above are clustered at 99% sequence identity to create FL-OTUs using

usearch -cluster_smallmem command with the *-id 0.99, -maxrejects 0, -centroids,* and *-sortedby*

size options. The *-maxrejects 0* option specified a complete database search, thereby providing more

confident clusters. Potential chimeras are identified and extracted using the *usearch -uchime2_ref*

command with the *-strand plus, -mode sensitive*, and -chimeras options using the FL-ASVs as a

reference database. The sensitive mode was chosen as it captures more chimeras (on the expense of

more false positives), as these are detrimental to the final reference database. The FL-ASVs were

used as the reference database because it contains exact references for the abundant 16S rRNA

genes, which are most likely to form chimeras. Chimeras are finally removed to create the chimera-

filtered FL-OTUs using the *usearch -search_exact* command with the *-strand plus -dbnotmatched*

options, using the identified chimera sequences as the query, and the pre-filtered FL-OTUs as the

reference database.

Taxonomy assignment

For the taxonomy assignment, we first create two independent taxonomies. The first taxonomy is

based on the most recent version of the SILVA SSURef Nr99 database and reflects the current state

of microbial taxonomy. The second is a robust, although not necessarily evolutionary correct, *de*

novo taxonomy. The latter will be used as a taxonomic placeholder for taxonomic ranks without

information in the SILVA-based taxonomy.

The first step in both cases is the alignment of each FL-ASV to the global alignment in the SILVA

reference database in ARB-format using SINA. The aligned sequences are hereafter trimmed to

position 1048 to 41788 in the global SILVA alignment using the generic Linux command awk. The

purpose of this trimming is two-fold. Firstly, it ensures that flanking sequences do not result in

artificially low identities when the FL-ASVs are mapped to reference sequences in the SILVA

database produced with the popular 27F and 1391R primers (2). Secondly, it allows a more robust

de novo clustering (3). After trimming, gaps are removed from the alignment of FL-ASVs using the

usearch -fasta_stripgaps command. Finally, the FL-ASVs are sorted based on the FL-ASV

numbers in R.

The SILVA based taxonomy is obtained by mapping each of the trimmed FL-ASVs to the most

recent version of the SILVA SSURef Nr99 and the type strain database in FASTA format and

obtaining the taxonomic information from the closest relative as well as the percent identity.

Mappings are done using the *usearch -usearch_global* command with the *-maxrejects 0, -*

maxaccepts 0, -top_hit_only, -strand plus, -id 0, and *-blast6out* options for the complete SILVA

database and the *-maxrejects 0, -maxaccepts 0, -strand plus, -id 0.987,* and *-blast6out* for the type

strains. The *-maxrejects 0* and *-maxaccepts 0* options ensures that a comprehensive search is

performed. The *-top_hits_only* and *-id 0* options provide the best hit in the complete SILVA

database, whereas the *-id 0.987* without the *-top_hit_only* option provides all hits within the species-

level threshold. The output files from the SILVA reference mapping are loaded into R, and a data

frame with columns for FL-ASV number, percent identity, and SILVA taxonomy of the closest

relative is created. The taxonomy field is then split into the seven main taxonomic ranks from

kingdom to species. Fields that contain any of the following whole words are cleared: "uncultured",

"unknown", "unidentified", "incertae sedis", "metagenome", "bacterium", and "possible". In

addition, we replace all "candidatus" with "Ca" and all white spaces with underscores. Finally, all

characters except letters, numbers and period, dash, and underscore are removed.

1011

The taxonomy obtained from the closest relative in the SILVA database does not necessarily match

that of the FL-ASV. The taxonomy is therefore trimmed based on the percent identity between the

FL-ASV and its closest relative. For this trimming we use the taxonomic thresholds proposed by

Yarza *et al.* (4) (94.5% for genus-level, 86.5% for family-level, 82.0%, for order-level, 78.5% for

class-level, and 75.0% for phylum-level). Species-level classifications are only obtained from the

mapping against type strains, and no classifications are provided if reference sequences from more

than one species are within the species-level threshold.

1019

To generate a comprehensive *de novo* taxonomy, FL-ASVs are clustered based on the taxonomic

thresholds proposed by Yarza *et al.* (4) using the *usearch -cluster_smallmem* command with the *-id*

x, -maxrejects 0, -uc, and *-sortedby other* options. Where x represent the threshold for the given

taxonomic rank. The cluster_smallmem algorithm with the *-sortedby other* option clusters the FL-

ASVs based on when they appear in the input FASTA file, and the same clusters are therefore

formed even though additional FL-ASVs are appended to the FL-ASV-database in the future. We

confirmed this by processing only the first half of the FL-ASVs from this study, which provided

identical clustering. The *-maxrejects 0* ensures that the best reference is found among all clustered

sequences. The output file is a UCLUST-format tabbed text, which describes the clusters formed.

1029

The six UCLUST output files (species- to phylum-level) are loaded into R, and each are converted

into a data frame with two columns. The first column with clusters is named based on the

taxonomic clustering rank, and the second column with the input sequences is named as the

taxonomic rank below. Subsequently, the data frames are merged from species to phylum level.

This results in a comprehensive taxonomy, where the clustered centroids determine affiliation to the

taxonomic ranks above.

1036

1037 The SILVA-based taxonomy and the *de novo* taxonomy are finally merged in R by replacing empty

1038 fields in the SILVA-based taxonomy with the *de novo* taxonomic information. This results in a

1039 comprehensive taxonomy, where all FL-ASVs are described on all seven taxonomic ranks.

1040

1041 Merging of the two taxonomies results in a few cases where a taxon has more than one parent (e.g.,

1042 sequences from the same species affiliate more than one genus). In these cases, the taxonomy of the

1043 FL-ASV with the lowest ASV-number within the taxa is assigned to all members. A log file is

1044 written with details of all such cases.

1045

Calculation of FL-ASV error-rate:

The Unoise3 algorithm predicts which sequences represent true biological variants (ASVs) based on

their relative abundance compared to those of closely related sequences with one or a few errors.

Based on the error-rate profile of full-length 16S rRNA genes found in Karst et al. (5) and the

assumption that sequencing-errors are randomly distributed we can calculate the relative chance of

obtaining a correct ASV compared to an ASV with a single sequencing error when we only

consider sequence seen at least twice:

1053

1054 The probability of obtaining a correct sequence for FL-ASV$_n$ is the product of the probability of

1055 obtaining a sequence without any errors and relative abundance of FL-ASV$_n$, and for the sequence

1056 to be seen twice, we need to square this number.

1057

1058
$$P_{correct}(FL.ASV_n) = \left(P_{no\ error} \times P_{FL.ESV_n}\right)^2$$

1059

The probability of obtaining a sequence with a single error for FL-ASV$_n$ is the product of the

probability of obtaining a sequence with one error and relative abundance of FL-ASV$_n$. However, to

calculate the probability of picking a sequence with the same error, we need to divide this

probability with three times the length of FL-ASV$_n$ because the error could occur at any position in

the sequence and could be any of the three wrong nucleotides. The probability for seeing a specific

incorrect sequence twice therefore becomes:

1066

1067
$$P_{one\ error}(FL.ASV_n) = P_{one\ error} \times P_{FL.ASV_n} \times \frac{P_{one\ error} \times P_{FL.ASV_n}}{3 \times length_{FL.ASV_n}(bp)}$$

1068

1069   The relative chance of obtaining a correct FL-ASV compared to an FL-ASV with a single

1070   sequencing error is therefore:

1071

1072
$$\frac{P_{correct}(FL.ASV_n)}{P_{one\ error}(FL.ASV_n)} = \frac{P_{no\ error}^2 \times 3 \times length_{FL.ASV_n}(bp)}{P_{one\ error}^2} = \frac{0.112^2 \times 3 \times 1450}{0.209^2} = 1249$$

1073

1074   The results show that for each time we obtain an FL-ASV with a single error, we should expect

1075   approximately 1250 times more sequences of the correct FL-ASV. The difference should be large

1076   enough for Unoise3 to be able to remove almost all sequences with errors even though the errors are

1077   not entirely randomly distributed.

1078

1079

To confirm this, we reanalyzed the mock data used in Karst *et al.* (5). This data contained 7816

trimmed sequences (27f to 1492r) from a balanced mixture of DNA from *Escherichia coli* MG1655

(NC_000913), *Bacillus subtilis* subsp. subtilis str. 168 (NC_000964) and *P. aeruginosa* PAO1

(NC_002516). Denoising with *usearch -unoise3* command and the *-minsize 2* option produced 16

error-corrected sequences, of which 13 mapped perfectly to the 16S rRNA reference sequences, two

mapped with a single error, and one mapped with five errors. As the two FL-ASVs with single

errors were observed 14 and 4 times, respectively, it is highly unlikely that they represent

sequencing error based on the previous error-rate calculations. The same is holds for the FL-ASV

with five errors, although this sequence was observed only twice.

1089

1090   **References:**

1091   1.   Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glöckner FO. 2013.

1092   The SILVA ribosomal RNA gene database project: Improved data processing and web-based

1093   tools. Nucleic Acids Res 41:D590-6.

1094   2.   Klindworth A, Pruesse E, Schweer T, Peplies J, Quast C, Horn M, Glöckner FO. 2013.

1095   Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-

1096   generation sequencing-based diversity studies. Nucleic Acids Res 41:1–11.

1097   3.   Schloss PD. 2013. Secondary structure improves OTU assignments of 16S rRNA gene

1098   sequences. ISME J 7:457–460.

1099   4.   Yarza P, Yilmaz P, Pruesse E, Glöckner FO, Ludwig W, Schleifer K-H, Whitman WB,

1100   Euzéby J, Amann R, Rosselló-Móra R. 2014. Uniting the classification of cultured and

1101   uncultured bacteria and archaea using 16S rRNA gene sequences. Nat Rev Microbiol

1102   12:635–645.

1103   5.   Karst SM, Dueholm MS, McIlroy SJ, Kirkegaard RH, Nielsen PH, Albertsen M. 2018.

1104   Retrieval of a million high-quality, full-length microbial 16S and 18S rRNA gene sequences

1105   without primer bias. Nat Biotechnol 36:190–195.

1106

1107