JARA: 'Just Another Red-List Assessment'

Henning Winker[1,], Nathan Pacoureau[2] & Richard B. Sherley[3,*]*

[1] Joint Research Centre (JRC), European Commission, Via E. Fermi 2749, 21027 Ispra, Italy.

[2] Earth to Ocean Research Group, Department of Biological Sciences, Simon Fraser University, Burnaby, British

Columbia V5A1S6, Canada.

[3] Centre for Ecology and Conservation, College of Life and Environmental Sciences, University of Exeter,

Penryn Campus, Cornwall, TR10 9FE, UK.

*Correspondence: Henning Winker. E-mail: henning.winker@ec.europa.eu; Richard B. Sherley. Email:

r.sherley@exeter.ac.uk.

Abstract

Identifying species at risk of extinction is necessary to prioritise conservation efforts. The International

Union for Conservation of Nature's (IUCN) Red List of Threatened Species is the global standard for

quantifying extinction risk, with many species categorised according to population reduction thresholds.

We introduce the Bayesian state-space framework 'JARA' (Just Another Red-List Assessment). Designed

as decision-support tool, JARA allows both process error and uncertainty to be incorporated into IUCN

Red List assessments under criterion A. JARA is implemented via an R package that is designed to be easy

to use, rapid and widely applicable, so conservation practitioners can apply it to their own count or relative

abundance data. JARA outputs display easy to interpret graphics of the posterior probability of the

population trend against the corresponding IUCN Red List categories, as well as additional graphics to

describe the timeseries and model diagnostics. We illustrate JARA using three real-world examples: (1)

relative abundance indices for two elasmobranchs, Yellowspotted Skate *Leucoraja wallacei* and Whitespot

Smoothhound *Mustelus palumbes*; (2) a comparison of standardized abundance indices for Atlantic Blue

Marlin *Makaira nigricans* and (3) absolute abundance data for Cape Gannets *Morus capensis*. Finally, using

a simulation experiment, we demonstrate how JARA provides greater accuracy than two approaches

commonly used to assigning a Red List Status under criterion A. Tools like JARA can help further

standardise Red List evaluations, increasing objectivity and lowering the risk of misclassification, with

substantial benefits for global conservation efforts.

Introduction

Quantifying trends in population abundance is central to ecological research and to conservation

biology in particular, where understanding extinction risk is necessary to prioritise effort in the

face of ever-increasing biodiversity loss [1,2]. Although a number classification protocols exist to

assess a species' extinction risk [3], the International Union for the Conservation of Nature's

(IUCN) Red List of Threatened Species is viewed widely as the global standard [1,4].

To list a species in a threatened category (Critically Endangered [CR], Endangered [EN] or

Vulnerable [VU]) on the Red List, expert assessors consider the risks associated with both the

small-population paradigm and the declining population paradigm [5] under five assessment

criteria (A to E). However, quantitative information on species abundance – as a direct measure of

the status of a population – is often preferred over less direct measures of extinction risk, like

changes in habitat extent or quality [4,6]. Moreover, the IUCN guidelines allow for assignment to

a category (ranging from Least Concern [LC] to Extinct) based only on the criterion that produces

the highest estimated risk [4]. Therefore, species are often listed only on the basis of a reduction

in population size (Criterion A); as of today > 5,000 species are classified as threatened on this

basis alone [7]. Criterion A is thus considered to be the most widely used stand-alone criterion for

assigning a Red List status to a wide range of animal taxa, including mammals, birds, reptiles,

fishes and insects [2,4,8–10].

Although the IUCN Red List provides a set of unified quantitative decision rules for assigning

threatened categories based on population decline thresholds, data quality and analytical

approaches can differ vastly [4,6]. Estimating the decline between just two points in time remains

widely applied as a rapid assessment approach [6,11,12]. Even when model-predicted abundance

indices or population trajectories with associated uncertainty estimates exist, it appears common

practice to simply extract the two points that span the required IUCN assessment time frame of the

longest period between three generation lengths (GL) or 10 years and then calculate the decline as

the ratio between these two points. If the observation period is shorter than three generation

lengths, a simple extrapolation formula may be applied [11,12]. Fitting linear or log-linear

regression models to timeseries of abundance observations is a second widely-applied approach

[6,8,10,11]. An advantage of using a regression is that estimates of the significance of the estimated

slope, parameter uncertainty, and goodness-of-fit can be obtained, which is not possible with only

two points [6]. However, regression implicitly assumes that the "true" population follows a

constant deterministic trend, with all deviations from this trend attributable to statistically-

independent observation errors [13]. This assumption does not agree well with our understanding

of the real world. For example, a population diminished due to a severe drought year or a spike in

poaching will start from a lower state than would be expected from an average long-term trend. In

reality, animal population trajectories can diverge substantially from this deterministic expectation

due to persistent variations in environmental regimes or human impacts [14–16].

In contrast to deterministic regression models, state-space models provide a general framework

for analysing dynamical systems that considers process (year-to-year variation) and observation

(or reporting) error simultaneously [17,18]. State-space formulations add more biological realism

by assuming a Markovian process, such that the population size in the next time step is conditioned

on its current state, but independent of past and future states [18]. From a statistical perspective,

the observations are independent given the unobservable (latent) states of population. This can

help in preventing violation of independence in observation errors, which often manifest

themselves as serial residual correlations when fitting simple regressions to timeseries data. Given

these desirable properties, it is not surprising that there has been a rapid uptake of state-space

approaches for modelling population dynamics [17,19,20], including more recent applications of

quantifying population decline [6,21–23]. State-space models have also been specifically applied

in simulation studies that highlight the need for a more rigorous quantification of process and

observation uncertainties in the context of threat category (mis)classification [3,6,16].

Bayesian implementations of state-space models offer a powerful framework to improve the

characterization and communication of uncertainty during IUCN Red List assessments [24]. The

posterior probabilities provide an intuitive and transparent way to express uncertainty about

population declines to conservation practitioners [22,25], which can be translated directly into

probabilistic statements about a population falling into a threatened category [22,23]. However,

developing customized Bayesian state-space models can be technically demanding and time

consuming, especially when dealing with often case-specific, 'noisy' abundance data that are

subject to missing values, irregular spacing or multiple indices that are measured at different

scales. These issues may therefore dissuade some conservation practitioners and hamper broader

applications.

To address this, we have developed the Bayesian state-space framework 'Just Another Red-List

Assessment' (JARA). Designed as an easy to use, rapid and widely applicable decision-support

tool, JARA allows both process error and uncertainty to be incorporated into Red List assessments.

A key output of JARA is an easy to interpret graphic in which the probability distribution of the

population decline is displayed against the IUCN Red List categories, and where each category is

assigned a probability given process and observation uncertainty. To ensure a high degree of

transparency, reproducibility and to allow further development, we provide the R package JARA

on the global open-source platform GitHub (https://github.com/henning-winker/JARA), so that

JARA can be readily applied by conservation practitioners to their own count or relative abundance

data. On GitHub, we provide a number of worked examples. Here, we illustrate the main features

of JARA using three real-world examples and a simulation experiment to examine how accurately

JARA classifies complex population declines under criterion A, relative to the regression and

"two-points" approaches discussed above.

Methods

JARA is a generalized Bayesian state-space decision-support tool for trend analysis of abundance

indices with direct applications to IUCN Red List assessments. The name 'Just Another Red-List

Assessment' acknowledges JAGS ('Just Another Gibbs Sampler', [26]), the software used to run

the Bayesian state-space model application. The name reference, together with user-friendly R

interface and modulated coding structure follows the example of the new open source fisheries

stock assessment software 'Just Another Bayesian Biomass Assessment' (JABBA, [27]). JARA

enables analysis of one or multiple abundance indices simultaneously, where each index can have

a different scale, contain missing years and span different periods. JARA provides the option for

fitting relative abundance indices to estimate a mean trend or absolute abundance indices (from

e.g. different subpopulations) to produce a summed population trend for the total population.

State-Space Model Formulation of JARA

A central assumption of the state-space approach is that the abundance (I_t) trend follows a

Markovian process, such that I_t in year t will be conditioned on I_{t-1} in the previous year. For

generality, it is assumed that the underlying population trend follows a conventional exponential

growth model $I_{t+1} = I_t \lambda_t$, where λ_t is the growth rate in year t [18]. The growth rate λ_t can vary

annually to accommodate fluctuations in reproductive success and survival as a result of

environmental conditions, anthropogenic activity or other latent (unobservable) impacts. State-

space models are hierarchical models that explicitly decompose an observed time-series into a

process variation and an observation error component [21]. On the log scale, the process equation

becomes $\mu_{t+1} = \mu_t + r_t$, where $\mu_t = \log(I_t)$ and $r_t = \log(\lambda_t)$ is the year-to-year rate of change,

with variations in log-growth rates following a random normal walk: $r_t \sim Normal(\bar{r}_p, \sigma_\eta^2)$, given

the estimable process error variance σ_η^2 and the estimable mean population rate of change $\bar{r}_p$ (i.e.

the underlying trend). Because the process error is lognormally distributed, r_t is adjusted for

lognormal bias by subtracting half the variance (otherwise the stochastic lognormal error induces

a small positive bias) at each time step: $r_t = \bar{r}_p + r_{t-1} - (0.5 * \sigma_\eta^2)$ [28]. The corresponding

observation equation is of the form $\log(y_t) = \mu_t + \epsilon_t$, where y_t denotes the abundance

observation for year t, ϵ_t is observation residual for year t, which is assumed to be normally

distributed on log-scale $\varepsilon_t \sim Normal(0, \sigma_\varepsilon^2)$ as a function of the observation variance σ_ε^2.

The model notation for relative abundance indices builds on the approach in JABBA for averaging

relative abundance indices [27] and assumes that the mean underlying abundance trend is an

unobservable state variable. The corresponding observation equation is then modified, such that

$\log(y_{t,i}) = \mu_t + \log(q_i) + \epsilon_{t,i}$, where $y_{t,i}$ is the relative abundance value for year t and index i,

μ_t is the natural logarithm of the mean abundance trend, $\varepsilon_{t,i}$ is the lognormal observation error

term for index i and year t, and q_i is a scaling parameter for index i. The abundance index with the

chronologically oldest record is taken as a reference index by fixing $q_1 = 1$ and the other indices

are scaled to this reference index, respectively, with $q_{2,\dots,n}$ being estimable model parameters. The

estimated posterior of the population trend for year t is then given by $I_{p,t} = \exp(\mu_t)$.

To estimate a total abundance trajectory for the 'global' population from multiple timeseries of

absolute abundance observations, we assume that each absolute abundance index represents a

'subpopulation' that may increase or decline independently from other subpopulations. The

process equation is therefore modified to $\mu_{t,i} = \mu_{t,i} + r_{t,i}$, where $r_{t,i} = \log(\lambda_{t,i})$ is the year-to-

year rate of change specific to index i that is assumed to vary around $\bar{r}_i$ to represent the underlying

mean rate of change for the subpopulation (instead of a global population mean $\bar{r}_p$), but with a

process variance σ_η^2 that is common to all subpopulations: $r_{t,i} \sim Normal(\bar{r}_i, \sigma_\eta^2)$. Again, we

adjusted $r_{t,i}$ for lognormal bias by subtracting half the process variance at each time step:

$r_t = \bar{r}_i + r_{t-1,i} - (0.5 * \sigma_\eta^2)$ [28]. The corresponding observation equation is adjusted to

$\log(y_{t,i}) = \mu_{t,i} + \epsilon_{t,i}$, so that abundance trend $\mu_{t,i}$ and the error terms $\varepsilon_{t,i} \sim Normal(0, \sigma_{\varepsilon_i}^2)$ now

become the specific to subpopulation i. The estimated posterior of the global population trajectory

$\hat{I}_{p,t}$ for year t is then computed from the sum of all individual subpopulation trajectory posteriors,

after correcting them for the lognormal bias by subtracting half their variance [28,29]: $\hat{I}_{p,t} =$

$\sum_i \exp(\mu_{t,i} - 0.5 * \sigma^2_{\mu_{t,i}})$. Although the sum of multiple lognormal distributions are not themselves

lognormally distributed [30], such that the posterior median or uncorrected mean of $\hat{I}_{p,t}$ would

yield inflated estimates (particularly) when population counts are large [29], our corrected mean

is unbiased [31].

Bayesian Framework

JARA is run from the statistical environment R [32] and executed in JAGS [26], using a wrapper

function from the R library 'r2jags' [33]. The Bayesian posterior distributions are estimated by

means of Markov Chain Monte Carlo (MCMC) simulation. In JAGS, all estimable hyper-

parameters are assigned to a prior distribution. JARA uses vague (uninformative) prior

distributions throughout, so all inferences are drawn from the information in the data. The

estimation of annual growth rate deviates r_t, is implemented through hierarchical priors [34],

where r_t is informed by the population mean $\bar{r}_p$ for relative abundance indices and $r_{t,i}$ is informed

by $\bar{r}_i$ for absolute abundance indices. Vague normal priors with a mean of 0 and variance of 1000,

$Normal(0,1000)$, are assumed for both $\bar{r}_p$ or $\bar{r}_i$. The initial population size in the first year $I_{t=1,i}$

is drawn in log-space from a normal distribution with the mean equal to the log of the first available

count $y_{t=1,i}$ and a standard deviation of 1000. Priors for the process variance (σ^2_η) can be either

fixed or estimated. If estimated (default), the process variance prior is implemented via a vague

inverse-gamma distribution by setting both scaling parameters to 0.001: $\sigma^2 \sim 1/$

$gamma(0.001,0.001)$ [35–37], which yields an approximately uniform prior on the log scale [27].

The total observation variance σ^2_ε can be separated into three components: (1) externally derived

standard errors $\sigma^2_{SE_{y,i}}$ for each abundance index *i*, (2) a fixed input variance σ^2_{fix} and (3) estimable

variance σ_{est}^2, where the prior for σ_{est}^2 assumes an uninformative inverse-gamma distribution with

both scaling parameters set to 0.001. All three variance components are additive in their squared

form [38], so the total observation variance for abundance index i and year y is: $\sigma_{\varepsilon_{y,i}}^2 = \sigma_{SE_{y,i}}^2 +$

$\sigma_{fix}^2 + \sigma_{est}^2$ [39]. Each variance component can be individually switched on or off by the user

[27]. Adding a fixed observation error is common practice to account for additional sampling error

associated with abundance indices [40]. Setting a minimum plausible observation error in this way

informs the estimate of the process variance as a portion of total variance is assigned *a priori* to

the observation variance [27] and helps to increase model stability and convergence of state-space

models [41].

Estimating probabilities of population decline

A posterior probability for the percentage change (*C%*) associated with each abundance index can

be conveniently calculated from the posteriors of $\hat{I}_{p,y}$, the model predicted population trajectory.

If $\hat{I}_{p,y}$ represents a longer time span than the assumed three GL, *C%* is automatically calculated as

the difference between the median of three years around the final observed data point T, and a

three-year median around the year corresponding to $T - (3 \times GL)$. The year $T + 1$ is always

projected to obtain a three-year average around T to reduce the influence of short-term fluctuations

[42]. When the span of $\hat{I}_t$ is $< 3 \times$ GL, JARA projects forward, by passing the number of desired

future years without observations to the state-space model until $\hat{I}_t > (3 \times GL) + 2$. The

projections for a single population or multiple subpopulations i are based on all possible posterior

realizations of $r_{t,i}$ across all T years in the observed timeseries: $\tilde{r}_i = \frac{1}{T}\sum_{t=1}^{T} r_{t,i}$.

Model Diagnostics

To evaluate the model fit, JARA provides the user with four plots. The first illustrates the unscaled

input data and uncertainty estimates around each observation in the form 95% Confidence Intervals

(Figure 1a-b). The second shows the observed and predicted abundance values for each timeseries

together with the 95% posterior predictive credibility intervals (Figure 1c-d). The third shows

individual fits on the log-scale, as well as the 95% Bayesian credible intervals (CI) derived from

the observation variance $\sigma^2_{\varepsilon_{y,i}}$ (Figure 2). The fourth is a residual plot (Figure 1e-f) to illustrate

potential data conflict when fitting multiple timeseries [27]. This plot includes: (1) colour-coded

lognormal residuals of observed versus predicted abundance indices i, (2) boxplots indicating the

median and quantiles of all residuals available for each year; the area of each box indicates the

strength of the discrepancy between the abundance indices (larger box means higher degree of

conflicting information), and (3) a loess smoother through all residuals which highlights

systematically auto-correlated residual patterns (Figure 1e-f). In addition, the Root-Mean-

Squared-Error (RMSE) and the deviance information criterion (DIC) are provided for the

comparison of goodness-of-fit and model selection purposes respectively. Convergence of the

MCMC chains is diagnosed using the 'coda' package [43], adopting minimal thresholds of p =

0.05 for Heidelberger and Welch [44] and Geweke's [45] diagnostics called from R. Unless

otherwise specified, JARA reports medians and 95% credible intervals (CI).

JARA decision support plots

To facilitate decision-making, JARA routinely produces four key plots (Figures 3–5) for each input

dataset showing: (1) the combined state-space model fit and 95% CI to abundance timeseries

where multiple indices relative abundance (e.g. Figure 4a) and stock assessment results (Figure

4b) are used, or the individual state-space models fits to each count dataset for absolute abundance

indices (e.g. individual colony counts for Cape Gannet; Figure 5a); (2) the overall observed and

projected (±95% CI) population trajectory over three GL (e.g. Figure 3a and b); (3) the median

and posterior probabilities for the percentage annual population change calculated from all the

observed data, and from each of the most recent 1 GL, 2 GL, and 3 GL (depending on the length

of the observed timeseries), shown relative to a stable population ($\%C = 0$) (e.g. Figure 5c); and

(4) how the posterior distribution for the percentage change in abundance ($\%C$) over 3 GL aligns

against the thresholds for the Red List categories (Critically Endangered CR, Endangered EN,

Vulnerable VU , Near Threatened NT or Least Concern LC) under criteria A2–A4 (e.g. Figure 4e

and f) or under A1. In addition, JARA can be used to undertake a retrospective analysis through

the sequential removal of terminal years and subsequent forward projections to attain 3 GL

("retrospective peel"). This enables to user to identify points in time where the $\%C$ crosses over

into different Red List categories (Figure 5f).

Simulation experiment

We conducted a simulation experiment to compare the performance of JARA against two

conventional approaches suggested in the IUCN Red List guidelines: (1) the 'two-point' approach

that calculates $\%C$ between two years with observations that are $3 \times$ GL apart and (2) regression

analysis that extrapolates $\%C$ from the estimated slope over a given assessment horizon of $3 \times$ GL

[11]. We first used an Operating Model (OM) to generate a 'true' population trend and an

'observed' abundance index (Figure S1). The observed abundance index is then passed to the three

estimation models (EMs: 'Two-Points', 'Regression', 'JARA') to determine the likely threat

category (Figure 6).

We modelled the 'true' population numbers using a log-linear Markovian process, such that

$\log(N_t) = \log(N_{t-1}) + r_t$, where r_t determines the annual rate of change in the population

numbers N_t. We assumed that r_t can be decomposed into three processes: (i) an underlying mean

population trend $\bar{r}$, (ii) stochastic annual variation in the population v_t and (iii) annual variation in

anthropogenic impact τ_t, such that $r_t = \bar{r} + v_t - \tau_t$ (Fig. S1). The process error deviations

associated with v_t and τ_t were assumed to be normally distributed and temporally autocorrelated

[15,16] by implementing first-order autoregressive functions (AR1) of the form: $v_t = v_{t-1}\phi_v +$

$\delta_t\sqrt{1 - \phi_v^2}$ and $\tau_t = v_{t-1}\phi_\tau + \gamma_t\sqrt{1 - \phi_\tau^2}$ where ϕ_v and ϕ_t are the AR1 autocorrelation

coefficients and $\delta_t \sim Normal(0, \sigma_\delta^2)$ and $\gamma_t \sim Normal(0, \sigma_\gamma^2)$ are normally distributed random

variables representing uncorrelated process errors with variances σ_δ^2 and σ_γ^2, respectively [46]. The

observed abundance index I_y is then generated as a function of the 'true' population numbers and

a lognormal observation error of the form: $I_y = N_t e^{\varepsilon_t}$, where $\varepsilon_t \sim Normal(0, \sigma_\varepsilon^2)$ and σ_ε^2 is the an

observation variance σ_ε^2. A generic R function of the population simulator is provided in the

Supplementary Information.

We simulated 200 random sets of population timeseries of 35 years with an assumed assessment

horizon of 30 years corresponding to a GL of 10 years (Figure 6). The initial population size was

$N_1 = 10,000$ and random deviates of $\bar{r}$, describing the underlying trend, were randomly generated

using a uniform distribution between -0.05 and 0 to cover a wide range of population trends and

associated *%C* (Figure S1-2). We set the standard deviation for the stochastic variation to a

moderate value of $\sigma_v = 0.1$ with a process error autocorrelation of $\phi_v = 0.3$. This choice of values

was intended to describe a population where annual reproductive success is fairly variable and

autocorrelation arises from individual age-classes of varying strengths growing through the

population. By comparison, the variation in anthropogenic impact was assumed to be less variable

$\sigma_{\upsilon} = 0.05$ and more persistent over longer, regime-like periods by assuming a fairly large

autocorrelation coefficient of $\phi_{\tau} = 0.7$. The observation error for the observed abundance was set

to $\sigma_{\varepsilon} = 0.15$ (i.e. CV ~ 15%). The resulting simulations produced a wide range of plausible

population trajectories (Figure 6).

We evaluated model performances in terms of the accuracy of the estimated 'true' %C over 3 GL

and the proportion of correctly classified threat categories according to the IUCN Red List A2

decline criterion. Accordingly, the 'true' %C over 30 years (3 GL) was calculated as

%$C = (N_{35}/N_6 - 1) \times 100$. For the 'Two-Point' EM we calculated %$\hat{C} = (I_{35}/I_6 - 1) \times 100$.

For the 'Regression' EM we closely followed the IUCN guidelines by: (i) fitting a log-linear

regression of the form $\log(I_y) = \alpha + \beta y$ to all observations I_{1-35}, (ii) predicting the expected

values $\hat{I}_{1-35}$, (iii) and then extrapolating the expected %$\hat{C}$ over 3 GL as: %$\hat{C} = (\hat{I}_{35}/\hat{I}_1)^{(3GL/n_y)} -$

$1) \times 100$ where n_y is the number of years. As described above (*Estimating probabilities of*

population change), for the JARA EM the %$\hat{C}$ was calculated as the median of the posterior %C.

Accuracy and bias were calculated using Median Absolute Error (MAE) and Median Error (ME),

where the error is the difference between the estimated %$\hat{C}$ and the 'true' %C.

Results and Discussion

Case study applications

We illustrate applications of JARA using the following worked examples for different abundance

data types: (1) multiple relative abundance indices (1991–2017) from South African scientific

demersal trawl surveys for two elasmobranchs, Yellowspotted Skate *Leucoraja wallacei* and

Whitespot Smoothhound *Mustelus palumbes*; (2) a comparison of model-predicted biomass

trajectories from the 2018 Atlantic Blue Marlin *Makaira nigricans* stock assessments and

standardized catch-per-unit-effort (CPUE) abundance indices from various fishing fleet, which

were used as input into the assessment model and (3) census data for a large seabird, the Cape

Gannet *Morus capensis*, in the form of total breeding size estimates for its six global breeding

localities in Namibia and South Africa (1979–2017). All worked examples and input data are

available on GitHub (https://github.com/henning-winker/JARA).

(1) Whitespot Smoothhound and Yellowspotted Skate are both endemic to southern Africa and

had been previously listed as Data Deficient by the IUCN [47,48]. During the 2018 Southern

African IUCN Red List workshop, hosted by the IUCN Shark Specialist Group (SSG), both species

were reassessed using JARA as a decision-support tool. Based on available life-history

information, the GLs were empirically estimated as 14 years for Whitespot smoothhound and 12

years for Yellowspotted Skate. Long-term abundance indices were obtained from analysis of

demersal research trawl survey data that were conducted with alternating trawl gear configurations

(old/new gear) during summer along South Africa's west coast (1991–2017) and during autumn

and spring along the south coast (1991–2016) by the Department of Agriculture, Forestry and

Fisheries. Yellowspotted Skate was only encountered during South Coast Surveys. Annual density

estimates (kg per nm^2 area swept) and associated standard errors on log-scale were estimated using

a geostatistical delta-GLMM [49,50]. We only considered the period from 1991 onwards due to

evident learning effects for identifying Chondrichthyes during initial survey years. The estimated

abundance indices were separately derived for each gear type, survey (coast/season). The input

data are characteristic of multiple indices with irregular spacing due to missing survey years

(Figure 1 and 2), and are estimated at varying spatial scales. They are therefore treated as relative

abundance indices.

JARA combined six abundance indices for Whitespot Smoothhound (Figure 1a) and four

abundance indices for Yellowspotted Skate (Figure 1b). Both sets of abundance indices spanned

~2 GL, so JARA automatically projected over ~ 1 GL to provide %C estimates of +20.2% with

95% of the posterior falling in LC for Whitespot Smoothhound and −47.1% for Yellowspotted

Skate, with EN the best-supported category (Figure 3). The plots of the percentage annual

population change (Figure 3c and d) nicely illustrate the contrasting situations for these two

species, with Whitespot Smoothhound showing an improving population growth rate in the most

recent 1 GL (relative to the whole dataset, Figure 3c), while for Yellowspotted Skate the decline

appears to have worsened over the last 1 GL (Figure 3d).

(2) The Blue Marlin, one of the most iconic gamefishes, is a highly migratory billfish species

distributed circumglobally in mostly tropical waters. The last IUCN assessment in 2010 classified

the species globally as VU [51]. The Atlantic population was assessed by fitting linear regressions

to biomass trend estimates from stock assessments undertaken by the International Commission

for the Conservation of Atlantic Tunas (ICCAT). GL was assumed to be up to 6 years,

corresponding to an IUCN trend assessment horizon of 18 years (3 x GL). Here, we use the 2018

ICCAT stock assessment of Atlantic Blue Marlin to illustrate how JARA can objectively convert

such stock assessment outputs into corresponding IUCN Red List decline thresholds and threat

categories recommendations. The assessment output used as input for JARA is the median of the

estimated biomass for the period 1950–2016 across all accepted candidate models (Figure 4b).

Uncertainty was represented by annual standard errors (SEs) of the log-biomass estimates [52].

We then compared the JARA results with a complementary JARA analysis of multiple CPUE

indices that served as input data into the 2018 stock assessment. This CPUE dataset comprised

standardized CPUE indices (+SEs) from ten different fishing fleets that covered different periods

between 1959 and 2016 (Figure 4a).

JARA estimated similar median $\%C$ for the Atlantic Blue Marlin whether using the multiple CPUE

indices (-29.3%) or the stock biomass (-21.8%) as input data. In both cases, the best-supported

category was LC, but uncertainty was (unsurprisingly) lower when using the biomass estimates

from the stock assessment (80.7% of the posterior in LC) versus the much more variable and

partially conflicting CPUE indices (51.4% in LC; Figure 4e and f). Although either input series

would likely lead to the same outcome (a classification of LC) if JARA were used a decision-

support tool to assess Atlantic Blue Marlin, the analysis of the stock assessment data would result

in greater confidence in the assessment's conclusion.

(3) The Cape Gannet is a large seabird endemic to southern Africa, where it breeds during the

austral spring and summer at six islands. Aerial photographs and on the ground measurements of

nesting density at all six breeding colonies have been used by the Department of Environmental

Affairs (South Africa) and the Ministry of Fisheries and Marine Resources (Namibia) to estimate

the total breeding population since the summer of 1956/57 [53]. Counts were sporadic until

1978/79 but undertaken approximately annually thereafter (Figure 5). Cape Gannets generally

have high adult survival, strong breeding site fidelity [54] and low reproductive success (≤ 0.82

chicks fledged per pair annually; [55]. Accordingly, their GL is estimated as 20.2 years [56] and

we use annual count data from 1956/57 to 2016/2017, which spans 3 GL or the recommended Red

List assessment period. Of a possible 366 annual counts between 1956/57 to 2016/2017, 173 were

not made for various reasons. Thus, the Cape Gannet dataset is chosen an example of multiple

absolute abundance indices (with missing data) that in sum represents the trend of the global

population.

The output for Cape Gannet nicely exemplifies the decision-support element of JARA. Based on

only the median $\%C$ (−51.3%), the species meets the criteria for classification as EN (Figure 5d).

However, the uncertainty spans LC to EN; assessors may, therefore, also want to consider that

while 55.6% of the posterior probability distribution falls into EN, 41.8% also falls into VU (Figure

5d). Moreover, the percentage annual population change plots show that the decline has slowed

and then accelerated over the last 2 GL (Figure 5c), while the retrospective plots show that the

threshold for EN was first exceeded in 2013, but the situation has not worsened markedly to 2016.

Assessors could then make an informed decision on whether to list the Cape Gannet as VU or EN,

depending on how risk prone or adverse they wanted to be in their assessment and the presence of

other mitigating factors.

Simulation experiment results

Median Absolute Errors (MAEs) between the estimated and 'true' percentage change over 3

generation lengths (GL) using JARA as the EM was 3.63%. The Regression EM (7.87%) and Two-

Point EM (10.96%) had MAEs >2 and >3 times larger, respectively, showing a distinct decrease

in the accuracy of the estimated $\%C$ in comparison to the JARA EM (Figure 7). In addition,

although all three EMs showed a tendency towards a positive bias (Median Errors > 0), this bias

was ~3.7 times larger for the regression EM (0.71%) and ~12 times larger for the Two-Point EM

(2.26%) than for JARA (0.19%), which was approximately unbiased. Moreover, the median %C

from JARA correctly identified the 'true' IUCN threat category in 89.5% of cases; > 10% more

than the regression EM (which was based on the current IUCN guidelines) and >20% more than

the Two-Point EM (Figure 7). Although these might seem like relatively small improvements,

misclassification on the IUCN Red List can have real consequences for species conservation

[7,57], particularly resource allocation if species move in and out of the threatened categories [58].

There will be many real-world cases where the observed or estimated decline is close to a Red List

category threshold (e.g. >50% decline for EN), as was the case for Yellowspotted Skate (median

%C −47.1%, Figure 3f) and Cape Gannet (median %C −51.3%, Figure 5d) in our examples. Not

only would we would expect JARA to indicate the 'true' IUCN threat category more reliably in

these sorts of situations, but the additional information provided by JARA – particularly the plot

of the probabilities that the rate of change falls within the IUCN Red List categories (e.g. Figure

5d) – offers assessors insight into the weight of evidence underpinning their assignment.

Current limitations and real-world applications

Although still being refined, JARA has already been used to support expert decision making in the

assessment of a terrestrial mammal [59], a number of elasmobranchs [31,60] and two seabirds

[31,53]. We designed JARA with the aim of supporting the global effort to provide robust,

consistent, and transparent assessments for all species under the IUCN Red List criteria [24]. And

we designed it to be free, open-source, and easy to use, to support access for conservation

practitioners, researchers and assessors working in places with limited opportunities to access

proprietary software [12]. While we certainly hope that more conservation practitioners will use

JARA, we caution against applying it to unfamiliar timeseries or blindly accepting the status

suggested by decision-support tools [12]. Undertaking Red List assessments requires assessors to

understand their target species and the rules specified in the Red List guidelines. The data used for

assessments should be proofed and verified by specialists (e.g. an IUCN Specialist Group) to

ensure their reliability, users should make informed ecological judgments regarding the length of

the timeseries relative to a species' GL when assessing against criterion A, and should remain

cognisant of biases in data availability or survey effort when making their recommendations

[11,12,24]. Moreover, JARA has been designed to be used as a decision-support tool and its

outputs are not necessarily indicative of a final classification of extinction risk [24]. Rather,

assessors should use tools like JARA to make an informed interpretation based on all of the Red

List guidelines before coming to a decision on the final assessment outcome [12,24]. Nevertheless,

when coupled with expert oversight, JARA has wide potential applicability to other studies of

wildlife populations threatened with extinction [61,62].

Acknowledgments

H.W. and R.B.S. thank Marc Kéry and Michael Schaub for introducing us to Bayesian state-space

models, and we thank Nick Dulvy for helpful conversations about the Red List process during the

development of JARA. RBS was supported by the Pew Fellows Program in Marine Conservation

at The Pew Charitable Trusts. The views expressed are those of the authors and do not necessarily

reflect the views of The Pew Charitable Trusts. N.P. was funded by the Shark Conservation Fund,

a philanthropic collaborative pooling expertise and resources to meet the threats facing the world's

sharks and rays. The Shark Conservation Fund is a project of Rockefeller Philanthropy Advisors.

References

1.   Hoffmann M, Brooks TM, Da Fonseca GAB, Gascon C, Hawkins AFA, James RE, et al.

Conservation planning and the IUCN Red List. Endanger Species Res. 2008; 6: 113–125.

doi:10.3354/esr00087

2.   Butchart SHM, Stattersfield AJ, Bennun LA, Shutes SM, Akçakaya HR, Baillie JEM, et

al. Measuring Global Trends in the Status of Biodiversity: Red List Indices for Birds. PLoS

Biol. 2004; 2: e383. doi:10.1371/journal.pbio.0020383

3.   Regan TJ, Taylor BL, Thompson GG, Cochrane JF, Ralls K, Runge MC, et al. Testing

decision rules for categorizing species' extinction risk to help develop quantitative listing

criteria for the U.S. endangered species act. Conserv Biol. 2013; 27: 821–831.

doi:10.1111/cobi.12055

4.   Mace GM, Collar NJ, Gaston KJ, Hilton-Taylor C, Akçakaya HR, Leader-Williams N, et

al. Quantification of extinction risk: IUCN's system for classifying threatened species.

Conserv Biology. 2008; 22: 1424–1442. doi:10.1111/j.1523-1739.2008.01044.x

5.   Caughley G. Directions in Conservation Biology. J Anim Ecol. 1994; 63: 215–244.

6.   Wilson HB, Kendall BE, Possingham HP. Variability in Population Abundance and the

Classification of Extinction Risk. Conserv Biol. 2011; 25: 747–757. doi:10.1111/j.1523-

1739.2011.01671.x

7.   Rueda-Cediel P, Anderson KE, Regan TJ, Regan HM. Effects of uncertainty and variability

on population declines and IUCN Red List classifications. Conserv Biol. 2018; 32: 916–

925. doi:10.1111/cobi.13081

8.   Dulvy NK, Jennings S, Rogers SI, Maxwell DL. Threat and decline in fishes: an indicator

of marine biodiversity. Can J Fish Aquat Sci. 2006; 63: 1267–1275. doi:10.1139/f06-035

9.   Seminoff JA, Shanker K. Marine turtles and IUCN Red Listing: A review of the process,

the pitfalls, and novel assessment approaches. J Exp Mar Biol Ecol. 2008; 356: 52–68.

doi:10.1016/j.jembe.2007.12.007

10.   Fox R, Harrower CA, Bell JR, Shortall CR, Middlebrook I, Wilson RJ. Insect population

trends and the IUCN Red List process. J Insect Conserv. 2018; 23: 269–278.

doi:10.1007/s10841-018-0117-1

11.   IUCN Standards and Petitions Subcommittee. Guidelines for Using the IUCN Red List

Categories and Criteria. IUCN, Gland, Switzerland and Cambridge, UK; 2019.

12.   Lee CKF, Keith DA, Nicholson E, Murray NJ. REDLISTR: Tools for the IUCN Red Lists

of Ecosystems and Threatened Species in R. Ecography. 2019; 42: 1–6.

doi:10.1111/ecog.04143

13.   Hilborn R, Mangel M. The Ecological Detective: Confronting Models with Data.

Princeton, NJ: Princeton University Press; 1997.

14.   D'Eon-Eggertson F, Dulvy NK, Peterman RM. Reliable Identification of Declining

Populations in an Uncertain World. Conserv Lett. 2015; 8: 86–96. doi:10.1111/conl.12123

15.   Keith D, Akçakaya HR, Butchart SHM, Collen B, Dulvy NK, Holmes EE, et al. Temporal

correlations in population trends: Conservation implications from time-series analysis of

diverse animal taxa. Biol Conserv. 2015; 192: 247–257. doi:10.1016/j.biocon.2015.09.021

16.   Connors BM, Cooper AB, Peterman RM, Dulvy NK. The false classification of extinction

risk in noisy environments. Proc Biol Sci. 2014; 281: 20132935.

doi:10.1098/rspb.2013.2935

17.   de Valpine P, Hastings A. Fitting Population Models Incorporating Process Noise and

Observation Error. Ecol Monogr. 2002; 72: 57–76. doi:10.1890/0012-

9615(2002)072[0057:fpmipn]2.0.co;2

18.   Kéry M, Schaub M. Bayesian population analysis using WinBUGS: A hierarchical

perspective. Oxford: Academic Press; 2012.

19.   Buckland ST, Newman KB, Thomas L, Koesters NB. State-space models for the dynamics

of wild animal populations. Ecol Modell. 2004; 171: 157–175.

doi:10.1016/j.ecolmodel.2003.08.002

20.   Thorson JT, Minto C. Mixed effects: a unifying framework for statistical modelling in

fisheries biology. ICES J Mar Sci. 2015; 72: 1245–1256. doi:10.1093/icesjms/fsu213.

doi:10.1093/icesjms/fst048

21.   Simmons RE, Kolberg H, Braby R, Erni B. Declines in migrant shorebird populations from

a winter-quarter perspective. Conserv Biol. 2015; 29: 877–887. doi:10.1111/cobi.12493

22.   Bauer H, Chapron G, Nowell K, Henschel P, Funston P, Hunter LTB, et al. Lion (Panthera

leo) populations are declining rapidly across Africa, except in intensively managed areas.

Proc Natl Acad Sci U S A. 2015; 112: 14894–14899. doi:10.1073/pnas.1500664112

23.   Boyd C, DeMaster DP, Waples RS, Ward EJ, Taylor BL. Consistent Extinction Risk

Assessment under the U.S. Endangered Species Act. Conserv Lett. 2017; 10: 328–336.

doi:10.1111/conl.12269

24.   Sherley RB, Winker H, Rigby CL, Kyne PM, Pollom R, Pacoureau N, et al. Estimating

IUCN Red List population reduction: JARA—A decision-support tool applied to pelagic

sharks. Conserv Lett. 2020; 13: e12688. doi:10.1111/conl.12688

25.   Sherley RB, Barham BJ, Barham PJ, Campbell KJ, Crawford RJM, Grigg J, et al. Bayesian

inference reveals positive but subtle effects of experimental fishery closures on marine

predator demographics. Proc Biol Sci. 2018; 285: 20172443. doi:10.1098/rspb.2017.2443

26.   Plummer M. JAGS: A Program for Analysis of Bayesian Graphical Models using Gibbs

511   Sampling. Vienna: 3rd International Workshop on Distributed Statistical Computing (DSC

512   2003); 2003.

513   27.   Winker H, Carvalho F, Kapur M. JABBA: Just Another Bayesian Biomass Assessment.

514   Fish Res. 2018; 204: 275–288. doi:10.1016/j.fishres.2018.03.010

515   28.   Methot RD, Taylor IG. Adjusting for bias due to variability of estimated recruitments in

516   fisheries assessment models. Can J Fish Aquat Sci. 2011; 68: 1744–1760.

517   29.   Che-Castaldo C, Jenouvrier S, Youngflesh C, Shoemaker KT, Humphries G, McDowall P,

518   et al. Pan-Antarctic analysis aggregating spatial estimates of Adélie penguin abundance

519   reveals   robust   dynamics   despite   stochastic   noise.   Nat   Commun.   2017;   8:   832.

520   doi:10.1038/s41467-017-00890-0

521   30.   Dufresne D. The log-normal approximation in financial and other computations. Adv Appl

522   Probab. 2004; 36: 747–773.

523   31.   Sherley RB, Crawford RJM, de Blocq AD, Dyer BM, Geldenhuys D, Hagen C, et al. The

524   conservation status and population decline of the African penguin deconstructed in space

525   and time. Ecol Evol. 2020; Forthcoming.

526   32.   R Core Team. R: A language and environment for statistical computing. Vienna: R

527   Foundation for Statistical Computing; 2018. Available: http://www.r-project.org/

528   33.   Su Y-S, Yajima M. R2jags: A Package for Running jags from R; 2012.

529   34.   Jiao Y, Hayes C, Corte E. Hierarchical Bayesian approach for population dynamics

530   modelling of fish complexes without species-specific data. ICES J Mar Sci. 2009; 66: 367–

531   377. doi:10.1093/icesjms/fsn162

35.   Chaloupka M, Balazs G. Using Bayesian state-space modelling to assess the recovery and

harvest potential of the Hawaiian green sea turtle stock. Ecol Modell. 2007; 205: 93–109.

doi:10.1016/j.ecolmodel.2007.02.010

36.   Brodziak J, Ishimura G. Development of Bayesian production models for assessing the

North Pacific swordfish population. Fish Sci. 2012; 77: 23–34. doi:10.1007/s12562-010-

0300-0

37.   Carvalho F, Ahrens R, Murie D, Ponciano JM, Aires-da-Silva A, Maunder MN, et al.

Incorporating specific change points in catchability in fisheries stock assessment models:

An alternative approach applied to the blue shark (Prionace glauca) stock in the south

Atlantic Ocean. Fish Res. 2014; 154: 135–146. doi:10.1016/j.fishres.2014.01.022

38.   Francis RICC, Hurst RJ, Renwick JA. Quantifying annual variation in catchability for

commercial and research fishing. Fish Bull. 2003; 101: 293–304.

39.   Conn PB. Hierarchical analysis of multiple noisy abundance indices. Can J Fish Aquat Sci.

2010; 67: 108–120. doi:10.1139/F09-175

40.   Maunder MN, Piner KR. Dealing with data conflicts in statistical inference of population

assessment models that integrate information from multiple diverse data sets. Fish Res.

2017; 192: 16–27.

41.   Auger-Méthé M, Field C, Albertsen CM, Derocher AE, Lewis MA, Jonsen ID, et al. State-

space models' dirty little secrets: even simple linear Gaussian models can have estimation

problems. Sci Rep. 2016; 6: 26677. doi:10.1038/srep26677

42.   Froese R, Demirel N, Coro G, Kleisner KM, Winker H. Estimating fisheries reference

points from catch and resilience. Fish Fish. 2017; 18: 506–526. doi:10.1111/faf.12190

43.   Plummer M, Nicky Best, Cowles K, Vines K. CODA: Convergence Diagnosis and Output

Analysis for MCMC. R News. 2006; 6: 7–11.

44.   Heidelberger P, Welch PD. Simulation run length control in the presence of an initial

557   transient. Oper Res. 1992; 31: 1109–1144. doi:10.1287/opre.31.6.1109

558   45.   Geweke J. Evaluating the accuracy of sampling-based approaches to the calculation of

559   posterior moments. In: Berger JO, Bernardo JM, Dawid AP, Smith AFM, editors. Bayesian

560   Statistics 4: Proceedings of the Fourth Valencia International Meeting. Oxford: Clarendon

561   Press; 1992. pp. 169–193.

562   46.   Thorson JT, Jensen OP, Zipkin EF. How variable is recruitment for exploited marine

563   fishes? A hierarchical model for testing life history theory. Can J Fish Aquat Sci. 2014; 71:

564   973–983.

565   47.   Smale M. Mustelus palumbes. In: The IUCN Red List of Threatened Species 2006:

566   e.T60247A12333813   [Internet].   2006   [cited   13   Jun   2020].   Available:

567   https://dx.doi.org/10.2305/IUCN.UK.2006.RLTS.T60247A12333813.en.

568   48.   Smale M. Leucoraja wallacei. In: The IUCN Red List of Threatened Species 2009:

569   e.T161492A5436168   [Internet].   2009   [cited   13   Jun   2020].   Available:

570   https://dx.doi.org/10.2305/IUCN.UK.2009-2.RLTS.T161492A5436168.en

571   49.   Thorson JT, Shelton AO, Ward EJ, Skaug HJ. Geostatistical delta-generalized linear mixed

572   models improve precision for estimated abundance indices for West Coast groundfishes.

573   ICES J Mar Sci. 2015; 72: 1297–1310. doi:10.1093/icesjms/fsu243

574   50.   Currie JC, Thorson JT, Sink KJ, Atkinson LJ, Fairweather TP, Winker H. A novel approach

575   to assess distribution trends from fisheries survey data. Fish Res. 2019; 214: 98–109.

576   doi:10.1016/j.fishres.2019.02.004

577   51.   Collette B, Acero A, Amorim AF, Boustany A, Canales Ramirez C, Cardenas G, et al.

578   Makaira   nigricans.   In: The   IUCN   Red   List   of   Threatened   Species   2011:

e.T170314A6743776 [Internet]. 2011 [cited 13 Jun 2020]. Available:

https://dx.doi.org/10.2305/IUCN.UK.2011-2.RLTS.T170314A6743776.en.

52. ICCAT. Report of the 2018 ICCAT Blue Marlin Stock Assessment Meeting.

SCRS/2018/008. International Commission for the Conservation of Atlantic Tunas

Collective Volume of Scientific Papers. 2018; 75: 813–888.

53. Sherley RB, Crawford RJM, Dyer BM, Kemper J, Makhado AB, Masotla M, et al. The

status and conservation of Cape Gannets Morus capensis. Ostrich. 2019; 90: 335–346.

doi:10.2989/00306525.2019.1684396

54. Distiller G, Altwegg R, Crawford RJM, Klages NTW, Barham B. Factors affecting adult

survival and inter-colony movement at the three South African colonies of Cape gannet.

Mar Ecol Prog Ser. 2012; 461: 245–255. doi:10.3354/meps09807

55. Crawford RJM, Sydeman WJ, Thompson SA, Sherley RB, Makhado AB. Food habits of

an endangered seabird indicate recent poor forage fish availability off western South

Africa. ICES J Mar Sci. 2019; 76: 1344–1352. doi:10.1093/icesjms/fsz081

56. BirdLife-International. Morus capensis. In: The IUCN Red List of Threatened Species

2018: e.T2269666 [Internet]. 2018 [cited 13 Jun 2020]. Available:

http://dx.doi.org/10.2305/IUCN.UK.2018-2.RLTS.T22696668A132587992.en.

57. Ale SB, Mishra C. The snow leopard's questionable comeback. Science. 2018; 359: 1110–

1110. doi:10.1126/science.aas9893

58. Akçakaya HR, Keith DA, Burgman M, Butchart SHM, Hoffmann M, Regan HM, et al.

Inferring extinctions III: A cost-benefit framework for listing extinct species. Biol Conserv.

2017; 214: 336–342. doi:10.1016/j.biocon.2017.07.027

59. Winker H, Novellie P, Selier J, Birss C, Hraber H. Population trends and management

strategy tools for Cape Mountain Zebra. South African National Biodiversity Institute;

2016 pp. 1–46. Report Number: SANBI/BAM/STATS/2016/MZ/Final.

60. da Silva C, Winker H, Parker D, Kerwath SE. Assessment of smoothhound shark Mustelus

mustelus in South Africa. Department of Agriculture, Forestry and Fisheries; 2019 pp. 1–

29. Report No.: DEFF/FISHERIES/LSWG/AUG 2019.

61. Godfrey M, Godley B. Seeing past the red: flawed IUCN global listings for sea turtles.

Endanger Species Res. 2008; 6: 155–159. doi:10.3354/esr00071

62. Boyd IL. Assessing the effectiveness of conservation measures: Resolving the "wicked"

problem of the Steller sea lion. Biol Conserv. 2010; 143: 1664–1674.

doi:10.1016/j.biocon.2010.04.006

Figures

Figure 1: Input datasets of standardized relative abundance indices (± 95% CIs) from South African trawl

surveys for two chondrichthyes species, (a) Whitespot Smoothhound *Mustelus palumbes* and (b)

Yellowspotted Skate *Leucoraja wallacei* and Model diagnostics plots produced by JARA showing (c) – (d)

scaled observed relative abundance (coloured points) for each catch per unit effort (CPUE) timeseries with

model fits (predicted relative abundance, black line) and 95% posterior predictive credibility intervals (grey

polygons) and (e) – (f) colour-coded lognormal residuals of observed versus predicted abundance indices

(coloured points), boxplots (grey bars) indicating the median and quantiles of all residuals available for any

given year, and a loess smoother (black line) through all residuals to highlight any systematically auto-

correlated residual patterns.

Year

Figure 2: Model diagnostics plots produced by JARA using data from Whitespot Smoothhound *Mustelus palumbes*. Each panel shows the state-space model fit (blue line) to the natural logarithm (log) of the observed relative abundance data (white points), shown with 95% credible intervals (black lines) derived from the observation variance.

Figure 3: JARA decision-support plots for (left) Whitespot Smoothhound *Mustelus palumbes* and (right)
Yellowspotted Skate *Leucoraja wallacei* showing (a) – (b) the overall JARA fit (black line) to the observed
timeseries and the projected (red line) population trajectory over three generation length (GL); (c) – (d) the
posterior probability for the percentage annual population change calculated from all the observed data (in
black), from the last 1 generation length (in blue), from the last 2 generation lengths (in green) with the

medians (solid lines) shown relative to a stable population (% change = 0, black dashed line); (e) – (f) the

median change over three generation lengths ("Change = xx") and corresponding probabilities for rates of

population change falling within the IUCN Red List categories.

Figure 4: JARA plots for trend analyses of two types of abundance information for Atlantic Blue Marlin

Makaira nigricans: (left) ten standardized catch per unit effort (CPUE) indices from multiple fishing fleets

(1959–2016) and (right) estimated biomass estimates from the 2018 Atlantic stock assessment conducted

by the International Commission for the Conservation of Atlantic Tuna (ICCAT), showing (a) – (b) scaled

observed relative abundance with model fits (predicted relative abundance, black line) and 95% posterior

predictive credibility intervals (grey polygons); (c) – (d) the overall JARA fit (black line) over the observed

timeseries with vertical dashed lines denoting the periods corresponding to 1–3 GL; (e) – (f) the median

change over three generation lengths ("Change = xx") and corresponding probabilities for rates of

population change falling within the IUCN Red List categories.

Figure 5: JARA decision-support plots for Cape Gannet *Morus capensis* showing (a) the JARA fits

(coloured lines) and 95% credible intervals (grey polygons) to each observed abundance timeseries

(coloured points); (b) the overall JARA fit (black line) to the observed timeseries and the projected (red

line) population trajectory over three generation times (GLs); (c) the posterior probability for the percentage

annual population change calculated from all the observed data (in black), from the last 1 generation length

(in blue), from the last 2 generation lengths (in green), and from the last 3 generation lengths (in red) with

the means (solid lines) shown relative to a stable population (% change = 0, black dashed line); (e) the

median change over three GLs ("Change = xx") and corresponding probabilities for rates of population

change falling within the IUCN Red List categories; (e) retrospective pattern of total abundance estimates

obtained through sequential removal of terminal years and subsequent forward projections to attain 3 GL

("retrospective peel"); and corresponding retrospective status posteriors of change over three generation

lengths - coloured according to the IUCN Red List category thresholds for the Red List criteria A2.

Figure. 6: Simulated 'true' abundance trajectories and observations for the first 15 of 200 simulation runs

together with estimated abundance trends using three alternative estimation models (EMs): (1) 'Two-

Point", (2) "Regression", and (3) 'JARA'. The vertical red dashed line denotes the assessment horizon of

30 years (3 GL) and grey polygons show the 95% credible intervals for the JARA fit.

Figure. 7: Simulation testing results across 200 runs showing (a) boxplots of errors between the estimated
and 'true' percentage change over 3 generation lengths (GL) with Median Absolute Errors and Median
Errors (in brackets) displayed of each box and (b) barplots illustrating the percentages of correctly classified
IUCN threat categories (LC, VU, EN, CR) for the three alternative estimation models (EMs): (1) 'Two-
Point", (2) "Regression", and (3) 'JARA'.

Supplementary Information

Figure S1: Illustrations of (a) a simulated 'true' population trajectory, the corresponding observed

values given the observation error and the underlying deterministic mean trend $\bar{r}$; and (b)

decomposed first-order autoregressive process error deviations simulating the natural variation in

population numbers (Stochasticity) and a more persistent, regime-like variation in anthropogenic

impact (Impact), together with a random observation deviations from the 'true' abundance due to

random sampling error.

Figure S2: Simulated distribution of the 'true' %*C* generated from 200 simulation replicates of

abundance trajectories over a 30-year assessment horizon.

```
688   #-----------------------------------------------------------------------------------------------------------
689   # Generic Population Simulator
690   #-----------------------------------------------------------------------------------------------------------
691
692   popsim = function(n0 = 1000 ,simyrs = 40,r.mu = runif(1,-0.05,0),proc.pop =
693         0.1,AR1.pop=0.3,proc.imp=0.05,AR1.imp=0.8,CV.obs=0.15,GT=10,Plot=TRUE){
694   # Process error
695   # (1) stochastic variation in popdyn
696   pop_var = rnorm(simyrs,0,proc.pop)
697   # Add AR1 (serial auto-correlation)
698   pop_dev = pop_var[1]
699   for(y in 2:simyrs) pop_dev[y] = AR1.pop * pop_dev[y-1]+pop_var[y]*sqrt(1-AR1.pop^2)
700
701   # (2) random variation in impact (e.g. offtake)
702   imp_var = rnorm(simyrs,0,proc.imp)
703   # Add AR1 (serial auto-correlation)
704   imp_dev = imp_var[1]
705   for(y in 2:simyrs) imp_dev[y] = AR1.imp * imp_dev[y-1]+imp_var[y]*sqrt(1-AR1.imp^2)
706
707   # Combine all as latent effect in r vector given r.mu
708   r = r.mu+pop_dev-imp_dev
709
710   # Add random observation error
711   obs_dev = rnorm(simyrs,0,CV.obs)
712
713   # True Population
714   n_t = n0*exp(r[1])
715   for(t in 2:simyrs){
716   n_t[t] = n_t[t-1]*exp(r[t-1])
717   if(n_t[t]>n0*1.5) n_t[t] = n0*1.5}
718
719   # Deterministic trend
720   nbar = n0
721   for(t in 2:simyrs){
722   nbar[t] = nbar[t-1]*exp(r.mu)}
723
724   # Imperfect observation
725   y_t =exp(log(n_t)+obs_dev)
726
727   if(Plot==TRUE){
728   Par = list(mfrow=c(1,2),mar = c(4, 4, 1, 1), mgp =c(2.5,1,0),mai = c(0.6, 0.6, 0.1, 0.1),mex=0.8, tck = -0.02,cex=0.8)
729   par(Par)
730   # Error
731   plot(n_t,ylim=c(0,max(n_t,y_t,n0*1.05)),type="l",col=2,lwd=2,ylab="Abundance",xlab="Year")
732   points(y_t,pch=21,bg="white",type="p")
733   lines(nbar,lty=2)
734   legend("topright",c("True","Observed","Mean trend"),pch=c(-1,1,1),lwd=c(2,-1,1),col=c(2,1,1),lty=c(1,-1,2),bty="n",cex=0.8)
735   legend("topleft","a)",bty="n",cex=1.1,x.intersp = -0.5,y.intersp = -0.2)
736   plot(pop_dev,ylim=range(c(pop_dev,imp_dev,obs_dev)),type="l",col=3,lwd=2,ylab="Deviations",xlab="Year")
737   lines(imp_dev,col=4,lwd=2)
738   legend("topright",c("Stochasticity","Impact","Observation"),pch=c(-1,-1,1),lwd=c(2,2,1),col=c(3,4,1),bty="n",cex=0.8)
739   points(obs_dev,type="b")
740   abline(h=0,lty=2)
741   legend("topleft","b)",bty="n",cex=1.1,x.intersp = -0.5,y.intersp = -0.2)
742
743   }
744   return(list(dat=data.frame(yr=1:simyrs,y=y_t),N=data.frame(yr=1:simyrs,N=n_t),r.mu=r.mu,perc.change=(n_t[simyrs]/n_t[simyrs-3*GT+1]-
745         1)*100))
746   }
747
748   # Run function
749   popsim()
750
751
752
```