1 **CRISPR-Cas immunity repressed by a biofilm-activating pathway in *Pseudomonas***
2 ***aeruginosa***
3

4 Adair L. Borges#, Bardo Castro#, Sutharsan Govindarajan, Tina Solvik, Veronica
5 Escalante, Joseph Bondy-Denomy*
6

Department of Microbiology & Immunology and the Quantitative Biosciences Institute,
University of California, San Francisco, San Francisco, CA 94143
#Equal contribution
*Corresponding author: joseph.bondy-denomy@ucsf.edu

CRISPR-Cas systems are adaptive immune systems that protect bacteria from
bacteriophage (phage) infection. To provide immunity, RNA-guided protein
surveillance complexes recognize foreign nucleic acids, triggering their
destruction by Cas nucleases. While the essential requirements for immune
activity are well understood, the physiological cues that regulate CRISPR-Cas
expression are not. Here, a forward genetic screen identifies a two-component
system (KinB/AlgB), previously characterized in regulating *Pseudomonas*
***aeruginosa* virulence and biofilm establishment, as a regulator of the biogenesis**
and activity of the Type I-F CRISPR-Cas system. Downstream of the KinB/AlgB
system, activators of biofilm production AlgU (a σ^E orthologue) and AlgR, act as
repressors of CRISPR-Cas activity during planktonic and surface-associated
growth. AmrZ, another biofilm activator, functions as a surface-specific repressor
of CRISPR-Cas immunity. *Pseudomonas* phages and plasmids have taken
advantage of this regulatory scheme, and carry hijacked homologs of AmrZ,
which are functional CRISPR-Cas repressors. This suggests that while CRISPR-
Cas regulation may be important to limit self-toxicity, endogenous repressive
pathways represent a vulnerability for parasite manipulation.

Clustered regularly interspaced short palindromic repeats (CRISPR) and CRISPR-
associated (cas) genes are RNA-guided nucleases found in nearly half of all bacteria[1].
CRISPR-Cas systems are mechanistically diverse, with six distinct types (I-VI) identified,
based on signature genes, mechanisms of action, and the type of nucleic acid target[1].
Our strong understanding of the basic components to enable sequence specific DNA

and RNA cleavage have enabled functional transplantation into heterologous bacterial[2,3]

and eukaryotic[4,5] hosts. Given that many Cas proteins encode nucleases[6] the fine-tuned

regulation of these systems to avoid toxicity is likely a key factor to enable safe retention

of a CRISPR-Cas immune system[7]. Multiple signals have been shown to activate

CRISPR-Cas function in diverse organisms, such as quorum sensing[8,9], temperature[10],

membrane stress[11,12], altered host metabolite levels[13,14], and phage infection[15-18].

However, relatively little is known regarding the factors and/or signals that serve to

temper CRISPR-Cas activity and mitigate the risk of acquiring and expressing a

nucleolytic immune system.

Type I CRISPR-Cas systems are comprised of a multi-subunit RNA-guided surveillance

complex, a trans-acting nuclease (Cas3)[19-21], and proteins dedicated to spacer

acquisition, Cas1 and Cas2[22,23]. *Pseudomonas aeruginosa* has become a powerful

model organism for studying Type I CRISPR-Cas mechanisms[24-29], functions[3,23,30,31],

evolution[32-34], and interactions with phages utilizing anti-CRISPR proteins[35-38]. The *P.*

aeruginosa strain PA14 possesses an active Type I-F CRISPR-Cas immune system that

naturally targets many phages. The surveillance complex is composed of four Cas

proteins (Csy1, Csy2, Csy3, and Csy4), which are encoded together in a single

operon[26]. The surveillance complex is guided by CRISPR RNAs (crRNAs) originating

from one of two chromosomally encoded CRISPR arrays[26]. Finally, a separate operon is

responsible for the production of Cas1 and a Cas2-3 fusion protein, the latter of which

mediates nucleolytic destruction of target DNA.

To discover regulators controlling the expression of the CRISPR-Cas surveillance

complex in *P. aeruginosa*, we utilized a *P. aeruginosa* strain PA14 engineered to

express *lacZ* in place of the *csy3* gene (*csy3::lacZ*)[39]. The *csy3*::*lacZ* reporter strain was

subjected to transposon mutagenesis and colonies isolated and screened on X-gal

plates. ~40,000 colonies were visually examined for increased or decreased levels of β-

galactosidase. Multiple independent insertions were identified that abolished LacZ levels

within *lacZ* and upstream genes (*csy1* and *csy2*), indicating successful mutagenesis and

screening. Thirty distinct mutants with transposon insertions outside of this region, with

altered β-galactosidase levels, were isolated and mapped. These mutants are listed in

Supplementary Table 1 with β-galactosidase levels quantified for mutants that grew well

in culture. Four independent insertions were identified in a single gene, *kinB*, which

resulted in apparent mucoidy plating morphology and decreased β-galactosidase

production (Supplementary Fig. 1a). Quantification of β-galactosidase levels in liquid

growth experiments revealed ~50% less *csy3::lacZ* expression compared to wild-type

(WT) PA14 for each mutant (Supplementary Table 1, Supplementary Fig. 1b). We

focused on *kinB* (a sensor kinase/phosphatase) because it was the only gene with >1

independent transposon insertion and displayed the largest β-galactosidase activity

change.

To determine the consequences of *kinB* disruption and decreased *csy* expression on

CRISPR-Cas function, we measured the ability of *kinB::Tn* insertions to limit the

replication of phage, when introduced into a wild-type strain background. Phages used

to assay activity are: DMS3m$_{acrIE3}$ which is an untargeted control phage, DMS3m$_{acrIE3}$ [40],

which is fully targeted by the PA14 Type I-F CRISPR-Cas system, and phage

DMS3m$_{acrIF4}$, which is partially targeted, by virtue of encoding a "weak" anti-CRISPR,

acrIF4, that binds to the surveillance complex to inhibit CRISPR-Cas function[35,37,41]. The

kinB::Tn strains remained resistant to DMS3m$_{acrIE3}$ infection, but DMS3m$_{acrIF4}$ formed 10-

fold more plaques relative to WT, demonstrating attenuated CRISPR-Cas function

(Figure 1a, Supplementary Fig. 1c). This defect was complemented by expression of

kinB in trans (Supplementary Fig. 1c). Growth of control phage DMS3$_{acrIE3}$ was not

impacted in the absence of KinB (Figure 1a). Furthermore, two other targeted phages,

JBD26 (naturally possessing *acrIF4*) and JBD25 (a partially targeted phage with no Acr)

also showed increased replication in the *kinB:Tn* strain (Supplementary Fig. 1d) relative

to WT PA14. Replication of a phage with a weak anti-CRISPR or one that is targeted by

a less active spacer is therefore a sensitive barometer for perturbations in CRISPR-Cas

levels. Together, these data confirm that in the absence of *kinB, csy* gene expression

and phage targeting are decreased.

KinB is a sensor kinase/phosphatase in a two-component system with response

regulator AlgB. The KinB/AlgB system has a large regulon within *P. aeruginosa*, but is

well-known for its role in biofilm regulation[42]. In the absence of KinB function, the activity

of the transcription factor AlgU (a σ^E orthologue and an activator of biofilm formation) is

enhanced[43,44]. This activation of AlgU manifests solely due to the lack of KinB

phosphatase activity and the resulting accumulation of the phosphorylated form of the

response regulator AlgB (P-AlgB). The phosphorylation of AlgB has been attributed to

unknown kinases[45,46] (Fig. 1b). The resulting accumulation of P-AlgB activates the

periplasmic protease AlgW (a DegS homolog), which degrades MucA, liberating sigma

factor AlgU[47,48,49] (Fig. 1b). AlgU positively regulates many factors involved in biofilm

production, including AlgR, AlgD, AlgB, and AmrZ[50-52].

To determine if this KinB-controlled biofilm activation pathway also regulates CRISPR-

Cas immunity, we tested the ability of phosphatase and kinase dead KinB mutants to

complement an in-frame Δ*kinB* deletion, using the replication of the partially-targeted

DMS3m$_{acrIF4}$ phage as a metric for CRISPR-Cas activity. WT *kinB* or kinase inactive

H385A *kinB* complemented the CRISPR defect (Fig. 1c), indicating KinB kinase activity

is dispensable for CRISPR-Cas activation. However, a P390S *kinB* mutant incapable of

dephosphorylating the response regulator AlgB did not complement, and in fact

decreased CRISPR-Cas activity further (Fig. 1c). This suggests that CRISPR-Cas

expression is inhibited by the accumulation of P-AlgB. A Δ*kinB*Δ*algB* double mutant

restored CRISPR-Cas targeting to WT levels (Supplementary Fig. 2a), confirming the

role of this signaling pathway. A strain lacking *algB* (Δ*algB*) or possessing a D59N

mutant that cannot be phosphorylated had elevated CRISPR-Cas activity, supporting the

repressive role of P-AlgB (Supplementary Fig. 2b). Together, these data suggest that

KinB-mediated dephosphorylation of AlgB lifts CRISPR-Cas inhibition, and conversely

accumulation of high levels of P-AlgB (achieved in Δ*kinB* or *kinB P390S*) leads to

CRISPR-Cas repression.

To assess CRISPR-Cas regulation downstream of P-AlgB, anti-phage immunity was

assayed in Δ*algU* and Δ*algR* backgrounds, revealing enhanced targeting of DMS3m$_{acrIF4}$

but not control phage DMS3$_{acrIE3}$ in each knockout (Fig. 1d). Complementation with *algR*

and *algU* restored CRISPR-Cas levels in their respective knockout backgrounds (Fig.

1d). This demonstrates that these proteins repress CRISPR-Cas immunity. Double

knockouts of each gene combined with Δ*kinB* each resembled the respective Δ*algU* and

Δ*algR* single knockouts, consistent with these factors acting downstream of KinB

(Supplementary Fig. 2a). The observed plaquing defects seen with the DMS3m$_{acrIF4}$

phage are CRISPR-dependent, as double knockouts (*kinB, algB, algU, algR* mutants

combined with *csy3::lacZ*) revealed plaquing equivalent to *csy3::lacZ* alone

(Supplementary Fig. 2c).

Phages that use anti-CRISPRs to inhibit CRISPR-Cas immunity must cooperatively
infect the same cell in order to produce a sufficient dose of anti-CRISPR to permit phage
replication[41,53]. The multiplicity of infection (MOI) required to overwhelm immunity is
dependent on the affinity of the anti-CRISPR for the CRISPR-Cas machinery and the
intracellular concentration of CRISPR-Cas components. To understand how this
regulatory pathway controls phage cooperation requirements on a population level by
potentially shifting the intracellular concentration of CRISPR-Cas complexes, liquid
cultures of ∆algU and ∆kinB mutants were infected with a virulent derivative of phage
DMS3m$_{acrIF4}$ at both high (where the WT cells are sensitive) and low (where the WT cells
are resistant) MOI (Figure 1e). The ∆algU mutant was able to limit phage replication
relative to WT and continue growing when infected with concentration of phages
sufficiently high to overwhelm WT immunity (10^6 PFU), indicating increased levels of
CRISPR-Cas machinery (Fig 1e). Conversely, when infected with a low MOI of phages
(10^4 PFU), all bacterial genotypes grew to high cell density, however the ∆kinB mutant
permitted increased phage replication relative to WT, demonstrating a decreased level of
immune system components (Fig. 1f). Importantly, these changes in phage replication
requirements were not seen in the absence of CRISPR-Cas immunity, as phage
replication in double knockouts algU csy3::lacZ and kinB csy3::lacZ resembles
csy3::lacZ (Supplementary Fig. 3a-d, f). In contrast, ∆algR demonstrated CRISPR-
independent phage resistance during liquid culture infection, and phage cooperation
requirements could not be reliably measured for this mutant (Supplementary Fig. 3e).
Taken together, these data demonstrate that a well-characterized biofilm activation
pathway controlled by alternate sigma factor, AlgU, represses CRISPR-Cas immunity in
P. aeruginosa.
161
Our screen focused on finding regulators of the csy operon, however the recruited Cas3
nuclease (encoded in a separate, nearby operon) is also an important player in Type I
CRISPR-Cas immunity. To measure the relative impact of the KinB/AlgB pathway in
controlling Cas3 and Csy complex levels, we fused an N-terminal sfCherry tag to the
endogenous csy1 or cas3 gene in the mutant backgrounds. We observed decreased
levels of both Cas3-sfCherry and Csy1-sfCherry in ∆kinB and increased levels in ∆algR
and ∆algU relative to WT, demonstrating that this pathway controls the levels of both
Cas3 and the Csy complex in the bacterial cell (Fig. 2a).

As Cas3-sfCherry was expressed at low levels relative to Csy1-sfCherry, and is also
known to be subject to post-translational control by Cas1[28], we sought to dissect the
relative contribution of nuclease versus surveillance complex disregulation in driving the
immune phenotypes of the KinB/AlgB pathway mutants. To specifically measure the
anti-phage activity of the Csy complex, we developed a Cas3-indepedent bioassay to
read out the activity of the surveillance complex in the cell. Through the rational design
of crRNAs to target an early phage promoter (P_{E}_1, P_{E}_2), we observed inhibition of
phage replication in a *P. aeruginosa* strain with a nuclease dead Cas3 ($Cas3_{D124A}$, Fig.
2a). This CRISPR-based transcriptional interference (CRISPRi) effect was also seen
when infecting with a phage that expressed the inhibitor of Cas3 recruitment, AcrIF3, but
not an inhibitor that blocks Csy complex-phage DNA binding[37], AcrIF1 (Fig. 2b).
CRISPRi was mediated by crRNAs targeting phage promoters, including one (P_{E}_2) that
completely limited phage replication in the absence of Cas3 activity, while an ORF-
targeting crRNA (ORF_1) was ineffective (Fig. 2b). We selected P_{E}_1 as a moderately-
functional CRISPRi spacer and expressed it in various mutant backgrounds generated in
this study. We observed decreased CRISPRi activity against phage DMS3m*acrIF3* in the
ΔkinB background, but increased CRISPRi-mediated phage repression in Δ*algR* and
Δ*algU,* (Fig. 2c), demonstrating that modulation of *csy* gene expression by this pathway
is sufficient to impact phage targeting, in a Cas3-independent manner. We conclude that
while the KinB/AlgB pathway regulates both Cas3 and Csy complex levels, repression of
surveillance complex biogenesis alone represents a powerful mechanism by which the
KinB/AlgB pathway controls CRISPR-Cas immunity.

AlgU and AlgR are proteins that regulate hundreds of genes in *P. aeruginosa*[54-56],
ultimately repressing acute virulence factors (e.g. pyocyanin) and activating chronic
virulence factors (e.g. biofilm production). To identify downstream CRISPR-Cas
regulators, we considered the large AlgU regulon[56], but focused on a factor involved in
biofilm production, whose expression is AlgU-dependent, *amrZ*[51,57]. AmrZ is a ribbon-
helix-helix type transcription factor, and acts as both a repressor and activator of
transcription in *Pseudomonas*[58]. We generated a knockout of *amrZ* and observed a
CRISPR-dependent decrease in DMS3m*acrIF4* plaque formation (Fig. 3a, Supplementary
Fig. 2c). This was complemented when *amrZ* was expressed *in trans* (Fig. 3a), indicating
that AmrZ is also a repressor of CRISPR-Cas immunity. However, when we measured
phage cooperation requirements in the Δ*amrZ* strain, phage replication and immune

protection did not differ substantially from WT (Fig. 3b,c). The most obvious difference
between the phage plaque assay and the assay to measure phage cooperation is that a
plaque assay is performed on solid plates whereas phage cooperation requirements
were measured in liquid culture. To test if AmrZ was a surface-specific repressor of
CRIPSR-Cas immunity, we measured the levels of Csy complex during surface
association and planktonic growth in WT and Δ*amrZ* cells using an endogenous *csy1-*
sfCherry reporter. In WT cells, the levels of Csy complex were attenuated during
surface-association relative to planktonic growth, but in the absence of AmrZ, Csy
complex levels during surface association increased to levels comparable to those in
planktonic growth (Fig. 3d). Deletion of *amrZ* did not impact Csy complex levels during
planktonic growth (Fig. 3e). To artificially increase the levels of AmrZ during planktonic
growth, we ectopically expressed AmrZ from a high copy plasmid. In this scenario, high
levels of AmrZ were sufficient to completely inhibit Csy complex expression, suggesting
that low AmrZ activity in planktonic growth underlies its surface-specific control of
CRISPR-Cas (Fig. 3f). In contrast to AmrZ, overexpression of the AlgU moderately
impacted Csy complex levels and AlgR was unable to repress *csy1-sfCherry* when
overexpressed (Fig. 3f). This is likely because AlgU and AlgR are master regulators and
many redundant mechanisms are in place to control their activity. Taken together, these
data demonstrate that AmrZ is a surface-specific repressor of the *csy* operon.

We next considered if phages and other mobile genetic parasites naturally antagonized
by CRISPR-Cas immunity had evolved mechanisms to manipulate this repressive
pathway. Inspired by the discovery of a *Paraburkholderia* phage that carried a distant
homolog of AmrZ[59], we searched the NCBI database for AmrZ homologs on
Pseudomonas mobile genetic elements (MGE). Excitingly, we identified 14 diverse
Pseudomonas genetic parasites carrying AmrZ homologs (Supplementary Table 2).
These MGEs included obligately lytic and temperate Myoviridae, temperate
Siphoviridae, and plasmids. AmrZ has been structurally characterized in complex with
operator DNA[58], and these mobile AmrZ homologs showed perfect conservation of
critical DNA-interacting residues in the ribbon-helix-helix domain, suggesting conserved
binding specificity (Fig. 4a, b, red residues/arrowheads). To test if these mobilized AmrZ
variants were capable of regulating CRISPR-Cas activity in *Pseudomonas aeruginosa*,
we synthesized 6 MGE-borne *amrZ* homologs and tested them for their ability to
complement the Δ*amrZ* strain. 5/6 homologs complemented the Δ*amrZ* mutant,

indicating they were active in the PA14 transcriptional network and were *bona fide* AmrZ
orthologs (Fig. 4c). Next, each gene was expressed in WT cells carrying an endogenous
csy1-sfCherry reporter, revealing 3 *P. aeruginosa* phage homologs (*amrZ$_{PaBG,}$ amrZ$_{phi3,}$*
amrZ$_{JBD68}$) that inhibited Csy complex biogenesis at varying efficiencies (Fig. 4d). This
suggests the possibility that *amrZ* has been hijacked for its capacity to inhibit CRISPR-
Cas expression. Thus, *amrZ* acts on both sides of the battle between bacteria and
mobile genetic elements.

Discussion

The factors that govern expression of bacterial processes are highly variable across
species, reflecting niche-specific adaptations. CRISPR-Cas immune systems limit the
replication of foreign genetic elements, but few reports address how the regulation of
CRISPR-Cas impacts phage targeting. Here, we utilized a genetic screen that identified
KinB as a positive regulator of CRISPR-Cas function in *P. aeruginosa.* This sensor
kinase/phosphatase operates in a two-component system with AlgB that regulates a
well-characterized biofilm-stimulatory pathway, though the ligand for KinB remains
unknown. Removal of KinB or inactivation of its phosphatase activity leads to the
accumulation of P-AlgB, which stimulates the activity of CRISPR-Cas repressors AlgU,
AlgR, and AmrZ. This pathway is well-studied due to the recurrent isolation of alginate-
overproducing (mucoid) *P. aeruginosa* from the lungs of cystic fibrosis patients, with
mutations in *mucA*[60] and *kinB*[47,61] that lead to the constitutive activation of AlgU. This
biofilm pathway generates the characteristic mucoidy phenotype of CF *P. aeruginosa*
isolates[43,62,63], which we also observed in *kinB:Tn* mutants (Supplementary Fig. 1a).

This pathway is well described in its capacity to activate biofilm production, and here we
show that this transcriptional network also functions to dampen CRISPR-Cas activity in
non-biofilm conditions. In WT cells, KinB phosphatase activity increases CRISPR-Cas
function, though the effect is relatively mild in our experimental conditions. Relieving
transcriptional repression via deletion of *algU* or *algR* however leads to a hyper-
activation of the CRISPR-Cas immune response. We further demonstrate that *amrZ* acts
in this pathway to specifically repress CRISPR-Cas when cells are surface-associated,
leading to lower levels of Csy complex during surface-association relative to planktonic
growth. Phages and plasmids have taken advantage of this regulatory scheme, and
some *Pseudomonas* genetic parasites carry hijacked copies of the CRISPR-Cas

repressor *amrZ*. We show that these mobilized *amrZ* homologs retain their capacity to
transcriptionally repress CRISPR-Cas, demonstrating a dual role for this gene in the
host-pathogen arms race between bacteria and their parasites.

Considering the physiology of *P. aeruginosa* can put these findings into context with a
phage and CRISPR-Cas-centric view. During exponential growth is when we and others
observe CRISPR-Cas activation[8,9], and is also when phage infection risk is likely high
(i.e. metabolically active, well-mixed planktonic culture[33]). Conversely, surface-
association lessens phage infection risk, as spatial structure limits phage dispersal and
prevents a phage bloom from overtaking the entire bacterial population[64]. Though we do
not measure phage sensitivity in a biofilm here, biofilm production likely provides an
even greater level of intrinsic phage resistance, due to high levels of spatial stratification
and abundant polysaccharide secretion.

Supporting our proposal that *P. aeruginosa* surface-association/biofilm formation and
CRISPR-Cas expression are inversely correlated are two independent observations: **i)**
Our analysis of a previously published PA14 RNAseq data set[65] and proteomic data set[66]
revealed activation of CRISPR-Cas expression in exponential phase, and strong
transcriptional repression during stationary phase and biofilm growth (Supplementary
Fig. 4a). Interestingly Cas proteins are still detected in stationary phase and biofilm
growth, suggesting the cells retain some immune-competency even after transcriptional
shutdown (Supplementary Fig. 4b). **ii)** Previous studies on the *P. aeruginosa* CRISPR-
Cas system have revealed that the genome is hyper-sensitive to CRISPR-induced DNA
damage during surface-association and biofilm formation, leading to cell death when a
mismatched prophage sequence target is present in the chromosome[30,31]. This suggests
that CRISPR auto-immunity costs are also dependent on the growth state and physical
environment of the cell, and that CRISPR-Cas regulation in *P. aeruginosa* is tuned to
reflect this intrinsic sensitivity.

Here, we report the identification of a CRISPR-Cas repressive pathway in *P. aeruginosa*.
Phages bearing distinct anti-CRISPRs are employed in bioassays to read out the activity
of the CRISPR-Cas surveillance complex. Using a CRISPRi phage-repression assay,
coupled with translational reporters, we conclude that *algU*, *algR*, and *amrZ* control the
production of the CRISPR surveillance complex (Csy complex), which is where the
screen was focused. Cas3 levels, though low, are also modulated by the KinB/AlgB

pathway. Interestingly, our observations show that CRISPR-Cas activity in *P. aeruginosa* does not operate at full strength, despite the capacity of a hyper-activated immune system to severely limit the replication of phages relying on anti-CRISPRs for survival. We speculate that the ability to control CRISPR-Cas activity during lifestyle transitions may be essential for *P. aeruginosa* to safely maintain a CRISPR-Cas system by limiting self-toxicity. Furthermore, we reveal an unexpected cost to negative regulation of CRISPR-Cas immunity in our discovery of MGE-encoded CRISPR-Cas repressors: though CRISPR-Cas control is likely essential for safe retention of the immune system, the evolution of potent mechanisms of CRISPR-Cas repression has created an Achilles Heel that is exploited by genetic parasites.

Acknowledgements

We thank members of the Bondy-Denomy lab for conversations and support relating to this project. We also thank Kristine Trotta (Seemay Chou's lab, UCSF) for assistance and advice in the development of fluorescent reporter assays, and Andrew Santiago-Frangos (Blake Wiedenheft's lab, MSU) for advice and consultation. The Bondy-Denomy lab was supported by the University of California, San Francisco Program for Breakthrough in Biomedical Research, funded in part by the Sandler Foundation, and an NIH Office of the Director Early Independence Award (DP5-OD021344), and R01GM127489. We thank Deborah Hung's lab for providing Δ*kinB,* Δ*algR,* Δ*algU* strains and associated mutants, and George O'Toole's lab providing the *csy3::lacZ* PA14 strain.

Author Contributions

J.B.D., A.L.B, and B.C. formulated study design and plans. A.L.B. designed and conducted bacteriophage plaque assays, liquid infection assays, and CRISPRi assays, performed CRISPR-Cas expression profiling, and conducted bioinformatics analyses. B.C. conducted the genetic screen, isolated and constructed bacterial mutants, conducted LacZ expression profiling, and performed bacteriophage plaque assays. S.G. designed, constructed, and validated sfCherry reporter strains. T.S. conducted CRISPRi assays and V.E. assisted in establishing fluorescent reporter assays. J.B.D. and A.L.B wrote the manuscript.

Competing Interests

335 J.B.-D. is a scientific advisory board member of SNIPR Biome and Excision
336 Biotherapeutics and a scientific advisory board member and co-founder of Acrigen
337 Biosciences.
338
339 **Materials & Correspondence**
340 Correspondence should be directed to joseph.bondy-denomy@ucsf.edu
341
Methods
Bacterial strains and bacteriophages. *P. aeruginosa* UCBPP-PA14 (PA14) strains
and *E. Coli* strains (Sup. Table) were grown on lysogeny broth (LB) agar or liquid at 37
^{o}C. Media was supplemented with gentamicin (50 μg ml^{-1} for *P. aeruginosa* and 30 μg ml^{-}
^{1} for *E. Coli*) to maintain the pHERD30T plasmid or carbenicillin (250 μg ml^{-1}) for *P.*
aeruginosa or ampicillin (100 μg ml^{-1}) for *E. coli* containing the pHERD20T plasmids.
pHERD plasmids were induced with 0.1% arabinose. Bacteriophages stocks were
prepared as described previously[40]. In brief, 3 ml of SM buffer was added to plate
lysates of the desired purified bacteriophage and incubated at room temperature for 15
minutes. SM buffer containing phages was collected and 100 μl of chloroform was
added. This was centrifuged at 10,000 x g for 5 minutes and supernatant containing
phages was transferred to a screw cap storage tube and incubated at 4 ^{o}C.
354
Tn mutagenesis screen. The *csy*::*lacZ* reporter strain was subjected to transposon
mutagenesis and colonies were isolated on plates containing X-gal (5-bromo-4-chloro-3-
indolyl-β-D-galactopyranoside). ~50,000 colonies were visually examined for increased
or decreased levels of β-galactosidase and insertions mapped by semi-random PCR. To
conduct transposon mutagenesis, overnights of PA14 *csy3::lacZ* and *E. coli* containing
the pBTK30 Tn suicide vector were mixed in a 1:2 ratio (donor : recipient) for
conjugation. A second control group containing PA14wt and *E. Coli* containing the
pBTK30 Tn suicide vector was set up in parallel. Mixed cells were centrifuged at 4,000 x
g for 10 minutes to pellet cells. 100 μl of resuspended conjugation pellet was then
spotted on LB agar plates and incubated at 37 ^{o}C for 6hrs. Conjugation spots were
collected and resuspended in LB liquid media. Conjugation was then screened on an LB
agar plates supplemented with nalidixic acid (30 μg ml^{-}1) and gentamicin (50 μg ml^{-1}).
Surviving colonies containing Tn insertions were collected into 1ml of LB liquid media.
Serial dilution of were prepared and plated on LB agar plates supplemented with x-gal

(200 µg ml^{-1}) and gentamicin (50 µg ml^{-1}) and nalidixic acid (30 µg ml^{-1}). Plates were

incubated at 37 $^{\circ}$C for 24 hours to allow for colonies to change color. Colonies displaying

changed expression levels as compared to our positive control (PA14 csy3::lacZ no

pBTK30) were then isolated onto secondary LB agar plates with X-gal and gentamicin

and nalidixic acid at the stated concentrations. Genomic DNA (gDNA) was collected

from isolated single colonies by resuspending bacterial colonies in 0.02% SDS and

boiling the sample for 15 minutes. Samples were then centrifuged at 10,000 x g and

supernatants containing gDNA were collected. Semi-random PCR was then performed

using PCR primers listed on sup table_. PCR samples were sequenced and reads were

then matched to the *P. aeruginosa* UCBPP-PA14 genome using BLAST. Isolated

containing genes of interest were saved as stocks (s. table 1). Transcription changes

were then verified via modified β-galactosidase assay.

Plaque assays. Plaque assays were performed on LB agar plates (1.5% agar) with LB

top agar (0.7% agar), both supplemented MgSO$_4$ (10 mM final concentration) and

gentamicin (50 µg ml^{-1}) as needed for plasmid maintenance. Spot titrations were done by

mixing 150 µl of a *P. aeruginosa* overnight culture with 3 ml of top agar, which was

dispersed evenly on a LB MgSO$_4$ plate. 3 µl of 10-fold phage dilutions were then spotted

on the surface. Plates were incubated overnight at 30 $^{\circ}$C. To count plaques, full plate

assays were used, except when CRISPR-targeting was so strong that discrete plaques

could not be accurately measured. In this case, spot titrations are shown. For full plate

assays, 3 µl of the selected phage dilution was incubated with 150 µl of *P. aeruginosa*

overnight for 15 minutes at 37 $^{\circ}$C. 3 ml of top agar was then added and the mixture was

dispersed evenly on a LB MgSO$_4$ plate. Individual plaques were then counted to assess

differences in plaquing efficiency.

β-galactosidase assay. A modified version of the previously described β-galactosidase

assay was used to measure *lacZ* activity in transcriptional fusions. Bacterial cultures

were grown overnight at 37 $^{\circ}$C. Cultures were then diluted 1:100 into LB liquid medium

supplemented with the desired antibiotic. Diluted cultures were then incubated at 37 $^{\circ}$C

until the desired time point was reached. Culture density was measured with a

spectrophotometer (OD$_{600}$) and 200 µl of the sample was added 800 µl to

permeabilization solution (components listed in sup table XX). Cells were mixed via

inversion and vortexed for 1 minute to permeabilize the cells. 200 µl of ONPG (4 mg ml^{-}

^{1}) was added and samples were incubated for 27 minutes at 30 ^{o}C. Enzymatic reaction

was stopped by addition of 300 µl of 1M Na_2CO_3. Samples were centrifuged at 13,000 x

g for 5 minutes to remove debris and 200 µl of supernatant was moved to a 96-well plate

to read absorbance at 420 nm and at 550 nm. Miller units were calculated using the

Miller equation. 3 technical replicates per sample per experiment.

Phage transduction of _kinB::Tn_ alleles. Transposon insertions in _kinB_ from a

csy3::lacZ background were transduced into WT PA14 to enable testing of CRISPR-Cas

function with the same transposon insertion. Phage phiPA3 was used to infect the donor

strain (_csy3::lacZ, kinB::Tn_), on plates with top agar overlays, using ~10^4 PFU to

generate near confluent lysis. Plates were soaked in 3-4 mL of phage SM buffer and 2

mL collected over chloroform, vortexed, and pelleted to isolate transducing phage in the

supernatant. Lysates were used to infect recipient strains (WT PA14). ~10^8 PFU were

used to infect a culture at an MOI of 1. After 30 minutes of static incubation on the

bench, cultures were gently shaken at 37 ^{o}C for 20 min and then pelleted at 5000 x g.

Cells were washed twice with LB, and subsequently incubated at 37 ^{o}C for 1 hour to

allow recombination and gentamicin resistance outgrowth. Cultures were pelleted and

resuspended in 200 µL of LB, and plated on LB plates containing gentamicin. Controls

included uninfected cells and cells infected with phages not propagated on a gentamicin

resistant donor strain. Additionally, phage lysate was directly plated under selection to

confirm no residual donor strain in the phage preparation. Plates were incubated

overnight at 37 ^{o}C and their identity (i.e. CRISPR-Cas intact) confirmed with a plaque

assay using DMS3m as the target phage and PCR of the kinB locus.

Introduction of _csy3::lacZ P. aeruginosa_ UCBPP-PA14 strains. The _lacZ_ gene was

introduced into PA14 strains of interest via homologous recombination. Recombination

vector containing lacZ flanked by homology arms matching csy2 and csy4 was

introduced via conjugation. PA14 strains and _E. coli_ containing vector were mixed at a

ratio of 1:2 (recipient:donor). Mixture was heat shocked at 42 ^{o}C for 10 min. Mating spot

was then plated on a LB agar plate and incubated overnight for 30 ^{o}C. Mating spot was

then collected, resuspended in 1 ml of LB liquid media and plated on VBMM plates

supplemented with 50 ug/mL gentamicin to select for colonies with the integrated

homology plasmid. Colonies were cultured overnight in LB in the absence of selection at

37 ^{o}C, and were then diluted and counterselected on no salt LB (NSLB) agar plates

supplemented with 15% sucrose. Surviving colonies were then grown on LB agar plates

supplemented with gentamicin and X-gal to check for *lacZ* insertion via color change and

lacZ insertion was further verified via PCR.

Generation of an PA14Δ*amrZ* strain using the endogenous I-F CRISPR-Cas

system. Complementary oligonucleotides encoding a crRNA targeting the *amrZ* gene of

PA14 were annealed and ligated into the multiple cloning site of the pHERD30T vector.

A fragment possessing homology arms flanking the desired mutation (500 bp upstream

and 500 bp downstream) around *amrZ* was cloned into a distinct location (NheI site) of

the same vector via Gibson assembly. The new plasmid containing both a crRNA and

homology region was introduced into WT PA14 via electroporation. Transformation

efficiency dropped dramatically in the presence of the crRNA due to the toxicity caused

by self-targeting. All surviving colonies had the desired clean deletion of the *amrZ* gene.

Deletions were confirmed by PCR of the region of interest and subsequent Sanger

sequencing of the amplicon. A 2000 bp region flanking *amrZ* was PCR amplified and

sequencing primers were designed to sequence both the deletion junction and outside of

the original 500 bp flanking regions to confirm the removal of the *amrZ* gene.

Liquid phage cooperation assay. Liquid phage infections were performed as

described in[41]. In brief, an overnight culture of cells was diluted 1:100 into fresh media,

and infected with virulent phage DMS3m$_{acrIF4}$ in biological triplicate in a 96 well Costar

plate. Cells were incubated at 37 $^{\circ}$C with constant rotation and OD600 measured every 5

minutes in a Biotek H4 Synergy plate reader. Phage were harvested from each well and

quantified by plaque assay after 24 hours. In these experiments, all strains used in the

assay carried 2 spacers against the DMS3m$_{acrIF4}$ phage to prevent phage escape: one

endogenous spacer (CRISPR2_sp1), and the other spacer was provided on a

pHERD30T plasmid.

Generation of endogenous Csy1-sfCherry and Cas3-sfCherry reporters

Endogenous Csy1-sfCherry and Cas3-sfCherry reporters were constructed similar to the

construction of *csy3::lacZ*. We initially verified that tagging of sfCherry at the N terminus

of Csy1 and Cas3 are functional, when expressed from a plasmid. pMQ30-sfCherry-

Csy1, which contains sfCherry sequence flanked by 657 bp upstream of csy1 and 701

bp downstream of *csy1* start codon, was cloned in pMQ30 plasmid between HindIII and

BamHI sites using Gibson assembly. pMQ30-sfCherry-Cas3, which contains sfCherry

sequence flanked by 353 bp upstream of cas3 and 350 bp downstream of cas3 start

codon, was cloned in pMQ30 plasmid between HindIII and BamHI sites using Gibson

assembly. 4 bp that overlap between the end of *cas1* and the beginning of *cas3* were

duplicated in the final construct. Both pMQ30-sfCherry-Csy1 and pMQ30-sfCherry-Cas3

contains ggaggcggtggagcc sequence (encoding GGGGA) as linker between sfCherry

and the respective tagged proteins. The Csy1-sfCherry and Cas3-sfCherry construct

were introduced into PA14 strains of interest via homologous recombination. Strains

containing appropriate insertion were verified via PCR.

sfCherry reporter profiling. **Liquid**: Cells were diluted 1:100 from an overnight culture

into fresh LB (with antibiotics and inducer if required), and grown for the indicated

number of hours in biological triplicate. 500 µl of each sample was then spun down at

8,000xg for 2 minutes, and resuspended in 500 µl of M9 media. Samples were loaded in

to a 96 well plate (150 µl /well) in technical triplicate and red fluorescence and OD600

were measured using a Biotek H4 Synergy. **Solid:** Cells were diluted 1:100 from an

overnight culture into fresh LB and 20 µl plated onto individual wells in biological

triplicate in a 24 well plate with each well containing solidified 1.5% LB Agar with

antibiotics and inducer if required. The 24 well plate was then covered with a breathable

Aeraseal, and incubated at 37 $^{\circ}$C with no shaking. At the indicated timepoint, cells were

harvested by flooding each well with 500 µl of M9 buffer, and were spun down at

8,000xg for 2 minutes, and resuspended in 500 µl of M9 media. Samples were loaded in

to a 96 well plate (150 µl /well) in technical triplicate and red fluorescence (excitation

580 nm, emission 610 nm) and OD600 were measured using a Biotek H4 Synergy. To

calculate the relative fluorescence units for each sample, the background fluorescence

and background OD600 values obtained were subtracted from the sample values, and

the sample fluorescence was then normalized to the sample OD600.

AmrZ homolog discovery and characterization. BLASTp was used to search the

nonredundant protein database for AmrZ homologs (accession: ABJ12639.1) in

Pseudmonas sp. (taxid: 286) in May 2019. This homolog list (e value > 0.001) was then

examined for homologs found on phage or plasmid genomes. Representative homologs

were aligned using Clustal and the alignment visualized in Jalview, and key conserved

residues were mapped onto the structure in Pymol (PDB ID: 3QOQ). Select homologs

were synthesized (TWIST Biosciences) and cloned into the SacI/PstI site of the
arabinose-inducible plasmid pHERD30T using Gibson assembly. Vectors were
electroporated into *Pseudomonas aeruginosa* strains for functional testing, where they
were induced with 0.1% arabinose and maintained with 50 ug/mL gentimicin.

Figure 1

**Figure 1: *A forward genetic screen identifies a role for a biofilm activating
pathway in repressing CRISPR-Cas immunity.* a**. Efficiency of immunity (EOI) against
isogenic phages DMS3$_{acrIE3}$ (non-targeted) and DMS3m$_{acrIF4}$ (CRISPR-targeted). Plaque
forming units (PFUs) were quantified on two independent *kinB* transposon mutants
(*kinB::Tn1* and *kinB::Tn2*), then represented as a ratio relative to the number of PFUs
measured on WT PA14. **b.** A cartoon summarizing the KinB/AlgB two component
system and downstream effects, based on prior work, with CRISPR-Cas regulation
added. **c,d** EOI measurements for indicated Δ*kinB,* Δ*algR,* and Δ*algU* strains with
complementation. **e,f**. Growth curves of PA14 WT, Δ*algU*, or Δ*kinB* strains infected with
either a high (e,10^6 PFU) or low (f, 10^4 PFU) MOI of virulent DMS3m$_{acrIF4}$. Phage

523　replication was quantified as PFUs after 24 hours, and the efficiency of immunity
524　expressed as a ratio of the number of PFUs harvested from the mutant strain compared
525　to the number of PFUs obtained from WT. All data (OD600 and EOI measurements) are
526　represented as the mean of 3 biological replicates +/- SD.

Figure 2

Figure 2. *The kinB/algB pathway modulates Csy complex levels and activity.*

a. Measurement of the fluorescence levels of Cas3-sfCherry (grey) or Csy1-sfCherry
(black) reporter strains after 10 hours of growth in liquid culture. Fluorescence
measurements are represented as the mean of 3 biological replicates +/- SD. **b**. Spot
titration of DMS3m$_{acrIF3}$ (F3) and DMS3m$_{acrIF1}$ (F1) on *cas3*$_{D124A}$ (dead Cas3) or Δcsy3
(active Cas3, no Csy complex). Phages are targeted by natural spacer CR2_1, as well
as crRNAs designed to target DMS3m genome in positions designated on ORF map. **c**.
Spot titration of DMS3m$_{acrIF3}$ and DMS3m$_{acrIF1}$ phages on deletion mutants expressing
the indicated crRNA.

Figure 3

Figure 3: *AmrZ is a surface-specific repressor of CRISPR-Cas immunity.*

a. Efficiency of immunity (EOI) against isogenic phages DMS3$_{acrIE3}$ (non-targeted) and

DMS3m$_{acrIF4}$ (CRISPR-targeted). Plaque forming units (PFUs) were quantified on ΔamrZ

or the complemented strain, then represented as a ratio of the number of PFUs

measured on WT PA14. EOI measurements are represented as the mean of 2 biological

replicates +/- SD. **b,c**. Growth curves of PA14 WT and Δ*amrZ* infected with either a high

(d,10^6 PFU) or low (e, 10^4 PFU) MOI of virulent DMS3m$_{acrIF4}$. Phage replication was

quantified as PFUs after 24 hours, and the efficiency of immunity expressed as a ratio of

the number of PFUs harvested from Δ*amrZ* to the number of PFUs obtained from WT.

OD600 and EOI measurements are represented as the mean of 3 biological replicates

+/- SD. **d, e**. Timecourse of the fluorescence levels of Csy1-sfCherry reporter strains

during surface-association (d) or planktonic growth (e). **f**. Overexpression of the

indicated gene in a WT *csy1-sfCherry* background, with fluorescence measurements at

10 hours in liquid culture and normalized to empty vector. Fluorescence measurements

are represented as the mean of 3 biological replicates +/- SD.

Figure 4. **Phage AmrZ homologs control CRISPR-Cas immunity. a.** Structure of an
AmrZ tetramer bound to 18bp of operator DNA[58] with DNA-contacting residues
highlighted in red. **b.** Protein alignment of six mobile AmrZ homologs and the native
PA14 AmrZ homolog, with the ribbon-helix-helix DNA binding domain secondary
structure schematized and DNA-contacting residues indicated with red arrows and text.
c. Efficiency of immunity (EOI) against isogenic phages DMS3$_{acrIE3}$ (non-targeted) and
DMS3m$_{acrIF4}$ (CRISPR-targeted). Plaque forming units (PFUs) were quantified on ΔamrZ
or the strains complemented with bacterial or MGE AmrZ homologs, and represented as
a ratio to the number of PFUs measured on WT PA14. EOI measurements as
represented as the mean of 3 biological replicates +/- SD. **d.** Measurement of the
normalized fluorescence levels of Csy1-sfCherry reporter strains transformed with
bacterial or MGE AmrZ homologs after 10 hours of growth in liquid culture. Fluorescence
measurements are represented as the mean of 3 biological replicates +/- SD.

Supplementary Table 1

Gene with Tn insertion	Transposon location	β-gal activity (% of unmutagenized)
kinB (1)	6448519	59%
kinB (2)	6447945	50%
kinB (3)	6447811	53%
kinB (4)	6449373	54%
kinB (5)	6449345	98%
purM	4618060	134%
lasR	4085810	135%
minD	1915545	N/D (growth defect)
gltB	5943637	108%
S-type pyocin	1196879	145%
pyoS3A	4404303	145%
tolA	4595505	158%
glycosyl transferase	5889967	81%
bacA	3490006	75%
Intergenic; zipA and smc +	3979746	N/D (growth defect)
gidB	6530337	88%
deaD	2373899	N/D (growth defect)
polA	6457284	N/D (growth defect)
pchH	798159	131%
crc	6275250	91%
putative plasmid stabilization protein	5347104	N/D (growth defect)
paraquat inducible protein	5532785	108%
pchH	797705	127%
yhiH/yhiI	6162134	108%
oxidoreductase FMN binding	4188602	N/D (growth defect)
putative membrane protein	1400718	118%
Intergenic; fstA and fstZ	5104077	111% (growth defect)
putative Zn-dependent oxidoreductase	2820446	115% (growth defect)
gnyL	3434168	87%
cytochrome c1 precursor	5126449	94%

Supplementary Table 1

All independent transposon insertions identified and mapped by visual screening with

increased or decreased *csy3::lacZ* β-galactosidase activity at 8 hours of growth. The

insertion location in the PA14 genome is shown, along with the measured level of β-

galactosidase enzyme. These measurements were not determined (N/D) for strains with

a growth defect.

Supplementary Table 2

Name	Accession	MGE type
PA14 AmrZ	ABJ12639.1	
Pseudomonas phage Noxifer	ARV77275.1	Lytic Myovirus
Pseudomonas phage phi3	YP_009276432.1	Integrated prophage
Pseudomonas phage PaBG	YP_008433620.1	Lytic Myovirus
Pseudomonas phage SM1	ALT58107.1	Siphoviridae (temperate)
Pseudomonas phage F10	YP_001293379.1	Siphoviridae (temperate)
Pseudomonas phage JBD68	ARM70500.1	Siphoviridae (temperate)
Pseudomonas sp. VLB120 plasmid pSTY	AGZ38169.1	Plasmid
Pseudomonas putida plasmid pKF715B	BAW27310.1	Plasmid
Pseudomonas veronii plasmid PVE_plasmid	SBW85251.1	Plasmid
Pseudomonas koreensis plasmid p3	AVX93364.1	Plasmid
Pseudomonas sp. Leaf58 plasmid pBASL58	AYG48213.1	Plasmid
Pseudomonas sp. XWY-1 plasmid	AUZ62175.1	Plasmid
Pseudomonas putida KF715C_pA870	BAW26592.1	Plasmid
Pseudomonas putida S12 plasmid pTTS12	AJA17154.1	Plasmid
Pseudomonas putida p12969-DIM	ALZ46341.1	Plasmid

577 **Supplementary Table 2:**

578 AmrZ homologs listed by the genome that encodes them, the accession number, and

579 the mobile genetic element type.

Supplementary Figure 1

Supplementary Figure 1

a. A streak plate on X-gal plates, showing strains involved in this study and isolated transposon (Tn) insertions. **b**. β-galactosidase measurements at the indicated time points for the unmutagenized (*csy3::lacZ*) strain and three isolated *kinB* transposon mutants (*kinB::Tn1-3*). **c**. Phage titration on lawns of the *kinB::Tn1* mutant transformed with empty vector or *kinB.* **d.** Spot titration of phages JBD26 (CR2_sp17, sp20-targeted, possessing *acrIF4*), JBD25 (CR1_sp1 targeted) on *kinB::Tn* mutants and ΔCRISPR-Cas.

Supplementary Figure 2

589 **Supplementary Figure 2:**

590 **a-c.** Efficiency of immunity measurements for indicated mutants relative to WT. **a**. Δ*algB*

591 mutant complemented. **b.** Double knockouts show Δ*kinB* combined with *algB*, *algU*, or

592 *algR*. **c**. Indicated knockouts were combined with *csy3::lacZ*.

Supplementary Figure 3

Supplementary Figure 3:

a-e. Growth curves of the indicated knockouts combined with *csy3::lacZ*. infected with virulent DMS3m*acrIF4* at multiplicity of infection (MOI, rainbow colors) increasing in 10-fold steps from 2×10^{-4} to 2×10^{2}. **f.** Phage replication was quantified as PFUs after 24 hours for the MOIs used in Figure 1 and 3, and the efficiency of immunity expressed as a ratio of the number of PFUs harvested from the mutants combined with *csy3::lacZ* under the number of PFUs obtained from WT *csy::lacZ*. OD600 and EOI measurements are represented as the mean of 3 biological replicates +/- SD.

Supplementary Figure 4

Supplementary Figure 4.

a. Log2 of Fragments Per Kilobase of transcript per Million mapped reads (FPKM) shown for each I-F *cas* gene in PA14 in the indicated growth condition. **b.** Log2 of protein levels for each of the I-F Cas proteins in PA14 in the indicated growth condition.

REFERENCES

1.   Makarova, K. S. *et al.* An updated evolutionary classification of CRISPR-Cas systems. *Nat Rev Micro* **13,** 722–736 (2015).
2.   Sapranauskas, R. *et al.* The Streptococcus thermophilus CRISPR/Cas system provides immunity in Escherichia coli. *Nucleic Acids Research* **39,** 9275–9282 (2011).
3.   Vorontsova, D. *et al.* Foreign DNA acquisition by the I-F CRISPR-Cas system requires all components of the interference machinery. *Nucleic Acids Research* **43,** 10848–10860 (2015).
4.   Mali, P. *et al.* RNA-guided human genome engineering via Cas9. *Science* **339,** 823–826 (2013).
5.   Cong, L. *et al.* Multiplex Genome Engineering Using CRISPR/Cas Systems. *Science* **339,** 819–823 (2013).
6.   Leon, L. M., Mendoza, S. D. & Bondy-Denomy, J. How bacteria control the CRISPR-Cas arsenal. *Curr Opin Microbiol* **42,** 87–95 (2018).
7.   Patterson, A. G., Yevstigneyeva, M. S. & Fineran, P. C. Regulation of CRISPR-Cas adaptive immune systems. *Curr Opin Microbiol* **37,** 1–7 (2017).
8.   Høyland-Kroghsbo, N. M. *et al.* Quorum sensing controls the Pseudomonas aeruginosa CRISPR-Cas adaptive immune system. *Proceedings of the National Academy of Sciences* **114,** 201617415–135 (2016).
9.   Patterson, A. G. *et al.* Quorum Sensing Controls Adaptive Immunity through the

633    Regulation of Multiple CRISPR-Cas Systems. *Mol Cell* **64,** 1102–1108 (2016).

634 10. Høyland-Kroghsbo, N. M., Muñoz, K. A. & Bassler, B. L. Temperature, by
635    Controlling Growth Rate, Regulates CRISPR-Cas Activity in Pseudomonas
636    aeruginosa. *mBio* **9,** 1709–12 (2018).

637 11. Perez-Rodriguez, R. *et al.* Envelope stress is a trigger of CRISPR RNA-mediated
638    DNA silencing in Escherichia coli. *Molecular Microbiology* **79,** 584–599 (2011).

639 12. Serbanescu, M. A. *et al.* Role of the Streptococcus mutans CRISPR/Cas systems
640    in immunity and cell physiology. (2014). doi:10.1128/JB.02333-14

641 13. Shinkai, A. *et al.* Transcription activation mediated by a cyclic AMP receptor
642    protein from Thermus thermophilus HB8. *J. Bacteriol.* **189,** 3891–3901 (2007).

643 14. Patterson, A. G., Chang, J. T., Taylor, C. & Fineran, P. C. Regulation of the Type
644    I-F CRISPR-Cas system by CRP-cAMP and GalM controls spacer acquisition and
645    interference. *Nucleic Acids Research* **43,** gkv517–6048 (2015).

646 15. Agari, Y. *et al.* Transcription profile of Thermus thermophilus CRISPR systems
647    after phage infection. *Journal of Molecular Biology* **395,** 270–281 (2010).

648 16. Young, J. C. *et al.* Phage-induced expression of CRISPR-associated proteins is
649    revealed by shotgun proteomics in Streptococcus thermophilus. *PLoS ONE* **7,**
650    e38077 (2012).

651 17. Quax, T. E. F. *et al.* Massive activation of archaeal defense genes during viral
652    infection. *Journal of Virology* **87,** 8419–8428 (2013).

653 18. Fusco, S. *et al.* Transcriptome analysis of Sulfolobus solfataricus infected with two
654    related fuselloviruses reveals novel insights into the regulation of CRISPR-Cas
655    system. *Biochimie* (2015). doi:10.1016/j.biochi.2015.04.006

656 19. Brouns, S. J. J. *et al.* Small CRISPR RNAs guide antiviral defense in prokaryotes.
657    *Science* **321,** 960–964 (2008).

658 20. Hochstrasser, M. L. *et al.* CasA mediates Cas3-catalyzed target degradation
659    during CRISPR RNA-guided interference. *Proceedings of the National Academy*
660    *of Sciences* **111,** 6618–6623 (2014).

661 21. Westra, E. R. *et al.* CRISPR Immunity Relies on the Consecutive Binding and
662    Degradation of Negatively Supercoiled Invader DNA by Cascade and Cas3. *Mol*
663    *Cell* **46,** 595–605 (2012).

664 22. Levy, A. *et al.* CRISPR adaptation biases explain preference for acquisition of
665    foreign DNA. **520,** 505–510 (2015).

666 23. Cady, K. C., Bondy-Denomy, J., Heussler, G. E., Davidson, A. R. & O'Toole, G. A.
667    The CRISPR/Cas Adaptive Immune System of *Pseudomonas aeruginosa*
668    Mediates Resistance to Naturally Occurring and Engineered Phages. 1–40
669    (2012).

670 24. Haurwitz, R. E., Jinek, M., Wiedenheft, B., Zhou, K. & Doudna, J. A. Sequence-
671    and Structure-Specific RNA Processing by a CRISPR Endonuclease. *Science*
672    **329,** 1355–1358 (2010).

673 25. Wiedenheft, B. *et al.* Structural basis for DNase activity of a conserved protein
674    implicated in CRISPR-mediated genome defense. *Structure* **17,** 904–912 (2009).

675 26. Wiedenheft, B. *et al.* RNA-guided complex from a bacterial immune system
676    enhances target recognition through seed sequence interactions. *Proceedings of*
677    *the National Academy of Sciences* **108,** 10092–10097 (2011).

678 27. Rollins, M. F., Schuman, J. T., Paulus, K., Bukhari, H. S. T. & Wiedenheft, B.
679    Mechanism of foreign DNA recognition by a CRISPR RNA-guided surveillance
680    complex from Pseudomonas aeruginosa. *Nucleic Acids Research* (2015).
681    doi:10.1093/nar/gkv094

682 28. Rollins, M. F. *et al.* Cas1 and the Csy complex are opposing regulators of Cas2/3
683    nuclease activity. *Proceedings of the National Academy of Sciences* **23,**

684  201616395–E5121 (2017).

685 29. Chowdhury, S. *et al.* Structure Reveals Mechanisms of Viral Suppressors that
686  Intercept a CRISPR RNA-Guided Surveillance Complex. *Cell* **169,** 47–57.e11
687  (2017).

688 30. Zegans, M. E. *et al.* Interaction between bacteriophage DMS3 and host CRISPR
689  region inhibits group behaviors of Pseudomonas aeruginosa. *J. Bacteriol.* **191,**
690  210–219 (2009).

691 31. Cady, K. C. & O'Toole, G. A. Non-identity-mediated CRISPR-bacteriophage
692  interaction mediated via the Csy and Cas3 proteins. *J. Bacteriol.* **193,** 3433–3445
693  (2011).

694 32. van Belkum, A. *et al.* Phylogenetic Distribution of CRISPR-Cas Systems in
695  Antibiotic-Resistant Pseudomonas aeruginosa. *mBio* **6,** e01796–15 (2015).

696 33. Westra, E. R. *et al.* Parasite Exposure Drives Selective Evolution of Constitutive
697  versus Inducible Defense. *Curr. Biol.* **25,** 1043–1049 (2015).

698 34. van Houte, S. *et al.* The diversity-generating benefits of a prokaryotic adaptive
699  immune system. *Nature* **532,** 385–388 (2016).

700 35. Bondy-Denomy, J., Pawluk, A., Maxwell, K. L. & Davidson, A. R. Bacteriophage
701  genes that inactivate the CRISPR/Cas bacterial immune system. *Nature* **493,**
702  429–432 (2013).

703 36. Pawluk, A., Bondy-Denomy, J., Cheung, V. H. W., Maxwell, K. L. & Davidson, A.
704  R. A new group of phage anti-CRISPR genes inhibits the type I-E CRISPR-Cas
705  system of Pseudomonas aeruginosa. *mBio* **5,** e00896–e00896–14 (2014).

706 37. Bondy-Denomy, J. *et al.* Multiple mechanisms for CRISPR-Cas inhibition by anti-
707  CRISPR proteins. *Nature* **526,** 136–139 (2015).

708 38. Pawluk, A. *et al.* Inactivation of CRISPR-Cas systems by anti-CRISPR proteins in
709  diverse bacterial species. *Nature Microbiology* **1,** 1–6 (2016).

710 39. Cady, K. C. & O'Toole, G. Non-Identity-Mediated CRISPR-Bacteriophage
711  Interaction Mediated via the Csy and Cas3 Proteins. **193,** 3433–3445 (2011).

712 40. Cady, K. C., Bondy-Denomy, J., Heussler, G. E., Davidson, A. R. & O'Toole, G. A.
713  The CRISPR/Cas adaptive immune system of Pseudomonas aeruginosa
714  mediates resistance to naturally occurring and engineered phages. **194,** 5728–
715  5738 (2012).

716 41. Borges, A. L. *et al.* Bacteriophage Cooperation Suppresses CRISPR-Cas3 and
717  Cas9 Immunity. *Cell* **174,** 917–925.e10 (2018).

718 42. Damron, F. H. *et al.* Analysis of the Pseudomonas aeruginosa Regulon Controlled
719  by the Sensor Kinase KinB and Sigma Factor RpoN. **194,** 1317–1330 (2012).

720 43. Martin, D. W. *et al.* Mechanism of conversion to mucoidy in Pseudomonas
721  aeruginosa infecting cystic fibrosis patients. *Proc Natl Acad Sci USA* **90,** 8377–
722  8381 (1993).

723 44. Qiu, D., Eisinger, V. M., Rowen, D. W. & Yu, H. D. Regulated proteolysis controls
724  mucoid conversion in Pseudomonas aeruginosa. *Proc Natl Acad Sci USA* **104,**
725  8107–8112 (2007).

726 45. Chand, N. S. *et al.* The Sensor Kinase KinB Regulates Virulence in Acute
727  Pseudomonas aeruginosa Infection. **193,** 2989–2999 (2011).

728 46. Chand, N. S., Clatworthy, A. E. & Hung, D. T. The Two-Component Sensor KinB
729  Acts as a Phosphatase To Regulate Pseudomonas aeruginosa Virulence. **194,**
730  6537–6547 (2012).

731 47. Damron, F. H., Qiu, D. & Yu, H. D. The Pseudomonas aeruginosa Sensor Kinase
732  KinB Negatively Controls Alginate Production through AlgW-Dependent MucA
733  Proteolysis. **191,** 2285–2295 (2009).

734 48. Cezairliyan, B. O. & Sauer, R. T. Control of Pseudomonas aeruginosa AlgW

protease cleavage of MucA by peptide signals and MucB. *Molecular Microbiology*
72, 368–379 (2009).
49. Schurr, M. J., Yu, H., Martinez-Salazar, J. M., Boucher, J. C. & Deretic, V. Control
of AlgU, a member of the sigma E-like family of stress sigma factors, by the
negative regulators MucA and MucB and Pseudomonas aeruginosa conversion to
mucoidy in cystic fibrosis. *J. Bacteriol.* **178,** 4997–5004 (1996).
50. Tart, A. H., Blanks, M. J. & Wozniak, D. J. The AlgT-dependent transcriptional
regulator AmrZ (AlgZ) inhibits flagellum biosynthesis in mucoid, nonmotile
Pseudomonas aeruginosa cystic fibrosis isolates. *J. Bacteriol.* **188,** 6483–6489
(2006).
51. Baynham, P. J. & Wozniak, D. J. Identification and characterization of AlgZ, an
AlgT-dependent DNA-binding protein required for Pseudomonas aeruginosa algD
transcription. *Molecular Microbiology* **22,** 97–108 (1996).
52. Wozniak, D. J. & Ohman, D. E. Transcriptional analysis of the Pseudomonas
aeruginosa genes algR, algB, and algD reveals a hierarchy of alginate gene
expression which is modulated by algT. *J. Bacteriol.* **176,** 6007–6014 (1994).
53. Landsberger, M. *et al.* Anti-CRISPR Phages Cooperate to Overcome CRISPR-
Cas Immunity. *Cell* **174,** 908–916.e12 (2018).
54. Kong, W. *et al.* ChIP-seq reveals the global regulator AlgR mediating cyclic di-
GMP synthesis in Pseudomonas aeruginosa. *Nucleic Acids Research* **43,** 8268–
8282 (2015).
55. Lizewski, S. E. *et al.* Identification of AlgR-regulated genes in Pseudomonas
aeruginosa by use of microarray analysis. *J. Bacteriol.* **186,** 5672–5684 (2004).
56. Schulz, S. *et al.* Elucidation of sigma factor-associated networks in Pseudomonas
aeruginosa reveals a modular architecture with limited and function-specific
crosstalk. *PLoS Pathog* **11,** e1004744 (2015).
57. Wozniak, D. J. *et al.* Alginate is not a significant component of the extracellular
polysaccharide matrix of PA14 and PAO1 Pseudomonas aeruginosa biofilms.
Proc Natl Acad Sci USA **100,** 7907–7912 (2003).
58. Pryor, E. E. *et al.* The transcription factor AmrZ utilizes multiple DNA binding
modes to recognize activator and repressor sequences of Pseudomonas
aeruginosa virulence genes. *PLoS Pathog* **8,** e1002648 (2012).
59. Pratama, A. A. & van Elsas, J. D. A novel inducible prophage from the
mycosphere inhabitant Paraburkholderia terrae BS437. *Sci Rep* **7,** 9156 (2017).
60. Boucher, J. C., Yu, H., Mudd, M. H. & Deretic, V. Mucoid Pseudomonas
aeruginosa in cystic fibrosis: characterization of muc mutations in clinical isolates
and analysis of clearance in a mouse model of respiratory infection. *Infection and*
Immunity **65,** 3838–3846 (1997).
61. Martin, D. W., Holloway, B. W. & Deretic, V. Characterization of a locus
determining the mucoid status of Pseudomonas aeruginosa: AlgU shows
sequence similarities with a Bacillus sigma factor. *J. Bacteriol.* **175,** 1153–1164
(1993).
62. Damron, F. H., Qiu, D. & Yu, H. D. The Pseudomonas aeruginosa sensor kinase
KinB negatively controls alginate production through AlgW-dependent MucA
proteolysis. *J. Bacteriol.* **191,** 2285–2295 (2009).
63. Jones, A. K. *et al.* Activation of the Pseudomonas aeruginosa AlgU regulon
through mucA mutation inhibits cyclic AMP/Vfr signaling. **192,** 5709–5717 (2010).
64. Heilmann, S., Sneppen, K. & Krishna, S. Sustainability of virulence in a phage-
bacterial ecosystem. *Journal of Virology* **84,** 3016–3022 (2010).
65. Dötsch, A. *et al.* The Pseudomonas aeruginosaTranscriptional Landscape Is
Shaped by Environmental Heterogeneity and Genetic Variation. *mBio* **6,** e00749–

15–10 (2015).
66. Erdmann, J., Preusse, M., Khaledi, A., Pich, A. & Häussler, S. Environment-driven
changes of mRNA and protein levels in Pseudomonas aeruginosa. *Environmental*
Microbiology **20,** 3952–3963 (2018).
790