CRISPR-Cas immunity repressed by a biofilm-activating pathway in *Pseudomonas aeruginosa*

Adair L. Borges#, Bardo Castro#, Sutharsan Govindarajan, Tina Solvik, Veronica Escalante, Joseph Bondy-Denomy*

Department of Microbiology & Immunology and the Quantitative Biosciences Institute, University of California, San Francisco, San Francisco, CA 94158

#Equal contribution

*Corresponding author: joseph.bondy-denomy@ucsf.edu

CRISPR-Cas systems are adaptive immune systems that protect bacteria from bacteriophage (phage) infection. To provide immunity, RNA-guided protein surveillance complexes recognize foreign nucleic acids, triggering their destruction by Cas nucleases. While the essential requirements for immune activity are well understood, the physiological cues that regulate CRISPR-Cas expression are not. Here, a forward genetic screen identifies a two-component system (KinB/AlgB), previously characterized in regulating *Pseudomonas aeruginosa* virulence and biofilm establishment, as a regulator of the biogenesis of the Type I-F CRISPR-Cas surveillance complex. Downstream of the KinB/AlgB system, activators of biofilm production AlgU (a σ^E orthologue) and AlgR, act as repressors of CRISPR-Cas surveillance complex expression during planktonic and surface-associated growth. AmrZ, another biofilm activator, functions as a surface-specific repressor of CRISPR-Cas activity. *Pseudomonas* phages and plasmids have taken advantage of this regulatory scheme, and carry hijacked homologs of AmrZ, which are functional CRISPR-Cas repressors. This suggests that while CRISPR-Cas regulation may be important to limit self-toxicity, endogenous repressive pathways represent a vulnerability for parasite manipulation.

Clustered regularly interspaced short palindromic repeats (CRISPR) and CRISPR-associated (cas) genes are RNA-guided nucleases found in nearly half of all bacteria[1]. CRISPR-Cas systems are mechanistically diverse, with six distinct types (I-VI) identified, based on signature genes, mechanisms of action, and the type of nucleic acid target[1].

Our strong understanding of the basic components to enable sequence specific DNA
and RNA cleavage have enabled functional transplantation into heterologous bacterial[2,3]
and eukaryotic[4,5] hosts. Given that many Cas proteins encode nucleases[6] the fine-tuned
regulation of these systems to avoid toxicity is likely a key factor to enable safe retention
of a CRISPR-Cas immune system[7]. Multiple signals have been shown to activate
CRISPR-Cas function in diverse organisms, such as quorum sensing[8,9], temperature[10],
membrane stress[11,12], altered host metabolite levels[13,14], and phage infection[15-18].
However, relatively little is known regarding the factors and/or signals that serve to
temper CRISPR-Cas activity and mitigate the risk of acquiring and expressing a
nucleolytic immune system.
45
Type I CRISPR-Cas systems are comprised of a multi-subunit RNA-guided surveillance
complex, a trans-acting nuclease (Cas3)[19-21], and proteins dedicated to spacer
acquisition, Cas1 and Cas2[22,23]. *Pseudomonas aeruginosa* has become a powerful
model organism for studying Type I CRISPR-Cas mechanisms[24-29], functions[3,23,30,31],
evolution[32-34], and interactions with phages utilizing anti-CRISPR proteins[35-38]. The *P.*
aeruginosa strain PA14 possesses an active Type I-F CRISPR-Cas immune system that
naturally targets many phages. The surveillance complex is composed of four Cas
proteins (Csy1, Csy2, Csy3, and Csy4), which are encoded together in a single
operon[26]. The surveillance complex is guided by CRISPR RNAs (crRNAs) originating
from one of two chromosomally encoded CRISPR arrays[26]. Finally, a separate operon is
responsible for the production of Cas1 and a Cas2-3 fusion protein, the latter of which
mediates nucleolytic destruction of target DNA.
58
To discover regulators controlling the transcription of the CRISPR-Cas surveillance
complex in *P. aeruginosa*, we utilized a *P. aeruginosa* strain PA14 engineered to
express *lacZ* in place of the *csy3* gene (*csy3::lacZ*)[39]. The *csy3::lacZ* reporter strain was
subjected to transposon mutagenesis and colonies isolated and screened on X-gal
plates. ~40,000 colonies were visually examined for increased or decreased levels of β-
galactosidase. Multiple independent insertions were identified that abolished LacZ levels
within *lacZ* and upstream genes (*csy1* and *csy2*), indicating successful mutagenesis and
screening. Thirty distinct mutants with transposon insertions outside of this region, with
altered β-galactosidase levels, were isolated and mapped. These mutants are listed in
Supplementary Table 1 with β-galactosidase levels quantified for mutants that grew well

in culture. Four independent insertions were identified in a single gene, *kinB*, which

resulted in apparent mucoidy plating morphology and decreased β-galactosidase

production (Supplementary Fig. 1a). Quantification of β-galactosidase levels in liquid

growth experiments revealed ~50% less *csy3::lacZ* expression compared to wild-type

(WT) PA14 for each mutant (Supplementary Table 1, Supplementary Fig. 1b). We next

focus on *kinB* (a sensor kinase/phosphatase) because it was the only gene with >1

independent transposon insertion and displayed the largest β-galactosidase activity

change.

To determine the consequences of *kinB* disruption and decreased *csy* expression on

CRISPR-Cas function, we measured the ability of *kinB::Tn* insertions to limit the

replication of phage. Phages used to assay activity are: DMS3$_{acrIE3}$ which is an

untargeted control phage, DMS3m$_{acrIE3}$ [40], which is fully targeted by the PA14 Type I-F

CRISPR-Cas system, and phage DMS3m $_{acrIF4}$, which is partially targeted, by virtue of

encoding a "weak" anti-CRISPR, *acrIF4,* that binds to the surveillance complex to inhibit

CRISPR-Cas function[35,37,41]. The *kinB::Tn* strains remained resistant to DMS3m$_{acrIE3}$

infection, but DMS3m$_{acrIF4}$ formed 10-fold more plaques relative to WT, demonstrating

attenuated CRISPR-Cas function (Figure 1a, Supplementary Fig. 1c). This defect was

complemented by expression of *kinB in trans* (Supplementary Fig. 1c). Growth of control

phage DMS3$_{acrIE3}$ was not impacted in the absence of KinB (Figure 1a). Furthermore,

two other targeted phages, JBD26 (naturally possessing *acrIF4*) and JBD25 (a partially

targeted phage with no Acr) also showed increased replication in the *kinB:Tn* strain

(Supplementary Fig. 1d) relative to WT PA14. Replication of a phage with a weak anti-

CRISPR or one that is targeted by a less active spacer is therefore a sensitive

barometer for perturbations in CRISPR-Cas levels. Together, these data confirm that in

the absence of *kinB, csy* gene expression and phage targeting are decreased.

KinB is a sensor kinase/phosphatase in a two-component system with response

regulator AlgB. The KinB/AlgB system has a large regulon within *P. aeruginosa*, but is

well-known for its role in biofilm regulation[42]. In the absence of KinB function, the activity

of the transcription factor AlgU (a σ^E orthologue and an activator of biofilm formation) is

enhanced[43,44]. This activation of AlgU manifests solely due to the lack of KinB

phosphatase activity and the resulting accumulation of the phosphorylated form of the

response regulator AlgB (P-AlgB). The phosphorylation of AlgB has been attributed to

unknown kinases[45,46] (Fig. 1b). The resulting accumulation of P-AlgB activates the

periplasmic protease AlgW (a DegS homolog), which degrades MucA, liberating sigma

factor AlgU[47,48,49] (Fig. 1b). AlgU positively regulates many factors involved in biofilm

production, including AlgR, AlgD, AlgB, and AmrZ[50-52].

To determine if this KinB-controlled biofilm activation pathway also regulates CRISPR-

Cas expression, we tested the ability of phosphatase and kinase dead KinB mutants to

complement an in-frame Δ*kinB* deletion, using the replication of the partially-targeted

DMS3m$_{acrIF4}$ phage as a metric for CRISPR-Cas activity. WT *kinB* or kinase inactive

H385A *kinB* (Fig. 1c) complemented the CRISPR defect, indicating KinB kinase activity

is dispensable for CRISPR-Cas activation. However, a P390S *kinB* mutant incapable of

dephosphorylating the response regulator AlgB did not complement, and in fact

decreased CRISPR-Cas activity further (Fig. 1c). This suggests that CRISPR-Cas

expression is inhibited by the accumulation of P-AlgB. A Δ*kinB*Δ*algB* double mutant

restored CRISPR-Cas targeting to WT levels (Supplementary Fig. 2a), confirming the

role of this signaling pathway. A strain lacking *algB* (Δ*algB*) or possessing a D59N

mutant that cannot be phosphorylated had elevated CRISPR-Cas activity, supporting the

repressive role of P-AlgB (Supplementary Fig. 2b). Together, these data suggest that

KinB-mediated dephosphorylation of AlgB lifts CRISPR-Cas inhibition, and conversely

accumulation of high levels of P-AlgB (achieved in Δ*kinB* or *kinB P390S*) leads to

CRISPR-Cas repression.

To assess CRISPR-Cas regulation downstream of P-AlgB, anti-phage immunity was

assayed in Δ*algU* and Δ*algR* backgrounds, revealing enhanced targeting of DMS3m$_{acrIF4}$

but not control phage DMS3$_{acrIE3}$ in each knockout (Fig. 1d). Complementation with *algR*

and *algU* restored CRISPR-Cas levels in their respective knockout backgrounds (Fig.

1d). This demonstrates that these proteins repress CRISPR-Cas immunity. Double

knockouts of each gene combined with Δ*kinB* each resembled the respective Δ*algU* and

Δ*algR* single knockouts, consistent with these factors acting downstream of KinB

(Supplementary Fig. 2a). The observed plaquing defects seen with the DMS3m$_{acrIF4}$

phage are CRISPR-dependent, as double knockouts (*kinB, algB, algU, algR* mutants

combined with *csy3::lacZ*) revealed plaquing equivalent to *csy3::lacZ* alone

(Supplementary Fig. 2c).

Phages that use anti-CRISPRs to inhibit CRISPR-Cas immunity must cooperatively

infect the same cell in order to produce a sufficient dose of anti-CRISPR to permit phage

replication[41,53]. The multiplicity of infection (MOI) required to overwhelm immunity is

dependent on the affinity of the anti-CRISPR for the CRISPR-Cas machinery and the

intracellular concentration of CRISPR-Cas complexes. To understand how this

regulatory pathway controls phage cooperation requirements on a population level by

potentially shifting the intracellular concentration of CRISPR-Cas complexes, liquid

cultures of $\Delta algU$ and $\Delta kinB$ mutants were infected with a virulent derivative of phage

DMS3m$_{acrIF4}$ at both high (where the WT cells are sensitive) and low (where the WT cells

are resistant) MOI (Figure 1e). The $\Delta algU$ mutant was able to limit phage replication

relative to WT and continue growing when infected with concentration of phages

sufficiently high to overwhelm WT immunity (10^6 PFU), indicating increased levels of

CRISPR-Cas machinery (Fig 1e). Conversely, when infected with a low MOI of phages

(10^4 PFU), all bacterial genotypes were able to grow to high cell density, however the

$\Delta kinB$ mutant permitted increased phage replication relative to WT, demonstrating a

decreased level of immune system components (Fig 1f). Importantly, these changes in

phage replication requirements were not seen in the absence of CRISPR-Cas immunity,

as phage replication in double knockouts *algU csy3::lacZ* and *kinB csy3::lacZ*

resembles *WT csy3::lacZ (*Supplementary Fig. 3a-d, f)*. In contrast, $\Delta algR$ demonstrated

CRISPR-independent phage resistance during liquid culture infection, and phage

cooperation requirements could not be reliably measured for this mutant (Supplementary

Fig. 3e). Taken together, these data demonstrate that a well-characterized biofilm

activation pathway controlled by alternate sigma factor, AlgU, represses CRISPR-Cas

immunity in *P. aeruginosa.*

As our screen was focused on finding regulators of the *csy* operon, and as Cas3 is

expressed at low levels and is furthermore subject to post-translational control by

Cas1[28], we hypothesized the KinB/AlgB pathway was controlling anti-phage immunity via

modulation of *csy* complex levels. To specifically measure the anti-phage activity of the

csy complex in the AlgU regulatory pathway mutants, we developed a Cas3-indepedent

bioassay to read out the activity of the surveillance complex in the cell. Through the

rational design of crRNAs to target an early phage phage promoter (P_E_1, P_E_2), we

observed inhibition of phage replication in a *P. aeruginosa* strain with a nuclease dead

Cas3 (Cas3$_{D124A}$, Figure 2a). This CRISPR-based transcriptional interference (CRISPRi)

effect was also seen when infecting with a phage that expressed the inhibitor of Cas3
recruitment, AcrIF3, but not an inhibitor that blocks Csy complex-phage DNA binding[37],
AcrIF1 (Figure 2a). CRISPRi was mediated by crRNAs targeting phage promoters,
including one (P_E_2) that completely limited phage replication in the absence of Cas3
activity, while an ORF-targeting crRNA was ineffective. We selected P_E_1 as a
moderately-functional CRISPRi spacer and expressed it in various mutant backgrounds
generated in this study. We observed decreased CRISPRi activity against phage
DMS3m$_{acrIF3}$ in the ΔkinB background, but increased CRISPRi-mediated phage
repression in Δ*algR* and Δ*algU,* (Figure 2b), demonstrating that modulation of *csy* gene
expression by this pathway impacts phage targeting in a Cas3-independent manner. To
confirm that this pathway controls the levels of *csy* complex in the cell, we fused an N-
terminal sfCherry tag to the endogenous *csy1* gene in the mutant backgrounds. We
observed decreased levels of red fluorescence in Δ*kinB* and increased levels in Δ*algR*
and Δ*algU* relative to WT, demonstrating that this pathway controls the levels of *csy*
complex in the bacterial cell (Fig. 2c).

AlgU and AlgR are proteins that regulate hundreds of genes in *P. aeruginosa*[54-56],
ultimately repressing acute virulence factors (e.g. pyocyanin) and activating chronic
virulence factors (e.g. biofilm production). To identify downstream CRISPR-Cas
regulators, we considered the large AlgU regulon[56], but focused on a factor involved in
biofilm production, whose expression is AlgU-dependent, *amrZ*[51,57]. AmrZ is a ribbon-
helix-helix type transcription factor, and acts as both a repressor and activator of
transcription in *Pseudomonas*[58]. We generated a knockout of *amrZ* and observed a
CRISPR-dependent (Supplementary Fig. 2c) decrease in DMS3m$_{acrIF4}$ plaque formation
(Fig. 3a). This was complemented when *amrZ* was expressed *in trans* (Fig. 3a),
indicating that AmrZ is also a repressor of CRISPR-Cas immunity. However, when we
measured phage cooperation requirements in the Δ*amrZ* strain, phage replication and
immune protection did not differ substantially from from WT (Fig. 3b,c). The most
obvious difference between the phage plaque assay and the assay to measure phage
cooperation is that a plaque assay is performed on solid plates whereas phage
cooperation requirements were measured in liquid culture. To test if AmrZ was a
surface-specific repressor of CRIPSR-Cas immunity, we measured the levels of Csy
complex during surface association and planktonic growth in WT and Δ*amrZ* cells using
the endogenous Csy1-sfCherry reporter. In WT cells, the levels of Csy complex were

attenuated during surface-association relative to planktonic growth, but in the absence of
AmrZ, Csy complex levels during surface association increased to levels comparable to
those in planktonic growth (Fig. 3d). Deletion of *amrZ* did not impact Csy complex levels
during planktonic growth (Fig. 3e). To artificially increase the levels of AmrZ during
planktonic growth, we ectopically expressed AmrZ from a high copy plasmid. In this
scenario, high levels of AmrZ were sufficient to repress Csy complex expression,
suggesting that low AmrZ activity in planktonic growth underlies its surface-specific
control of CRISPR-Cas (Fig. 3f). In contrast to AmrZ, the transcriptional repressors AlgU
and AlgR were unable to repress Csy complex levels when heterologously expressed
(Fig. 3f), likely because they are master regulators and many redundant mechanisms
are in place to control their activity. Taken together, these data demonstrate that AmrZ is
a uniquely potent surface-specific repressor of the *csy* operon.

We next considered if phages and other mobile genetic parasites naturally antagonized
by CRISPR-Cas immunity had evolved mechanisms to manipulate this repressive
pathway. Inspired by a recent discovery of a *Paraburkholderia* phage that carried a
distant homolog of AmrZ[59], we searched the NCBI database for AmrZ homologs on
Pseudomonas mobile genetic elements (MGE). Surprisngly, we identified 14 diverse
Pseudomonas genetic parasites carrying AmrZ homologs (Supplementary Table 2).
These MGEs included obligately lytic and temperate Myoviridae, temperate
Siphoviridae, and plasmids. AmrZ has been structurally characterized in complex with
operator DNA[58], and these mobile AmrZ homologs showed perfect conservation of
critical DNA-interacting residues in the ribbon-helix-helix domain, suggesting conserved
binding specificity (Fig. 4a, b, red residues/arrowheads). To test if these mobilized AmrZ
variants were capable of regulating CRISPR-Cas activity in *Pseudomonas aeruginosa*,
we synthesized 2 phage-borne *amrZ* homologs and tested them for their ability to
complement the Δ*amrZ* strain. Both homologs complemented the Δ*amrZ* mutant,
indicating they were active in the PA14 transcriptional network and were *bona fide* AmrZ
orthologs (Fig. 4c). Next, each phage gene was expressed in WT cells, revealing one
homolog (*amrZ_{phi3}*) limited the expression a Csy1-sfCherry reporter. This suggests the
possibility that *amrZ* has been hijacked for its capacity to inhibit CRISPR-Cas expression
(Fig. 4d). Thus, *amrZ* acts on both sides of the battle between bacteria and mobile
genetic elements.

Discussion

The factors that govern expression of bacterial processes are highly variable across
species, reflecting niche-specific adaptations. CRISPR-Cas immune systems limit the
replication of foreign genetic elements, but few reports address how the regulation of
CRISPR-Cas impacts phage targeting. Here, we utilized a genetic screen that identified
KinB as a positive regulator of CRISPR-Cas function in *P. aeruginosa.* This sensor
kinase/phosphatase operates in a two-component system with AlgB that regulates a
well-characterized biofilm-stimulatory pathway, though the ligand for KinB remains
unknown. Removal of KinB or inactivation of its phosphatase activity leads to the
accumulation of P-AlgB, which stimulates the activity of CRISPR-Cas repressors AlgU,
AlgR, and AmrZ. This pathway is well-studied due to the recurrent isolation of alginate-
overproducing (mucoid) *P. aeruginosa* from the lungs of cystic fibrosis patients, with
mutations in *mucA*[60] and *kinB*[47,61] that lead to the constitutive activation of AlgU. This
biofilm pathway generates the characteristic mucoidy phenotype of CF *P. aeruginosa*
isolates[43,62,63], which we also observed in *kinB:Tn* mutants (Supplementary Fig. 1a).

This pathway is well described in its capacity to activate biofilm production, and here we
show that this transcriptional network also functions to dampen CRISPR-Cas activity in
non-biofilm conditions. In WT cells, KinB phosphatase activity increases CRISPR-Cas
function, though the effect is relatively mild in our experimental conditions. Relieving
transcriptional repression via deletion of *algU* or *algR* however leads to a hyper-
activation of the CRISPR-Cas immune response. We further demonstrate that *amrZ* acts
in this pathway to specifically repress CRISPR-Cas when cells are surface-associated,
leading to lower levels of *csy* complex during surface-association relative to planktonic
growth. Phages and plasmids have taken advantage of this regulatory scheme, and
some *Pseudomonas* genetic parasites carry hijacked copies of the CRISPR-Cas
repressor *amrZ*. We show that these mobilized *amrZ* homologs retain their capacity to
transcriptionally repress CRISPR-Cas, demonstrating a dual role for this gene in the
host-pathogen arms race between bacteria and their parasites.

Considering the physiology of *P. aeruginosa* can put these findings into context with a
phage and CRISPR-Cas-centric view. During exponential growth is when we and others
observe CRISPR-Cas activation[8,9], and is also when phage infection risk is likely high
(i.e. metabolically active, well-mixed planktonic cultures at high cell density[33]).

Conversely, surface-association lessens phage infection risk, as spatial structure limits
phage diffusion and prevents a phage bloom from overtaking the entire bacterial
population[64]. Though we do not measure phage sensitivity in a biofilm here, biofilm
production likely provides an even greater level of intrinsic phage resistance, due to high
levels of spatial stratification and abundant polysaccharide secretion.

Supporting our proposal that *P. aeruginosa* surface-association/biofilm formation and
CRISPR-Cas expression are inversely correlated are two independent observations: **i)**
our analysis of a previously published PA14 RNAseq data set[65] and proteomic data set[66]
revealed activation of CRISPR-Cas expression in exponential phase, and strong
transcriptional repression during stationary phase and biofilm growth (Supplementary
Fig. 4a). Interestingly Cas proteins are still detected in stationary phase and biofilm
growth, suggesting the cells retain some immune-competency even after transcriptional
shutdown (Supplementary Fig. 4b), and **ii)** previous studies on the *P. aeruginosa*
CRISPR-Cas system have revealed that the genome is hyper-sensitive to CRISPR-
induced DNA damage during surface-association and biofilm formation, leading to cell
death when a mismatched prophage sequence target is present in the chromosome[30,31].
This suggests that CRISPR auto-immunity costs are also dependent on the growth state
and physical environment of the cell, and that CRISPR-Cas regulation in *P. aeruginosa*
is tuned to reflect this intrinsic sensitivity.

Here, we report the identification of a CRISPR-Cas repressive pathway in *P. aeruginosa*.
Phages bearing distinct anti-CRISPRs are employed in bioassays to read out the activity
of the CRISPR-Cas surveillance complex. Using a CRISPRi phage-repression assay,
coupled with translational reporters, we conclude that *algU*, *algR*, and *amrZ* control the
production of the CRISPR surveillance complex (*csy* complex), which is where the
screen was focused. Interestingly, our observations show that CRISPR-Cas activity in *P.*
aeruginosa does not operate at full strength, despite the capacity of a hyper-activated
immune system to severely limit the replication of phages relying on anti-CRISPRs for
survival. We speculate that the ability to control CRISPR-Cas activity during lifestyle
transitions may be essential for *P. aeruginosa* to safely maintain a CRISPR-Cas system
by limiting self-toxicity. Furthermore, we reveal an unexpected cost to negative
regulation of CRISPR-Cas immunity in our discovery of MGE-encoded CRISPR-Cas
repressors: though CRISPR-Cas control is likely essential for safe retention of the
immune system, the evolution of potent mechanisms of CRISPR-Cas repression has

created an Achilles Heel that is exploited by genetic parasites.

Acknowledgements
The Bondy-Denomy lab was supported by the University of California, San Francisco
Program for Breakthrough in Biomedical Research, funded in part by the Sandler
Foundation, and an NIH Office of the Director Early Independence Award (DP5-
OD021344), and R01GM127489. We thank Deborah Hung's lab for providing Δ*kinB,*
Δ*algR,* Δ*algU* strains and associated mutants. George O'Toole's lab provided the
csy3::lacZ PA14 strain.

Author Contributions
J.B.D., A.L.B, and B.C. formulated study design and plans. A.L.B. designed and
conducted bacteriophage plaque assays, liquid infection assays, and CRISPRi assays,
performed. B.C. conducted the genetic screen, isolated and constructed bacterial
mutants, conducted LacZ expression profiling, and performed bacteriophage plaque
assays. CRISPR-Cas expression profiling, and conducted bioinformatics analyses. S.G.
designed, constructed, and validated sfCherry reporter strains. T.S. conducted CRISPRi
assays and V.E. assisted in establishing fluorescent reporter assays. J.B.D. and A.L.B
wrote the manuscript.

Competing Interests
J.B.-D. is a scientific advisory board member of SNIPR Biome and Excision
Biotherapeutics and a scientific advisory board member and co-founder of Acrigen
Biosciences.

Materials & Correspondence
Correspondence should be directed to joseph.bondy-denomy@ucsf.edu

Methods
Bacterial strains and bacteriophages. *P. aeruginosa* UCBPP-PA14 (PA14) strains
and *E. Coli* strains (Sup. Table) were grown on lysogeny broth (LB) agar or liquid at 37
°C. Media was supplemented with gentamicin (50 µg ml^{-1} for *P. aeruginosa* and 30 µg ml^{-}
^{1} for *E. Coli*) to maintain the pHERD30T plasmid or carbenicillin (250 µg ml^{-1}) for *P.*
aeruginosa or ampicillin (100 µg ml^{-1}) for *E. coli* containing the pHERD20T plasmids.

pHERD plasmids were induced with 0.1% arabinose. Bacteriophages stocks were
prepared as described previously[40]. In brief, 3 ml of SM buffer was added to plate
lysates of the desired purified bacteriophage and incubated at room temperature for 15
minutes. SM buffer containing phages was collected and 100 µl of chloroform was
added. This was centrifuged at 10,000 x g for 5 minutes and supernatant containing
phages was transferred to a screw cap storage tube and incubated at 4 $^{\circ}$C.

Tn mutagenesis screen. The *csy*::*lacZ* reporter strain was subjected to transposon
mutagenesis and colonies were isolated on plates containing X-gal (5-bromo-4-chloro-3-
indolyl-β-D-galactopyranoside). ~50,000 colonies were visually examined for increased
or decreased levels of β-galactosidase and insertions mapped by semi-random PCR. To
conduct transposon mutagenesis, overnights of PA14 *csy3::lacZ* and *E. coli* containing
the pBTK30 Tn suicide vector were mixed in a 1:2 ratio (donor : recipient) for
conjugation. A second control group containing PA14wt and *E. Coli* containing the
pBTK30 Tn suicide vector was set up in parallel. Mixed cells were centrifuged at 4,000 x
g for 10 minutes to pellet cells. 100 µl of resuspended conjugation pellet was then
spotted on LB agar plates and incubated at 37 $^{\circ}$C for 6hrs. Conjugation spots were
collected and resuspended in LB liquid media. Conjugation was then screened on an LB
agar plates supplemented with nalidixic acid (30 µg ml^{-1}) and gentamicin (50 µg ml^{-1}).
Surviving colonies containing Tn insertions were collected into 1ml of LB liquid media.
Serial dilution of were prepared and plated on LB agar plates supplemented with x-gal
(200 µg ml^{-1}) and gentamicin (50 µg ml^{-1}) and nalidixic acid (30 µg ml^{-1}). Plates were
incubated at 37 $^{\circ}$C for 24 hours to allow for colonies to change color. Colonies displaying
changed expression levels as compared to our positive control (PA14 csy3::lacZ no
pBTK30) were then isolated onto secondary LB agar plates with X-gal and gentamicin
and nalidixic acid at the stated concentrations. Genomic DNA (gDNA) was collected
from isolated single colonies by resuspending bacterial colonies in 0.02% SDS and
boiling the sample for 15 minutes. Samples were then centrifuged at 10,000 x g and
supernatants containing gDNA were collected. Semi-random PCR was then performed
using PCR primers listed on sup table_. PCR samples were sequenced and reads were
then matched to the *P. aeruginosa* UCBPP-PA14 genome using BLAST. Isolated
containing genes of interest were saved as stocks (s. table 1). Transcription changes
were then verified via modified β-galactosidase assay.

Plaque assays. Plaque assays were performed on LB agar plates (1.5% agar) with LB
top agar (0.7% agar), both supplemented $MgSO_4$ (10 mM final concentration) and
gentamicin (50 µg ml^{-1}) as needed for plasmid maintenance. Spot titrations were done by
mixing 150 µl of a *P. aeruginosa* overnight culture with 3 ml of top agar, which was
dispersed evenly on a LB $MgSO_4$ plate. 3 µl of 10-fold phage dilutions were then spotted
on the surface. Plates were incubated overnight at 30 ^{o}C. To count plaques, full plate
assays were used, except when CRISPR-targeting was so strong that discrete plaques
could not be accurately measured. In this case, spot titrations are shown. For full plate
assays, 3 µl of the selected phage dilution was incubated with 150 µl of *P. aeruginosa*
overnight for 15 minutes at 37 ^{o}C. 3 ml of top agar was then added and the mixture was
dispersed evenly on a LB $MgSO_4$ plate. Individual plaques were then counted to assess
differences in plaquing efficiency.

β-galactosidase assay. A modified version of the previously described β-galactosidase
assay was used to measure *lacZ* activity in transcriptional fusions. Bacterial cultures
were grown overnight at 37 ^{o}C. Cultures were then diluted 1:100 into LB liquid medium
supplemented with the desired antibiotic. Diluted cultures were then incubated at 37 ^{o}C
until the desired time point was reached. Culture density was measured with a
spectrophotometer (OD_{600}) and 200 µl of the sample was added 800 µl to
permeabilization solution (components listed in sup table XX). Cells were mixed via
inversion and vortexed for 1 minute to permeabilize the cells. 200 µl of ONPG (4 mg ml^{-}
^{1}) was added and samples were incubated for 27 minutes at 30 ^{o}C. Enzymatic reaction
was stopped by addition of 300 µl of 1M Na_2CO_3. Samples were centrifuged at 13,000 x
g for 5 minutes to remove debris and 200 µl of supernatant was moved to a 96-well plate
to read absorbance at 420 nm and at 550 nm. Miller units were calculated using the
Miller equation. 3 technical replicates per sample per experiment.

Phage transduction of *kinB::Tn* alleles. Transposon insertions in *kinB* from a
csy3::lacZ background were transduced into WT PA14 to enable testing of CRISPR-Cas
function with the same transposon insertion. Phage phiPA3 was used to infect the donor
strain (*csy3::lacZ, kinB::Tn*), on plates with top agar overlays, using ~10^4 PFU to
generate near confluent lysis. Plates were soaked in 3-4 mL of phage SM buffer and 2
mL collected over chloroform, vortexed, and pelleted to isolate transducing phage in the
supernatant. Lysates were used to infect recipient strains (WT PA14). ~10^8 PFU were

used to infect a culture at an MOI of 1. After 30 minutes of static incubation on the

bench, cultures were gently shaken at 37 ^{o}C for 20 min and then pelleted at 5000 x g.

Cells were washed twice with LB, and subsequently incubated at 37 ^{o}C for 1 hour to

allow recombination and gentamicin resistance outgrowth. Cultures were pelleted and

resuspended in 200 µL of LB, and plated on LB plates containing gentamicin. Controls

included uninfected cells and cells infected with phages not propagated on a gentamicin

resistant donor strain. Additionally, phage lysate was directly plated under selection to

confirm no residual donor strain in the phage preparation. Plates were incubated

overnight at 37 ^{o}C and their identity (i.e. CRISPR-Cas intact) confirmed with a plaque

assay using DMS3m as the target phage and PCR of the kinB locus.

Introduction of *csy3::lacZ P. aeruginosa* UCBPP-PA14 strains. The *lacZ* gene was

introduced into PA14 strains of interest via homologous recombination. Recombination

vector containing lacZ flanked by homology arms matching csy2 and csy4 was

introduced via conjugation. PA14 strains and *E. coli* containing vector were mixed at a

ratio of 1:2 (recipient:donor). Mixture was heat shocked at 42 ^{o}C for 10 min. Mating spot

was then plated on a LB agar plate and incubated overnight for 30 ^{o}C. Mating spot was

then collected, resuspended in 1 ml of LB liquid media and plated on VBMM plates

supplemented with 50 ug/mL gentamicin to select for colonies with the integrated

homology plasmid. Colonies were cultured overnight in LB in the absence of selection at

37 ^{o}C, and were then diluted and counterselected on no salt LB (NSLB) agar plates

supplemented with 15% sucrose. Surviving colonies were then grown on LB agar plates

supplemented with gentamicin and X-gal to check for *lacZ* insertion via color change and

lacZ insertion was further verified via PCR.

Generation of an PA14Δ*amrZ* strain using the endogenous I-F CRISPR-Cas

system. Complementary oligonucleotides encoding a crRNA targeting the *amrZ* gene of

PA14 were annealed and ligated into the multiple cloning site of the pHERD30T vector.

A fragment possessing homology arms flanking the desired mutation (500 bp upstream

and 500 bp downstream) around *amrZ* was cloned into a distinct location (NheI site) of

the same vector via Gibson assembly. The new plasmid containing both a crRNA and

homology region was introduced into WT PA14 via electroporation. Transformation

efficiency dropped dramatically in the presence of the crRNA due to the toxicity caused

by self-targeting. All surviving colonies had the desired clean deletion of the *amrZ* gene.

Deletions were confirmed by PCR of the region of interest and subsequent Sanger
sequencing of the amplicon. A 2000 bp region flanking *amrZ* was PCR amplified and
sequencing primers were designed to sequence both the deletion junction and outside of
the original 500 bp flanking regions to confirm the removal of the *amrZ* gene.

Liquid phage cooperation assay. Liquid phage infections were performed as
described in[41]. In brief, an overnight culture of cells was diluted 1:100 into fresh media,
and infected with virulent phage DMS3m$_{acrIF4}$ in biological triplicate in a 96 well Costar
plate. Cells were incubated at 37 °C with constant rotation and OD600 measured every 5
minutes in a Biotek H4 Synergy plate reader. Phage were harvested from each well and
quantified by plaque assay after 24 hours.

Generation of endogenous Csy1-sfCherry reporter. Endogenous Csy1-sfCherry
reporter was constructed similar to the construction of *csy3::lacZ*. We initially verified
that tagging of sfCherry at the N-terminus of *csy1* is functional, when expressed from a
plasmid. pMQ30-sfCherry-Csy1, which contains sfCherry sequence flanked by 657 bp
upstream of csy1 and 701 bp downstream of csy1 start codon, was cloned in pMQ30
plasmid between HindIII and BamHI sites using Gibson assembly. The Csy1-sfCherry
construct was introduced into PA14 strains of interest via homologous recombination.
Strains containing Csy1-sfCherry insertion were verified via PCR.

Csy1-sfCherry reporter profiling. **Liquid**: Cells were diluted 1:100 from an overnight
culture into fresh LB (with antibiotics and inducer if required), and grown for the indicated
number of hours in biological triplicate. 500 µl of each sample was then spun down at
8,000xg for 2 minutes, and resuspended in 500 µl of M9 media. Samples were loaded in
to a 96 well plate (150 µl /well) in technical triplicate and red fluorescence and OD600
were measured using a Biotek H4 Synergy. **Solid:** Cells were diluted 1:100 from an
overnight culture into fresh LB and 20 µl plated onto individual wells in biological
triplicate in a 24 well plate with each well containing solidified 1.5% LB Agar with
antibiotics and inducer if required. The 24 well plate was then covered with a breathable
Aeraseal, and incubated at 37 °C with no shaking. At the indicated timepoint, cells were
harvested by flooding each well with 500 µl of M9 buffer, and were spun down at
8,000xg for 2 minutes, and resuspended in 500 µl of M9 media. Sample were loaded in
to a 96 well plate (150 µl /well) in technical triplicate and red fluorescence (excitation

580 nm, emission 610 nm) and OD600 were measured using a Biotek H4 Synergy. To
calculate the relative fluorescence units for each sample, the background fluorescence
and background OD600 values obtained were subtracted from the sample values, and
the sample fluorescence was then normalized to the sample OD600.
477
AmrZ homolog discovery and characterization. BLASTp was used to search the
nonredundant protein database for AmrZ homologs (accession: ABJ12639.1) in
Pseudmonas sp. (taxid: 286) in May 2019. This homolog list (e value > 0.001) was then
examined for homologs found on phage or plasmid genomes. Representative homologs
were aligned using Clustal and the alignment visualized in Jalview, and key conserved
residues were mapped onto the structure in Pymol (PDB ID: 3QOQ). Select homologs
were synthesized (TWIST Biosciences) and cloned into the SacI/PstI site of the
arabinose-inducible plasmid pHERD30T using Gibson assembly. Vectors were
electroporated into *Pseudomonas aeruginosa* strains for functional testing, where they
were induced with 0.1% arabinose and maintained with 50 ug/mL gentimicin.
488
489
490

Figure 1

Figure 1: *A forward genetic screen identifies a role for a biofilm activating pathway in repressing CRISPR-Cas immunity.* **a**. Efficiency of immunity (EOI) against isogenic phages DMS3$_{acrIE3}$ (non-targeted) and DMS3m$_{acrIF4}$ (CRISPR-targeted). Plaque forming units (PFUs) were quantified on two independent *kinB* transposon mutants (*kinB::Tn1* and *kinB::Tn2*), then represented as a ratio relative to the number of PFUs measured on WT PA14. **b.** A cartoon summarizing the KinB/AlgB two component system and downstream effects, based on prior work, with CRISPR-Cas regulation added. **c**,**d** EOI measurements for indicated Δ*kinB,* Δ*algR,* and Δ*algU* strains with complementation. **e,f**. Growth curves of PA14 WT, Δ*algU*, or Δ*kinB* strains infected with either a high (e,10^6 PFU) or low (f, 10^4 PFU) MOI of virulent DMS3m$_{acrIF4}$. Phage replication was quantified as PFUs after 24 hours, and the efficiency of immunity expressed as a ratio of the number of PFUs harvested from the mutant strain compared to the number of PFUs obtained from WT. All data (OD600 and EOI measurements) are

  represented   as   the   mean   of   3   biological   replicates   +/-   SD.

Figure 2

Figure 2. *The kinB/algB pathway modulates Csy complex levels and activity.*

a. Spot titration of DMS3m$_{acrIF3}$ (F3) and DMS3m$_{acrIF1}$ (F1) on *cas3*$_{D124A}$ (dead Cas3) or

Δcsy3 (active Cas3, no Csy complex). Phages are targeted by natural spacer CR2_1, as

well as crRNAs designed to target DMS3m genome in positions designated on ORF

map. **b**. Spot titration of DMS3m$_{acrIF3}$ and DMS3m$_{acrIF1}$ phages on deletion mutants

expressing the indicated crRNA. **c.** Measurement of the normalized fluorescence levels

of Csy1-sfCherry reporter strains after 10 hours of growth in liquid culture. Fluorescence

measurements are represented as the mean of 3 biological replicates +/- SD.

Figure 3

Figure 3: *AmrZ is a surface-specific repressor of CRISPR-Cas immunity.* **a**.

Efficiency of immunity (EOI) against isogenic phages DMS3$_{acrIE3}$ (non-targeted) and

DMS3m$_{acrIF4}$ (CRISPR-targeted). Plaque forming units (PFUs) were quantified on ΔamrZ

or the complemented strain, then represented as a ratio of the number of PFUs

measured on WT PA14. EOI measurements are represented as the mean of 2 biological

replicates +/- SD. **b,c**. Growth curves of PA14 WT and Δ*amrZ* infected with either a high

(d,10^6 PFU) or low (e, 10^4 PFU) MOI of virulent DMS3m$_{acrIF4}$. Phage replication was

quantified as PFUs after 24 hours, and the efficiency of immunity expressed as a ratio of

the number of PFUs harvested from Δ*amrZ* to the number of PFUs obtained from WT.

OD600 and EOI measurements are represented as the mean of 3 biological replicates

+/- SD. **d, e**. Timecourse of the relative fluorescence levels of Csy1-sfCherry reporter

strains during surface-association (d) or planktonic growth (e). Fluorescence

measurements are represented as the mean of 3 biological replicates +/- SD. **f**.

Overexpression of the indicated gene in a WT *csy3::lacZ* background, with Miller units

measured at 8h and normalized to empty vector.

Figure 4. **Phage AmrZ homologs control CRISPR-Cas immunity. a.** Structure of an

AmrZ tetramer bound to 18bp of operator DNA[58] with DNA-contacting residues

highlighted in red. **b.** Protein alignment of seven mobile AmrZ homologs and the native

PA14 AmrZ homolog, with the ribbon-helix-helix DNA binding domain secondary

structure schematized and DNA-contacting residues indicated with red arrows and text.

c. Efficiency of immunity (EOI) against isogenic phages DMS3$_{acrIE3}$ (non-targeted) and

DMS3m$_{acrIF4}$ (CRISPR-targeted). Plaque forming units (PFUs) were quantified on ΔamrZ

or the strains complemented with bacterial or phage AmrZ homologs, and represented

as a ratio under the number of PFUs measured on WT PA14. EOI measurements as

represented as the mean of 3 biological replicates +/- SD. **d.** Measurement of the

normalized fluorescence levels of Csy1-sfCherry reporter strains transformed with

bacterial or phage AmrZ homologs after 10 hours of growth in liquid culture or 20 hours

of growth on solid media. Fluorescence measurements are represented as the mean of

3 biological replicates +/- SD.

Supplementary Table 1

Gene with Tn insertion	Transposon location	β-gal activity (% of unmutagenized)
kinB (1)	6448519	59%
kinB (2)	6447945	50%
kinB (3)	6447811	53%
kinB (4)	6449373	54%
kinB (5)	6449345	98%
purM	4618060	134%
lasR	4085810	135%
minD	1915545	N/D (growth defect)
gltB	5943637	108%
S-type pyocin	1196879	145%
pyoS3A	4404303	145%
tolA	4595505	158%
glycosyl transferase	5889967	81%
bacA	3490006	75%
Intergenic; zipA and smc +	3979746	N/D (growth defect)
gidB	6530337	88%
deaD	2373899	N/D (growth defect)
polA	6457284	N/D (growth defect)
pchH	798159	131%
crc	6275250	91%
putative plasmid stabilization protein	5347104	N/D (growth defect)
paraquat inducible protein	5532785	108%
pchH	797705	127%
yhiH/yhiI	6162134	108%
oxidoreductase FMN binding	4188602	N/D (growth defect)
putative membrane protein	1400718	118%
Intergenic; fstA and fstZ	5104077	111% (growth defect)
putative Zn-dependent oxidoreductase	2820446	115% (growth defect)
gnyL	3434168	87%
cytochrome c1 precursor	5126449	94%

Supplementary Table 1

All independent transposon insertions identified and mapped by visual screening with

increased or decreased *csy3::lacZ* β-galactosidase activity at 8 hours of growth. The

insertion location in the PA14 genome is shown, along with the measured level of β-

galactosidase enzyme. These measurements were not determined (N/D) for strains with

a growth defect.

Supplementary Table 2

Name	Accession	MGE type
PA14 AmrZ	ABJ12639.1	
Pseudomonas phage Noxifer	ARV77275.1	Lytic Myovirus
Pseudomonas phage phi3	YP_009276432.1	Integrated prophage
Pseudomonas phage PaBG	YP_008433620.1	Lytic Myovirus
Pseudomonas phage SM1	ALT58107.1	Siphoviridae (temperate)
Pseudomonas phage F10	YP_001293379.1	Siphoviridae (temperate)
Pseudomonas phage JBD68	ARM70500.1	Siphoviridae (temperate)
Pseudomonas sp. VLB120 plasmid pSTY	AGZ38169.1	Plasmid
Pseudomonas putida plasmid pKF715B	BAW27310.1	Plasmid
Pseudomonas veronii plasmid PVE_plasmid	SBW85251.1	Plasmid
Pseudomonas koreensis plasmid p3	AVX93364.1	Plasmid
Pseudomonas sp. Leaf58 plasmid pBASL58	AYG48213.1	Plasmid
Pseudomonas sp. XWY-1 plasmid	AUZ62175.1	Plasmid
Pseudomonas putida KF715C_pA870	BAW26592.1	Plasmid
Pseudomonas putida S12 plasmid pTTS12	AJA17154.1	Plasmid
Pseudomonas putida p12969-DIM	ALZ46341.1	Plasmid

Supplementary Table 2:

AmrZ homologs listed by the genome that encodes them, the accession number, and

the mobile genetic element type.

Supplementary Figure 1

a. A streak plate on X-gal plates, showing strains involved in this study and isolated

transposon (Tn) insertions. **b**. β-galactosidase measurements at the indicated time

points for the unmutagenized (*csy3::lacZ*) strain and three isolated *kinB* transposon

mutants (*kinB::Tn1-3*). **c**. Phage titration on lawns of the *kinB::Tn1* mutant transformed

with empty vector or *kinB.* **d.** Spot titration of phages JBD26 (CR2_sp17, sp20-targeted,

possessing *acrIF4*), JBD25 (CR1_sp1 targeted) on *kinB::Tn* mutants and ΔCRISPR-Cas.

Supplementary Figure 2:

a-c. Efficiency of immunity measurements for indicated mutants relative to WT. **a**. Δ*algB*

mutant complemented. **b.** Double knockouts show Δ*kinB* combined with *algB*, *algU*, or

algR. **c**. Indicated knockouts were combined with *csy3::lacZ*.

Supplementary Figure 3:

a-e. Growth curves of the indicated knockouts combined with *csy3::lacZ*. infected with virulent DMS3m*acrIF4* at multiplicity of infection (MOI, rainbow colors) increasing in 10-fold steps from 2×10^{-4} to 2×10^{2}. **f.** Phage replication was quantified as PFUs after 24 hours for the MOIs used in Figure 1 and 3, and the efficiency of immunity expressed as a ratio of the number of PFUs harvested from the mutants combined with *csy3::lacZ* under the number of PFUs obtained from WT *csy::lacZ*. OD600 and EOI measurements are represented as the mean of 3 biological replicates +/- SD.

Supplementary Figure 4

Supplementary Figure 4.

a. Log2 of Fragments Per Kilobase of transcript per Million mapped reads (FPKM) shown for each I-F *cas* gene in PA14 in the indicated growth condition. **b.** Log2 of protein levels for each of the I-F Cas proteins in PA14 in the indicated growth condition.

REFERENCES

1. Makarova, K. S. *et al.* An updated evolutionary classification of CRISPR-Cas systems. *Nat Rev Micro* **13,** 722–736 (2015).
2. Sapranauskas, R. *et al.* The Streptococcus thermophilus CRISPR/Cas system provides immunity in Escherichia coli. *Nucleic Acids Research* **39,** 9275–9282 (2011).
3. Vorontsova, D. *et al.* Foreign DNA acquisition by the I-F CRISPR-Cas system requires all components of the interference machinery. *Nucleic Acids Research* **43,** 10848–10860 (2015).
4. Mali, P. *et al.* RNA-guided human genome engineering via Cas9. *Science* **339,** 823–826 (2013).
5. Cong, L. *et al.* Multiplex Genome Engineering Using CRISPR/Cas Systems. *Science* **339,** 819–823 (2013).
6. Leon, L. M., Mendoza, S. D. & Bondy-Denomy, J. How bacteria control the CRISPR-Cas arsenal. *Curr Opin Microbiol* **42,** 87–95 (2018).
7. Patterson, A. G., Yevstigneyeva, M. S. & Fineran, P. C. Regulation of CRISPR-Cas adaptive immune systems. *Curr Opin Microbiol* **37,** 1–7 (2017).
8. Høyland-Kroghsbo, N. M. *et al.* Quorum sensing controls the Pseudomonas aeruginosa CRISPR-Cas adaptive immune system. *Proceedings of the National Academy of Sciences* **114,** 201617415–135 (2016).
9. Patterson, A. G. *et al.* Quorum Sensing Controls Adaptive Immunity through the

613   Regulation of Multiple CRISPR-Cas Systems. *Mol Cell* **64,** 1102–1108 (2016).

10.   Høyland-Kroghsbo, N. M., Muñoz, K. A. & Bassler, B. L. Temperature, by Controlling Growth Rate, Regulates CRISPR-Cas Activity in Pseudomonas aeruginosa. *mBio* **9,** 1709–12 (2018).

11.   Perez-Rodriguez, R. *et al.* Envelope stress is a trigger of CRISPR RNA-mediated DNA silencing in Escherichia coli. *Molecular Microbiology* **79,** 584–599 (2011).

12.   Serbanescu, M. A. *et al.* Role of the Streptococcus mutans CRISPR/Cas systems in immunity and cell physiology. (2014). doi:10.1128/JB.02333-14

13.   Shinkai, A. *et al.* Transcription activation mediated by a cyclic AMP receptor protein from Thermus thermophilus HB8. *J. Bacteriol.* **189,** 3891–3901 (2007).

14.   Patterson, A. G., Chang, J. T., Taylor, C. & Fineran, P. C. Regulation of the Type I-F CRISPR-Cas system by CRP-cAMP and GalM controls spacer acquisition and interference. *Nucleic Acids Research* **43,** gkv517–6048 (2015).

15.   Agari, Y. *et al.* Transcription profile of Thermus thermophilus CRISPR systems after phage infection. *Journal of Molecular Biology* **395,** 270–281 (2010).

16.   Young, J. C. *et al.* Phage-induced expression of CRISPR-associated proteins is revealed by shotgun proteomics in Streptococcus thermophilus. *PLoS ONE* **7,** e38077 (2012).

17.   Quax, T. E. F. *et al.* Massive activation of archaeal defense genes during viral infection. *Journal of Virology* **87,** 8419–8428 (2013).

18.   Fusco, S. *et al.* Transcriptome analysis of Sulfolobus solfataricus infected with two related fuselloviruses reveals novel insights into the regulation of CRISPR-Cas system. *Biochimie* (2015). doi:10.1016/j.biochi.2015.04.006

19.   Brouns, S. J. J. *et al.* Small CRISPR RNAs guide antiviral defense in prokaryotes. *Science* **321,** 960–964 (2008).

20.   Hochstrasser, M. L. *et al.* CasA mediates Cas3-catalyzed target degradation during CRISPR RNA-guided interference. *Proceedings of the National Academy of Sciences* **111,** 6618–6623 (2014).

21.   Westra, E. R. *et al.* CRISPR Immunity Relies on the Consecutive Binding and Degradation of Negatively Supercoiled Invader DNA by Cascade and Cas3. *Mol Cell* **46,** 595–605 (2012).

22.   Levy, A. *et al.* CRISPR adaptation biases explain preference for acquisition of foreign DNA. **520,** 505–510 (2015).

23.   Cady, K. C., Bondy-Denomy, J., Heussler, G. E., Davidson, A. R. & O'Toole, G. A. The CRISPR/Cas Adaptive Immune System of *Pseudomonas aeruginosa* Mediates Resistance to Naturally Occurring and Engineered Phages. 1–40 (2012).

24.   Haurwitz, R. E., Jinek, M., Wiedenheft, B., Zhou, K. & Doudna, J. A. Sequence- and Structure-Specific RNA Processing by a CRISPR Endonuclease. *Science* **329,** 1355–1358 (2010).

25.   Wiedenheft, B. *et al.* Structural basis for DNase activity of a conserved protein implicated in CRISPR-mediated genome defense. *Structure* **17,** 904–912 (2009).

26.   Wiedenheft, B. *et al.* RNA-guided complex from a bacterial immune system enhances target recognition through seed sequence interactions. *Proceedings of the National Academy of Sciences* **108,** 10092–10097 (2011).

27.   Rollins, M. F., Schuman, J. T., Paulus, K., Bukhari, H. S. T. & Wiedenheft, B. Mechanism of foreign DNA recognition by a CRISPR RNA-guided surveillance complex from Pseudomonas aeruginosa. *Nucleic Acids Research* (2015). doi:10.1093/nar/gkv094

28.   Rollins, M. F. *et al.* Cas1 and the Csy complex are opposing regulators of Cas2/3 nuclease activity. *Proceedings of the National Academy of Sciences* **23,**

201616395–E5121 (2017).
29. Chowdhury, S. *et al.* Structure Reveals Mechanisms of Viral Suppressors that
Intercept a CRISPR RNA-Guided Surveillance Complex. *Cell* **169,** 47–57.e11
(2017).
30. Zegans, M. E. *et al.* Interaction between bacteriophage DMS3 and host CRISPR
region inhibits group behaviors of Pseudomonas aeruginosa. *J. Bacteriol.* **191,**
210–219 (2009).
31. Cady, K. C. & O'Toole, G. A. Non-identity-mediated CRISPR-bacteriophage
interaction mediated via the Csy and Cas3 proteins. *J. Bacteriol.* **193,** 3433–3445
(2011).
32. van Belkum, A. *et al.* Phylogenetic Distribution of CRISPR-Cas Systems in
Antibiotic-Resistant Pseudomonas aeruginosa. *mBio* **6,** e01796–15 (2015).
33. Westra, E. R. *et al.* Parasite Exposure Drives Selective Evolution of Constitutive
versus Inducible Defense. *Curr. Biol.* **25,** 1043–1049 (2015).
34. van Houte, S. *et al.* The diversity-generating benefits of a prokaryotic adaptive
immune system. *Nature* **532,** 385–388 (2016).
35. Bondy-Denomy, J., Pawluk, A., Maxwell, K. L. & Davidson, A. R. Bacteriophage
genes that inactivate the CRISPR/Cas bacterial immune system. *Nature* **493,**
429–432 (2013).
36. Pawluk, A., Bondy-Denomy, J., Cheung, V. H. W., Maxwell, K. L. & Davidson, A.
R. A new group of phage anti-CRISPR genes inhibits the type I-E CRISPR-Cas
system of Pseudomonas aeruginosa. *mBio* **5,** e00896–e00896–14 (2014).
37. Bondy-Denomy, J. *et al.* Multiple mechanisms for CRISPR-Cas inhibition by anti-
CRISPR proteins. *Nature* **526,** 136–139 (2015).
38. Pawluk, A. *et al.* Inactivation of CRISPR-Cas systems by anti-CRISPR proteins in
diverse bacterial species. *Nature Microbiology* **1,** 1–6 (2016).
39. Cady, K. C. & O'Toole, G. Non-Identity-Mediated CRISPR-Bacteriophage
Interaction Mediated via the Csy and Cas3 Proteins. **193,** 3433–3445 (2011).
40. Cady, K. C., Bondy-Denomy, J., Heussler, G. E., Davidson, A. R. & O'Toole, G. A.
The CRISPR/Cas adaptive immune system of Pseudomonas aeruginosa
mediates resistance to naturally occurring and engineered phages. **194,** 5728–
5738 (2012).
41. Borges, A. L. *et al.* Bacteriophage Cooperation Suppresses CRISPR-Cas3 and
Cas9 Immunity. *Cell* **174,** 917–925.e10 (2018).
42. Damron, F. H. *et al.* Analysis of the Pseudomonas aeruginosa Regulon Controlled
by the Sensor Kinase KinB and Sigma Factor RpoN. **194,** 1317–1330 (2012).
43. Martin, D. W. *et al.* Mechanism of conversion to mucoidy in Pseudomonas
aeruginosa infecting cystic fibrosis patients. *Proc Natl Acad Sci USA* **90,** 8377–
8381 (1993).
44. Qiu, D., Eisinger, V. M., Rowen, D. W. & Yu, H. D. Regulated proteolysis controls
mucoid conversion in Pseudomonas aeruginosa. *Proc Natl Acad Sci USA* **104,**
8107–8112 (2007).
45. Chand, N. S. *et al.* The Sensor Kinase KinB Regulates Virulence in Acute
Pseudomonas aeruginosa Infection. **193,** 2989–2999 (2011).
46. Chand, N. S., Clatworthy, A. E. & Hung, D. T. The Two-Component Sensor KinB
Acts as a Phosphatase To Regulate Pseudomonas aeruginosa Virulence. **194,**
6537–6547 (2012).
47. Damron, F. H., Qiu, D. & Yu, H. D. The Pseudomonas aeruginosa Sensor Kinase
KinB Negatively Controls Alginate Production through AlgW-Dependent MucA
Proteolysis. **191,** 2285–2295 (2009).
48. Cezairliyan, B. O. & Sauer, R. T. Control of Pseudomonas aeruginosa AlgW

protease cleavage of MucA by peptide signals and MucB. *Molecular Microbiology*
72, 368–379 (2009).
49. Schurr, M. J., Yu, H., Martinez-Salazar, J. M., Boucher, J. C. & Deretic, V. Control
of AlgU, a member of the sigma E-like family of stress sigma factors, by the
negative regulators MucA and MucB and Pseudomonas aeruginosa conversion to
mucoidy in cystic fibrosis. *J. Bacteriol.* **178,** 4997–5004 (1996).
50. Tart, A. H., Blanks, M. J. & Wozniak, D. J. The AlgT-dependent transcriptional
regulator AmrZ (AlgZ) inhibits flagellum biosynthesis in mucoid, nonmotile
Pseudomonas aeruginosa cystic fibrosis isolates. *J. Bacteriol.* **188,** 6483–6489
(2006).
51. Baynham, P. J. & Wozniak, D. J. Identification and characterization of AlgZ, an
AlgT-dependent DNA-binding protein required for Pseudomonas aeruginosa algD
transcription. *Molecular Microbiology* **22,** 97–108 (1996).
52. Wozniak, D. J. & Ohman, D. E. Transcriptional analysis of the Pseudomonas
aeruginosa genes algR, algB, and algD reveals a hierarchy of alginate gene
expression which is modulated by algT. *J. Bacteriol.* **176,** 6007–6014 (1994).
53. Landsberger, M. *et al.* Anti-CRISPR Phages Cooperate to Overcome CRISPR-
Cas Immunity. *Cell* **174,** 908–916.e12 (2018).
54. Kong, W. *et al.* ChIP-seq reveals the global regulator AlgR mediating cyclic di-
GMP synthesis in Pseudomonas aeruginosa. *Nucleic Acids Research* **43,** 8268–
8282 (2015).
55. Lizewski, S. E. *et al.* Identification of AlgR-regulated genes in Pseudomonas
aeruginosa by use of microarray analysis. *J. Bacteriol.* **186,** 5672–5684 (2004).
56. Schulz, S. *et al.* Elucidation of sigma factor-associated networks in Pseudomonas
aeruginosa reveals a modular architecture with limited and function-specific
crosstalk. *PLoS Pathog* **11,** e1004744 (2015).
57. Wozniak, D. J. *et al.* Alginate is not a significant component of the extracellular
polysaccharide matrix of PA14 and PAO1 Pseudomonas aeruginosa biofilms.
Proc Natl Acad Sci USA **100,** 7907–7912 (2003).
58. Pryor, E. E. *et al.* The transcription factor AmrZ utilizes multiple DNA binding
modes to recognize activator and repressor sequences of Pseudomonas
aeruginosa virulence genes. *PLoS Pathog* **8,** e1002648 (2012).
59. Pratama, A. A. & van Elsas, J. D. A novel inducible prophage from the
mycosphere inhabitant Paraburkholderia terrae BS437. *Sci Rep* **7,** 9156 (2017).
60. Boucher, J. C., Yu, H., Mudd, M. H. & Deretic, V. Mucoid Pseudomonas
aeruginosa in cystic fibrosis: characterization of muc mutations in clinical isolates
and analysis of clearance in a mouse model of respiratory infection. *Infection and*
Immunity **65,** 3838–3846 (1997).
61. Martin, D. W., Holloway, B. W. & Deretic, V. Characterization of a locus
determining the mucoid status of Pseudomonas aeruginosa: AlgU shows
sequence similarities with a Bacillus sigma factor. *J. Bacteriol.* **175,** 1153–1164
(1993).
62. Damron, F. H., Qiu, D. & Yu, H. D. The Pseudomonas aeruginosa sensor kinase
KinB negatively controls alginate production through AlgW-dependent MucA
proteolysis. *J. Bacteriol.* **191,** 2285–2295 (2009).
63. Jones, A. K. *et al.* Activation of the Pseudomonas aeruginosa AlgU regulon
through mucA mutation inhibits cyclic AMP/Vfr signaling. **192,** 5709–5717 (2010).
64. Heilmann, S., Sneppen, K. & Krishna, S. Sustainability of virulence in a phage-
bacterial ecosystem. *Journal of Virology* **84,** 3016–3022 (2010).
65. Dötsch, A. *et al.* The Pseudomonas aeruginosaTranscriptional Landscape Is
Shaped by Environmental Heterogeneity and Genetic Variation. *mBio* **6,** e00749–

15–10 (2015).

66.   Erdmann, J., Preusse, M., Khaledi, A., Pich, A. & Häussler, S. Environment-driven
changes of mRNA and protein levels in Pseudomonas aeruginosa. *Environmental*
Microbiology **20,** 3952–3963 (2018).
770