High-throughput multiplexed tandem repeat genotyping using targeted long-read sequencing

Devika Ganesamoorthy[1#], Mengjia Yan[1], Valentine Murigneux[1], Chenxi Zhou[1], Minh Duc Cao[1], Tania PS Duarte[1] and Lachlan JM Coin[1#].

[1]Institute for Molecular Bioscience, The University of Queensland, Brisbane, Australia

[#] Correspondence to d.ganesamoorthy@imb.uq.edu.au or l.coin@imb.uq.edu.au

Devika Ganesamoorthy	d.ganesamoorthy@imb.uq.edu.au
Mengjia Yan	mengjia.y2@gmail.com
Valentine Murigneux	v.murigneux@imb.uq.edu.au
Chenxi Zhou	chenxi.zhou@imb.uq.edu.au
Minh Duc Cao	minhduc.cao@gmail.com
Tania PS Duarte	t.duarte@imb.uq.edu.au
Lachlan JM Coin	l.coin@imb.uq.edu.au

ABSTRACT

Background:

Tandem repeats (TRs) are highly prone to variation in copy numbers due to their repetitive and unstable nature, which makes them a major source of genomic variation between individuals. However, TRs are not been explored in the context of genomic variation, due to the limitations of existing tools, which are either low-throughput or restricted to a small subset of TRs. Here, we report a targeted sequencing approach combined with long-read sequencing technologies to overcome these limitations. Our approach described here presents an accurate and cost-effective method to explore TRs at the population level.

Results:

We report targeted enrichment of 142 TR loci using sequence capture combined with Nanopore and PacBio sequencing. We achieved 50-60% on-target sequence reads and an average of 100X sequence coverage per sample with Nanopore sequencing using 200ng of input DNA. Genotyping estimates from targeted long-read sequencing were equivalent to the accuracy of gold standard PCR sizing analysis with correlation greater than 0.98 for both Nanopore and PacBio sequencing. Genotyping estimates from targeted long-read sequencing were determined using two different methods (VNTRTyper and Tandem-genotypes) and results were comparable. Furthermore, genotyping estimates from targeted long-read sequencing were highly correlated with genotyping estimates from whole genome long-read sequencing.

Conclusions:

40   We demonstrate a targeted long-read sequencing approach that enables simultaneous analysis

41   of hundreds of TRs and accuracy is comparable to PCR sizing analysis. Our approach is

42   feasible to scale for more targets and more samples facilitating large-scale analysis of TRs.

44   **KEYWORDS**

45   Tandem repeats, targeted sequencing, long-read sequencing

BACKGROUND

Repeated sequences occur in multiple copies throughout the genome, and they make up almost half of the human genome [1]. Repeat sequences can be divided into two categories, namely interspersed repeats and tandem repeats (TRs). Interspersed repeats are scattered throughout the genome and are remnants of transposons [2]. TRs consists of repeat units that are located adjacent to each other (*i.e.* in tandem). There are almost 1 million TRs in the human genome covering 10.6% of the entire genome [3]. TRs can be further divided into two types based on the length of the repeat unit; repeats with less than six base pair repeat units are classified as microsatellites or short tandem repeats (STRs) and those with more than 6 base pair repeat units are known as minisatellites [4]. TRs that have variable copy numbers are known as variable number tandem repeats (VNTRs).

VNTRs are prone to high rates of copy number variation and mutation due to the repetitive unstable nature, which makes them a major source of genomic variation between individuals. Variation in VNTRs may explain some of the phenotypic variation observed in complex diseases as it is poorly tagged by single nucleotide variation [5, 6]. Recent studies have shown that 10% to 20% of coding and regulatory regions contain TRs and suggested that variations in TRs could have phenotypic effect [7]. Although TRs represent a highly variable fraction of the genome, analysis of TRs so far are limited to known pathogenic regions, mainly STRs due to the limitations in analysis techniques.

Traditionally, TR analysis has been carried out via restriction fragment length polymorphism

(RFLP) analysis [8] or PCR amplification of the target loci followed by fragment length

analysis [9]. These techniques are only applicable to a specific target region and not scalable

to high-throughput analysis, which limits the possibility of genome-wide TR analysis. In the

recent decade, significant progress has been made in utilising high throughput short-read

sequencing data for genotyping STRs [10]. Our group and others have also demonstrated

targeted sequencing approaches using short-read sequencing for TR analysis [11, 12]. Several

computational tools have been developed to improve the accuracy of TR genotyping from

short-read sequencing data with varying performance [13-17]. Yet, most of these tools have

focused mainly on the analysis of STRs and analysis of longer TRs remains a hurdle for these

approaches. We reported GtTR in Ganesamoorthy *et. al.* (2018) [12], which utilizes short-read

sequencing data to genotype longer TRs. GtTR reports absolute copy number of the TRs, but

it does not report the exact genotype of two alleles due to the use of short-read sequencing data.

Sequencing reads that span the entire repeat region are informative to accurately genotype TRs

[11], and therefore are ideal for genome-wide TR analysis. Long-read sequencing technologies

have the potential to span all TRs in human genome, including long TRs. There have been few

reports on the use of long-read sequencing for the analysis of specific TRs implicated in

diseases [18-20]. Genotyping tools utilizing long-read sequencing data, such as

Nanosatellite[19], RepeatHMM [21] and Tandem-genotypes [22] have been reported in the

recent years with varying performance across different length of repeat units and repeat length.

We reported VNTRTyper in Ganesamoorthy *et. al.* (2018) [12] to genotype TRs from long-read

whole genome sequencing data. Despite the availability of genotyping tools, long-read

sequencing is not widely used for TR analysis, due to the high costs associated with whole

genome long-read sequencing.

In this study, we demonstrate a targeted sequence capture approach combined with long-read

sequencing for large-scale analysis of TRs. We used a hybridization based target enrichment

system to enrich regions of interest followed with long-read sequencing [23]. We have

previously demonstrated that targeted sequence capture of repetitive TR sequences are feasible

using short-read sequencing technologies [12]. In this study, we demonstrate the targeted

sequence capture of repetitive TRs using both Oxford Nanopore and PacBio long-read

sequencing technologies. We demonstrate the accuracy of genotype estimates from targeted

long-read sequencing by comparison with gold-standard PCR sizing analysis. Our targeted

long-read sequencing approach presented in this study provides an accurate and cost effective

approach for large-scale analysis of TRs.

RESULTS

We demonstrate a targeted sequence capture approach combined with long-read sequencing to

genotype hundreds of TRs. We evaluated this approach using both Nanopore and PacBio long-

read sequencing technologies.

Targeted Capture Sequencing of Tandem repeats

We performed sequence enrichment of targeted TRs followed by Nanopore or PacBio long-

read sequencing as described in Methods. Table 1 summarises the metrics for targeted

sequencing on both Nanopore and PacBio sequencing technologies. Sequencing yield per

sample was lower for Nanopore compared to PacBio due to multiplexing of samples. Nanopore

multiplexing (See Methods) group 1 samples had similar yield between samples, however,

Nanopore multiplexing group 2 samples had varying yield per sample. Despite the differences

in sequencing yield, on target capture rate was approximately 50% for Nanopore sequencing,

however only 8% of reads were on target for PacBio sequence capture. Hence, the fold

enrichment of targets were lower for PacBio sequence capture compared to Nanopore capture.

It is worth noting that, the sequencing capture protocols used were different between Nanopore

and PacBio sequencing as described in Methods.

Figure 1a shows the read length distribution observed in targeted capture sequencing data using

Nanopore and PacBio sequencing technologies. The median read length was higher for PacBio

sequence capture protocol compared to Nanopore capture sequencing protocol. However, it is

worth noting that lengths of targeted TR range from 112bp to 25236bp, with a median length

of 667bp (Additional File 1: Figure 1). The read length in this study was sufficient to analyse

majority of targeted TRs. Figure 1b shows the sequence coverage (*i.e.* average per base

coverage in the region) distribution in targeted TR regions for Nanopore and PacBio

sequencing. On average 80X and 100X sequence coverage was achieved for PacBio and

Nanopore sequencing respectively. Furthermore, 8% and 14% of targets had less than 1X

coverage in PacBio and Nanopore capture sequencing respectively and majority of these targets

had less than 25% GC content (Additional File 1: Figure 2).

Genotyping of Tandem repeats using targeted long-read sequencing

Genotype estimates from targeted long-read sequencing datasets were estimated using our tool

VNTRTyper [12] with the improvements described in Methods. We also applied Tandem-

genotypes [22] for genotyping TRs from long-read sequencing data. We used a minimum of

two reads as read threshold to determine the repeat number for each alleles.

Genotype estimates by VNTRTyper were compared to the PCR sizing results on 8 targets which

had sufficient coverage (Table 2). Genotype estimates from both Nanopore and PacBio targeted

sequencing correlated well with PCR results and the correlation values were greater than 0.980

for all samples. Genotype estimates by Tandem-genotypes were also compared to the PCR

sizing results (Additional File 1: Table 1) and correlation values were greater than 0.984 for all

samples, however fewer targets had sufficient data to compare with PCR sizing results.

The three samples (NA12877, NA12878 and NA12879) which were sequenced using both

Nanopore and PacBio sequencing technologies were used to assess the performance of different

sequencing technologies. Both VNTRTyper and Tandem-genotypes correlated well with PCR

sizing results with correlation values greater than 0.984 for both sequencing technologies

(Table 2 and Additional File 1: Table 1). However, Nanopore targeted sequencing had

marginally better correlation than PacBio targeted sequencing (Additional File 1: Figure 3 and

4).

We were able to determine the genotype on 60% of the targets (range 48% to 75%) with

Nanopore targeted sequencing and on 54% of the targets (range 54% to 55%) with PacBio

targeted sequencing using VNTRTyper. On the other hand, Tandem-genotypes was only able

to genotype 51% of the targets (range 34% to 70%) with Nanopore sequencing and 45% of the

targets (range 44% to 46%) with PacBio sequencing. Both genotyping approaches failed to

genotype targets with low GC sequence content (< 25% GC content) and targets which are

greater than 2Kb in length, which accounts for 20% of the targets. Targets with low GC

sequence content (< 25% GC content) didn't have sufficient sequence coverage for analysis

due to inefficient sequence enrichment in these regions (Additional File 1: Figure 2). Targets

which are greater than 2Kb in length didn't have sufficient spanning reads for genotyping

analysis (See Methods and Additional File 1: Figure 1).

Tandem-genotypes had lower genotyping rate compared to VNTRTyper and this is likely due

to the difference in algorithms. Targets which were only genotyped by VNTRTyper, had less

than 5 reads supporting the alleles, which indicates that the coverage in these targets were not

sufficient for Tandem-genotypes analysis. Despite these differences, genotype estimates by

VNTRtyper and Tandem-genotypes correlate well and the correlation values range from 0.950

to 0.995 for Nanopore (except 0.904 for sample NA12881, which had low sequence coverage)

and 0.982 to 0.999 for PacBio targeted sequencing

Genotype estimates from VNTRTyper and Tandem-genotypes for all 142 targets from

Nanopore capture sequencing samples and PacBio capture sequencing samples are provided in

Additional File 2: Table 1 and Table 2 respectively.

Genotyping of Tandem repeats using long-read whole genome sequencing

To investigate the accuracy of genotype estimates of TRs from targeted sequence capture

compared to WGS, we performed genotyping analysis on targeted regions using VNTRTyper

and Tandem-genotypes on whole genome long-read sequencing data. We downloaded whole

genome long-read Nanopore and PacBio sequencing data on CEPH Pedigree 1463 NA12878

sample. We have previously reported genotyping estimates by VNTRTyper on PacBio

NA12878 WGS data [12]. Here we use the genotype estimates by VNTRTyper on PacBio

NA12878 WGS data to compare genotype estimates by Tandem-genotype and the results of

PacBio targeted sequencing analysis.

We compared the accuracy of genotype estimates from WGS data with PCR sizing analysis.

Genotype estimates by VNTRTyper and Tandem-genotypes on WGS data were compared with

PCR sizing results on 10 targets (Table 3). VNTRTyper and Tandem-genotypes had comparable

correlation with PCR sizing analysis for both Nanopore and PacBio WGS.

We also compared genotype estimates from targeted capture sequencing data and WGS data

with PCR sizing analysis on 7 targets which had sufficient coverage (Figure 2 and Table 3).

Nanopore WGS data correlated precisely with 13/14 genotype estimates on Nanopore capture

sequencing and 12/14 genotype estimates correlated accurately between PacBio WGS and

capture sequencing using VNTRTyper. Genotype estimates using Tandem-genotypes on

Nanopore WGS data correlated precisely with 9/10 genotype estimates on Nanopore capture

sequencing and 10/12 genotype estimates correlated accurately between PacBio WGS and

capture sequencing. Genotype estimates for all 142 targets from Nanopore and PacBio WGS

data determined using VNTRTyper and Tandem-genotypes are provided in Additional File 2:

Table 3.

We compared the genotype estimates on all targets between WGS data and targeted capture

sequencing data for both Nanopore and PacBio sequencing. Genotype estimates by

VNTRTyper correlated well between WGS data and targeted capture sequencing data with a

correlation of 0.9782 and 0.9378 for Nanopore and PacBio sequencing respectively (Figure 3).

Genotype estimates by Tandem-genotypes correlated well for PacBio WGS data and targeted

capture sequencing with a correlation value of 0.9954, however had low correlation (r^2=0.8618)

for Nanopore WGS and targeted capture sequencing data (Additional File 1: Figure 5).

Variation in Tandem repeats

To assess the extent of variation in repeat numbers between individuals, we compared the

genotype estimates to the reported reference (hg19) repeat number. Genotype estimates

determined by VNTRTyper on Nanopore capture sequencing on 7 members of CEPH pedigree

1463 were used to assess the variation. On average, 49% (range 40% - 55%) of the targets were

similar to the reported reference genotype and 51% (range 45% - 60%) of the targets differed

from the reference (Figure 4). Furthermore, more deletions (28%) than duplications (23%)

were observed in repeat number.

DISCUSSION

In this study, we presented a targeted sequencing approach combined with long-read

sequencing technology to genotype TRs. To our knowledge, this is the first report on

genotyping analysis of hundreds of TRs using targeted long-read sequencing approach.

Sequencing reads that span the entire repeat region and flanking region are often useful in

providing an accurate estimation of the repeat genotype. Long-read sequencing technologies

have the ability to generate reads which can span the entire repeat region and flanking regions.

However, whole genome long-read sequencing analysis is still expensive for large-scale

population analysis; hence, we developed a targeted long-read sequencing approach for TR

analysis.

We showed that 1) target enrichment of repetitive sequences followed by long-read sequencing

is feasible, 2) both Nanopore and PacBio long-read sequencing technologies can be utilised to

genotype TRs and 3) genotype predictions on targeted TR sequencing are comparable to the

accuracy of PCR sizing analysis of repeats. We modified target enrichment protocol between

Nanopore and PacBio sequencing technologies to minimize sequencing cost and to achieve

required input DNA requirements (See Methods). Hence, there were differences in sequencing

output between Nanopore and PacBio targeted sequencing experiments. On target capture was

less than 10% for PacBio sequencing, but greater than 50% for Nanopore sequencing. However,

median read length was greater than 1.1Kb for PacBio but less than 770bp for Nanopore

sequencing. These differences are likely due to the difference in probe hybridization conditions

and reagents between Nanopore and PacBio capture experiments. Despite these biases,

estimated genotypes from both Nanopore and PacBio capture sequencing were highly

concordant.

On average, 60% of the targets could be genotyped by targeted long-read sequencing analysis.

However, this improved to 75% with increased sequence output, indicating more sequencing

could improve genotyping rate. We also observed targets greater than 2Kb in length could not

be genotyped due to the lack of spanning reads for genotyping analysis. This is primarily due

to the limitation in sequence read length observed from the capture process. Streptavidin beads

used during the capture process has limitations on the size of the fragments it can bind to, which

limits the fragment length attainable with the capture protocol. Furthermore, we noticed targets

with low GC sequence content (< 25% GC content) didn't have sufficient sequence coverage

in both Nanopore and PacBio sequencing experiments. We have previously performed short-

read target capture on these regions [12] and did not observe any significant bias in sequence

coverage due to GC sequence content. Furthermore, both Nanopore and PacBio WGS data

didn't have any bias in sequence coverage in low GC regions. Hence, the lack of sequence

coverage in low GC region for targeted sequencing is likely due to the capture protocol used

in this study.

We used VNTRTyper, an in-house genotyping tool described in Ganesamoorthy et. al. (2018)

[12] to determine the repeat number of TRs from long-read sequencing technologies. For

comparison, we used Tandem-genotypes [22], recently reported genotyping tool for the

detection of TR expansions from long-read sequencing. Both genotyping methods were

comparable to PCR sizing analysis and genotyping estimates were comparable between the

approaches. However, Tandem-genotypes genotyped less targets than VNTRTyper. The

differences is likely due to the different algorithms used between the methods. Both

VNTRTyper and Tandem-genotypes uses reads spanning the repeat region. However, for

Tandem-genotypes the flanking length used for analysis is depended on the length of the repeat

unit, with a maximum of 100bp on both sides of the repeat unit. On the other hand, VNTRTyper

uses a default 30bp flanking length for analysis, but it is feasible to change the flanking length.

Due to the longer flank length requirement, Tandem-genotypes could have possibly failed to

genotype more targets compared to VNTRTyper.

Variations in TRs are a major source of genomic variation between individuals. TRs targeted

in this study were initially selected due to the variation observed between case and control

samples for obesity analysis [12] and these TRs are variable in the population. We show that

approximately 50% of the targeted TRs differ from reported reference copy number. However,

the major limitation in this analysis is that the sample size is small and the individuals are

related, which introduces a bias in the analysis. Nevertheless, these findings indicate the

possibility of variation in TR copy number between individuals and further large-scale studies

are required to ascertain the extent of variation.

We demonstrated that the accuracy of genotype estimates between WGS and targeted capture

sequencing were comparable to the accuracy of PCR sizing analysis. However, targeted capture

enrichment protocols used in this study have amplification steps, which can introduce errors in

TR analysis. This could possibly explain the differences in genotype estimates observed

between WGS and targeted capture sequencing for some targets.

An amplification free targeted analysis with long-read sequencing is an ideal option for

accurate genotyping TRs. Targeted cleavage with Cas9 enzyme followed by Nanopore

sequencing [24] or PacBio sequencing [25] has been recently reported as alternative option for

enrichment of regions of interest. This method does not have any amplifications and can be

adapted for multiple targets in a single assay. However, currently the DNA input requirements

are high and sequencing output are low, which currently restricts wide use of this technique for

large-scale analysis.

CONCLUSIONS

In summary, we present a targeted approach combined with long-read sequencing to enable

cost-effective and accurate approach to genotype TRs using long-read sequencing. Using this

method, we have successfully demonstrated the feasibility of targeted capture sequencing of

repetitive sequences and genotyping TRs using both Nanopore and PacBio long-read

sequencing technologies. Our targeted long-read sequencing approach would provide a cost-

effective tool for large-scale population analysis of tandem repeats.

METHODS

Samples for sequencing:

DNA samples of CEPH/UTAH pedigree 1463 were purchased from Coriell Institute for

Medical Research (USA). Seven family members from the pedigree used for sequencing

analysis were NA12877, NA12878, NA12879, NA12881, NA12882, NA12889 and NA12890.

Selection of Tandem Repeats and Probe design

The selection of TRs and design of probes were described in Ganesamoorthy *et. al*. (2018) [12].

Briefly, 142 TRs were selected; they range from 112bp to 25236bp in length in the reference

human genome (hg19) and the number of repeat units range from 2 to 2300 repeats. Agilent

SureSelect DNA design was used to design target probes to capture the targeted regions

(including 100bp flanking regions) and regions flanking the TRs (at least 1000bp).

Nanopore targeted sequencing of TRs:

All 7 family members from the CEPH pedigree 1463 were used for Nanopore targeted

sequencing analysis (NA12877, NA12878, NA12879, NA12881, NA12882, NA12889 and

NA1289). Target sequence capture for Nanopore sequencing was performed using Agilent

SureSelect XT HS Target Enrichment System according to the manufacturer's instructions with

slight modifications. Briefly, 200ng of DNA was fragmented to 3Kb using Covaris Blue

miniTUBE. Greater than 90% of the targeted TRs are less than 3Kb and SureSelect capture

protocol works effectively on fragments less than 4Kb in length; therefore, DNA products were

sheared to 3Kb. Fragmented DNA was end repaired, adapter ligated and amplified prior to

target capture. Extension time for pre-capture amplification was increased to 4 minutes to allow

for the amplification of long fragments and 14 cycle amplification was used. Purified pre-

capture PCR products were hybridized to the designed capture probes for 2 hours. Streptavidin

beads were used to pull down the DNA fragments bound to the probes. Finally, captured DNA

was amplified with long extension time (4 minutes) using Illumina Index adapters provided in

the enrichment kit. Post capture PCR products were purified using 0.8X - 1X Agencourt

AMPure XP beads.

Nanopore sequencing library preparation was performed using 1D Native barcoding genomic

DNA (with EXP-NBD103 and SQK-LSK108) protocol according to the manufacturer's

instructions with minor modifications. Briefly, 100ng – 200ng of post capture PCR products

were end repaired and incubated at 20°C for 15mins and 65°C for 15mins. End repaired products

were ligated with unique native barcodes. Purification steps after end repair and barcode

ligation were avoided to minimize the loss of DNA. Barcoded samples were pooled in

equimolar concentrations prior to adapter ligation. Adapter ligated samples were purified using

0.4X Agencourt AMPure XP beads. Samples were split into 2 sequencing groups; NA12877,

NA12878, NA12879 and NA12890 – group 1; NA12881, NA12882 and NA12889 – group 2.

Sequencing was performed using Oxford Nanopore Technologies MinION sequencer using

R9.5 flow cell. Both groups were sequenced for 48 hours. Nanopore sequencing data were base

called using Albacore (version 2.2.7) and reads were demultiplexed using Albacore (version

2.2.7) based on the barcode sequences.

PacBio targeted sequencing of TRs:

Three family members from the CEPH pedigree 1463 were selected for PacBio targeted

sequencing analysis (NA12877, NA12878 and NA12879). Target capture process for PacBio

sequencing was similar to Nanopore target capture process as described above. However, to

facilitate high input requirement for PacBio sequencing a high input target capture protocol

was used. Agilent SureSelect XT Target Enrichment System was used according to the

manufacturer's instructions with slight modifications. Briefly, 3000ng of DNA was used as

input and fragmented to 3Kb using Covaris Blue miniTUBE. Fragmented DNA was end

repaired, adapter ligated and amplified prior to target capture. Purified pre-capture PCR

products were hybridized to the designed capture probes for 24 hours. Streptavidin beads were

used to pull down the DNA fragments bound to the probes. Captured DNA was amplified using

InPE 1.0 short and InPE 2.0 short primers provided by Agilent Technologies. Amplified

products (approximately 1000ng) were used as input for PacBio library preparation.

PacBio sequencing library preparation was performed using PacBio® Template Prep Kit

protocol according to the manufacturer's instructions. Briefly, amplified DNA fragments were

end repaired, blunt adapter ligated and SMRTbell templates were prepared. Purified SMRTbell

libraries were sequenced on PacBio RS sequencing system using one SMRTcell per sample.

Sequencing data were processed using SMRTportal and sub reads were extracted for

downstream analysis.

Public data used in the study:

Nanopore WGS data on CEPH Pedigree 1463 sample NA12878 were obtained from Nanopore

WGS consortium [26]. PacBio WGS data on NA12878 sample were downloaded from SRA

with accession numbers SRX627421 and SRX638310 [27]

VNTRTyper:

Both Nanopore and PacBio sequencing data were mapped to hg19 reference genome using

Minimap2 (version 2.13) [28]. For Nanopore sequencing reads '-ax map-ont' and for PacBio

sequencing reads '-ax map-pb' parameters were used. VNTRTyper, our in-house tool described

in Ganesamoorthy *et. al.* (2018) [12] was used to genotype TRs from long-read sequencing

data. Briefly, VNTRTyper takes advantage of the long-read sequencing to identify the number

of repeat units in the TR regions. Firstly, the tool identifies reads that span the repeat region

and applies Hidden Markov Models (HMM) to align the repetitive portion of each read to the

repeat unit. Then it estimates the multiplicity of the repeat units in a read using a profile HMM.

Recently, we further improved the accuracy of genotyping estimates by clustering the copy

number counts from reads to identify the likely genotypes per target. A minimum threshold of

two supporting reads per genotype was used to assign genotypes. Furthermore, for

heterozygous alleles, both alleles should have at least 10% of reads supporting the allele, if not

allele with less than 10% of reads was excluded during the analysis.

The updated version of VNTRTyper can be accessed from https://github.com/mdcao/japsa

release 1.9-3c and can be deployed using script name jsa.tr.longreads. Details of VNTRTyper

analysis are previously reported in Ganesamoorthy *et. al.* (2018) [12].

Tandem-genotypes:

We also used another independent method Tandem-genotypes to estimate genotypes from long-

read sequencing data. Tandem-genotypes was recently reported for analysis of TR genotypes

from long-read sequencing data [22] and it can be utilised for both Nanopore and PacBio

sequencing technologies.

Nanopore and PacBio sequencing data were mapped to the hg19 reference genome using LAST

v959 [29]. Calculation of repeat length per sequencing read was performed with Tandem-

genotypes as reported in Mitsuhashi et. al. (2019) [22]. Copy number changes in reads covering

the repeat's forward and reverse strands were merged and the two alleles with the highest

number of supporting reads for each VNTR were extracted. A minimum threshold of two

supporting reads per genotype was used to assign genotypes.

PCR analysis of VNTRs:

Ten targeted VNTR regions which are less than 1Kb in repetitive sequence were validated by

PCR sizing analysis in this study. These ten targets include various repeat unit length and repeat

sequence combinations to assess the accuracy of the genotypes determined from sequencing

data. Majority of these targets were tested in our previous study [12] and the results from the

previous PCR anlaysis were used for these regions. PCRs were performed using HotStar Taq

DNA Polymerase (Qiagen) and PCR conditions were optimized for each PCR target. PCR

products were purified and subjected to capillary electrophoresis on an ABI3500xL Genetic

Analyzer (Applied BioSystems). Fragment sizes were analyzed using GeneMapper 4.0

(Applied BioSystems).

Statistical analysis:

Linear regression analysis was used to determine correlation between genotype estimates. All

plots were generated using GraphPad Prism version 7.00 for Windows, GraphPad Software,

La Jolla California USA, www.graphpad.com

LIST OF ABBREVIATIONS

TR – Tandem Repeats

STR _Short Tandem Repeats

VNTR – Variable Number of Tandem Repeats

PCR – Polymerase Chain Reaction

HMM - Hidden Markov Model

WGS – Whole Genome Sequencing

DECLARATIONS

Ethics approval and consent to participate: Not applicable

Consent for publication: Not applicable

Availability of data and materials: The sequencing datasets generated during the study are available in NCBI SRA repository, project number PRJNA422490.

Competing interests: The authors declare that they have no competing interests.

Funding: This study is supported by the funding from Australian National Health and Medical Research Council (APP1052303).

Authors' contributions: DG and LC conceived the study. DG, MY and TD performed the experiments. DG, MY, VM, CZ, MDC and LC performed the sequencing and statistical analysis. DG and MY wrote the paper with input from the other authors. All authors read and approved the final manuscript.

Acknowledgements: Not Applicable.

REFERENCES:

1.   Lander ES, Linton LM, Birren B, Nusbaum C, Zody MC, Baldwin J, Devon K, Dewar K, Doyle M, FitzHugh W, et al: **Initial sequencing and analysis of the human genome.** *Nature* 2001, **409:**860-921.

2.   Jurka J, Kapitonov VV, Kohany O, Jurka MV: **Repetitive sequences in complex**

genomes: structure and evolution. *Annu Rev Genomics Hum Genet* 2007, **8:**241-259.

3.   Gelfand Y, Rodriguez A, Benson G: **TRDB--the Tandem Repeats Database.** *Nucleic*

Acids Res 2007, **35:**D80-87.

4.   Gemayel R, Cho J, Boeynaems S, Verstrepen KJ: **Beyond junk-variable tandem**

repeats as facilitators of rapid evolution of regulatory and coding sequences. *Genes*

2012, **3:**461-480.

5.   Armour JA: **Tandemly repeated DNA: why should anyone care?** *Mutat Res* 2006,

598:6-14.

6.   Hannan AJ: **TRPing up the genome: Tandem repeat polymorphisms as dynamic**

sources of genetic variability in health and disease. *Discov Med* 2010, **10:**314-321.

7.   Gemayel R, Vinces MD, Legendre M, Verstrepen KJ: **Variable tandem repeats**

accelerate evolution of coding and regulatory sequences. *Annual review of genetics*

2010, **44:**445-477.

8.   Bidwell JL, Bignon JD: **DNA-RFLP methods and interpretation scheme for HLA-**

DR and DQ typing. *Eur J Immunogenet* 1991, **18:**5-22.

9.   Tagliabracci A, Buscemi L, Sassaroli C, Paoli M, Rodriguez D: **Allele typing of short**

tandem repeats by capillary electrophoresis. *Int J Legal Med* 1999, **113:**26-32.

10.   Bahlo M, Bennett MF, Degorski P, Tankard RM, Delatycki MB, Lockhart PJ: **Recent**

advances in the detection of repeat expansions with short-read next-generation

sequencing. *F1000Res* 2018, **7**.

11.   Duitama J, Zablotskaya A, Gemayel R, Jansen A, Belet S, Vermeesch JR, Verstrepen

KJ, Froyen G: **Large-scale analysis of tandem repeat variability in the human**

genome. *Nucleic acids research* 2014, **42:**5728-5741.

12.   Ganesamoorthy D, Cao MD, Duarte T, Chen W, Coin L: **GtTR: Bayesian estimation of absolute tandem repeat copy number using sequence capture and high throughput sequencing.** *BMC Bioinformatics* 2018, **19:**267.

13.   Gymrek M, Golan D, Rosset S, Erlich Y: **lobSTR: A short tandem repeat profiler for personal genomes.** *Genome Res* 2012, **22:**1154-1162.

14.   Highnam G, Franck C, Martin A, Stephens C, Puthige A, Mittelman D: **Accurate human microsatellite genotypes from high-throughput resequencing data using informed error profiles.** *Nucleic Acids Res* 2013, **41:**e32.

15.   Cao MD, Tasker E, Willadsen K, Imelfort M, Vishwanathan S, Sureshkumar S, Balasubramanian S, Boden M: **Inferring short tandem repeat variation from paired-end short reads.** *Nucleic Acids Res* 2014, **42:**e16.

16.   Willems T, Zielinski D, Yuan J, Gordon A, Gymrek M, Erlich Y: **Genome-wide profiling of heritable and de novo STR variations.** *Nat Methods* 2017, **14:**590-592.

17.   Dashnow H, Lek M, Phipson B, Halman A, Sadedin S, Lonsdale A, Davis M, Lamont P, Clayton JS, Laing NG, et al: **STRetch: detecting and discovering pathogenic short tandem repeat expansions.** *Genome Biol* 2018, **19:**121.

18.   Schule B, McFarland KN, Lee K, Tsai YC, Nguyen KD, Sun C, Liu M, Byrne C, Gopi R, Huang N, et al: **Parkinson's disease associated with pure ATXN10 repeat expansion.** *NPJ Parkinsons Dis* 2017, **3:**27.

19.   De Roeck A, De Coster W, Bossaerts L, Cacace R, De Pooter T, Van Dongen J, D'Hert S, De Rijk P, Strazisar M, Van Broeckhoven C, Sleegers K: **Accurate characterization**

of expanded tandem repeat length and sequence through whole genome long-read sequencing on PromethION. *bioRxiv* 2018**:**439026.

20. Ebbert MTW, Farrugia SL, Sens JP, Jansen-West K, Gendron TF, Prudencio M, McLaughlin IJ, Bowman B, Seetin M, DeJesus-Hernandez M, et al: **Long-read sequencing across the C9orf72 'GGGGCC' repeat expansion: implications for clinical use and genetic discovery efforts in human disease.** *Mol Neurodegener* 2018, **13:**46.

21. Liu Q, Zhang P, Wang D, Gu W, Wang K: **Interrogating the "unsequenceable" genomic trinucleotide repeat disorders by long-read sequencing.** *Genome Med* 2017, **9:**65.

22. Mitsuhashi S, Frith MC, Mizuguchi T, Miyatake S, Toyota T, Adachi H, Oma Y, Kino Y, Mitsuhashi H, Matsumoto N: **Tandem-genotypes: robust detection of tandem repeat expansions from long DNA reads.** *Genome Biol* 2019, **20:**58.

23. Karamitros T, Magiorkinis G: **Multiplexed Targeted Sequencing for Oxford Nanopore MinION: A Detailed Library Preparation Procedure.** *Methods Mol Biol* 2018, **1712:**43-51.

24. Gilpatrick T, Lee I, Graham JE, Raimondeau E, Bowen R, Heron A, Sedlazeck FJ, Timp W: **Targeted Nanopore Sequencing with Cas9 for studies of methylation, structural variants and mutations.** *bioRxiv* 2019**:**604173.

25. Hafford-Tear NJ, Tsai YC, Sadan AN, Sanchez-Pintado B, Zarouchlioti C, Maher GJ, Liskova P, Tuft SJ, Hardcastle AJ, Clark TA, Davidson AE: **CRISPR/Cas9-targeted enrichment and long-read sequencing of the Fuchs endothelial corneal dystrophy-**

associated TCF4 triplet repeat. *Genet Med* 2019.

26.   Jain M, Koren S, Miga KH, Quick J, Rand AC, Sasani TA, Tyson JR, Beggs AD,

Dilthey AT, Fiddes IT, et al: **Nanopore sequencing and assembly of a human genome**

with ultra-long reads. *Nat Biotechnol* 2018, **36:**338-345.

27.   Pendleton M, Sebra R, Pang AW, Ummat A, Franzen O, Rausch T, Stutz AM, Stedman

W, Anantharaman T, Hastie A, et al: **Assembly and diploid architecture of an**

individual human genome via single-molecule technologies. *Nat Methods* 2015,

12:780-786.

28.   Li H: **Minimap2: pairwise alignment for nucleotide sequences.** *Bioinformatics* 2018,

34:3094-3100.

29.   Kielbasa SM, Wan R, Sato K, Horton P, Frith MC: **Adaptive seeds tame genomic**

sequence comparison. *Genome Res* 2011, **21:**487-493.

Figure 1: Targeted sequence capture (a) read length and (b) sequence coverage distribution for

Nanopore and PacBio sequencing.

Figure 2: Comparison of genotype estimates between WGS and Target capture sequencing for NA12878 sample using (a) VNTRTyper and (b) Tandem-genotypes. Red line indicates PCR sizing results. Targets with no genotype estimates are shown as a gap for the corresponding column.

Figure 3: Correlation of VNTRTyper genotype estimates between WGS and Targeted sequencing on (a) Nanopore and (b) PacBio sequencing data for NA12878 sample.

555 **Figure 4:** Percentage difference between reported repeat number in reference genome (hg19)

556 and estimated repeat number based on genotype estimates using VNTRTyper on Nanopore

557 targeted sequencing.

Table 1: Targeted Sequencing metrics for Nanopore and PacBio Capture sequencing of tandem repeats

Sample	Number of reads	Number of Aligned bases	On-bait bases (%)^	Off-bait bases (%)^	Fold enrichment^	Median Read length (bp)	Max Read length (bp)
NA12877_Nanopore*	207976	63384429	59%	41%	3497	568	11207
NA12878_Nanopore*	189050	70210003	62%	38%	3566	646	6781
NA12879_Nanopore*	209341	72340656	56%	44%	3284	614	24657
NA12890_Nanopore*	206077	79329086	52%	48%	2955	633	8040
NA12881_Nanopore#	44363	21288912	48%	52%	2739	702	5720
NA12882_Nanopore#	106259	52366827	49%	51%	2823	718	7383
NA12889_Nanopore#	1021376	539165551	45%	55%	2573	776	10127
NA12877_PacBio	742698	740195633	8%	92%	325	1225	44272
NA12878_PacBio	651117	649109211	8%	92%	362	1206	38765
NA12879_PacBio	725437	691795401	7%	93%	321	1192	40661

^ Output from Picard CollectHS Metrics (Version 2.18.29) (On-bait bases - Number of bases that are uniquely mapped to the targeted regions of the genome; Off-bait bases - Number of bases that are uniquely mapped away from the targeted regions of the genome; Fold enrichment - The fold by which the targeted region has been amplified above genomic background)

*Nanopore Multiplexing Group – 1; #Nanopore Multiplexing Group – 2

Table 2: Genotype estimates on Nanopore and PacBio targeted capture sequencing using VNTRTyper

Sample	Method	Genotype of Target*								Correlation with PCR
		VNTR_8	VNTR_57	VNTR_86	VNTR_87	VNTR_93	VNTR_109	VNTR_112	VNTR_120	
NA12877	PCR	12.0/13.0	10.6/13.6	2.0/2.0	6.0/8.0	2.0/2.0	15.3/17.3	3.0/4.0	2.2/2.2	
	Nanopore_Capture	12.0/13.0	ND	2.0/2.0	6.2/8.4	2.0/2.0	15.3/17.3	3.0/4.1	2.2/3.2	0.9988
	PacBio_Capture	12.0/12.0	10.6/13.6	2.0/2.0	3.0/7.4	2.0/2.0	15.3/15.3	4.0/4.0	2.2/3.2	0.9845
NA12878	PCR	12.0/12.0	10.6/12.6	2.0/2.0	6.0/9.0	2.0/2.0	15.3/17.3	3.0/4.0	2.2/2.2	
	Nanopore_Capture	11.0/12.1	ND	2.0/2.0	6.0/8.8	2.0/2.0	14.8 /17.1	3.0/4.0	2.2/3.2	0.9978
	PacBio_Capture	12.0/12.0	10.6/12.6	2.0/2.0	6.0/6.0	2.0/2.0	15.5/15.5	3.0/4.0	2.2/2.2	0.9880
NA12879	PCR	12.0/13.0	10.6/10.6	2.0/2.0	8.0/9.0	2.0/2.0	15.3/17.3	3.0/3.0	2.2/2.2	
	Nanopore_Capture	12.4/12.4	10.6/10.6	2.0/2.0	6.0/8.6	2.0/2.0	14.8/16.9	3.0/4.0	2.2/3.2	0.9927
	PacBio_Capture	ND	ND	2.0/2.0	8.0/8.0	2.0/2.0	17.3/17.3	3.0/3.0	2.2/2.2	0.9944
NA12881	PCR	12.0/12.0	10.6/10.6	2.0/2.0	8.0/9.0	2.0/2.0	15.3/15.3	3.0/4.0	2.2/2.2	
	Nanopore_Capture	12.0/12.0	10.6/10.6	2.0/2.0	8.0/9.0	ND	15.3/17.3	3.0/4.0	2.2/3.2	0.9944
NA12882	PCR	12.0/13.0	10.6/10.6	2.0/2.0	6.0/6.0	2.0/2.0	15.3/17.3	3.0/3.0	2.2/2.2	
	Nanopore_Capture	11.8/13.0	10.6/10.6	2.0/2.0	6.0/8.5	2.0/2.0	15.3/17.3	3.0/4.0	2.2/3.2	0.9919
NA12889	PCR	12.0/12.0	13.6/17.6	2.0/2.0	8.0/8.0	2.0/2.0	17.3/17.3	4.0/4.0	2.2/2.2	
	Nanopore_Capture	12.1/12.1	10.6/15.8	2.0/2.0	6.1/8.1	2.0/2.0	14.8/17.2	3.0/4.0	2.2/3.2	0.9800
NA12890	PCR	12.0/13.0	10.6/10.6	2.0/2.0	6.0/9.0	2.0/2.0	15.3/15.3	3.0/3.0	2.2/2.2	
	Nanopore_Capture	12.2/12.2	10.6/10.6	2.0/2.0	6.0/9.0	2.0/2.0	13.3/15.3	3.0/4.0	2.2/3.2	0.9936

* Repeat numbers that do not agree with PCR results are highlighted in red.

ND – Sufficient data not available for genotype analysis

Table 3: Genotype estimates on NA12878 WGS and Capture sequencing data using VNTRTyper and Tandem-genotypes

Method^	Genotype of Target*										Corr. with PCR
	VNTR_8	VNTR_32	VNTR_57	VNTR_64	VNTR_86	VNTR_87	VNTR_93	VNTR_109	VNTR_112	VNTR_120	
PCR	12.0/12.0	9.4/10.4	10.6/12.6	17.3/18.3	2.0/2.0	6.0/9.0	2.0/2.0	15.3/17.3	3.0/4.0	2.2/2.2	
NPW_VT	12.0/12.0	8.2/10.7	10.6/12.6	17.3/18.6	2.0/2.0	6.0/9.0	2.0/2.0	15.3/17.3	3.0/4.0	2.2/3.2	0.9980
NPW_TG	11.0/12.0	5.4/9.4	10.6/12.6	18.3/19.3	2.0/2.0	6.0/9.0	2.0/2.0	15.3/17.3	3.0/4.0	2.2/2.2	0.9861
NPC_VT	11.0/12.1	ND	ND	ND	2.0/2.0	6.0/8.8	2.0/2.0	14.8/17.1	3.0/4.0	2.2/3.2	0.9978
NPC_TG	11.0/12.0	ND	ND	ND	2.0/2.0	6.0/9.0	ND	13.3/17.3	3.0/4.0	ND	0.9942
PBW_VT	12.0/12.0	11.0/12.4	10.6/12.6	18.3/19.3	2.0/2.0	6.0/9.0	2.0/2.0	15.3/17.3	3.0/4.0	2.2/2.2	0.9957
PBW_TG	12.0/13.0	9.4/10.4	12.6/12.6	18.3/18.3	2.0/2.0	6.0/9.0	2.0/2.0	15.3/18.3	3.0/4.0	2.2/2.2	0.9966
PBC_VT	12.0/12.0	ND	10.6/12.6	ND	2.0/2.0	6.0/6.0	2.0/2.0	15.5/15.5	3.0/4.0	2.2/2.2	0.9880
PBC_TG	12.0/13.0	ND	10.6/12.6	ND	ND	6.0/6.0	2.0/2.0	15.3/16.3	3.0/4.0	2.2/2.2	0.9865

^NPW_VT – Nanopore WGS VNTRTyper; NPW_TG – Nanopore WGS Tandem-genotypes; NPC_VT – Nanopore Capture sequencing VNTRTyper; NPC_TG – Nanopore Capture sequencing Tandem-genotypes; PBW_VT – PacBio WGS VNTRTyper; PBW_TG – PacBio WGS Tandem-genotypes; PBC_VT – PacBio Capture sequencing VNTRTyper; PBC_TG – PacBio Capture sequencing Tandem-genotypes

*Repeat numbers that do not agree with PCR results are highlighted in red.

ND – Sufficient data not available for genotype analysis