1 **Worker body size but not its age influences phototaxis in a bumblebee (*Bombus***
2 ***terrestris*, L.)**

4 Michal Merling, Shmuel Eisenmann, and Guy Bloch

5 Department of Ecology, Evolution and Behavior

6 The Hebrew University of Jerusalem

Abstract

We studied phtototaxis, the directional movement relative to light in the bumblebee *Bombus terrestris*.
We first developed and validated a MATLAB based system enabling reliable high-resolution tracking of a
bumblebee relative to a changing LED light source. Our tracking protocol enables us to separate the
phototaxis response from simple directional movement, overall levels of locomotor activity, or arousal state.
We next used this system to compare the phototactic response of workers varying in their body size, age,
or task performance. In all our experiments the bees showed a positive phototaxis. The strength of the
phototactic response was influenced by body size but not age, and this effect was significant when the light
source was weak. In one of two trials foragers that were larger showed stronger phototactic response than
nurses when tested with the weak light source. Taken together, the evidence that phototaxis is associated
with size-based division of labor in the bumblebee and with age-related division of labor in the honeybee,
lend credence to response threshold models implicating the response to light in the organization of division
of labor in cavity dwelling social insect.

Introduction

Phototaxis is a behavior in which an animal moves towards (positive phototaxis) or away (negative
phototaxis) from a source of elevated light intensity. The phototactic behavior is thought to be functionally
adaptive because it regulates light exposure and facilitate orientation in space. For example, positive
phototaxis may facilitate foraging behavior or escape in flying animals (e.g., Minot 1988; Reisenman et
al.,1998), and negative phototaxis may protect animals by guiding them towards places hidden from
predators (Hunte & Myers 2007 ;Minot 1988), or direct them towards burrowed nutrients (Sawin *et al.* 2004)
Phototactic behavior is sensitive to light intensity and is commonly used to assess the animal capacity to
discriminate between lights of different intensity. Pioneering studies with fruit flies have already shown early
in the 20th century that strong light can inhibit the phtototactic response that is induced by weaker light
intensities (Carpenter 1905). In regard to the underlying mechanisms, phototaxis is one of the first behaviors
for which mutations were identified (McEwen 1925). However, the phototactic behavior is not hardwired
and is modulated by external factors such as temperature and odors (Shimizu and Kato 1978; Inoko et al.

1981), and internal factors such as diet, nutritional state, the endogenous circadian clock or mating state
(Barker and Cohen, 1971; Inoko et al. 1981; Reisenman et al.,1998; Mazzoni *et al.* 2005; Bernadou and
Heinze 2013). The phototactic behavior is also commonly modulated during development. For example,
bird chicks switch from negative to positive phototaxis along the time they are ready to fledge (Minot 1988)
and in Drosophila the phototaxis of young larvae is negative but changes to positive near the end of the
third instar and at the final instar, during which it prepares for pupation seeking dry surfaces, rather than
the moist depths of fruits (Sawin *et al.* 2004; Mazzoni 2005).

The honeybee has been one of the first model systems with which to study phototactic behavior and its
underlying mechanisms (Bertholf 1927). As in other animals, phototaxis in honeybees was shown to be
sensitive to the light intensity (Bertholf 1927; Kaiser et al., 1977; Menzel & Greggers, 1985). The phototactic
behavior in bees can be mediated by multiple photoreceptors and is not restricted to a narrow wavelength
(Kaiser et al., 1977; Menzel & Greggers, 1985). In addition to the compound eyes, there is also evidence
that the ocelli are important for proper phototactic behavior in bees (Kastberger, 1990). The honeybee
provides some of the most striking examples for developmental modulation of phototaxis. First, the
developmentally determined caste of female bees affects their phototactic behavior during the adult stage.
Whereas phototaxis is negative in queens it is strongly positive in foraging worker bees. Additional
modulation occurs during the lifetime of worker bees. The phototactic behavior of the worker changes with
age and experience (e.g., orientation flights; Vollbehr, 1971). Later studies further linked this association to
age-related division of labor. Young workers typically tend brood inside the nest and show weaker
phototaxis compared with their older sisters that typically take foraging activities outside the nest (Ben-
Shahar et al., 2003). Pharmacological studies support this association by showing that treatment with
neurochemicals that modulate age-related division of labor or increase with age, such as cGMP, seretonin
(5-HT) and tyramine, can enhance the positive phototactic behavior as expected for a trait that is associated
with foraging activities (Ben-Shahar et al., 2003; Thamm et al., 2010, Scheiner et al., 2014). There is also
evidence suggesting an even finer task-related modulation of the phototactic behavior, which is associated
with the specialization of foragers in the pollen or nectar collection (Scheiner et al., 2014, Erber et al., 2006,
Tsuruda & Page, 2009). However, given that the differences between pollen and nectar foragers were not
consistent across studies, this interesting hypothesis needs to be further tested.

The association of task performance and phototaxis in honeybees is thought to be functionally significant
because strong positive phototaxis can guide bees towards the nest entrance and position them where they
may be induced to forage by exposure to additional foraging related stimuli. Weak or negative phototaxis
may be helpful for nurses because the brood is typically located deep in the dark cavity of the nest (Ben
Shahar et al. 2003). Here we study phototaxis in the bumblebee *Bombus terrestris,* in which the division of
labor relates to body size, with relatively little influence of age (e.g., Yerusalmi et al. 2006). Body size in
bumblebees is also correlated with anatomical, physiological and morphological differences that appear to
improve foraging performance. For example, large workers have larger eyes (Kapustjanskij et al. 2007;
Spaethe and Chittka 2003), and a greater density of antennal sensilla (Spaethe et al. 2007), which are

associated with better visual and olfactory accuracy. Their brain is larger (Mares et al. 2005; Riveros and
Gronenberg 2010) and they perform better in some learning assays (Riveros and Gronenberg, 2012;
Worden et al. 2005). They also have more cells expressing the circadian neuropeptide Pigment Dispersing
Factor (PDF; Weiss et al. 2009) and stronger circadian rhythms (Yerushalmi et al. 2006). It has been also
suggested that smaller bees are better suited to performing some in-nest activities (Brian 1952; Couvillon
and Dornhaus 2010). This individual variation is largely independent of genetic effects, because workers in
a bumblebee colony are typically the daughters of one singly-mated queen and are therefore closely
related. Given this association between body size, function and task performance, we hypothesized that
large bumblebee workers will have stronger phototaxis compared to their smaller sister bees. We also
hypothesized that age which has lower influence on task-performance will have a weaker effect, if at all, on
the phototactic response. To test these hypotheses we developed a new phototaxis monitoring system,
allowing continues tracking of the phototactic response of individually isolated bumblebees exposed to
various light intensities. We next used this system to compare the phototactic behavior of nurse and forager
workers, workers differing in body size, or workers differing in age. Our results show that body size but not
worker age is positively associated with phototaxis.

Materials and methods

Bees

We purchased *Bombus terrestris* colonies from Polyam Industries, Kibbutz Yad Mordechai (Israel).
Colonies were obtained a few days up to two weeks after the emergence of the first worker and contained
a queen, 1-50 workers (dependent on the experiment, see below), and brood at all stages of development.
We housed the colonies in wooden nest-boxes (30 x 23 x 20 cm) with a transparent Plexiglas cover. We
performed all treatments and inspections under dim red light which bumblebees do not see well (Peitsch et
al., 1992). We measured the length of each forewing twice as an index for body size (Yerushalmi et al.
2006).

Monitoring phototactic behavior

The phototaxis monitoring system included a stand with a perpendicular wooden board to which we
fixed a removable cage holder. The cage holders were made of 5 mm wide foamed cardboard (Kapa), and
each contained four parallel slots in which we placed the monitoring cages. The distance between the cages
was 5 cm. The monitoring cages (30x5x5 cm) were made of a similar foamed cardboard and had front and
side walls made of a transparent glass. In order to optimize contrast between the bee and the background,
we used white foamed cardboard for the posterior part of the cage that faced the camera. The bottom and
top walls were covered with opaque black cardboard to minimize exposure to light from the other cages. A
white LED light with spectral wavelength ranging from 440 to 700nm (main peak at 460 nm, and a lower

107 one at ~550) was placed in each side of each of the four cages (**Supp Fig. 1**). During the operation of the

108 LEDs, the temperature varied by ±1^{O}C. The LED lights were tested with a Polaroid polarizer to confirm that

109 LEDs do not produced polarized light. We positioned the LED lights such that they faced a 40 degree angle

110 downward. This positioning resulted in the light spot hitting a point next to the cage's edge, with no bleaching

111 to adjacent cages. To measure illumination intensity and spectrum we placed the sensor of an Ocean Optic

112 Spectrometer inside the cage in a distance of 1.5 cm from the LED light. A stand to which we fixed an

113 infrared sensitive camcorder (Sony TRV 75E) was placed at a distance of 1 meter from the monitoring

114 arena; the height of the camcorder was similar to that of the board with the cages (**supp Fig. 1**). The camera

115 produced dim infrared light that illuminated the arena during the experiment. The phototaxis experiments

116 were conducted in a dark environmental chamber (28 ± 1°C; 50-60% relative humidity).

118 ***Algorithm for automatic data acquisition and for phtototaxis***

119 For the automatic monitoring of worker position relative to the LED light source, and for the analysis of

120 the phototactic response we developed custom made tracking code using MATLAB (version 7.0; the Math

121 Works, Natick, Massachusetts, USA). The code identified an object with strong contrast relative to its

122 background within a defined arena; each of the four cages was defined as an arena. The algorithm identifies

123 the object (bee), tracks its location (center of mass), and records its position into a text file. It is also possible

124 to divide an arena into distinct "zones" and to calculate how much time the object (bee) stayed within each

125 zone (see *"Validation of the automatic phototaxis monitoring system" below)*.

126 We first converted the video records from the experiments into digital (AVI) formats with Pinaccle Studio

127 9 (Pinnacle, Systems, Mountain View, California). Next, we used the AVIedit software (AM Software,

128 Russia) to convert the digital video file into a series of JPEG digital 720 x 576 pixels photos (12.5 frames/

129 second). In order to minimize the influence of background noise we used Adobe Photoshop 7.0 (Adobe

130 Systems Inc, San Jose, California) to digitally remove the bees from a representative photo. This way we

131 obtained a photo of the background. This background photo, which included the image of the cages but not

132 the bees, was subtracted from all the digital photos (with bees) of the same experiment. We use the Image

133 J image analysis software (US National Institutes of Health, Bethesda, Maryland, USA) to record the X, Y

134 coordinates defining each one of the arenas (cages), and prepared two text files. In the first we recorded

135 for each frame the position (right or left) and state (on or off) of the light sources for each of the four arenas.

136 The second included the following information: photograph resolution, range of frames in which the light is

137 on, range of frame in which the light is off, X Y coordinates for the object (bee) in each arena, and the

138 identity of the focal bee (i.e., forager or nurse, large or small, young of old).

139 The algorithm was composed of several scripts. In the main script *(mmain)* the software removes the

140 background photo (without the bees, see above) from each frame in the file and performs an integral of the

141 contrast on the Y-axis for each one of the arenas using a predefined coordinates. The second script *(getbee)*

142 *extracted the* bee position on the x-axis for each frame was the average of the two most similar estimates

out of three possible position estimates that were calculated. The first was simply the X value corresponding
to the highest integral on the Y axis. Given that the integral over the Y-axis sums the entire range of the
cage, the shadow of the bee may be identify as the stronger contrast when the bee stands next to the light
source. Therefore, the second and third estimates were designed to correct for this potential bias. The
second estimate employs a specified script to identify the pixels with the sharpest change (the integral,
calculated using the script "*deriv*") in contrast value relative to neighboring pixels on the x-axis – this
produced two points corresponding to the front and back parts of the bee. The position of the bee is
calculated as the median between these two points (the "center" of the bee). The third estimate also uses
the *"deriv"* script but calculates an accumulative integral. This estimate was specifically useful for cases of
high background noise (for example, when the bee stands next to the light and causes apparent blinking in
light intensity. An addition script (*DecideX)* compares the three position estimates and send to the main
script the average for the two most similar estimates (see Supplementary Information for more details).

The positions of the bee at each frame were used to calculate the total path length she made during the
experiment and the Phototaxis Index (PTI). The PTI was calculated by assigning for each frame one of
three possible light position ("L" in Equation 1) values: -1 when the light is on the left side of the arena; 1
when the light is on the right side of the arena; 0 when the light is off on both sides. The position of the bee
on the x axis ("X" in Equation 1) was normalized such the -1≤X≤1, with -1, and 1 assigned to the left and
right edges of the cage respectively. Equation 1 was used for calculating the PTI, in which n is the total
number of frames analyzed by the algorithm for a specific bee:

Equation 1. $\text{PTI} = \frac{1}{n} \sum_{i=1}^{n} X_i L_i$.

The PTI value varied between 1 and – 1, with positive values indicating positive phototaxis (i.e. the bee
is close to the light), and negative values indicate negative phototaxis (i.e. the bee is far from the light). We
omitted from the phototaxis analyses bees that were not active or show very little level of activity during the
experiment.

Validation of the automatic phototaxis monitoring system

To confirm that our algorithm reliably capture the bee position we compared the automatically collected
data with the actual path for ten bees monitored in different experiments (6 in high light and 4 in low light
intensity). To determine the actual position of the bee, we used the ImageJ image analyses software to
analyses the first 400 frames in each of these 10 records. We inspected every forth frame in the set of the
first 400 frames and precisely determined the position of the bee center on the X-axis for each frame. We
recorded the bee position (in pixels) for each of the 100 frames and normalized the X values such that -1 ≤
X ≤ 1, similar to the normalization of our monitoring algorithm. Using this information, we calculated the
correlation coefficient between the automatically and manually estimated paths. We also used Equation 2
to calculate the percentage error for the automatically collected path.

Equation 2 $Err(\%) = \dfrac{\frac{100}{n} \sum_{i=1}^{n} \left| X_i^{manual} - X_i^{automatic} \right|}{D}$

Where n is the number of frames, Xi(Manual) is the normalized manually-determined position of the bee,

Xi(automatic) is the normalized software-determined position of the bee, and D is the length of the cage. D

= 2 because after normalization the X values ranged between -1 ≤ X ≤ 1.

In a second set of validations we compared the manual analyses of the phototactic response for the

bees investigated in Experiment 1 in which the bee position was assigned to one of five zones (rather than

a continuous measure as above). We first captured the video file into the computer (with Pinnacle Studio

9), and then divided each arena (cage) into five similar size zones ranging from the illuminated to the dark

side of the cage. For each bee we recorded the number of frames in which it was detected in each of the

five frames. As an index for the phototactic response of the bee we divided the number of times the bee

was observed in the zone next to the light source by the number of frames she was in the zone in the

opposite side of the cage. In this index, that we termed the "Light/ Dark index" values >1 indicate positive

phototaxis, and values <1 indicate negative phototaxis.

Experiment 1. The influence of worker task on the phototactic response

The bumble bee colonies were obtained about two weeks after the emergence of the first worker, and

contained approximately 50 workers. Immediately after receiving the colony, we marked all the workers

with individual numbered tag (Graze, Germany), and transferred the entire colony into an observation hive

made of transparent Plexiglas (32x24x13cm, described in Yerushalmi et al. 2006). The observation hives

have a cover with an opening allowing the removal and introduction of bees, and a cardboard bottom. We

placed the observation hive on shock absorbers to minimize vibrations to which bumblebees are very

sensitive. The observation hive was placed in an environmental chamber (28+-1C; 50-60% relative

humidity). All manipulations and observations were done under dim red light. During the first few days the

colony was fed *ad libitum* with commercial syrup (Polyam Pollination Services) and fresh pollen (collected

by honeybees). The hive was connected to the outside by a clear plastic tube (length = ~1 m, diameter = 2

cm). Food provision was gradually reduced and eventually stopped when a focal colony contained 10 - 30

workers (7-13 days from the emergence of the first worker). However, in cases in which the colonies did

not stored sufficient nectar or pollen, we supplemented the nest as needed in order to prevent larvae culling

by the nest bees. We performed detailed daily observations starting approximately a week after connecting

the colony to the outside. Each day we observed the entrance to the colony between 06:30 – 08:30 during

the morning, and 17:00 – 18:00 during the afternoon, for a total of three hours a day, and recorded the

number tags for bees returning from pollen or nectar foraging trips. Additional observations on the activity

inside the hive were performed daily between 09:00- 10:00 at the morning, and 16:00-17:00 at the

afternoon. During these observations we recorded the following behaviors: entering the hive with pollen or

nectar, building or manipulating wax cells, opening brood cells, inspecting egg cells, and feeding larvae

(see Yerushalmi et al. 2006; Shpigler et al. 2013 for more details).

We carried out the phototaxis experiments when the colony contained at least 30 active foragers. We

classified a worker bee as a forager if she was recorded performing ≥20 foraging trips, and no or very few

brood care activities during the three days preceding the phototaxis assay. We classified a bee as a nurse

if during the during this period she was recorded inspecting or manipulating brood cells, or feeding larvae,

and had never been observed flying outside the hive. All the phototaxis tests were done over three days,

always between 09:00 – 12:00. We made the experiments and observations at the same time of day in

order to prevent confounding influences of time of day. We collected foragers at the hive entrance, and a

similar number of nurses from inside the nest. Two nurses and two foragers were assigned randomly to

each of the four monitoring cages. To prevent a bias due to cage position, we confirmed that nurses and

foragers were not placed more often at a specific cage (position). The cages with the bees were placed in

the monitoring arena and were allowed to acclimate and relax for 10 minutes before beginning the

phototaxis test.

We preformed two trials with bees from colonies ET and HT. The protocols for these two trials differed

in several parameters. In the first trial with colony ET we used a light intensity which ranged between13.5

Lux (measured at the end of the cage closest to the lit LED) to 1.7 Lux (measured in the middle of the cage)

to 0.98 Lux (measured at the end of the cage furthest from the lit LED). These trials included 6 sessions in

which the light is on in one side of the cage. Each session lasted 3 minutes and was followed by 5 seconds

of darkness for the bee the recover. At the end of the day we determined the size of the bees and returned

them to the colony and the cages were cleaned with 70% ethanol. In the second trial with colony HT we

adjusted the protocol based on the lessons learned from the first trial. We shortened the sessions to 2

minutes each (since it takes a bee only about 5 second to walk the entire length of the cage) and limited

the number of sessions to 5 (lowering the duration of the entire test to about 10 minutes, instead of more

than 18 minutes in the first trial). We also lowered the light intensity to 0.3375 Lux (measured at the end of

the cage closest to the lit LED), 0.0425 Lux (measured at the middle of the cage) and 0.0245 Lux (measured

at the end of the cage furthest from the lit LED).

Experiment 2. The influence of body size on the phototactic response

The colonies from which we collected bees for this experiment were obtained a few days after the

emergence of the first worker and contained a queen, 1-10 workers, and brood at all stages. The bees were

marked and transferred to wooden cages (30x23x20 cm) with a transparent Plexiglass top. The colonies

were fed ad libitum and had no access to the outside. Every day we collected all the newly emerging bees

(easily identified by their lack of yellow pigmentation), recorded their size (front wing length), and marked

each individually with a number tag. We designated bees as "large" or "small" if they were larger or smaller

than the median of their colony, respectively. We attempted to collect bees on the edges of the size

spectrum (i.e., the largest and smallest in the colony) but the size comparison is nevertheless relative and

not in absolute terms. We tested only bees 2-5 days of age, and repeated the experiment with bees from

five different colonies.

For each phototaxis assay we collected 2 large and 2 small bees that were assigned randomly to the

monitoring cages (as in Experiment 1). The phototaxis assay included two parts; each part was composed

of six sessions of 1 min each, separated by 5 min of darkness. In the first part of the test we used low light

intensity (0.25µA), and in the second part high light intensity (10µA). The two parts of the test were

separated by 15 seconds of darkness.

257 *Experiment 3. The influence of worker age on the phototactic response.*

258   The experimental design and phototaxis test were overall similar to Experiment 2. In each trial of this

259 experiment we compared the phototactic response of two young bees (2-5 days of age) and two older bees

260 (16 – 19 days of age). We repeated this experiment with bees from three different colonies: EA, HA, GS

261 (that was also used in Experiment 2; each bee was tested only once).

263 *Statistical analyses*

We used parametric statistics for both the PTI and activity level for which the values fit normal

distribution. For each bee we calculated an average PTI value and therefore the average for each treatment

group was the mean of the averages which is normally distributed. In Exp. 1 in which we compared nurses

and foragers, we used t-tests, and in Experiments 2 and 3 we used three-way analysis of variance (ANOVA)

with treatment (size, or age), colony, and cage position as factors. For the statistical analyses of the Light/

Dark position in the first experiment we used the non-parametric Mann-Whitney test.

271 **Results**

272 *Validation of the automatic phototaxis monitoring system*

Our custom made algorithm for automatically tracking the bee position produced a track that was very

similar to that actually made by the bee (as determined by manual inspection of the bees position; **Fig. 1**).

The percentage error (see Material and Methods for details) was very low under both low light (mean %Error

= 0.71%, range = 0.45 – 0.84%), and high light (mean %Error = 1.81%, range = 1.13 – 2.27%; **Fig. 1C**).

The manually and software determined tracks were highly correlated in all 10 tracks we analyzed, with the

software produced record explaining in average 96% of the variation in the bee position (linear regression

analyses; P< 0.001 for all analyses; R^2 range = 0.809 – 0.998; **Fig. 1**).

In the second validation experiment we used the Light/Dark index and compared records that were determined manually and with the automatic monitoring system for bees from colony HT in Exp. 1. The results of the two analyses were overall very similar with a more positive phototaxis for foragers than the nurses (compare Colony HT in Fig. 1A and 1B). However, the differences in Light/Dark values were statistically significant only for the data collected automatically (**Fig. 2A, B**; Mann-Whitney test, two-tailed, P=0.155, P=0.024, for the manual and software collected data, respectively). The analysis with the software recorded data was probably more powerful because sampling rate was 25 times higher. Complementary analyses using the Phototaxis Index produced overall similar results for the two colonies (**Fig. 2C**). These two validation experiments indicate that our custom-made algorithm reliably tracks the bee position relative to the light source and produces a reliable and precise estimation of the phototactic behavior of the bee.

Experiment 1. The influence of worker task on the phototactic response.

The two repetitions of this experiment differed in the details of the phototactic response protocol. In the first repetition with bees from colony HT in which we used low light intensity (see Materials and Methods), the foragers, but not the nurses showed a positive phototaxis (single sample t-test compared with 0, P<0.001, P=0.14, respectively; **Fig. 2C**). The PTI value was significantly higher for the foragers (unpaired t-test, P<0.05, see also **Figs 2A** and **2B** for manual and automatic comparisons using the Light/Dark index, respectively). In this trial the foragers were significantly larger than the nurses (Two-tailed t-test, P<0.001). In the repetition with bees from colony ET in which the bees were tested with strong light, both the nurses and foragers showed a positive phototaxis (P<0.001), that was similar to the two task groups (**Fig.2C;** P=0.54; similar results were obtained using the manual Light/ Dark index, **Fig.2A**). In this trial, the foragers and nurses did not differ in body size (data not shown, perhaps due to strong bee-eaters predation that forced also relatively small bees to forage). This experiment shows that the phototactic behavior of bees can be modulated and suggest that body size is important because significant differences were obtained only in the trial in which the foragers were significantly larger than the nurses. To further asses the influence of body size on the phototactic behavior we next compared the phototactic response of same age small and large bees in colonies with no access to the outside.

Experiment 2. The influence of body size on the phototactic response of workers

We repeated this experiment with bees from five colonies that were tested with both low and high light intensities. Both the large and the small bees had a positive phototaxis in the two light intensities (P≤0.001 for all the analyses). Nevertheless, the positive phototactic response of large bees was significantly stronger when tested with low light intensity (**Fig. 3**; two-way ANOVA with colony and size as factors; colony, P=0.41; size, P=0.003; interaction, P=0.14). A similar trend was seen when the bees were assayed with strong light intensity, but the differences were not statistically significant (**Fig. 3**; colony, P=0.11; size, P=0.6; interaction, P=0.2).

Experiment 3. The influence of worker age on the phototactic response

We repeated this experiment with bees from three different colonies that were assayed with both low and high light intensities. Both the young and the old bees had a positive phototaxis when tested in the two light intensities ($P<0.01$). Overall, there was no significant influence of age on the phototactic response of workers (low light intensity: colony, $P=0.09$; age, $P=0.49$; interaction, $P=0.13$).

Discussion

We developed and validated a MATLAB based tracking system enabling reliable high-resolution tracking of individually isolated bumblebees. To estimate the phototactic response we exposed the bees to LED lights that can emit light over a range of intensities. Our algorithm continuously tracks the bee location and calculates the distance between the bee and the light source and thus provides precise quantification of the bee location relative to the light source. Importantly, we switched the light source between the two sides of the monitoring cage several times in each assay. This procedure enables us to separate the phototaxis response from simple directional movement, overall levels of locomotor activity, or arousal state. Our validation experiments in which we compared our tracking system to a visual frame-by-frame inspection of the bee location confirmed the precision and reliability of our system. This system can be easily adapted to study phototaxis in other insect species.

Following the validation of our phototaxis analysis system, we used it to begin investigating the modulation of the phototactic behavior of bumblebees. We focused on three factors that were shown to affect phototaxis in animals; age, body size, and task performance (see Introduction for references). In all the trials in all three experiments the bees moved toward the light, showing a positive phototaxis. However, the intensity of the phototactic response was modulated, with the stronger effect for body size. Larger bees showed overall stronger phototactic response compared with their smaller sisters. This difference was statistically significant only for bees tested with the weaker light intensity. The effect of light intensity is not surprising, because the phototactic response of animals is typically influenced by the intensity of the light source (Carpenter 1905; Yamaguchi et al. 2011; Menzel and Greggers, 1985; Erber et al. 2006). Thus, although small bees can show relatively strong positive phototaxis when light intensity is high, their weaker phototaxis relative to that of large bee suggest that they have a higher response threshold. A better description of the effect of body size on phototaxis can be achieved by studies using a broader range of light intensities. By contrast to the overall effect of body size, we did not find an affect of age. Pthototaxis is commonly influenced by age, specifically in animals that show ontogenic changes in behavior (Minot 1988; Sawin *et al.* 2004; Mazzoni 2005). These include the honeybee in which older bees show stronger phototactic response (Vollbehr, 1971; Ben Shahar et al., 2003). In honeybee colonies, workers show a form of behavioral development that underlies colony level division of labor. Young bees perform mostly

tasks inside the hive such as care for the brood, whereas older bees typically perform activities outside the nest such as guarding the hive entrance and foraging on flowers. It is assumed that a strong positive phototaxis helps guide foragers to the hive entrance and orient them outside, whereas weak or negative phototaxis is beneficial for nest bees that perform activities in the dark cavity of the nest (Ben-Shahar et al., 2003). Given that in bumblebees worker task relates to its body size, with only little effect to age (Yerushalmi et al. 2006; Michener et al. 1974; Alfrod 1975), the association of strong positive phototaxis with size in bumblebees and with age in honeybees is consistent with this hypothesis. An additional support for an association between task and phototaxis came from Exp. 1 in which we explicitly compared bees that specialized in nursing and foraging activities. Foragers showed stronger phototactic response only in the trial with bees from colony HT in which phototaxis was tested with the lower light intensity and in which the tested foragers were significantly larger than nurses. We do not know whether the lack of influence of task performance in the trial with colony ET is because this colony was tested with a stronger light intensity, because in this colony nurses and foragers did not differ in size, or because genetic differences between colonies ET and HT affect their phototactic behavior. Nevertheless, the results of the trial with colony HT strengthen the positive association between body size and phototaxis that we found in the second experiment. Additional studies with bees representing a diverse genetic background, that are beyond the scope of the current paper, should test whether and how foraging experience itself affects the phototacitc response.

In a broader view, our findings are consistent with predictions of response threshold models for the division of labor in insect societies. Workers in social insect colonies specialize in different tasks such as foraging, brood care, guarding or nest cleaning. This specialization forms a division of labor which is one of the most important organization principles of insect societies (Wilson 1971). But how does a worker knows which task it should take? Response threshold models state that workers have internal thresholds for responding to task-specific stimuli and that variation in task thresholds among workers in a colony generates division of labor (Beshers and Fewell, 2001; Beshers et al. 1999; Bonabeau and Theraulaz 1999). For social insects such as the Western honeybee, bumblebees and many species of ants that nest in dark cavities, light stimuli are associated with foraging outside the nest. Thus, response threshold models predict that individuals with strong positive phototaxis (i.e., lower response threshold to light) are more likely to approach the nest entrance and take foraging tasks compared to individuals showing a negative or weak positive phototaxis (Ben-Shahar et al. 2003; Erber et al. 2006). Given that in bumblebees larger bees are more likely to take foraging tasks, response threshold models predict that they will be more sensitive to foraging related stimuli compared to their smaller sisters, even when they have no previous foraging experience. The finding of Exp. 2 showing that large bees that never forage outside the nest nevertheless showed overall a stronger positive phototaxis than smaller bees meet this prediction. The influence of body size on phototaxis in the current study also adds to previous studies showing the large bees typically perform better in behaviors that are associated with foraging activities (Kapustjanskij et al. 2007; Spaethe and Chittka 2003; Riveros and Gronenberg, 2012; Worden et al. 2005; Yerushalmi et al.

2006). Taken together, these studies suggest that developmental processes that affect that final size of the

bumblebee larva (Shpigler et al. 2013) determine not only the size of the emerging worker bee, but also

many behavioral and physiological qualities that affect it propensity to specialize in a certain task.

References

Barker, R.J. & Cohen, C.F. (1971) Factors influencing phototaxis of houseflies. *J. Insect. Physiol.* **17**, 133-
142.

Ben-Shahar, Y., Leung, H.T., Pak, W.L., Sokolowski, M.B. and Robinson, G.E. (2003) cGMP-dependent
changes in phototaxis: a possible role for the foraging gene in honey bee division of labor.
Journal of Experimental Biology, **206**, 2507-2515.

Bernadou, A and Heinze, J (2013) Mating-Associated Changes in the Behaviour of *Leptothorax gredleri*
Ant Queens. ETHOLOGY 119**:** 634-643 **DOI:** 10.1111/eth.12103.

Bertholf L. M. (1927) The Relative Sensitivity of Honeybees to Light of Different Wave-Lengths, *Journal of*
Economic Entomology, 20: 521, https://doi.org/10.1093/jee/20.3.521

Beshers, S. N. and Fewell, J. H. (2001). Models of division of labor in social insects. *Annu. Rev.*
Entomol. 46,413 -440

Beshers S.N., Robinson G.E., Mittenthal J.E. (1999) Response thresholds and division of labor in
insect colonies. In: Detrain C., Deneubourg J.L., Pasteels J.M. (eds) Information Processing in
Social Insects. Birkhäuser, Basel

Bonabeau E., Theraulaz G. (1999) Role and variability of response thresholds in the regulation of
division of labor in insect societies. In: Detrain C., Deneubourg J.L., Pasteels J.M. (eds)
Information Processing in Social Insects. Birkhäuser, Basel

Brian AD (1952) Division of Labour and Foraging in Bombus agrorum Fabricius. Journal of Animal
Ecology 21:223-240.

Carpenter, F.W. (1905) The reactions of the pomace fly (*Drosophila ampelophilia* Loew*)* to light, gravity,
and mechanical stimulation. *Amer. Nat.* **39,** 157-171

Couvillon MJ, Dornhaus A, (2010) Small worker bumble bees (*Bombus impatiens*) are hardier against
starvation than their larger sisters. Insectes Sociaux 57:193-197. doi: 10.1007/s00040-010-0064-
7

Erber, J., Hoormann, J. and Scheiner, R. (2006) Phototactic behaviour correlates with gustatory
responsiveness in honey bees (Apis mellifera L.). *Behavioural Brain Research,* **174**, 174-180.

Hunte, W. & Myers R.A. (1984) Phototaxis and cannibalism in gammaridean amphipods. *Mar. Biol.* **81,**
75-79.

Inoko, H., Katsuki, M. & Watanabe, I. (1981) Suppression of Phototaxis in silkworm larvae. *Anim. Behav.*
29, 873-877.

Kaiser, W., Seidl, R. and Vollmar, J. (1977) Participation of all 3 color receptors in phototactic behavior of
fixed walking honeybees. *Journal of Comparative Physiology,* **122**, 27-44.

Kapustjanskij A, Streinzer M, Paulus HF, Spaethe J, (2007) Bigger is better: implications of body size for
flight ability under different light conditions and the evolution of alloethism in bumblebees.
Functional Ecology 21:1130-1136. doi: 10.1111/j.1365-2435.2007.01329.x.

Kastberger, G. (1990) The ocelli control the flight course in honeybees. *Physiol. Ent.* **15**, 337–346**.**

Mares S, Ash L, Gronenberg W, (2005) Brain Allometry in Bumblebee and Honey Bee Workers. Brain,
Behavior and Evolution 66:50-61.

Mazzoni, E. O., Despian, C. & Blau, J. (2005) Circadian pacemaker neurons transmit and modulate visual
information to control a rapid behavioural response. *Neuron* **45**, 293-300.

McEwen R.S. (1925) Concerning the relative phototropism of vestigial and wild type *Drosophila.* Biol. Bull.
Woods Hole, 49: 354-364

Menzel, R. and Greggers, U. (1985) Natural phototaxis and its relationship to color-vision in honeybees.
Journal of Comparative Physiology a-Sensory Neural and Behavioral Physiology, **157**, 311-321.

Minot, E.O. (1988) Sudden Reversal of Phototaxis during the development of nestling birds. *Anim. Behav.*
36, 1619-1625

Peitsch, D; Fietz, A; Hertel, H; Desouza, J; Ventura, Df; Menzel, R (1992) The spectral input systems of
hymenopteran insects and their receptor-based color-vision Journal Of Comparative Physiology
A-Sensory Neural And Behavioral Physiology 170: 23-40; DOI: 10.1007/BF00190398

Reisenman, C. E., Lazzari C. R. & Giurfa M. (1998) Circadian control of photonegative sensitivity in the
haematophgous bug *Triatoma infestans. J. Comp. Physiol. A.* **183,** 533-541.

Riveros AJ, Gronenberg W, (2010) Brain Allometry and Neural Plasticity in the Bumblebee Bombus
occidentalis. Brain, Behavior and Evolution 75:138-148. doi: 10.1159/000306506.

Riveros AJ, Gronenberg W (2012) Decision-making and associative color learning in harnessed
bumblebees (Bombus impatiens). Animal Cognition 15**:** 1183 1193; **DOI:** 10.1007/s10071-012-
0542-6

Sawin, E. P., Harris, L. R, Campos, A.R. & Sokolowski M.B. (1994) Sensorimotor transformation from
light reception to phototactic behaviour in *Drosophila* larvae (Diptera: Drosophilidae). *J. Insect Behav.*
7, 553-567.

Scheiner, R., Toteva, A., Reim, T., Sovik, E. and Barron, A.B. (2014) Differences in the phototaxis of
pollen and nectar foraging honey bees are related to their octopamine brain titers. *Frontiers in*
Physiology, **5**.

Shimizu, I. & Kato. M. (1978) Loss of phototaxis in silkworm larvae after smelling mulberry leaves, and
recovery after electroconvulsive shock. Nature, Lond., 272, 248-249.

Shpigler, H., Tamarkin, M., Gruber, Y. et al. Behav Ecol Sociobiol (2013) 67: 1601.
https://doi.org/10.1007/s00265-013-1571-0

Spaethe J, Chittka L, (2003) Interindividual variation of eye optics and single object resolution in
bumblebees. J Exp Biol 206:3447-3453.

Spaethe J, Brockmann A, Halbig C, Tautz J, (2007) Size determines antennal sensitivity and behavioral
threshold to odors in bumblebee workers. Naturwissenschaften 94:733-739. doi: 10.1007/s00114-
007-0251-1.

Thamm, M., Balfanz, S., Scheiner, R ,.Baumann, A. and Blenau, W. (2010) Characterization of the 5-
HT1A receptor of the honeybee (Apis mellifera) and involvement of serotonin in phototactic
behavior. *Cellular and Molecular Life Sciences,* **67**, 2467-2479.

Tsuruda, J.M. and Page, R.E. (2009) The effects of foraging role and genotype on light and sucrose
responsiveness in honey bees (Apis mellifera L.). *Behavioural Brain Research,* **205**, 132-137.

Vollbehr, J. (1971) Phototactic run behavior of honey bees (*Apis mellifica* l.). *Experientia,* **27**, 1174.&-

Weiss R, Dov A, Fahrbach SE, Bloch G, (2009) Body size-related variation in Pigment Dispersing Factor-
immunoreactivity in the brain of the bumblebee Bombus terrestris (Hymenoptera, Apidae). J
Insect Physiol 55:479-487. doi: 10.1016/j.jinsphys.2009.01.016.

Wilson E.O. (1971) The insect societies

Worden BD, Skemp AK, Papaj DR (2005) Learning in two context: the effects of interference and body
size in bumblebees. J Exp Biol 208:2045–2053

Yamaguchi, S; Heisenberg, M (2011) Photoreceptors and neural circuitry underlying phototaxis in insects.
FLY Volume: 5 Issue: 4 Pages: 333-336.

Yerushalmi S, Bodenhaimer S, Bloch G (2006) Developmentally determined attenuation in circadian
rhythms links chronobiology to social organization in bees. Journal of Experimental
Biology 209: 1044-1051; doi: 10.1242/jeb.02125

485 Figure 1.

(A)   **(B)**

(C)

Trail #	Light intensity	R^{2} [a]	% Error [b]
1	Low	0.994	0.64 %
2	Low	0.998	0.46 %
3	Low	0.993	084 %
4	Low	0.994	0.45 %
5	Low	0.994	0.72 %
6	Low	0.809	1.13 %
7	High	0.989	2.16 %
8	High	0.997	2.27%
9	High	0.865	1.27%
10	High	0.858	1.54%

a - Linear regression analysis

b - See text for details.

Figure 1. Validation of the automatic phototaxis monitoring system. We video recorded 10 bees
during the phototaxis assay and determined their position with the automatic tracking system and
manually by visually examining each frame. We determined the position of every 4th frame in the set of
the first 400 frames (n=100 frames). The Y-axis is the normalized position of the bee. **(A)** and **(B)** –
comparison of the automatically and manually recorded positions of two representative bees . **(A)** A nurse
bees assayed with low light intensity (Trial #4 in the table below). **(B)** A young bee assayed with high light
intensity (Trial #7 in the table below). **(C)** A table summarizing all 10 trials. The trials highlighted with grey
background correspond to the plots shown in (A) and (B) above.

Figure 2

Figure 2. The phototactic response of foragers and nurses. (A) Manual determination of the bee
position relative to the light source using the Light/ Dark index for the trails with colony ET (assayed with
low light intensity) and colony HT (assayed with high light intensity). **(B)** Calculation of the Light/ Dark
index based on the automatic phototaxis monitoring records for the trail with bees from Colony HT. **(C)**
Determination of the phototactic response using the Phototaxis Index. The bars show mean ± SE, sample
size within bars; * - P< 0.05. See text for more details and statistical analyses

Figure 3

Figure 3. The phototactic response of large and small bumble bee workers. The plots summarize
the results of five trials, each with bees from a different colonies, that were assayed with low (right) and
high (left) light intensities. We used two-way ANOVA with Trial (colony) and body size as factors. ** -
P<0.01. The Trial effect was not statistically significant in the assays with both low and high light intensity
(P>0.1, see text for details). Other details as in Fig. 2.

 Figure 4.

Figure 4. The phototactic response of young and old bumble bee workers. The plots summarize the
results of three trials, each with bees from a different colonies, that were assayed with low (left) and high
(right) light intensities. We used two-way ANOVA with trial (colony) and age as factors. The Trial effect
was not statistically significant in the assays with both low and high light intensity ($P > 0.1$, see text for
details). Other details as in Fig. 3.

Supplementary Figure S1

Supplementary Figure S1. Setup for monitoring the phototactic response of insects. (A) A general
scheme of the monitoring system. (B) A photo showing 4 cages, each with a single bee. The LED light on
the left side of the cages is on. C) Representative traces produced by the tracking algorithm.