Transcriptome analysis reveals the importance of the immune system during early pregnancy in sheep (*Ovis aries*)

Kisun Pokharel[1], Jaana Peippo[1], Melak Weldenegodguad[1,2], Mervi Honkatukia[3], Meng-Hua
Li[4*], Juha Kantanen[1*]

[1]Natural Resources Institute of Finland (Luke), Jokioinen, Finland

[2]Department of Environmental and Biological Sciences, Faculty of Science and Forestry, University
of Eastern Finland, Kuopio, Finland

[3]NordGen – The Nordic Genetic Resources Center, Ås, Norway

[4]CAS Key Laboratory of Animal Ecology and Conservation Biology, Institute of Zoology, Chinese
Academy of Sciences (CAS), Beijing, China

* Correspondence:
Prof. Meng-Hua Li
menghua.li@ioz.ac.cn
Prof. Juha Kantanen
juha.kantanen@luke.fi

Keywords: Finnsheep, Texel, progesterone, immunoglobulins, Siglec, endogenous retrovirus,
endometrium, corpus luteum, prolificacy, preimplantation

Running title: Sheep preimplantation transcriptome

1   Abstract

The majority of pregnancy loss in ruminants occurs during the preimplantation stage, which is thus
the most critical period determining reproductive success. While ovulation rate is the major
determinant of litter size in sheep, interactions among the conceptus, corpus luteum and endometrium
are essential for pregnancy success. To evaluate the role of reproductive tract function in sheep
fertility, we performed a comparative transcriptome study by sequencing total RNA (mRNA and
miRNA) from corpus luteum (CL) and endometrium tissues collected during the preimplantation
stage of pregnancy in Finnsheep, Texel ewes and F1 crosses. A total of 21,287 genes and 531
miRNAs were expressed in our dataset. Ten out of the top 25 most highly expressed genes were
shared, indicating the complementary functions of the CL and endometrium. Moreover, highly
expressed autosomal genes in the endometrium and CL were associated with biological processes
such as progesterone formation (*STAR* and *HSD3B1*) in the CL and facilitation of maternal
recognition of pregnancy, trophoblast elongation and implantation (*LGALS15*, *CST3*, *CST6*, and
EEF1A1) in the endometrium. In the CL, a group of sialic acid-binding immunoglobulin (Ig)-like
lectins (Siglecs), solute carriers (*SLC13A5*, *SLC15A2*, *SLC44A5*) and chemokines (*CCL5*, *CXCL13*,
CXCL9) were upregulated in Finnsheep, while several multidrug resistance-associated proteins
(MRPs) were upregulated in Texel ewes. Similarly, a number of Igs were upregulated in the
endometria of Finnsheep, while endogenous retroviruses (ERVs) were upregulated in the endometria
of Texel ewes. We also identified a novel ERV gene located in a reduced FecL locus that is
associated with prolificacy and is upregulated in prolific Finnsheep. Moreover, we report, for the first
time in any species, several genes that are active in the CL during early pregnancy (including

SIGLEC13, *SIGLEC14*, *SIGLEC6*, *MRP4,* and *CA5A*). Importantly, functional analysis of
differentially expressed genes suggested that Finnsheep have a better immune system than Texel
ewes and that high prolificacy in Finnsheep is governed by immune system regulation. Taken
together, the findings of this study provide new insights into the interplay between the CL and the
endometrium in gene expression dynamics during early pregnancy. The data and results will serve as
a basis for studying this highly critical period of pregnancy, which has wide significance in
mammalian fertility and reproduction.

2   Introduction

Litter size, a key determinant for the profitability of sheep production systems, is highly dependent
on ovulation rate and embryo development in the uterus. Earlier studies have shown that the trait of
high prolificacy can result due to the action of either a single gene with a major effect, as in the
Chinese Hu, Boorola Merino, Lacaune and small-tailed Han breeds (Chu et al., 2007; Davis et al.,
2002, 2006; Drouilhet et al., 2013; Mulsant et al., 2001; Souza et al., 2001), or different sets of genes,
as in the Finnsheep and Romanov breeds (Ricordeau et al., 1990; Xu et al., 2018). The Finnsheep or
Finnish landrace, one of the most highly prolific breeds, has been exported to more than 40 countries
to improve local breeds, although the heritability of ovulation rate is low (Hanrahan and Quirke,
1984). In recent years, a FecG^F (V371M) mutation in gene *GDF9* has been identified to be strongly
associated with litter size in Finnsheep and breeds such as the Norwegian White Sheep, Cambridge
and Belclare breeds, which were developed using Finnsheep (Hanrahan et al., 2004; Mullen and
Hanrahan, 2014; Pokharel et al., 2018; Våge et al., 2013).

The success of pregnancy establishment in sheep and other domestic ruminants is determined at the
preimplantation stage and involves coordination among pregnancy recognition, implantation and
placentation, in which the corpus luteum (CL) and endometrium play vital roles (Geisert et al., 1992;
Spencer et al., 2004b, 2007). The preimplantation stage of pregnancy is the most critical period in
determining the litter size because of the high embryo mortality during this period. It has been shown
that most embryonic deaths occur before day 18 of pregnancy in sheep (Bolet, 1986; Quinlivan et al.,
1966). However, due to the biological complexity of the process and to technical difficulties, embryo
implantation is still not well understood.

The CL is an endocrine structure whose main function is to synthesize and secrete the hormone
progesterone. Progesterone production is essential for the establishment of pregnancy. However, if
pregnancy is not established, the CL will regress as a result of luteolysis, and a new cycle will begin.
The endometrium is the site of blastocyst implantation, but its function is not limited to implantation.
The outer lining of the endometrium secretes histotroph, a complex mixture of enzymes, growth
factors, hormones, transport proteins and other substances that are key to conceptus survival and
implantation, pregnancy recognition signal production and placentation (Forde et al., 2013; Spencer
and Bazer, 2004). In addition, the endometrium also plays an important role in regulating the estrous
cycle (Spencer et al., 2008).

The whole-transcriptome profiling approach enables a deeper understanding of the functions of both
the CL and endometrium, which may allow the identification of genes and markers that are
differentially expressed, for example, between breeds showing different litter size phenotypes.
Although most of the studies associated with early pregnancy have been performed in sheep, only a
few studies have applied transcriptomic approaches to the endometrium and CL. A microarray-based
transcriptomic study conducted by Gray et al. (2006) identified a number of endometrial genes
regulated by progesterone (from the CL) and interferon tau (*IFNT*; from the conceptus) in pregnant

vs uterine gland knockout (UGKO) ewes. In a more comprehensive study conducted by Brooks et al.
(2016), transcriptome analysis of uterine epithelial cells during the peri-implantation period of
pregnancy identified various regulatory pathways and biological processes in sheep. Moore et al.
(2016) combined gene expression data with genome-wide association studies (GWASs) to
understand the roles of CL and endometrium transcriptomes in dairy cattle fertility. A study by Kfir
et al. (2018) identified differentially expressed genes (DEGs) between Day 4 and Day 11 in the CL in
cattle. Recently, a study on endometrial gene expression differences between Finnsheep and
European mouflon identified several genes associated with reproductive processes (Yang et al.,
2018). Though these studies have certainly enhanced our understanding of the roles of the CL and
endometrium during early pregnancy and in ruminant fertility in general, none of the studies have
conducted specific comparisons between breeds with different reproductive potential. Thus, in this
study, a comparison of transcriptome profiles between two breeds was conducted to provide insight
into the similarities in developmental events in early pregnancy between the breeds. More
importantly, we used F1 crosses in our experiment to understand the heritability of the genetic
markers.

3   Materials and Methods

3.1   Experimental design

All procedures for the feeding experiment and sheep sampling were approved by the Southern
Finland Animal Experiment Committee (approval no. ESAVI/5027/04.10.03/2012). The animals
were kept at Pusa Farm in Urjala, located in the province of Western Finland, during the
experimental period. A total of 31 ewes representing three breed groups (Finnsheep, Texel ewes and
F1 crosses) were included. Analyses were conducted for two different time points during the
establishment of pregnancy: the follicular growth phase (Pokharel et al., 2018) and early pregnancy
prior to implantation. After ovary removal (Pokharel et al., 2018), the ewes were mated using two
Finnsheep rams, and the pregnant ewes were slaughtered during the preimplantation phase of the
pregnancy when the embryos were estimated to be one to three weeks old. At the slaughterhouse, a
set of tissue samples (the pituitary gland, a CL, oviductal and uterine epithelial cells, and
preimplantation embryos) were collected and stored in RNAlater reagent (Ambion/Qiagen, Valencia,
CA, USA) following the manufacturer's instructions. Of the collected tissue samples, CL and
endometrium tissues were subjected to RNA-Seq. Endometrial samples were collected from the
uterine horns with a cytobrush, which was rinsed in a tube containing RNAprotect Cell Reagent
(Qiagen, Valencia, CA, USA). One of the CLs was dissected from each ovary. For the present study,
six ewes each from the Finnsheep, Texel and F1 cross groups were included. The experimental
animals and feeding experiments have been described in more detail in an earlier study (Pokharel et
al., 2018).

3.2   mRNA sequencing and data analysis

3.2.1 Library preparation and sequencing

We used an RNeasy Plus Mini Kit (Qiagen, Valencia, CA, USA) to extract mRNA from the tissues
according to the manufacturer's instructions. RNA libraries were prepared according to the Illumina
TruSeq® Stranded mRNA Sample Preparation Guide (part # 15031047). Unique Illumina TruSeq
indexing adapters were ligated to each sample during an adapter ligation step to enable pooling of
multiple samples into one flow cell lane. Only high-quality libraries were sequenced after the quality
and concentrations of the libraries were assessed with an Agilent Bioanalyzer 2100 and by Qubit®
Fluorometric Quantitation, Life Technologies, respectively. All samples were normalized and pooled

for automated cluster preparation at an Illumina cBot station. The samples were sequenced with an
Illumina HiSeq 2000 instrument using TruSeq SBS v3 sequencing chemistry. Paired-end sequencing
with a 2 x 100 bp read length was performed with a 6 bp index run.

3.2.2 Data preprocessing and mapping

The raw reads were assessed for errors and the presence of adapters using FastQC v0.11.6 (Simon
Andrews). As we noticed the presence of adapters, Trim Galore v0.5.0 (Felix Krueger; Martin, 2011)
was used to remove the adapters and low-quality reads and bases. The transcripts were quantified
under the quasi-mapping-based mode in Salmon v0.11.2 (Patro et al., 2017). We extracted the
FASTA sequences (oar31_87.fa) of the sheep transcriptome (oar31_87.gtf) using the gffread utility
(Trapnell et al., 2010) and built the transcriptome index. The resulting index was used for mapping
the RNA-Seq reads.

3.2.3 Differential expression

The results from Salmon were used to identify DEGs. DESeq2 (Love et al., 2013) was used to
compare gene expression differences. We started by considering both tissues, but after observing the
high variation between the endometrial samples, we decided to analyze the two tissues separately. All
replicates were collapsed before running DESeq. We set the filtering criteria for significant DEGs to
an adjusted p-value of 0.1 (*padj <1*) and an absolute log2(fold change) of greater than 1
(*abs(log2Foldchange)>1*). All the significant DEGs were annotated with Bioconductor biomaRt
(Durinck et al., 2005) for additional information (gene name, gene description, Entrez ID, human
ortholog and chromosome number).

3.2.4 Gene ontology and pathway analysis

The ClueGO v2.5.3 (Bindea et al., 2009) plugin in Cytoscape v3.7.0 (Shannon et al., 2003) was
employed for gene functional analysis. Prior to performing the analyses, we downloaded the latest
versions of the Kyoto Encyclopedia of Genes and Genomes (KEGG) pathways and Gene Ontology
(GO) terms. In addition, we retrieved Entrez gene IDs for all expressed genes in our dataset using the
biomaRt Bioconductor package. The enrichment analysis was based on a two-sided hypergeometric
test with the Bonferroni step-down correction method. We used a custom reference set that included
a list of all the expressed genes in our dataset. We also modified the default GO and pathway
selection criteria in such a way that a minimum of three genes and three percent of genes from a
given GO or KEGG pathway should be present in the query list. Furthermore, GO terms with a
minimum level of three and a maximum level of 5 were selected.

3.2.5 Manual annotation of select genes

When we noticed that many genes lacked gene annotations, we manually annotated those among the
top 25 most highly expressed genes in each tissue and the significant DEGs. First, we extracted the
coding sequence of each novel gene using Ensembl BioMart. All genes that had coding sequences
were BLASTed against the nonredundant (NR) nucleotide database. For the BLAST-based
annotation, we chose the hit with the highest coverage and the highest percentage of sequence
identity to the query sequence. Gene IDs that lacked coding sequences (CDSs) were queried back to
the Ensembl database to retrieve existing information. Throughout the paper (including in the
supplementary data files), the genes that were annotated based on the BLAST results have been
marked with an asterisk (*), while those that were annotated based on information available in
Ensembl are marked with a hash (#).

3.3   Small RNA sequencing and data analysis

3.3.1 Library preparation and sequencing

Both mRNA and miRNA were extracted from the tissues using an RNeasy Plus Mini Kit (Qiagen,
Valencia, CA, USA) following the manufacturer's protocol. The details on RNA extraction have
been described previously (Hu et al., 2015; Pokharel et al., 2018). RNA quality (RNA concentration
and RNA integrity number) was measured using a Bioanalyzer 2100 (Agilent Technologies,
Waldbronn, Germany) before sending the samples to the Finnish Microarray and Sequencing Center,
Turku, Finland, where library preparation and sequencing were performed. Library preparation for
mRNA and miRNA were performed according to Illumina's TruSeq® Stranded mRNA and TruSeq®
Small RNA sample preparation guides, respectively. The high quality of the libraries was confirmed
with an Agilent Bioanalyzer 2100, and the concentrations of the libraries were quantified by Qubit©
Fluorometric Quantification, Life Technologies. High-quality libraries of mRNA and miRNA were
sequenced with an Illumina HiSeq 2000 instrument using paired-end (2x100 bp) and single-end
(1x50) sequencing strategies, respectively.

3.3.2 Data preprocessing and analysis

The raw sequence data were initially screened to obtain an overview of the data quality, including the
presence or absence of adapters, using FastQC v0.11.6 (Simon Andrews). Next, the Illumina adapters
and low-quality bases were removed using Trim Galore v0.5.0 (Felix Krueger; Martin, 2011). In
addition, reads that were too short (having fewer than 18 bases) after trimming were also discarded.
To reduce downstream computational time, high-quality reads were collapsed using Seqcluster
v1.2.4a7 (Pantano et al., 2011). The FASTQ output from Seqcluster was first converted into a
FASTA file. The FASTA header was reformatted by including a sample-specific three letter code,
which is also a requirement for miRDeep2 analysis. For instance, ">A01_1_x446 A01" represents
sample C1033, whose first read was repeated 446 times.

The collapsed reads were mapped against the ovine reference genome (oar v3.1) using Bowtie
(Langmead et al., 2009). The Bowtie parameters were adjusted so that (1.) the resulting alignments
had no more than 1 mismatch (-v 1); (2.) the alignments for a given read were suppressed if more
than 8 alignments existed for it (-m 8); and (3.) the best-aligned read was reported (--strata, --best).
The alignment outputs (in SAM format) were coordinate-sorted and converted to BAM files. The
sorted BAM files were converted to the miRDeep2 ARF format using the "bwa_sam_converter.pl"
script.

miRDeep2 v2.0.0.5 (Friedländer et al., 2012) was used to identify known ovine miRNAs and to
predict conserved (known in other species) and novel ovine miRNAs. Before running the miRDeep2
pipeline, we merged both the collapsed FASTA files and the mapped ARF files. Furthermore, hairpin
and mature sequences of all species were extracted from miRBase v22 (Kozomara and Griffiths-
Jones, 2011, 2014). The extracted sequences were grouped into mature ovine sequences, ovine
hairpin sequences, and mature sequences for all species except sheep. The results from miRDeep2
were further processed to compile a list of all known and novel miRNAs. For novel and conserved
miRNAs, we designated provisional IDs that included the genomic coordinates of the putative mature
and star sequences.

3.3.3 Differential analysis

From the list of miRNAs discovered with miRDeep2, those with a minimum count of 10 reads across
all samples were considered for expression analysis. We used DESeq2 for expression analysis, in
which the technical replicates of three samples (C107, C4271 and C312) were collapsed prior to
running the DESeq command. We compared the expression levels between the breeds in each tissue

separately. The differentially expressed miRNAs with adjusted p-values less than 0.1 were regarded
as significant.

4   Results and Discussion

4.1   Phenotypic observations

After removal of the remaining ovary, we counted the number of CLs visually in each animal. With
an average of 4.09, Finnsheep had the highest number of CLs, whereas Texel ewes had an average of
1.7 CLs (Supplementary Table S1). On the other hand, F1 ewes showed phenotypes closer to those of
Finnsheep than those of Texel ewes, having 3.75 CLs on average (Supplementary Table S1). We did
not observe more than 2 CLs in the Texel group or fewer than 3 CLs in Finnsheep or F1 cross-bred
ewes. Similarly, on average, Finnsheep had the highest number of embryos (n=2.6), followed by F1
crosses (n=1.8) and Texel ewes (n=1.5). The F1 crosses displayed phenotypes similar to those of
Finnsheep; this was unsurprising, as we observed a similar pattern in an earlier study (Pokharel et al.,
2018). It would be of great interest to determine if productivity follows the same pattern in F2 (i.e.,
F1 x F1) crosses and backcrosses.

4.2   RNA-Seq data

From the 42 libraries (21 from each tissue, including three technical replicates), 4.4 billion raw reads
were sequenced, of which 4.2 billion clean reads were retained after trimming. The summary
statistics from Trim Galore revealed that up to 3.6% of the reads were trimmed, with reverse-strand
reads having a comparatively higher percentage of trimmed bases. However, the percentage of reads
that were excluded for being shorter than 18 bp was always less than 1% across all samples
(Supplementary table S2). Up to 70% of the high-quality reads were mapped to the ovine reference
transcriptome (Ensembl release 92).

4.3   Gene expression

A total of 21,287 gene transcripts were expressed in the whole data set, of which 1,019 and 959 were
specific to the endometrium and CL, respectively. Genes such as cytochrome P450, family 11,
subfamily A, polypeptide 1 (*CYP11A1*), C-C motif chemokines (*CCL21*, *CCL26*), serpin family A
members (*SERPINA1*, *SERPINA5*), inhibin subunit alpha (*INHA*) and paternally expressed 10
(*PEG10*) were specific to CL while genes related with solute carriers (*SLC44A4*, *SLC7A9*,
SLC34A2), ERVs were endometrium-specific (Table 1). Further grouping of the expressed genes
showed that the most genes (n =19,440) were expressed in endometrium samples of Texel ewes,
while the fewest genes (n =19,305) were expressed in endometrium samples of F1 crosses. The total
number of genes expressed in these tissues is comparatively higher than that in ovaries (Pokharel et
al., 2018). As shown in Fig. 1, the highest number of breed-specific genes expressed in the CL was
found in Finnsheep (n=254), followed by F1 crosses (n=204) and Texel ewes (n=199). Similarly,
from endometrium samples, we observed the highest number of unique genes in F1 crosses (n=284),
followed by Finnsheep (n=244) and Texel ewes (n=201). In a pairwise comparison, based on overall
gene expression, Finnsheep and Texel ewes shared a higher number of genes (n=260) in the CL than
in the endometrium. Moreover, Finnsheep and F1 crosses were found to share relatively more
common genes (n=278) than the other pairs (Fig. 1).

Although we observed considerable overlap of genes between tissues, principal component analysis
(PCA) of the 500 most highly expressed genes clearly indicated two distinct groups (Fig. 2A). In
addition to distinctiveness in terms of gene expression, the PCA plot also revealed the natures of the

tissue samples; the CL samples appeared to be more homogeneous than the endometrium samples.
The two sub-clusters within the endometrium cluster is linked to the age of the embryo (i.e. days
after mating). In addition, a heatmap plot based on the top 25 genes with the highest levels of gene
expression variation across all samples showed a similar pattern. However, we did not observe any
breed-specific clusters in either of the tissues, which was also the case in our earlier ovarian
transcriptome study. This result probably reflects the existence of only a few genes with breed-
specific differences. In addition, there is likely within-breed variation due to the stage of the estrous
cycle and other factors (e.g., the age of the animal). Because of the high variation between tissues, we
opted to perform gene expression analysis on each tissue separately.

4.4   Most highly expressed genes

One of the most interesting findings was the significant interplay between the CL and endometrium
during the preimplantation phase, as revealed by the most highly expressed genes. To obtain an
overview of the most abundant genes in each tissue, we selected the top 25 genes (Table 2). We
noticed that fifteen out of the top 25 genes were shared in both tissues, and the majority (9 out of 15)
were mitochondrial genes. Mitochondrial genes play prominent roles during reproduction. We have
also observed high levels of expression of mitochondrial genes in ovaries during the follicular growth
phase (Pokharel et al., 2018). Six shared autosomal genes also appeared to play substantial roles
during the preimplantation stage. Translationally controlled tumor protein (*TCTP*) is a highly
conserved, multifunctional protein that plays essential roles in development and other biological
processes in different species (Branco and Masle, 2019; Brioudes et al., 2010; Chen et al., 2007; Li et
al., 2011; Tuynder et al., 2002). With a maximum level of expression on Day 5 of pregnancy, this
protein has been shown to play a significant role in embryo implantation in mice (Li et al., 2011).
Consistent with these earlier studies, *TCTP* appeared to have the highest level of expression during
the embryo implantation period. Matrix Gla protein (*MGP*) is a vitamin K-dependent extracellular
matrix protein whose expression has been shown to be correlated with development and maturation
processes (Zhao and Nishimoto, 1996; Zhao and Warburton, 1997) and receptor-mediated adhesion
to the extracellular matrix (Loeser and Wallin, 1992). Several studies have reported that *MGP* is
highly expressed in the bovine endometrium (Forde et al., 2013; Mamo et al., 2012; Spencer et al.,
1999). The high level of expression of *MGP* in our study is consistent with the results of earlier
studies in which this gene was found to be elevated during the preimplantation stage in sheep (Gray
et al., 2006; Spencer et al., 1999) and cattle (Mamo et al., 2012). Similarly, Casey et al. (2005)
reported that *MGP* was significantly upregulated in nonregressed compared to regressed bovine CLs.
Our data and supporting results from earlier studies on cattle show that *MGP* is highly expressed in
both tissues during the preimplantation stage and plays important roles in superficial implantation
and placentation in sheep. In summary, gene regulation in the CL and endometrium is highly
coordinated and interlinked.
291
Six genes (*NUPR1*, *BCL2L15*, *CST3*, *CST6*, *S100G*, and *OST4*; see Table 2 for descriptions) specific
to the endometrium and one gene (*B2M*) common to both the CL and endometrium were also found
to be highly abundant in a recent study in which the authors compared gene expression changes in
the luteal epithelium and glandular epithelium during the peri-implantation stage in sheep (Brooks et
al., 2016). Galectin 15 (*LGALS15*) is induced by *IFNT* and is involved in conceptus development and
implantation (Gray et al., 2004; Kim et al., 2003; Lewis et al., 2007). *LGALS15* mRNA has been
detected in ewes from Day 9 until Day 12 (Satterfield et al., 2006). *IFNT* is secreted by the ovine
conceptus trophectoderm during the middle to late luteal phase and acts as the signal for maternal
recognition of pregnancy. Furthermore, *LGALS15* is an important gene that facilitates adhesion of the
trophectoderm to the endometrial luminal epithelium (Lewis et al., 2007; Spencer et al., 2007). Two

cystatin (CST) family members, namely, cystatin C (*CST3*) and cystatin E/M (*CST6*), were highly
expressed in the endometrium. Known for their importance during the elongation and implantation of
the conceptus, CSTs are protease inhibitors that are initiated by progesterone, and their high
expression levels are attributable to stimulation by *IFNT* (Spencer et al., 2008, 2015). Elongation
factor 1-alpha (*EEF1A1*) is an important component of the protein synthesis machinery because it
transports aminoacyl tRNA to the A sites of ribosomes in a GTP-dependent manner (Mateyak and
Kinzy, 2010; Tatsuka et al., 1992). The high levels of expression of *EEF1A1* in the endometrium
most likely correspond to the production and transport of progesterone and other molecules that are
essential during the implantation stage. The exact function of BCL2-like 15 (*BCL2L15*) in the sheep
endometrium is not known, nor has it been reported in the endometria of other species, but its high
expression has been reported previously (Brooks et al., 2016; Koch et al., 2010; Romero et al., 2017).

Oxytocin (*OXT*) was one of the most highly expressed genes in the CL. In cyclic ewes, *OXT* secreted
from the CL and posterior pituitary is widely known to bind with oxytocin receptor (*OTR*) from the
endometrium to concomitantly release prostaglandin $F_{2\alpha}$ (*PGF*) pulses and induce luteolysis (Bazer,
2013; Flint and Sheldrick, 2004; Spencer et al., 2004a, 2004b). However, for noncyclic ewes, *OXT*
plays an important role during peri-implantation and throughout pregnancy (Kendrick, 2000). *OXT*
signaling is known to be influenced by progesterone, but the mechanism underlying the regulation is
not yet clear due to conflicting findings (Bishop, 2013; Fleming et al., 2006; Gimpl et al., 2002;
Grazzini et al., 1998). *OTR* expression in both the CL and endometrium was almost negligible
compared to OXT expression. Steroidogenic acute regulatory protein (*STAR*) plays an important role
in mediating the transfer of cholesterol to sites of steroid production (Christenson and Devoto, 2003;
Stocco, 2000). Post ovulation, the expression of the majority of genes associated with progesterone
synthesis starts to increase and peaks around the late luteal phase, when the CL has fully matured
(Davis and LaVoie, 2018; Devoto et al., 2001; Juengel et al., 1995). *STAR*, together with the
cytochrome P450 side chain cleavage (P450cc) complex and 3b-hydroxysteroid
dehydrogenase/delta5 delta4-isomerase (*HSD3B1*), are the three most important actors involved in
progesterone biosynthesis. *STAR* is involved in transporting free cholesterol to the inner
mitochondrial membrane. The P450cc complex, composed of a cholesterol side chain cleavage
enzyme (*CYP11A1*), ferredoxin reductase (*FDXR*) and ferredoxin (*FDX1*), converts the newly arrived
cholesterol into pregnenolone (King and LaVoie, 2009). Finally, *HSD3B1* helps in converting
pregnenolone to progesterone (Davis and LaVoie, 2018; Hu et al., 2010; Plant et al., 2015; Stouffer
and Hennebold, 2015). Two of these major genes involved in progesterone synthesis (*STAR* and
HSD3B1) were ranked among the top 25 most highly expressed autosomal genes, while *CYP11A1*
(TPM=2,080), *FDXR* (TPM=86) and *FDX1* (TPM=1,206) were also highly expressed.

4.5   Differentially expressed genes

Overall, the CL appeared to display higher levels of gene expression differences between the breeds
than the endometrium (Table 3). This tissue-specific difference is consistent with the findings of a
recent study in cattle in which Moore et al. (2016) identified nine and 560 DEGs in the endometrium
and CL, respectively, between fertile and infertile cows. In the three possible pairwise comparisons
for each tissue, the highest number of DEGs (n=199) was found between the pure breeds in the CL,
while the fewest DEGs (n=2) were found between the Finnsheep and F1 crosses in the endometrium.
In both tissues, the pure breed comparisons had the highest numbers of DEGs, but the two
comparisons (for both the endometrium and CL) that involved the F1 crosses had the lowest. In other
words, in the CL, Finnsheep had more DEGs (n=67) than F1 crosses, whereas in the endometrium,
Texel ewes had more DEGs (n=17) than F1 crosses.

4.6   Differentially expressed genes: Corpus luteum

Out of the 199 significant DEGs in the CL, 140 were upregulated in Finnsheep (Supplementary table
S3). However, 91 out of the 199 genes lacked annotations (i.e., a gene name and gene description).
We were able to retrieve the CDSs for 82 genes and employed a BLAST search against the NR
database. Based on the Ensembl ncRNA prediction system, two out of nine genes that lacked CDSs
were predicted to be miRNAs, and the rest were lincRNAs.

In the list of DEGs, we observed a few cases in which more than one gene from the same family was
present. All eight genes related to multidrug resistance-associated proteins (*MRPs*) were upregulated
in Texel ewes, of which seven genes were type 4, while one was type 1. Both *MRP1* and *MRP4* are
lipophilic anion transporters. Earlier reports have suggested a role of *MRP4* in transporting
prostaglandins in the endometrium (Lacroix-Pépin et al., 2011), and *MRP4* has been found to be
upregulated in the endometrium in infertile cows compared to fertile cows (Moore et al., 2016).
Although there are no reports regarding the existence and roles of both *MRP4* and *MRP1* in the CL,
we speculate that the comparatively lower levels of these prostaglandin (PG) transporters in
Finnsheep provide a luteoprotective effect. Six sialic acid-binding Ig-like lectins (Siglecs) were
upregulated in Finnsheep. Based on a BLAST search and on the information available in Ensembl,
the sequences were related to *SIGLEC-5* (*ENSOARG00000002701*), *SIGLEC-13*
(*ENSOARG00000014846* and *ENSOARG00000014850*) and *SIGLEC-14* (*ENSOARG00000014875*,
ENSOARG00000002909, and *ENSOARG00000001575*). Siglecs are transmembrane molecules that
are expressed on immune cells and mediate inhibitory signaling (Varki and Angata, 2006). So far,
SIGLEC-13 has been reported only in nonhuman primates; it was deleted during the course of human
evolution (Angata et al., 2004). The importance of Siglecs in immune system regulation has been
reviewed elsewhere (Pillai et al., 2012). Siglecs constantly evolve through gene duplication events
and may vary between species and even within a species (Bornhöfft et al., 2018; Cao and Crocker,
2011; Pillai et al., 2012). We can conclude from the observations made in this study that Siglecs also
play highly important roles during early pregnancy (in preimplantation and pregnancy recognition)
through immunoregulatory activities and may even dictate prolificacy in sheep.

Similarly, three genes related to phospholipase A2 inhibitor and Ly6/PLAUR domain-containing
protein-like (*PINLYP*) were upregulated in Finnsheep. Other genes with more than one member
included major histocompatibility complexes (MHCs) (*BOLA-DQA5*, *HLA-DMA*, *HLA-DRA*, *MICA*,
*BOLA-DQB*2001*, etc.), chemokines (*CCL5*, *CXCL13*, and *CXCL9*), solute carriers (*SLC13A5*,
SLC44A5, and *SLC15A2*), interleukin receptors (*IL2RG*, *IL12RB1*, and *IL12RB2*), cluster of
differentiation factors (CDs) (*CD52*, *CD74*, and *CD300H*), granzymes (*GZMM* and
LOC114109030/*GZMH*), calcium homeostasis modulators (*CALHM3* and *CALHM5*) and
neurofilaments (*NEFL* and *NEFM*). Five out of seven significantly differentially expressed lincRNAs
were upregulated in Finnsheep. *GZMM* and *GZMH* have been found to be upregulated in Yakutian
cattle compared to Finncattle and Holstein cattle (Pokharel et al., 2019).

Out of 140 genes that were upregulated in Finnsheep, 50 genes (35.71%) were associated with 48
different GO terms (Fig. 3, Supplementary table S6) within the biological processes category,
whereas 90 genes lacked GO annotations. The upregulated genes were associated with positive
regulation of different processes ("T cell migration", "cytokine production", "defense response",
"immune effector process", "immune system process", "interferon-gamma production", "leukocyte
chemotaxis", "response to external stimulus", "smooth muscle cell proliferation", "adaptive immune
response based on somatic recombination of immune receptors built from immunoglobulin
superfamily domains", "T cell activation", "behavior", "cell–cell adhesion", "cytokine-mediated
signaling pathway", "leukocyte activation", and "lymphocyte proliferation"). Some biological
processes potentially associated with implantation were "maintenance of location", "plasma

membrane invagination", "import into cell", "chemotaxis", and "receptor internalization". Other
biological processes, such as "response to bacterium", "response to lipopolysaccharide",
"lymphocyte-mediated immunity" and "chemotaxis", could be associated with adaptation of
Finnsheep to the rugged Finnish climate and with disease resistance.

Similarly, only 40 of the 140 genes upregulated in Finnsheep were associated with 29 KEGG
pathways (Supplementary fig. 1, Supplementary table S5). The majority of the pathways were
associated with diseases; "tryptophan metabolism", "cell adhesion molecules (CAMs)", "Th1 and
Th2 cell differentiation" and "Th17 cell differentiation" appeared to play roles in implantation. Out
of 59 genes that were downregulated in Finnsheep, 17 and 14 genes were associated with GO IDs
and KEGG pathways, respectively. However, after applying our selection criteria, only one
biological process, "negative regulation of endopeptidase activity" (associated genes: *COL28A1*,
LOC101104482, and *SLPI*) and one KEGG pathway, "bile secretion" (associated genes:
LOC101106409, *LOC101107772*, and *LOC101112460*), were identified.

Altogether, 67 genes were differentially expressed between Finnsheep and F1 crossbred ewes, of
which 49 genes were upregulated in Finnsheep (Supplementary table S6). *CA5A* is a member of the
carbonic anhydrase family of zinc-containing metalloenzymes, whose primary function is to catalyze
the reversible conversion of carbon dioxide to bicarbonate. The mitochondrial enzyme *CA5A* plays
an important role in supplying bicarbonate (HCO_{3-}) to numerous other mitochondrial enzymes. In a
previous study, we observed downregulation of *CA5A* in the ovaries of Texel ewes compared to F1
ewes (Pokharel et al., 2018). More recently, *CA5A* was also identified in the ovaries of the Pelibuey
breed of sheep; the gene was upregulated in a subset of ewes that gave birth to two lambs compared
to uniparous animals (Hernández-Montiel et al., 2019). However, there are no reports regarding the
expression and function of *CA5A* in the CL. Based on the results from our current and earlier reports
(Hernández-Montiel et al., 2019; Pokharel et al., 2018), *CA5A* appears to have an important function,
at least until the preimplantation stage of reproduction. The level of expression in F1 crosses in the
CL and endometrium followed the same pattern as that in the ovary, which led us to conclude that
CA5A is heritable and potentially an imprinted gene. Further experiments are needed to determine
whether the gene is associated with high prolificacy. Out of the 49 upregulated genes, 24 genes had
available functional annotations and were associated with nine different GO terms (Fig. 5,
Supplementary table S7). The majority of the GO terms were related to transport ("anion transport",
"lipid transport", "organic anion transport", and "fatty acid transport") and regulation ("regulation of
lipid transport", "regulation of homotypic cell–cell adhesion", "negative regulation of T cell
activation", and "regulation of lipid localization").

Altogether, 20 out of the 49 genes upregulated in Finnsheep vs F1 crosses were linked to KEGG
pathways. Based on the selection criteria, only two KEGG pathways, namely, "complement and
coagulation cascades" (associated genes *C5AR1*, *F13A1*, and *VSIG4*) and "Fc gamma R-mediated
phagocytosis" (associated genes: *FCGR1A*, *SCIN*, and *SYK*), were identified. The lowest number
(n=22) of DEGs was observed between Texel and F1 crossbred ewes, with 13 genes being
upregulated in Texel ewes (Supplementary table S8).

4.7   Differentially expressed genes: Endometrium

The highest number of significant DEGs (n=43) was present in the comparison of the pure breeds; of
the DEGs, 22 were upregulated in Finnsheep (Supplementary table S9). The majority of the genes
(29/43) lacked Ensembl annotations and were labeled as "novel genes". Therefore, we performed a
BLAST search using the cDNA sequences of these genes against the NR database. Homology-based
annotation revealed that half of the genes upregulated in Finnsheep were associated with

immunoglobulins (Igs). Igs are heterodimeric proteins that belong to the Ig superfamily (IgSF)
(Williams and Barclay, 1988). Igs are composed of two heavy and two light chains, and the light
chain may further consist of a κ or λ chain (Williams and Barclay, 1988). Interestingly, the structure
and organization of the genes enable Igs to be receptive to a virtually unlimited array of antigens
rather than being limited to a fixed set of ligands (Honjo, 1983). This feature is particularly important
for adaptation to changing environments and may have contributed to enabling Finnsheep, for
example, to survive in the harsh Finnish climate. Studies on humans have shown that Igs, in general,
improve pregnancy success (Coulam and Goodman, 2000; De Placido et al., 1994). In addition to 11
Ig genes representing both the light and heavy chains, the joining chain of multimeric IgA and IgM
(*JCHAIN*) was also upregulated in Finnsheep. *JCHAIN* is a small polypeptide containing eight
cysteine residues that makes disulfide (C-C) bonds with IgA and IgM to form multimers. Two of the
eight cysteines are linked with cysteines available on the heavy chain of IgA or IgM to result in
dimer or pentamer forms, respectively (Bastian et al., 1995). We also identified four genes associated
with endogenous retroviruses (ERVs) that were upregulated in Texel ewes. ERVs are copies of
retroviral genomes that have been integrated into the host genome during evolution. Sheep ERVs
share sequence similarity with exogenous and pathogenic Jaagsiekte sheep retrovirus (JSRV)
(DeMartini et al., 2003). The genome of sheep contains at least 32 ERVs related to JSRV (Sistiaga-
Poveda and Jugo, 2014), and these ERVs are essential during pregnancy, including during placental
morphogenesis and conceptus elongation (Dunlap et al., 2006b; Palmarini et al., 2001; Spencer and
Palmarini, 2012). A number of earlier studies have suggested critical roles of *enJSRV*s in uterine
protection from viral infection, preimplantation conceptus development and placental morphogenesis
(Denner, 2016; Dunlap et al., 2005, 2006b, 2006a). We speculate that the upregulation of ERVs in
Texel ewes is possibly due to breeding and might be linked to certain phenotypes (e.g., meat quality)
that have been desired in Texel ewes. The novel gene (ENSOARG00000009959) upregulated in
Finnsheep is located on the reverse strand of chromosome X and has 24 paralogs. This gene is not
listed for 162 (out of 184) species available in the Ensembl database. Although Ensembl lists 71
orthologs of this gene, none of them have even 50% sequence homology. A BLAST search against
the NR database showed that 97% of the bases matched to the region of the reduced FecL locus
(GenBank ID KC352617.1), which was recently characterized (Drouilhet et al., 2013). So far, only
two genes, beta-1,4 N-acetylgalactosaminyltransferase 2 (*B4GALNT2*) and insulin-like growth factor
2 mRNA-binding protein 1 (*IGF2BP1*), and a pseudogene, ezrin-like protein, have been identified to
exist in that region; our results have added one more gene. In addition to the finding that the best hit
was related to the FecL locus, the gene appeared to be an ERV, as we noticed that the query gene had
98% sequence identity with a partial sequence of the endogenous-virus beta-2 pro/pol region (see
also Fig. 4).
474
Six ncRNAs, including four lincRNAs, one snoRNA and one miRNA, were also in the list of DEGs
between the pure breeds; among these, U3 snoRNA (*RF00012*), a miRNA (*ENSOARG00000022916*)
and two lincRNAs (*ENSOARG00000026131* and *ENSOARG00000026482*) were upregulated in
Finnsheep, while the other two lincRNAs (*ENSOARG00000025875* and *ENSOARG00000026176*)
were upregulated in Texel ewes. LincRNAs are long ncRNAs (lncRNAs) that originate from
intergenic regions and do not overlap a protein-coding transcript. LincRNAs have a wide array of
functions, including transcriptional regulation, biogenesis, epigenetic regulation, tissue specificity
and developmental patterning (see reviews by Deniz and Erman, 2017; Pauli et al., 2011; Ransohoff
et al., 2018; Ulitsky and Bartel, 2013).

Only two genes (*OLFM1* and *ENSOARG00000022916,* which encodes a miRNA) were significantly
differentially expressed between Finnsheep and F1 crosses, with both upregulated in Finnsheep
(Supplementary table S10). Similarly, 17 genes were significantly differentially expressed between

Texel ewes and F1 crosses, among which five were upregulated in Texel ewes (Supplementary table
S11). Due to the small number of genes and, more importantly, due to a lack of gene annotations, we
were not able to perform GO and KEGG pathway analyses for the DEGs in the endometrium.

4.8 Overlapping differentially expressed genes

We noticed that as many as 54 genes were differentially expressed in more than one comparison,
increasing our confidence in the identification of these DEGs. Few DEGs were exclusively up- or
downregulated in a breed irrespective of the tissue. For example, a transcript encoding a miRNA
(*ENSOARG00000022916*) was found to be significantly differentially expressed in four different
comparisons in both tissues and was always upregulated in Finnsheep compared to both Texel ewes
and F1 crosses. *HNRNPK* was always downregulated in Finnsheep, and *CA5A* was always
upregulated in F1 crosses compared to the other two breeds. Similarly, coiled-coil domain-containing
73 (*CCDC73*) and a pseudogene (*ENSOARG00000020196*) were upregulated in Texel ewes
compared to F1 crosses. A lincRNA (*ENSOARG00000025875*) was downregulated in Finnsheep
compared to Texel ewes. *MICA** was upregulated in Finnsheep compared to Texel ewes.

In endometrium samples, seven Igs were significantly differentially expressed in two comparisons
(Finnsheep vs Texel ewes and Texel ewes vs F1 crosses), and in all cases, they were downregulated
in Texel ewes. This finding clearly indicates that Igs play more important roles in Finnsheep than in
Texel ewes, and importantly, these roles were inherited by F1 crosses. Genes that were exclusively
upregulated in Texel ewes were centrosomal protein 162 kDa (*CEP162**), kallikrein-related
peptidase 14 (*KLK14*), Plexin A4 (*PLXNA4*), contactin 1 (*CNTN1*), butyrophilin like 2 (*BTNL2*), and
antileukoproteinase precursor (*SLPI*). In addition to seven Igs, phospholipase A2 group IB
(*PLA2GIB*) and lincRNA (*ENSOARG00000026482*) were also downregulated in Texel ewes.

Oxidized low-density lipoprotein receptor 1 (*OLR1*), NLR family apoptosis inhibitory protein
(*NAIP*), macrophage scavenger receptor 1 (*MSR1*), high-affinity Ig gamma Fc receptor 1 precursor
(*FCGR1A*), hemoglobin subunit alpha-1/2, folate receptor 3 (*FLOR3**), Fc gamma 2 receptor,
chromogranin B (*CHGB*), Siglec-14, clavsin 2 (*CLVS2*), copine 4 (*CPNE4*), EPH receptor B6
(*EPBH6**) and *MICA** were exclusively upregulated in the CLs of Finnsheep. Gastrula zinc finger
XICGF17.1-like (*LOC10562107*), crystallin mu (*CRYM*), myeloid-associated differentiation marker-
like (*LOC101119079*) and tolloid-like 2 (*TLL2*) were downregulated in the CLs of Texel ewes
compared to the other two breeds. These results also indicated that F1 crosses were more similar to
Finnsheep than Texel crosses. *CA5A* appeared to be upregulated in F1 crosses compared to both
Finnsheep and Texel ewes in all three analyses (i.e., in the CL and endometrium in this study and in
the ovary in our earlier study). A miRNA, *ENSOARG00000022916*, was significantly differentially
expressed in four different comparisons (Finnsheep vs Texel ewe CL, Finnsheep vs F1 CL,
Finnsheep vs Texel ewe endometrium, and Finnsheep vs F1 endometrium), and it was always
upregulated in Finnsheep. *HNRNPK* was always downregulated in Finnsheep (Finnsheep vs Texel
ewe CL, Finnsheep vs F1 CL, and Finnsheep vs Texel ewe endometrium).

4.9 miRNAs expressed in the dataset

A total of 336.6 M reads were sequenced, of which approximately 42% contained adapters and/or
low-quality bases. After trimming, more than 92% of the reads (n=311.3 M) were retained as high-
quality clean reads. On average, collapsing of duplicate reads revealed 483,096 unique reads per
sample, of which 54.4% of the unique sequences (collapsed reads) were mapped to the ovine
reference genome. The detailed summary statistics for each sample are shown in Supplementary table
S12. There were more collapsed reads and uniquely mapped reads for endometrial samples than CL

samples despite the similar numbers of raw and clean reads in both tissues. After filtering out low-
count and ambiguous reads, a total of 531 miRNAs were included in the expression analysis. The
majority of the expressed miRNAs (n=456) were shared by both tissues, with 43 and 32 miRNAs
being unique to the CL and endometrium, respectively. Currently, 153 miRNAs are available in the
miRBase database (Kozomara et al., 2019). The database was updated to the current version (22)
from an earlier version (miRBase 21) after four years, and the overall number of miRNA sequences
increased by over a third. However, the number of sheep miRNAs remained the same. Moreover,
studies that produce miRNA datasets have been scarce. As of April 2019, miRNA datasets from only
three studies were available in the European Nucleotide Archive (ENA) database, with accession
codes PRJNA308631 (n=3), PRJEB22101 (n=37) and PRJNA414087 (n=40); the PRJEB22101
dataset was from the first phase of this study (Pokharel et al., 2018). In the current study, we
quantified over threefold more sheep miRNAs (n=453) than are available in miRBase. Therefore,
these miRNAs will certainly improve the existing resources and will be valuable in future studies.

Two miRNAs, both upregulated in Finnsheep, were significantly differentially expressed between the
pure breeds in the CL, while the other comparisons did not reveal any significantly differentially
expressed miRNAs (Table 4). Of these two significantly differentially expressed miRNAs, rno-miR-
451-5p is a conserved miRNA similar to one found in rats (*Rattus norvegicus*). The other, oar-
18_757_mt, is a novel miRNA expressed on chromosome 18. For the endometrium dataset, one
miRNA in the Finnsheep *vs* Texel ewe and Texel ewe *vs* F1 cross comparisons and two miRNAs in
the Finnsheep *vs* F1 cross comparison were significantly differentially expressed (Table 4). All four
significantly differentially expressed miRNAs in the endometrium are novel miRNAs, three of which
are on chromosome 18, while one is located on an unplaced scaffold (JH921839.1) in the current
assembly (oar v3.1) of the sheep genome. Moreover, we identified that oar-18_882 was exclusively
significantly upregulated in the endometrium samples of Texel ewes. Chromosomal placement of the
quantified miRNAs revealed a large cluster of miRNAs on chromosome 9 that we also observed in
the ovaries (Fig. 7).

4.10 Limitations and thoughts for future studies

We acknowledge certain limitations of this study. We believe that with the availability of a better
annotated reference genome, the data from this study will reveal additional information that we may
have missed in this paper. With sequencing costs becoming increasingly inexpensive, increasing the
sample size of each breed group would certainly add statistical power. In particular, given that time-
series experiments are not feasible with the same animal, sampling could be performed with a larger
group of animals at different stages of pregnancy to obtain an overview of gene expression changes.
It should be noted that overall gene expression and, more specifically, differential expression
between breeds is inherently a stochastic process; thus, there is always some level of bias caused by
individual variation (Hansen et al., 2011). Including more individuals in future experiments will
minimize such bias. The results from breeding experiments have shown that productivity traits such
as litter sizes may not carry on to F2 crosses (F1 x F1) and/or backcrosses. Therefore, future
experiments that involve F2 crosses and backcrosses would provide more valuable findings related to
prolificacy. In addition, replicating such experiments in different environments would be even more
exciting. To minimize or alleviate noise from tissue heterogeneity, single-cell experiments may prove
beneficial in future studies. While we observed interplay between the endometrium and the CL, it
would be equally interesting to examine the overall impacts of these tissues on the embryo. Finally,
the application of gene-modifying technologies such as CRISPR/Cas9 to edit certain regions (such as
the region in the FecL locus homologous to the partial retrovirus sequence) may provide important
insights into phenotypes associated with infertility, prolificacy and other traits of interest.

5   Conclusion

We compiled the most comprehensive list thus far of genes (n=21,287) and miRNAs (n=531)
expressed in the CL and endometrium, which are the most important tissues during the
preimplantation stage and therefore determine the success of pregnancy in sheep. Our results agree
well with the (limited) existing reports, which are mainly focused on the interplay of the
endometrium and conceptus, but we have shown that the CL plays an equally important role. The
relative scarcity of genetic information about the CL means that its functional importance is
underrated. We identified several actors, including coding genes (producing mRNA) and noncoding
genes (miRNAs, snoRNAs, and lincRNAs), that are essential during early pregnancy. A total of
21,287 gene transcripts and 531 miRNAs were expressed in the dataset. Functional analysis primarily
based on literature searches and earlier studies revealed the significant roles of the most highly
expressed genes in pregnancy recognition, implantation and placentation. F1 crosses were more
closely related to Finnsheep than to Texel, as indicated by phenotypic and gene expression results
that need to be validated with additional experiments (with F2 crosses and backcrosses). Several
genes with potential importance during early pregnancy (including *SIGLEC13*, *SIGLEC14*,
SIGLEC6, *MRP4*, and *CA5A*) were reported in the CL for the first time in any species. The roles of
retroviruses during early pregnancy and in breed-specific phenotypes were clearly indicated by the
observed gene expression dynamics, especially in the endometrium. A novel gene upregulated in
Finnsheep and sharing similarity with an ERV was identified in the FecL locus. The results from this
study show the importance of the immune system during early pregnancy. We also highlight the need
for improved annotation of the sheep genome and emphasize that our data will certainly contribute to
such improvement. We observed a cluster of miRNAs on chromosome 18 homologous to that found
on chromosome 14 in humans. Taken together, our data provide new information to aid in
understanding the complex reproductive events during the preimplantation period in sheep and may
also have implications for other ruminants (such as goats and cattle) and mammals, including
humans.

6   Abbreviations

Ig (immunoglobulin), Siglec (sialic acid-binding Ig-like lectin), ERV (endogenous retrovirus), CDS
(coding sequence), OXT (oxytocin), MRP (multidrug resistance-associated protein), CL (corpus
luteum), lincRNA (long intergenic noncoding RNA)

7   Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial
relationships that could be construed as potential conflicts of interest.

8   Author Contributions

J.K. and M.H.L. conceived and designed the project. J.K., M.H. and J.P. collected the samples. K.P.
analyzed the data and wrote the manuscript. J.P. contributed substantially to revising the manuscript.
M.W. and J.K. contributed to the data analysis and manuscript writing, respectively. All authors
revised and approved the final manuscript.

8.1   Funding

This study was funded by the Academy of Finland (decisions 250633, 250677 and 285017). This
study is part of the ClimGen ("Climate Genomics for Farm Animal Adaptation") project funded by

FACCE-JPI ERA-NET Plus on Climate Smart Agriculture. K.P. acknowledges financial support
from the Niemi Foundation.

9 Acknowledgments

We are grateful to Anu Tuomola for providing the experimental facilities on her farm in Urjala and to
Kati Kaisaioki from Lallin Lammas Ltd. for providing the slaughtering facilities. We thank Johanna
Rautiainen, Arja Seppälä, Annukka Numminen, Kalle Saastamoinen, Ilma Tapio, Tuula Marjatta
Hamama, Anneli Virta and Magnus Andersson for their valuable assistance during this study. The
authors wish to acknowledge the CSC – IT Center for Science, Finland, for computational resources.
This study was supported by the Finnish Functional Genomics Centre of the University of Turku,
Åbo Akademi and Biocenter Finland.

10 References

Angata, T., Margulies, E. H., Green, E. D., and Varki, A. (2004). Large-scale sequencing of the
CD33-related Siglec gene cluster in five mammalian species reveals rapid evolution by multiple
mechanisms. *Proc. Natl. Acad. Sci.* 101, 13251–13256. doi:10.1073/pnas.0404833101.

Bastian, A., Kratzin, H., Fallgren-Gebauer, E., Eckart, K., and Hilschmann, N. (1995). "Intra- and
Inter-Chain Disulfide Bridges of J Chain in Human S-IgA," in (Springer, Boston, MA), 581–
583. doi:10.1007/978-1-4615-1941-6_122.

Bazer, F. W. (2013). Pregnancy recognition signaling mechanisms in ruminants and pigs. *J. Anim.
Sci. Biotechnol.* doi:10.1186/2049-1891-4-23.

Bindea, G., Mlecnik, B., Hackl, H., Charoentong, P., Tosolini, M., Kirilovsky, A., et al. (2009).
ClueGO: a Cytoscape plug-in to decipher functionally grouped gene ontology and pathway
annotation networks. *Bioinformatics* 25, 1091–3. doi:10.1093/bioinformatics/btp101.

Bishop, C. V. (2013). Progesterone inhibition of oxytocin signaling in endometrium. *Front.
Neurosci.* doi:10.3389/fnins.2013.00138.

Bolet, G. (1986). "Timing and Extent of Embryonic Mortality in Pigs Sheep and Goats: Genetic
Variability," in *Embryonic Mortality in Farm Animals* (Dordrecht: Springer Netherlands), 12–
43. doi:10.1007/978-94-009-5038-2_2.

Bornhöfft, K. F., Goldammer, T., Rebl, A., and Galuska, S. P. (2018). Siglecs: A journey through the
evolution of sialic acid-binding immunoglobulin-type lectins. *Dev. Comp. Immunol.* 86, 219–
231. doi:10.1016/J.DCI.2018.05.008.

Branco, R., and Masle, J. (2019). Systemic signalling through TCTP1 controls lateral root formation
in Arabidopsis. *J. Exp. Bot.* doi:10.1093/jxb/erz204.

Brioudes, F., Thierry, A.-M., Chambrier, P., Mollereau, B., and Bendahmane, M. (2010).
Translationally controlled tumor protein is a conserved mitotic growth integrator in animals and
plants. *Proc. Natl. Acad. Sci.* 107, 16384–16389. doi:10.1073/pnas.1007926107.

Brooks, K., Burns, G. W., Moraes, J. G. N., and Spencer, T. E. (2016). Analysis of the Uterine
Epithelial and Conceptus Transcriptome and Luminal Fluid Proteome During the Peri-

Implantation Period of Pregnancy in Sheep. *Biol. Reprod.* 95, 88–88.
doi:10.1095/biolreprod.116.141945.

Cao, H., and Crocker, P. R. (2011). Evolution of CD33-related siglecs: regulating host immune
functions and escaping pathogen exploitation? *Immunology* 132, 18–26. doi:10.1111/j.1365-
2567.2010.03368.x.

Casey, O. M., Morris, D. G., Powell, R., Sreenan, J. M., and Fitzpatrick, R. (2005). Analysis of gene
expression in non-regressed and regressed bovine corpus luteum tissue using a customized
ovarian cDNA array. *Theriogenology* 64, 1963–1976.
doi:10.1016/j.theriogenology.2005.04.015.

Chen, S. H., Wu, P.-S., Chou, C.-H., Yan, Y.-T., Liu, H., Weng, S.-Y., et al. (2007). A Knockout
Mouse Approach Reveals that TCTP Functions as an Essential Factor for Cell Proliferation and
Survival in a Tissue- or Cell Type–specific Manner. *Mol. Biol. Cell* 18, 2525–2532.
doi:10.1091/mbc.e07-02-0188.

Christenson, L. K., and Devoto, L. (2003). Cholesterol transport and steroidogenesis by the corpus
luteum. *Reprod. Biol. Endocrinol.* 1, 90. doi:10.1186/1477-7827-1-90.

Chu, M. X., Mu, Y. L., Fang, L., Ye, S. C., and Sun, S. H. (2007). Prolactin receptor as a candidate
gene for prolificacy of small tail Han sheep. *Anim. Biotechnol.* 18, 65–73.
doi:10.1080/10495390601090950.

Coulam, C. B., and Goodman, C. (2000). Increased pregnancy rates after IVF/ET with intravenous
immunoglobulin treatment in women with elevated circulating C56+ cells. *Early Pregnancy* 4,
90–8. Available at: http://www.ncbi.nlm.nih.gov/pubmed/11723539 [Accessed May 31, 2019].

Davis, G. H., Farquhar, P. A., O'Connell, A. R., Everett-Hincks, J. M., Wishart, P. J., Galloway, S.
M., et al. (2006). A putative autosomal gene increasing ovulation rate in Romney sheep. *Anim.
Reprod. Sci.* 92, 65–73. doi:10.1016/j.anireprosci.2005.05.015.

Davis, G. H., Galloway, S. M., Ross, I. K., Gregan, S. M., Ward, J., Nimbkar, B. V, et al. (2002).
DNA tests in prolific sheep from eight countries provide new evidence on origin of the
Booroola (FecB) mutation. *Biol. Reprod.* 66, 1869–74. Available at:
http://www.ncbi.nlm.nih.gov/pubmed/12021074 [Accessed January 25, 2013].

Davis, J. S., and LaVoie, H. A. (2018). *Molecular Regulation of Progesterone Production in the
Corpus Luteum*. 3rd ed. Elsevier Inc. doi:10.1016/b978-0-12-813209-8.00015-7.

De Placido, G., Zullo, F., Mollo, A., Cappiello, F., Nazzaro, A., Colacurci, N., et al. (1994).
Intravenous immunoglobulin (IVIG) in the prevention of implantation failures. *Ann. N. Y. Acad.
Sci.* 734, 232–4. Available at: http://www.ncbi.nlm.nih.gov/pubmed/7978921 [Accessed May
31, 2019].

DeMartini, J. C., Carlson, J. O., Leroux, C., Spencer, T., and Palmarini, M. (2003). Endogenous
retroviruses related to jaagsiekte sheep retrovirus. *Curr. Top. Microbiol. Immunol.* 275, 117–37.
Available at: http://www.ncbi.nlm.nih.gov/pubmed/12596897 [Accessed June 3, 2019].

Deniz, E., and Erman, B. (2017). Long noncoding RNA (lincRNA), a new paradigm in gene

expression control. *Funct. Integr. Genomics* 17, 135–143. doi:10.1007/s10142-016-0524-x.

Denner, J. (2016). Expression and function of endogenous retroviruses in the placenta. *APMIS* 124,
31–43. doi:10.1111/apm.12474.

Devoto, L., Kohen, P., Gonzalez, R. R., Castro, O., Retamales, I., Vega, M., et al. (2001). Expression
of Steroidogenic Acute Regulatory Protein in the Human Corpus Luteum throughout the Luteal
Phase. *J. Clin. Endocrinol. Metab.* 86, 5633–5639. doi:10.1210/jcem.86.11.7982.

Drouilhet, L., Mansanet, C., Sarry, J., Tabet, K., Bardou, P., Woloszyn, F., et al. (2013). The Highly
Prolific Phenotype of Lacaune Sheep Is Associated with an Ectopic Expression of the
B4GALNT2 Gene within the Ovary. *PLoS Genet.* 9, e1003809.
doi:10.1371/journal.pgen.1003809.

Dunlap, K. A., Palmarini, M., Adelson, D. L., and Spencer, T. E. (2005). Sheep Endogenous
Betaretroviruses (enJSRVs) and the Hyaluronidase 2 (HYAL2) Receptor in the Ovine Uterus
and Conceptus. *Biol. Reprod.* 73, 271–279. doi:10.1095/biolreprod.105.039776.

Dunlap, K. A., Palmarini, M., and Spencer, T. E. (2006a). Ovine Endogenous Betaretroviruses
(enJSRVs) and Placental Morphogenesis. *Placenta* 27, 135–140.
doi:10.1016/j.placenta.2005.12.009.

Dunlap, K. A., Palmarini, M., Varela, M., Burghardt, R. C., Hayashi, K., Farmer, J. L., et al. (2006b).
Endogenous retroviruses regulate periimplantation placental growth and differentiation. *Proc.*
Natl. Acad. Sci. U. S. A. 103, 14390–5. doi:10.1073/pnas.0603836103.

Durinck, S., Moreau, Y., Kasprzyk, A., Davis, S., De Moor, B., Brazma, A., et al. (2005). BioMart
and Bioconductor: a powerful link between biological databases and microarray data analysis.
Bioinformatics 21, 3439–40. doi:10.1093/bioinformatics/bti525.

Felix Krueger Trim Galore. Available at:
http://www.bioinformatics.babraham.ac.uk/projects/trim_galore/ [Accessed January 9, 2019].

Fleming, J. A. G. W., Spencer, T. E., Safe, S. H., and Bazer, F. W. (2006). Estrogen regulates
transcription of the ovine oxytocin receptor gene through GC-rich SP1 promoter elements.
Endocrinology. doi:10.1210/en.2005-1120.

Flint, A. P. F., and Sheldrick, E. L. (2004). Ovarian oxytocin and the maternal recognition of
pregnancy. *Reproduction*. doi:10.1530/jrf.0.0760831.

Forde, N., Mehta, J. P., McGettigan, P. A., Mamo, S., Bazer, F. W., Spencer, T. E., et al. (2013).
Alterations in expression of endometrial genes coding for proteins secreted into the uterine
lumen during conceptus elongation in cattle. *BMC Genomics* 14, 321. doi:10.1186/1471-2164-
14-321.

Friedländer, M. R., Mackowiak, S. D., Li, N., Chen, W., and Rajewsky, N. (2012). miRDeep2
accurately identifies known and hundreds of novel microRNA genes in seven animal clades.
Nucleic Acids Res. 40, 37–52. doi:10.1093/nar/gkr688.

Geisert, R. D., Morgan, G. L., Short, E. C., and Zavy, M. T. (1992). Endocrine events associated with

endometrial function and conceptus development in cattle. *Reprod. Fertil. Dev.* 4, 301–5.
Available at: http://www.ncbi.nlm.nih.gov/pubmed/1438962 [Accessed April 3, 2019].

Gimpl, G., Wiegand, V., Burger, K., and Fahrenholz, F. (2002). Cholesterol and steroid hormones:
Modulators of oxytocin receptor function. in *Progress in Brain Research* doi:10.1016/S0079-
6123(02)39006-X.

Gray, C. A., Abbey, C. A., Beremand, P. D., Choi, Y., Farmer, J. L., Adelson, D. L., et al. (2006).
Identification of Endometrial Genes Regulated by Early Pregnancy, Progesterone, and
Interferon Tau in the Ovine Uterus. *Biol. Reprod.* 74, 383–394.
doi:10.1095/biolreprod.105.046656.

Gray, C. A., Adelson, D. L., Bazer, F. W., Burghardt, R. C., Meeusen, E. N. T., and Spencer, T. E.
(2004). Discovery and characterization of an epithelial-specific galectin in the endometrium that
forms crystals in the trophectoderm. *Proc. Natl. Acad. Sci.* 101, 7982–7987.
doi:10.1073/pnas.0402669101.

Grazzini, E., Guillon, G., Mouillac, B., and Zingg, H. H. (1998). Inhibition of oxytocin receptor
function by direct binding of progesterone. *Nature*. doi:10.1038/33176.

Hanrahan, J. P., Gregan, S. M., Mulsant, P., Mullen, M., Davis, G. H., Powell, R., et al. (2004).
Mutations in the genes for oocyte-derived growth factors GDF9 and BMP15 are associated with
both increased ovulation rate and sterility in Cambridge and Belclare sheep (Ovis aries). *Biol.*
Reprod. 70, 900–9. doi:10.1095/biolreprod.103.023093.

Hanrahan, J. P., and Quirke, J. F. (1984). Contribution of variation in ovulation rate and embryo
survival to within breed variation in litter size. *Genet. Reprod. sheep / Ed. by R.B. Land, D.W.*
Robinson. Available at: http://agris.fao.org/agris-search/search.do?recordID=US201301472510
[Accessed April 2, 2019].

Hansen, K. D., Wu, Z., Irizarry, R. A., and Leek, J. T. (2011). Sequencing technology does not
eliminate biological variability. *Nat. Biotechnol.* 29, 572–573. doi:10.1038/nbt.1910.

Hernández-Montiel, W., Collí-Dula, R. C., Ramón-Ugalde, J. P., Martínez-Núñez, M. A., and
Zamora-Bustillos, R. (2019). RNA-seq Transcriptome Analysis in Ovarian Tissue of Pelibuey
Breed to Explore the Regulation of Prolificacy. *Genes (Basel).* 10, 358.
doi:10.3390/genes10050358.

Honjo, T. (1983). Immunoglobulin Genes. *Annu. Rev. Immunol.* 1, 499–528.
doi:10.1146/annurev.iy.01.040183.002435.

Hu, J., Zhang, Z., Shen, W.-J., and Azhar, S. (2010). Cellular cholesterol delivery, intracellular
processing and utilization for biosynthesis of steroid hormones. *Nutr. Metab. (Lond).* 7, 47.
doi:10.1186/1743-7075-7-47.

Hu, X., Pokharel, K., Peippo, J., Ghanem, N., Zhaboyev, I., Kantanen, J., et al. (2015). Identification
and characterization of miRNAs in the ovaries of a highly prolific sheep breed. *Anim. Genet.*,
n/a-n/a. doi:10.1111/age.12385.

Juengel, J. L., Meberg, B. M., Turzillo, A. M., Nett, T. M., and Niswender, G. D. (1995). Hormonal

regulation of messenger ribonucleic acid encoding steroidogenic acute regulatory protein in
ovine corpora lutea. *Endocrinology* 136, 5423–5429. doi:10.1210/endo.136.12.7588291.

Kendrick, K. M. (2000). Oxytocin, motherhood and bonding. *Exp. Physiol.* 85, 111s-124s.
doi:10.1111/j.1469-445x.2000.tb00014.x.

Kfir, S., Basavaraja, R., Wigoda, N., Ben-Dor, S., Orr, I., and Meidan, R. (2018). Genomic profiling
of bovine corpus luteum maturation. *PLoS One* 13, e0194456.
doi:10.1371/journal.pone.0194456.

Kim, S., Choi, Y., Bazer, F. W., and Spencer, T. E. (2003). Identification of Genes in the Ovine
Endometrium Regulated by Interferon τ Independent of Signal Transducer and Activator of
Transcription. *Endocrinology* 144, 5203–5214. doi:10.1210/en.2003-0665.

King, S. R., and LaVoie, H. A. (2009). "Regulation of the Early Steps in Gonadal Steroidogenesis,"
in *Reproductive Endocrinology* (Boston, MA: Springer US), 175–193. doi:10.1007/978-0-387-
88186-7_16.

Koch, J. M., Ramadoss, J., and Magness, R. R. (2010). Proteomic profile of uterine luminal fluid
from early pregnant ewes. *J. Proteome Res.* 9, 3878–3885. doi:10.1021/pr100096b.

Kozomara, A., Birgaoanu, M., and Griffiths-Jones, S. (2019). miRBase: from microRNA sequences
to function. *Nucleic Acids Res.* 47, D155–D162. doi:10.1093/nar/gky1141.

Kozomara, A., and Griffiths-Jones, S. (2011). miRBase: integrating microRNA annotation and deep-
sequencing data. *Nucleic Acids Res.* 39, D152–D157. doi:10.1093/nar/gkq1027.

Kozomara, A., and Griffiths-Jones, S. (2014). miRBase: annotating high confidence microRNAs
using deep sequencing data. *Nucleic Acids Res.* 42, D68–D73. doi:10.1093/nar/gkt1181.

Lacroix-Pépin, N., Danyod, G., Krishnaswamy, N., Mondal, S., Rong, P.-M., Chapdelaine, P., et al.
(2011). The Multidrug Resistance-Associated Protein 4 (MRP4) Appears as a Functional Carrier
of Prostaglandins Regulated by Oxytocin in the Bovine Endometrium. *Endocrinology* 152,
4993–5004. doi:10.1210/en.2011-1406.

Langmead, B., Trapnell, C., Pop, M., and Salzberg, S. L. (2009). Ultrafast and memory-efficient
alignment of short DNA sequences to the human genome. *Genome Biol.* 10, R25.
doi:10.1186/gb-2009-10-3-r25.

Lewis, S. K., Farmer, J. L., Burghardt, R. C., Newton, G. R., Johnson, G. A., Adelson, D. L., et al.
(2007). Galectin 15 (LGALS15): A Gene Uniquely Expressed in the Uteri of Sheep and Goats
that Functions in Trophoblast Attachment1. *Biol. Reprod.* 77, 1027–1036.
doi:10.1095/biolreprod.107.063594.

Li, S., Chen, X., Ding, Y., Liu, X., Wang, Y., and He, J. (2011). Expression of translationally
controlled tumor protein (TCTP) in the uterus of mice of early pregnancy and its possible
significance during embryo implantation. *Hum. Reprod.* 26, 2972–2980.
doi:10.1093/humrep/der275.

Loeser, R. F., and Wallin, R. (1992). Cell adhesion to matrix Gla protein and its inhibition by an Arg-

Gly-Asp-containing peptide. *J. Biol. Chem.* 267, 9459–62. Available at:
http://www.ncbi.nlm.nih.gov/pubmed/1577789 [Accessed May 21, 2019].

Love, M., Anders, S., and Huber, W. (2013). Differential expression of RNA-Seq data at the gene
level – the DESeq2 package.

Madeira, F., Park, Y., M., Lee, J., Buso, N., Gur, T., Madhusoodanan, N., et al. (2019). The EMBL-
EBI search and sequence analysis tools APIs in 2019. *Nucleic Acids Res.*, 2–4.

Mamo, S., Mehta, J. P., Forde, N., McGettigan, P., and Lonergan, P. (2012). Conceptus-
Endometrium Crosstalk During Maternal Recognition of Pregnancy in Cattle. *Biol. Reprod.* 87,
6, 1–9. doi:10.1095/biolreprod.112.099945.

Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet.journal 17, 10–12. Available at:
http://journal.embnet.org/index.php/embnetjournal/article/view/200/479 [Accessed November
25, 2013].

Mateyak, M. K., and Kinzy, T. G. (2010). eEF1A: thinking outside the ribosome. *J. Biol. Chem.* 285,
21209–13. doi:10.1074/jbc.R110.113795.

Moore, S. G., Pryce, J. E., Hayes, B. J., Chamberlain, A. J., Kemper, K. E., Berry, D. P., et al.
(2016). Differentially Expressed Genes in Endometrium and Corpus Luteum of Holstein Cows
Selected for High and Low Fertility Are Enriched for Sequence Variants Associated with
Fertility1. *Biol. Reprod.* 94. doi:10.1095/biolreprod.115.132951.

Mullen, M. P., and Hanrahan, J. P. (2014). Direct evidence on the contribution of a missense
mutation in GDF9 to variation in ovulation rate of Finnsheep. *PLoS One* 9, e95251.
doi:10.1371/journal.pone.0095251.

Mulsant, P., Lecerf, F., Fabre, S., Schibler, L., Monget, P., Lanneluc, I., et al. (2001). Mutation in
bone morphogenetic protein receptor-IB is associated with increased ovulation rate in Booroola
Mérino ewes. *Proc. Natl. Acad. Sci. U. S. A.* 98, 5104–9. doi:10.1073/pnas.091577598.

Palmarini, M., Gray, C. A., Carpenter, K., Fan, H., Bazer, F. W., and Spencer, T. E. (2001).
Expression of Endogenous Betaretroviruses in the Ovine Uterus: Effects of Neonatal Age,
Estrous Cycle, Pregnancy, and Progesterone. *J. Virol.* 75, 11319–11327.
doi:10.1128/JVI.75.23.11319-11327.2001.

Pantano, L., Estivill, X., and Martí, E. (2011). A non-biased framework for the annotation and
classification of the non-miRNA small RNA transcriptome. *Bioinformatics* 27, 3202–3203.
doi:10.1093/bioinformatics/btr527.

Patro, R., Duggal, G., Love, M. I., Irizarry, R. A., and Kingsford, C. (2017). Salmon provides fast
and bias-aware quantification of transcript expression. *Nat. Methods* 14, 417–419.
doi:10.1038/nmeth.4197.

Pauli, A., Rinn, J. L., and Schier, A. F. (2011). Non-coding RNAs as regulators of embryogenesis.
Nat. Rev. Genet. 12, 136–49. doi:10.1038/nrg2904.

Pillai, S., Netravali, I. A., Cariappa, A., and Mattoo, H. (2012). Siglecs and immune regulation.
Annu. Rev. Immunol. 30, 357–92. doi:10.1146/annurev-immunol-020711-075018.

Plant, T. M., Zeleznik, A. J., and Auchus, R. J. (2015). "Chapter 8 – Human Steroid Biosynthesis," in
Knobil and Neill's Physiology of Reproduction, 295–312. doi:10.1016/B978-0-12-397175-
3.00008-9.

Pokharel, K., Peippo, J., Honkatukia, M., Seppälä, A., Rautiainen, J., Ghanem, N., et al. (2018).
Integrated ovarian mRNA and miRNA transcriptome profiling characterizes the genetic basis of
prolificacy traits in sheep (Ovis aries). *BMC Genomics* 19. doi:10.1186/s12864-017-4400-4.

Pokharel, K., Weldenegodguad, M., Popov, R., Honkatukia, M., Huuki, H., Lindeberg, H., et al.
(2019). Whole blood transcriptome analysis reveals footprints of cattle adaptation to sub-arctic
conditions. *Anim. Genet.* 50, 217–227. doi:10.1111/age.12783.

Quinlivan, T. D., Martin, C. A., Taylor, W. B., and Cairney, I. M. (1966). Estimates of pre- and
perinatal mortality in the New Zealand Romney Marsh ewe. I. Pre- and perinatal mortality in
those ewes that conceived to one service. *J. Reprod. Fertil.* 11, 379–90. Available at:
http://www.ncbi.nlm.nih.gov/pubmed/5940535 [Accessed May 31, 2019].

Ransohoff, J. D., Wei, Y., and Khavari, P. A. (2018). The functions and unique features of long
intergenic non-coding RNA. *Nat. Rev. Mol. Cell Biol.* 19, 143–157. doi:10.1038/nrm.2017.104.

Ricordeau, G., Thimonier, J., Poivey, J. P., Driancourt, M. A., Hochereau-De-Reviers, M. T., and
Tchamitchian, L. (1990). I.N.R.A. research on the Romanov sheep breed in France: A review.
Livest. Prod. Sci. 24, 305–332. doi:10.1016/0301-6226(90)90009-U.

Romero, J. J., Liebig, B. E., Broeckling, C. D., Prenni, J. E., and Hansen, T. R. (2017). Pregnancy-
induced changes in metabolome and proteome in ovine uterine flushings. *Biol. Reprod.* 97, 273–
287. doi:10.1093/biolre/iox078.

Satterfield, M. C., Bazer, F. W., and Spencer, T. E. (2006). Progesterone Regulation of
Preimplantation Conceptus Growth and Galectin 15 (LGALS15) in the Ovine Uterus. *Biol.*
Reprod. 75, 289–296. doi:10.1095/biolreprod.106.052944.

Shannon, P., Markiel, A., Ozier, O., Baliga, N. S., Wang, J. T., Ramage, D., et al. (2003). Cytoscape:
a software environment for integrated models of biomolecular interaction networks. *Genome*
Res. 13, 2498–504. doi:10.1101/gr.1239303.

Simon Andrews FastQC A Quality Control tool for High Throughput Sequence Data. Available at:
http://www.bioinformatics.babraham.ac.uk/projects/fastqc/ [Accessed January 18, 2013].

Sistiaga-Poveda, M., and Jugo, B. M. (2014). Evolutionary dynamics of endogenous Jaagsiekte sheep
retroviruses proliferation in the domestic sheep, mouflon and Pyrenean chamois. *Heredity*
(Edinb). 112, 571–8. doi:10.1038/hdy.2013.136.

Souza, C. J., MacDougall, C., MacDougall, C., Campbell, B. K., McNeilly, A. S., and Baird, D. T.
(2001). The Booroola (FecB) phenotype is associated with a mutation in the bone
morphogenetic receptor type 1 B (BMPR1B) gene. *J. Endocrinol.* 169, R1-6. Available at:
http://www.ncbi.nlm.nih.gov/pubmed/11312159 [Accessed April 2, 2019].

Spencer, T. ., Burghardt, R. ., Johnson, G. ., and Bazer, F. . (2004a). Conceptus signals for
establishment and maintenance of pregnancy. *Anim. Reprod. Sci.* 82–83, 537–550.
doi:10.1016/j.anireprosci.2004.04.014.

Spencer, T. E., and Bazer, F. W. (2004). Conceptus signals for establishment and maintenance of
pregnancy. *Reprod. Biol. Endocrinol.* 2, 49. doi:10.1186/1477-7827-2-49.

Spencer, T. E., Forde, N., and Lonergan, P. (2015). The role of progesterone and conceptus-derived
factors in uterine biology during early pregnancy in ruminants. *J. Dairy Sci.* 99, 5941–5950.
doi:10.3168/jds.2015-10070.

Spencer, T. E., Johnson, G. A., Bazer, F. W., and Burghardt, R. C. (2004b). Implantation
mechanisms: insights from the sheep. *Reproduction* 128, 657–68. doi:10.1530/rep.1.00398.

Spencer, T. E., Johnson, G. A., Bazer, F. W., Burghardt, R. C., and Palmarini, M. (2007). Pregnancy
recognition and conceptus implantation in domestic ruminants: roles of progesterone,
interferons and endogenous retroviruses. *Reprod. Fertil. Dev.* 19, 65–78. Available at:
http://www.ncbi.nlm.nih.gov/pubmed/17389136 [Accessed April 3, 2019].

Spencer, T. E., and Palmarini, M. (2012). Endogenous retroviruses of sheep: a model system for
understanding physiological adaptation to an evolving ruminant genome. *J. Reprod. Dev.* 58,
33–7. Available at: http://www.ncbi.nlm.nih.gov/pubmed/22450282 [Accessed June 3, 2019].

Spencer, T. E., Sandra, O., and Wolf, E. (2008). Genes involved in conceptus–endometrial
interactions in ruminants: insights from reductionism and thoughts on holistic approaches.
REPRODUCTION 135, 165–179. doi:10.1530/REP-07-0327.

Spencer, T. E., Stagg, A. G., Joyce, M. M., Jenster, G., Wood, C. G., Bazer, F. W., et al. (1999).
Discovery and Characterization of Endometrial Epithelial Messenger Ribonucleic Acids Using
the Ovine Uterine Gland Knockout Model [1]. *Endocrinology* 140, 4070–4080.
doi:10.1210/endo.140.9.6981.

Stocco, D. M. (2000). The role of the StAR protein in steroidogenesis: challenges for the future. *J.*
Endocrinol. 164, 247–53. Available at: http://www.ncbi.nlm.nih.gov/pubmed/10694364
[Accessed May 2, 2019].

Stouffer, R. L., and Hennebold, J. D. (2015). "Structure, Function, and Regulation of the Corpus
Luteum," in *Knobil and Neill's Physiology of Reproduction*, 1023–1076. doi:10.1016/B978-0-
12-397175-3.00023-5.

Tatsuka, M., Mitsui, H., Wada, M., Nagata, A., Nojima, H., and Okayama, H. (1992). Elongation
factor-1α gene determines susceptibility to transformation. *Nature* 359, 333–336.
doi:10.1038/359333a0.

Trapnell, C., Williams, B. A., Pertea, G., Mortazavi, A., Kwan, G., van Baren, M. J., et al. (2010).
Transcript assembly and quantification by RNA-Seq reveals unannotated transcripts and isoform
switching during cell differentiation. *Nat. Biotechnol.* 28, 511–5. doi:10.1038/nbt.1621.

Tuynder, M., Susini, L., Prieur, S., Besse, S., Fiucci, G., Amson, R., et al. (2002). Biological models
and genes of tumor reversion: cellular reprogramming through tpt1/TCTP and SIAH-1. *Proc.*

Natl. Acad. Sci. U. S. A. 99, 14976–81. doi:10.1073/pnas.222470799.

Ulitsky, I., and Bartel, D. P. (2013). lincRNAs: genomics, evolution, and mechanisms. *Cell* 154, 26–
46. doi:10.1016/j.cell.2013.06.020.

Våge, D. I., Husdal, M., Kent, M. P., Klemetsdal, G., and Boman, I. a (2013). A missense mutation
in growth differentiation factor 9 (GDF9) is strongly associated with litter size in sheep. *BMC*
Genet. 14, 1. doi:10.1186/1471-2156-14-1.

Varki, A., and Angata, T. (2006). Siglecs—the major subfamily of I-type lectins. *Glycobiology* 16,
1R-27R. doi:10.1093/glycob/cwj008.

Waterhouse, A. M., Procter, J. B., Martin, D. M. A., Clamp, M., and Barton, G. J. (2009). Jalview
Version 2--a multiple sequence alignment editor and analysis workbench. *Bioinformatics* 25,
1189–1191. doi:10.1093/bioinformatics/btp033.

Williams, A. F., and Barclay, A. N. (1988). The Immunoglobulin Superfamily—Domains for Cell
Surface Recognition. *Annu. Rev. Immunol.* 6, 381–405.
doi:10.1146/annurev.iy.06.040188.002121.

Xu, S.-S., Gao, L., Xie, X.-L., Ren, Y.-L., Shen, Z.-Q., Wang, F., et al. (2018). Genome-Wide
Association Analyses Highlight the Potential for Different Genetic Mechanisms for Litter Size
Among Sheep Breeds. *Front. Genet.* 9, 118. doi:10.3389/fgene.2018.00118.

Yang, J., Li, X., Cao, Y.-H., Pokharel, K., Hu, X.-J., Chen, Z.-H., et al. (2018). Comparative mRNA
and miRNA expression in European mouflon (Ovis musimon) and sheep (Ovis aries) provides
novel insights into the genetic mechanisms for female reproductive success. *Heredity (Edinb).*
doi:10.1038/s41437-018-0090-1.

Zhao, J., and Nishimoto, S. K. (1996). Matrix Gla protein gene expression is elevated during
postnatal development. *Matrix Biol.* 15, 131–40. Available at:
http://www.ncbi.nlm.nih.gov/pubmed/8837014 [Accessed May 21, 2019].

Zhao, J., and Warburton, D. (1997). Matrix Gla protein gene expression is induced by transforming
growth factor-beta in embryonic lung culture. *Am. J. Physiol. Cell. Mol. Physiol.* 273, L282–
L287. doi:10.1152/ajplung.1997.273.1.L282.

11   FIGURE LEGENDS

Figure 1 Venn diagram showing the distribution of genes expressed in the A) CL and B)
endometrium of Finnsheep, Texel and F1 crosses.

Figure 2 Sample relatedness. (A) PCA plot of the top 500 expressed genes in the CL (left) and endometrium (right) and (B) heatmap of the top 25 most variable genes across all samples. Tissue-specific samples are denoted with a trailing c (for CL) or e (for endometrium). Legend: FS – Finnsheep; TX – Texel ewes, F1 – F1 crosses of Finnsheep and Texel sheep

Figure 3 GO terms associated with the list of genes that were upregulated in the CLs of Finnsheep
compared to Texel ewes

Figure 4 Multiple sequence alignment of the novel endogenous retrovirus gene. We generated the
reverse complements of ENSOARG00000009959 and endogenous-virus beta-2 pro/pol and aligned
them with a partial sequence of the reduced FecL locus using Clustal Omega (Madeira et al., 2019).
The bases are colored based on the nucleotide coloring scheme in Jalview (Waterhouse et al., 2009).

Figure 5 miRNA clusters in sheep (Chr. 18, top) and humans (Chr. 14, bottom).

12 TABLES

Table 1: Top 25 genes (ranked by TPM) exclusively expressed in CL and endometrium.

	TPM	Gene name	Chr.	Gene description
CL				
ENSOARG00000003867	2080.2	CYP11A1	18	cytochrome P450, family 11, subfamily A, polypeptide 1
ENSOARG00000009107	1233.5	CCL21	2	C-C motif chemokine ligand 21
ENSOARG00000013402	1004.3	PTGFR	1	prostaglandin F receptor
ENSOARG00000013340	622.3	CCL26	24	C-C motif chemokine ligand 26
ENSOARG00000019663	424.4	PTHLH	3	parathyroid hormone like hormone
ENSOARG00000015144	273.6	SERPINA5	18	serpin family A member 5
ENSOARG00000004774	248.0	LOC101114790*	11	C-C motif chemokine 15
ENSOARG00000014882	225.6	SERPINA1	18	serpin family A member 1
ENSOARG00000009230	201.1	DPT	12	dermatopontin
ENSOARG00000016052	188.9	AOX1	2	aldehyde oxidase 1
ENSOARG00000008189	177.9	PKIB	8	cAMP-dependent protein kinase inhibitor beta
ENSOARG00000004455	177.6	LHCGR	3	luteinizing hormone/choriogonadotropin receptor
ENSOARG00000020188	160.0	FAM110B*	9	Ovis aries family with sequence similarity 110 member B
ENSOARG00000009597	151.2	HS6ST2*	X	heparan sulfate 6-O-sulfotransferase 2
ENSOARG00000000474	143.3	GJA4	1	gap junction protein alpha 4
ENSOARG00000002475	131.9	PEG10	4	paternally expressed 10
ENSOARG00000010344	130.1	LTBP1	3	latent transforming growth factor beta binding protein 1
ENSOARG00000020976	118.9	GLDN	7	gliomedin
ENSOARG00000014966	116.3	INSL3	5	insulin like 3

ENSOARG00000009887	111.7	CHST15	22	carbohydrate sulfotransferase 15
ENSOARG00000017273	108.7	PROS1	1	protein S
ENSOARG00000020243	104.9	INHA	2	Ovis aries inhibin subunit alpha (INHA), mRNA.
ENSOARG00000019971	99.6	PLN	8	phospholamban
ENSOARG00000005118	92.8	KCNK12	3	potassium two pore domain channel subfamily K member
ENSOARG00000016448	90.1	TFR2	24	transferrin receptor 2
Endometrium				
ENSOARG00000004013	2378.2	FXYD4	25	FXYD domain containing ion transport regulator 4
ENSOARG00000012900	1429.8	IGFBP1	4	insulin like growth factor binding protein 1
ENSOARG00000002255	1191.2		19	acyl-coenzyme A thioesterase THEM4-like
ENSOARG00000019845	1078.1		3	enJSRV-20 endogenous virus Jaagsiekte sheep retrovirus
ENSOARG00000015448	654.7	LOC114115259*	1	endogenous retrovirus group K member 6 Pro protein-like
ENSOARG00000014187	589.7	HamM*	19	endogenous virus Jaagsiekte sheep retrovirus
ENSOARG00000011311	544.3	CLDN7	11	claudin 7
ENSOARG00000010828	436.5	HAVCR1*	5	hepatitis A virus cellular receptor 1
ENSOARG00000004021	420.7	GRP*	23	gastrin releasing peptide
ENSOARG00000003992	389.0	SLC44A4	20	solute carrier family 44 member 4
ENSOARG00000019903	367.3		14	endogenous retrovirus group K member 9 Pol protein-like
ENSOARG00000004279	333.7	SLC7A9	14	solute carrier family 7 member 9
ENSOARG00000018755	327.1	TNNI1	12	troponin I1, slow skeletal type
ENSOARG00000013020	316.0	HSD11B1	12	hydroxysteroid 11-beta dehydrogenase 1
ENSOARG00000003224	300.3	SMPDL3B	2	sphingomyelin phosphodiesterase acid like 3B
ENSOARG00000012115	266.9		15	Jaagsiekte sheep retrovirus-like element
ENSOARG00000013158	263.8	SPDYC	21	speedy/RINGO cell cycle regulator family member C
ENSOARG00000006119	257.5	MMP7	15	matrix metallopeptidase 7
ENSOARG00000004751	246.5	TMEM92	11	transmembrane protein 92
ENSOARG00000010191	234.5	PEBP4	2	phosphatidylethanolamine binding protein 4
ENSOARG00000007680	232.5	SLC34A2	6	solute carrier family 34 member 2
ENSOARG00000004297	216.7	ALOX12*	11	arachidonate 12-lipoxygenase, 12S type
ENSOARG00000007592	211.8	PTGS2	12	prostaglandin-endoperoxide synthase 2
ENSOARG00000008009	211.1	CDH17	9	cadherin 17
ENSOARG00000014451	204.7	IFI27L2	18	interferon alpha-inducible protein 27-like protein 2

Table 2: List of the 25 most abundant genes in the CL and endometrium. Fifteen of the top 25
genes were shared by both tissues and were dominated by mitochondrial genes. The table includes
the Ensembl gene ID, chromosome number (Chr.), gene ID (GeneID) and gene description. The table
is divided into three sections; the first section lists the 25 genes that were shared by the two tissues,
and the other two list the remaining 10 genes in the endometrium and CL. Gene IDs and annotations
that were not available in BioMart were retrieved based on a homology search using the nucleotide
BLAST (marked with an asterisk, "*") or on information available in Ensembl (marked with a hash,
"#").

	Chr.	GeneID	Description
Common			
ENSOARG00000007815	6	LOC105580399*	Cercocebus atys 60S ribosomal protein L41-like
ENSOARG00000000019	MT	COX2	cytochrome c oxidase subunit II

ENSOARG00000018666	7	RPLP1	ribosomal protein lateral stalk subunit P1
ENSOARG00000000035	MT	CYTB	cytochrome b
ENSOARG00000000021	MT	ATP8	ATP synthase F0 subunit 8
ENSOARG00000007617	10	TCTP	tumor protein, translationally controlled 1
ENSOARG00000000023	MT	COX3	cytochrome c oxidase subunit III
ENSOARG00000003793	23	TSMB4X	thymosin beta-4
ENSOARG00000000037	MT	Mt tRNA#	mitochondrial tRNA
ENSOARG00000020724	3	MGP	matrix Gla protein
ENSOARG00000000016	MT	COX1	cytochrome c oxidase subunit I
ENSOARG00000000022	MT	ATP6	ATP synthase F0 subunit 6
ENSOARG00000000028	MT	ND4	NADH dehydrogenase subunit 4
ENSOARG00000003782	7	B2M	beta-2-microglobulin
ENSOARG00000000006	MT	ND1	NADH dehydrogenase subunit 1
Endometrium			
ENSOARG00000019088	AMGL01125506.1	LGALS15	lectin, galactoside-binding, soluble, 15
ENSOARG00000003184	24	NUPR1	nuclear protein 1, transcriptional regulator
ENSOARG00000019924	1	BCL2L15	BCL2-like 15
ENSOARG00000006202	13	CST3	cystatin C
ENSOARG00000021079	1	S100A11	S100 calcium-binding protein A11
ENSOARG00000016080	13	ATP5F1E	PRELI domain-containing 3B
ENSOARG00000013018	X	S100G	S100 calcium-binding protein G
ENSOARG00000019491	3	OST4	oligosaccharyltransferase complex subunit 4, noncatalytic
ENSOARG00000001346	21	CST6	cystatin E/M
ENSOARG00000006149	8	EEF1A1	eukaryotic translation elongation factor 1 alpha 1
CL			
ENSOARG00000004595	13	OXT	oxytocin/neurophysin I prepropeptide
ENSOARG00000022293	13	RF02216#	misc. RNA
ENSOARG00000000027	MT	ND4L	NADH dehydrogenase subunit 4L
ENSOARG00000002586	15	APOA1	apolipoprotein A1
ENSOARG00000002472	25	MSMB	microseminoprotein beta
ENSOARG00000001269	26	STAR	steroidogenic acute regulatory protein
ENSOARG00000000010	MT	ND2	NADH dehydrogenase subunit 2
ENSOARG00000013157	X	TIMP1	TIMP metallopeptidase inhibitor 1
ENSOARG00000020402	1	HSD3B1	hydroxy-delta-5-steroid dehydrogenase, 3 beta- and steroid delta-isomerase 1
ENSOARG00000000033	MT	ND6	NADH dehydrogenase subunit 6

977 Table 3: Numerical summary of differentially expressed genes in the CL and endometrium. Legend:
978 FS – Finnsheep, TX – Texel, F1 – F1-cross

Comparison	CL		Endometrium	
	Upregulated	Downregulated	Upregulated	Downregulated

FS vs TX	140	59	22	21
FS vs F1	49	18	2	0
TX vs F1	13	9	5	12

980 **Table 4 Significantly differentially expressed miRNAs.** The last column indicates the breedwise
981 comparisons, where FS represents Finnsheep, TX represents Texel ewes and F1 represents F1 crosses
982 of Finnsheep and Texel sheep. In addition, the CL and endometrium are denoted with the prefixes c
983 and e.

miRNA	BaseMean	Log2FoldChange	Padj	Comparison
rno-miR-451-5p	1189.41	0.86	0.05	cFS-TX
oar-18_757_mt	20.48	1.16	0.05	cFS-TX
oar-18_882_mt	73.39	-3.33	0.04	eFS-TX
oar-18_882_mt	73.39	3.18	0.06	eTX-F1
oar-JH921839.1_2216_mt	365.99	4.19	0.02	eFS-F1
oar-18_870_mt	92.35	4.4	0.02	eFS-F1

13 Supplementary Material

Fig. S1: KEGG pathways associated with the DEGs upregulated in Finnsheep compared to Texel
ewes in the CL

Fig. S2: GO terms associated with DEGs upregulated in Finnsheep compared to F1 crosses

Fig. S3: PCA of the top 500 expressed miRNAs in the CL (left) and endometrium (right)

Table S1: Phenotype data of the samples

Table S2: Summary of the samples included in mRNA-Seq

Table S5S3: List of DEGs between the CLs of Finnsheep and Texel ewes

Table S4: List of GO terms associated with the upregulated genes in the CLs of Finnsheep compared
to Texel ewes

Table S5: List of KEGG pathways associated with the upregulated genes in the CLs of Finnsheep
compared to Texel ewes

Table S6: List of DEGs between the CLs of Finnsheep and those of F1 crosses

Table S7: List of GO terms associated with the upregulated genes in the CLs of Finnsheep compared
to F1 crosses

Table S8: List of DEGs between the CLs of Texel ewes and F1 crosses

Table S9: List of DEGs between the endometria of Finnsheep and Texel ewes

Table S10: List of DEGs between the endometria of Finnsheep and F1 crosses

Table S11: List of DEGs between the endometria of Texel and F1 crosses

Table S12: Summary of the miRNA-Seq data

Data Availability Statement

The raw FASTQ sequence data (for both mRNAs and miRNAs) from this study have been deposited
in the European Nucleotide Archive (ENA) database under accession code PRJEB32852. The
accession codes for each sample are included in the sample summary tables (Supplementary table S3
and S12 for mRNA and miRNA, respectively).