Gene expression profiling of *corpus luteum* reveals the importance of immune system during early pregnancy in domestic sheep.

Kisun Pokharel[1], Jaana Peippo[2] Melak Weldenegodguad[1], Mervi Honkatukia[2], Meng-Hua Li[3*], Juha Kantanen[1*]

[1] Natural Resources Institute Finland (Luke), Jokioinen, Finland
[2] Nordgen – The Nordic Genetic Resources Center, Ås, Norway
[3] College of Animal Science and Technology, China Agriculture University, Beijing, China
* Correspondence: MHL, menghua.li@cau.edu.cn; JK, juha.kantanen@luke.fi

Abstract: The majority of pregnancy loss in ruminants occurs during the preimplantation stage, which is thus the most critical period determining reproductive success. While ovulation rate is the major determinant of litter size in sheep, interactions among the conceptus, *corpus luteum* and endometrium are essential for pregnancy success. Here, we performed a comparative transcriptome study by sequencing total mRNA from corpus luteum (CL) collected during the preimplantation stage of pregnancy in Finnsheep, Texel and F1 crosses, and mapping the RNA-Seq reads to the latest Rambouillet reference genome. A total of 21,287 genes were expressed in our dataset. Highly expressed autosomal genes in the CL were associated with biological processes such as progesterone formation (*STAR*, *CYP11A1*, and *HSD3B1*) and embryo implantation (eg. *TIMP1*, *TIMP2* and *TCTP*). Among the list of differentially expressed genes, a group of sialic acid-binding immunoglobulin (Ig)-like lectins (Siglecs), solute carriers (*SLC13A5*, *SLC15A2*, *SLC44A5*) and chemokines (*CCL5*, *CXCL13*, *CXCL9*) were upregulated in Finnsheep, while four multidrug resistance-associated proteins (MRPs) were upregulated in Texel ewes. A total of 17 genes and two non-coding RNAs (ncRNA) were differentially expressed in breed-wise comparisons owing to flushing diet effect. Moreover, we report, for the first time in any species, several genes that are active in the CL during early pregnancy (including *SIGLEC13*, *SIGLEC14*, *SIGLEC6*, *MRP4*, and *CA5A*). Importantly, functional analysis of differentially expressed genes suggested that Finnsheep have a better immune system than Texel and that high prolificacy in Finnsheep might be governed by immune system regulation.

Keywords: Finnsheep, Texel, progesterone, preimplantation

1. Introduction

Litter size, a key determinant for the profitability of sheep production systems, is highly dependent on ovulation rate and embryo development in the uterus. Earlier studies have shown that the trait of high prolificacy can result due to the action of either a single gene with a major effect, as in the Chinese Hu, Boorola Merino, Lacaune and small-tailed Han breeds (Mulsant et al., 2001; Souza et al., 2001; Davis et al., 2002, 2006; Chu et al., 2007; Drouilhet et al., 2013), or different sets of genes, as in the Finnsheep and Romanov breeds (Ricordeau et al., 1990; Xu et al., 2018). The Finnsheep or Finnish landrace, one of the most highly prolific breeds, has been exported to more than 40 countries to improve local breeds, although the heritability of ovulation rate is low (Hanrahan and Quirke, 1984). In recent years, a FecG^F (V371M) mutation in gene *GDF9* has been identified to be strongly associated with litter size in Finnsheep and breeds such as the Norwegian White Sheep, Cambridge and Belclare breeds, which were developed using Finnsheep (Hanrahan et al., 2004; Våge et al., 2013; Mullen and Hanrahan, 2014; Pokharel et al., 2018).

The success of pregnancy establishment in sheep and other domestic ruminants is determined at the preimplantation stage and involves coordination among pregnancy recognition, implantation and placentation, in which the corpus luteum (CL) and endometrium play vital roles (Geisert et al., 1992; Spencer et al., 2004b, 2007). The preimplantation stage of pregnancy is the most critical period in determining the litter size because of the high embryo mortality during this period. It has been shown that most embryonic deaths occur before day 18 of pregnancy in sheep (Quinlivan et al., 1966; Bolet, 1986; Rickard et al., 2017). However, due to the biological complexity of the process and to technical difficulties, embryo implantation is still not well understood.

The CL is an endocrine structure whose main function is to synthesize and secrete the hormone
progesterone. Progesterone production is essential for the establishment of pregnancy. However, if pregnancy is
not established, the CL will regress as a result of luteolysis, and a new cycle will begin.

The whole-transcriptome profiling approach enables a deeper understanding of the functions of the CL,
which may allow the identification of genes and markers that are differentially expressed, for example, between
breeds showing different litter size phenotypes. Although most of the studies associated with early pregnancy
have been performed in sheep (Spencer et al., 2004b, 2007; Mamo et al., 2012; Bazer, 2013; Raheem, 2017),
only a few studies have applied transcriptomic approaches to the CL. A microarray-based transcriptomic study
conducted by Gray et al. (2006) identified a number of genes regulated by progesterone (from the CL) and
interferon tau (*IFNT*; from the conceptus) in pregnant vs uterine gland knockout (UGKO) ewes (Gray et al.,
2006). In a more comprehensive study conducted by Brooks et al. (2016), transcriptome analysis of uterine
epithelial cells during the peri-implantation period of pregnancy identified various regulatory pathways and
biological processes in sheep (Brooks et al., 2016). Moore et al. (2016) combined gene expression data with
genome-wide association studies (GWASs) to understand the roles of CL and endometrium transcriptomes in
dairy cattle fertility (Moore et al., 2016). Another study by identified differentially expressed genes (DEGs)
between Day 4 and Day 11 in the CL in cattle (Kfir et al., 2018). Though these studies have certainly enhanced
our understanding of the roles of the CL during early pregnancy and in ruminant fertility in general, none of the
studies have conducted specific comparisons between breeds with different reproductive potential. Thus, in this
study, a comparison of transcriptome profiles between two breeds (high prolific Finnsheep and low-prolific
Texel) was conducted to provide insight into the similarities in developmental events in early pregnancy
between the breeds. Using F1 crosses we were able to better understand the heritability of the genetic markers.
Here, the main goal of this study was to build a global picture of transcriptional complexity in Cl and examine
differences in developmental profiles during early pregnancy in sheep breeds showing contrasting fertility
phenotypes. Thus, this study has relevance to sheep breeding towards achieving improved reproductive
capacity.
74
75

2.   Materials and Methods

2.1. Experimental design

All procedures for the experiment and sheep sampling were approved by the Southern Finland Animal
Experiment Committee (approval no. ESAVI/5027/04.10.03/2012). The animals were kept at Pusa Farm in
Urjala, located in the province of Western Finland, during the experimental period. A total of 31 ewes
representing three breed groups (Finnsheep (n=11), Texel (n=11) and F1 crosses (n=9) were included in the
main experiment (please note that only 18 of the 31 ewes have been included in this study). Analyses were
conducted for two different time points during the establishment of pregnancy: the follicular growth phase
(Pokharel et al., 2018) and early pregnancy prior to implantation (current study). After ovary removal, the ewes
were mated using two Finnsheep rams, and the pregnant ewes were slaughtered during the preimplantation
phase of the pregnancy when the embryos were estimated to be one to three weeks old (Supplementary Table
S1). At the slaughterhouse, a set of tissue samples (the pituitary gland, a CL, oviductal and uterine epithelial
cells, and preimplantation embryos) were collected and stored in RNAlater reagent (Ambion/Qiagen, Valencia,
CA, USA) following the manufacturer's instructions. Of the collected tissue samples, CL was subjected to
current study. One of the CLs was dissected from each ovary. For the present study, and particularly for the
RNA-Seq of the CL, six ewes each from the Finnsheep, Texel and F1 cross groups were included. Therefore,
out of 31 ewes that were originally included in the main experiment, only 18 have been considered here. The
experimental design have been described in more detail in an earlier study (Pokharel et al., 2018).

2.2. Library preparation and sequencing

RNA was extracted from the tissues using an RNeasy Plus Mini Kit (Qiagen, Valencia, CA, USA)
following the manufacturer's protocol. The details on RNA extraction have been described previously
(Pokharel et al., 2018). RNA quality (RNA concentration and RNA integrity number) was measured using a
Bioanalyzer 2100 (Agilent Technologies, Waldbronn, Germany) before sending the samples to the Finnish
Microarray and Sequencing Center, Turku, Finland, where library preparation and sequencing were performed.

RNA libraries were prepared according to the Illumina TruSeq® Stranded mRNA Sample Preparation Guide
(part # 15031047) which included poly-A selection step. Unique Illumina TruSeq indexing adapters were
ligated to each sample during an adapter ligation step to enable pooling of multiple samples into one flow cell
lane. The quality and concentrations of the libraries were assessed with an Agilent Bioanalyzer 2100 and by
Qubit® Fluorometric Quantitation, Life Technologies, respectively. All samples were normalized and pooled
for automated cluster preparation at an Illumina cBot station. High-quality libraries of mRNA were sequenced
with an Illumina HiSeq 2000 instrument using paired-end (2x100 bp) sequencing strategy.

2.3. Data preprocessing and mapping

The raw reads were assessed for errors and the presence of adapters using FastQC v0.11.6 (Simon
Andrews). As we noticed the presence of adapters, Trim Galore v0.5.0 (Felix Krueger; Martin, 2011) was used
to remove the adapters and low-quality reads and bases. Clean RNA-Seq reads were aligned to the latest sheep
reference genome using STAR v2.6.1a (Dobin et al., 2013). The reference genome (oar_rambouillet_v1.0) and
annotation (NCBI Ovis aries Annotation Release 103) were used to construct "Star genome" prior to mapping
step. In order to facilitate the differential expression analysis, "*--quantMode GeneCounts*" option was included
in the STAR mapping command. Alternatively, we performed transcript quantification under the
quasi-mapping-based mode in Salmon v1.1.0 (Patro et al., 2017) transcriptome index built using
oar_rambouillet_v1.0 transcriptome (https://www.ncbi.nlm.nih.gov/assembly/GCF_002742125.1/).

2.4. Differential gene expression analysis

The raw counts quantified as part of STAR mapping were considered for gene expression analysis
whereas the Salmon-based transcript estimates (TPM, transcripts per million) were summarized to gene level
estimates to prepare a table of genes with their abundance using tximport v1.12.3 (Soneson et al., 2016)
Bioconductor package. Prior to gene level summarization, customized "*tx2gene*" data frame was created using
the annotation (*.gtf file*) of rambouillet v1.0 transcriptome. We used DESeq2 (Love et al., 2014) for differential
gene expression analysis. Transcripts with less than 5 read counts were discarded and technical replicates
(samples representing same animal) were collapsed/merged before running the DESeq command. Pairwise
differential expression analysis was performed between Finnsheep, Texel and F1 crosses. Differentially
expressed genes with adjusted p-value of 0.05 (*padj<0.05*) and absolute log 2(fold change) of 1.5
(*abs(log2FoldChange)>1.5*) were regarded as significant in this study. In DESeq2, Benjamini & Hochberg's
method is used for estimating the adjusted p-values. For identifying a subset of genes potentially differentially
expressed due to flushing diet effect, we employed a separate design by including diet as a secondary factor.
The statistical analyses were performed in R v3.6.0.

2.5. Functional analysis of differentially expressed genes

The ClueGO v2.5.5 (Bindea et al., 2009) plugin in Cytoscape v3.7.0 (Shannon et al., 2003) was employed
for gene functional analysis. Prior to performing the analyses, we downloaded the latest versions of the Kyoto
Encyclopedia of Genes and Genomes (KEGG) pathways and Gene Ontology (GO) terms. The enrichment
analysis was based on a one-sided hypergeometric test with the Benjamini-Hochberg correction method. We
used a custom reference set that included a list of all the expressed genes in our dataset. We also modified the
default GO and pathway selection criteria in such a way that a minimum of three genes and four percent of
genes from a given GO or KEGG pathway should be present in the query list. Furthermore, GO terms with a
minimum level of three and a maximum level of eight were retained. Finally, GO terms and/or KEGG pathways
sharing similar genes were grouped together using kappa statistics with kappa score threshold of 0.4.

3. Results

3.1. Phenotypic observations

After removal of the remaining ovary, we counted the number of CLs visually in each animal. With an
average of 4.09, Finnsheep had the highest number of CLs, whereas Texel had an average of 1.7 CLs (Table 1,
Supplementary Table S1). F1 showed phenotypes closer to those of Finnsheep than those of Texel, having 3.75
CLs on average (Supplementary Table S1); this was unsurprising, as we observed a similar pattern in an earlier
study (Pokharel et al., 2018). We did not observe more than 2 CLs in the Texel or fewer than 3 CLs in Finnsheep

or F1. Similarly, on average, Finnsheep had the highest number of embryos (n=2.6), followed by F1 crosses
(n=1.8) and Texel (n=1.5). Interestingly, the embryo survival rate in Texel was highest, where 1.5 embryos
were present from 1.7 CLs (88%) on average. On the other hand, Finnsheep (63%) and F1 cross (48%) had
remarkably low embryo survival rate. While these findings are based on fewer animals, the results are in line
with earlier studies (Rhind et al., 1980; Silva et al., 2016) where typically higher litter size is associated with
higher embryo mortality and vice versa. It would be of great interest to determine if productivity follows the
same pattern in F2 (i.e., F1 x F1) crosses, backcrosses and presumably also in a reciprocal cross.

Table 1. Average number of CL and embryos in Finnsheep, Texel and their F1 crosses. The full details of the
phenotypes are available in supplementary table S1.

Breed	# CL	# Embryos
Finnsheep	4.1	2.6
Texel	1.7	1.5
F1	3.75	1.8

3.2. RNA-Seq data

From the 42 libraries (21 from each tissue, including three technical replicates), around 1,104 million (M)
raw reads were sequenced, of which 1,094 M clean reads were retained after trimming (Table 2). The summary
statistics from Trim Galore revealed that up to 3.6% of the reads were trimmed, with reverse-strand reads
having a comparatively higher percentage of trimmed bases. However, the percentage of reads that were
excluded for being shorter than 18 bp was always less than 1% across all samples. More than 80% of the reads
from all samples were aligned to the Rambouillet reference genome and transcriptome using STAR and
Salmon, respectively. The raw RNA-Seq data (Fastq files) from this study are available in European Nucleotide
Archive (ENA) under project accession code PRJEB32852. The details of sample summary including ENA
accession code of individual samples is available in Table 2.

Table 2. Sample Summary. Here, FS, TX and F1 represent Finnsheep, Texel and F1 crosses, respectively. Diet
is either normal diet (C) or flushing diet (F). Alignment statistics from STAR include only the uniquely aligned
reads whereas those from salmon include all reads that mapped to reference transcriptome.

Sample	Breed	Diet	Raw reads (M)	Clean reads (M)	Uniquely aligned reads (STAR, M)	Aligned reads (salmon, M)	ENA accession code
1033	TX	C	56.4	55.9	48.0	50.7	ERR3349023
107A	TX	F	55.0	54.5	46.5	49.3	ERR3349025
107B	TX	F	55.7	55.2	48.3	49.8	ERR3349027
251	TX	F	41.8	41.2	35.4	36.9	ERR3349029
302	TX	C	54.5	54.1	47.1	48.5	ERR3349031
312A	FS	F	54.9	54.4	47.0	49.3	ERR3349033
312B	FS	F	79.2	78.4	69.2	71.1	ERR3349035
3609	FS	C	54.5	54.1	47.0	49.0	ERR3349037
379	TX	F	43.1	42.7	36.0	38.3	ERR3349039
4208	F1	C	54.3	53.8	44.8	49.0	ERR3349041
4271A	F1	F	47.2	46.8	40.1	42.4	ERR3349043
4271B	F1	F	46.9	46.4	40.0	42.1	ERR3349045
4519	F1	F	51.6	51.1	44.5	46.3	ERR3349047
4563	F1	F	54.1	53.7	45.7	48.6	ERR3349049
4590	F1	F	48.5	48.1	40.0	43.5	ERR3349051
4823	F1	C	52.6	52.1	42.6	47.2	ERR3349053
48	FS	C	53.4	53.0	44.7	47.9	ERR3349055

554	FS	F	56.0	55.5	48.1	49.7	ERR3349057
73	TX	C	46.1	45.7	39.1	41.3	ERR3349059
897	FS	F	53.3	52.9	46.1	47.9	ERR3349061
974	FS	C	45.1	44.7	38.6	40.4	ERR3349063

3.3. Gene expression in the CL

STAR-based alignment revealed 23,327 genes and pseudogenes expressed in the whole data set which makes up approximately 70% of the list (n=33,372) available for the recent Rambouillet reference genome. Further grouping of the expressed genes showed that the most genes (n=21,052) were expressed in Finnsheep followed by Texel (n=20,957) and F1 crosses (n=20,928). The cumulative difference in the number of genes in different samples and breeds might be due to transcriptional noise. As shown in Fig. 1, the highest number of breed-specific genes was found in Finnsheep (n=399), followed by Texel (n=363) and F1 crosses (n=303). In a pairwise comparison, based on overall gene expression, Finnsheep and F1 crosses shared a higher number of genes (n=338) than other pairs, indicating the closer relatedness of F1 crosses to Finnsheep (Fig. 1). As indicated by the principal component analysis (PCA), we did not observe any breed-specific clusters in either of the tissues and was also the case in our earlier ovarian transcriptome study (Pokharel et al., 2018). Alternatively, salmon-based quantification revealed 27,072 genes expressed in the CL of which 18,463 had TPM greater than 0.1 (Supplementary Table S2).

Figure 1: Graphical summary of gene expression in the CL. (A) Shared and unique genes expressed in the CL of Finnsheep, Texel and their F1. (B) PCA plot based on variance stabilized transformation (VST) of gene expression counts derived from DESeq2 in three breeds.

3.4. Highly expressed genes

To obtain an overview of the most abundant genes in the tissue, we selected the top 25 genes expressed in the CL (Table 3) derived from Salmon quantification. We noticed that 10 out of the top 25 genes were ribosomal proteins. Moreover, top expressed genes also appeared to play substantial roles during the preimplantation stage. Steroidogenic acute regulatory protein (*STAR*), the second most highly expressed gene, plays an important role in mediating the transfer of cholesterol to sites of steroid production (Stocco, 2000; Christenson and Devoto, 2003). Post ovulation, the expression of the majority of genes associated with progesterone synthesis starts to increase and peaks around the late luteal phase, when the CL has fully matured (Juengel et al., 1995; Devoto et al., 2001; Davis and LaVoie, 2018). *STAR*, together with the cytochrome P450 side chain cleavage (P450cc) complex and 3b-hydroxysteroid dehydrogenase/delta5 delta4-isomerase (*HSD3B1*), are the three most important actors involved in progesterone biosynthesis. *STAR* is involved in transporting free cholesterol to the inner mitochondrial membrane. The P450cc complex, composed of a cholesterol side chain cleavage enzyme (*CYP11A1*), ferredoxin reductase (*FDXR*) and ferredoxin (*FDX1*), converts the newly arrived

cholesterol into pregnenolone (King and LaVoie, 2009). Finally, *HSD3B1* helps in converting pregnenolone to
progesterone (Hu et al., 2010; Plant et al., 2015; Stouffer and Hennebold, 2015; Davis and LaVoie, 2018). Three
of these major genes involved in progesterone synthesis (*STAR, HSD3B1,* and *HSD3B1*) were ranked among the
top 25 most highly expressed autosomal genes, while *FDX1* (TPM=972.8) and *FDXR* (TPM=221.5) were also
highly expressed.
Oxytocin (*OXT*) was one of the most highly expressed genes in the CL. In cyclic ewes, *OXT* secreted
from the CL and posterior pituitary is widely known to bind with oxytocin receptor (*OXTR*) from the
endometrium to concomitantly release prostaglandin $F_{2\alpha}$ (*PGF*) pulses and induce luteolysis (Flint and
Sheldrick, 2004; Spencer et al., 2004a, 2004b; Bazer, 2013). However, for noncyclic ewes, *OXT* plays an
important role during peri-implantation and throughout pregnancy (Kendrick, 2000). *OXT* signaling is known to
be influenced by progesterone, but the mechanism underlying the regulation is not yet clear due to conflicting
findings (Grazzini et al., 1998; Gimpl et al., 2002; Fleming et al., 2006; Bishop, 2013). Meanwhile *OXTR*
(TPM=0.2) expression was almost negligible compared to OXT expression.

Table 3. List of top 25 most abundant genes expressed in the CL. Gene names and gene descriptions for gene IDs
starting with "LOC" were retrieved from NCBI Genbank (https://www.ncbi.nlm.nih.gov/genbank/) and,
GeneCards (https://www.genecards.org/), respectively.

GeneID	Gene Description	Chr	mean TPM
TIMP1	Tissue inhibitor of metallopeptidase 1	X	16705.6
STAR	Steroidogenic acute regulatory protein	26	14226.6
OXT	oxytocin/neurophysin I prepropeptide	13	13113.2
MGP	matrix Gla protein	3	12294.0
LOC114112617	ncRNA (uncharacterized)	un	12155.9
LOC114113966	translationally controlled tumour protein (*TCTP*)	3	8341.7
HSD3B1	hydroxy-delta-5-steroid dehydrogenase, 3 beta- and steroid delta-isomerase 1	1	8249.3
LOC101117785	Apolipoprotein A1 (*APOA1*)	15	8213.2
LOC101110773	elongation factor 1-alpha 1 (*EEF1A1*)	10	7564.5
LOC101103639	uncharacterized (protein coding)	3	7272.8
LOC114116158	thymosin beta-4 pseudogene	1	6689.3
LOC114118103	40S ribosomal protein S29	14	5599.0
CYP11A1	Cytochrome P450 Family 11 Subfamily A Member 1	18	5096.8
RPLP1	ribosomal protein lateral stalk subunit P1	7	5032.2
LOC114116632	60S ribosomal protein L23a (*RPL23a*)	10	4998.9
FTH1	Ferritin heavy chain 1	21	4498.0
RPS8	ribosomal protein S8	1	4402.1
LOC105606567	60S ribosomal protein L39	4	4215.9
MSMB	microseminoprotein beta	25	4093.6
RPS24	ribosomal protein S24	13	4029.1
RPS17	ribosomal protein S17	18	3950.0
LOC114108766	60S ribosomal protein L17 (*RPL17*)	4	3832.5
RPS18	ribosomal protein S18	20	3657.2
TIMP2	Tissue inhibitor of metallopeptidase 2	11	3587.4
LOC114114908	40S ribosomal protein S27 (*RPS27*)	5	3570.4

219
Two members of tissue-derived matrix metalloproteinase (*MMP*) inhibitors, also referred to as tissue
inhibitors of metallopeptidases (TIMPs) were among the top 25 most abundant genes of which *TIMP1* was
ranked number 1 and *TIMP2*, number 24. In addition to *TIMP1* and *TIMP2*, at least 20 MMPs (including two
isoforms of *MMP9* and *MMP24*) two TIMPs, *TIMP3* (TPM=265) and *TIMP4* (TPM=0.5) were expressed in our
samples. The high level of TIMPs indicated successful pregnancies as these inhibitors have low level of

expression while their target MMPs are elevated during luteolysis (Curry and Osteen, 2001; Kliem et al., 2007;
Nissi et al., 2013). TIMPs have important role in regulation of several processes relevant to uterine physiology
including angiogenesis (Moses et al., 1990), cell differentiation (Docherty et al., 1985), and embryo
development (Nakanishi et al., 1986). Northern blot analyses in several tissues of sheep showed that both
TIMP1 and *TIMP2* had greatest abundance in CL during early pregnancy (Hampton et al., 1995). Moreover,
MMPs and TIMPs have important roles in implantation and that the high level expression of *TIMP1* and *TIMP2*
indicated active invasion of trophoblast cells during implantation (Librach et al., 1991; Bai et al., 2005).
Another study in bovine oviduct showed that *TIMP1* and *TIMP2* were highly expressed during the time of
ovulation (Gabler et al., 2001). Hence, these two genes may play significant role during the whole reproduction
process – from ovulation to implantation.

Translationally controlled tumor protein (*TCTP*) is a highly conserved, multifunctional protein that plays
essential roles in development and other biological processes in different species (Tuynder et al., 2002; Chen et
al., 2007; Brioudes et al., 2010; Li et al., 2011; Branco and Masle, 2019). With a maximum level of expression
on Day 5 of pregnancy, this protein has been shown to play a significant role in embryo implantation in mice (Li
et al., 2011). Consistent with these earlier studies, *TCTP* appeared to have the highest level of expression during
the embryo implantation period. Matrix Gla protein (*MGP*) is a vitamin K-dependent extracellular matrix
protein whose expression has been shown to be correlated with development and maturation processes (Zhao
and Nishimoto, 1996; Zhao and Warburton, 1997) and receptor-mediated adhesion to the extracellular matrix
(Loeser and Wallin, 1992). Several studies have reported that *MGP* is highly expressed in the bovine
endometrium (Spencer et al., 1999; Mamo et al., 2012; Forde et al., 2013) and we have shown that it's also the
case with CL. The high level of expression of *MGP* in our study is consistent with the results of earlier studies in
which this gene was found to be elevated during the preimplantation stage in sheep (Spencer et al., 1999; Gray
et al., 2006) and cattle (Mamo et al., 2012). Similarly, (Casey et al., 2005) reported that *MGP* was significantly
upregulated in non-regressed compared to regressed bovine CLs. Our data and supporting results from earlier
studies on cattle show that *MGP* is highly expressed in CL during the preimplantation stage and plays important
roles in superficial implantation and placentation in sheep.

3.5. Breed wise gene expression differences in the CL

Out of the 133 significant DEGs in the CL of pure breeds (i.e Finnsheep *vs* Texel), 90 were upregulated in
Finnsheep (Supplementary table S4) and the rest were downregulated. In the list of DEGs, we observed a few
cases in which more than one gene from the same family was present. Four isoforms of multidrug
resistance-associated protein 4-like (*MRP4*) were upregulated in Texel ewes. Earlier reports have suggested a
role of *MRP4* in transporting prostaglandins in the endometrium (Lacroix-Pépin et al., 2011), and *MRP4* has
been found to be upregulated in the endometrium in infertile cows compared to fertile cows (Moore et al.,
2016). Although there are no reports regarding the existence and roles of *MRP4* in the CL, we speculate that the
comparatively lower levels of these prostaglandin (PG) transporters in Finnsheep provide a luteoprotective
effect. Five sialic acid-binding Ig-like lectins (Siglecs) including three isoforms of *SIGLEC-14* were
upregulated in Finnsheep. Siglecs are transmembrane molecules expressed on immune cells and mediate
inhibitory signaling (Varki and Angata, 2006). So far, *SIGLEC-13* has been reported only in nonhuman
primates; it was deleted during the course of human evolution (Angata et al., 2004). The importance of Siglecs
in immune system regulation has been reviewed elsewhere (Pillai et al., 2012). Siglecs constantly evolve
through gene duplication events and may vary between species and even within a species (Cao and Crocker,
2011; Pillai et al., 2012; Bornhöfft et al., 2018). Here we have reported the expression of Siglecs in CL which
are known to play a role in the immune response during early pregnancy (preimplantation).

Figure 2. Volcano plot of 23327 genes expressed in the pure breeds. Top 23 of 133 genes significantly differentially expressed between Finnsheep and Texel are denoted by gene names except three unknown ncRNAs. All DE genes with absolute log2 fold change greater than 2 and adjusted P value lower than 0.001 have been highlighted here. For additional details and complete list of 133 DE genes please refer supplementary table. Note that *TXNL1* gene (marked by an asterisk) has an adjusted P-value of 4.84E-16. Volcano plot created using Bioconductor package EnhancedVolcano (Blighe et al., 2019) and some adjustments in the figure were made in Adobe Illustrator (Adobe Inc.). The x-axis represents log2 fold change and the y-axis represents adjusted p-values(-log10). Legends in top: NS = non-significant; Log2 FC = genes having absolute log2 fold change greater than 2; P = genes with adjusted P value less than 0.001; and P & Log2 FC: genes passing the criteria for log2 fold change and adjusted P value.

Several cytokines including four chemokines (*CCL5*, *CCL24*, *CXCL9*, and *CXCL13*), two interleukin receptors (*IL1RN*, *IL12RB1*) and two interferons (*ISG20*, *GVINP1*) were upregulated in Finnsheep. Other genes

with more than one member included major histocompatibility complexes (MHCs) (*HA1B*, *MICB*, *MICA*,
BOLA-DQB*0101, etc.), tripartite motif containing proteins (two isoforms of *TRIM5*, and *TRIM10*), cluster of
differentiation factors (*CD4*, *CD69*), CD300 family molecules (*CD300C*, and *CD300H*). A novel protein
coding gene (*LOC114116052*) was significantly upregulated in Texel. Nucleotide blast search of this novel
gene's CDS revealed its sequence similarity with *SENP6* in several mammalian species including transcript
variants of sheep *SENP6* (*LOC101118793*, 95% query coverage and 98.7% sequence identity). Therefore, we
concluded that *LOC114116052* is one of the isoforms of *SENP6*. All four ribosomal proteins (*MRPS33*, *RPS3A*,
RPL17, and *RPL34*) differentially expressed between the pure breeds were upregulated in Finnsheep. Similarly,
16 of the differentially expressed genes were classified as uncharacterized ncRNAs of which nine were
upregulated in Finnsheep.

Out of 90 genes that were significantly upregulated in Finnsheep compared to Texel, 62 were recognized
by ClueGO. However, 22 of the recognized genes lacked functional annotation. Thus, enrichment analysis for
GO terms and KEGG pathways is based on 40 genes. In the end, only 16 genes passed the selection criteria (see
materials and methods section) and were associated with 16 representative terms and pathways (Fig3). It turned
out that several of the terms and pathways comprised similar genes and grouping based on common genes
revealed four different types of terms and/or pathways. The enriched terms and pathways belonged to four main
categories based on the way genes were shared. The representative terms and/or pathways associated with
upregulated genes were "negative regulation of viral life cycle", "Th1 and Th2 cell differentiation",
"chemokine receptor binding" and, "dicarboxylic acid transport". In summary, genes involved in the immune
response were upregulated in Finnsheep CL during early pregnancy.

Figure 3. Go terms and KEGG pathways associated with significantly differentially expressed genes between
Finnsheep and Texel. The functional grouping option in ClueGO categorized 16 GO terms and pathways into
four groups using kappa score where the genes shared between the terms are iteratively compared to form
functional groups.

Altogether, 31 genes were differentially expressed between Finnsheep and F1 crossbred ewes, of which 19
genes were upregulated in Finnsheep (Supplementary table S7).
The lowest number (n=27) of DEGs was observed between Texel and F1 crossbred ewes, with two-third
genes being upregulated in F1 (Supplementary table S9).

3.6. Uniquely differentially expressed genes

We noticed that several genes (see Fig. 4) were differentially expressed in more than one comparison,
increasing our confidence in the identification of these DEGs. Few DEGs were exclusively up- or
downregulated in particular breed compared to other two breeds. *ANOS1*, *CLVS2*, *FOLR3*, *EEF1A1* and
LOC114115287 (uncharacterized ncRNA) were always upregulated in Finnsheep whereas seven genes were
exclusively downregulated. *MEPE* was found to be downregulated in Finnsheep also during the ovulation
(Pokharel et al., 2018). Seven genes (*DNER*, *LOC106990106* (*EZR*), *LOC114109364* (*H3.3*), *LOC114115573*
(*COX7C*), *LOC114115623* (*TOMM20*) and *LOC114115666* (*EEF2*)) were exclusively upregulated and 13
genes including three ncRNAs were exclusively downregulated in Texel (Fig. 4).

Figure 4. List of uniquely differentially expressed genes in Finnsheep, Texel and F1. Gene names that were not
part of Rambouillet reference transcriptome are marked with an asterisk (*) while gene IDs starting with LOC
are "uncharacterized ncRNA".

Similarly, *CA5A*, *FGF23* and *LOC114109527* (*NCLP*) were exclusively upregulated in F1 and *STX7* was
downregulated. Among these, *CA5A* appeared to be upregulated in F1 crosses compared to both Finnsheep and
Texel from both phases (i.e., in the CL in this study and in the ovary in our earlier study). *CA5A* is a member of
the carbonic anhydrase family of zinc-containing metalloenzymes, whose primary function is to catalyze the
reversible conversion of carbon dioxide to bicarbonate. The mitochondrial enzyme *CA5A* plays an important
role in supplying bicarbonate (HCO_3^-) to numerous other mitochondrial enzymes. In a previous study, we
observed downregulation of *CA5A* in the ovaries of Texel compared to F1 (Pokharel et al., 2018). More
recently, *CA5A* was also shown to be expressed in the ovaries of the Pelibuey breed of sheep; the gene was
upregulated in a subset of ewes that gave birth to two lambs compared to uniparous animals
(Hernández-Montiel et al., 2019). However, there are no reports regarding the expression and function of *CA5A*
in the CL. Based on the results from our current and earlier reports (Pokharel et al., 2018; Hernández-Montiel et

al., 2019), *CA5A* appears to have an important function, at least until the preimplantation stage of reproduction.
The level of expression in F1 crosses in the CL followed the same pattern as that in the ovary, which led us to
conclude that *CA5A* is heritable and potentially an imprinted gene. Further experiments are needed to determine
whether the gene is associated with high prolificacy.

3.7. Influence of flushing in gene expression

Four genes were significantly differentially expressed between Finnsheep and Texel of which three were
upregulated in Texel. The only gene upregulated in Finnsheep was cytochrome c oxidase subunit 7c,
mitochondrial (*COX7C*). The three genes significantly upregulated in Texel were *PAPPA2*, *LOC114116073*
(ncRNA, uncharacterized) and *LOC114118704* (Golgi apparatus membrane protein *TVP23* homolog B
pseudogene).
Out of 12 genes differentially expressed between Finnsheep and F1, five were upregulated in Finnsheep
while seven were upregulated in F1. *TVP23* pseudogene was always downregulated in Finnsheep as a result of
flushing diet. None of the six significantly differentially expressed genes were upregulated in Finnsheep
suggesting null influence of flushing diet in this breed. Similarly, *HOXD10* (see XU 2014, important for
regulation of embryonic implantation), *LOC105611004* (ncRNA, uncharacterized) and *LOC114116073* were
upregulated in F1 compared to Finnsheep.

Table 4. List of genes differentially expressed due to the effect of flushing diet in Finnsheep (FS), Texel (TX)
and F1 crosses (F1).

Gene	Gene*	BaseMean	LFC	Padj	Condition
COX7C		101.753584	1.93	0.03584019	FS vs TX
LOC114116073	un_ncRNA	832.4	-4.60	1.26E-28	FS vs TX
PAPPA2		24.4	-2.27	0.00091	FS vs TX
LOC114118704	TVP23(p)	38.8	-2.51	1.70E-06	FS vs TX
ADAMTS4		892.8005278	-1.80	0.02409534	FS vs F1
SHOX2		74.00640751	1.95	0.0200087	FS vs F1
HOXD10		99.29244359	-2.17	0.00279421	FS vs F1
SCTR		87.95488542	1.70	0.0146455	FS vs F1
HOXC11		24.11123058	-1.81	0.01918401	FS vs F1
LOC105611004	un_ncRNA	40.93791063	-2.06	0.00279421	FS vs F1
LOC114116073	un_ncRNA	832.360494	-4.82	5.58E-31	FS vs F1
LRP11		124.9792312	-1.84	0.02732988	FS vs F1
LOC114118704	TVP23(p)	38.85436938	-1.92	0.00279421	FS vs F1
BTN1A1		22.44434414	1.86	0.02926403	FS vs F1
KCNQ1		856.9910538	1.77	0.04977007	FS vs F1
HNR1PA1		81.09239312	1.93	0.0200087	FS vs F1
COX7C		101.753584	-1.88	0.03094773	TX vs F1
PCSK5		469.246216	1.69	0.01712337	TX vs F1
SCTR		87.9548854	1.79	0.00816462	TX vs F1
HOXC11		24.1112306	-1.92	0.00816462	TX vs F1
PPM1H		575.325443	1.67	0.00684851	TX vs F1
HOXA9		74.2026779	-1.96	0.00816462	TX vs F1
RDH10		2603.9004	1.56	0.00816462	TX vs F1
CA5A		983.421994	-1.50	0.01137482	TX vs F1

3.8. Limitations and thoughts for future studies

We acknowledge certain limitations of this study. With sequencing costs becoming increasingly
inexpensive, increasing the sample size of each breed group would certainly add statistical power. Given that
time-series experiments are not feasible with the same animal, sampling could be performed with a larger group
of animals at different stages of pregnancy to obtain an overview of gene expression changes. One ovary from

each ewe was removed earlier (Pokharel et al., 2018) and all the CLs for this study was collected from the
remaining ovary. Therefore, there might be some impact due to possible negative feedback effects on overall
gene expression. It should be noted that overall gene expression and, more specifically, differential expression
between breeds is inherently a stochastic process; thus, there is always some level of bias caused by individual
variation (Hansen et al., 2011). The results from breeding experiments have shown that productivity traits such
as litter sizes may not carry on to F2 crosses (F1 x F1) and/or backcrosses. Therefore, future experiments that
involve F2 crosses and backcrosses would provide more valuable findings related to prolificacy. Moreover, by
doing a reciprocal cross experiment, we might be able to get insight into POE and measure the potential
contribution of the Texel and Finnsheep in each cross. In addition, replicating such experiments in different
environments would be relevant for breeding strategies to mitigate the effects of climate change. To minimize
or alleviate noise from tissue heterogeneity, single-cell experiments may prove beneficial in future studies.

4. Conclusions

Our phenotypic records indicated lower embryo survival rate in Finnsheep compared to Texel. The
relative scarcity of transcriptomic information about the CL means that its functional importance is underrated.
We identified several key transcripts, including coding genes (producing mRNA) and noncoding genes, that are
essential during early pregnancy. Functional analysis primarily based on literature searches and earlier studies
revealed the significant roles of the most highly expressed genes in pregnancy recognition, implantation and
placentation. F1 crosses were more closely related to Finnsheep than to Texel, as indicated by phenotypic and
gene expression results that need to be validated with additional experiments (with F2 crosses and backcrosses).
Several genes with potential importance during early pregnancy (including *SIGLEC14*, *SIGLEC8*, *MRP4,* and
CA5A) were reported in the CL for the first time in any species. The results from this study show the importance
of the immune system during early pregnancy and may even have greater significance to high prolificacy as
revealed by significant upregulation of immune-related genes in Finnsheep. We also highlight the need for
improved annotation of the sheep genome and emphasize that our data will certainly contribute to such
improvement. Taken together, our data provide new information to aid in understanding the complex
reproductive events during the preimplantation period in sheep and may also have implications for other
ruminants (such as goats and cattle) and mammals, including humans.

Author Contributions: Conceptualization, J.K., and M.H.L; methodology, K.P., and M.W.; software, K.P.; formal analysis, K.P.; investigation, K.P.; resources, J.K., M.H., and J.P.; data curation, K.P.; writing—original draft preparation, K.P.; writing—review and editing, K.P., J.P., and J.K; visualization, K.P.; supervision, J.K., and M.H.L.; project administration, J.K., and M.H.; funding acquisition, J.K., and M.H.L. All authors have read and agreed to the published version of the manuscript.

Funding: This study was funded by the Academy of Finland (decisions 250633, 250677 and 285017). This study is part of the ClimGen ("Climate Genomics for Farm Animal Adaptation") project funded by FACCE-JPI ERA-NET Plus on Climate Smart Agriculture. K.P. acknowledges financial support from the Niemi Foundation.

Acknowledgments: We are grateful to Anu Tuomola for providing the experimental facilities on her farm in Urjala and to Kati Kaisajoki from Lallin Lammas Ltd. for providing the slaughtering facilities. We thank Johanna Rautiainen, Arja Seppälä, Annukka Numminen, Kalle Saastamoinen, Ilma Tapio, Tuula Marjatta Hamama, Anneli Virta and Magnus Andersson for their valuable assistance during this study. The authors wish to acknowledge the CSC – IT Center for Science, Finland, for computational resources. This study was supported by the Finnish Functional Genomics Centre of the University of Turku, Åbo Akademi and Biocenter Finland.

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

References

Angata, T., Margulies, E. H., Green, E. D., and Varki, A. (2004). Large-scale sequencing of
the CD33-related Siglec gene cluster in five mammalian species reveals rapid
evolution by multiple mechanisms. *Proc. Natl. Acad. Sci.* 101, 13251–13256.
doi:10.1073/pnas.0404833101.

Bai, S. X., Wang, Y. L., Qin, L., Xiao, Z. J., Herva, R., and Piao, Y. S. (2005). Dynamic
expression of matrix metalloproteinases (MMP-2, -9 and -14) and the tissue
inhibitors of MMPs (TIMP-1, -2 and -3) at the implantation site during tubal
pregnancy. *Reproduction* 129, 103–113. doi:10.1530/rep.1.00283.

Bazer, F. W. (2013). Pregnancy recognition signaling mechanisms in ruminants and pigs. *J.*
Anim. Sci. Biotechnol. doi:10.1186/2049-1891-4-23.

Bindea, G., Mlecnik, B., Hackl, H., Charoentong, P., Tosolini, M., Kirilovsky, A., et al.
(2009). ClueGO: a Cytoscape plug-in to decipher functionally grouped gene ontology
and pathway annotation networks. *Bioinforma. Oxf. Engl.* 25, 1091–3.
doi:10.1093/bioinformatics/btp101.

Bishop, C. V. (2013). Progesterone inhibition of oxytocin signaling in endometrium. *Front.*
Neurosci. doi:10.3389/fnins.2013.00138.

Blighe, K., Rana, S., and Myles, L. (2019). *EnhancedVolcano: Publication-ready volcano*
plots with enhanced colouring and labeling. Available at:
https://github.com/kevinblighe/EnhancedVolcano.

Bolet, G. (1986). "Timing and Extent of Embryonic Mortality in Pigs Sheep and Goats:
Genetic Variability," in *Embryonic Mortality in Farm Animals* (Dordrecht: Springer
Netherlands), 12–43. doi:10.1007/978-94-009-5038-2_2.

Bornhöfft, K. F., Goldammer, T., Rebl, A., and Galuska, S. P. (2018). Siglecs: A journey
through the evolution of sialic acid-binding immunoglobulin-type lectins. *Dev.*
Comp. Immunol. 86, 219–231. doi:10.1016/J.DCI.2018.05.008.

Branco, R., and Masle, J. (2019). Systemic signalling through TCTP1 controls lateral root
formation in Arabidopsis. *J. Exp. Bot.* doi:10.1093/jxb/erz204.

Brioudes, F., Thierry, A.-M., Chambrier, P., Mollereau, B., and Bendahmane, M. (2010).
Translationally controlled tumor protein is a conserved mitotic growth integrator in
animals and plants. *Proc. Natl. Acad. Sci.* 107, 16384–16389.
doi:10.1073/pnas.1007926107.

Brooks, K., Burns, G. W., Moraes, J. G. N., and Spencer, T. E. (2016). Analysis of the
Uterine Epithelial and Conceptus Transcriptome and Luminal Fluid Proteome During
the Peri-Implantation Period of Pregnancy in Sheep. *Biol. Reprod.* 95.
doi:10.1095/biolreprod.116.141945.

Cao, H., and Crocker, P. R. (2011). Evolution of CD33-related siglecs: regulating host
immune functions and escaping pathogen exploitation? *Immunology* 132, 18–26.
doi:10.1111/j.1365-2567.2010.03368.x.

Casey, O. M., Morris, D. G., Powell, R., Sreenan, J. M., and Fitzpatrick, R. (2005). Analysis
of gene expression in non-regressed and regressed bovine corpus luteum tissue using
a customized ovarian cDNA array. *Theriogenology* 64, 1963–1976.
doi:10.1016/j.theriogenology.2005.04.015.

Chen, S. H., Wu, P.-S., Chou, C.-H., Yan, Y.-T., Liu, H., Weng, S.-Y., et al. (2007). A
Knockout Mouse Approach Reveals that TCTP Functions as an Essential Factor for
Cell Proliferation and Survival in a Tissue- or Cell Type–specific Manner. *Mol. Biol.*
Cell 18, 2525–2532. doi:10.1091/mbc.e07-02-0188.

Christenson, L. K., and Devoto, L. (2003). Cholesterol transport and steroidogenesis by the
corpus luteum. *Reprod. Biol. Endocrinol. RBE* 1, 90. doi:10.1186/1477-7827-1-90.

Chu, M. X., Mu, Y. L., Fang, L., Ye, S. C., and Sun, S. H. (2007). Prolactin receptor as a
candidate gene for prolificacy of small tail Han sheep. *Anim. Biotechnol.* 18, 65–73.
doi:10.1080/10495390601090950.

Curry, T. E., and Osteen, K. G. (2001). Cyclic Changes in the Matrix Metalloproteinase
System in the Ovary and Uterus. *Biol. Reprod.* 64, 1285–1296.
doi:10.1095/biolreprod64.5.1285.

Davis, G. H., Farquhar, P. A., O'Connell, A. R., Everett-Hincks, J. M., Wishart, P. J.,
Galloway, S. M., et al. (2006). A putative autosomal gene increasing ovulation rate in
Romney sheep. *Anim. Reprod. Sci.* 92, 65–73.
doi:10.1016/j.anireprosci.2005.05.015.

Davis, G. H., Galloway, S. M., Ross, I. K., Gregan, S. M., Ward, J., Nimbkar, B. V, et al.
(2002). DNA tests in prolific sheep from eight countries provide new evidence on
origin of the Booroola (FecB) mutation. *Biol. Reprod.* 66, 1869–74.

Davis, J. S., and LaVoie, H. A. (2018). *Molecular Regulation of Progesterone Production in*
the Corpus Luteum. 3rd ed. Elsevier Inc. doi:10.1016/b978-0-12-813209-8.00015-7.

Devoto, L., Kohen, P., Gonzalez, R. R., Castro, O., Retamales, I., Vega, M., et al. (2001).
Expression of Steroidogenic Acute Regulatory Protein in the Human Corpus Luteum
throughout the Luteal Phase. *J. Clin. Endocrinol. Metab.* 86, 5633–5639.
doi:10.1210/jcem.86.11.7982.

Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., et al. (2013).
STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* 29, 15–21.
doi:10.1093/bioinformatics/bts635.

Docherty, A. J., Lyons, A., Smith, B. J., Wright, E. M., Stephens, P. E., Harris, T. J., et al.
(1985). Sequence of human tissue inhibitor of metalloproteinases and its identity to
erythroid-potentiating activity. *Nature* 318, 66–69. doi:10.1038/318066a0.

Drouilhet, L., Mansanet, C., Sarry, J., Tabet, K., Bardou, P., Woloszyn, F., et al. (2013). The
Highly Prolific Phenotype of Lacaune Sheep Is Associated with an Ectopic
Expression of the B4GALNT2 Gene within the Ovary. *PLoS Genet.* 9, e1003809.
doi:10.1371/journal.pgen.1003809.

Felix Krueger Trim Galore. Available at:
http://www.bioinformatics.babraham.ac.uk/projects/trim_galore/ [Accessed January
9, 2019].

Fleming, J. A. G. W., Spencer, T. E., Safe, S. H., and Bazer, F. W. (2006). Estrogen regulates
transcription of the ovine oxytocin receptor gene through GC-rich SP1 promoter
elements. *Endocrinology*. doi:10.1210/en.2005-1120.

Flint, A. P. F., and Sheldrick, E. L. (2004). Ovarian oxytocin and the maternal recognition of
pregnancy. *Reproduction*. doi:10.1530/jrf.0.0760831.

Forde, N., Mehta, J. P., McGettigan, P. A., Mamo, S., Bazer, F. W., Spencer, T. E., et al.
(2013). Alterations in expression of endometrial genes coding for proteins secreted
into the uterine lumen during conceptus elongation in cattle. *BMC Genomics* 14, 321.
doi:10.1186/1471-2164-14-321.

Gabler, C., Killian, G. J., and Einspanier, R. (2001). Differential expression of extracellular
matrix components in the bovine oviduct during the oestrous cycle. *Reprod. Camb.*
Engl. 122, 121–130.

Geisert, R. D., Morgan, G. L., Short, E. C., and Zavy, M. T. (1992). Endocrine events
associated with endometrial function and conceptus development in cattle. *Reprod.*
Fertil. Dev. 4, 301–5.

Gimpl, G., Wiegand, V., Burger, K., and Fahrenholz, F. (2002). Cholesterol and steroid
hormones: Modulators of oxytocin receptor function. in *Progress in Brain Research*
doi:10.1016/S0079-6123(02)39006-X.

Gray, C. A., Abbey, C. A., Beremand, P. D., Choi, Y., Farmer, J. L., Adelson, D. L., et al.
(2006). Identification of Endometrial Genes Regulated by Early Pregnancy,
Progesterone, and Interferon Tau in the Ovine Uterus. *Biol. Reprod.* 74, 383–394.
doi:10.1095/biolreprod.105.046656.

Grazzini, E., Guillon, G., Mouillac, B., and Zingg, H. H. (1998). Inhibition of oxytocin
receptor function by direct binding of progesterone. *Nature*. doi:10.1038/33176.

Hampton, A. L., Butt, A. R., Riley, S. C., and Salamonsen, L. A. (1995). Tissue inhibitors of
metalloproteinases in endometrium of ovariectomized steroid-treated ewes and
during the estrous cycle and early pregnancy. *Biol. Reprod.* 53, 302–311.
doi:10.1095/biolreprod53.2.302.

Hanrahan, J. P., Gregan, S. M., Mulsant, P., Mullen, M., Davis, G. H., Powell, R., et al.
(2004). Mutations in the genes for oocyte-derived growth factors GDF9 and BMP15
are associated with both increased ovulation rate and sterility in Cambridge and
Belclare sheep (Ovis aries). *Biol. Reprod.* 70, 900–9.
doi:10.1095/biolreprod.103.023093.

Hanrahan, J. P., and Quirke, J. F. (1984). Contribution of variation in ovulation rate and
embryo survival to within breed variation in litter size. *Genet. Reprod. Sheep Ed. RB*
Land DW Robinson.

Hansen, K. D., Wu, Z., Irizarry, R. A., and Leek, J. T. (2011). Sequencing technology does
not eliminate biological variability. *Nat. Biotechnol.* 29, 572–573.
doi:10.1038/nbt.1910.

Hernández-Montiel, W., Collí-Dula, R. C., Ramón-Ugalde, J. P., Martínez-Núñez, M. A.,
and Zamora-Bustillos, R. (2019). RNA-seq Transcriptome Analysis in Ovarian
Tissue of Pelibuey Breed to Explore the Regulation of Prolificacy. *Genes* 10, 358.
doi:10.3390/genes10050358.

Hu, J., Zhang, Z., Shen, W.-J., and Azhar, S. (2010). Cellular cholesterol delivery,
intracellular processing and utilization for biosynthesis of steroid hormones. *Nutr.*
Metab. 7, 47. doi:10.1186/1743-7075-7-47.

Hu, X., Pokharel, K., Peippo, J., Ghanem, N., Zhaboyev, I., Kantanen, J., et al. (2015).
Identification and characterization of miRNAs in the ovaries of a highly prolific
sheep breed. *Anim. Genet.*, n/a-n/a. doi:10.1111/age.12385.

Juengel, J. L., Meberg, B. M., Turzillo, A. M., Nett, T. M., and Niswender, G. D. (1995).
Hormonal regulation of messenger ribonucleic acid encoding steroidogenic acute
regulatory protein in ovine corpora lutea. *Endocrinology* 136, 5423–5429.
doi:10.1210/endo.136.12.7588291.

Kendrick, Keith. M. (2000). Oxytocin, motherhood and bonding. *Exp. Physiol.* 85,
111s–124s. doi:10.1111/j.1469-445x.2000.tb00014.x.

Kfir, S., Basavaraja, R., Wigoda, N., Ben-Dor, S., Orr, I., and Meidan, R. (2018). Genomic
profiling of bovine corpus luteum maturation. *PLOS ONE* 13, e0194456.
doi:10.1371/journal.pone.0194456.

King, S. R., and LaVoie, H. A. (2009). "Regulation of the Early Steps in Gonadal
542   Steroidogenesis," in *Reproductive Endocrinology* (Boston, MA: Springer US),
543   175–193. doi:10.1007/978-0-387-88186-7_16.

544   Kliem, H., Welter, H., Kraetzl, W. D., Steffl, M., Meyer, H. H. D., Schams, D., et al. (2007).
545   Expression and localisation of extracellular matrix degrading proteases and their
546   inhibitors during the oestrous cycle and after induced luteolysis in the bovine corpus
547   luteum. *Reprod. Camb. Engl.* 134, 535–547. doi:10.1530/REP-06-0172.

548   Lacroix-Pépin, N., Danyod, G., Krishnaswamy, N., Mondal, S., Rong, P.-M., Chapdelaine,
549   P., et al. (2011). The Multidrug Resistance-Associated Protein 4 (MRP4) Appears as
550   a Functional Carrier of Prostaglandins Regulated by Oxytocin in the Bovine
551   Endometrium. *Endocrinology* 152, 4993–5004. doi:10.1210/en.2011-1406.

552   Li, S., Chen, X., Ding, Y., Liu, X., Wang, Y., and He, J. (2011). Expression of translationally
553   controlled tumor protein (TCTP) in the uterus of mice of early pregnancy and its
554   possible significance during embryo implantation. *Hum. Reprod.* 26, 2972–2980.
555   doi:10.1093/humrep/der275.

556   Librach, C. L., Werb, Z., Fitzgerald, M. L., Chiu, K., Corwin, N. M., Esteves, R. A., et al.
557   (1991). 92-kD type IV collagenase mediates invasion of human cytotrophoblasts. *J.*
558   *Cell Biol.* 113, 437–449. doi:10.1083/jcb.113.2.437.

559   Loeser, R. F., and Wallin, R. (1992). Cell adhesion to matrix Gla protein and its inhibition by
560   an Arg-Gly-Asp-containing peptide. *J. Biol. Chem.* 267, 9459–62.

561   Love, M. I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and
562   dispersion for RNA-seq data with DESeq2. *Genome Biol.* 15, 550.
563   doi:10.1186/s13059-014-0550-8.

564   Mamo, S., Mehta, J. P., Forde, N., McGettigan, P., and Lonergan, P. (2012).
565   Conceptus-Endometrium Crosstalk During Maternal Recognition of Pregnancy in
566   Cattle. *Biol. Reprod.* 87, 6, 1–9. doi:10.1095/biolreprod.112.099945.

567   Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing
568   reads. *EMBnet.journal* 17, 10–12.

569   Moore, S. G., Pryce, J. E., Hayes, B. J., Chamberlain, A. J., Kemper, K. E., Berry, D. P., et al.
570   (2016). Differentially Expressed Genes in Endometrium and Corpus Luteum of
571   Holstein Cows Selected for High and Low Fertility Are Enriched for Sequence
572   Variants Associated with Fertility1. *Biol. Reprod.* 94.
573   doi:10.1095/biolreprod.115.132951.

574   Moses, M. A., Sudhalter, J., and Langer, R. (1990). Identification of an inhibitor of
575   neovascularization from cartilage. *Science* 248, 1408–1410.
576   doi:10.1126/science.1694043.

Mullen, M. P., and Hanrahan, J. P. (2014). Direct evidence on the contribution of a missense
mutation in GDF9 to variation in ovulation rate of Finnsheep. *PloS One* 9, e95251.
doi:10.1371/journal.pone.0095251.

Mulsant, P., Lecerf, F., Fabre, S., Schibler, L., Monget, P., Lanneluc, I., et al. (2001).
Mutation in bone morphogenetic protein receptor-IB is associated with increased
ovulation rate in Booroola Mérino ewes. *Proc. Natl. Acad. Sci. U. S. A.* 98, 5104–9.
doi:10.1073/pnas.091577598.

Nakanishi, Y., Sugiura, F., Kishi, J., and Hayakawa, T. (1986). Collagenase inhibitor
stimulates cleft formation during early morphogenesis of mouse salivary gland. *Dev.*
Biol. 113, 201–206. doi:10.1016/0012-1606(86)90122-3.

Nissi, R., Talvensaari-Mattila, A., Kotila, V., Niinimäki, M., Järvelä, I., and
Turpeenniemi-Hujanen, T. (2013). Circulating matrix metalloproteinase MMP-9 and
MMP-2/TIMP-2 complex are associated with spontaneous early pregnancy failure.
Reprod. Biol. Endocrinol. RBE 11, 2. doi:10.1186/1477-7827-11-2.

Patro, R., Duggal, G., Love, M. I., Irizarry, R. A., and Kingsford, C. (2017). Salmon: fast and
bias-aware quantification of transcript expression using dual-phase inference. *Nat.*
Methods 14, 417–419. doi:10.1038/nmeth.4197.

Pillai, S., Netravali, I. A., Cariappa, A., and Mattoo, H. (2012). Siglecs and immune
regulation. *Annu. Rev. Immunol.* 30, 357–92.
doi:10.1146/annurev-immunol-020711-075018.

Plant, T. M., Zeleznik, A. J., and Auchus, R. J. (2015). "Chapter 8 – Human Steroid
Biosynthesis," in *Knobil and Neill's Physiology of Reproduction*, 295–312.
doi:10.1016/B978-0-12-397175-3.00008-9.

Pokharel, K., Peippo, J., Honkatukia, M., Seppälä, A., Rautiainen, J., Ghanem, N., et al.
(2018). Integrated ovarian mRNA and miRNA transcriptome profiling characterizes
the genetic basis of prolificacy traits in sheep (Ovis aries). *BMC Genomics* 19.
doi:10.1186/s12864-017-4400-4.

Quinlivan, T. D., Martin, C. A., Taylor, W. B., and Cairney, I. M. (1966). Estimates of pre-
and perinatal mortality in the New Zealand Romney Marsh ewe. I. Pre- and perinatal
mortality in those ewes that conceived to one service. *J. Reprod. Fertil.* 11, 379–90.

Raheem, K. A. (2017). An insight into maternal recognition of pregnancy in mammalian
species. *J. Saudi Soc. Agric. Sci.* 16, 1–6. doi:10.1016/j.jssas.2015.01.002.

Rhind, S. M., Robinson, J. J., Fraser, C., and McHattie, I. (1980). Ovulation and embryo
survival rates and plasma progesterone concentrations of prolific ewes treated with
PMSG. *J. Reprod. Fertil.* 58, 139–144. doi:10.1530/jrf.0.0580139.

Rickard, J. P., Ryan, G., Hall, E., Graaf, S. P. de, and Hermes, R. (2017). Using transrectal
ultrasound to examine the effect of exogenous progesterone on early embryonic loss
in sheep. *PLOS ONE* 12, e0183659. doi:10.1371/journal.pone.0183659.

Ricordeau, G., Thimonier, J., Poivey, J. P., Driancourt, M. A., Hochereau-De-Reviers, M. T.,
and Tchamitchian, L. (1990). I.N.R.A. research on the Romanov sheep breed in
France: A review. *Livest. Prod. Sci.* 24, 305–332.
doi:10.1016/0301-6226(90)90009-U.

Shannon, P., Markiel, A., Ozier, O., Baliga, N. S., Wang, J. T., Ramage, D., et al. (2003).
Cytoscape: a software environment for integrated models of biomolecular interaction
networks. *Genome Res.* 13, 2498–504. doi:10.1101/gr.1239303.

Silva, C. L. A. D., Brand, H. van den, Laurenssen, B. F. A., Broekhuijse, M. L. W. J., Knol,
E. F., Kemp, B., et al. (2016). Relationships between ovulation rate and embryonic
and placental characteristics in multiparous sows at 35 days of pregnancy. *animal* 10,
1192–1199. doi:10.1017/S175173111600015X.

Simon Andrews FastQC A Quality Control tool for High Throughput Sequence Data.
Available at: http://www.bioinformatics.babraham.ac.uk/projects/fastqc/ [Accessed
January 18, 2013].

Soneson, C., Love, M. I., and Robinson, M. D. (2016). Differential analyses for RNA-seq:
transcript-level estimates improve gene-level inferences. *F1000Research* 4, 1521.
doi:10.12688/f1000research.7563.2.

Souza, C. J., MacDougall, C., MacDougall, C., Campbell, B. K., McNeilly, A. S., and Baird,
D. T. (2001). The Booroola (FecB) phenotype is associated with a mutation in the
bone morphogenetic receptor type 1 B (BMPR1B) gene. *J. Endocrinol.* 169, R1-6.

Spencer, T. E., Burghardt, R. C., Johnson, G. A., and Bazer, F. W. (2004a). Conceptus
signals for establishment and maintenance of pregnancy. *Anim. Reprod. Sci.* 82–83,
537–550. doi:10.1016/j.anireprosci.2004.04.014.

Spencer, T. E., Johnson, G. A., Bazer, F. W., and Burghardt, R. C. (2004b). Implantation
mechanisms: insights from the sheep. *Reprod. Camb. Engl.* 128, 657–68.
doi:10.1530/rep.1.00398.

Spencer, T. E., Johnson, G. A., Bazer, F. W., Burghardt, R. C., and Palmarini, M. (2007).
Pregnancy recognition and conceptus implantation in domestic ruminants: roles of
progesterone, interferons and endogenous retroviruses. *Reprod. Fertil. Dev.* 19,
65–78.

Spencer, T. E., Stagg, A. G., Joyce, M. M., Jenster, G., Wood, C. G., Bazer, F. W., et al.
(1999). Discovery and Characterization of Endometrial Epithelial Messenger

Ribonucleic Acids Using the Ovine Uterine Gland Knockout Model [1]. *Endocrinology*
140, 4070–4080. doi:10.1210/endo.140.9.6981.

Stocco, D. M. (2000). The role of the StAR protein in steroidogenesis: challenges for the
future. *J. Endocrinol.* 164, 247–53.

Stouffer, R. L., and Hennebold, J. D. (2015). "Structure, Function, and Regulation of the
Corpus Luteum," in *Knobil and Neill's Physiology of Reproduction*, 1023–1076.
doi:10.1016/B978-0-12-397175-3.00023-5.

Tuynder, M., Susini, L., Prieur, S., Besse, S., Fiucci, G., Amson, R., et al. (2002). Biological
models and genes of tumor reversion: cellular reprogramming through tpt1/TCTP
and SIAH-1. *Proc. Natl. Acad. Sci. U. S. A.* 99, 14976–81.
doi:10.1073/pnas.222470799.

Våge, D. I., Husdal, M., Kent, M. P., Klemetsdal, G., and Boman, I. a (2013). A missense
mutation in growth differentiation factor 9 (GDF9) is strongly associated with litter
size in sheep. *BMC Genet.* 14, 1. doi:10.1186/1471-2156-14-1.

Varki, A., and Angata, T. (2006). Siglecs—the major subfamily of I-type lectins.
Glycobiology 16, 1R-27R. doi:10.1093/glycob/cwj008.

Xu, S.-S., Gao, L., Xie, X.-L., Ren, Y.-L., Shen, Z.-Q., Wang, F., et al. (2018).
Genome-Wide Association Analyses Highlight the Potential for Different Genetic
Mechanisms for Litter Size Among Sheep Breeds. *Front. Genet.* 9, 118.
doi:10.3389/fgene.2018.00118.

Zhao, J., and Nishimoto, S. K. (1996). Matrix Gla protein gene expression is elevated during
postnatal development. *Matrix Biol. J. Int. Soc. Matrix Biol.* 15, 131–40.

Zhao, J., and Warburton, D. (1997). Matrix Gla protein gene expression is induced by
transforming growth factor-beta in embryonic lung culture. *Am. J. Physiol.-Lung
Cell. Mol. Physiol.* 273, L282–L287. doi:10.1152/ajplung.1997.273.1.L282.

672