Somatic aging pathways regulate reproductive plasticity resulting from early life starvation in *Caenorhabditis elegans*

Maria C. Ow, Alexandra M. Nichitean, and Sarah E. Hall[1]

Department of Biology, Syracuse University, Syracuse, NY 13244 USA

[1]Corresponding author: shall@syr.edu

Abstract

Adverse experiences early in the life of an animal often result in changes in gene

expression that are followed by changes in their adult phenotype. In the nematode *C.*

elegans, starvation during the early larval stage promotes the entry into a non-feeding,

stress-resistant stage where they remain as dauer larvae until environmental conditions

improve. *C. elegans* adults that have experienced dauer (postdauers) retain a cellular

memory of early life starvation that results in global changes in gene expression and

decreased fertility. Here we show that the endocrine pathways attributed to the germline

signaling regulation of somatic aging in *C. elegans* adults lacking a functional germline

also regulate the reproductive phenotypes of adults that have experienced dauer as a

result of early life starvation. This is in part due to the modulation of fatty acid metabolic

pathways that serve to re-allocate fat reserves in postdauer adults in a manner

benefitting the progeny at the expense of the parental somatic fat reservoir. Our results

also show that the metabolic plasticity in postdauer animals and the inheritance of

ancestral starvation memory is a result of crosstalk between somatic and reproductive

tissues mediated by the HRDE-1 nuclear RNAi Argonaute.

Introduction

Evidence indicating that experiences during early development affect behavior and physiology in a stress-specific manner later in life is abundant throughout the animal kingdom (Telang and Wells, 2004; Weaver et al. 2004; Binder et al. 2008; Pellegroms et al. 2009; van Abeelen et al. 2012; Zhao et al. 2014; Canario et al. 2017; Dantzer et al. 2019; Vitikainen et al. 2019). In humans, individuals that experienced trauma early in infancy or childhood are adversely affected by stress-related psychiatric or chronic illness later in their lifetime (Danese and McEwen, 2012; Ehlert 2013; Vaiserman 2015; Teicher and Samson, 2016). In addition, epidemiological studies and studies using mammalian animal models have supported the "thrifty" phenotype hypothesis which proposes that fetal or postnatal malnutrition results in the increased risk for metabolic disorders in the offspring (Neel, 1962; Hales and Barker, 1992; Vaag et al. 2012; Smith and Ryckman, 2015). For instance, individuals exposed to the WWII Dutch Winter Hunger during gestation had lower glucose tolerance and increased risk of obesity, diabetes, and cardiovascular diseases in adulthood compared to siblings born before the famine (Painter et al. 2008; Lumey et al. 2011; Veenendaal et al. 2013). Genome-scale analyses have revealed that individuals exposed pre-natally to the Dutch famine had differential DNA methylation signatures in genes with functions related to growth and lipid metabolism (Tobi et al. 2014). While some studies have been conducted to understand the cellular mechanism by which stress early in life affect adult phenotypes in animals (Weaver et al. 2004, Klengel et al. 2013), the process by which stress perception is translated into tissue specific changes in gene expression remains largely unknown. Thus, understanding the relationship between fetal malnutrition and

the increased susceptibility to metabolic disorders in adulthood is critical for

development of intervention and therapeutic strategies.

The nematode *C. elegans* makes a critical decision regarding its developmental

trajectory based on the environmental conditions experienced during its first larval stage

(L1). If conditions are poor (e.g. low food availability, crowding, or high temperatures),

decreased insulin and TGF-β signaling in L1 larvae promote entry into an alternative,

stress-resistant, non-aging, diapause stage named dauer where they can survive for

months. Once conditions improve, dauer larvae resume development as postdauer L4

larvae and continue through reproductive adulthood as postdauer adults (Cassada and

Russell, 1975). Alternatively, if conditions are favorable, L1 larvae proceed through

additional larval stages (L2-L4) until reaching reproductive adulthood (control adults)

(Sulston and Horvitz, 1977). Although postdauer adults are morphologically similar to

control adults, we previously showed that postdauer adults retained a cellular memory

of their early life experience that resulted in genome-wide changes in their chromatin,

transcriptome, and life history traits (Hall et al. 2010, 2013; Ow et al. 2018).

Remarkably, postdauer adults also encoded the nature of their L1 environmental

experience and gauged their adult reproductive phenotypes and genome-wide gene

expression based on this memory (Ow et al. 2018). Postdauer adults that experienced

crowding or high pheromone conditions (PD$_{Phe}$) exhibited increased fertility and

upregulated expression of genes involved in reproduction relative to control adults that

never experienced crowding (CON$_{Phe}$). In contrast, postdauer adults that experienced

starvation (PD$_{Stv}$) exhibited decreased fertility and an enrichment in somatic gene

expression compared to control adults that never experienced starvation (CON$_{Stv}$) (Ow

et al. 2018). Moreover, the reproductive and gene expression plasticity observed in

PD_{Phe} and PD_{Stv} with respect to their control adults is dependent on signaling between

somatic and reproductive tissues by distinct mechanisms: systemic RNAi signals (for

PD_{Phe}) or a functional germ line (for PD_{Stv}) (Ow et al. 2018).

The requirement of crosstalk between somatic and reproductive tissues for the

reproductive phenotypes is also a key regulatory feature governing adult lifespan and

stress response (Kenyon 2010, 2010b). In *C. elegans*, endocrine signaling has emerged

as one of the principal pathways extending the lifespan of animals lacking a germ line

due to either ablation of germ line precursor cells or a mutation in the *glp-1*/Notch

receptor gene. The two main effectors of endocrine signaling, the FOXO transcription

factor DAF-16 and the nuclear hormone receptor (NHR) DAF-12, are required for the

increased lifespan of germline-less animals (Hsin and Kenyon, 1999; Kenyon 2010,

2010b; Murphy and Hu, 2013). The nuclear localization of DAF-16 is governed by its

phosphorylation state in response to the physiological state of the animal. When an

animal experiences reproductive stress, including sterility, DAF-16 is dephosphorylated

and translocated to the nucleus where it can effect changes in target gene expression to

promote the extended longevity of germ line-less animals (Kenyon 2010, 2010b;

Murphy and Hu, 2013). The activity of KRI-1 (an ortholog of the human intestinal

ankyrin-repeat protein KRIT1/CCM1) and TCER-1 (a homolog of the human

transcription elongation factor TCERG1) are also required for DAF-16 nuclear

localization and the increased lifespan attributable to germ line loss (Berman and

Kenyon, 2006; Ghazi et al. 2009).

Nuclear hormone receptors are ligand bound receptors that recognize lipid

moieties and hormone ligands to mediate transcription of their target genes. In *C.*

elegans, 284 members of the NHR family have been identified. One of the best

characterized NHRs in *C. elegans* is DAF-12, a homolog of the mammalian vitamin D

receptor. Under favorable environmental conditions, DAF-12 binds to bile acid-like

steroid ligands (dafachronic acids or DA) to promote the expression of genes involved in

reproduction and growth. Under unfavorable conditions lacking DA, DAF-12 binds to its

co-repressor, DIN-1, to promote the expression of genes for dauer entry (Antebi 2014).

Additional NHRs contribute to the increased adult lifespan in germ line-less animals by

modulating the fatty acid metabolic status of an animal. Most of *C. elegans* NHRs are

derived from a common ancestor, the hepatocyte nuclear factor 4 alpha (HNF4α) that

acts as a lipid sensor in mammals (Antebi 2014). Of this large family of *C. elegans*

NHRs, NHR-49 and NHR-80 function as regulators of fatty acid metabolism by

promoting fatty acid desaturation in long-lived germline-less animals (Van Gilst et al.

2005a; Goudeau et al. 2011; Pathare et al. 2012; Ratnappan et al. 2014). Three genes

whose expression are promoted by NHR-49 and NHR-80 are the *fat-5*, *fat*-6, and *fat-7*

delta-9 (Δ9) desaturase genes, which carry out the first catalytic step in the production

of polyunsaturated fatty acids (PUFAs) that serve as energy reserves, membrane

building blocks, and signaling molecules (Brock et al. 2006; Watts and Ristow, 2017).

In this study, we show that steroid hormone signaling, reproductive longevity

signaling, and nuclear hormone receptors contribute to the decreased fertility of

postdauer animals that experienced early life starvation by regulating fatty acid

metabolism. The reproductive plasticity of starvation-induced postdauers is a result of

crosstalk between somatic and reproductive tissues, the effect of which is an increase in

lipid metabolic pathway function in an animal that has experienced dauer, resulting in

decreased fertility but endowing metabolic robustness to their progeny. We also show

that the signals between the soma and the germ line are dependent on HRDE-1, a

germline-specific nuclear RNAi Argonaute. Given the principal role of HRDE-1 in RNAi-

mediated transgenerational inheritance, this pathway may transmit the ancestral

starvation memory to an ensuing generation and provide the necessary hardiness to

survive future famine.

Results

GLP-4 is required globally to affect reproductive plasticity resulting from early life

starvation

We have previously shown that postdauer adults exhibit gene expression and

reproductive phenotypes that reflect their environmental and developmental history. In

adults that experienced starvation-induced dauer formation (PD_{Stv}), we showed that

gene expression changes of somatically-expressed seesaw genes and their

characteristic reduced brood size relative to control adults (CON_{Stv}) required a functional

glp-4 gene (Ow et al. 2018). The *glp-4* gene encodes a valyl aminoacyl tRNA

synthetase that is expressed in the intestine, somatic gonad, and the germ line (Rastogi

et al. 2015). The temperature-sensitive *bn2* allele is a partial loss-of-function lesion that

likely results in decreased translation in both somatic and germline tissues (Beanan and

Strome 1992; Rastogi et al. 2015). *glp-4(bn2)* mutants exhibit increased adult lifespan

and stress resistance that require crosstalk between the germ line and the soma via

endocrine signaling pathways (Arantes-Oliveira et al. 2002).

To address whether the expression of *glp-4* in the intestine, somatic gonad, or

the germ line affect reproductive plasticity between control and PD_{Stv} adults, we

constructed single copy rescuing transgenes with tissue specific promoters using Mos1-

mediated single copy insertion (MosSCI). We performed brood size assays of control

and PD_{Stv} adults at 15°C, the permissive temperature at which *glp-4(bn2)* is fertile.

While the brood size of PD_{Stv} wild-type N2 adults was decreased to near significance (*p*

= 0.0525) at the low temperature, the brood size of *glp-4(bn2)* PD_{Stv} adults increased

significantly, consistent with our previous observation (Supplemental Fig. S1;

Supplemental Table S11) (Ow et al. 2018). Expression of *glp-4* under its endogenous

promoter partially rescued the increased brood size of PD_{Stv} *glp-4* adults (Supplemental

Fig. S1; Supplemental Table S11). However, limited expression of *glp-4* in the intestine,

somatic gonad, or the germ line resulted in a brood size phenotype similar to that of *glp-*

4(bn2) mutants (Supplemental Fig. S1; Supplemental Table S11). These results

suggest that the contribution of GLP-4 to the reproductive plasticity of PD_{Stv} adults is not

due to its function in a singular tissue (germ line, somatic gonad or intestine) but rather

from multiple locations to promote inter-tissue crosstalk.

Dafachronic acid-dependent DAF-12 signaling is required for decreased fecundity after

starvation-induced dauer formation

Given that endocrine signaling across tissues is a prominent feature of reproductive

longevity, we examined whether wild-type PD_{Stv} adults shared any gene expression

signatures with animals lacking a functional germ line. In *glp-1* mutants, increased

longevity is dependent on TOR (target of rapamycin) signaling, DAF-16/FOXO gene

regulation, steroid hormone signaling, and fatty acid metabolism regulation (Lapierre

and Hansen, 2012). With the exception of TOR signaling, we found significant gene

expression changes in PD_{Stv} adults of key genes in each of these regulatory pathways.

In the steroid signaling pathway, dafachronic acid (DA) biosynthesis requires the

cytochrome P450 DAF-9, the Reiske-like oxygenase DAF-36, and the hydroxysteroid

dehydrogenase HSD-1 (Gerisch et al. 2004; Rottiers et al. 2006; Patel et al. 2008;

Mahanti et al. 2014). In animals lacking a functional germ line, the levels of *daf-36*

mRNA and the Δ^7 form of DA (Δ^7-DA) are significantly increased compared to wild-type

(Shen et al. 2012). Interestingly, we found that in wild-type PD_{Stv} adults with a germ line,

daf-36 mRNA is also increased 3-fold (p = 3.25e-04; FDR = 0.01) compared control

adults (Ow et al. 2018). To investigate whether DA signaling plays a role in mediating

reproductive plasticity as a result of early life experience, we asked whether mutations

in DA biosynthesis genes would affect the reduced brood size observed in PD_{Stv} adults.

We found that *daf-9(rh50)* and *daf-36(k114),* but not *hsd-1(mg433),* exhibited a

significant increase in brood size in PD_{Stv} adults, opposite of what we observed in wild-

type animals (Fig. 1A; Supplemental Table S1).

We asked whether the steroid signaling that contributed to the reduced fertility of

PD_{Stv} adults acted through related NHRs, DAF-12 and NHR-8. Since null mutants of

daf-12 are dauer defective, we used two *daf-12* alleles, *rh284* (Class 5 mutant) and

rh285 (Class 4 mutant), with lesions in the ligand binding domain (LBD) that affect

steroid signaling activity but otherwise have nearly normal dauer phenotype (Antebi et

al. 2000). Surprisingly, *daf-12(rh284)* and *daf-12(rh285)* exhibited a significant increase

in brood size, similar to that observed for the DA biosynthesis mutants (Fig. 1A;

Supplemental Table S1). These results suggest that DA-dependent activity of DAF-12 is

required for reduced fertility in PD$_{Stv}$ adults. The gene *nhr-8* is also up-regulated 2.5-fold

in PD$_{Stv}$ adults (p = 7.28e-03; FDR = 0.15); thus, we asked whether a mutant disrupting

the LBD of NHR-8, would similarly affect the fecundity of PD$_{Stv}$ adults (Lindblom et al.

2001; Magner et al. 2013; Ow et al. 2018). However, *nhr-8(ok186)* exhibited a

significant brood size reduction between control and PD$_{Stv}$ animals at levels comparable

to wild-type N2 (Fig. 1A; Supplemental Table S1). These results indicate that the

reduced fertility of PD$_{Stv}$ adults is dependent on DA biosynthesis through DAF-9 and

DAF-36 and the subsequent DA-binding activity of DAF-12.

We next asked whether the exogenous addition of DA was sufficient to reduce

the brood size of PD$_{Stv}$ adults. To test this hypothesis, we induced larva into dauer by

starvation with or without exogenously added 40 nM of Δ^7-DA and measured their brood

size. The addition of Δ^7-DA did not affect the fertility of wild-type and *daf-36(k114)* (Fig.

1B; Supplemental Table S2). However, *daf-9(rh50)* PD$_{Stv}$ adults exhibited a significant

decrease in brood size in the presence of exogenous Δ^7-DA (Fig. 1B; Supplemental

Table S2). DAF-9 is reported to promote a feedback loop of DA production between the

neuroendocrine XXX cells and the hypodermis; thus, *daf-9* mutants may be more

sensitive to small changes in DA concentration (Antebi 2014). Taken together, these

results suggest that increased DA-dependent DAF-12 signaling is necessary, but

perhaps not sufficient, for the reduced brood size phenotype of PD$_{Stv}$ adults.

The DAF-16 reproductive longevity pathway mediates reproductive plasticity

DAF-16 acts as a major effector of reduced insulin/IGF-1-like signaling (IIS) by

promoting the expression of a group of stress response genes (Class I targets) (Tepper

et al. 2013). PQM-1, a C2H2 zinc finger transcription factor, acts in a mutually

antagonistic manner to DAF-16 by up-regulating genes associated with growth and

reproduction (Class II targets), as its cellular localization in the nucleus is anti-correlated

to that of DAF-16 (Tepper et al. 2013). Interestingly, we previously found that the

expression of *pqm-1* is up-regulated 3-fold (p = 1.14e-04; FDR = 5.75e-03) in PD_{Stv}

adults compared to control adults (Ow et al. 2018). In addition, we observed that

18.82% (313 genes; p < 2.52e-91, hypergeometric test) of the reported 1,663 DAF-16

Class I target genes and 13.21% (229 genes; p < 1.96e-37) of the 1,733 PQM-1 Class II

target genes significantly overlapped with the 1,121 somatically-enriched genes that we

previously reported to be up-regulated in PD_{Stv} adults. We also observed significant

overlaps between our previously reported 551 down-regulated genes in PD_{Stv} adults

with DAF-16 Class I genes (0.54%; 9 genes; p < 1.03e-11, hypergeometric test) and

PQM-1 Class II genes (4.79%; 83 genes; p < 2.83e-07) (Tepper et al. 2013; Ow et al.

2018) (Fig. 2A; Supplemental Table S3). In addition, *daf-18*, the functional ortholog of

the human PTEN tumor suppressor gene that promotes the nuclear localization of DAF-

16 (Ogg and Ruvkun, 1998; Gil et al. 1999; Mihaylova et al. 1999; Solari et al. 2005),

was down-regulated 2.8-fold (p = 6.6e-06; FDR = 5.72e-04) in PD_{Stv} adults compared to

control adults (Ow et al. 2018). The overlap in gene targets and the significant changes

in the expression of *pqm-1* and *daf-18* prompted us to ask whether PQM-1 and DAF-18

contribute to the reduced fertility in PD_{Stv} adults by altering the regulation of genes

involved in reproduction and sequestering DAF-16 in the cytoplasm. We investigated

this hypothesis by performing brood assays on adults carrying mutations in *pqm-1* and

daf-18. Surprisingly, the brood size of control and PD$_{Stv}$ adults in *pqm-1(ok485)* and *daf-*

18(e1375) strains was similar to wild type, indicating that gene regulation by PQM-1 is

unlikely to contribute to the PD$_{Stv}$ decreased brood size phenotype (Fig. 2B;

Supplemental Table S4). Since *daf-18(e1375)* is a hypomorph, we next tested the

possibility that DAF-16 localization may play a role in regulating postdauer reproduction.

daf-16 null mutants are dauer defective; thus, we asked whether a *daf-16(mu86)* null

mutant expressing a rescuing transgene (*daf-16a^{AM}::gfp*) that constitutively localizes

DAF-16 to the nucleus would show a brood size plasticity phenotype (Lin et al. 2001).

Similar to what was observed for *daf-18(e1375)*, the *daf-16(mu86); daf-16a^{AM}::gfp*

transgenic strain displayed a reduced brood size in PD$_{Stv}$ similar to wild-type N2 (Fig.

2B; Supplemental Table S4). These results suggest that the nuclear localization of DAF-

16, and not PQM-1, may be contributing to the diminished fertility phenotype of PD$_{Stv}$

adults.

To further investigate whether the nuclear localization of DAF-16 is required for

the reproductive plasticity between control and PD$_{Stv}$ animals, we measured the brood

size of animals with mutations in genes that regulate the cellular localization of DAF-16.

First, we tested SMK-1/PPP4R3A, which promotes the nuclear localization of DAF-16

when animals are exposed to pathogenic bacteria, ultraviolet irradiation, and oxidative

stress, but does not affect the dauer development and reproductive timing activities of

DAF-16 (Wolff et al. 2006). We found that *smk-1(mn156)* mutants continued to exhibit a

decreased PD$_{Stv}$ fertility compared to controls. Secondly, we examined two genes, *kri-1*

and *tcer-1,* that are required for DAF-16 nuclear localization and extend adult lifespan in

germ line-less animals, but do not regulate dauer formation (Berman and Kenyon, 2006;

Ghazi et al. 2009). We found that decreased brood size in PD_{Stv} was abrogated in the

kri-1(ok1251) and *tcer-1(tm1452)* mutants (Fig. 2B; Supplemental Table S4). Because

TCER-1 regulates the transcription of its target genes in a DAF-16-dependent and

independent manner (Armit et al. 2016), these results indicate that KRI-1, TCER-1, and

possibly DAF-16 are required for the reproductive plasticity in PD_{Stv} animals.

Increased fatty acid metabolism promotes PD_{Stv} fertility

In animals lacking a germ line, DAF-16 and TCER-1 are required to bolster the

expression of lipid biosynthesis, storage, and hydrolysis genes to promote adult

longevity (Armit et al. 2016). Similar to germ line-less *glp-1* mutants, PD_{Stv} adults

exhibited a significant increase in the expression of fatty acid metabolic genes targeted

by DAF-16 and TCER-1 (Fig. 2C; Supplemental Table S5). Specifically, 47% (7 out of

15) of DAF-16 and TCER-1 target genes predicted to regulate lipid synthesis and

storage are up-regulated in PD_{Stv} adults ($p < 6.86e-06$, hypergeometric test). Similarly,

50% (12 out of 24) of lipid hydrolysis genes targeted by DAF-16 and TCER-1 are up-

regulated in PD_{Stv} adults ($p < 1.16e-09$) (Amrit et al. 2016; Ow et al. 2018) (Fig. 2C;

Supplemental Table S5). Additionally, 27% (4 out of 15) of lipid catabolism genes ($p <$

0.008) and 12% (9 out of 74) of lipid anabolism genes ($p < 0.021$) that were not

identified as targets of DAF-16 or TCER-1 were up-regulated in PD_{Stv} adults (Amrit et al.

2016; Ow et al. 2018). Notably, none of the genes down-regulated in PD_{Stv} adults were

represented by the DAF-16 and TCER-1 lipid metabolic target genes (Amrit et al. 2016;

Ow et al. 2018) (Fig. 2C). This observation was also true for lipid anabolism and

catabolism genes that are not targets of DAF-16 or TCER-1. These results suggest that

DAF-16 and TCER-1 play a role in regulating reproductive plasticity between controls

and PD$_{Stv}$ by up-regulating fatty acid metabolism.

To investigate this hypothesis, we asked if mutations in known regulators of lipid

metabolism would exhibit changes in brood size in PD$_{Stv}$ adults when compared to

controls. One of the genes jointly up-regulated by DAF-16 and TCER-1 is *nhr-49*, a

nuclear hormone receptor homologous to the mammalian HNF4α lipid sensing nuclear

receptor involved in the regulation of fatty acid metabolism and the oxidative stress

response (Ratnappan et al. 2014; Amrit et al. 2016; Moreno-Arriola et al. 2016; Goh et

al. 2018; Hu et al. 2018). Additional nuclear hormone receptors, NHR-80, NHR-13, and

NHR-66, and the Mediator complex subunit, MDT-15, partner with NHR-49 and co-

regulate the expression of genes involved in various aspects of lipid metabolism such

as fatty acid β-oxidation, transport, remodeling, and desaturation (Van Gilst et al. 2005a,

2005b; Taubert et al. 2006; Nomura et al. 2010; Pathare et al. 2012; Ratnappan et al.

2014; Folick et al. 2015; Amrit et al. 2016). In addition, SBP-1 (homolog of mammalian

SREBP) and NHR-49 are co-regulated by MDT-15 as part of a transcriptional network

coordinating the expression of delta-9 (Δ9) fatty acid desaturase genes (Yang et al.

2006; Taubert et al. 2006).

The reduced fertility of PD$_{Stv}$ characteristic of wild-type animals was also

observed in *nhr-80(tm1011)*, *nhr-13(gk796)*, *nhr-66(ok940)*, and *mdt-15(tm2182)*

strains, but not in *sbp-1(ep79)* (Fig. 3A; Supplemental Table S6). In addition, two *nhr-49*

loss-of-function (*lof*) mutants, *nhr-49(nr2014)* and *nhr-49(gk405)*, failed to exhibit the

decreased fertility in PD$_{Stv}$ adults compared to CON$_{Stv}$, while a third *nhr-49(ok2165) lof*

allele continued to display a PD$_{Stv}$ reduction in brood size (Fig. 3A; Supplemental Table

S6). Because the three *nhr-49* mutant alleles, *nr2014*, *gk405*, and *ok2165*, differ in the

nature and the location of their lesions (Lee et al. 2016), their biological function may

vary and hence result in phenotypic differences (*nr2041* and *gk405* vs. *ok2165*).

Animals expressing *nhr-49* gain-of-function alleles (*et7*, *et8*, and *et13*) with missense

lesions located at or near the LBD may augment NHR-49 activity resulting in a

reproductive plasticity phenotype comparable to that of wild-type N2 (Svensk et al.

2013; Lee et al. 2016) (Fig. 3A; Supplemental Table S6). Because of the interaction

between NHR-49, NHR-80, NHR-13, and NHR-66, we also examined if double and

triple mutants of these NHRs would have any reproductive plasticity phenotypes.

Double and triple mutant combinations of alleles exhibited an abolishment of the PD$_{Stv}$

decreased brood size phenotype only if they expressed the *nhr-49(nr2041)* mutation

(Fig. 3B; Supplemental Table S7). Thus far, our results indicate that NHR-49 and SBP-1

are important in the reproduction program of PD$_{Stv}$ adults, likely by up-regulating fat

metabolism genes.

NHR-49 and SBP-1 up-regulate the expression of genes involved in fatty acid

biosynthesis, including the Δ9 desaturases, *fat-5*, *fat-6*, and *fat-7*, and the delta-12 (Δ12)

desaturase *fat-2* (Nomura et al. 2010; Han et al. 2017). FAT-5, FAT-6, and FAT-7

convert saturated fatty acids (SFAs) to mono-unsaturated fatty acids (MUFAs), while

FAT-2 catalyzes the conversion of MUFAs to poly-unsaturated fatty acids (PUFAs)

(Watts and Ristow, 2017). Our previous mRNA-Seq results showed that the expression

of *fat-5*, *fat-6*, *fat-7*, and *fat-2* increased significantly between 3.8- and 26.6-fold in wild-

type PD$_{Stv}$ adults compared to controls (Ow et al. 2018). While strains expressing a

single *fat-5(tm331)*, *fat-6(tm420)*, *fat-7(wa36)*, or *fat-2(wa17)* mutation continued to

exhibit the reduced brood size phenotype in PD$_{Stv}$ adults, double mutants expressing a

mutation in *fat-6* (*fat-6(tm420); fat-5(tm331)* and *fat-6(tm420); fat-7(wa36))* no longer

displayed the reduced PD$_{Stv}$ phenotype. The double mutant *fat-7(wa36); fat-5(tm331)*

continued to display a significant fertility difference between controls and PD$_{Stv}$ (Fig. 3C;

Supplemental Table S8). These results suggest: 1) a functional redundancy between

the Δ9 fatty acid desaturases in modulating lipid homeostasis of PD$_{Stv}$ adults, with FAT-

6 playing a more principal role than FAT-5 and FAT-7; and 2) MUFAs are required for

the decreased fertility phenotype in PD$_{Stv}$ adults compared to controls.

In *C. elegans*, MUFAs are essential for viability as a triple *fat-5; fat-6; fat-7*

mutant is lethal (Brock et al. 2006). MUFAs, such as oleic acid, can be remodeled to

become PUFAs, phospholipids, and neutral lipids such as triacylglycerols (TAGs), which

serve as energy storage molecules in the intestine, hypodermis, and germ line (Watts

and Ristow, 2017). In addition to acting as key regulators of fat metabolism, FAT-5,

FAT-6, and FAT-7 are also essential in promoting the long lifespan of adult worms

lacking a germ line (Van Gilst et al. 2005a; Brock et al. 2006; Goudeau et al. 2011;

Ratnappan et al. 2014). Given that MUFAs are required for the reduced fertility of PD$_{Stv}$

adults compared to controls, we asked whether the dietary addition of oleic acid to PD$_{Stv}$

animals would further reduce their brood size such that they would produce even fewer

progeny compared to PD$_{Stv}$ animals not supplemented with oleic acid. To test this

hypothesis, we compared the brood size of PD$_{Stv}$ adults fed *E. coli* OP50 grown with

oleic acid and PD$_{Stv}$ adults whose bacterial diet was not pre-loaded with oleic acid. We

tested the N2 wild-type, *nhr-49(nr2041)*, *nhr-49(gk405)*, and *nhr-*

80(tm1011) strains, and found that for all genotypes, the supplementation of oleic acid

significantly increased the progeny number produced by PD_{Stv} compared to PD_{Stv} adults

on the control diet (Fig. 3D; Supplemental Table S9). These results indicate that oleic

acid is a limiting factor for reproduction, even for wild-type animals, after passage

through the starvation-induced dauer stage. Altogether, these results suggest that NHR-

49 and SBP-1 are critical for the decreased fertility in PD_{Stv} adults by mediating the up-

regulation of fatty acid desaturases and promoting the synthesis of sufficient levels of

oleic acid needed for reproduction after the animals experienced starvation-induced

dauer formation.

Starvation-induced postdauer adults have reduced fat stores

In long-lived *glp-1* mutants lacking a germ line, fat stores are increased relative to wild-

type (Steinbaugh et al. 2015; Amrit et al. 2016). Given that PD_{Stv} adults also have an

increased expression of lipid metabolism genes similar to *glp-1* adults, but have limited

quantities of oleic acid for reproduction, we questioned what the status of fat stores

would be in wild-type adults with an intact germ line that had passed through dauer due

to starvation. Using Oil Red O (ORO) staining, we compared the amounts of neutral

triglycerides and lipids (O'Rourke et al. 2009) in PD_{Stv} one-day-old adults and age-

matched control adults. Interestingly, PD_{Stv} adults have a significantly reduced amount

of stored lipids relative to control adults in their intestine, but still displayed ORO

staining in the embryos within the uterus (Fig. 4A, 4B). These results are consistent with

a model that PD$_{Stv}$ adults have increased expression of lipid metabolism genes in order

to produce sufficient levels of MUFAs needed for reproduction rather than lipid storage.

Next, we investigated whether the decreased lipid stores in PD$_{Stv}$ adults was

dependent upon DAF-12 and TCER-1. Both *daf-12(rh284)* and *daf-12(rh285)* mutants

displayed a significant increase in lipid storage in PD$_{Stv}$ adults relative to control adults,

opposite to what was observed in wild-type, indicating that DA-mediated DAF-12 activity

is required for regulation of fat stores in PD$_{Stv}$ adults (Fig. 4A, 4B). Interestingly, the

ORO staining positively correlated with brood sizes in wild-type N2 and the *daf-12*

mutants: fat stores and fertility decreased in wild-type PD$_{Stv}$ adults, while fat stores and

fertility increased in *daf-12* PD$_{Stv}$ adults compared to *daf-12* CON$_{Stv}$ adults (Fig. 1A;

Supplemental Table S1; Fig. 4A, 4B). In *tcer-1(tm1452)* adults, lipid staining was

diminished in PD$_{Stv}$ compared to controls, similar to what is seen in wild-type (Fig. 4A,

4B). However, since both *tcer-1* control and PD$_{Stv}$ adults have reduced fat stores

compared to their wild-type counterparts, and TCER-1 is known to positively regulate

NHR-49 and fatty acid metabolism, this result is likely due to the inability of these

animals to store fat and not because the decreased fat stores in wild-type PD$_{Stv}$ adults is

TCER-1 independent. Furthermore, the brood size of *tcer-1(tm1452)* did not positively

correlate with their ORO staining: a very modest increase ($p = 0.0546$) in PD$_{Stv}$ brood

size and a significant decrease in PD$_{Stv}$ ORO staining (Fig. 4A, 4B). In addition to

regulation of fat metabolism genes in the intestine, TCER-1 also functions redundantly

with PUF-8, a member of an evolutionarily conserved stem cell proliferation modulatory

family, to potentiate germ cell proliferation in animals undergoing continuous

development (Pushpa et al. 2013). Thus, our *tcer-1(tm1452)* results suggest that TCER-

1 activity may also be necessary in regulating germline mechanisms that curtail fertility

following exit from dauer diapause.

During *C. elegans* reproduction, intestinal fat stores are reallocated into low-

density lipoproteins (LDL)-like particles (yolk lipoproteins or vitellogenins) that are

incorporated into oocytes through receptor-mediated endocytosis in a process called

vitellogenesis, and serve as the only supply of nutrients to the developing embryos

(Kimble and Sharrock, 1983; Grant and Hirsch, 1999). Six vitellogenins homologous to

the human ApoB proteins are encoded in the *C. elegans* genome, *vit-1* through *vit-6*,

and concomitant multiple RNAi knockdown of the *vit* genes increases adult lifespan in a

process that requires NHR-49 and NHR-80 (Spieth et al. 1991; Seah et al. 2016). The

transition from dauer to reproductive adulthood is an energetically costly process that

likely drains the already reduced fat reservoirs in dauers that experienced early-life

starvation. Because vitellogenesis mobilizes intestinal fat resources for reproduction

and depletes somatic lipid stores (Kimble and Sharrock, 1983), we hypothesized that

PD_{Stv} adults have reduced fat reservoirs because they prioritize vitellogenesis as a

reproductive investment over intestinal storage. To test this hypothesis, we examined

the fat stores in control and PD_{Stv} adults following RNAi knockdown of *vit-1,* which also

results in the knockdown of *vit-2*, *vit-3*, *vit-4*, and *vit-5* due to the high sequence

homology amongst the *vit* genes (Seah et al. 2016) (Supplemental Fig. S3A). In the

RNAi treated animals, PD_{Stv} adults continued to exhibit decreased fat stores compared

to control adults, similar to the empty vector negative controls. However, PD_{Stv} adults

treated with *vit* RNAi have significantly greater intestinal fat deposits than PD_{Stv} negative

controls ($p < 0.0001$), indicating that increased vitellogenesis in PD_{Stv} adults may be one

contributing factor to the lack of stored intestinal fat (Fig. 4C, 4D; Supplemental Fig.

S3A). These results are consistent with our previous finding that oleic acid is a limiting

resource for reproduction in PD_{Stv} animals, and suggests a model where PD_{Stv} adults

utilize fat accumulated after diapause for reproduction and not storage.

Steroid signaling, reproductive longevity, and fatty acid metabolic pathways act

synergistically at different developmental time points to regulate reproductive plasticity

Thus far, our results indicate that both DA-dependent DAF-12 activity and TCER-1 are

required for the reduced fertility phenotype in PD_{Stv} adults. However, the *daf-12* and

tcer-1 PD_{Stv} mutant phenotypes are distinct, suggesting that they act in different

pathways, tissues, or developmental time points to regulate PD_{Stv} fertility. Furthermore,

the brood sizes of *kri-1(ok1251)* or *tcer-1(tm1425)* PD_{Stv} adults supplemented with

exogenous Δ^7-DA were unaffected, indicating that the effect of DA on PD_{Stv} brood size

is independent of the reproductive longevity pathway (Supplemental Fig. S2;

Supplemental Table S12). To further examine the genetic interactions of DAF-12 and

TCER-1, we performed epistasis analysis by measuring the brood sizes of control and

PD_{Stv} adults in *kri-1(ok1251); daf-12(rh284)* and *tcer-1(tm1425); daf-12(rh284)* double

mutants, and in *kri-1(ok1251); daf-12(rh285)* and *tcer-1(tm1425); daf-12(rh285)* double

mutants. For all four double mutants, we continued to observe a significantly increased

brood size in PD_{Stv} adults compared to controls, similar to the *daf-12(rh284)* and *daf-*

12(rh285) single mutants (Fig. 1A; Supplemental Table S1; Fig. 5A; Supplemental Table

S10). The combination of mutations between *kri-1(ok1251)* and *tcer-1(tm1425)* with the

daf-12(rh284) allele synergistically exacerbated the fertility phenotype of PD_{Stv} adults to

a more than 600-fold average compared to each of the single mutants, suggesting that

these pathways act in parallel (Fig. 1A; Supplemental Table S1; Fig. 5A; Supplemental

Table S10). However, the brood sizes of the *kri-1(ok1251); daf-12(rh285)* and *tcer-*

1(tm1425); daf-12(rh285) double mutants were statistically indistinguishable from the

daf-12(rh285) single mutant, instead suggesting that DAF-12 acts downstream of

TCER-1 and KRI-1 in the same pathway (Fig. 1A, Supplemental Table S1; Fig. 5A;

Supplemental Table S10). Since the *daf-12* and *tcer-1* mutant strains used in our

experiment are not null alleles, we must interpret these epistasis results cautiously.

However, we can make some conclusions with respect to the phenotypes of *daf-*

12(rh284) and *daf-12(rh285)*, which have previously been characterized for

developmental timing and dauer formation defects (Antebi et al. 1998; 2000). The *daf-*

12(rh284) mutant (P746S lesion in helix 12 of the LBD) displays delayed gonadal

development while the *daf-12(rh285)* mutant (Q707stop mutation in the LBD after helix

9) has penetrant heterochronic phenotypes that include delayed gonadal and

extragonadal developmental events (Antebi et al. 1998; 2000). In addition, our brood

assay results show that the *rh285* allele has a more severe phenotype than *rh284* in

terms of reproduction (Fig. 1A; Supplemental Table S1). These phenotypes are perhaps

due to the differences in the nature of LBD disruption that have not been characterized.

Although the epistasis results could be interpreted as DAF-12 acting in parallel to KRI-1

and TCER-1, we favor a model in which DAF-12 acts downstream of KRI-1 and TCER-1

due to the greater severity of the *daf-12(rh285)* allele over the *daf-12(rh284)* allele.

To further investigate the developmental mechanism of steroid signaling,

reproductive longevity, and fatty acid metabolism pathways in the regulation of

reproductive plasticity, we examined if DAF-12, TCER-1, and NHR-49 play a direct role

in the timing of germline proliferation in postdauer larvae. We previously demonstrated

that wild-type PD$_{Stv}$ animals delay the onset of germline proliferation compared to

control animals, resulting in a reduced brood size (Ow et al. 2018). *C. elegans* begins

germ cell proliferation after hatching and continues throughout larval development.

Germ cells at the most distal ends of the gonad undergo mitotic self-renewal while germ

cells most proximal to the gonad undergo meiosis (Kimble and Crittenden, 2005). In *C.*

elegans hermaphrodites, undifferentiated germ cells initiate spermatogenesis during the

L4 larval stage, followed by a transition to oogenesis at the adult stage (L'Hernault,

2006). Therefore, reproduction in the hermaphroditic *C. elegans* is sperm-limited (Byerly

et al. 1976; Ward and Carrel, 1979; Kimble and Ward, 1988). In previous results, when

control and PD$_{Stv}$ somatic development was synchronized using vulva morphology, we

observed significantly fewer germ cell rows in PD$_{Stv}$ larvae compared to control larvae,

correlating with fewer sperm available for self-fertilization in adulthood (Ow et al. 2018).

To determine if DAF-12, TCER-1, and NHR-49 play a direct role in germline

proliferation as a mechanism to regulate reproductive plasticity, we counted the number

of germ cell rows in control and PD$_{Stv}$ mutant larvae that were developmentally

synchronized by their somatic morphology. Because *daf-12(rh284)* and *daf-12(rh285)*

have altered germline morphologies that prevent accurate synchronization, we used the

daf-36(k114) mutant as a representative of the steroid signaling pathway. First, we

recapitulated our previous results showing that wild-type PD$_{Stv}$ larvae have significantly

fewer total germ cell rows compared to control larvae due to reduced cell rows in the

meiotic transition zone (Fig. 5B; Supplemental Fig. S4A). In contrast, *daf-36(k114)*

control and PD$_{Stv}$ larvae did not exhibit a significant difference in total (mitotic and

meiotic) cell rows, indicating that DAF-36-dependent DA is required during early

germline development for the delay in PD$_{Stv}$ germ cell proliferation (Fig. 5B;

Supplemental Fig. S4). This result is consistent with the increased brood size of PD$_{Stv}$

adults in *daf-12(rh284)* and *daf-12(rh285)* mutants, which express DAF-12 proteins

unable to bind to DA.

In contrast to the *daf-36* mutant, the *nhr-49(nr2041)* mutant exhibited a significant

decrease in the number of total germ cell rows in PD$_{Stv}$ larvae relative to control larvae,

similar to wild-type animals (Fig. 5B). Since NHR-49 is required for the reduced fertility

in PD$_{Stv}$ adults, these results are consistent with the role of NHR-49 in regulating fatty

acid metabolism in the intestine to support vitellogenesis. Like NHR-49, TCER-1 plays a

role in the intestine acting to upregulate fatty acid metabolism genes in animals lacking

germ cells, but also plays a role in the regulation of germ cell proliferation (Pushpa et al.

2013; Amrit et al. 2016). We found that a mutation in *tcer-1* resulted in a similar number

of total germ cell rows in control and PD$_{Stv}$ larvae, indicating that TCER-1 plays a role in

PD$_{Stv}$ reproductive plasticity by regulating germ cell proliferation in addition to its

intestinal role (Fig. 5B). Interestingly, *tcer-1* PD$_{Stv}$ larvae also appear to have a defect in

the onset of meiosis, since PD$_{Stv}$ larvae have significantly great number of mitotic germ

cell rows compared to controls (Supplemental Fig. S4). Together, these results indicate

that DAF-12 steroid signaling and TCER-1 act early in germline development to delay

the onset of germ cell proliferation in PD$_{Stv}$ animals, while NHR-49 functions later in

development to promote the reduced fertility of PD$_{Stv}$ adults.

Generational transmission of early life starvation memory

Our results suggest that PD_{Stv} upregulate their fatty acid metabolism to increase fat

transport to embryos. In humans, nutritional stress *in utero* not only promotes metabolic

syndrome in adulthood of the affected individuals, but also promotes obesity in

subsequent generations (Painter et al. 2008; Veenendaal et al. 2013). Therefore, to

examine whether the starvation-induced physiology of PD_{Stv} adults is inherited, we used

ORO staining to quantitate the fat storage in adult F_1 progeny of control and PD_{Stv}

adults that did not experience starvation themselves. Surprisingly, the adult F_1 progeny

of PD_{Stv} adults had an increased level of stored fat compared to F_1 progeny of control

adults (Fig. 6), but the difference was abolished by the F_2 generation. These results

indicate that the F_1 progeny of PD_{Stv} adults inherit a starvation memory that results in

metabolic reprogramming to increase their stored fat reserves.

In *C. elegans*, inheritance of gene expression states is often mediated by small

RNA pathways (Feng and Guang, 2013; Rechavi and Lev, 2017). The germline-specific,

nuclear Argonaute HRDE-1/WAGO-9 has been shown to promote transgenerational

silencing via the formation of heterochromatin at targeted genomic loci (Buckley et al.

2012; Feng and Guang, 2013; Rechavi and Lev, 2017). In particular, starvation-induced

L1 diapause has been shown to alter the physiology and small RNA populations of

subsequent generations, some of which is dependent upon HRDE-1 (Rechavi et al.

2014). We examined whether HRDE-1 was required for the generational inheritance of

starvation memory by quantifying fat storage in *hrde-1(tm1200)* mutant control and

PD_{Stv} adults and their F_1 and F_2 progeny. First, we observed that the stored fat levels

between *hrde-1* control and PD_{Stv} adults were similar, suggesting that the germline-

specific Argonaute is required for decreased intestinal fat storage of PD_{Stv} adults (Fig.

6). In addition, the ORO staining of the F_1 progeny of control and PD_{Stv} adults was also

similar, indicating that HRDE-1 is required for the generational inheritance of starvation

memory (Fig. 6). Remarkably, in the F_2 generation, the grand-progeny of *hrde*-1 PD_{Stv}

mutants displayed increased ORO staining like that observed for the F_1 generation of

wild-type adults, suggestive of a generational delay in the physiological response to

ancestral early-life starvation upon the loss of HRDE-1 activity (Fig. 6). Our genetic

evidence suggests that the germline nuclear RNAi pathway plays two roles: it

contributes to the metabolic programming of individuals that experienced early life

starvation as well as promote the generational inheritance of starvation memory in the

progeny. Our results show that *C. elegans*, like humans, inherit the disposition for

increased adiposity from parents that experienced early life starvation.

Discussion

The trade-off between reproduction and longevity has long been associated with

the notion that in the absence of reproduction, the fat stores of an animal would be

redistributed to promote somatic maintenance (Williams 1966; Kirkwood 1977;

Westendorp and Kirkwood, 1998; Partridge et al. 2005). As such, reducing or

suppressing reproduction in animals often extends lifespan, in part due to enhanced

lipogenic processes (Fowler and Partridge 1989; Tatar et al. 2001; Kenyon 2010,

2010b; Judd et al. 2011). What remains unclear is how reproduction and somatic

maintenance is balanced by developmental signals. In this study, we show that

starvation early in the life of a *C. elegans* animal with an intact reproductive system

programs its transcriptome to promote reproduction. Because dauers are non-feeding,

starvation-induced dauers have limited stored fat to survive the diapause stage and

remodel out of dauer. We propose a model whereby steroid signaling, reproductive

longevity, and fatty acid metabolic pathways are reprogrammed in animals that

experienced starvation-induced dauer in order to delay the onset of germline

proliferation and redistribute intestinal fat to developing oocytes (Fig. 7). These changes

allow PD$_{Stv}$ adults to delay reproduction until energy thresholds are met to provide

adequate levels of nutrition to fewer but albeit viable embryos at the expense of somatic

survival and maintenance. In the absence of a functional germ line, the fat that would

normally be allocated to the progeny, is channeled to nurture somatic tissues and

consequently extend adult lifespan. Thus, the mechanisms that prolong lifespan in the

absence of a functional germ line, are tantamount to the cellular programs deployed to

respond to developmental signals triggered by starvation early in life history in an

animal capable of reproduction.

The extended lifespan associated with animals missing a functional germ line is

specifically dependent on the lack of proliferating germ cells and not due to sterility

resulting from sperm, oocyte, or meiotic precursor cell deficiency (Hsin and Kenyon,

1999; Arantes-Oliveira et al. 2002). Animals without a functional germ line have up-

regulated fat metabolism pathways and exhibit increased levels of intestinal fat stores

that are associated with a longer lifespan (Kenyon 2010, 2010b; Amrit et al. 2016).

Postdauer adults that experienced early life starvation have a delay in the onset of germ

cell proliferation after dauer exit and similarly have upregulated fat metabolism

pathways in adulthood. These results indicate that the cellular mechanisms that

587 promote germline proliferation during a critical period in development also negatively

588 regulate lipid metabolism, leading to altered somatic fat accumulation and longevity

589 (Wang et al. 2008). While a number of genes and cellular components affecting

590 germline proliferation have been extensively investigated (Kimble and Crittenden,

591 2005), what might the actual signals communicating the state of proliferating germ cells

592 be that arbitrate lipid levels and the aging process? Because upregulation of DAF-36

593 and Δ^7-DA is crucial for aging extension in germline-less animals and reproductive

594 plasticity in PD_{Stv} adults, it is likely that the signal communicating the state of germ cell

595 proliferation may include dafachronic acids. Dafachronic acids mediating increased

596 longevity are produced in the somatic gonad, which includes the stem cell niche site of

597 the germ line (Yamawaki et al. 2010). Based on our results, DA is required for the delay

598 in germline proliferation (Fig. 5B), but DA-dependent DAF-12 activity is not necessary

599 for the increase in fatty acid metabolism observed in PD_{Stv} adults (Fig. 4A, 4B),

600 suggesting the possibility that dafachronic acids act in a DAF-12-independent manner to

601 regulate germ cell proliferation and fat metabolism. In addition, our results showing that

602 *hrde-1* is required for the altered fat metabolism in PD_{Stv} adults (Fig. 6) raise the

603 possibility that small RNAs could be another type of signaling molecule that

604 communicates the state of the germ line to the soma. Small RNA pathways are potent

605 surveillance mechanisms that guard against the threat of foreign genetic invaders such

606 as viruses and transposons that can compromise germline integrity (Almeida et al.

607 2019). Because germ line proliferation occurs throughout the lifetime of *C. elegans*, it is

608 likely that small RNA pathways survey the proliferative nature of the germ line at all

stages of animal development and potentially act as signaling molecules to regulate

somatic processes dependent upon the germ line.

One of the most intriguing findings of this study is that the starvation memory of

the parent is "bequeathed" to the F_1 progeny, triggering elevated levels of fat stores,

presumably as a physiological defense against ancestral famine (Fig. 6). Interestingly,

the increased intestinal fat inherited by the F_1 progeny of wild-type postdauer parents

bypassed the F_1 generation in the *hrde-1* mutant and was only evident in the F_2

generation (Fig. 6). One potential explanation is that small RNA signals are transmitted

to the F_1 generation via the HRDE-1 nuclear RNAi pathway to effect somatic

phenotypes. However, with the exception of DAF-16, none of the germline longevity

pathway genes or the vitellogenesis genes examined in this study are categorized as

direct HRDE-1 targets (Buckley et al. 2012). Given that the life stage (adulthood) at

which the HRDE-1 targets were identified is the same life stage that were used in this

study, we speculate that HRDE-1 may be indirectly targeting endocrine and

vitellogenins genes by: 1) targeting germ line genes that then affect somatic gene

expression; or 2) indirectly regulating the function of the endocrine and vitellogenin

genes by targeting a different repertoire of somatic targets. Interestingly, we find that

62% of small RNAs associated with HRDE-1 are target genes (984 out of 1587)

expressed in somatic tissues (neuronal, intestine, hypodermis, and muscle), while 31%

of HRDE-1 targets (486 out of 1587) are germline genes; and of these, 17% uniquely

overlap between HRDE-1 targets (272 out of 1587) and germline genes only (Ortiz et al.

2014; Kaletsky et al. 2016). While silencing somatic genes in the germ line is consistent

with the described function of HRDE-1, our results raise the possibility that HRDE-1

may also directly, or indirectly, target somatic genes. Accordingly, HRDE-1 is known to

contribute to the heritability of a cohort of small RNAs targeting nutrition and lipid

transporter genes that was inherited for at least three generations from populations that

experienced L1 larval arrest (Rechavi et al. 2014). In addition, HRDE-1 is required for

the repression of a group of genes activated upon multi-generational high temperature

stress that is inherited for at least two generations in the absence of the stress (Ni et al.

2016). Thus, it is not unlikely that HRDE-1 and the nuclear RNAi pathway could serve

as a signaling referees between the soma and the germ line to effect changes due to

environmental and developmental signals to perdure ancestral starvation memory.

Our study showed that PD_{Stv} adults have upregulated expression of lipid

metabolism genes as a means to load embryos with increased fat and potentially

protect progeny against future starvation conditions. Not only are fatty acids critical for

cell and organelle maintenance and act as efficient energy storage molecules but they

can regulate aging by acting as signaling molecules (e.g. as nuclear hormone receptor

ligands). Indeed, the lipid profiles in a number of species change as animals age

resulting in the acceleration of cell and tissue damage such that levels of certain lipids

can serve as aging biomarkers (Papsdorf and Brunet, 2019). Also, studies have

reported that fatty acids can alter animal physiology through the modification of

chromatin states either by the direct addition of lipid moieties to chromatin or by

activating signaling pathways that affect chromatin landscapes (Hansen et al. 2013;

Papsdorf and Brunet, 2019). Furthermore, MUFAs have been shown to be the link

between H3K4me3 modifiers and lifespan in *C. elegans*, while the inheritance of

epigenetic memory through soma-germline crosstalk in response to cellular stresses are

associated with the H3K4me3 modification complex (Tauffenberger and Parker, 2014;

Kishimoto et al. 2016; Han et al. 2017). Altogether, these studies provide an exciting

groundwork to examine how the chromatin states in PD_{Stv} animals correlate with

metabolic plasticity due to early life starvation across generations.

During the course of its natural history, *C. elegans* occupies ephemeral

environments such as rotting fruit or decomposing vegetation, where conditions and

food availability are highly unpredictable. The dauer larvae affords *C. elegans* a survival

and dispersal strategy to escape harsh environmental conditions by often associating

with passing invertebrate carriers. Accordingly, *C. elegans* is often found in the wild as

dauers. Once a food source is found, dauers resume reproductive development to

colonize the new habitat. Upon exhaustion of resources and population expansion,

young larvae enter dauer and thereby repeating the boom and bust life cycle

(Schulenburg and Félix, 2017). Because of frequent environmental perturbations, an

adopted phenotypic plasticity strategy would ensure an advantage in species survival.

The generation following a bust period would inherit the cellular programs for increased

somatic lipid stores. It is thus remarkable that the cellular mechanisms to ensure

survival of the species are fundamentally similar between humans and nematodes, two

species that have diverged hundreds of millions of years ago, once again underscoring

the relevance of a simple roundworm in understanding basic animal physiology.

Materials and methods

C. elegans *strains and husbandry*

N2 Bristol wild-type strain was used as the reference strain. Worms were grown in

Nematode Growth Medium (NGM) seeded with *Escherichia coli* OP50 using standard

methods at 20°C unless otherwise indicated (Brenner et al. 1974; Stiernagle 2006).

Mutants that were not previously backcrossed were backcrossed at least four times to

our laboratory N2 wild-type before use. Strains used in this study are listed under

Supplemental Material and Methods (Supplemental Table S14).

Transgenic strains

Single copy insertions, *pdrSi1* (*Pglp-4::glp-4 cDNA::gfp::glp-4 3'UTR*), *pdrSi2* (*Pnhx-*

2::glp-4 cDNA::gfp::glp-4 3'UTR), *pdrSi3* (*Pzfp-2 ::glp-4 cDNA::gfp::glp-4 3'UTR*), and

pdrSi4 (*Ppgl-1::glp-4 cDNA::gfp::glp-4 3'UTR*), were generated by Mos1-mediated

single copy insertion (MosSCI) (Frokjaer-Jensen et al. 2008). Information on plasmid

construction are provided in Supplemental Material and Methods. Primers sequences

are listed in Supplemental Table S15. Following integration into *unc-119(ed3)* animals

segregated from EG6699 [*ttTi5605 II; unc-119(ed3) III; oxEx1578* (*eft-3p::gfp + Cbr-unc-*

119)] that have lost the *oxEx1578* array, MosSCI insertions were genetically crossed

into the *glp-4(bn2)* background.

Brood assays

Brood assays were done as previously described using at least three biological

independent replicates (Ow et al. 2018). Statistical significance was determined using

Prism v.8 (GraphPad Software; La Jolla, California, USA).

Oil Red O staining

Fat stores were stained using Oil Red O (ORO) dye as previously described (O'Rourke

et al. 2009). Images were captured with either a Nikon Eclipse Ci with Spot 5.2 software

or with an iPhone through iDu Optics® equipped with a LabCam® adapter (New York,

NY, USA). Color images were separated into their RGB channel components and the

intensity measured on the green channel (Yen et al. 2010) using ImageJ (NIH).

RNA interference

Gravid adults were treated with hypochlorite to obtain embryos using standard methods

(Stiernagle 2006). Embryos were placed on NGM plates supplemented with 1mM IPTG

and 50 μg/ml carbenicillin seeded with a 10x concentrated bacterial culture expressing

the *k09f5.2* (*vit-1*) RNAi clone obtained from the Ahringer library (Kamath et al. 2001).

Embryos were allowed to grow until adulthood at which time they were treated again

with hypochlorite to obtain embryos. The recovered embryos were grown until day one

of adulthood and collected for ORO staining.

Dafachronic acid supplementation

The Δ^7 form of dafachronic acid (Δ^7-DA) (a kind gift from Frank Schroeder) was added

to a freshly grown culture of *E. coli* OP50 at a concentration of 40 nM and immediately

seeded onto NGM plates. NGM plates supplemented with an equivalent volume of

ethanol (the Δ^7-DA solvent) to those of the Δ^7-DA-supplemented NGM plates were used

as the control plates. Seeded plates were allowed to dry overnight before use.

Oleic acid supplementation

Animals were induced into dauer by starvation on peptone-less NGM plates seeded

with *E. coli* OP50 pre-loaded with oleic acid (NuChek Prep, Inc.; Elysian, Minnesota,

USA) as described by Devkota et al. 2017. Briefly, OP50 was grown overnight at 37°C

in liquid LB supplemented with 600 μM of oleic acid or with an equivalent volume of

ethanol (the oleic acid solvent) to serve as the control. Cultures were pelleted and

washed several times with M9 buffer (Stiernagle, 2006) and resuspended at a 10x

concentration. The 10x OP50 was seeded onto peptone-less NGM plates and allowed

to dry overnight before use. Brood assays were performed as described before (Ow et

al. 2018).

Germ cell rows

All worm strains used for germ cell row counting have the integrated transgene *gld-*

1::*gfp*::*3xFLAG* from the BS1080 strain in their genetic background. Germ cell row

assays were done as previously described (Ow et al. 2018). Worms with L3 vulva

morphology were identified by using Nomarski DIC microscopy at 630x (Seydoux et al.

1993) and DAPI stained using the standard whole worm DAPI staining protocol. The

stained worms were imaged using a Leica DM5500B microscope with a Hamamatsu

camera controller C10600 ORCA-R2. When performing the germ cell row counts, the

start of the transition was identified when at least two cells in a row exhibited the

crescent-shaped nuclei morphology (Shakes et al. 2009).

Acknowledgments

We are grateful for the generous gift of dafachronic acid from Frank Schroeder and

Pooja Gubibanda for advice on its use. We thank Leszek Kotula and Angelina Regua

for access to their microscope, and Eleanor Maine and Arjumand Ghazi for thoughtful

comments on this manuscript. Strain BS1080 was kindly provided by Tim Schedl. We

thank the CGC, which is funded by the NIH Office of Research Infrastructure Programs

(P40 OD010440), for providing strains.

Author contributions: M.C.O. and S.E.H. conceived and designed experiments.

M.C.O. and A.M.N. performed experiments. M.C.O., A.M.N., and S.E.H. analyzed the

data. M.C.O. and S.E.H. wrote the manuscript.

Competing interests

The authors declare no competing interests.

References

Almeida MV, Andrade-Navarro MA, Ketting RF. 2019. Function and evolution of

nematode RNAi pathways. *RNA* **5:** 1-25.

Amrit FR, Steenkiste EM, Ratnappan R, Chen SW, McClendon TB, Kostka D, Yanowitz

J, Olsen CP, Ghazi A. 2016. DAF-16 and TCER-1 facilitate adaptation to

germline loss by restoring lipid homeostasis and repressing reproductive

physiology in *C. elegans*. *PLoS Genet* **12:** e1005788.

Antebi A, Culotti JG, Hedgecock EM. 1998. *daf-12* regulates developmental age and

the dauer alternative in *Caenorhabditis elegans*. *Development* **125:** 1191-1205.

Antebi A, Yeh WH, Tait D, Hedgecock EM, Riddle DL. 2000. *daf-12* encodes a nuclear

receptor that regulates the dauer diapause and developmental age in *C.*

elegans. *Genes Dev* **14:** 1512-1527.

Antebi A. 2014. Nuclear receptor signal transduction in *C. elegans*. in *WormBook* (ed.

The *C. elegans* research community), doi/10.1895/wormbook.1.64.2.

Arantes-Oliveira N, Apfeld J, Dillin A, Kenyon C. 2002. Regulation of life-span by

germ-line stem cells in *Caenorhabditis elegans*. *Science* **295:** 502-505.

Beanan MJ, Strome S. 1992. Characterization of a germ-line proliferation

mutation in *C. elegans*. *Development* **116:** 755-766.

Berman JR, Kenyon C. 2006. Germ-cell loss extends *C. elegans* life span through

regulation of DAF-16 by *kri-1* and lipophilic-hormone signaling. *Cell* **124:** 1055-

1068.

Bouret SG, Draper SJ, Simerly RB. 2004. Trophic action of leptin on hypothalamic

neurons that regulate feeding. *Science* **304:** 108-110.

Brenner S. 1974. The genetics of *Caenorhabditis elegans*. *Genetics* **77:** 71-94.

Brock TJ, Browse J, Watts JL. 2006. Genetic regulation of unsaturated fatty acid

composition in *C. elegans*. *PLoS Genet* **2:** e108.

Buckley BA, Burkhart KB, Gu SG, Spracklin G, Kershner A, Fritz H, Kimble J, Fire A,

Kennedy S. 2012. A nuclear Argonaute promotes multigenerational

epigenetic inheritance and germline immortality. *Nature* **489:** 447-451.

Byerly L, Cassada RC, Russell RL. 1976. Life-cycle of nematode *Caenorhabditis*

elegans. I. Wild-type growth and reproduction. *Dev Biol* **51:** 23-33.

Canario L, Lundeheim N, Bijma P. 2017. The early-life environment of a pig shapes the

phenotypes of its social partners in adulthood. *Heredity* **118:** 534-541.

Cassada RC, Russell RL. 1975. The dauerlarva, a post-embryonic developmental

variant of the nematode *Caenorhabditis elegans*. *Dev Biol* **46:** 326-342.

Cutter AD. 2008. Divergence times in *Caenorhabditis* and *Drosophila* inferred from

direct estimates of the neutral mutation rate. *Mol Biol Evol* **25:** 778-786.

Dantzer B, Dubuc C, Goncalves IB, Cram DL, Bennett NC, Ganswindt A, Heistermann

M, Duncan C, Gaynor D, Clutton-Brock TH. 2019. The development of

individual differences in cooperative behaviour: maternal glucocorticoid

hormones alter helping behaviour of offspring in wild meerkats. *Phil Trans*

R Soc B **374:** 20180117.

Devkota R, Svensk E, Ruiz M, Ståhlman M, Borén J, Pilon M. 2017. The adiponectin

receptor AdipoR2 and its *Caenorhabditis elegans* homolog PAQR-2

prevent membrane rigidification by exogenous saturated fatty acids. *PLoS*

Genet **13:** e1007004.

Feng X, Guang S. 2013. Small RNAs, RNAi and the inheritance of gene silencing in

Caenorhabditis elegans. *J Genet Genomics* **40:** 153-160.

Folick A, Oakley HD, Yu Y, Armstrong EH, Kumari M, Sanor L, Moore DD, Ortlund EA,

Zechner R, Wang MC. 2015. Lysosomal signaling molecules regulate

longevity in *Caenorhabditis elegans*. *Science* **347:** 83-86.

Fowler K, Partridge L. 1989. A cost of mating in female fruitflies. *Nature* **338:** 760-761.

Frokjaer-Jensen C, Davis MW, Hopkins CE, Newman BJ, Thummel JM, Olesen SP,

Grunnet M, Jorgensen EM. 2008. Single-copy insertion of transgenes in

Caenorhabditis elegans. *Nat Genet* **40:** 1375-1383**.**

Gerisch B, Antebi A. 2004. Hormonal signals produced by DAF-9/cytochrome P450

regulate *C. elegans* dauer diapause in response to environmental cues.

Development **131:** 1765-1776.

Ghazi A, Henis-Korenblit S, Kenyon C. 2009. A transcription elongation factor that links

signals from the reproductive system to lifespan extension in *Caenorhabditis*

elegans. *PLoS Genet* **5:** e1000639.

Gil EB, Malone Link E, Liu LX, Johnson CD, Lees JA.1999. Regulation of the insulin-like

developmental pathway of *Caenorhabditis elegans* by a homolog of the

PTEN tumor suppressor gene. *Proc Natl Acad Sci USA* **96:** 2925-2930.

Goh GYS, Winter JJ, Bhanshali F, Doering KRS, Lai R, Lee K, Veal EA, Taubert S.

2018. NHR-49/HNF4 integrates regulation of fatty acid metabolism with a

protective transcriptional response to oxidative stress and fasting. *Aging*

Cell **17:** e12743.

Goudeau J, Bellemin S, Toselli-Mollereau E, Shamalnasab M, Chen Y, Aguilaniu H.

2011. Fatty acid desaturation links germ cell loss to longevity through

NHR 80/HNF4 in *C. elegans*. *PLoS Biol* **9:** e1000599.

Grant B, Hirsh D. 1999. Receptor-mediated endocytosis in the *Caenorhabditis elegans*

oocyte. *Mol Biol Cell* **10:** 4311-4326.

Hales CN, Barker DJ. 1992. Type 2 (non-insulin-dependent) diabetes mellitus: the thrifty

phenotype hypothesis. *Diabetologia* **35:** 595-601.

Hall SE, Beverly M, Russ C, Nusbaum C, Sengupta P. 2010. A cellular memory of

developmental history generates phenotypic diversity in *C. elegans*. *Curr*

Biol **20:** 149-155.

Hall SE, Chirn GW, Lau NC, Sengupta P. 2013. RNAi pathways contribute to

developmental history-dependent phenotypic plasticity in *C. elegans*. *RNA*

19: 306-319.

Han S, Schroeder EA, Silva-García CG, Hebestreit K, Mair WB, Brunet A. 2017. Mono-

saturated fatty acids link H3K4me3 modifiers to *C. elegans* lifespan.

Nature **544:** 185-190.

Hansen M, Flatt T, Aguilaniu H. 2013. Reproduction, fat metabolism, and life span: what

is the connection? *Cell Metab* **17:** 10-19.

Hsin C, Kenyon C. 1999. Signals from the reproductive system regulate the lifespan

of *C. elegans*. *Nature* **399:** 362-366.

Hu Q, D'Amora DR, MacNeil LT, Walhout AJM, Kubiseski TJ. 2018. The *Caenorhabditis*

elegans oxidative stress response requires the NHR-49 transcription

factor. *G3 (Bethesda)* **8:** 3857-3863.

Kaletsky R, Yao V, Williams A, Runnels AM, Tadych A, Zhou S, Troyanskaya OG,

Murphy CT. 2018. Transcriptome analysis of adult *Caenorhabditis elegans*

cells reveals tissue-specific gene and isoform expression. *PLoS Genet*.

14: e1007559.

Kamath RS, Martinez-Campos M, Zipperlen P, Fraser AG, Ahringer J. 2001.

Effectiveness of specific RNA-mediated interference through ingested

double-stranded RNA in *Caenorhabditis elegans*. *Genome Biol* **2:**

RESEARCH0002.

Kenyon C. 2010. A pathway that links reproductive status to lifespan in *Caenorhabditis*

elegans. *Ann N Y Acad Sci* **1204:** 156-162.

Kenyon CJ. 2010b. The genetics of ageing. *Nature* **464:** 504-512.

Kimble J, Sharrock WJ. 1983. Tissue-specific synthesis of yolk proteins in

Caenorhabditis elegans. *Dev Biol* **96:** 189-196.

Kimble J, Ward S. 1988. Germ-line development and fertilization. in *The nematode*

Caenorhabditis elegans (ed. WB Wood), pp. 191-213. Cold Spring Harbor

Laboratory Press, New York, NY.

Kimble J, Crittenden SL. 2005. Germline proliferation and its control. in *WormBook* (ed.

The *C. elegans* research community) doi/10.1895/ wormbook.1.13.1.

Kirkwood TB. 1977. Evolution of aging. *Nature* **270:** 301-304.

Kishimoto S, Uno M, Okabe E, Nono M, Nishida E. 2017. Environmental stresses

induce transgenerationally inheritable survival advantages via

germline-to-soma communication in *Caenorhabditis elegans*. *Nat*

Commun **8:** 14031-14043.

Klengel T, Mehta D, Anacker C, Rex-Haffner M, Pruessner JC, Pariante CM, Pace TW,

Mercer KB, Mayberg HS, Bradley B, et al. 2013. Allele-specific FKBP5

DNA demethylation mediates gene-childhood trauma interactions. *Nat*

Neurosci **16:** 33-41.

Lapierre LR, Hansen M. 2012. Lessons from *C. elegans*: signaling pathways for

longevity. *Trends Endocrinol Metab* **23:** 637-644.

Lee K, Goh GY, Wong MA, Klassen TL, Taubert S. 2016 Gain-of-function alleles in

Caenorhabditis elegans nuclear hormone receptor *nhr-49* are functionally

distinct. *PLoS One* **11:** e0162708.

L'Hernault, SW. 2006. Spermatogenesis. in *WormBook* (ed. The *C. elegans* research

community) doi/10.1895/ wormbook.1.85.1.

Lin K, Hsin H, Libina N, Kenyon C. 2001. Regulation of the *Caenorhabditis elegans*

longevity protein DAF-16 by insulin/IGF-1 and germline signaling. *Nat*

Genet **28:** 139-145.

Lindblom TH, Pierce GJ, Sluder AE. 2001. A *C. elegans* orphan nuclear receptor

contributes to xenobiotic resistance. *Curr Biol* **11:** 864-868.

Lumey LH, Stein AD, Susser E. 2011. Prenatal famine and adult health. *Annu Rev*

Public Health **32:** 237-262.

Magner DB, Wollam J, Shen Y, Hoppe C, Li D, Latza C, Rottiers V, Hutter H, Antebi A.

2013. The NHR-8 nuclear receptor regulates cholesterol and bile acid

homeostasis in *C. elegans*. *Cell Metab* **18:** 212-224.

Mahanti P, Bose N, Bethke A, Judkins JC, Wollam J, Dumas KJ, Zimmerman AM,

Campbell SL, Hu PJ, Antebi A, Schroeder FC. 2014. Comparative metabolomics

reveals endogenous ligands of DAF-12, a nuclear hormone receptor, regulating

C. elegans development and lifespan. *Cell Metab* **19:** 73-83.

Mihaylova VT, Borland CZ, Manjarrez L, Stern MJ, Sun H. 1999. The PTEN tumor

suppressor homolog in *Caenorhabditis elegans* regulates longevity and

dauer formation in an insulin receptor-like signaling pathway. *Proc Natl*

Acad Sci USA **96:** 7427-7432.

Moreno-Arriola E, El Hafidi M, Ortega-Cuéllar D, Carvajal K. 2016. AMP-activated

protein kinase regulates oxidative metabolism in *Caenorhabditis elegans*

through the NHR-49 and MDT-15 transcriptional regulators. *PLoS One*

11: e0148089.

Murphy CT, Hu PJ. 2013. Insulin/insulin-like growth factor signaling in *C. elegans*. in

WormBook (ed. The *C. elegans* research community)

doi/10.1895/wormbook.1.164.1.

Neel JV. 1962. Diabetes mellitus: a "thrifty" genotype rendered detrimental by

"progress"? *Am J Hum Genet* **14:** 353-362.

Ni JZ, Kalinava N, Chen E, Huang A, Trinh T, Gu SG. 2016. A transgenerational role of

the germline nuclear RNAi pathway in repressing heat stress-induced

transcriptional activation in *C. elegans*. *Epigenetics Chromatin* **9:** 3-15.

Nomura T, Horikawa M, Shimamura S, Hashimoto T, Sakamoto K. 2010. Fat

accumulation in *Caenorhabditis el*egans is mediated by SREBP homolog SBP-1

Genes Nutr **5:** 17-27.

Ogg S, Ruvkun G. 1998. The *C. elegans* PTEN homolog, DAF-18, acts in the insulin

receptor-like metabolic signaling pathway. *Mol Cell* **2:** 887-893.

O'Rourke EJ, Soukas AA, Carr CE, Ruvkun G. 2009. *C. elegans* major fats are stored in

vesicles distinct from lysosome-related organelles. *Cell Metab* **10:** 430-435.

Ortiz MA, Noble D, Sorokin EP, Kimble J. 2014. A new dataset of spermatogenic vs.

oogenic transcriptomes in the nematode *Caenorhabditis elegans*. *G3*

(Bethesda) **4:** 1765-1772.

Ow MC, Borziak K, Nichitean AM, Dorus S, Hall SE. 2018. Early experiences mediate

distinct adult gene expression and reproductive programs in *Caenorhabditis*

elegans. *PLoS Genet* **14:** e1007219.

Painter R, Osmond C, Gluckman P, Hanson M, Phillips D, Roseboom T. 2008.

Transgenerational effects of prenatal exposure to the Dutch famine on

neonatal adiposity and health in later life. *BJOG* **115:** 1243-1249.

Patel DS, Fang LL, Svy DK, Ruvkun G, Li W. 2008. Genetic identification of HSD-1, a

conserved steroidogenic enzyme that directs larval development in

Caenorhabditis elegans. *Development* **135:** 2239-2249.

Papsdorf K, Brunet A. 2019. Linking lipid metabolism to chromatin regulation in aging.

Trends Cell Biol **29:** 97-116.

Partridge L, Gems D, Withers DJ. 2005. Sex and death: what is the connection? *Cell*

120: 461-472.

Pathare PP, Lin A, Bornfeldt KE, Taubert S, Van Gilst MR. 2012. Coordinate regulation

of lipid metabolism by novel nuclear receptor partnerships. *PLoS Genet* **8:**

e1002645.

Pellegroms B. Van Dongen S, Van Dyck H, Luc L. 2009. Larval food stress differentially

affects flight morphology in male and female speckled woods (*Pararge*

aegeria). *Ecol Entomol* **34:** 387-393.

Pushpa K, Kumar GA, Subramaniam K. 2013. PUF-8 and TCER-1 are essential for

normal levels of multiple mRNAs in the *C. elegans* germline. *Development*

140: 1312-1320.

Rastogi S, Borgo B, Pazdernik N, Fox P, Mardis ER, Kohara Y, Havranek J, Schedl T.

2015. *Caenorhabditis elegans glp-4* encodes a valyl aminoacyl tRNA

synthetase. *G3 (Bethesda)* **5:** 2719-2728.

Ratnappan R, Amrit FR, Chen SW, Gill H, Holden K, Ward J, Yamamoto KR, Olsen CP,

Ghazi A. 2014. Germline signals deploy NHR-49 to modulate fatty-acid β-

oxidation and desaturation in somatic tissues of *C. elegans*. *PLoS Genet* **10:**

e1004829.

Rechavi O, Houri-Ze'evi L, Anava S, Goh WSS, Kerk SY, Hannon GJ, Hobert O. 2014.

Starvation-induced transgenerational inheritance of small RNAs in *C.*

elegans. *Cell* **158:** 277-287.

Rechavi O, Lev I. 2017. Principles of transgenerational small RNA inheritance in

Caenorhabditis elegans. *Curr Biol* **27:** R720-R730.

Rottiers V, Motola DL, Gerisch B, Cummins CL, Nishiwaki K, Mangelsdorf DJ, Antebi A.

2006. Hormonal control of *C. elegans* dauer formation and life span by a

Rieske-like oxygenase. *Dev Cell* **10:** 473-482.

Schulenburg H, Félix MA. 2017. The natural biotic environment of *Caenorhabditis*

elegans. *Genetics* **206:** 55-86.

Seah NE, de Magalhaes Filho, CD, Petrashen AP, Henderson HR, Laguer J, Gonzalez

J, Dillin A, Hansen M, Lapierre LR. 2016. Autophagy-mediated longevity is

modulated by lipoprotein biogenesis. *Autophagy* **12:** 261-272.

Seydoux G, Savage C, Greenwald I. 1993. Isolation and characterization of mutations

causing abnormal eversion of the vulva in *Caenorhabditis elegans*. *Dev*

Biol. **157:** 423-436.

Shakes DC, Wu JC, Sadler PL, Laprade K, Moore LL, Noritake A, Chu DS. 2009.

Spermatogenesis-specific features of the meiotic program in *Caenorhabditis*

elegans. PLoS Genet. **5:** e1000611.

Shen Y, Wollam J, Magner D, Karalay O, Antebi A. 2012. A steroid receptor-microRNA

switch regulates life span in response to signals from the gonad. *Science* **338:**

1472-1476.

Smith CJ, Ryckman KK. 2015. Epigenetic and developmental influences on the risk of

obesity, diabetes, and metabolic syndrome. *Diabetes Metab Syndr Obes*

8: 295-302.

Solari F, Bourbon-Piffaut A, Masse I, Payrastre B, Chan AM, Billaud M. 2005. The

human tumour suppressor PTEN regulates longevity and dauer formation

in *Caenorhabditis elegans*. *Oncogene* **24:** 20-27.

Spieth J, Nettleton M, Zuckeraprison E, Lea K, Blumenthal T. 1991. Vitellogenin motifs

conserved in nematodes and vertebrates. *J Mol Evol* **32:** 429-438.

Stein LD, Bao Z, Blasiar D, Blumenthal T, Brent MR, Chen N, Chinwalla A, Clarke L,

Clee C, Coghlan A, et al. 2003. The genome sequence of *Caenorhabditis*

briggsae: a platform for comparative genomics. *PLoS Biol* **1:** e45.

Steinbaugh MJ, Narasimhan SD, Robida-Stubbs S, Moronetti Mazzeo LE, Dreyfuss JM,

Hourihan JM, Raghavan P, Operaña TN, Esmaillie R, Blackwell TK. 2015.

Lipid mediated regulation of SKN-1/Nrf in response to germ cell absence.

Elife **4:** e07836.

Stiernagle T. 2006. Maintenance of *C. elegans*. in *WormBook* (ed. The *C. elegans*

research community) doi/10.1895/wormbook.1.101.1.

Sulston JE, Horvitz HR. 1977. Post-embryonic cell lineages of the nematode,

Caenorhabditis elegans. *Dev Biol* **56:** 110-156.

Svensk E, Ståhlman M, Andersson CH, Johansson M, Borén J, Pilon M. 2013. PAQR-2

regulates fatty acid desaturation during cold adaptation in *C. elegans*.

PLoS Genet **9:** e1003801.

Taubert S, Van Gilst MR, Hansen M, Yamamoto KR. 2006. A Mediator subunit, MDT-

15, integrates regulation of fatty acid metabolism by NHR-49-dependent

and -independent pathways in *C. elegans*. *Genes Dev* **20:** 1137-1149.

Tauffenberger A, Parker JA. 2014. Heritable transmission of stress resistance by high

dietary glucose in *Caenorhabditis elegans*. *PLoS Genet* **10:** e1004346.

Telang A, Wells MA. 2004. The effect of larval and adult nutrition on successful

autogenous egg production by a mosquito. *J Insect Physiol* **50:** 677-685.

Tepper RG, Ashraf J, Kaletsky R, Kleemann G, Murphy CT, Bussemaker HJ. 2013.

PQM-1 complements DAF-16 as a key transcriptional regulator of DAF-2-

mediated development and longevity. *Cell* **154:** 676-690.

Tobi EW, Goeman JJ, Monajemi R, Gu H, Putter H, Zhang Y, Slieker RC, Stok AP,

Thijssen PE, Müller F, et al. 2014. DNA methylation signatures link

prenatal famine exposure to growth and metabolism. *Nat Commun* **5:**

5592.

Vaag AA, Grunnet LG, Arora GP, Brøns C. 2012. The thrifty phenotype hypothesis

revisited. *Diabetologia* **55:** 2085-2088.

Van Abeelen AF, Elias SG, Bossuyt PM, Grobbee DE, van der Schouw YT, Roseboom

TJ, Uiterwaal CS. 2012. Famine exposure in the young and the risk of

type 2 diabetes in adulthood. *Diabetes* **61:** 2255-2260.

Van Gilst MR, Hadjivassiliou H, Jolly A, Yamamoto KR. 2005a. Nuclear hormone

receptor NHR-49 controls fat consumption and fatty acid composition in *C.*

elegans. *PLoS Biol* **3:** e53.

Van Gilst MR, Hadjivassiliou H, Yamamoto KR. 2005b. A *Caenorhabditis elegans*

nutrient response system partially dependent on nuclear receptor NHR-49. *Proc*

Natl Acad Sci USA **102:** 13496-13501.

Veenendaal M, Painter R, de Rooij S, Bossuyt P, van der Post J, Gluckman P, Hanson

M, Roseboom T. 2013. Transgenerational effects of prenatal exposure to

the 1944-45 Dutch famine. *BJOG* **120:** 548-554.

Vitikainen EIK, Thompson FJ, Marshall HH, Cant MA. 2019. Live long and prosper:

durable benefits of early-life care in banded mongooses. *Phil Trans R Soc*

B **374:** 20180114.

Wang MC, O'Rourke E, Ruvkun G. 2008. Fat metabolism links germline stem

cells and longevity in *C. elegans*. *Science* **322:** 957-960.

Wang H, Jiang, X, Wu J, Zhang L, Huang J, Zhang Y, Zou X, Liang B. 2016. Iron

overload coordinately promotes ferritin expression and fat accumulation in

Caenorhabditis elegans. *Genetics* **203:** 241-253.

Ward S, Carrel JS. 1979. Fertilization and sperm competition in the nematode

Caenorhabditis elegans. *Dev Biol* **73:** 304-321.

Watts JL, Ristow M. 2017. Lipid and carbohydrate metabolism in *Caenorhabditis*

elegans. *Genetics* **207:** 413-446.

Weaver IC, Cervoni N, Champagne FA, D'Alessio AC, Sharma S, Seckl JR, Dymov S,

Szyf M, Meaney MJ. 2004. Epigenetic programming by maternal behavior.

Nat Neurosci **7:** 847-854.

Westendorp RGJ, Kirkwood TBL. 1998. Human longevity at the cost of reproductive

success. *Nature* **396:** 743-746.

Williams GC. 1966. Natural selection, the costs of reproduction, and a refinement of

Lack's principle. *Am Nat* **100:** 687-690.

Wolff S, Ma H, Burch D, Maciel GA, Hunter T, Dillin A. 2006. SMK-1, an essential

regulator of DAF-16-mediated longevity. *Cell* **124:** 1039-1053.

Yamawaki TM, Berman JR, Suchanek-Kavipurapu M, McCormick M, Gaglia MM, Lee

SJ, Kenyon C. 2010. The somatic reproductive tissues of *C. elegans* promote

longevity through steroid hormone signaling. *PLoS Biol.* **8:** e1000468.

Yang F, Vought BW, Satterlee JS, Walker AK, Jim Sun ZY, Watts JL, DeBeaumont R,

Saito RM, Hyberts SG, Yang S, et al. 2006. An ARC/Mediator subunit

required for SREBP control of cholesterol and lipid homeostasis. *Nature*

442: 700-704.

Yen K, Le TT, Bansal A, Narasimhan SD, Cheng JX, Tissenbaum HA. 2010. A

comparative study of fat storage quantitation in nematode *Caenorhabditis*

elegans using label and label-free methods. *PLoS One* **5:** e12810.

Zhao LQ, Zhu DH. 2014. Effects of environmental factors and appendage injury on the

wing variation in the cricket *Velarifictorus ornatus*. *J Insect Sci* **14:** 117.

Figure legends

Figure 1. Adult reproductive plasticity is dependent on steroid signaling. (*A*) Brood

size comparison of PD_{Stv} relative to CON_{Stv} in wild-type N2, *daf-12(rh284)*, *daf-*

12(rh285), *daf-9(rh50)*, *daf-36(k114)*, *hsd-1(mg433)*, and *nhr-8(ok186)* strains. Assays

were performed at 20°C from at least three biologically independent trials. Significance

asterisks (** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$; Student's *t*-test) refer to the

comparison of PD_{Stv} to CON_{Stv} of a given genotype. Significance ampersands ([&&&] $p <$

0.001, [&&&&] $p < 0.0001$; one-way ANOVA with Fisher's LSD test) refer to the comparison

of PD_{Stv}/CON_{Stv} brood size change between a given mutant genotype to wild-type N2.

Additional brood size data is provided in Supplemental Table S1. (*B*) Brood size

comparison of PD_{Stv} grown in the presence of 40 nM Δ^7-dafachronic acid (Δ^7-DA)

relative to the brood size of PD_{Stv} grown in the absence of Δ^7-DA of wild-type N2, *daf-*

9(rh50), and *daf-36(k114)* strains. Assays were conducted at 20°C from four biologically

independent trials. **** $p < 0.0001$ (Student's *t*-test) indicates the comparison between

PD_{Stv} with Δ^7-DA to PD_{Stv} without Δ^7-DA for *daf-9(rh50)* adults. [&] $p < 0.05$ (one-way

ANOVA with Fisher's LSD test) refers to the comparison of brood size fold change

(PD_{Stv} with Δ^7-DA to PD_{Stv} without Δ^7-DA) between *daf-9(rh50)* relative to wild-type N2.

Additional brood size data can be found in Supplemental Table S2.

1076

Figure 2. DAF-16 and TCER-1 modulation of fatty acid metabolism is important for

adult reproductive plasticity. (*A*) Venn diagram depicting overlap of DAF-16 and PQM-1

targets (Tepper et al. 2013) with differentially expressed N2 PD_{Stv} genes compared to

CON_{Stv} adults (Ow et al. 2018). See Supplemental Table S3. (*B*) Brood size comparison

of PD_{Stv} relative to CON_{Stv} in wild-type N2, *daf-18(e1375), pqm-1(ok485), daf-16(mu86);*

daf-16a^AM^::gfp, smk-1(mn156), kri-1(ok1251), and *tcer-1(tm1425)* strains. Assays were

performed at 20°C from at least three biologically independent replicates. ** $p < 0.01$, ***

$p < 0.001$, **** $p < 0.0001$ (Student's *t*-test) refer to the comparison of PD_{Stv} brood size

relative to CON_{Stv} of a given genotype. & $p < 0.05$, &&&& $p < 0.0001$ (one-way ANOVA

with Fisher's LSD test) indicate the comparison of the PD_{Stv}/CON_{Stv} brood size change

between a given mutant genotype to wild-type N2. Additional brood size data is

provided in Supplemental Table S4. (*C*) Venn diagram comparing DAF-16 and TCER-1

targets with functions in lipid metabolism from germ line-less *glp-1(e2141ts)* animals

(Amrit et al. 2016) with differentially expressed N2 PD_{Stv} genes compared to CON_{Stv}

adults (Ow et al. 2018). See Supplemental Table S5.

1092

Figure 3.   NHR-49, SBP-1, and FAT-6 modulate adult reproductive plasticity. (*A*)

Brood size comparison of PD_{Stv} relative to CON_{Stv} for wild-type N2, *nhr-49(nr2041), nhr-*

49(ok2165), nhr-49(gk405), nhr-49(et8), nhr-49(et7), nhr-49(et13), nhr-80(tm1011), nhr-

13(gk796), nhr-13(gk796), nhr-66(ok940), mdt-15(tm2182), and *sbp-1(ep79)* strains.

Additional brood size data is provided in Supplemental Table S6. (*B*) Brood size

comparison of PD_{Stv} relative to CON_{Stv} for wild-type N2, *nhr-49(nr2041); nhr-*

80(tm1011), nhr-80(tm1011); nhr-13(gk796), nhr-49(nr2041); nhr-13(gk796), nhr-

49(nr2041); nhr-80(tm1011); nhr-13(gk796), and *nhr-49(nr2041); nhr-66(ok940)* strains.

Additional brood size data is provided in Supplemental Table S7. (*C*) Brood size

comparison of PD_{Stv} relative to CON_{Stv} for wild-type N2, *fat-6(tm331), fat-5(tm420), fat-*

7(wa36), fat-2(wa17), fat-6(tm331); fat-5(tm420), fat-7(wa36); fat-5(tm420), and *fat-*

6(tm420); *fat-7(wa36)* strains. Additional brood size data is provided in Supplemental

Table S8. (*D*) Brood size of PD_{Stv} wild-type N2, *nhr-49(nr2041)*, *nhr-49(gk405)*, and *nhr-*

49(nr2041); nhr-80(tm1011) animals fed with *E. coli* OP50 pre-loaded with or without

600 μM of oleic acid (OA). Additional brood size data is provided in Supplemental Table

S9. All assays were performed at 20°C from at least three biologically independent

replicates. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$ (Student's *t*-test)

indicate brood size comparison within a given genotype. & $p < 0.05$, &&&& $p < 0.0001$ and

n.s. (no significance) (one-way ANOVA with Fisher's LSD test) indicates the comparison

of PD_{Stv}/CON_{Stv} brood size change between a given genotype to wild-type N2.

1113

Figure 4.   Fat stores are reduced in wild-type postdauer adults that experienced

starvation-induced dauer. (*A*) Neutral lipid staining with Oil Red O (ORO) in CON_{Stv} and

PD_{Stv} wild-type N2, *daf-12(rh284)*, *daf-12(rh285)*, and *tcer-1(tm1452)* one-day-old adults

grown at 20°C. **** $p < 0.0001$; Student's *t*-test comparison between CON_{Stv} and PD_{Stv}

of the same genotype. &&&& $p < 0.0001$; one-way ANOVA with Fisher's LSD test

between wild-type N2 CON_{Stv} and mutant CON_{Stv}. \$\$\$\$ $p < 0.0001$; one-way ANOVA

between wild-type N2 PD_{Stv} and mutant PD_{Stv}. n.s. indicates no significance between

wild-type N2 PD_{Stv} and mutant PD_{Stv} (one-way ANOVA with Fisher's LSD test). Total

sample size is indicated by *n* from at least 3 biologically independent trials. (*B*)

Representative micrographs of one-day-old adults stained with ORO. *(C)* ORO staining

of CON_{Stv} and PD_{Stv} wild-type N2 fed with empty RNAi vector or *vit-1* RNAi, which

knocks down *vit-1, 2, 3, 4,* and *5* (see Results). **** $p < 0.0001$; Student's *t*-test

comparison between CON_{Stv} and PD_{Stv} of the same RNAi. && $p = 0.0038$ and &&&& $p <$

0.0001 (Student's *t*-test) denote comparison between *vit-1* RNAi to empty RNAi vector

control of the same condition. *(D)* Representative micrographs of ORO staining of (C)

showing whole worms and their embryos.

1130

Figure 5.   DAF-12 acts downstream of KRI-1 and TCER-1 to regulate adult

reproductive plasticity. (*A*) Brood size comparison of PD_{Stv} relative to CON_{Stv} for wild-

type N2, *kri-1(ok1251); daf-12(rh284)*, *tcer-1(tm1425); dat-12(rh284)*, *kri-1(ok1251); daf-*

12(rh285), and *tcer-1(tm1425); dat-12(rh285)*. * $p < 0.05$; *** $p < 0.001$, **** $p < 0.0001$

(Student's *t*-test) indicates the brood size comparison between PD_{Stv} relative to CON_{Stv}

of a given genotype. &&&& $p < 0.0001$ (one-way ANOVA with Fisher's LSD test) indicates

the comparison of PD_{Stv}/CON_{Stv} brood size change between a given genotype to wild-

type N2. Assays were performed at 20°C from at least three independent biological

replicates. Additional brood size data is provided in Supplemental Table S10. (*B*) Total

germ cell rows in CON_{Stv} and PD_{Stv} wild-type N2, *daf-36(k114), nhr-49(nr2041),* and *tcer-*

1(tm1452) larva exhibiting L3 vulva morphology (see Materials and methods). ** $p <$

0.01; **** $p < 0.0001$ (Student's *t*-test) indicate comparison between CON_{Stv} and PD_{Stv}

within a genotype. Total sample size is indicated by *n* from at least three biologically

independent trials.

1145

Figure 6.   Transgenerational inheritance of starvation memory. *(A)* ORO lipid

staining of wild-type N2 and *hrde-1(tm1200)* control and postdauer P_0, F_1, and F_2

generations. Only the P_0 PD_{Stv} generation experienced starvation during early

development. **** $p < 0.0001$; Student's *t*-test comparison between CON_{Stv} and PD_{Stv} of

the same genotype and generation. n.s. indicates no significance. Assays were

performed at 20°C from three independent biological replicates. Total sample size is

indicated by *n*. (*B*) Representative micrographs of N2 and *hrde-1(tm1200)* CON_{Stv} and

PD_{Stv} P_0, F_1, and F_2 worms stained with ORO.

1154

1155   **Figure 7.**   Model depicting genetic pathways acting as contributors to fatty acid

1156   metabolism of postdauer animals that experienced early life starvation. Signaling

1157   between the soma and the germ line establishing metabolic plasticity and the

1158   transmission of ancestral starvation memory are dependent on HRDE-1.

Ow et al. Figure 1

A

B

A

B

C

A

B

C

D

Ow et al. Figure 4

A

B

C

D

A

B

Ow et al. Figure 6

A

B