Pathognomonic and epistatic genetic alterations in B-cell non-Hodgkin lymphoma

Man Chun John Ma[1][¥], Saber Tadros[1][¥], Alyssa Bouska[2], Tayla B. Heavican[2], Haopeng Yang[1], Qing Deng[1], Dalia Moore[3], Ariz Akhter[4], Keenan Hartert[3], Neeraj Jain[1], Jordan Showell[1], Sreejoyee Ghosh[1], Lesley Street[5], Marta Davidson[5], Christopher Carey[6], Joshua Tobin[7], Deepak Perumal[8], Julie M. Vose[9], Matthew A. Lunning[9], Aliyah R. Sohani[10], Benjamin J. Chen[11], Shannon Buckley[12], Loretta J. Nastoupil[1], R. Eric Davis[1], Jason R. Westin[1], Nathan H. Fowler[1], Samir Parekh[8], Maher K. Gandhi[7], Sattva S. Neelapu[1], Douglas Stewart[5], Javeed Iqbal[2], Timothy Greiner[2], Scott J. Rodig[13], Adnan Mansoor[5], Michael R. Green[1,14,15]*

[1]Department of Lymphoma and Myeloma, Division of Cancer Medicine, The University of Texas MD Anderson Cancer Center, Houston, TX, USA; [2]Department of Pathology and Microbiology, University of Nebraska Medical Center, Omaha, NE, USA; [3]Eppley Institute for Research in Cancer and Allied Diseases, University of Nebraska Medical Center, Omaha, NE, USA; [4]Department of Pathology and Laboratory Medicine, University of Calgary, Calgary, AB, Canada; [5]Section of Hematology, Department of Medicine, University of Calgary, Calgary, AB, Canada; [6]Northern Institute for Research, Newcastle University, Newcastle upon Tyne, England; [7]Mater Research, University of Queensland, QLD, Australia; [8]Division of Hematology and Medical Oncology, Icahn School of Medicine at Mount Sinai, New York, NY, USA; [9]Department of Internal Medicine, Division of Hematology-Oncology, University of Nebraska Medical Center, Omaha, NE, USA; [10]Department of Pathology, Massachusetts General Hospital and Harvard Medical School, Boston, MA, USA; [11]Department of Pathology, University of Massachusetts Medical School, UMass Memorial Medical Center, Worcester, MA, USA; [12]Department of Genetics, Cell Biology and Anatomy, University of Nebraska Medical Center, Omaha, NE, USA; [13]Department of Pathology, Brigham and Womens Hospital, Boston, MA, USA; [14]Department of Genomic Medicine, University of Texas MD Anderson Cancer Center, Houston, TX, USA; [15]Center for Cancer Epigenetics, University of Texas MD Anderson Cancer Center, Houston, TX, USA.

[¥]Equally contributed

*Corresponding Author

Michael R. Green, Ph.D.

Departments of Lymphoma/Myeloma and Genomic Medicine,

University of Texas MD Anderson Cancer Center,

1515 Holcombe Blvd, Unit 903,

Houston, TX 77030, USA

Ph.:   +1-713-745-4244

Email: mgreen5@mdanderson.org

KEY POINTS

1. Genetic perturbation of the ubiquitin proteasome system is an emerging hallmark of B-cell non-Hodgkin lymphoma (B-NHL).

2. Co-occurring sets of genetic alterations define B-NHL subtypes and likely represent epistatic interactions.

ABSTRACT

B-cell non-Hodgkin lymphoma (B-NHL) encompasses multiple clinically and phenotypically distinct subtypes of malignancy with unique molecular etiologies. Common subtypes of B-NHL such as diffuse large B-cell lymphoma (DLBCL) have been comprehensively interrogated at the genomic level, but other less common subtypes such as mantle cell lymphoma (MCL) remain sparsely characterized. Furthermore, multiple B-NHL subtypes have thus far not been comprehensively compared to identify conserved or subtype-specific patterns of genomic alterations. Here, we employed a large targeted hybrid-capture sequencing approach encompassing 380 genes to interrogate the genomic landscapes of 755 B-NHL tumors at high depth; primarily including DLBCL, MCL, follicular lymphoma (FL), and Burkitt lymphoma (BL). We identified conserved hallmarks of B-NHL that were deregulated across major subtypes, such as the frequent genetic deregulation of the ubiquitin proteasome system (UPS). In addition, we identified subtype-specific patterns of genetic alterations, including clusters of co-occurring mutations that are pathognomonic. The cumulative burden of mutations within a single cluster were more significantly discriminatory of B-NHL subtypes than individual mutations, implicating likely patterns of genetic epistasis that contribute to disease etiology. We therefore provide a framework of co-occurring mutations that deregulate genetic hallmarks and likely cooperate in lymphomagenesis of B-NHL subtypes.

INTRODUCTION

Non-Hodgkin lymphomas (NHL) are a heterogeneous group of lymphoid malignancies that predominantly arise from mature B-cells (B-NHL). Although mature B-cell neoplasms encompass 38 unique diagnostic subtypes, over 85% of cases fall within only 7 subtypes[1,2]. Recent next generation sequencing (NGS) studies have shed light onto the key driver mutations in many of these NHL subtypes; for example, large studies of diffuse large B-cell lymphoma (DLBCL) have led to proposed genomic subtypes that have unique etiologies[3-5]. However, many less-common NHL subtypes such as mantle cell lymphoma (MCL) have not been as extensively characterized[6,7]. Furthermore, until recently[3,4] genetic alterations have been considered in a binary fashion as either driver events, which directly promote disease genesis or progression, or passenger events, which have little or no impact on disease biology. In contrast to this principle, most B-NHLs do not result from a single dominant driver but instead result from the serial acquisition of genetic alterations that cooperate in lymphomagenesis. The genetic context of each mutation therefore likely determines its oncogenic potential, and groups of mutations should therefore be considered collectively rather than as singular events. For example, the 'C5' and 'MCD' clusters identified in DLBCL by Chapuy *et al.* and Schmitz *et al.*, respectively, are characterized by the co-occurrence of *CD79B* and *MYD88* mutations[3,4]. In animal models, the *Myd88* L265P mutation was found to promote down-regulation of surface IgM and a phenotype resembling B-cell anergy[8]. However, this effect could be rescued by *Cd79b* mutation, showing that these co-occurring mutations are epistatic[8]. The characterization of other significantly co-occurring genetic alterations are therefore likely to reveal additional important epistatic relationships.

We approached this challenge by performing genomic profiling of 755 B-NHLs across different subtypes. Through this cross-sectional analysis we characterized genomic hallmarks of B-NHL

and identified pathognomonic genetic alterations, including disease-specific mechanisms for

deregulating hallmark processes and protein complexes, and sets of significantly co-associated

events that represent subtype-specific epistatic genetic alterations. This study therefore

provides new insight into how cooperating genetic alterations may contribute to molecularly and

phenotypically distinct subtypes of B-NHL.

METHODS

An overview of our approach is shown in Figure S1. For detailed methods, please refer to

supplementary information.

<u>Tumor DNA samples</u>

We collected DNA for 755 B-NHL tumors, including 199 follicular lymphoma (FL), 196 mantle

cell lymphoma (MCL), 148 diffuse large B-cell lymphoma (DLBCL), 108 Burkitt's lymphoma

(BL), 45 chronic lymphocytic leukemia / small lymphocytic lymphoma (CLL/SLL), 24 marginal

zone lymphoma (MZL), 21 high-grade B-cell lymphoma not otherwise specified (HGBL-NOS),

and 14 high-grade B-cell lymphoma with MYC, BCL2 and/or BCL6 rearrangement (DHL) (Table

S1). A total of 502 samples were obtained from the University of Nebraska Medical Center, and

were prioritized for inclusion in this study if they had fresh/frozen tissue available (n=577) and

been previously interrogated by Affymetrix U133 Plus 2.0 gene expression microarrays

(n=290)[9-11]. An additional series of 178 formalin-fixed paraffin-embedded (FFPE) tumors were

collected from other centers. Samples were de-identified but accompanied by their diagnosis

from medical records, plus overall survival time and status (alive, dead) when available. Medical

record diagnosis was used in all cases except for those with fluorescence *in situ* hybridization

showing translocations in MYC and BCL2 and/or BCL6, which were amended to DHL.

Sequencing results for a subset of 52 BL tumors was described previously[12]. All MCL samples

were either positive for *CCND1* translocation by FISH or positive for CCND1 protein expression

by IHC, depending on the diagnostic practices of the contributing institution.

<u>Next generation sequencing</u>

A total of 100-1000ng of gDNA was sonicated using a Covaris S2 Ultrasonicator, and libraries

prepared using KAPA Hyper Prep Kits (Roche) with TruSeq Adapters (Bioo Scientific) and a

maximum of 8 cycles of PCR (average of 6 cycles). Libraries were qualified by TapeStation

4200, quantified by Qubit and 10- to 12-plexed for hybrid capture. Each multiplexed library was

enriched using our custom LymphoSeq panel encompassing the full coding sequences of 380

genes that have previously been reported to be somatically mutated in B-cell lymphoma[6,7,13-32]

(Table S2, Supplementary Methods), as well as tiling recurrent translocation breakpoints.

Enrichments were amplified with 4-8 cycles of PCR and sequenced on a single lane of an

Illumina HiSeq 4000 with 100PE reads in high-output mode at the Hudson Alpha Institute for

Biotechnology or the MD Anderson Sequencing and Microarray Facility. Variants were called

using our previously validated ensemble approach[12,21], copy number alterations identified using

off-target reads with the CopyWriteR tool[33], and translocation called using FACTERA[34].

Germline polymorphisms were filtered using dbSNP annotation and the EXAC dataset

containing 60,706 healthy individuals[35]. Significantly mutated genes were defined by

MutSig2CV[36], significant DNA copy number alterations by GISTIC2[37], and the clonal

representation of mutations by ABSOLUTE[38]. Mutation and CNA data are publicly viewable

through cBioPortal: https://www.cbioportal.org/study/summary?id=mbn_mdacc_2013. For

further details, refer to supplementary methods.

RESULTS

<u>Recurrently mutated genes highlight conserved functional hallmarks of B-NHL</u>

We used a 380 gene custom targeted sequencing approach to interrogate the genomes of 755 mature B-NHLs, sequencing to an average depth of 599X (Min, 101X; Max, 1785X; Table S1) for a total yield of 2.06 Tbp. Somatic nucleotide variants (SNVs) and small insertions/deletions (InDels) were identified using an ensemble approach that we have previously validated[21] (Table S3) and significantly mutated genes were identified using MutSig2CV (Table S4). Genes that were significantly mutated in the full cohort or in any one of the 4 subtypes with >100 tumors (BL, DLBCL, FL, MCL), as well as frequently mutated genes that are targets of AID (Table S5, **Figure 1**), were included in downstream analyses. Mutation distributions were classified using a novel metric that classifies them into hotspot, clustered or diffuse (Figure S2). The residue of mutational hotspots and conserved domains targeted by clustered mutations are shown on figure 1. As a proof of principle, genes such as *EZH2* and *MYD88* with known mutational hotspots were classified as 'hotspot', and genes such as *CREBBP* and *TCF3* that accumulate mutations within a single domain were classified as 'clustered'. The mutational burden calculated from our targeted region significantly correlated with that from the whole exome (Figure S3A) and was significantly higher in DLBCL and other high-grade tumors compared to FL and MCL (**Figure 1**; Figure S3B). The mutational signatures were also different between malignancies, but were predominated by mutations attributable to failure of double-strand break repair by homologous recombination[39] (Signature 3; Table S6).

To identify key hallmarks that are deregulated by somatic mutations, we grouped genes using DAVID functional annotation clustering[40]. BCL6 function is not annotated in gene sets within public databases, but was an obvious feature of the recurrently mutated genes, so this hallmark

was assigned according to literature support. The most frequently perturbed processes were

chromatin modification and cell signaling, with most of the genes in these categories having

been well described in prior reports. Chromatin modifying genes were mutated in 68% of BL,

66% of DLBCL, 91% of FL and 45% of MCL, and included those that encode proteins that

catalyze post-translational modifications of histones (*KMT2D*, *CREBBP*, *EZH2*, *EP300*,

WHSC1, *ASHL1L* and *KMT2A*), components of the SWI/SNF chromatin remodeling complex

(*ARID1A*, *SMARCA4*, *BCL7A*), linker histones (*HIST1H1E*, *HIST1H1C*, *HIST1H1B*), and the

TET2 gene. Genes with a role in signaling included those involved in B-cell receptor (*CD79B*,

ID3, *TCF3, RFTN1*), toll-like receptor (*MYD88*), NFκB (*TNFAIP3*, *CARD11*, *NFKBIE*), Notch

(*NOTCH1*, *NOTCH2*), JAK/STAT (*SOCS1*, *STAT6*), PI3K/mTOR (*FOXO1*, *ATP6V1B2*,

APT6AP1) and G-protein (GNA13, GNAI2) signaling. The *CD79A* and *BCL10* genes were also

mutated at a lower frequency that was not significant by MutSig (Figure S4A-B). Among these,

the *RFTN1* gene (Figure S4C) is a novel recurrently mutated gene that was mutated in 7.4% of

DLBCL and encodes a lipid raft protein that is critical for B-cell receptor signaling[41].

Other processes that are not as well described as genetic hallmarks of B-NHL included BCL6

function and the ubiquitin proteasome system (UPS). Genes associated with BCL6 function

encompassed BCL6 itself and its interacting proteins (*BCL6, TBL1XR1*, *BCOR*, *SPEN*),

regulators of BCL6 activity or expression (*MEF2B*, *IRF8*, *IRF4*), and a critical BCL6 target gene

(*PRDM1*). Recurrent mutations of the *NCOR1* and *NCOR2* genes that encode BCL6 co-

repressor proteins were also observed (Figure S4D-E), but were not significant by MutSig2CV.

Multiple recurrently mutated chromatin modifying genes have also been implicated in BCL6

function and therefore also contribute to this hallmark, including *CREBBP* and *EZH2*[42,43]. The

recurrently mutated UPS genes included the *CDC27* gene, which encodes an E3 ligase for

CCND1[44] that is mutated in 14% of MCL and has not been previously described in the literature.

The *TMEM51* gene was also identified as a novel recurrently mutated gene in 9% of BL, though

the function of this gene is poorly defined. The cross-sectional analysis of multiple lymphoma

subtypes therefore identified conserved hallmarks that are recurrently targeted by somatic

mutations in B-NHL.

<u>Enrichment of functional hallmarks by structural alterations</u>

The hybrid capture probes utilized in our design also targeted recurrent breakpoint regions in

the immunoglobulin heavy- and light-chain loci, as well as the regions of recurrent breakpoints

in or near the *BCL2*, *MYC* and *BCL6* genes. Translocations were called using a method that

detects discordantly mapped reads[34] and our prior validation of this approach in cases with

matched fluorescence in situ hybridization (FISH) data for *MYC* showed that it is 100% specific,

but only ~40% sensitive for translocation detection[12]. In addition, we failed to detect *CCND1*

translocations using this approach. However, we did observe a significantly higher fraction of

BCL6 translocations (57% [27/47]) partnered to non-immunoglobulin loci (eg. *CIITA*, *RHOH*,

EIF4A2, *ST6GAL1*; Table S7) compared to *BCL2* (1% [1/114]) and *MYC* (5% [2/38])

translocations (**Figure 2A**; Fisher P-value <0.001). These were more frequent in FL (88%

[15/17] of *BCL6* translocations) as compared to DLBCL (39% [9/23] of *BCL6* translocations),

presumably because the two immunoglobulin loci in FL are either translocated with the *BCL2*

gene or functioning in immunoglobulin expression[45]. We also employed off-target reads to

detect DNA copy number alterations (CNAs) in a manner akin to low-pass whole genome

sequencing, identified significant peaks of copy gain and losses using GISTIC2[37] (**Figure 2A**;

Figure S5; Table S8-9), and defined the likely targets of these CNAs by integrative analysis of

matched gene expression profiling (GEP) data from 290 tumors (**Figure 2B-C**, Figure S5, Table

S10-11). This identified known CNA targets (**Figure 2D**), including but not limited to deletion of

TNFAIP3 (6q24.2)[46], *ATM* (11q22.3)[47], *B2M* (15q15.5)[48] and *PTEN* (10q23.21)[49], and copy gain

of *REL* and *BCL11A* (2p15), and *TCF4* (18q23)[50]. In addition, we identified novel targets such

as deletion of *IBTK* (6q14.1), *UBE3A* (11q22.1) and *FBXO25* (8p23.3), and copy gain of *ATF7*

(12q13.13), *UCHL5* (1q31.3), and *KMT2A* (11q23.3). Several CNA peaks, defined as the

smallest and most statistically significant region, included genes that were significantly mutated

(**Figure 2E**) as well as other genes for which we detected mutations at lower frequencies that

were not significant by MutSig (*POU2AF1*, *TP53BP1*, *FAS*, *PTEN*). Furthermore, for several of

these genes (*ATM*, *B2M, BIRC3* and *TNFRSF14*), deletions were significantly co-associated

with mutations, suggesting that deletion and mutation are complementary mechanisms

contributing to biallelic inactivation.

Analyzing targets of translocations and CNAs added to enrichment of functional categories that

we observed within recurrently mutated genes, including chromatin modification (*BCL11A*) and

signaling (*TCF4*, *PTEN*). This was most prominently observed for genes with a role in apoptosis

and the cell cycle (*CDKN2A*, *CDKN2B*, *FAS*, *RPL5*, *DFFB*, *MDM2*, *CDK6*) and the UPS

(*CUL4A*, *FBXO25*, *IBTK*, *RNF38*, *UBAP1, UBE3A*, *UBQLN1*, *UCHL5*). Recurrently altered UPS

genes collectively promote or suppress the abundance or activity of proteins with roles in other

hallmark characteristics (**Figure 3A**). For example; *SOCS1*[51], *BIRC3*[52], *DTX1*[53,54], *IBTK*[55],

TRIM13[56], *TNFAIP3*[53] and *UBE3A*[53] all inhibit the function of proteins with a role in important B-

cell signaling pathways (**Figure 3B**), thereby establishing a functional link between genetic

alteration of the UPS and deregulation of signaling and adding an additional layer to the

mechanisms by which B-cell receptor signaling is perturbed in B-NHL. The combined analysis of

mutations and structural alterations therefore identified large sets of genes that are targeted by

genetic alterations and collectively contribute to hallmark features that are commonly

deregulated across multiple B-NHL subtypes.

Disease-specific patterns of genetic alterations

Some interesting patterns of disease-specificity were obvious for both mutations and structural

alterations. We therefore formally tested the over- or under-representation of these events in

each of the 4 subtypes with >100 tumors (BL, DLBCL, FL, MCL), compared to all other tumors

in the study (**Figure 4**; Table S12). Using matched GEP data for a subset of cases (Table S1),

we also tested the association between genetic alterations and molecularly-defined Burkitt's

lymphoma subtypes (n=154; Figure S6-7, Table S13), as well as DLBCL cell of origin subtypes

(n=98; Figure S7, Table S14). These analyses showed that the genetic alterations associated

with clinically-defined BL were also associated with molecularly-defined BL. Moreover, genetic

alterations that we and others have characterized as being over-represented in ABC-like DLBCL

such as *CD79B* mutation[57], *MYD88* mutation[58] and *TCF4* copy gain[50] were significantly over-

represented in this analysis also.

We observed some interesting patterns within hallmark characteristics that differ between

subtypes. For example, linker histone mutations were present in germinal center B (GCB)-cell

derived malignancies (BL, DLBCL, FL) at variable frequencies, but were largely absent from

MCL (Figure S8). In contrast, mutations of genes encoding the H3K36 methyltransferases,

WHSC1 (aka *MMSET*/*NSD2*) and *ASH1L*, were frequent in MCL and largely absent from GCB-

derived malignancies. We also noted that the SWI/SNF complex was perturbed in different ways

in different diseases (**Figure 5**). Specifically, mutations of the *SMARCA4* (aka. BRG1)

component of the ATPase module were significantly enriched in BL (24%) compared to other

subtypes (4%, Q-value<0.001), while mutations of the *BCL7A* component of the ATPase

module were significantly enriched in FL (11%) compared to other subtypes (4%, Q-

value=0.007). In contrast, mutations of *ARID1A* were frequent in both BL (19%) and FL (15%),

and DNA copy number gains of *BCL11A* were frequent in both DLBCL (28%) and FL (22%).

The SWI/SNF complex is therefore a target of recurrent genetic alterations, as previously

suggested[59], but the manner in which this complex is perturbed varies between B-NHL subtypes

(**Figure 5**). Similar disease-specific patterns were also observed for signaling genes; for

example, *TCF3* and *ID3* have important functions in normal germinal center B-cells[60], but

mutations of these genes are specifically enriched within BL and are rarely found in the other

GCB-derived malignancies, DLBCL and FL. Similarly, the *ATP6AP1* and *ATP6V1B2* genes that

function in mTOR signaling[61,62] are specifically mutated in FL, and the *DUSP2* gene which

inactivates ERK1/2[63] and STAT3[64] is specifically mutated in DLBCL. The disease-specific

patterns of genetic alterations therefore reveal subtle but important differences in how each

subtype of B-NHL perturbs hallmark features.

Pathognomonic sets of co-associated genomic alterations in B-NHL subtypes

We next defined how each genetic alteration co-associated with or mutually-excluded other

genetic alterations by pairwise assessments using false-discovery rate (FDR)-corrected Fisher's

tests (Table S15). A matrix of the transformed FDR Q-values (-logQ) was used for unsupervised

hierarchical clustering to identify clusters of co-associated genetic alterations. Together with

patterns of disease-specificity, unsupervised clustering revealed clear groupings of co-

associated events for BL, DLBCL, FL and MCL (**Figure 4**). We identified a single cluster that

was specifically enriched in DLBCL, included co-associated genetic alterations that were over-

represented in the ABC-like, and overlapped with the previously described C5/MCD clusters[3,4].

We also identified a cluster consisting of TP53 mutations and multiple CNAs similar to the C2

subtype reported in DLBCL, but which was enriched in both FL and DLBCL. A cluster with

features similar to the C3/EZB DLBCL clusters was also observed, but was specifically enriched

in FL. The remaining two previously reported DLBCL clusters (C1/N2 and C4) were not

observed in this analysis, which may be a result of the different clustering strategies used and/or

the lower number of DLBCL tumors in this study compared to other prior DLBCL-focused

efforts[3,4]. We therefore focused our subsequent description on BL, FL and MCL.

The core set of co-associated genetic alterations in BL were *MYC* translocation and mutation,

and mutations of *SMARCA4*, *CCND3* and *ID3*. Mutations of *GNAI2*, *GNA13*, *FOXO1*, *TCF3*,

ARID1A and *TMEM51* were also clustered with this core set, but were present at lower

frequencies. Mutations of *TP53* and copy gain of 1q were frequent in BL, but did not cluster

together with other BL-associated genetic alterations. The core set of genetic alterations in FL

were *BCL2* translocations and mutations, and mutations of *KMT2D*, *CREBBP* and *EZH2*, in line

with prior observations[16,21]. Copy loss of 10q23.31 encompassing the *FAS* and *PTEN* genes,

and mutations *EEF1A1*, *TNFRSF14*, *IRF8*, *BCL7A*, *ATP6V1B2* and *ATP6AP1* were present at

lower frequencies and clustered with this core set. The MCL cluster included mutations and

deletions of *ATM* and *BIRC3*, as well as frequent deletions of 9p21.32 (*CDKN2A* and *CDKN2B*),

9q21.3 (*UBQLN1*), 13q14.2 (miR-15a/16 and TRIM13), 13q34 (*CUL4A* and *ING1*), and 1p21

(*RPL5*). Mutation of *BCOR*, *UBR5*, *SP140*, *CCND1* and *WHSC1* were also significantly over-

represented in MCL compared to other subtypes, and were associated with this cluster. These

data show that different subtypes of B-NHL are defined by characteristic sets of co-associated

genetic alterations that affect multiple hallmarks.

Combinations of clonal genetic alterations define B-NHL subtypes

Our data have revealed statistical enrichment of individual genetic alterations in subtypes of B-

NHL, and pairwise relationships between different genetic alterations that define clusters of

subtype-specific events. However, individual genetic alterations from disease-specific clusters

are also observed at variable frequencies in other B-NHL subtypes (**Figure 6A**). We therefore

investigated whether the combination of multiple genetic alterations from each cluster, rather

than the presence or absence of a single alteration, were pathognomonic. By enumerating the

number of genetic alterations from each disease-specific cluster in each tumor, we found that

this was indeed the case. Specifically over half of all BL, FL and MCL tumors possessed ≥3

genetic alterations defined by their respective subtype-specific clusters (**Figure 6**). These rates

were significantly higher than those observed in other diseases (Fisher P <0.001 for all

comparisons). For example, DLBCL and FL share many of the same recurrently mutated genes,

particularly when considering the C3/EZB subtype of DLBCL[3,4]. Thus, one or more FL cluster

alterations are found in 93% of FL tumors, but also in 54% of DLBCL tumors. However, FL

tumors have a significantly higher rate of accumulation of these mutations, with 68% of FL

tumors bearing ≥3 of the FL cluster alterations but only 18% of DLBCL tumors bearing ≥3 of

these genetic alterations (Fisher P < 0.001). This pattern is also conserved for the BL and MCL

clusters, for which the rate of acquiring ≥3 of the cluster alterations was significantly higher in BL

or MCL compared to other subtypes of B-NHL, respectively. Thus, B-NHL subtypes are defined

by the acquisition of multiple genetic alterations from subtype-associated clusters.

The co-association of multiple mutations may be suggestive of a clonal structure in which early

driver mutations are clonal and later driver mutations are present in sub-clones. We therefore

utilized ABSOLUTE (Table S16-18) to determine whether co-associated genetic alterations are

present at different clonal fractions, and could therefore be phased to create clonal hierarchies.

This analysis failed for a small subset of cases and could not be performed for X-linked genes

or translocations. However, it clearly revealed the majority of co-associated genetic alterations

were present at cancer cell fractions (CCF) >0.9, which is indicative that each mutation is

clonally represented in every cancer cell at the time of sampling. These data suggest that

combinations of mutations may therefore be required for effective expansion of lymphoma cells,

such that common precursor cells (CPCs) are not detectable within the clonal structure of the

clinically-detected tumor. These data therefore support the premise of epistatic interactions

between co-occurring genetic alterations in B-NHL.

DISCUSSION

By performing cross-sectional genomic profiling of a large cohort of tumors, we have defined

genes and functional hallmarks that are recurrently targeted by genetic alterations and showed

that combinations of genetic alterations are disease-defining features of B-NHL subtypes. Some

of the functional hallmarks that we identified have been previously appreciated in other

diseases, with a few exceptions. For example, the mutation of genes with roles in chromatin

modification are known to be a hallmark of FL[65] and we observed that over 90% of FLs

possessed mutations in one or more of the genes in this category. However, these mutations

were also observed in two thirds of BL and DLBCL tumors and nearly half of MCLs. There are

subtype-specific patterns of chromatin modifying gene alterations, such as those that we

described for H3K36 methylation in MCL and the unique patterns of SWI/SNF mutations across

GCB-derived B-NHLs. But we suggest that the genetic deregulation of chromatin modification

should be considered a general hallmark of B-NHL. In addition, we suggest that the

deregulation of BCL6 function and perturbation of the ubiquitin proteasome system are

hallmarks of B-NHL that require further investigation. We have highlighted some of the known

substrates of recurrently altered UPS genes from the literature, which shows the potential for

these genetic alterations to contribute to aberrant B-cell receptor signaling and proliferation.

However, many of the genes within these categories have not been functionally studied in the

context of B-cell lymphoma, which represents a significant gap in our understanding of disease

etiology.

The role of epistatic interactions between co-occurring genetic alterations is also an emerging

field that requires further investigation. These interactions are not uncommon in cancer[66], but

our data show that they are pervasive and disease-defining features of the B-NHL genetic

landscape. Epistasis between co-associated genetic alterations identified in this study requires

formal validation in cell line and/or animal models. However, there are many instances in which

co-occurring genetic alterations that we observed have already been shown to cooperate in

lymphomagenesis. In addition to the aforementioned example of *MYD88* and *CD79B* mutations,

transgenic mouse models of *Ezh2* activating mutations or conditional deletion of *Crebbp* or

Kmt2d have shown that these events are not alone sufficient for lymphomagenesis[42,67-71]. We

and others have observed a co-association between the mutation of these genes and *BCL2*

translocations[16,21], and the addition of a *Bcl2* transgene to these murine models indeed

promoted lymphoma at a significantly higher rate than that observed with the *Bcl2* transgene

alone[42,67-71]. These genetic alterations are therefore significantly more lymphomagenic in

combination than they are alone, which provides proof of principal that an epistatic relationship

exists between these co-occurring genetic alterations. Future studies focusing on other co-

occurring mutations, such as *MYC* translocation and *SMARCA4* mutation in BL, *CREBBP* and

KMT2D mutation in FL, *TCF4* copy gain and *MYD88* mutation in DLBCL, and *ATM* mutation and

RPL5 deletion in MCL, should therefore be performed to further explore these concepts and

define their underlying functional relationship. We suggest that combinations of genetic

alterations are likely to more accurately recapitulate the biology of B-NHL than single gene

models, and may reveal contextually different functional roles of genetic alterations depending

on the co-occurring events.

In conclusion, we have provided a framework of functional hallmarks and co-occurring genetic

alterations that are enriched within B-NHL subtypes. These genetic alterations likely represent

epistatic interactions that underpin the biology of these tumors, and represent an opportunity for

better understanding lymphoma etiology so that we can identify novel rational approaches for

therapeutic targeting of the underlying biology.

ACKNOWLEDGEMENTS

This research was supported by NCI R01CA201380 (MRG), the Nebraska Department of

Health and Human Services (LB506 2016-17; MRG), the Jaime Erin Follicular Lymphoma

Research Consortium (MRG and SSN), the Futcher Foundation (LN and MRG), the Schweitzer

Family Fund (JRW, RED, MRG), a Leukemia and Lymphoma Society Fellow Award (HY), and

NCI cancer center support grants to the University of Texas MD Anderson Cancer Center (P30

CA016672) and the Fred & Pamela Buffet Cancer Center (P30 CA036727).

AUTHOR CONTRIBUTIONS

MCJM and ST performed experiments, analyzed data and wrote the manuscript. AB analyzed

data. AB, TBH, HY, QD, DM, KH, NJ, JS, and SG performed experiments. AA, LS, MD, CC, JT,

DP, KMV, MAL, ARS, BJC, RB, SSN, LN, RED, JW, SP, MKG, DS, JI, TG, SR, and AM

provided samples and/or clinical data. MRG conceived and supervised the study, performed

experiments, analyzed the data and wrote the manuscript. All authors reviewed and approved

the manuscript.

DISCLOSURES OF CONFLICTS OF INTEREST

The authors have no conflicts of interest related to this work.

REFERENCES

Swerdlow, S. H. *et al.* The 2016 revision of the World Health Organization classification of lymphoid neoplasms. *Blood* **127**, 2375-2390, doi:10.1182/blood-2016-01-643569 (2016).

Armitage, J. O., Gascoyne, R. D., Lunning, M. A. & Cavalli, F. Non-Hodgkin lymphoma. *Lancet* **390**, 298-310, doi:10.1016/S0140-6736(16)32407-2 (2017).

Chapuy, B. *et al.* Molecular subtypes of diffuse large B cell lymphoma are associated with distinct pathogenic mechanisms and outcomes. *Nature medicine* **24**, 679-690, doi:10.1038/s41591-018-0016-8 (2018).

Schmitz, R. *et al.* Genetics and Pathogenesis of Diffuse Large B-Cell Lymphoma. *The New England journal of medicine* **378**, 1396-1407, doi:10.1056/NEJMoa1801445 (2018).

Reddy, A. *et al.* Genetic and Functional Drivers of Diffuse Large B Cell Lymphoma. *Cell* **171**, 481-494 e415, doi:10.1016/j.cell.2017.09.027 (2017).

Bea, S. *et al.* Landscape of somatic mutations and clonal evolution in mantle cell lymphoma. *Proceedings of the National Academy of Sciences of the United States of America* **110**, 18250-18255, doi:10.1073/pnas.1314608110 (2013).

Zhang, J. *et al.* The genomic landscape of mantle cell lymphoma is related to the epigenetically determined chromatin state of normal B cells. *Blood* **123**, 2988-2996, doi:10.1182/blood-2013-07-517177 (2014).

Wang, J. Q. *et al.* Synergistic cooperation and crosstalk between MYD88L265P and mutations that dysregulate CD79B and surface IgM. *The Journal of experimental medicine* **214**, 2759-2776, doi:10.1084/jem.20161454 (2017).

Lenz, G. *et al.* Stromal gene signatures in large-B-cell lymphomas. *The New England journal of medicine* **359**, 2313-2323, doi:10.1056/NEJMoa0802885 (2008).

Dave, S. S. *et al.* Molecular diagnosis of Burkitt's lymphoma. *The New England journal of medicine* **354**, 2431-2442, doi:10.1056/NEJMoa055759 (2006).

Iqbal, J. *et al.* Genome-wide miRNA profiling of mantle cell lymphoma reveals a distinct subgroup with poor prognosis. *Blood* **119**, 4939-4948, doi:10.1182/blood-2011-07-370122 (2012).

Bouska, A. *et al.* Adult High Grade B-cell Lymphoma with Burkitt Lymphoma Signature: Genomic features and Potential Therapeutic Targets. *Blood*, doi:10.1182/blood-2017-02-767335 (2017).

Khodadoust, M. S. *et al.* Antigen presentation profiling reveals recognition of lymphoma immunoglobulin neoantigens. *Nature* **543**, 723-727, doi:10.1038/nature21433 (2017).

Schmitz, R. *et al.* Burkitt lymphoma pathogenesis and therapeutic targets from structural and functional genomics. *Nature* **490**, 116-120, doi:10.1038/nature11378 (2012).

Love, C. *et al.* The genetic landscape of mutations in Burkitt lymphoma. *Nature genetics* **44**, 1321-1325, doi:10.1038/ng.2468 (2012).

447	16	Morin, R. D. *et al.* Frequent mutation of histone-modifying genes in non-Hodgkin lymphoma. *Nature* **476**, 298-303, doi:10.1038/nature10351 (2011).
449	17	Morin, R. D. *et al.* Mutational and structural analysis of diffuse large B-cell lymphoma using whole-genome sequencing. *Blood* **122**, 1256-1265, doi:10.1182/blood-2013-02-483727 (2013).
452	18	Pasqualucci, L. *et al.* Analysis of the coding genome of diffuse large B-cell lymphoma. *Nature genetics* **43**, 830-837, doi:10.1038/ng.892 (2011).
454	19	Lohr, J. G. *et al.* Discovery and prioritization of somatic mutations in diffuse large B-cell lymphoma (DLBCL) by whole-exome sequencing. *Proceedings of the National Academy of Sciences of the United States of America* **109**, 3879-3884, doi:10.1073/pnas.1121343109 (2012).
458	20	Green, M. R. *et al.* Hierarchy in somatic mutations arising during genomic evolution and progression of follicular lymphoma. *Blood* **121**, 1604-1611, doi:10.1182/blood-2012-09-457283 (2013).
461	21	Green, M. R. *et al.* Mutations in early follicular lymphoma progenitors are associated with suppressed antigen presentation. *Proceedings of the National Academy of Sciences of the United States of America* **112**, E1116-1125, doi:10.1073/pnas.1501199112 (2015).
464	22	Okosun, J. *et al.* Integrated genomic analysis identifies recurrent mutations and evolution patterns driving the initiation and progression of follicular lymphoma. *Nature genetics* **46**, 176-181, doi:10.1038/ng.2856 (2014).
467	23	Pasqualucci, L. *et al.* Genetics of follicular lymphoma transformation. *Cell reports* **6**, 130-140, doi:10.1016/j.celrep.2013.12.027 (2014).
469	24	Landau, D. A. *et al.* Evolution and impact of subclonal mutations in chronic lymphocytic leukemia. *Cell* **152**, 714-726, doi:10.1016/j.cell.2013.01.019 (2013).
471	25	Puente, X. S. *et al.* Whole-genome sequencing identifies recurrent mutations in chronic lymphocytic leukaemia. *Nature* **475**, 101-105, doi:10.1038/nature10113 (2011).
473	26	Wang, L. *et al.* SF3B1 and other novel cancer genes in chronic lymphocytic leukemia. *The New England journal of medicine* **365**, 2497-2506, doi:10.1056/NEJMoa1109016 (2011).
476	27	Kiel, M. J. *et al.* Whole-genome sequencing identifies recurrent somatic NOTCH2 mutations in splenic marginal zone lymphoma. *The Journal of experimental medicine* **209**, 1553-1565, doi:10.1084/jem.20120910 (2012).
479	28	Rossi, D. *et al.* The coding genome of splenic marginal zone lymphoma: activation of NOTCH2 and other pathways regulating marginal zone development. *The Journal of experimental medicine* **209**, 1537-1551, doi:10.1084/jem.20120904 (2012).
482	29	Chapman, M. A. *et al.* Initial genome sequencing and analysis of multiple myeloma. *Nature* **471**, 467-472, doi:10.1038/nature09837 (2011).
484	30	Treon, S. P. *et al.* MYD88 L265P somatic mutation in Waldenstrom's macroglobulinemia. *The New England journal of medicine* **367**, 826-833, doi:10.1056/NEJMoa1200710 (2012).
487	31	Gunawardana, J. *et al.* Recurrent somatic mutations of PTPN1 in primary mediastinal B cell lymphoma and Hodgkin lymphoma. *Nature genetics* **46**, 329-335, doi:10.1038/ng.2900 (2014).
490	32	Vater, I. *et al.* The mutational pattern of primary lymphoma of the central nervous system determined by whole-exome sequencing. *Leukemia* **29**, 677-685, doi:10.1038/leu.2014.264 (2015).
493	33	Kuilman, T. *et al.* CopywriteR: DNA copy number detection from off-target sequence data. *Genome biology* **16**, 49, doi:10.1186/s13059-015-0617-1 (2015).
495	34	Newman, A. M. *et al.* FACTERA: a practical method for the discovery of genomic rearrangements at breakpoint resolution. *Bioinformatics* **30**, 3390-3393, doi:10.1093/bioinformatics/btu549 (2014).

35   Lek, M. *et al.* Analysis of protein-coding genetic variation in 60,706 humans. *Nature* **536**,
285-291, doi:10.1038/nature19057 (2016).
36   Lawrence, M. S. *et al.* Mutational heterogeneity in cancer and the search for new
cancer-associated genes. *Nature* **499**, 214-218, doi:10.1038/nature12213 (2013).
37   Mermel, C. H. *et al.* GISTIC2.0 facilitates sensitive and confident localization of the
targets of focal somatic copy-number alteration in human cancers. *Genome biology* **12**,
R41, doi:10.1186/gb-2011-12-4-r41 (2011).
38   Carter, S. L. *et al.* Absolute quantification of somatic DNA alterations in human cancer.
Nature biotechnology **30**, 413-421, doi:10.1038/nbt.2203 (2012).
39   Alexandrov, L. B. *et al.* Signatures of mutational processes in human cancer. *Nature*
500, 415-421, doi:10.1038/nature12477 (2013).
40   Huang, D. W. *et al.* DAVID Bioinformatics Resources: expanded annotation database
and novel algorithms to better extract biology from large gene lists. *Nucleic acids*
research **35**, W169-175, doi:10.1093/nar/gkm415 (2007).
41   Saeki, K., Miura, Y., Aki, D., Kurosaki, T. & Yoshimura, A. The B cell-specific major raft
protein, Raftlin, is necessary for the integrity of lipid raft and BCR signal transduction.
EMBO J **22**, 3015-3026, doi:10.1093/emboj/cdg293 (2003).
42   Jiang, Y. *et al.* CREBBP Inactivation Promotes the Development of HDAC3-Dependent
Lymphomas. *Cancer discovery* **7**, 38-53, doi:10.1158/2159-8290.CD-16-0975 (2017).
43   Beguelin, W. *et al.* EZH2 and BCL6 Cooperate to Assemble CBX8-BCOR Complex to
Repress   Bivalent   Promoters,   Mediate   Germinal   Center   Formation   and
Lymphomagenesis. *Cancer cell* **30**, 197-213, doi:10.1016/j.ccell.2016.07.006 (2016).
44   Pawar, S. A. *et al.* C/EBP{delta} targets cyclin D1 for proteasome-mediated degradation
via induction of CDC27/APC3 expression. *Proceedings of the National Academy of*
Sciences   of   the   United   States   of   America **107**,   9210-9215,
doi:10.1073/pnas.0913813107 (2010).
45   Akasaka, T., Lossos, I. S. & Levy, R. BCL6 gene translocation in follicular lymphoma: a
harbinger of eventual transformation to diffuse aggressive lymphoma. *Blood* **102**, 1443-
1448, doi:10.1182/blood-2002-08-2482 (2003).
46   Kato, M. *et al.* Frequent inactivation of A20 in B-cell lymphomas. *Nature* **459**, 712-716,
doi:10.1038/nature07969 (2009).
47   Greiner, T. C. *et al.* Mutation and genomic deletion status of ataxia telangiectasia
mutated (ATM) and p53 confer specific gene expression profiles in mantle cell
lymphoma. *Proceedings of the National Academy of Sciences of the United States of*
America **103**, 2352-2357, doi:10.1073/pnas.0510441103 (2006).
48   Challa-Malladi, M. *et al.* Combined genetic inactivation of beta2-Microglobulin and CD58
reveals frequent escape from immune recognition in diffuse large B cell lymphoma.
Cancer cell **20**, 728-740, doi:10.1016/j.ccr.2011.11.006 (2011).
49   Pfeifer, M. *et al.* PTEN loss defines a PI3K/AKT pathway-dependent germinal center
subtype of diffuse large B-cell lymphoma. *Proceedings of the National Academy of*
Sciences   of   the   United   States   of   America **110**,   12420-12425,
doi:10.1073/pnas.1305656110 (2013).
50   Jain N, H. K., Tadros S, Fiskus W, Havranek O, Ma M, Bouska A, Heavican T, Kumar D,
Deng Q, Moore D, Pak C, Liu C, Gentles A, Hartmann E, Kridel R, Ekstrom-Smedby K,
Juliusson G, Rosenquist R, Gascoyne R, Rosenwald A, Giancotti F, Neelapu S, Westin
J, Vose J, Lunning M, Greiner T, Rodig S, Iqbal J, Alizadeh A, Davis RE, Bhalla K,
Green M. Targetable genetic alterations of TCF4 (E2-2) drive immunoglobulin
expression in the activated B-cell subtype of diffuse large B-cell lymphoma. *Sci. Transl.*
Med. **In Press** (2019).

51   Ingley, E. *et al.* Csk-binding protein mediates sequential enzymatic down-regulation and
degradation of Lyn in erythropoietin-stimulated cells. *The Journal of biological chemistry*
281, 31920-31929, doi:10.1074/jbc.M602637200 (2006).

52   Hu, S., Alcivar, A., Qu, L., Tang, J. & Yang, X. CIAP2 inhibits anigen receptor signaling
by targeting Bcl10 for degredation. *Cell cycle* **5**, 1438-1442, doi:10.4161/cc.5.13.2866
(2006).

53   Compagno, M. *et al.* Mutations of multiple genes cause deregulation of NF-kappaB in
diffuse large B-cell lymphoma. *Nature* **459**, 717-721, doi:10.1038/nature07968 (2009).

54   Liu, W. H. & Lai, M. Z. Deltex regulates T-cell activation by targeted degradation of
active MEKK1. *Molecular and cellular biology* **25**, 1367-1378,
doi:10.1128/MCB.25.4.1367-1378.2005 (2005).

55   Liu, W. *et al.* Direct inhibition of Bruton's tyrosine kinase by IBtk, a Btk-binding protein.
Nature immunology **2**, 939-946, doi:10.1038/ni1001-939 (2001).

56   Joo, H. M. *et al.* Ret finger protein 2 enhances ionizing radiation-induced apoptosis via
degradation of AKT and MDM2. *Eur J Cell Biol* **90**, 420-431,
doi:10.1016/j.ejcb.2010.12.001 (2011).

57   Davis, R. E. *et al.* Chronic active B-cell-receptor signalling in diffuse large B-cell
lymphoma. *Nature* **463**, 88-92, doi:10.1038/nature08638 (2010).

58   Ngo, V. N. *et al.* Oncogenically active MYD88 mutations in human lymphoma. *Nature*
470, 115-119, doi:10.1038/nature09671 (2011).

59   Krysiak, K. *et al.* Recurrent somatic mutations affecting B-cell receptor signaling pathway
genes in follicular lymphoma. *Blood* **129**, 473-483, doi:10.1182/blood-2016-07-729954
(2017).

60   Gloury, R. *et al.* Dynamic changes in Id3 and E-protein activity orchestrate germinal
center and plasma cell development. *The Journal of experimental medicine* **213**, 1095-
1111, doi:10.1084/jem.20152003 (2016).

61   Okosun, J. *et al.* Recurrent mTORC1-activating RRAGC mutations in follicular
lymphoma. *Nature genetics* **48**, 183-188, doi:10.1038/ng.3473 (2016).

62   Wang, F. *et al.* Follicular lymphoma-associated mutations in vacuolar ATPase
ATP6V1B2 activate autophagic flux and mTOR. *The Journal of clinical investigation* **130**,
1626-1640, doi:10.1172/JCI98288 (2019).

63   Hu, J. *et al.* MiR-361-3p regulates ERK1/2-induced EMT via DUSP2 mRNA degradation
in pancreatic ductal adenocarcinoma. *Cell Death Dis* **9**, 807, doi:10.1038/s41419-018-
0839-8 (2018).

64   Lu, D. *et al.* The phosphatase DUSP2 controls the activity of the transcription activator
STAT3 and regulates TH17 differentiation. *Nature immunology* **16**, 1263-1273,
doi:10.1038/ni.3278 (2015).

65   Green, M. R. Chromatin modifying gene mutations in follicular lymphoma. *Blood* **131**,
595-604, doi:10.1182/blood-2017-08-737361 (2018).

66   Ashworth, A., Lord, C. J. & Reis-Filho, J. S. Genetic interactions in cancer progression
and treatment. *Cell* **145**, 30-38, doi:10.1016/j.cell.2011.03.020 (2011).

67   Beguelin, W. *et al.* EZH2 is required for germinal center formation and somatic EZH2
mutations promote lymphoid transformation. *Cancer cell* **23**, 677-692,
doi:10.1016/j.ccr.2013.04.011 (2013).

68   Garcia-Ramirez, I. *et al.* Crebbp loss cooperates with Bcl2 over-expression to promote
lymphoma in mice. *Blood* **129**, 2645-2656, doi:10.1182/blood-2016-08-733469 (2017).

69   Zhang, J. *et al.* The Crebbp Acetyltransferase is a Haploinsufficient Tumor Suppressor in
B Cell Lymphoma. *Cancer discovery*, doi:10.1158/2159-8290.CD-16-1417 (2017).

70   Zhang, J. *et al.* Disruption of KMT2D perturbs germinal center B cell development and
promotes lymphomagenesis. *Nature medicine* **21**, 1190-1198, doi:10.1038/nm.3940
(2015).

71   Ortega-Molina, A. *et al.* The histone lysine methyltransferase KMT2D sustains a gene
expression program that represses B cell lymphoma development. *Nature medicine* **21**,
1199-1208, doi:10.1038/nm.3943 (2015).

FIGURE LEGENDS

Figure 1: Recurrently mutated genes in B-NHL subtypes. An oncoplot shows significantly

mutated genes across our cohort of 755 B-NHL tumors, arranged according to functional

category and frequency. The mutational burden and distribution of mutation types for each case

are shown at the top. Amino acid residues for mutational hotspots and/or domains targeted by

clustered mutations are shown on the right.

Figure 2: Structural alterations in B-NHL subtypes. A) A circos plot shows translocations of

the MYC (purple), BCL2 (orange) and BCL6 (green) genes, and GISTIC tracks of DNA copy

number gains (red) and losses (blue). **B-C)** Volcano plots of integrative analysis results showing

the changes in gene expression of genes within peaks of DNA copy number gain (B) or loss (C).

The shaded region in B marks genes with significantly increased expression in tumors with

increased copy number compared to those without; the converse is true for the shaded region in

C. **D)** An oncoplot with DNA copy number losses and gains ranked according to their frequency

shows the distribution of structural alterations across tumors. E) Oncoplots show the overlap of

structural alterations and mutations that target the same genes. P-values are derived from a

Fisher's exact test (ns, not significant).

Figure 3: Genetic alterations of the ubiquitin proteasome system (UPS) and potential interplay with BCR signaling. A) A graphical illustration of UPS gene genetic alterations (mutations, teal; deletions, blue; copy gain, red) and literature supported targets of the UPS genes. The function of UPS gene targets is categorized according to the hallmarks shown in figure 1. **B)** A schematic of the different layers of genetic alterations (mutations, teal; deletions, blue; copy gain, red) affecting signaling pathways in B-NHL. UPS genes are highlighted with a yellow halo.

Figure 4: Subtype-specific clusters of co-occurring genetic alterations. The frequency (bar graph) and over/under-representation (blue to yellow scale) of mutations and structural alterations is shown on the left for BL, DLBCL, FL and MCL. Over/under-representation in molecular Burkitt and cell of origin molecular subtypes is shown for each alteration at the bottom. The correlation matrix of co-associated (green) and mutually-exclusive (purple) relationships was clustered and identified groups of co-occurring genetic alterations that were predominantly over-represented in a single B-NHL subtype.

Figure 5: Subtype-specific patterns of SWI/SNF complex mutations. A) An oncoplot shows the frequency of genetic alterations in genes that encode components of the SWI/SNF complex. **B)** A schematic of the SWI/SNF complex shows recurrently mutated genes, *ARID1A*, *SMARCA4* and *BCL7A,* and the *BCL11A* gene that is targeted by 2p15 DNA copy number gains. **C-E)** Lollipop plots show the distribution of mutations in the SWI/SNF components ARID1A (C), SMARCA4 (D), and BCL7A (E). **F)** A heatplot shows the location of chromosome 2p DNA copy number gains (red) in all tumors (rows) with 2p15 copy gain (n=92, copy number > 2.2) ordered from highest DNA copy number (top) to lowest (bottom). The *BCL11A* gene is in the peak focal copy gain.

Figure 6: Cumulative burden of clonal co-occurring mutations. A) A bar plot shows the frequency of tumors with genetic alterations from each subtype-defining cluster, grouped by the number of co-occurring alterations observed per tumor. The frequency of tumors with 3 or more genetic alterations per tumor is significantly higher in the subtype for which each cluster is named (***Fisher P-value < 0.001). **B-D)** An oncoplot, colored by cancer cell fraction (CCF) from ABSOLUTE, shows the representation and clonality of BL cluster (B), FL cluster (C) and MCL cluster (D) mutations across the dataset. The rate of co-occurrence of multiple genetic alterations in a single tumor (columns) is observably higher in the subtypes for which the

667   clusters are named, and the co-occurring genetics are most often clonal (CCF > 0.9). BL, Burkitt

668   Lymphoma. HG, other high-grade lymphoma (DHL/THL, HGBL-NOS). DLBCL, diffuse large B-

669   cell lymphoma. FL, follicular lymphoma. MCL, mantle cell lymphoma. LG, other low-grade

670   lymphoma (CLL/SLL, MZL).

Figure 1

Figure 2

Figure 3

A

B

Figure 4

Figure 5

A

	BL	DLBCL	FL	MCL
ARID1A	19%	7%	15%	6%
SMARCA4	24%	3%	8%	6%
BCL7A	7%	7%	11%	0%
BCL11A	11%	21%	18%	2%

Alteration Type: ■ Multiple ■ Nonsense/Frameshift ■ Missense ■ Copy gain

Figure 6