1 **Temporal and spatial limitations in global surveillance for bat filoviruses and**
2 **henipaviruses**
3

4 Daniel J. Becker[1-3*], Daniel E. Crowley[1], Alex D. Washburne[1], Raina K. Plowright[1]
5
[1]Department of Microbiology and Immunology, Montana State University, Bozeman, Montana
[2]Department of Biology, Indiana University, Bloomington, Indiana
[3]Center for the Ecology of Infectious Diseases, University of Georgia, Athens, Georgia
[*]danbeck@iu.edu

Running head: Spatiotemporal bat virus dynamics
Keywords: Chiroptera; spillover; sampling design; zoonotic virus; phylogenetic meta-analysis;
spatiotemporal

Abstract
Sampling reservoir hosts over time and space is critical to detect epizootics, predict spillover,
and design interventions. However, because sampling is logistically difficult and expensive,
researchers rarely perform spatiotemporal sampling of many reservoir hosts. Bats are reservoirs
of many virulent zoonotic pathogens such as filoviruses and henipaviruses, yet the highly mobile
nature of these animals has limited optimal sampling of bat populations. To quantify the
frequency of temporal sampling and to characterize the geographic scope of bat virus research,
we here collated data on filovirus and henipavirus prevalence and seroprevalence in wild bats.
We used a phylogenetically controlled meta-analysis to next assess temporal and spatial variation
in bat virus detection estimates. Our analysis shows that only one in four bat virus studies report
data longitudinally, that sampling efforts cluster geographically (e.g., filovirus data are available
across much of Africa and Asia but are absent from Latin America and Oceania), and that
sampling designs and reporting practices may affect some viral detection estimates (e.g.,
filovirus seroprevalence). Within the limited number of longitudinal bat virus studies, we
observed high heterogeneity in viral detection estimates that in turn reflected both spatial and
temporal variation. This suggests that spatiotemporal sampling designs are essential to
understand how zoonotic viruses are maintained and spread within and across wild bat
populations, which in turn could help predict and preempt risks of zoonotic viral spillover.

Introduction

Risks of pathogen spillover vary across time and space [1,2], in part because pathogen shedding from reservoir hosts is a dynamic spatiotemporal processes [3,4]. Metapopulation dynamics and other spatial processes characterize many reservoir hosts [5], where populations connectivity can determine the spatiotemporal distribution of a pathogen [6,7] and degree of spatial synchrony structuring infection dynamics [8]. Temporal pulses of shedding driven by seasonality in birth and climate are also common [9,10]. Understanding how infection in reservoir hosts varies over space and time is thus a critical need for predicting and managing zoonotic disease risks.

However, surveillance strategies often do not sample this underlying spatiotemporal process, as spatially and temporally explicit designs present logistical challenges when studying mobile and gregarious species [3,11,12]. For hosts such as birds and bats, surveillance is often opportunistic or relies on convenience sampling [13]. These non-probabilistic and often single sampling events cannot characterize spatial and temporal fluctuations in infection, can over- or under-represent times or locations of high prevalence, and can result in non-randomly missing data [3,14]. These challenges cannot be fixed with statistical modeling and can bias estimates of prevalence and epidemiological parameters such as the basic reproductive number [13,15].

Given a fixed cost, difficult decisions must be made about how to allocate sampling efforts. Sampling over space facilitates detecting geographic clusters of disease and predictive mapping [16,17], while sampling over time can identify periods of intensive pathogen shedding and enable inference about dominant transmission routes [18,19]. Researchers often treat this as a tradeoff between sampling over either time or space, rather than allocating effort to both [20]. Implicit here is that the temporal component is constant over space or that the spatial component is constant over time, and such sampling designs result in no data to assess this assumption.

We here quantify the temporal and spatial data limitations for two taxa of high-profile zoonotic viruses of bats: the family *Filoviridae* and genus *Henipavirus*. Bats have been widely studied as reservoirs for zoonotic pathogens and host more viruses with zoonotic potential than other mammals [21,22]. Henipaviruses and some filoviruses (e.g., Marburg virus) can be shed from bats into the environment [23,24] and can cause fatal disease in humans by environmental exposure or from contact with intermediate hosts such as horses, wild primates, or pigs [25–30]. Current evidence suggests many filo- and henipaviruses show variable dynamics in space and time, including shedding pulses from bats [6,25,31–33], which implies that spatiotemporal sampling is critical to capture viral dynamics in bat reservoirs. Yet while past efforts have focused on bat virus discovery [34], the determinants of reservoir status [35], and experimental mechanisms of viral transmission [36], spatiotemporal studies of bat–virus dynamics are rare [37]. This limits understanding how zoonotic viruses are maintained and spread within and across bat populations and impairs improving future sampling designs and ecological interventions [20,38]. We here systematically collated data on the prevalence and seroprevalence of filo- and henipaviruses in wild bats to (*i*) quantify the frequency of temporal studies and (*ii*) assess the geographic scope of current research. We used phylogenetic meta-analysis to (*iii*) quantify how sampling designs and reporting practices may influence viral detection estimates. Single snapshots could miss pulses of viral shedding from bats, whereas pooling data over time could under- or overestimate viral presence [18,20]. Lastly, we (*iv*) characterized the degree of temporal and spatial variation in bat virus detection estimates.

Methods

To systematically identify studies quantifying the proportion of wild bats positive for filoviruses
and henipaviruses using PCR or serology, we searched Web of Science, CAB Abstracts, and
PubMed (see Fig. S1). Our dataset included 1176 records from 68 studies. Viruses included not
only Hendra, Nipah, Ebola, and Marburg virus but also Lloviu and Reston virus. We grouped
viruses by taxa given our sample sizes and known issues of serological cross-reactivity [39,40].

From each study, we defined sampling subunits: a temporally defined sampling event of
one bat species in one location per viral detection estimate. Each subunit is the lowest spatial,
temporal, and phylogenetic scale (of bats and their viruses) reported. We classified subunits into
three sampling designs and reporting practices: one sampling event, multiple events, or pooled
events over time. Records of a single prevalence or seroprevalence estimate from a population
sampled from a period less than or equal to one month were classified as single sampling events,
whereas records of a population over multiple monthly timepoints were classified as spanning
multiple events (i.e., a longitudinal study). For example, every monthly prevalence estimate per
population of *Pteropus lylei* in Thailand would represent a unique subunit and be classified as
longitudinal [41]. Records of a period longer than one month were classified as pooled events,
where researchers may have sampled a population across more than one timepoint but reported
data as a single viral detection estimate. A schematic of these categorizations is provided in
Figure 1A. One month was selected because this timeframe was the lowest common temporal
unit and because bat shedding of these viruses can occur within a month [36,42]. These data
were reported for most records (1121/1176 subunits; three publications did not report these data
and three additional publications did not always report such data for all records). For each
subunit, we also recorded the bat species, virus taxon, coarse detection method (i.e., PCR or
serology), number of bats sampled, proportion of bats positive, sampling timepoints, sampling
location, and country (recoded to the United Nations geoscheme for our descriptive analyses).

We quantified the proportion of studies using each sampling and reporting design, both
across all data and stratified by virus taxon. To assess how the frequency of longitudinal studies
(i.e., those with repeated sampling) has changed over time, we fit a generalized additive model
with the *mgcv* package in R and a smooth term for publication year [43]. We also calculated the
duration of repeat sampling for these longitudinal studies. For studies that pooled data over time,
we quantified days represented per subunit. To describe geographic biases in bat virus studies,
we assessed sampling gaps according to region (United Nations geoscheme). We used a χ^2 test to
assess if sampling designs and reporting practices were differently distributed across regions.

To assess the contribution of sampling designs and reporting practices to viral detection
estimates and to quantify the degree of spatial and temporal variation in bat–virus interactions,
we used the *metafor* package to calculate logit-transformed proportions and sampling variances
and to fit hierarchical meta-analysis models [44,45]. To account for phylogenetic dependence,
we included bat species as a random effect [46], for which the covariance structure used the
phylogenetic correlation matrix; we obtained our phylogeny from the Open Tree of Life with the
rotl and *ape* packages [47,48]. We excluded subunits that pooled data across or within bat genera
(*n*=102). As few subunits (*n*=14) pooled data across specified species in a genus, we randomly
selected one species to retain these records. Our final dataset included 1019 subunits from 60
studies and 215 bat species (Fig. S2). Our models also included subunit nested within study as a
random effect and weighting by sampling variances. To first assess heterogeneity among viral
detection estimates, we fit a random-effects model (REM; intercept only) and stratified this
analysis per viral taxon and detection method. We used restricted maximum likelihood to obtain
unbiased estimates of the variance components, from which we derived I^2 to quantify the

contribution of true heterogeneity to total variance in viral detection estimates [49]. We used
these estimates to partition variance attributed to each random effect; in the case of bat species,
we derived phylogenetic heritability (H^2) as a measure of phylogenetic signal [46]. We used
Cochran's Q to test if such heterogeneity was greater than expected by sampling error alone [50].

To next test how sampling designs and reporting practices may influence viral detection
estimates, we fit a mixed-effects model (MEM) with the same random effects and an interaction
between sampling design and reporting practices, detection method, and virus taxon. We tested
significance of moderators and interactions using the Q test [44] and derived a pseudo-R^2 as the
proportional reduction in the summed variance components compared with those of a REM [51].

To test if viral detection estimates showed spatiotemporal variation, we fit models with
the same random effects to our data subset reporting multiple events (n=273). We fit a REM to
quantify I^2 for longitudinal studies. We then fit MEMs with location and month as univariate
moderators to test if viral detection estimates varied across space and time. Because this subset
of the data included many unique locations (n=28) and months (n=12), we did not use interaction
terms and instead fit an additional set of MEMs to each viral taxon–detection method strata.

Results
Only 26% of bat virus studies reported data longitudinally (10 filo- and 9 henipavirus studies;
Fig. 1). However, the frequency of such studies has weakly increased over time (Fig. S3,
χ^2_1=2.75, p=0.1). Eleven studies reported sampling populations 2–3 times while 12 reported
sampling populations over four times. The duration of longitudinal studies ranged from 150 days
to over 10 years, on average spanning 2.5 years of repeat sampling (Fig. S4). In contrast, half of
our studies (n=34) instead reported estimates across multiple timepoints as pooled proportions,
which on average represented 644 days of temporally aggregated data (SD=492; Fig. S5).

Bat sampling also showed geographic biases (Fig. 1, Table 1). Filovirus studies were
conducted across much of Africa and Asia but not in Latin America and Oceania. PCR and
serology have been used in the same region in most areas, but only one or the other have been
used in Southern Africa for filoviruses and in Europe, Eastern and Middle Africa, and Eastern
Asia for henipaviruses (Table 1). Geography was also associated with sampling design and
reporting practices (χ^2=369.3, p=0.001). Longitudinal data were only reported from Central,
Eastern, Middle, and Southern Africa for filoviruses and only reported from Southeastern Asia,
Eastern Africa, and Oceania for henipaviruses (Table 1).

We observed significant heterogeneity across viral detection estimates (I^2=0.91,
Q_{1017}=6929, p<0.001). Bat species and study accounted for most variation ($I^2_{species}$=0.41,
I^2_{study}=0.34, H^2=0.45; Table S2). We also found significant heterogeneity within each viral
taxon–detection strata, although I^2 and H^2 values varied across these subsets (Table S1). Viral
detection estimates for henipaviruses had much stronger phylogenetic signal than filoviruses.

Our MEM showed that viral detection estimates broadly varied with detection method
(Q_1=5.41, p=0.02; seroprevalence was generally higher than prevalence) and were associated
with sampling design and reporting; however, the effect tended to depended on virus taxa and
detection method (three-way interaction: Q_2=5.36, p=0.07, R^2=0.06; Table S2). A post-hoc
analysis with MEMs fit to each strata showed sampling design and reporting were associated
with filovirus seroprevalence (Q_2=10.30, p=0.006; Fig. 2), with longitudinal studies generally

showing higher proportions of positive bats. Sampling design and reporting had no effects on
henipavirus seroprevalence nor prevalence estimates for either virus taxon (Table S3).

We also detected high variation in viral detection estimates across longitudinal studies
(Q_{271}=2866, p<0.0001, I^2=0.94; Fig. 2). Study contributed more to residual variance than
phylogeny ($I^2_{species}$=0.28, I^2_{study}=0.52, $I^2_{subunit}$=0.13). Across these data, location did not predict
viral detection estimates (Q_{28}=17.67, p=0.91); however, MEMs fit to each strata showed high
spatial variation for all subsets except filovirus prevalence (Table S4). Month also had little
predictive power across all longitudinal data (Q_{11}=6.93, p=0.80), but separate MEMs revealed
high temporal variation for filovirus seroprevalence and henipavirus prevalence (Table S5).

Discussion

Our study provides a systematic synthesis of prevalence and seroprevalence for bat filoviruses
and henipaviruses that can guide future sampling. Only one in four studies reported longitudinal
data, although use of such approaches is increasing. Half of studies instead pooled data over time
(and space). Geographic limitations were also evident, especially for where longitudinal studies
have been conducted. This was especially evident for filoviruses; although the absence of studies
in Latin America and Oceania may reflect the lack of reported human cases, bat reservoirs are
predicted to occur in both regions [35]. Many studies also used either PCR or serology, although
using both may improve statistical inference about how zoonotic pathogens persist in hosts [18].

We found generally weak evidence that such variation in sampling design and reporting
affected viral detection estimates, although filovirus seroprevalence tended to be greatest from
longitudinal studies. Serological surveys of Marburg and Ebola virus have found strong temporal
dynamics that may reflect seasonality in bat reproduction or resource availability [31,52,53].
Detection estimates could be higher with repeated sampling, given that such studies are more
likely to detect shedding pulses and pooling of data could increase zeros in the numerator
(underestimating seroprevalence). The lack of a similar pattern for filovirus PCR data could
result from low prevalence and be biased by zero inflation. We also qualify that our low R^2,
alongside high contributions of bat phylogeny and study random effects, suggests other aspects
of bat ecology (e.g., seasonal birth [31,54]) or study idiosyncrasies (e.g., serological cutoffs
[39,40]) may play more critical roles in shaping viral detection estimates. High H^2 for
henipaviruses suggests cladistic or trait-based analyses of shedding could be insightful [35,55].
Yet given the potential for sampling design and reporting to affect viral detection estimates, we
encourage researchers to publish data at the lowest spatial, temporal, and phylogenetic scale
associated with sampling and to provide data at such scales to facilitate these future analyses.

Our analysis of longitudinal studies found significant spatial and temporal variation in
some bat virus data. This implies spatiotemporal sampling is critical to make inferences about
bat virus spillover. Although sampling over space and time is challenging, especially for highly
mobile animals like bats, sampling can be informed by spatiotemporal variation in prevalence
and seroprevalence and analyses of spatiotemporal autocorrelation [20,56]. Greater variation
over space can require more fine-scale spatial sampling, and greater variation over time can
require more fine-scale temporal sampling. Spatiotemporal designs, such as stratified random
sampling or rotating panels, can help capture spatial and temporal variation in virus shedding
while also addressing some logistical challenges [13,57,58]. The increased adoption of such
approaches, especially in the understudied regions identified in our analysis, will be key to
improve understanding bat virus dynamics and how spillover risk varies over time and space.

Data availability
Data are available in the Dryad Digital Repository [59].

Author contributions
DJB and DEC designed the study, DEC collected data, DJB analyzed data, and all authors contributed to writing the manuscript.

Funding
The authors were supported by the National Science Foundation (DEB-1716698), the Defense Advanced Research Projects Agency (Young Faculty Award D16AP00113 and PREEMPT award D18AC0003), and USDA National Institute of Food and Agriculture (Hatch project 1015891). The content of the information does not necessarily reflect the position or the policy of the U.S. government, and no official endorsement should be inferred.

Acknowledgements
We thank Megan Higgs and anonymous reviewers for helpful comments on this manuscript.

References

1. Plowright RK, Parrish CR, McCallum H, Hudson PJ, Ko AI, Graham AL, Lloyd-Smith JO. 2017 Pathways to zoonotic spillover. *Nat. Rev. Microbiol.* **15**, 502–510.

2. Schmidt JP, Park AW, Kramer AM, Han BA, Alexander LW, Drake JM. 2017 Spatiotemporal Fluctuations and Triggers of Ebola Virus Spillover. *Emerg. Infect. Dis.* **23**, 415–422. (doi:10.3201/eid2303.160101)

3. Hoye BJ, Munster VJ, Nishiura H, Klaassen M, Fouchier RAM. 2010 Surveillance of Wild Birds for Avian Influenza Virus. *Emerg. Infect. Dis.* **16**, 1827–1834. (doi:10.3201/eid1612.100589)

4. Plowright RK *et al.* 2015 Ecological dynamics of emerging bat virus spillover. *Proc R Soc B* **282**, 20142124. (doi:10.1098/rspb.2014.2124)

5. Vergara PM, Saravia-Zepeda A, Castro-Reyes N, Simonetti JA. 2016 Is metapopulation patch occupancy in nature well predicted by the Levins model? *Popul. Ecol.* , 1–9. (doi:10.1007/s10144-016-0550-5)

6. Plowright RK, Foley P, Field HE, Dobson AP, Foley JE, Eby P, Daszak P. 2011 Urban habituation, ecological connectivity and epidemic dampening: the emergence of Hendra virus from flying foxes (Pteropus spp.). *Proc. R. Soc. B Biol. Sci.* **278**, 3703–3712. (doi:10.1098/rspb.2011.0522)

7. Hess G. 1996 Disease in Metapopulation Models: Implications for Conservation. *Ecology* **77**, 1617–1632. (doi:10.2307/2265556)

8. Bjørnstad ON, Ims RA, Lambin X. 1999 Spatial population dynamics: analyzing patterns and processes of population synchrony. *Trends Ecol. Evol.* **14**, 427–432.

9. Altizer S, Dobson A, Hosseini P, Hudson P, Pascual M, Rohani P. 2006 Seasonality and the dynamics of infectious diseases. *Ecol. Lett.* **9**, 467–484.

10. Peel AJ, Pulliam JRC, Luis AD, Plowright RK, O'Shea TJ, Hayman DTS, Wood JLN, Webb CT, Restif O. 2014 The effect of seasonal birth pulses on pathogen persistence in wild mammal populations. *Proc. R. Soc. Lond. B Biol. Sci.* **281**, 20132962. (doi:10.1098/rspb.2013.2962)

11. Kuiken T, Leighton FA, Fouchier R a. M, LeDuc JW, Peiris JSM, Schudel A, Stöhr K, Osterhaus ADME. 2005 Pathogen Surveillance in Animals. *Science* **309**, 1680–1681. (doi:10.1126/science.1113310)

12. Stallknecht DE. 2007 Impediments to Wildlife Disease Surveillance, Research, and Diagnostics. In *Wildlife and Emerging Zoonotic Diseases: The Biology, Circumstances and Consequences of Cross-Species Transmission*, pp. 445–461. Springer, Berlin, Heidelberg. (doi:10.1007/978-3-540-70962-6_17)

13. Nusser SM, Clark WR, Otis DL, Huang L. 2008 Sampling Considerations for Disease
Surveillance in Wildlife Populations. *J. Wildl. Manag.* **72**, 52–60.

14. Venette RC, Moon RD, Hutchison WD. 2002 Strategies and statistics of sampling for rare
individuals. *Annu. Rev. Entomol.* **47**, 143–174.
(doi:10.1146/annurev.ento.47.091201.145147)

15. Hayman DTS, Suu-Ire R, Breed AC, McEachern JA, Wang L, Wood JLN, Cunningham AA.
2008 Evidence of Henipavirus Infection in West African Fruit Bats. *PLOS ONE* **3**, e2739.
(doi:10.1371/journal.pone.0002739)

16. Peterson AT. 2006 Ecologic Niche Modeling and Spatial Patterns of Disease Transmission.
Emerg. Infect. Dis. **12**, 1822–1826. (doi:10.3201/eid1212.060373)

17. Wang J-F, Stein A, Gao B-B, Ge Y. 2012 A review of spatial sampling. *Spat. Stat.* **2**, 1–14.
(doi:10.1016/j.spasta.2012.08.001)

18. Plowright RK, Peel AJ, Streicker DG, Gilbert AT, McCallum H, Wood J, Baker ML, Restif
O. 2016 Transmission or Within-Host Dynamics Driving Pulses of Zoonotic Viruses in
Reservoir–Host Populations. *PLoS Negl. Trop. Dis.* **10**, e0004796.
(doi:10.1371/journal.pntd.0004796)

19. Restif O *et al.* 2012 Model-guided fieldwork: practical guidelines for multidisciplinary
research on wildlife ecological and epidemiological dynamics. *Ecol. Lett.* **15**, 1083–1094.

20. Plowright RK, Becker DJ, McCallum H, Manlove KR. 2019 Sampling to elucidate the
dynamics of infections in reservoir hosts. *Philos. Trans. R. Soc. B*
(doi:10.1098/rstb.2018.0336)

21. Luis AD *et al.* 2013 A comparison of bats and rodents as reservoirs of zoonotic viruses: are
bats special? *Proc R Soc B* **280**, 20122753. (doi:10.1098/rspb.2012.2753)

22. Olival KJ, Hosseini PR, Zambrana-Torrelio C, Ross N, Bogich TL, Daszak P. 2017 Host and
viral traits predict zoonotic spillover from mammals. *Nature* **546**, 646–650.
(doi:10.1038/nature22975)

23. Amman BR *et al.* 2015 Oral shedding of marburg virus in experimentally infected egyptian
fruit bats (rousettus aegyptiacus). *J. Wildl. Dis.* **51**, 113–124. (doi:10.7589/2014-08-198)

24. Middleton DJ, Morrissy CJ, van der Heide BM, Russell GM, Braun MA, Westbury HA,
Halpin K, Daniels PW. 2007 Experimental Nipah Virus Infection in Pteropid Bats (Pteropus
poliocephalus). *J. Comp. Pathol.* **136**, 266–272. (doi:10.1016/j.jcpa.2007.03.002)

25. Amman BR *et al.* 2012 Seasonal Pulses of Marburg Virus Circulation in Juvenile Rousettus
aegyptiacus Bats Coincide with Periods of Increased Risk of Human Infection. *PLOS*
Pathog. **8**, e1002877. (doi:10.1371/journal.ppat.1002877)

26. McFarlane R, Becker N, Field H. 2011 Investigation of the Climatic and Environmental
Context of Hendra Virus Spillover Events 1994–2010. *PLOS ONE* **6**, e28374.
(doi:10.1371/journal.pone.0028374)

27. Luby SP *et al.* 2006 Foodborne Transmission of Nipah Virus, Bangladesh. *Emerg. Infect.*
Dis. **12**, 1888–1894. (doi:10.3201/eid1212.060732)

28. Pulliam JRC *et al.* 2012 Agricultural intensification, priming for persistence and the
emergence of Nipah virus: a lethal bat-borne zoonosis. *J. R. Soc. Interface* **9**, 89–101.
(doi:10.1098/rsif.2011.0223)

29. Glennon EE, Restif O, Sbarbaro SR, Garnier R, Cunningham AA, Suu-Ire RD, Osei-
Amponsah R, Wood JLN, Peel AJ. 2018 Domesticated animals as hosts of henipaviruses and
filoviruses: A systematic review. *Vet. J.* **233**, 25–34. (doi:10.1016/j.tvjl.2017.12.024)

30. Pourrut X, Kumulungui B, Wittmann T, Moussavou G, Délicat A, Yaba P, Nkoghe D,
Gonzalez J-P, Leroy EM. 2005 The natural history of Ebola virus in Africa. *Microbes Infect.*
7, 1005–1014.

31. Brook CE *et al.* 2019 Disentangling serology to elucidate henipa- and filovirus transmission
in Madagascar fruit bats. *J. Anim. Ecol.* **0**. (doi:10.1111/1365-2656.12985)

32. Pourrut X, Délicat A, Rollin PE, Ksiazek TG, Gonzalez J-P, Leroy EM. 2007 Spatial and
Temporal Patterns of Zaire ebolavirus Antibody Prevalence in the Possible Reservoir Bat
Species. *J. Infect. Dis.* **196**, S176–S183. (doi:10.1086/520541)

33. Páez DJ, Giles J, McCallum H, Field H, Jordan D, Peel AJ, Plowright RK. 2017 Conditions
affecting the timing and magnitude of Hendra virus shedding across pteropodid bat
populations in Australia. *Epidemiol. Infect.* , 1–11. (doi:10.1017/S0950268817002138)

34. Drexler JF *et al.* 2012 Bats host major mammalian paramyxoviruses. *Nat. Commun.* **3**, 796.
(doi:10.1038/ncomms1796)

35. Han BA, Schmidt JP, Alexander LW, Bowden SE, Hayman DT, Drake JM. 2016
Undiscovered bat hosts of filoviruses. *PLOS Negl Trop Dis* **10**, e0004815.

36. Schuh AJ *et al.* 2017 Modelling filovirus maintenance in nature by experimental
transmission of Marburg virus between Egyptian rousette bats. *Nat. Commun.* **8**, 14446.

37. Hayman DTS, Bowen RA, Cryan PM, McCracken GF, O'Shea TJ, Peel AJ, Gilbert A, Webb
CT, Wood JLN. 2013 Ecology of Zoonotic Infectious Diseases in Bats: Current Knowledge
and Future Directions. *Zoonoses Public Health* **60**, 2–21. (doi:10.1111/zph.12000)

38. Sokolow Susanne H. *et al.* 2019 Ecological interventions to prevent and manage zoonotic
pathogen spillover. *Philos. Trans. R. Soc. B Biol. Sci.* **374**, 20180342.
(doi:10.1098/rstb.2018.0342)

39. Peel AJ *et al.* 2013 Use of cross-reactive serological assays for detecting novel pathogens in
wildlife: Assessing an appropriate cutoff for henipavirus assays in African bats. *J. Virol.*
Methods **193**, 295–303. (doi:10.1016/j.jviromet.2013.06.030)

40. Gilbert AT *et al.* 2013 Deciphering serology to understand the ecology of infectious diseases
in wildlife. *Ecohealth* **10**, 298–313.

41. Wacharapluesadee S, Boongird K, Wanghongsa S, Ratanasetyuth N, Supavonwong P,
Saengsen D, Gongal G n., Hemachudha T. 2009 A Longitudinal Study of the Prevalence of
Nipah Virus in Pteropus lylei Bats in Thailand: Evidence for Seasonal Preference in Disease
Transmission. *Vector-Borne Zoonotic Dis.* **10**, 183–190. (doi:10.1089/vbz.2008.0105)

42. Halpin K *et al.* 2011 Pteropid bats are confirmed as the reservoir hosts of henipaviruses: a
comprehensive experimental study of virus transmission. *Am. J. Trop. Med. Hyg.* **85**, 946–
951. (doi:10.4269/ajtmh.2011.10-0567)

43. Wood S. 2006 *Generalized additive models: an introduction with R*. CRC press.

44. Viechtbauer W. 2010 Conducting meta-analyses in R with the metafor package. *J. Stat.*
Softw. **36**, 1–48.

45. Konstantopoulos S. 2011 Fixed effects and variance components estimation in three-level
meta-analysis. *Res. Synth. Methods* **2**, 61–76.

46. Nakagawa S, Santos ES. 2012 Methodological issues and advances in biological meta-
analysis. *Evol. Ecol.* **26**, 1253–1274.

47. Michonneau F, Brown JW, Winter DJ. 2016 rotl : an R package to interact with the Open
Tree of Life data. *Methods Ecol. Evol.* (doi:10.1111/2041-210X.12593)

48. Paradis E, Claude J, Strimmer K. 2004 APE: analyses of phylogenetics and evolution in R
language. *Bioinformatics* **20**, 289–290.

49. Senior AM, Grueber CE, Kamiya T, Lagisz M, O'Dwyer K, Santos ESA, Nakagawa S. 2016
Heterogeneity in ecological and evolutionary meta-analyses: its magnitude and implications.
Ecology **97**, 3293–3299. (doi:10.1002/ecy.1591)

50. Borenstein M, Hedges LV, Higgins JP, Rothstein HR. 2011 *Introduction to meta-analysis*.
John Wiley & Sons. See
http://books.google.com/books?hl=en&lr=&id=JQg9jdrq26wC&oi=fnd&pg=PT14&dq=Bor
enstein+M,+Hedges+L,+Higgins+J,+Rothstein+H+(2005).+Comprehensive+Meta-
Analysis&ots=VH_-OOjzjB&sig=obmoEp98npSeyQvbvsSC_jjdPMc.

51. López-López JA, Marín-Martínez F, Sánchez-Meca J, Noortgate W, Viechtbauer W. 2014
Estimation of the predictive power of the model in mixed-effects meta-regression: A
simulation study. *Br. J. Math. Stat. Psychol.* **67**, 30–48.

52. Changula K *et al.* 2018 Seroprevalence of filovirus infection of Rousettus aegyptiacus bats in
Zambia. *J. Infect. Dis.* **218**, S312–S317.

53. Pawęska JT, van Vuren PJ, Kemp A, Storm N, Grobbelaar AA, Wiley MR, Palacios G,
Markotter W. 2018 Marburg virus infection in Egyptian rousette bats, South Africa, 2013–
2014. *Emerg. Infect. Dis.* **24**, 1134.

54. Plowright RK, Field HE, Smith C, Divljan A, Palmer C, Tabor G, Daszak P, Foley JE. 2008
Reproduction and nutritional stress are risk factors for Hendra virus infection in little red
flying foxes (Pteropus scapulatus). *Proc. R. Soc. B* **275**, 861–869.
(doi:10.1098/rspb.2007.1260)

55. Washburne AD, Crowley DE, Becker DJ, Olival KJ, Taylor M, Munster VJ, Plowright RK.
2018 Taxonomic patterns in the zoonotic potential of mammalian viruses. *PeerJ* **6**, e5979.
(doi:10.7717/peerj.5979)

56. Cressie N, Wikle CK. 2015 *Statistics for Spatio-Temporal Data*. John Wiley & Sons.

57. Yoccoz NG, Nichols JD, Boulinier T. 2001 Monitoring of biological diversity in space and
time. *Trends Ecol. Evol.* **16**, 446–453.

58. Urquhart NS, Kincaid TM. 1999 Designs for detecting trend from repeated surveys of
ecological resources. *J. Agric. Biol. Environ. Stat.* , 404–414.

59. Becker DJ, Crowley DE, Washburne AD, Plowright RK. 2019 Data from: Temporal and
spatial limitations in global surveillance for bat filoviruses and henipaviruses.
(doi:10.5061/dryad.kkwh70s18)

Tables

Table 1. Summary of the temporal and spatial limitations for bat filovirus and henipavirus prevalence and seroprevalence data. Some studies had multiple diagnostic methods, sampling designs, and reporting methods. Diagnostic mismatch refers to geographic regions (United Nations geoscheme) where either PCR or serology have been used (but not together).

		Longitudinal virus studies	Geographic sampling gaps	Diagnostic mismatch	Regions with longitudinal data
Filoviruses	PCR	4/19	Latin America, Oceania, Southern Africa	Southern Africa	Central Africa, Eastern Africa
	Serology	7/25	Latin America, Oceania		Central Africa, Eastern Africa, Middle Africa, Southern Africa
Henipaviruses	PCR	4/13	Eastern Africa, Southern Africa Eastern Asia	Europe, Eastern Africa, Middle Africa, Eastern Asia	Southeastern Asia, Oceania
	Serology	5/27	Middle Africa, Southern Africa, Europe		Southeastern Asia, Oceania, Eastern Africa

Figures and legends

 Figure 1. Top: Conceptual schematic of how different sampling designs and reporting practices
 (colored points and lines) capture the underlying temporal dynamics of infection (black line),
 followed by observed proportions for studies of bat filoviruses and henipaviruses (grey shows
 the proportion of studies not reporting these data). Bottom: Countries sampled for bat filoviruses
 and henipaviruses and where wild bats have been found positive through PCR or serology.

405　Figure 2. Top: Influence of sampling designs and reporting practices on virus detection
406　estimates. Points show proportions of positive bats per subunit; lines and diamonds display back-
407　transformed predicted means and 95% confidence intervals from the MEM. Bottom:
408　Spatiotemporal variation in viral detection estimates for longitudinal studies. Points represent
409　each subunit virus detection estimate and are colored by locations and shaped by month.

411