Chromosome-scale assembly comparison of the Korean Reference Genome KOREF from PromethION and PacBio with Hi-C mapping information

Hui-Su Kim[1], Sungwon Jeon[1,2], Changjae Kim[1], Yeon Kyung Kim[1], Yun Sung Cho[3], Asta Blazyte[1], Andrea Manica[4], Semin Lee[1,2]*, Jong Bhak[1,2,3,5]*

1) KOGIC, Ulsan National Institute of Science and Technology (UNIST), Ulsan 44919, Republic of Korea

2) Department of Biomedical Engineering, School of Life Sciences, UNIST, Ulsan 44919, Republic of Korea

3) Clinomics Inc., Ulsan 44919, Republic of Korea

4) Department of Zoology, Cambridge, University, Cambridge, UK

5) Personal Genomics Institute, Genome Research Foundation, Cheongju 28160, Republic of Korea

JB: jongbhak@genomics.org

Abstract

Background: Long DNA reads produced by single molecule and pore-based sequencers are more suitable for assembly and structural variation discovery than short read DNA fragments. For *de novo* assembly, PacBio and Oxford Nanopore Technologies (ONT) are favorite options. However, PacBio's SMRT sequencing is expensive for a full human genome assembly and costs over 50,000 USD for 30x coverage as of 2018. ONT PromethION sequencing, on the other hand, is cost-effective at one-twentieth the price of PacBio for the same coverage and provides longer average and maximum read lengths, heralding the personalized reference genome era. We aimed to check the cost-effectiveness of PromethION sequencing product.

Findings: We performed whole genome *de novo* assemblies and comparison to construct the new version of KOREF, the Korean reference genome, using sequencing data produced by PromethION and PacBio. With PromethION, an assembly using sequenced reads with 64x coverage (193 Gb, 3 flowcell sequencing) resulted in 3,725 contigs with N50s of 16.7 Mbp and a total genome length of 2.8 Gbp. It was comparable to the KOREF assembly constructed using PacBio at 62x coverage (188 Gbp, 2,695 contigs and N50s of 17.9 Mbp). When we applied Hi-C-derived long-range mapping data, an even higher quality assembly for the 64x coverage was achieved, resulting in 3,179 scaffolds with an N50 of 56.4 Mbp.

Conclusion: In conclusion, the pore-based PromethION approach provides a good quality chromosome-scale human genome assembly at a low cost and is much more cost-effective than PacBio.

Keywords: Korean reference genome; PromethION; Hi-C; nanopore sequencing, single

molecule sequencing

Data Description

Next-generation sequencing (NGS) is a set of powerful sequencing technologies and a recent

trend in genomics is to use cost-effective long DNA reads for assembly and structural

variation discovery using single molecule sequencing methods. Oxford Nanopore

Technologies (ONT) and PacBio platforms have advantages of a short run time and long read

lengths over short fragmented reads by Illumina [1, 2]. Unfortunately, both methods share

high base-calling error rates [3, 4]. However, bioinformatics pipelines for self-error

correction and/or polishing sequences with short reads have become an effective option, and

the overall accuracy of long read based assemblies is approaching what is required to be a

viable option for personal reference genome construction [5]. Despite its excellent

performance, PacBio's SMRT sequencing is expensive for the effective coverage required for

a full human genome assembly, costing over 50,000 USD for 30x coverage as of 2018 [6].

On the other hand, the nanopore based single molecule, long read platform, PromethION

from ONT is highly cost-effective at one-twentieth the price of PacBio's for the same amount

of reads, with an advantage of even longer average and maximum read lengths [7]. Although

the two methods share some similarity, they are fundamentally different in that ONT uses a

minimal amount of reagents with small form factor devices, and can be a promising future

technology for a very broad scope of applications given its advantageous size and cost.

Hi-C is a well-established technology that provides very long-range chromosome

mapping information by digesting fresh cells and then cross-linking intact chromosomal

regions and is often used to extend the size of *de novo* genome scaffolds [8].

In this study, we performed benchmark tests of PromethION and PacBio with low

and high coverages of sequencing data and investigated the advantages of pairing these long

read technologies with Hi-C data, using the already existing high-quality Korean reference,

KOREF, as a benchmark [9].

Whole genome sequencing by ONT PromethION platform

Human KOREF cell lines (http://koref.net) were cultured at 37°C in 5% CO_2 in RPMI-1640

medium with 10% heat-inactivated fetal bovine serum. DNA was extracted from cells using

the DNeasy Blood & Tissue kit (Qiagen). The KOREF cells (5 x 10^6) were centrifuged at 300

g for 5 min; the pelleted cells were suspended in 200 µL of PBS and DNA was extracted

according to the manufacturer's instructions. To preserve large-sized DNA and purify DNA

fragments, we used Genomic DNA Clean & Concentrator kit (Zymo). The DNA quality and

size were assessed by running 1 µL of purified DNA on the Bioanalyzer system (Agilent).

Concentration of DNA was assessed using the dsDNA BR assay on a Qubit fluorometer

(Thermo Fisher).

DNA repair (NEBNext FFPE DNA Repair Mix, NEB M6630) and end-prep

(NEBNext End Repair/dA-tailing, NEB E7546) were performed using 1 µg human genomic

DNA. The mixture of 1 µL DNA CS, 3.5 µL FFPE Repair Buffer, 2 µL FFPE DNA Repair

Mix, 3.5 µL Ultra II End-prep reaction buffer, and 3 µL Ultra II End-prep enzyme mix was

added to 47 µL DNA sample. The final mixture was incubated at 20°C for 5 min and then at

65°C for 5 min, cleaned up using 60 µL AMPure XP beads, incubated on Hula mixer for 5

min at room temperature, and washed twice with 200 µL fresh 70% ethanol. The pellet was

allowed to dry for 30 s, and then DNA was eluted in 61 µL of nuclease-free water. An aliquot

of 1 µL was quantified by Qubit to ensure ≥ 1 µg DNA was retained.

Adaptor ligation was performed by adding 5 µL of Adaptor Mix (AMX, SQK-LSK109 Ligation Sequencing Kit 1D, Oxford Nanopore Technologies (ONT)), 25 µL Ligation Buffer (LNB, SQK-LSK109), and 10 µL NEBNext Quick T4 DNA Ligase (NEB, E6056) to 60 µL bead cleaned-up DNA, followed by gentle mixing and incubation for 10 min at room temperature.

The adaptor-ligated DNA was cleaned up by adding 40 µL of AMPure XP beads, incubating for 5 min at room temperature and re-suspending the pellet twice in 250 µL L Fragment Buffer (LFB, SQK-LSK109). The purified ligated DNA was re-suspended in 25 µL of Elution Buffer (ELB, SQK-LSK109), incubated for 10 min at room temperature, followed by pelleting the beads, and transferring the supernatant (pre-sequencing mix or PSM) to a new Eppendorf Lobind tube. A 1-µL aliquot was quantified by Qubit to ensure ≥ 500 ng DNA was retained.

To load the library, 75 µL of Sequencing Buffer (SQB, SQK-LSK109) was mixed with 51 µL of Loading Beads (LB, SQK-LSK109) and this mixture was added to 24 µL DNA library. This library was mixed by pipetting slowly and 150 µL of sample was loaded through the inlet port.

Whole genome sequencing by PacBio Sequel platform

Genomic DNA was extracted from human KOREF blood samples using QIAGEN Blood & Cell Culture DNA Kit (cat no 13323). A total of 5 µg of each sample was used as input for library preparation. The SMRTbell library was constructed using SMRTbell® Express Template Preparation Kit (101-357-000). Using the BluePippin Size selection system we removed the small fragments for large-insert library. After sequencing primer v4 was

annealed to the SMRTbell template, DNA polymerase was bound to the complex (Sequel Binding kit 2.0). We purified the complex using AMPure Purification to remove excess primer and polymerase prior to sequencing. The SMRTbell library was sequenced using SMRT cells (Pacific Biosciences) using Sequel Sequencing Kit v2.1 and 10 h movies were captured for each SMRT Cell 1M v2 using the Sequel (Pacific Biosciences) sequencing platform.

Short read sequencing by Illumina HiSeq

Short paired-end raw reads using Illumina HiSeq 2000 platform were acquired from a previous study, accession no. SRR2204706 (ftp://ftp.sra.ebi.ac.uk/vol1/srr/SRR220/006/SRR2204706).

Hi-C chromosome conformation captured reads sequencing

Long distance Hi-C chromosome conformation capture data were generated using the Arima-HiC kit (A160105 v01), and double restriction enzymes were used for chromatin digestion. To prepare KOREF cell line samples for Hi-C analysis, cells were harvested and cross-linked as instructed by the manufacturer. One million cross-linked cells were used as input in the Hi-C protocol. Briefly, chromatin from cross-linked cells or nuclei was solubilized, and then digested using restriction enzymes A1 and A2. The digested ends were then labeled using a biotinylated nucleotide, and ends were ligated to create ligation products. Ligation products were purified, fragmented, and selected by size using AMpure XP Beads. Biotinylated fragments were then enriched using Enrichment beads, and Illumina-compatible sequencing libraries were constructed on End Repair, dA-tailing, and Adaptor Ligation using a modified

1 workflow of the Hyper Prep kit (KAPA Biosystems, Inc.). The bead-bound library was then

2 amplified, and amplicons were purified using AMpure XP beads and subjected to deep

3 sequencing.

Short and long sequence reads processing

A total of 144 Gbp of short paired-end DNA raw reads were obtained from SRA2204706.

Adapter sequences were trimmed from sequenced raw reads using Trimmomatic v0.36 [10]

(ILLUMINACLIP:2:30:10   LEADING:5   TRAILING:5   SLIDINGWINDOW:4:20

HEADCROP:15 MINLEN:60), and screening for vectors and microbial contaminants were

performed using customized database from Refseq. After preprocessing, a total of 137 Gbp

cleaned reads were obtained.

A total of 80.7 Gbp and 193 Gbp raw reads (27x and 64x coverage) were obtained as

a result of PromethION nanopore sequencing using one and three flowcells. Removing

adapter sequences from the raw reads was performed using Porechop v0.2.4 [11]. We also

acquired 92.2 Gbp and 187.9 Gbp raw reads from PacBio Sequel sequencing resulting in 30x

and 62x coverage (Table 1).

Long-read sequence based *de novo* genome assemblies

19 *De novo* assemblies for the 27x and 64x PromethION raw reads were performed using

20 wtdbg2 v2.3 [12]. To compare the accuracy, two sets of raw reads with 30x and 62x coverage

21 of PacBio Sequel were also used employing the same assembler. Parameters for the

22 assembler were set optimally for each sequencing platform with multiple trials. For self-error

23 correction with long reads, we generated consensus sequences using Racon v1.3.2 [13]. To

improve the accuracy of assemblies, polishing consensus sequences with 48.2x coverage short reads was performed using Pilon v1.23 [14]. To assess the completeness of the long-read genome assemblies, BUSCO v3.0.2 [15] with the default AUGUSTUS model for human was used to locate presence and absence of 4,104 single copy orthologous genes from mammalian OrthoDB v9.

For constructing chromosome-scale assemblies for the PromethION long-reads data, map assembly with Hi-C reads was performed using SALSA2 v2.2 [16]. Duplicated Hi-C reads were removed using clumpify.sh program from BBTools suite v38.32 [17]. Mapping Hi-C reads to the assembled genome was conducted using the pipeline provided by Arima-Genomics (https://github.com/ArimaGenomics/mapping_pipeline).

Long read assemblies from 27x and 64x PromethION sequencing yielded total assembly sizes of 2,757 Mbp and 2,827 Mbp, with scaffold N50s of 7.6 Mbp and 16.7 Mbp, respectively (Table 2). Assemblies from PacBio sequencing at 30x and 60x coverage yielded the total assembly sizes of 2,800 Mbp and 2,815 Mbp, with scaffold N50s of 11.1 Mbp and 17.9 Mbp, respectively. Adding Hi-C reads to assemblies led to 3.4- to 4.3-fold increase in the scaffold N50 lengths of PromethION (32.7 Mbp for 27x coverage and 56.4 Mbp for 64x coverage). For the PacBio assemblies, 2.2- to 3.4-fold increase was achieved for the scaffold N50 lengths (38.1 Mbp for 30x coverage and 40.9 Mbp for 62x coverage). The longest scaffold from both PromethION and PacBio assemblies with Hi-C was two times the length of the assemblies without Hi-C.

Comparison between PromethION and PacBio assemblies

The comparison between PromethION and PacBio assemblies without Hi-C mapping

information using sequenced reads at 64x coverage indicated comparable quality. In terms

of N50, the PromethION assembly at 64x coverage yielded 1.45-fold and 0.93-fold longer

N50s compared with the PacBio assemblies at 30x and 62x coverage, respectively (Figure

1a). When we compared the longest contigs, the PromethION assembly at 64x coverage

yielded 1.7-fold and 1.1-fold length increase compared with the PacBio assemblies at 30x

and 62x coverage, respectively (Figure 1b). Comparing the number of scaffolds, PacBio

assembly at 30x coverage showed the fewest (2,443) compared with that of PromethION

assembly at 64x coverage (3,725).

When Hi-C mapping information was added to the assembly construction, the

PromethION assembly at 64x coverage showed the best statistics as N50s of 56.4 Mbp and

the longest scaffold length of 175.2 Mbp. The PromethION assembly at 27x coverage with

Hi-C mapping information yielded 32.7 Mbp for N50s, which was comparable to both 30x

and 62x coverage PacBio assemblies with Hi-C; 0.85-fold and 0.79-fold for N50s,

respectively (Table 2).

When we compared assessment results from BUSCO, all the assemblies that had

been polished with short reads showed good quality; around 92% completed orthologous

genes with less than 1.1% completed and duplicated orthologous genes. Comparing the

accuracy of the assemblies to KOREF_S, which is the current standard, both showed around

99.8% accuracy (Table 3). The accuracy comparison was performed using assess_assembly

program from Pomoxis [18].

Conclusions

We generated high-quality assemblies of a Korean human genome using ONT's PromethION

long-reads accompanied with Hi-C mapping information and compared them against PacBio

sequencing and assemblies of the same sample. Comparing the results from the PromethION

64x sequencing to the PacBio 62x sequencing, we found that the former provided high

contiguity and completeness at one-twentieth the cost of PacBio. Results from just 27x

PromethION sequencing combined with Hi-C mapping information were also comparable to

the 30x and even 62x coverage PacBio sequencing data. Therefore, to generate a

chromosome-scale assembly with a long-read technology, at present, the ONT's PromethION

sequencing is a good alternative to PacBio's, owing to its quality and cost-effectiveness.

Simple pore-based long read sequencing has potential to revolutionize sequencing and

subsequent bioinformatics analysis for personal genome projects and cancer genome analyses

where *de novo* assemblies are necessary for structural and copy number variations that cannot

be detected easily by conventional short read only methods.

Availability of supporting data

Raw long-read sequencing data from PromethION and PacBio is available at NCBI genbank

under the project accession number PRJNA549351. All genome assemblies of KOREF are

available at KOREF website (http://koref.net).

Abbreviations

BUSCO: Benchmakring Universal Single-Copy Orthologs; PacBio: Pacific Biosciences;

SMRT: single-molecule real-time

Competing interests

Y.S.C. is an employee, and J.B. is the CEO of Clinomics Inc. J.B. and Y.S.C. have an equity

interest in the company. All other coauthors have no conflicts of interest to declare.

Funding

This work was supported by U-K BRAND Research Fund (1.190007.01) of UNIST;

Research Project Funded by Ulsan City Research Fund (1.190033.01) of UNIST.

1 **Figure Legends**

3 **Figure 1.** Comparison of N50s and the longest contig/scaffold lengths for PromethION and

4 PacBio assemblies of the Korean human reference, KOREF

References

1. Mccarthy A. Third generation DNA sequencing: Pacific biosciences' single molecule real time technology. Chem Biol [Internet]. Elsevier Ltd; 2010;17:675–6.

2. Laver T, Harrison J, O'Neill PA, Moore K, Farbos A, Paszkiewicz K, et al. Assessing the performance of the Oxford Nanopore Technologies MinION. Biomol Detect Quantif [Internet]. Elsevier GmbH; 2015;3:1–8.

3. Chin C-S, Alexander DH, Marks P, Klammer AA, Drake J, Heiner C, et al. Nonhybrid, finished microbial genome assemblies from long-read SMRT sequencing data. Nat Methods [Internet]. 2013;10:563–9.

4. Laver T, Harrison J, O'Neill PA, Moore K, Farbos A, Paszkiewicz K, et al. Assessing the performance of the Oxford Nanopore Technologies MinION. Biomol Detect Quantif [Internet]. Elsevier GmbH; 2015;3:1–8.

5. Fu S, Wang A, Au KF. A comparative evaluation of hybrid error correction methods for error-prone long reads. Genome Biol [Internet]. Genome Biology; 2019;20:1–17.

6. AllSeq. Available from: https://allseq.com/knowledge-bank/sequencing-platforms/pacific-biosciences/

7. Nanopore tech. Available from: https://nanoporetech.com/products/comparison/

8. Korbel JO, Lee C. Genome assembly and haplotyping with Hi-C. Nat Biotechnol [Internet]. Nature Publishing Group; 2013;31:1099–101. Available from: http://dx.doi.org/10.1038/nbt.2764

9. Cho YS, Kim H, Kim HM, Jho S, Jun JH, Lee YJ, et al. Corrigendum: An ethnically relevant consensus Korean reference genome is a step towards personal reference genomes. Nat Commun [Internet]. 2017;8:16168.

10. Bolger AM, Lohse M, Usadel B. Trimmomatic: A flexible trimmer for Illumina

sequence data. Bioinformatics [Internet]. 2014;30:2114–20

11. Porechop, adapter trimmer for Oxford Nanopore reads. https://github.com/rrwick/Porechop

12. Ruan J, Li H. Fast and accurate long-read assembly with wtdbg2. bioRxiv [Internet]. 2019;530972.

13. Racon, ultrafast consensus module for raw de novo genome assembly of long uncorrected reads. https://github.com/isovic/racon

14. Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, et al. Pilon: An integrated tool for comprehensive microbial variant detection and genome assembly improvement. PLoS One [Internet]. 2014;9.

15. Simão FA, Waterhouse RM, Ioannidis P, Kriventseva E V., Zdobnov EM. BUSCO: Assessing genome assembly and annotation completeness with single-copy orthologs. Bioinformatics [Internet]. 2015;31:3210–2.

16. Ghurye J, Pop M, Koren S, Bickhart D, Chin CS. Scaffolding of long read assemblies using long range contact information. BMC Genomics [Internet]. BMC Genomics; 2017;18:1–11.

17. BBMap, short read aligner and other bioinformatics tools. https://sourceforge.net/projects/bbmap/

18. Pomoxis, bioinformatics tools for nanopore research. https://nanoporetech.github.io/pomoxis/

1   **Table 1. Statistics of raw sequenced reads**

	ONT PromethION		PacBio Sequel		Short read
	27x	**64x**	**30x**	**62x**	**Illumina HiSeq 2000**
Number of reads	15,004,723	47,591,997	11,195,434	20,683,965	1,433,779,680
Total length of reads	80,770,821,288	193,027,803,978	92,229,416,062	187,914,740,184	144,811,747,680
N50	12,736	9,190	13,426	14,568	101
Max contig length	774,322	1,160,324	65,865	169,910	101

Table 2. Statistics of KOREF genome assemblies using ONT PromethION and PacBio Sequel sequencing

	ONT PromethION				PacBio Sequel			
	27x assembly	**64x assembly**	**27x assembly with Hi-C**	**64x assembly with Hi-C**	**30x assembly**	**62x assembly**	**30x assembly with Hi-C**	**62x assembly with Hi-C**
Contigs / Scaffolds No.	3,262	3,725	2,313	3,179	2,443	2,695	1,476	2,338
Total length	2,757,297,803	2,827,624,042	2,757,776,303	2,827,900,542	2,800,962,512	2,815,311,932	2,801,450,512	2,818,181,112
Scaffold N50	7,655,153	16,706,773	32,758,624	56,457,651	11,137,362	17,931,968	38,113,117	40,953,073
Max contig / scaffold length	60,569,695	88,903,341	120,666,262	175,227,974	50,101,007	77,816,513	126,818,544	131,630,574
Gap	0.00%	0.00%	0.02%	0.01%	0.00%	0.00%	0.02%	0.02%
GC content	40.82%	40.81%	40.82%	40.81%	40.90%	40.92%	40.90%	40.92%

Table 3. Statistics of KOREF genome assembly assessment using BUSCO and accuracy comparison

BUSCO assessment	ONT PromethION				PacBio Sequel			
	27x assembly	64x assembly	27x assembly with Hi-C	64x assembly with Hi-C	30x assembly	62x assembly	30x assembly with Hi-C	62x assembly with Hi-C
Complete	92.50%	92.70%	92.60%	94.00%	93.80%	93.10%	93.80%	93.50%
Complete and single-copy	91.80%	91.60%	91.90%	93.20%	93.00%	92.00%	93.00%	92.70%
Complete and duplicated	0.70%	1.10%	0.70%	0.80%	0.80%	1.10%	0.80%	0.80%
Fragmented	3.10%	3.70%	3.20%	3.10%	3.00%	3.20%	3.00%	3.10%
Missing	4.40%	3.60%	4.20%	2.90%	3.20%	3.70%	3.20%	3.40%
Accuracy comparison*	99.81%	99.79%	99.78%	99.73%	99.86%	99.79%	99.84%	99.80%

* Compared with KOREF_S, Korean reference genome

1   **Figure 1**