K$^+$ uptake and root-to-shoot allocation in *Arabidopsis* require coordination of nitrate

transporter1/peptide transporter family member NPF6.3/NRT1.1

Xian Zhi Fang[1,2], Xing Xing Liu[1,2], Ya Xing Zhu[1], Jia Yuan Ye[1], Chong Wei Jin[1, *]

[1]Ministry of Education Key Laboratory of Environmental Remediation and Ecosystem

Health, College of Natural Resources and Environmental Science, Zhejiang University,

Hangzhou, 310058, China

[2]These authors contributed equally to this work.

*To whom correspondence should be addressed.

Email: jincw@zju.edu.cn

Tel: +(86) 571 88982478

Abstract

K^+ and NO_3^- are the major forms of potassium and nitrogen that are absorbed by the roots

of most terrestrial plants. In this study, we observed that the close relationship between

NO_3^- and K^+ homeostasis was mediated by nitrate transporter1 (NRT1.1) in *Arabidopsis*.

The *nrt1.1* mutants lacking NRT1.1 function showed disturbed K^+ uptake and root-to-

shoot allocation, especially under K^+-limited conditions, and had a yellow-shoot sensitive

phenotype on K^+-limited medium. The K^+ uptake and root-to-shoot allocation of these

mutants were partially rescued by expressing NRT1.1 in the root epidermis-cortex and

central vasculature by using *Sultr1;2* and *PHO1* promoters, respectively. Furthermore,

two-way analysis of variance based on the K^+ content in *nrt1.1-1/akt1*, *nrt1.1-1/hak5-3*,

nrt1.1-1/kup7, and *nrt1.1-1/skor-2* double mutants and their corresponding single mutants

and wild-type plants revealed physiological interactions between NRT1.1 and K^+

channels located in the root epidermis-cortex and central vasculature. Taken together,

these data suggest that the expression of NRT1.1 in the root epidermis-cortex coordinates

with K^+ uptake channels to improve K^+ uptake, whereas its expression in the root central

vasculature coordinates with the channels loading K^+ into the xylem to facilitate K^+

allocation from the roots to the shoot.

Introduction

Potassium (K^+) is essential for plant growth and development, and it determines the yield and quality of crops in agriculture production (Wang & Wu, 2013). However, the concentration of soluble K^+ in most soils is relatively low, which often limits plant growth (Maathuis, 2009). Crop production is increased by applying a large amount of potassic fertilizers to agricultural fields, but about one-half of the applied fertilizers is unavailable to plants and left behind as residues in soils, consequently leading to environmental contamination (Meena *et al*., 2016). Thus, improving the K^+ utilization efficiency of plants by thoroughly understanding the molecular mechanisms of K^+ transport and regulation is urgently required. Therefore, in the past few decades, researchers have been focusing on identifying K^+ channels and transporters as well as their regulation mechanisms in plants.

In *Arabidopsis thaliana*, 71 K^+ channels and transporters have been identified and categorized into three channel families (shaker, tandem-pore K^+, and Kir-like) and three transporter families (KUP/HAK/KT, HKT, and CPA families) (Wang & Wu, 2010). Among them, the shaker inward K^+ channel AKT1 and the KUP/HAK/KT K^+ transporter HAK5 have been characterized as the two major components that mediate K^+ uptake from the external environment into root cells, but they operate at different K^+ levels (Pyo *et al*., 2010; Wang & Wu, 2013). AKT1 functions in plant K^+ uptake over a wide range of K^+ concentrations. Comparatively, HAK5 shows a high-affinity K^+ transport activity (Gierth *et al*., 2005). After its uptake into root epidermal cells, K^+ is then distributed to different organs or tissues of a plant. An *Arabidopsis* shaker-like outward-rectifying K^+ channel, SKOR, was first identified to facilitate K^+ secretion from the stelar cells to the xylem sap in the roots, which is a critical step for long-distance K^+ transport from the roots to the shoots (Gaymard *et al*., 1998). Recently, *At*KUP7, one of the KT/HAK/KUP family members, was functionally characterized as a novel K^+ transporter participating in both K^+ uptake by roots from the growth medium and K^+ release into the xylem for its allocation from roots to shoots, especially under K^+-limited conditions (Han *et al*., 2016). However, the uptake affinity for K^+ of KUP7 is considerably lower than that of *At*HAK5 (Wang & Wu, 2017).

In addition to the several aforementioned K^+ channels and transporters, many other nutrients such as Na^+, Ca^{2+}, and N also remarkably affect K^+ homeostasis in plants,

particularly the association of N with K^+, which has been explored for a long time because

N is the mineral element that is required in the greatest quantity by most plants and is the

most widely used fertilizer nutrient in crop production (Fageria & Baligar, 2005; Wang &

Wu, 2013; Shin, 2017). Since the 1960s, physiological studies have revealed a close

relationship between NO_3^- and K^+ with regard to uptake and translocation (Zioni *et al.*,

1971; Blevins *et al.*, 1978; Barneix & Breteler, 1985; Drechsler *et al.*, 2015). However,

how these two nutrients coordinate with each other in the transport pathways has not yet

been clearly explored at the molecular level. Interestingly, lines of evidence have shown

that NRT1.5 from the nitrate transporter1 (NRT1)/peptide transporter (PTR) transporter

family (NPF) functions as a component in controlling K^+ allocation in plants (Lin *et al.*,

2008; Drechsler *et al.*, 2015; Li *et al.*, 2017; Du *et al.*, 2019), which was initially

identified as a pH-dependent bidirectional NO_3^- transporter and is mainly expressed in the

root pericycle cells mediating NO_3^- release into the xylem vessels (Lin *et al.*, 2008). In

this study, we showed that the loss of another nitrate transporter1 member NRT1.1 in

nrt1.1 mutants also led to the development of a more serious K-deficiency phenotype

under low-K stress. Further physiological and genetic evidences revealed that both the

uptake and root-to-shoot allocation of K^+ in plants require NRT1.1 in plants. However,

NRT1.1 acts as a coordinator rather than a K^+ channel/transporter in the above two K^+

nutrition issues, which could depend on its transport activity for NO_3^-. Our findings

suggest that the modification of NRT1.1 activity in crops by using biological engineering

approach might be a promising way for simultaneously improving the utilization

efficiencies of K and N fertilizers in agriculture production.

Material and Methods

Plant material

The mutants *chl1-5* (Huang et al., 1996), *nrt1.1-1* (salk_097431), *nrt1.2* (cs859605),

nrt2.1 (cs859604), *nrt2.2* (salk_043543), *nrt2.5* (GK-213H10.06), *akt1* (salk_071803),

hak5 (salk_130604), *kup7* (cs805085), and *skor* (cs2103489) and the transgenic plants

pNRT1.1::NRT1.1-GFP and *pNRT1.1::NRT1.1-GUS* (cs6513) are in a Columbia (Col-0)

background, whereas the *chl1.6* (cs6154) and *nrt2.4* (cs27332) lines are in a Landsberg

erecta (*Ler*) background. The seeds of the lines *chl1-5* and *pNRT1.1::NRT1.1-GFP* were

kindly gifted by Dr. Philippe Nacry (Biochimie et Physiologie Moléculaire des Plantes,

Montpellier, France). The double mutants *nrt1.1-1/akt1*, *nrt1.1-1/hak5-3*, *nrt1.1-1/kup7*,

and *nrt1.1-1/skor-2* were generated by crossing *nrt1.1-1* with *akt1*, *hak5-3*, *kup7*, and
skor-2, respectively. The mutants were verified using the primers listed in Table S1.

Plant culture

The seeds were surface-sterilized and sown on basal agar medium containing 1% sucrose
(w/v) and 0.9% (w/v) agar. The complete nutrients in the basal agar medium had the
following composition: 2.25 mM KNO_3, 500 μM $CaCl_2$, 500 μM $MgSO_4$, 750 μM
NaH_2PO_4, 375 μM K_2SO_4, 375 μM $(NH_4)_2SO_4$, 25 μM Fe-EDTA, 10 μM H_3BO_3, 0.5 μM
$MnSO_4$, 0.5 μM $ZnSO_4$, 0.1 μM $CuSO_4$, and 0.1 μM $(NH_4)_6Mo_7O_{24}$, pH 6.5. After 4 d of
germination, the seedlings were used in studies, and the final concentrations of K^+ and
NO_3^- were adjusted using KCl and $Ca(NO_3)_2$, as described in the figure legends.

Phenotypic analyses

After 8 d of growth in agar medium containing 2 mM K^+ or 0.05 mM K^+, the phenotype
and fresh weight of plants were recorded. The chlorophyll of fresh leaves was extracted
with 80% (v/v) acetone and was colorimetrically measured according to Choi *et al*. (2014).
In addition, the chlorophyll fluorescence was analyzed using a pulse amplitude-
modulating imaging fluorometer (IMAGING-PAM; Germany). Briefly, after a 20 min
dark-adaption, the leaves were arranged on the fluorometer, and the maximum quantum
efficiency (Fv/Fm) and electron transport rate through PSII (ETR) were monitored.

Measurement of K concentrations

The harvested plants were separated into shoots and roots. Samples were dried at 75°C for
48 h, and then wet-digested as described by Luo *et al*. (2012). The digests were diluted
with ultrapure water, and the K concentrations were analyzed using MP-AES (Agilent
Technologies; USA).

Green fluorescent protein and β-glucuronidase expression analysis

The NRT1.1-green fluorescent protein (GFP) expression in the roots of
pNRT1.1::NRT1.1-GFP plants was observed using an epifluorescence microscope
(excitation, 488 nm; emission, 525−550 nm; Ni-U; Nikon). Histochemical assay of β-
glucuronidase (GUS) expression in the roots of *pNRT1.1::NRT1.1-GUS* plants was
performed as described by Fang *et al*. (2016), and the distribution and intensity of the blue
product were observed under a microscope.

Complementation of *nrt1.1-1* with *pSultr1;2::NRT1.1* and *pPHO1::NRT1.1*

The 2 kb genomic fragments upstream of the *PHO1* and *Sultr1;2* initiation codons were, respectively, PCR-amplified on genomic DNA from Arabidopsis by using the primers listed in Table S1. Subsequently, the above two amplified products were cloned upstream of the NRT1.1 coding region by using the pCAMBIA1300 vector. The constructs were transformed in *nrt1.1-1* plants by using the Agrobacterium-mediated floral dip method. Homozygous T4 transgenic plants were used.

Measurement of NO_3^- and K^+ and Rb^+ uptake

Net fluxes of NO_3^- and K^+ were measured in the roots by using a non-invasive microelectrode ion flux measurement system (ipa-2; AE, USA). After 3 d of preculture in the nutrient medium containing 2 mM or 0.05 mM K^+, the flux of NO_3^- and K^+ in each roots section was monitored using NO_3^--selective and K^+-selective microelectrodes, as described by Hawkins *et al.* (2008) and Li *et al.* (2012), respectively. For Rb^+ uptake, the 4-d-old seedlings were grown on complete nutrient medium containing 2 mM or 0.05 mM K^+ replaced with Rb^+ for 12 h, and then these plants were harvested and used for Rb concentration analysis.

Transcription analysis

Total RNA in roots was extracted using RNAisoPlus (TaKaRa, Otsu, Japan) and was treated with DNase I to remove genomic DNA contamination. The first-strand cDNA was synthesized from genomic DNA-removed RNA by using PrimeScript™ RT reagent kit (TaKaRa). Next, the transcript levels of the corresponding genes were measured using quantitative real-time PCR by using TB Green™ Premix Ex Taq™ II (TaKaRa). The primers used are listed in Table S1.

RNA sequencing and functional enrichment analysis

Total RNA was extracted from the roots of Col-0 and *nrt1.1-1* plants by using the MagZol Reagent (Magen), according to manufacturer's protocol. Sequencing libraries were prepared using NEBNext Ultra RNA Library Prep Kit for Illumina (NEB, USA), and the quality was assessed using the Agilent 2100 Bioanalyzer. Sequencing was performed on a Hiseq Xten (Illumina) at RIBOBIO (Guangzhou, China). Raw sequencing reads were quality controlled and trimmed using Trimmomatic tools and FastQC, and then the clean

reads were aligned to the TAIR10 reference genome by using HISAT2. Significantly
differentially expressed genes were assessed using an adjusted P-value threshold of <0.05
and |log$_2$(fold change)| of >1 by using DEGseq. All the genes with significant differential
expression were grouped based on k-means clustering and selected for subsequent GO
analyses by using R package clusterProfiler.

Grafting of Arabidopsis plants

Grafts between *nrt1.1-1* mutants and Col-0 plants were performed following the protocol
described by Andersen *et al*. (2014). Briefly, seedlings to be grafted were grown on
complete nutrient agar medium for 5 days. After cotyledons were removed, the seedlings
were separated into scion and rootstock by using a sterilized blade. The scion was then
grafted to the corresponding rootstock on new agar plates by using forceps under a
stereomicroscope. The grafted seedlings were left undisturbed for another 5−7 days to
form graft union. The adventitious roots emerging at or above the graft union were
eliminated, and then the successful unified seedlings were used in the study as described
in the legend of figure 4.

Yeast complementation assay

The protein coding sequence of NRT1.1 was constructed into the pDR196 vector by using
primers listed in Table S1. Next, the pDR196-NRT1.1 vector was transformed into the
yeast strain R5421 (*trk1Δ, trk2Δ*) by using the lithium acetate method (Ito *et al*., 1983).
Yeast strain R757 was used as a positive control. The picked single yeast cells were
precultured overnight at 30°C in 2 mL liquid YPDA medium containing 100 mM KCl.
The cells were collected the following day and washed with ultrapure water three times.
Subsequently, the yeast cells were resuspended in ultrapure water to an OD$_{600}$ of 1, and
10-fold serial dilutions from OD$_{600}$ of 1 to 10^{-3} were incubated on AP agar plates
supplemented with different K$^+$ concentrations for the experiments.

Results

NO$_3^-$ facilitates K$^+$ uptake by roots

We first determined how the NO$_3^-$ supply affects the K levels in Arabidopsis wild-type
(Col-0) plants. The presence of NO$_3^-$ in the growth medium increased the K levels in both
roots and shoots compared with those in the NO$_3^-$-free treatment (Fig. 1a,b). However, a

further increase of NO_3^- supply over 6 mM decreased the K levels in plants. These results

indicate that an appropriate NO_3^- supply might favor the uptake of K^+ by roots. Therefore,

we calculated the amount of K^+ absorbed per weight of roots and found that these values

also increased along with the supply of NO_3^- when the level was less than 6 mM (Fig. S1).

To further evaluate how NO_3^- influences K^+ uptake by Col-0 roots, we used the non-

invasive micro-test technology (NMT system) to measure the transmembrane K^+ influx

and efflux by root cells in a non-invasive manner (Li *et al.*, 2012). Although the net K^+

influx by root cells was observed in the NO_3^--free medium (Fig. 1c), its average value was

only around -30 pmol cm^{-2} s^{-1} (Fig.1d, a negative value indicates influx). Interestingly,

addition of 6 mM NO_3^- to the rooting media resulted in a sudden increase in the net K^+

influx rate, which reached around -900 pmol cm^{-2} s^{-1} (Fig. 1c). Subsequently, the rate of

transmembrane K^+ influx gradually decreased to a relatively stable level, but the average

value was still about 7-fold higher than that of the NO_3^--free treatment (Fig. 1c,d). Taken

together, these results suggest that the uptake of K^+ by roots might require a coordination

of NO_3^- uptake.

Resistance to K deficiency requires NRT1.1

As the uptake of K^+ requires a coordination of NO_3^- uptake in roots, we intended to

determine whether the sensitivity of a plant to low K stress is also associated with the root

NO_3^- uptake. In *Arabidopsis*, six nitrate transporters, including NRT1.1, NRT1.2, NRT2.1,

NRT2.2, NRT2.4, and NRT2.5, have been characterized to be involved in the root NO_3^-

uptake (Wang *et al.*, 2012; Léran *et al.*, 2014; Lezhneva *et al.*, 2014). Here we found that,

in the growth medium containing 6 mM NO_3^-, the two NRT1.1 knockout mutants *nrt1.1-1*

and *chl1-5* exhibited severer leaf senescence than in the wild-type plants (Col-0) after 8 d

of low K treatment (0.05 mM K), but these three plant lines did not show distinguishable

phenotypes under sufficient K treatment (2 mM K; Fig. 2a) Considering that the decrease

of chlorophyll content would affect the photosynthesis of plants, the maximum quantum

efficiency of photosystem II (Fv/Fm) and photosynthetic electron transport rate (ETR)

were also evaluated as indicators of the photosynthetic reactions in leaves. In the leaves of

nrt1.1-1 and *chl1-5* plants, the Fv/Fm was almost completely abolished (Fig. 2c), and the

ETR also declined apparently compared with that in Col-0 plants under low-K conditions

(Fig. 2d). Therefore, the *nrt1.1* knockout mutants had both shorter roots and less biomass

than the Col-0 plants after low-K treatment (Figs 2e,f, S2). In addition, we examined a

third NRT1.1-null mutant, *chl1.6*, the fresh weight and root elongation of which were also

significantly reduced compared with those of its corresponding wild-type *Ler* under low-
K conditions (Fig. S3a,b). Since NRT1.1 can act as a high-affinity as well as low-affinity
system (Wang *et al*., 2012; Léran *et al*., 2014), the growth of these NRT1.1-null mutants
was also evaluated in low-K growth medium containing 0.2 mM NO_3^-. However, the
phenotypes of these mutants were indistinguishable from those of wild-type plants (Fig.
S4a,b). These results suggest that NRT1.1 plays an important role in plants to confer
resistance to K deficiency, but it requires a sufficient supply of NO_3^- in the growth
medium.

We next determined the roles of the other five NRTs in plant growth responses to low-K
stress. Since NRT2.1, NRT2.2, NRT2.4, and NRT2.5 act as high-affinity NO_3^-
transporters, the growth of these NRTs-null mutants was evaluated in a growth medium
containing 0.2 mM NO_3^-. However, the fresh weight and root elongation of *nrt2.1*, *nrt2.2*,
nrt2.4, and *nrt2.5* mutants were similar to those of the corresponding wild-type plants in
both low-K and sufficient-K treatments (Fig. S5). Although NRT1.2 is a low-affinity
NO_3^- transporter (Huang *et al*., 1999), the *nrt1.2* mutant showed similar fresh weight and
root elongation as those of wild-type plants in both low-K and sufficient-K growth
medium containing 6 mM NO_3^- (Fig. S5e,j). These results indicate that NRT1.2, NRT2.1,
NRT2.2, NRT2.4, and NRT2.5 do not act like NRT1.1 when conferring resistance to K
deficiency.

NRT1.1 is required for homeostasis of K in plants

We next emphasized on how NRT1.1 affects K nutrition in plants. In both sufficient-K
and low-K treatments, the K levels were significantly lower in *nrt1.1-1*, *chl1-5*, and *chl1-*
6 mutants than in their crossponding wild-type plants (Figs 3a,b, S3c,d), indicating that
the lack of NRT1.1 function disturbed the K homeostasis in plants. To provide further
support for the role of NRT1.1 in K homeostasis, we measured the K levels in the
transgenic plants *pNRT1.1::NRT1.1-GFP*, which are in the *chl1-5* mutant background
(Krouk *et al*., 2010). In both sufficient-K and low-K treatments, the *pNRT1.1::NRT1.1-*
GFP transgenic plants contained higher K levels than those in the *chl1-5* mutants and
similar K levels as those in Col-0 plants (Fig. 3c,d). Therefore, phenotype analysis
showed that complementation of NRT1.1 conferred the transgenic plants tolerance to K^+
deficiency, as evidenced by the similar chlorophyll content, root elongation, and fresh
weight as those in Col-0 plants (Figs 3e, S6).

As described above, the uptake of K^+ by roots requires a coordination of NO_3^- uptake.
However, when the plants were grown in the medium containing low NO_3^- (0.2 mM), no
significant difference in both growth and K levels was noted between *NRT1.1-null*
mutants and wild-type plants in both sufficient-K and low-K treatments (Fig. S4). The
result suggested that the role of NRT1.1 in maintaining K homeostasis in plants depends
on sufficient NO_3^- supply. We also investigated the role of the other NRTs in maintaining
K homeostasis. Nevertheless, the K levels in the roots and shoots of *nrt1.2, nrt2.1, nrt2.2,*
nrt2.4, and *nrt2.5* mutants were not obviously different from those of their corresponding
wild-type plants in sufficient-K or low-K treatments (Fig. S8). Thus, NRT1.1 might
improve the K nutrition in a relatively specific manner.

Root is the action point for NRT1.1 to improve K nutrition in plants, and low-K
stress stimulates the activity of NRT1.1 in roots

NRT1.1 is expressed in both roots and shoots (Guo *et al*., 2001). Therefore, homografted
(Col-0 scion/Col-0 stock and *nrt1.1-1* scion/*nrt1.1-1* stock) and heterografted (Col-0
scion/*nrt1.1-1* stock and *nrt1.1-1* scion/Col-0 stock) plants were generated to clarify
whether either or both the roots and shoots are the action point for NRT1.1 to improve K
nutrition in plants. The leaves of Col-0/*nrt1.1-1* and *nrt1.1-1*/*nrt1.1-1* plants developed
severely chlorotic phenotype after 8 d of being grown on low-K medium (Fig. 4a).
However, *nrt1.1-1*/Col-0 and Col-0/Col-0 did not exhibit any chlorotic symptom. This
result suggested that the absence of the root-part function of NRT1.1 was primarily
responsible for the sensitive symptom. This conclusion was further supported by the
observation that K levels were clearly reduced in the shoots of the plants grafted with
nrt1.1-1 root stock in low-K treatment (Fig. 4c). Furthermore, we found that the K levels
in both the shoots and roots of Col-0/*nrt1.1-1* and *nrt1.1-1*/*nrt1.1-1* plants were
significantly lower than those in the plants grafted with Col-0 rootstocks in the sufficient
K treatment (Fig. 4b). These results suggested that only the root part is the action point for
NRT1.1 to improve K nutrition in plants.

The role of root NRT1.1 in improving plant K nutrition prompted us to investigate how
NRT1.1 in roots responds to low-K stress. Real-time quantitative PCR analysis showed
that low-K treatment clearly increased the expression of *NRT1.1* in roots (Fig. 4d),
whereas the expression of the other five NO_3^- uptake transporter genes was either
decreased (*NRT2.1*) (Fig. S9a) or not significantly affected (*NRT1.2, NRT2.2, NRT2.4,*

and *NRT2.5*; Fig. S9b-e), suggesting that low-K stress might have a specific effect on the

action of NRT1.1. Consistent with gene expression, the NRT1.1-GFP protein expression

indicated by GFP-associated fluorescence in the roots of *pNRT1.1::NRT1.1-GFP*

transgenic plants was elevated in the low-K treatment than in the sufficient-K treatment

(Fig. 4e). The result was confirmed by GUS staining of the roots of *pNRT1.1::NRT1.1-*

GUS transgenic plants (Fig. 4e). These findings suggested that the activity of NRT1.1

could be up-regulated in response to low K^+ stress. Thus, we analyzed the rate of net NO_3^-

fluxes at the surface of the maturation, elongation, and meristematic zones of Col-0 roots

by using the NMT system. In all the three measured zones, the plants precultured with

low-K medium had a higher rate of net NO_3^- influx than in those precultured with

sufficient-K medium in the same testing medium (Fig. 4f), providing a direct evidence

that low-K stress stimulates NO_3^- uptake activity, which should be associated with an up-

regulation of NRT1.1 due to low-K stress.

Both uptake and root-to-shoot allocation of K require a coordination of NRT1.1

We next determined how NRT1.1 improves K nutrition in plants by investigating its role

in the root K uptake. The alkaline ion Rb^+, the closest analog of K^+, was first used to

evaluate the K^+ uptake by roots. The rates of root Rb^+ uptake in *chl1-5* and *nrt1.1-1*

mutants were considerably less than those in Col-0 plants in the medium containing either

2 mM Rb^+ or 0.05 mM Rb^+ (Fig. 5a), indicating that the loss of NRT1.1 function might

disturb the uptake of K^+ by roots. To further evaluate the contribution of NRT1.1 in root

K^+ uptake, we measured the root K^+ fluxes of the above three plant lines. We found that,

at the elongation and maturation zones, the rates of net K^+ influx of *nrt1.1-1* and *chl1-5*

mutants were less than 50% of those obtained in the Col-0 plants in both 2 mM K^+ and

0.05 mM K^+ testing medium. However, the net K^+ influx rate in the meristematic zone did

not differ significantly among these plants (Fig. 5b,c). These results suggest that the

elongation and maturation zones, but not the meristematic zone, are the target regions for

NRT1.1 to contribute to K^+ uptake by root cells.

Consistent with its role in NO_3^- uptake, NRT1.1 was expressed in the epidermis and

cortex of roots (Huang *et al*., 1996). However, NRT1.1 was also expressed in the root

central vasculature (Remans *et al*., 2006), revealing its another function: allocating root

NO_3^- to shoot parts through xylem NO_3^- loading in the root stele (Léran *et al*., 2013). This

additional function suggests that NRT1.1 might also play a role in root-to-shoot allocation

of K. Therefore, the rates of K distributed in the root and shoot were calculated. However,

the proportion of K distributed in either shoots or roots did not differ between the Col-0

plants and *nrt1.1* knockout mutants when they were grown in sufficient-K medium (Fig.

5d). Interestingly, both *chl1-5* and *nrt1.1-1* mutants had less proportion of K distributed in

the shoots, but greater proportion of K distributed in the roots compared with those in

Col-0 plants grown in the low-K medium (Fig. 5d). These results indicated that a K-

insufficient status of plants allows NRT1.1 to play the distinct role in root-to-shoot

allocation of K.

Action of NRT1.1 in favoring uptake and root-to-shoot allocation of K, respectively,

depends on its specific expression in the epidermis-cortex and central vasculature of

roots

As NRT1.1 is expressed in both the epidermis-cortex and central vasculature of roots

(Huang *et al.*, 1996; Remans *et al.*, 2006), we intended to determine whether such

expression pattern correlates with its dual roles in coordinating the uptake and root-to-

shoot allocation of K^+. To address this question, we generated transgenic plants

expressing NRT1.1 under the control of the promoters of *Sultr1;2* and *PHO1* genes,

respectively, in the *nrt1.1-1* mutant background. *Sultr1;2* is expressed in the root

epidermis and cortex (Yoshimoto, 2002), whereas *PHO1* is expressed in the root central

vasculature (Hamburger *et al.,* 2002; Wege & Poirier, 2014). Therefore, the uses of these

two promoters to drive *NRT1.1* gene in transgenic plants could, respectively, ensure

NRT1.1 expression in the epidermis-cortex and central vasculature of roots, although their

expression levels might not well match with those in the wild-type plants. The

determination of K levels in two independent *pSultr1;2::NRT1.1* transgenic lines in

comparison to those in Col-0 and *nrt1.1-1* plants revealed partial recovery of K to wild-

type levels in the plants in both sufficient-K and low-K treatments (Fig. 6a,b). However,

the calculation of the ratio of K distributed in the root and shoot showed that

pSultr1;2::NRT1.1 complementation in *nrt1.1-1* plants did not improve the root-to-shoot

allocation of K (Fig. 6c). Therefore, the expression of NRT1.1 in the epidermis and cortex

might only be responsible for improving the K uptake.

The measurement of K levels in other two independent *pPHO1::NRT1.1* transgenic lines

also showed partial recovery of K to Col-0 wild-type levels in the plants in both K

treatments, with the recovery being more distinct in the shoots (Fig. 6d,e). Furthermore,

the ratio of K distributed in the root and shoot revealed that the complementation of

pPHO1::NRT1.1 in *nrt1.1-1* plants increased the K transport from root to shoot (Fig. 6f),

suggesting that the expression of NRT1.1 in the central vasculature of roots is responsible

for supporting the root-to-shoot allocation of K.

Action of NRT1.1 in favoring the uptake and root-to-shoot allocation of K,

respectively, depends on the K$^+$ channels in the epidermis-cortex and central

vasculature of roots

Next, we investigated how the expression of NRT1.1 in the epidermis-cortex and central

vasculature, respectively, coordinated the uptake and root-to-shoot allocation of K. In

addition to a role in mediating nitrate transport across the plasmalemma, NRT1.1 was

found to act as a singling component in regulating the expression of numerous genes in

plants (Ho *et al*., 2009; Bouguyon *et al*., 2015; Wang *et al*., 2018). Therefore, the

difference in whole-genome gene expression profiles between Col-0 plants and *nrt1.1*

mutants in response to different K treatments were compared using RNA-seq. Gene

ontology (GO) analysis revealed that the differentially expressed genes (DEGs) were

involved in many physiological processes such as response to nutrient levels, response to

reactive oxygen species, and response to salicylic acid (Fig. S10). However, among the

genes associated with K uptake and trafficking, only the expression of *SKOR*, which

mediates K$^+$ loading into the xylem (Gaymard *et al*., 1998; Drechsler *et al*., 2015), was

inhibited by NRT1.1 mutation in both sufficient-K and low-K treatments, whereas that of

the other genes was either increased or not affected (Fig. 7a). This result cannot explain

the decreased root K uptake due to the loss of function of NRT1.1. Therefore, we also

investigated whether NRT1.1 acts a K$^+$ channel. To functionally test the role of NRT1.1

in K$^+$ transport activity in a heterologous expression system, we used the yeast

(*Saccharomyces cerevisiae*) mutant R5421, lacking the main K$^+$ uptake systems (*trk1Δ,*

trk2Δ; Gaber *et al*., 1988; Han *et al*., 2016). However, NRT1.1 could not complement the

growth retardation of R5421 under K deficiency (Fig. 7b), refuting the K$^+$ transport

activity of NRT1.1.

Theoretically, the flux of anions across the membrane should be accompanied by an

orthokinetic flux of cations so as to maintain the ionic balance, and this process might

depend on the cooperation between anion transporters/channels and cation

transporters/channels. In this context, the action of NRT1.1-mediated NO$_3^-$ flux across the

plasmalemma might collaborate wih K^+ channels. To test this hypothesis, we generated

nrt1.1-1/akt1, *nrt1.1-1/hak5-3*, *nrt1.1-1/skor-2*, and *nrt1.1-1/kup7* double mutants by

crossing the NRT1.1-null mutant *nrt1.1-1* with *akt1*, *hak5-3*, *skor-2*, and *kup7* mutants

and measured K levels in these plants. AKT1, HAK5, and KUP7 are the three

characterized components that contribute to K^+ uptake in *Arabidopsis* roots, and the latter

two channels only have a significant contribution in K uptake under K-limited condition

(Lagarde *et al*., 1996; Gierth *et al*., 2005; Han *et al.,* 2016). In addition, KUP7 was found

to be involved in K^+ loading into the xylem in addition to the SKOR channel (Han *et al*.,

2016). Two-way analysis of variance (ANOVA) showed that the difference of K levels in

the shoots and roots between *akt1* and *akt1/nrt1.1-1* mutants was significantly less than

that between Col-0 and *nrt1.1-1* in both K treatments (Fig. 8a,b), suggesting that the

reduced K levels due to the loss of NRT1.1 was associated with the action of AKT1.

Similarly, two-way ANOVA based on the data obtained in Col-0, *nrt1.1-1, hak5-3,*

nrt1.1-1/hak5-3, *kup7*, and *nrt1.1-1/kup7* plants showed that the reduced K levels due to

the loss of NRT1.1 were also associated with HAK5 and KUP7 in low-K medium (Fig.

8c-e). These results indicated that NRT1.1 should interplay with the K uptake channels in

the root epidermis-cortex to improve the plant K nutrition. Next, we determined whether

the KUP7 and SKOR channels are required for the action of NRT1.1 in favoring root-to-

shoot allocation of K under low K condition. Two-way ANOVA showed that the

difference in the proportion of K distributed from the roots to shoots either between *skor-*

2 and *nrt1.1-1/skor-2* or between *kup7* and *nrt1.1-1/kup7* was significantly less than that

between Col-0 and *nrt1.1-1* (Fig. 8f,h). These results indicate that the action of NRT1.1 in

favoring root-to-shoot allocation of K depends on the K^+ channels that mediate K^+

loading into the xylem.

Discussion

K^+ and NO_3^- are the major forms of potassium and nitrogen that are absorbed by roots of

most terrestrial plants. In addition, they, respectively, represent the most abundant

inorganic cation and anion sources in plant cells, and are thus also the major forms of

potassium and nitrogen that are transported within plants (Du *et al*., 2019). In this study,

we revealed a close relationship between NO_3^- and K^+ homeostasis, which was mediated

by the NO_3^- transporter NRT1.1, in the root tissues of *Arabidopsis*.

NRT1.1 was initially characterized as an NO_3^- transporter involved in root NO_3^- uptake

from the growth medium (Tsay *et al.*, 1993). Further, it was found to be a bidirectional

transporter (Remans *et al.*, 2006), thereby acting as a component involved in the xylem

NO_3^- loading in the root stele, which confers the root-to-shoot allocation of NO_3^- (Léran *et*

al., 2013). Several studies have revealed many additional functions such as regulation of

another NO_3^- transporter NRT2.1 (Ho *et al.*, 2009), auxin transport (Krouk *et al.*, 2010),

and regulation of tolerance to NH_4^+ toxicity (Jian *et al.*, 2018). These functions of NRT1.1

are independent of its NO_3^- transport activity, because the absence of NO_3^- supply in the

growth medium did not affect the above functions. However, our finding that wild-type

plants did not show higher K levels in both shoots and roots than in *nrt1.1* knockout

mutants when NH_4^+ was the sole nitrogen source shows that the activity of NO_3^- transport

across the plasmalemma is probably necessary for NRT1.1 to coordinate K^+ homeostasis

in plants (Fig. S7c,d). Theoretically, NO_3^- transport-coordinated K^+ homeostasis in plants

follows a general mechanism. However, the other nitrate uptake NRTs did not act like

NRT1.1 when coordinating K^+ homeostasis in plants. Previous studies showed that

NRT1.1 contributes to > 70% of the total root NO_3^- uptake in NO_3^--sufficient medium

(Huang *et al.*, 1996; Wang *et al.*, 1998), suggesting that the other NRTs (i.e., NRT1.2,

NRT2.1, NRT2.2, NRT2.4, and NRT2.5) might just be responsible for < 30% of NO_3^-

uptake. Therefore, an insufficient NO_3^- transport of any of these five NRTs might explain

why none of them plays an evident role in improving K^+ homeostasis in plants. This also

suggests that the action of NRT1.1 to improve K^+ homeostasis might require a higher

NO_3^- supply to ensure sufficient NO_3^- transport. This notion was supported by the

observation that the increase of NO_3^- supply elevated the K levels in wild-type plants that

have a normal NRT1.1 function (Fig. 1a,b). In addition, the finding that NRT1.1-null

mutants did not have distinguishable growth phenotype and K levels compared with those

of wild-type plants in low NO_3^- growth medium (0.2 mM NO_3^-; Fig S4) further supports

the above notion.

Interestingly, the coordination of K^+ homeostasis by NRT1.1 in *Arabidopsis*, including

root uptake and root-to-shoot allocation processes, depend on the nature of NRT1.1

expression pattern in the root epidermis-cortex and central vasculature, respectively.

Hence, determining how such expression pattern of NRT1.1 in root tissues improves the

uptake and root-to-shoot allocation of K^+ becomes necessary. Previously, NRT1.1 and

other NRT transporters were shown to be a kind of proton-coupled transporters that

transport NO_3^- across the plasmalemma together with an equal amount of proton (H^+)

influx via a symport mechanism (Tsay et $al.$, 1993; Fang et $al.$, 2016). However, the H^+

levels in the regular growth media were considerably less than those of NO_3^-. Thus, an H^+

efflux mechanism of root cells should be required to compensate for the consumed H^+ due

to NO_3^- uptake, which consequently ensures the subsequent NO_3^- uptake. The uptake of

K^+ across the plasmalemma by K^+ channels of the root cortex cells is known to be

electrically coupled to H^+-ATPase-mediated H^+ efflux (Behl & Raschke, 1987; Wang &

Wu, 2013). Since K^+ is the most abundant inorganic cation absorbed by roots, the amount

of H^+ efflux owing to K^+ uptake could be a key compensation to the consumed H^+ by

NO_3^- uptake, which might also be a mechanism to retain the charge balance during NO_3^-

$–K^+$ fluxes across the plasmalemma and thus to ensure a normal electrophysiological

environment in the cells. In this context, NRT1.1 is probably coordinated by K^+ channels

in the root epidermis-cortex. Our two-way ANOVA based on K levels in $nrt1.1/akt1$,

$nrt1.1/hak5$, and $nrt1.1/kup7$ double mutants and their corresponding monomutants and

wild-type plants supported the above notion, that is, the NRT1.1 interplay with K^+

channels in the root epidermis-cortex could improve the plant K uptake (Fig. 8a,b,c,e). As

all the investigated K^+ channels could physiologically interplay with NRT1.1 to improve

K uptake, we suggest that these interplays should function in a nonspecific manner.

The two-way ANOVA based on K distribution in the shoot and root tissues of $nrt1.1/skor$

and $nrt1.1/kup7$ double mutants and their corresponding monomutants and wild-type

plants also revealed a physiological interaction between NRT1.1 and K^+ channels in the

root central vasculature, which improves the root-to-shoot allocation of K (Fig. 8e-h). In

addition to this mechanism, transcript regulation by NRT1.1 might also play a role in

improving the root-to-shoot allocation of K, because the expression of $SKOR$ was

positively regulated by NRT1.1 (Fig. 7a). We also analyzed the K distribution in the shoot

and root tissues of $nrt1.1/akt1$ and $nrt1.1/hak5$ double mutants and their corresponding

monomutants and wild-type plants, but no physiological interaction was noted between

NRT1.1 and AKT1 or HAK5 in improving the root-to-shoot allocation of K, as revealed

by the two-away ANOVA (Fig. S11c,d), indicating that the K^+ uptake channels might not

be involved in the NRT1.1-coordinated root-to-shoot allocation of K. Notably, the

contribution of NRT1.1 to the root-to-shoot allocation of K^+ is only remarkable under the

K-limited condition, but not under the K-sufficient condition (Fig. 5d). Our finding that

insufficient NO_3^- supply did not efficiently support the role of NRT1.1 in the uptake and

root-to-shoot allocation of K^+ (Fig. 4e) indicated that a higher ratio of NO_3^- to K^+

transport across the cell plasmalemma might be a prerequisite for NRT1.1 to remarkably

contribute the root-to-shoot allocation of K^+ in plants. As the K^+ in plants could be easily

allocated to shoots after its root uptake from the growth medium (De Boer, 1999), the

amount of K^+ loading into the root xylem needs to be very higher under the K-sufficient

condition, which results in a lower ratio of NO_3^- to K^+ transport across the cell

plasmalemma in the root central vasculature. This might explain why the contribution of

NRT1.1 to the root-to-shoot allocation of K^+ is not remarkable under the K-sufficient

condition.

As both the uptake and root-to-shoot allocation of K^+ were significantly improved by

NRT1.1 under the K-limited condition (Figs 3b, 5d), the *nrt1.1* knockout mutants had

shorter roots, less biomass, and severer leaf senescence, compared with the wild-type

plants in the low-K growth medium (Fig. 2e,f,a), indicating that NRT1.1 is required for

plants to resist K deficiency. Owing to the limited availability of K^+ in most natural soils,

plants often suffer from K^+-deficiency stress (Wang & Wu, 2013). Therefore, plants have

to evolve a complex signaling and physiological regulatory network to adapt to K^+-

deficient environments, which helps them survive under K^+-deficiency stress (Ashley *et*

al., 2006; Tsay *et al.,* 2011). We found that the activity of NRT1.1-mediated NO_3^- uptake

was significantly up-regulated in response to low-K^+ stress (Fig. 4f). Considering a role of

NRT1.1 in improving both root uptake and root-to-shoot allocation of K^+ in plants, the

up-regulation of NRT1.1 activity by low-K^+ stress could be recognized as an adaption

mechanism for plants to help them survive in K^+-deficient environments. Therefore, for

crop cultivation, designing other more practical methods for improving NRT1.1-mediated

NO_3^- uptake and increasing the utilization efficiency of K fertilizers is necessary.

However, how low-K^+ stress up-regulates NRT1.1 activity is not yet known. In our study,

we observed that the rate of root NO_3^- uptake in low-K medium was clearly less than that

in sufficient-K medium (Fig. 4f). Therefore, an insufficient K supply might result in mild

nitrogen starvation in plants, which could favor an up-regulation of NRT1.1 to

compensate for the decreased NO_3^- uptake. Nevertheless, further studies are required to

experimentally verify theis assumption.

In conclusion, we showed that the NO_3^- transport activity of NRT1.1 plays an important

role in the K nutrition in plants: its expression in root epidermis-cortex coordinates with

the K^+ uptake channels such as AKT1, HAK5, and KUP7 to improve the root K^+ uptake

from the growth medium, whereas its expression in the root central vasculature coordinates with the channels that load K^+ into the xylem, such as SORK and KUP7, to facilitate the root-to-shoot allocation of K^+ (Fig. 9). In the current agricultural practices in China, excessive N and K fertilizers are usually applied by farmers; this reduces the utilization efficiencies of these fertilizers and results in environmental pollution (Guo *et al.*, 2010; Zhang, 2017). Our findings indicate that designing practical methods for enhancing NRT1.1 activity in crops might be possible to improve simultaneously the utilization efficiencies of N and K fertilizers in agricultural production.

Acknowledgments

The authors thank Dr. Philippe Nacry for sharing seeds and Dr. Yi Wang (China Agricultural University) for providing the yeast strains. This work was financially supported by the Natural Key R&D Program of China [2016YFD0200103], the Natural Science Foundation of China [31622051 and 31670258], and the Fundamental Research Funds for the Central Universities [2017XZZX002-06].

Author Contributions

X.Z.F., X.X.L. and C.W. J. conceived the project and designed the experiments; C.C. and X.X.L performed most of the experiments and analyzed data; Y.X.Z. and J.Y.Y. helped with generation of double mutants; X.Z.F. and C.W. J. wrote the article with contributions from all authors.

Supporting Information

Fig. S1 Effect of NO_3^- on the amount of K^+ absorbed per weight of roots in Col-0 plants.

Fig. S2 Root growth of *Arabidopsis thaliana* Col-0, *nrt1.1-1*, and *chl-5* plants to low-K^+ stress.

Fig. S3 Comparisons of growth and K concentration between *Ler* and *chl1-6* plants.

Fig. S4 Comparisons of growth and K concentrations between Col-0 and *nrt1.1* plants in low nitrate medium.

Fig. S5 Growth responses of Col-0, *nrt1.2*, *nrt2.1*, *nrt2.2*, *nrt2.5*, *Ler*, and *nrt2.4* plants to low-K^+ stress.

Fig. S6 Comparison of low-K sensitivity between *chl1-5* mutants and complementation line.

Fig. S7 Comparisons of growth and K concentration between Col-0 and *nrt1.1* plants in

nitrate-free medium.

Fig. S8 The K concentrations in Col-0, *nrt1.2*, *nrt2.1*, *nrt2.2*, *nrt2.5*, *Ler*, and *nrt2.4* plants.

Fig. S9 Effect of low-K$^+$ stress on the expression of *NRTs* in the root of Col-0 plants.

Fig. S10 Heatmap analysis of changes in gene expression regulated by K in the roots of

Col-0 and *nrt1.1-1* plants.

Fig. S11 Roles of K$^+$ channels in NRT1.1-imrpoved K nutrition is associated with the

biological function of K$^+$ channels.

Table S1 Primers used in this study.

References

Andersen TG, Liang D, Halkier BA, White R. 2014. Grafting *Arabidopsis*. Bio-protocol 4(13): e1164. DOI: 10.21769/BioProtoc.1164.

Ashley MK, Grant M, Grabov A. 2006. Plant responses to potassium deficiencies: a role for potassium transport proteins. *Journal of Experimental Botany* **57:** 425–36.

Barneix AJ, Breteler H. 1985. Effect of cations on uptake, translocation and reduction of nitrate in wheat seedlings. *New Phytologist* **99:** 367-379.

Behl R, Raschke K. 1987. Close coupling between extrusion of H^+ and uptake of K^+ by barley roots. *Planta* **172(4):** 531-538.

Blevins DG, Barnett NM, Frost WB. 1978. Role of potassium and malate in nitrate uptake and translocation by wheat seedlings. *Plant Physiology* **62:** 784-788.

Bouguyon E, Brun F, Meynard D, Kubeš M, Pervent M *et al.* 2015. Multiple mechanisms of nitrate sensing by *Arabidopsis* nitrate transceptor NRT1. 1. *Nature Plants* **1(3):** 15015.

Choi WG, Toyota M, Kim SH, Hilleary R, Gilroy S. 2014. Salt stress-induced Ca^{2+} waves are associated with rapid, long distance root-to-shoot signaling in plants. *Proc. Natl. Acad. Sci. USA* **111:** 6497–6502.

De Boer AH. 1999. Potassium translocation into the root xylem. *Plant Biology* **1(01):** 36-45.

Drechsler N, Zheng Y, Bohner A, Nobmann B, von Wirén N, Kunze R, Rausch C 2015. Nitrate-dependent control of shoot K homeostasis by the nitrate transporter1/peptide transporter family member NPF7. 3/NRT1. 5 and the stelar K^+ outward rectifier SKOR in Arabidopsis. *Plant Physiology* **169(4):** 2832-2847.

Du XQ, Wang FL, Li H, Jing S, Yu M *et al.* 2019. The transcription factor MYB59 regulates K^+/NO_3^- translocation in the *Arabidopsis* response to low K^+ stress. *The Plant Cell* tpc-00674.

Fageria NK, Baligar VC. 2005. Enhancing nitrogen use efficiency in crop plants. *Advances in Agronomy* **88:** 97-185.

Fang XZ, Tian WH, Liu XX, Lin XY, Jin CW, Zheng SJ. 2016. Alleviation of proton
toxicity by nitrate uptake specifically depends on nitrate transporter 1.1 in *Arabidopsis*.
New Phytologist **211(1):** 149-158.

Gaber RF, Styles CA, Fink GR. **1988.** TRK1 encodes a plasma membrane protein
required for high-affinity potassium transport in *Saccharomyces cerevisiae. Molecular*
and Cellular Biology **8:** 2848–2859.

Gaymard F, Pilot G, Lacombe B, Bouchez D, Bruneau D *et al*. **1998.** Identification
and disruption of a plant shaker-like outward channel involved in K^+ release into the
xylem sap. *Cell* **4:** 647–655.

Gierth M, Mäser P, Schroeder JI. 2005. The potassium transporter AtHAK5 functions
in K^+ deprivation-induced high-affinity K^+ uptake and AKT1 K^+ channel contribution to
K^+ uptake kinetics in *Arabidopsis* roots. *Plant Physiology* **137(3):** 1105-1114.

Guo FQ, Wang R, Chen M, Crawford NM. 2001. The *Arabidopsis* dualaffinity nitrate
transporter gene AtNRT1.1 (CHL1) is activated and functions in nascent organ
development during vegetative and reproductive growth. *The Plant Cell* **13:** 1761–1777.

Guo JH, Liu XJ, Zhang Y, Shen JL., Han WX *et al.* **2010.** Significant acidification in
major Chinese croplands. *Science* **327:** 1008-1010.

Hamburger D, Rezzonico E, Petétot JMC, Somerville C, Poirier Y. 2002.
Identification and characterization of the *Arabidopsis* PHO1 gene involved in phosphate
loading to the xylem. *The Plant Cell* **14(4):** 889-902.

Han M, Wu W, Wu WH, Wang Y. 2016. Potassium transporter KUP7 is involved in K^+
acquisition and translocation in *Arabidopsis* root under K^+-limited conditions. *Molecular*
Plant **9(3):** 437-446.

Hawkins BJ, Boukcim H, Plassard C. 2008. A comparison of ammonium, nitrate and
proton net fluxes along seedling roots of *Douglas-fir* and *lodgepole pine* grown and
measured with different inorganic nitrogen sources. *Plant, Cell and Environment* **31:**
278–287.

Ho CH, Lin SH, Hu HC, Tsay YF. 2009. CHL1 functions as a nitrate sensor in plants.
Cell **138(6):** 1184-1194.

Huang NC, Chiang CS, Crawford NM, Tsay YF. 1996. CHL1 encodes a component of
the low-affinity nitrate uptake system in *Arabidopsis* and shows cell type-specific
expression in roots. *The Plant Cell* **8:** 2183-2191.

Huang NC, Liu KH, Lo HJ, Tsay YF. 1999. Cloning and functional characterization of
an *Arabidopsis* nitrate transporter gene that encodes a constitutive component of low-
affinity uptake. *The Plant Cell* **11(8):** 1381-1392.

Ito H, Fukuda Y, Murata K, Kimura A. 1983. Transformation of intact yeast cells
treated with alkali cations. *Journal of Bacteriology* **153:** 163-168.

Jian S, Liao Q, Song H, Liu Q, Lepo JE, Guan C, Zhang J, Ismail AM, Zhang Z.
2018. NRT1. 1-Related NH_4^+ toxicity is associated with a disturbed balance between
NH_4^+ uptake and assimilation. *Plant physiology* **178(4):** 1473-1488.

Krouk G, Lacombe B, Bielach A, Perrine-Walker F, Malinska K *et al.* 2010. Nitrate-
regulated auxin transport by NRT1. 1 defines a mechanism for nutrient sensing in plants.
Developmental Cell **18(6):** 927-937.

Lagarde D, Basset M, Lepetit M, Conejero G, Gaymard F, Astruc S, Grignon C.
1996. Tissue-specific expression of *Arabidopsis* AKT1 gene is consistent with a role in
K^+ nutrition. *The Plant Journal* **9(2):** 195-203.

Léran S, Muños S, Brachet C, Tillard P, Gojon A, Lacombe B. 2013. *Arabidopsis*
NRT1. 1 is a bidirectional transporter involved in root-to-shoot nitrate translocation.
Molecular Plant **6(6):** 1984-1987.

Léran S, Varala K, Boyer JC, Chiurazzi M, Crawford N, Daniel-Vedele F *et al.* 2014.
A unified nomenclature of nitrate transporter1/peptide transporter family members in
plants. *Trends in Plant Science* **19(1):** 5-9.

Lezhneva L, Kiba T, Feria-Bourrellier A B, Lafouge F, Boutet-Mercey S *et al.* 2014.
The *Arabidopsis* nitrate transporter NRT 2.5 plays a role in nitrate acquisition and
remobilization in nitrogen-starved plants. *The Plant Journal* **80(2):** 230-241.

Li H, Yu M, Du XQ, Wang ZF, Wu WH, Quintero FJ, Jin XH, Dong JH, Wang Y.
2017. NRT1. 5/NPF7. 3 functions as a proton-coupled H^+/K^+ antiporter for K^+ loading
into the xylem in *Arabidopsis*. *The Plant Cell* **29(8):** 2016-2026.

Lin SH, Kuo HF, Canivenc G, Lin CS, Lepetit M *et al*. 2008. Mutation of the
Arabidopsis NRT1. 5 nitrate transporter causes defective root-to-shoot nitrate transport.
The Plant Cell **20(9):** 2514-2528.

Li S, Yu J, Zhu M, Zhao F, Luan S. 2012. Cadmium impairs ion homeostasis by
altering K^+ and Ca^{2+} channel activities in rice root hair cells. *Plant, Cell and Environment*
35(11): 1998-2013.

Luo BF, Du ST, Lu KX, Liu WJ, Lin XY, Jin CW. 2012. Iron uptake system mediates
nitrate-facilitated cadmium accumulation in tomato (*Solanum lycopersicum*) plants.
Journal of Experimental Botany **63:** 3127–3136.

Maathuis FJM. 2009. Physiological functions of mineral macronutrients. *Current*
Opinion in Plant Biology **12:** 250–258.

Meena VS, Maurya BR, Verma JP, Meena RS. 2016. Potassium solubilizing
microorganisms for sustainable agriculture. New Dehli: Springer.

Pyo YJ, Gierth M, Schroeder JI, Cho MH. 2010. High-affinity K^+ transport in
Arabidopsis: HAK5 and AKT1 are vital for seedling establishment and postgermination
growth under low-potassium conditions. *Plant Physiology* **153:** 863–875.

Remans T, Nacry P, Pervent M, Filleur S, Diatloff E, Mounier E, Tillard P, Forde
BG, Gojon A. 2006. The *Arabidopsis* NRT1. 1 transporter participates in the signaling
pathway triggering root colonization of nitrate-rich patches. *Proc. Natl. Acad. Sci. USA*
103(50): 19206-19211.

Shin R. 2017. Potassium sensing, signaling, and transport: toward improved potassium
use efficiency in plants. *In Plant Macronutrient Use Efficiency*. Academic Press, 149-163.

Tsay YF, Ho CH, Chen HY, Lin SH. 2011. Integration of nitrogen and potassium
signaling. *Annual Review of Plant Biology* **62:** 207–226.

Tsay YF, Schroeder JI, Feldmann KA, Crawford NM. 1993. The herbicide sensitivity
gene CHL1 of *Arabidopsis* encodes a nitrate-inducible nitrate transporter. *Cell* **72:** 705–
713.

Wang R, Liu D, Crawford NM. 1998. The Arabidopsis CHL1 protein plays a major role
in high-affinity nitrate uptake. *Proc. Natl. Acad. Sci. USA* **95:** 15134–15139.

Wang Y, Wu WH. 2010. Plant sensing and signaling in response to K^+-deficiency.
Molecular Plant **3:** 280–287.

Wang Y, Wu WH. 2013. Potassium transport and signaling in higher plants. *Annual*
Review of Plant Biology **64:** 451-476.

Wang Y, Wu WH. 2017. Regulation of potassium transport and signaling in
plants. *Current Opinion in Plant Biology* **39:** 123-128.

Wang YY, Cheng YH, Chen KE, Tsay YF. 2018. Nitrate transport, signaling, and use
efficiency. *Annual Review of Plant Biology* **69:** 85-122.

Wang, Y.Y., Hsu, P.K., and Tsay, Y.F. 2012. Uptake, allocation and signaling of nitrate.
Trends in Plant Science **17:** 458–467.

Wege S, Poirier Y. 2014. Expression of the mammalian Xenotropic Polytropic Virus
Receptor 1 (XPR1) in tobacco leaves leads to phosphate export. *FEBS Letters* **588:** 482-
489.

Yoshimoto N, Takahashi H, Smith FW, Yamaya T, Saito K. 2002. Two distinct high-
affinity sulfate transporters with different inducibilities mediate uptake of sulfate in
Arabidopsis roots. *The Plant Journal* **29(4):** 465-473.

Zhang X. 2017. A plan for efficient use of nitrogen fertilizers. *Nature* **543:** 322-323.

Zioni AB, Vaadia Y, Lips SH. 1971. Nitrate uptake by roots as regulated by nitrate
reduction products of the shoot. *Physiologia Plantarum* **24(2):** 288-290.

Figure 1. Increase of NO$_3^-$ supply improves K nutrition in *Arabidopsis* Col-0 plants

(A) and (B) The K concentration in the shoots and roots. After 4 d of germination on the basal agar media, as described in the Materials and Methods, the seedlings were transplanted to the basal agar media containing different NO$_3^-$ levels as indicated. The K concentrations were analyzed 8 d after seedling transfer. Bars represent the SD (n = 5 biological replicates). Different letters above bars indicate significant differences at $P < 0.05$ (LSD test). DW, dry weight. (C) Kinetics of net K$^+$ flux in Col-0 roots in response to NO$_3^-$ addition. The net K$^+$ flux in Col-0 roots was first measured for 5 min in the absence of NO$_3^-$, and then it was measured in the presence of 6 mM NO$_3^-$ (indicated by the red arrow). (D) The average values of root K$^+$ fluxes calculated from Figure c. Bars represent the mean ± SD. Different letters above bars indicate significant differences at $P < 0.05$ (LSD test). DW, dry weight.

Figure 2. Loss of NRT1.1 function in *Arabidopsis nrt1.1* mutant leads to hypersensitivity to low-K stress

(A) Phenotypic photographs of plants; (B) chlorophyll content of leaves; (C) maximal quantum efficiency of PS; (D) electron transport rate (ETR); (E) root elongation; (F) biomass of plants. The 4-d-old seedlings of Col-0, *nrt1.1-1*, and *chl1-5* were transferred to an agar medium containing 2 mM or 0.05 mM K^+, as described in the Materials and Methods section. The analyses were performed 8 d after seedling transfer. Bars represent the mean ± SD (n = 4–5). (B), (E), and (F) Different letters above bars indicate significant differences at $P < 0.05$ (LSD test); asterisks indicate a significant genotype by treatment interaction (*** $P < 0.001$, two-way ANOVA). (D) Asterisks show significant differences compared with the Col-0; ns, non-significant (* $P < 0.05$, *** $P < 0.001$, two-tailed Student's *t*-test).

Figure 3. NRT1.1 is required for ensuring K nutrition in *Arabidopsis*

(A) and (B) Shoot and root K concentrations of Col-0, *nrt1.1-1*, and *chl-5* plants. (C) and (D) Shoot and root K concentrations of Col-0, *chl1-5*, and *pNRT1.1::NRT1.1-GFP* plants. (E) Phenotypic photographs of Col-0, *chl-5*, and *pNRT1.1::NRT1.1-GFP* plants. The 4-d-old seedlings were transferred to an agar medium containing 2 mM K⁺ or 0.05 mM K⁺, as described in the Materials and Methods section. The analyses were performed 8 d after seedling transfer. Bars represent the mean ± SD (n = 5). Asterisks indicate significant differences between genotypes; ns, non-significant (* $P < 0.05$, ** $P < 0.01$, two-tailed Student's *t*-test). DW, dry weight.

Figure 4. Improvement of K nutrition by NRT1.1 in *Arabidopsis* is a root behavior, and low-K stress stimulates NRT1.1 activity in the roots

(A) The phenotypic photographs; (B) and (C) the K concentrations of grafted plants. The unified seedlings were transferred to an agar medium containing 2 mM K$^+$ or 0.05 mM K$^+$. The analyses were performed 8 d after seedling transfer.

(D) Relative expression of *NRT1.1*. (E) NRT1.1-GFP expression and GUS staining in *pNRT1.1::NRT1.1-GFP* and *pNRT1.1::NRT1.1-* GUS transgenic plants, respectively. (F) Average values of net NO$_3^-$ fluxes in the meristematic, elongation, and maturation zones of Col-0 roots. The 4-d-old seedlings were transferred to an agar medium containing 2 mM K$^+$ or 0.05 mM K$^+$, as described in the Materials and Methods section. The analyses were performed 5 d after seedling transfer. The relative expression levels were normalized to the levels of *UBQ10.* The negative value in (F) indicates net influx. Bars represent the mean ± SD (n = 5); different letters above bars indicate significant differences at *P* < 0.05 (LSD test). DW, dry weight.

Figure 5. Both the uptake and root-to-shoot allocation of K$^+$ require an action of NRT1.1 in *Arabidopsis*

(A) Rb$^+$ uptake rate of roots. The 4-d-old seedlings were transferred to an agar medium containing 2 mM Rb$^+$ or 0.05 mM Rb$^+$ for 12 h, as described in the Materials and Methods section. Bars represent the mean ± SD (n = 5).

(B) and (C) The net K$^+$ fluxes in root. The 4-day-old seedlings were used to measure the net K$^+$ fluxes in the meristematic, elongation, and maturation zones under 2 mM K$^+$ or 0.05 mM K$^+$ treatment. Negative indicates net influx; bars represent the mean ± SD (n = 15).

(D) Proportions of K distributed in the shoots and roots. The 4-d-old seedlings were transferred to an agar medium containing 2 mM K$^+$ or 0.05 mM K$^+$, as described in the Materials and Methods section. The analyses were performed 8 d after seedling transfer. Bars represent the mean ± SD (n = 5); asterisks indicate significant differences compared with Col-0 plants; ns, non-significant (* $P < 0.05$, ** $P < 0.01$, two-tailed Student's *t*-test).

Figure 6. *NRT1.1* expression driven by *Sultr1;2* and *PHO1* promoters partially rescued the uptake and root-to-shoot allocation of K in *nrt1.1* mutants, receptively.

(A) and (B) K concentrations and (C) distributions in Col-0, *nrt1.1-1*, and two independent *pSultr1;2:NRT1.1*-transformed *nrt1.1-1* lines (*pSultr1;2:NRT1.1-1/-2*).

(D) and (E) K concentrations and (F) distributions in Col-0, *nrt1.1-1*, and two independent *pPHO1:NRT1.1-* transformed *nrt1.1-1* lines (*pPHO1:NRT1.1-1/-2*). The 4-d-old seedlings were transferred to an agar media containing 2 mM K$^+$ or 0.05 mM K$^+$ for 8 d, as described in the Materials and Methods section. Bars represent the mean ± SD (n = 5); different letters above bars indicate significant differences at $P < 0.05$ (LSD test).

Figure 7. Both the regulation of gene expression and K⁺ transport activity might not explain the effect of NRT1.1 in improving K nutrition in *Arabidopsis*

(A) Heatmap analysis of genes enriched in the GO terms: K^+ homeostasis, K^+ transport, and other processes of K^+ metabolism in the roots of *Arabidopsis* Col-0 and *nrt1.1-1* plants. The 4-d-old seedlings were transferred to a media containing 2 mM or 0.05 mM K^+ for 5 d, and roots were collected to perform transcriptome analysis, as described in the Materials and Methods section. Red color represents significantly up- or down- regulated genes compared with those in the control group. Changes in response to K treatment with a *p*-value of <0.05 were selected for GO analysis.

(B) Functional analysis of NRT1.1 in *Saccharomyces cerevisiae*. K^+ uptake activity was analyzed in yeast R757 strain and the K^+ import mutant strain R5421 (*trk1Δ*, *trk2Δ*), which were transformed with the expression constructs, as described in the Materials and Methods section. pDR196, empty vector.

Figure 8. Physiological interactions between NRT1.1 and K⁺ channels

(A) and (B) The K concentrations in Col-0, *nrt1.1-1*, *akt1,* and *nrt1.1-1/akt1* plants. (C) and (D) The K concentrations in Col-0, *nrt1.1-1*, *hak5-3*, and *nrt1.1-1/hak5-3* plants. (E) K concentrations and (F) distributions in Col-0, *nrt1.1-1*, *kup7*, and *nrt1.1-1/kup7* plants. (G) K concentrations and (H) distributions in Col-0, *nrt1.1-1*, *skor-2*, and *nrt1.1-1/skor-2* plants. The 4-d-old seedlings were transferred to an agar medium containing 2 mM K⁺ or 0.05 mM K⁺ for 8 d and subsequently analyzed for K levels, as described in the Materials and Methods section. Bars represent the mean ± SD (n = 5). Different letters above bars indicate significant differences at $P < 0.05$ (LSD test). Asterisks indicate a significant genotype by genotype interaction; ns, non-significant (* $P < 0.05$, ** $P < 0.01$, two-way ANOVA).

Figure 9. **Schematic model of how NRT1.1 responds to low-K stress in *Arabidopsis*.**

Under insufficient K growth conditions, the expression of the *NRT1.1* gene and its encoding protein in the root epidermis-cortex and central vasculature was induced, which coordinated with the K^+ uptake channels (such as AKT1, HAK5, and KUP7) and xylem K^+ loaders (such as SORK and KUP7) to improve root K^+ uptake from the external environment and facilitate K^+ root-to-shoot allocation, respectively.