Maintenance of fertility in the face of meiotic drive

Lara Meade [a] [*], Sam Finnegan [a], Ridhima Kad [a], Kevin Fowler [a] & Andrew Pomiankowski [a, b]

[a] Department of Genetics, Evolution and Environment, University College London, Gower

Street, London, WC1E 6BT, UK

[b] CoMPLEX, University College London, Gower Street, London, WC1E 6BT, UK

Keywords: accessory gland, multiple mating, sex ratio distorter, sperm competition, testis

Word count: 3200

Online-only elements: Appendix

Submitted to The American Naturalist as a Note.

Second revision

Abstract

Selfish genetic elements that gain a transmission advantage through the destruction of

sperm have grave implications for drive male fertility. In the X-linked SR meiotic drive

system of a stalk-eyed fly, we found that drive males have greatly enlarged testes and

maintain high fertility despite the destruction of half their sperm, even when challenged

with fertilising large numbers of females. Conversely, we observed reduced allocation of

resources to the accessory glands that probably explains the lower mating frequency of SR

males. Body size and eyespan were also reduced, which are likely to impair viability and pre-

copulatory success. We discuss the potential evolutionary causes of these differences

between drive and standard males.

Introduction

Meiotic drive genes gain a transmission advantage through manipulation of meiosis or
gametogenesis and are likely to have profound ecological and evolutionary consequences,
ranging from the evolution of sex determination systems and changes in karyotype, to
impacts on population persistence and sexual selection (Hurst and Werren 2001; Jaenike
2001; Werren 2011; Lindholm et al. 2016). Drivers have been uncovered in a wide range of
taxa, with a preponderance of linkage to the sex chromosomes in the heterogametic sex
(Hurst and Pomiankowski 1991; Jaenike 2001; Taylor and Ingvarsson 2003). When meiotic
drive occurs in males, it severely disrupts the maturation and fertilisation capacity of non-
carrier sperm, imposing a fertility disadvantage to organismal fitness (Price and Wedell
2008) which is exaggerated under conditions of sperm competition (Taylor et al. 1999;
Angelard et al. 2008; Price et al. 2008*a*) and typically has pleiotropic viability costs in both
sexes (Burt and Trivers 2006).

The extent to which these and other detrimental effects of sperm-killer drive promote
adaptive responses in the host species has received limited attention. There is an extensive
literature on genetic elements that interfere and suppress the action of drive. For example,
in *Drosophila* species, suppressors of X-linked drive have been found on the Y chromosome
(Carvalho et al. 1997; Cazemajor et al. 1997; Branco et al. 2013) and throughout the rest of
the genome (Carvalho and Klaczko 1993; Atlan et al. 2003; Tao et al. 2007). A more recent
suggestion is that drive may promote the evolution of female polyandry in order to dilute
the ejaculates of drive males (Haig and Bergstrom 1995; Zeh and Zeh 1997; Wedell 2013).
There is some evidence for this from experimental evolution studies using populations

exposed to meiotic drive in *D. pseudoobscura* (Price et al. 2008*b*) and *Mus musculus*

(Manser et al. 2017), and from natural populations in which the rate of multiple mating

correlates negatively with the frequency of drive in *D. pseudoobscura* (Price et al. 2014) and

D. neotestacea (Pinzone and Dyer 2013). Female mate choice may additionally evolve in

response to drive. In stalk-eyed flies, meiotic drive has been linked to small eyespan, which

may allow females to avoid mating with carrier males through assessing eyespan (Wilkinson

et al. 1998*b*; Cotton et al. 2014). Female house mice could avoid mating with drive males

through detecting unique major histocompatibility alleles linked to the driving *t* complex

(Silver 1985; Lindholm et al. 2013), although evidence remains unclear (Lindholm and Price

2016).

Another, as yet unexplored, route by which males could adapt to drive is by increasing the

allocation of resources to sperm production, to offset the destructive effect of drive on

gametogenesis. Sperm number is positively correlated with testis size in many intra-specific

studies (Gage 1994; Fry 2006; Hettyey and Roberts 2006) and increased testis size is a well

characterised evolutionary response to heightened sperm competition favouring greater

sperm production (Hosken and Ward 2001; Pitnick et al. 2001; Simmons and García-

González 2008; Gay et al. 2009). The loss of sperm in drive males could be compensated for

by increased investment in testis. Meiotic drive elements are typically found within

inversions or other areas of low recombination that keep drive and insensitive responder

loci together (Palopoli and Wu 1996; Johns et al. 2005; Dyer et al. 2007), facilitate the

spread of modifiers that enhance transmission distortion (Hartl 1975; Larracuente and

Presgraves 2012) and are predicted to be enriched for male beneficial sexually-antagonistic

alleles (Rydzewski et al. 2016). For similar reasons, alleles that enable compensatory

investment in testes could become associated with the drive haplotype.

We test this idea using the Malaysian stalk-eyed fly species *Teleopsis dalmanni*. This species

harbours SR, an X-linked driver, which produces strongly female-biased broods due to the

destruction of Y-bearing sperm (Presgraves et al. 1997; Wilkinson and Sanchez 2001).

Meiotic drive arose around 2-3.5 Mya in the *Teleopsis* clade, and the X^{SR} drive chromosome

in *T. dalmanni* is estimated to have diverged from a non-driving ancestor (X^{ST}) around 1 Mya

(Swallow et al. 2005; Paczolt et al. 2017), and is characterised by a large inversion(s)

covering most of the X chromosome (Johns et al. 2005; Paczolt et al. 2017). X^{SR} is found at

appreciable frequencies (10 – 30%) across populations and generations (Wilkinson et al.

2003; Cotton et al. 2014) but appears to lack genetic suppressors (Reinhold et al. 1999;

Wolfenbarger and Wilkinson 2001; Paczolt et al. 2017). This means that there has been

ample time and opportunity for adaptive responses to selection to evolve in male carriers of

the drive chromosome.

We determined whether SR and standard (ST) males differed in their reproductive (testis

and accessory gland size) and morphological traits (eyespan and body size). Testis size

predicts the amount of sperm found within female storage (Fry 2006). Accessory glands

produce all non-sperm components of the ejaculate, and accessory gland size is positively

associated with male mating frequency (Baker et al. 2003; Rogers et al. 2005*a*, 2005*b*).

Body size and eyespan are also important predictors of male mating frequency (Wilkinson et

al. 1998*a*; Small et al. 2009; Cotton et al. 2010). We determined SR and ST sperm production

by mating them to low or high numbers of females over a 10-hour period, counting the

number of fertilized eggs produced. Males were also exposed to females over a short time

period (30-minutes) to compare the copulation rate of SR and ST males.

Methods

Details of stock collection and day-to-day upkeep can be found in the Appendix.

Experimental males were taken from the SR-stock population in which males are a ~50:50

mix of X^{SR} and X^{ST} genotypes. Experimental females were taken from the ST-stock

population, which lacks meiotic drive. Single non-virgin males were allowed to mate freely

with either one or five virgin ST-stock females, over a period of 10 hours. Mated females

were allowed to lay eggs for 14 days, by which time most females had stopped laying fertile

eggs. Fecundity was recorded through egg counts, and egg hatch was used as an estimate

for fertility. On the following day, experimental males, and a similar number of unmated

males, were anaesthetised on ice and their testes and accessory glands were removed (fig.

1A) and photographed under differential interference contrast microscopy. Organ area was

measured at x50 magnification by tracing the outline. Male eyespan (Hingle et al. 2001) and

a proxy for body size, thorax length, (Rogers et al. 2008) were measured.

In a second experiment, SR-stock males were introduced to two ST-stock non-virgin females

at artificial dawn. All copulations were counted during 30 minutes. To minimise any effects

on mating frequency due to female choice, the experimental males were standardised to

have a narrow range of eyespan (7.5 – 8.5 mm).

121 Males from both experiments were genotyped using either two X-linked INDEL markers,

122 *comp162710* and *cnv395*, or a microsatellite marker, *ms395*. Allele size of these markers

123 reliably indicates the SR genotype of the males in our laboratory stocks (Meade et al. 2018).

125 *Statistical analysis*

127 We tested if male genotypes differed in their morphological (body size and eyespan; linear

128 models) and reproductive traits (testis size and accessory gland size; linear mixed effects

129 models). Differences in relative trait sizes between genotypes, as well as in absolute trait

130 sizes (models where body size is excluded) are reported. The total number of fertile eggs

131 (Poisson generalised linear mixed effects model (GLMM)) and proportion fertility (fertile

132 eggs, non-fertile eggs; binomial GLMM) of females are compared when mated to SR (i.e.

133 XSR/Y genotype) or ST (i.e. XST/Y genotype) males. We also tested if male reproductive

134 traits, and their interaction with male genotype, were important predictors of fertility.

135 Lastly, we tested whether SR and ST males differed in their mating frequency over 30-

136 minutes by comparing the likelihood that SR and ST males mate at all (binomial GLMM), as

137 well as the total number of copulations among males that mated at least once (Poisson

138 GLMM).

140 To avoid collinearity of male morphological and reproductive traits with body size, models

141 used residual values (Dormann et al. 2013). Where appropriate, experimental batch was

142 included as a random effect. Further details and model effect sizes can be found in the

143 Appendix.

Results

SR trait size

SR males had small body size (mean ± s.e. 2.290 ± 0.013 mm) compared to ST males (2.336 ±

0.009 mm; $F_{1,357}$ = 8.745, P = 0.003; fig. 1B). SR males also had small absolute (SR: 8.048 ±

0.046mm; ST: 8.402 ± 0.031mm; $F_{1,357}$ = 42.631, P < 0.001; fig. 1B) and relative eyespan

($F_{1,355}$ = 0.713, P = 0.016), especially when body size was small (body size by genotype

interaction $F_{1,355}$ = 4.175, P = 0.042).

Despite their small body size, SR testis size was large (1.940 ± 0.050 mm^2) compared to ST

males (1.54 ± 0.028 mm^2; $F_{1,280.16}$ = 73.796, P < 0.001; fig. 1C). SR males also had large

relative testis size ($F_{1,282.78}$ = 99.982, P < 0.001). In contrast, SR males had small absolute (SR:

0.306 ± 0.011 mm^2; ST: 0.348 ± 0.010 mm^2; $F_{1,335.36}$ = 16.353, P < 0.001; fig. 1D) and relative

accessory gland size ($F_{1,334.03}$ = 7.801, P = 0.006). Taking relative values for each genotype,

eyespan ($F_{1,286}$ = 19.892, P <0.001) and accessory gland size ($F_{1,274.418}$ = 26.008, P <0.001)

increased with testes size, but the rate was reduced in SR males (interaction eyespan: $F_{1,286}$

= 5.261, P = 0.023, fig. A1; interaction accessory glands: $F_{1,268}$ = 8.375, P = 0.004, fig. A2).

SR fertility

SR males did not differ from ST males in total (mean ± s.e. SR: 112.047 ± 8.290, ST: 107.053

± 5.597; χ^2_1 = 2.416, P = 0.120, N = 215; fig. 2A, 2B) or proportion fertility (SR: 0.833 ± 0.025,

ST: 0.762 ± 0.019; χ^2_1 = 2.469, P = 0.116, N = 215) when kept with females over an extended

10-hour period. Males mating with five females achieved higher total fertility (one female:

79.231 ± 5.090, five females: 138.123 ± 6.653; χ^2_1 = 43.698, P < 0.001, N = 215) but a lower

proportion fertility (one female: 0.804 ± 0.024, five females: 0.763 ± 0.199; χ^2_1 = 6.021, P =

0.014, N = 215) than those mating with a single female. The interaction between mating

group (one or five females) and genotype did not influence total (χ^2_1 = 0.591, P = 0.442, N =

215) or proportion fertility (χ^2_1 = 1.377, P = 0.241, N = 215).

Male testis size was an important predictor of fertility. Both total (χ^2_1 = 5.897, P = 0.015, N =

165; fig. 2C, 2D) and proportion fertility (χ^2_1 = 18.837, P < 0.001, N = 165) were greater

amongst males with larger testis size, even when accounting for male body size (total: χ^2_1 =

6.216, P = 0.013, N = 165; proportion: χ^2_1 = 16.646, P < 0.001, N = 165). The addition of testis

size did not alter the relationship between genotype and total (χ^2_1 = 0.018, P = 0.895, N =

173) or proportion fertility (χ^2_1 = 0.260, P = 0.610, N = 173). There was no interaction

between testis size and genotype predicting total (χ^2_1 = 0.164, P = 0.686, N = 173) or

proportion fertility (χ^2_1 = 0.617, P = 0.432, N = 173). Accessory gland size did not predict

total (χ^2_1 = 0.032, P = 0.858, N = 165) or proportion fertility (χ^2_1 = 0.160, P = 0.689, N = 165).

SR mating frequency

A total of 493 copulations from 193 males were observed over the 30-minute mating trials.

SR males (mean ± s.e. 2.750 ± 0.175, N = 81) copulated fewer times on average than ST

males (3.550 ± 0.186, N = 76; χ^2_1 = 6.304, P = 0.012; fig. 1E), but were not less likely to mate

at least once (SR: 81/104, ST: 76/89; χ^2_1 = 1.665, P = 0.197, N = 193).

Discussion

One of the main features of drive in males is reduced sperm production due to the dysfunction of non-carrier sperm. This has been reported to cause a loss in fertility in a variety of species including *Drosophila* (Hartl et al. 1967; Jaenike 1996; Angelard et al. 2008; Price et al. 2012; Pinzone and Dyer 2013), house mice (Carroll et al. 2004), and *Silene alba* (Taylor et al. 1999). Here, we present evidence that SR males in *T. dalmanni* overcome this deficit by having greatly enlarged testes. SR males carry an extreme form of the X^{SR} drive chromosome, siring female-only broods due to the dysfunction of Y-bearing gametes. Despite gamete loss, SR males achieve fertility at a level equivalent to that of ST males, both when exposed to a single female or 5 females over a 10-hour period (fig. 2). Our results contradict a previous study which found an SR fertility deficit using a similar design (Wilkinson et al. 2006). But this study measured fertility as the number of adults that eclosed, compounding fertility with egg-to-adult survival. Recent work shows larval survival is reduced in drive heterozygous females (Finnegan et al. 2019), which could account for the drop in SR male fertility. The patterns in *T. dalmanni* are in contrast to other insect species with X-linked meiotic drive which generally show a deficiency in fertility of drive males either after a single or multiple matings (Jaenike 1996; Atlan et al. 2004; Angelard et al. 2008; Price et al. 2012; Pinzone and Dyer 2013).

These experiments were designed to test whether daily sperm reserves differ between SR and ST males, not to replicate normal levels of mating observed under natural conditions which occur at far lower rates (Cotton et al. 2015). On dissection, we discovered that SR males have greatly enlarged testes (fig. 1C), about 26% larger than ST males. This difference

remained after controlling for body size (fig. 1C). Our interpretation is that the increase in

testis size allowed SR males to compensate for the loss of sperm due to the action of

meiotic drive. This is supported by the finding that fertility increased with increasing testis

size, both for absolute and relative testis size, in both SR and ST males (fig. 2). Our

interpretation also aligns with previous findings that SR male ejaculates deliver similar

numbers of sperm as ST males, after single and multiple matings (Meade et al. 2018).

Despite the destruction of half their sperm, the increased investment in SR testis size (i.e.

sperm production) allows them to deliver sufficient sperm to achieve similar fertility as ST

males. To further understand the extent of this compensation, we need to assess SR male

success under sperm competition, which is the norm in *T. dalmanni* (Wilkinson et al. 1998*a*;

Baker et al. 2001; Corley et al. 2006). Previous work suggests that SR males perform poorly

under sperm competition (Wilkinson et al. 2006) but this assessment again does not take

account of the lower egg-to-adult viability of X^{SR} carriers (Finnegan et al. 2019) which could

simulate an advantage of ST males in sperm competition. In our experimental design,

autosomal background was standardised across SR and ST males. So, it seems likely that

control of testis size is linked to alleles that are located in the X^{SR} chromosomal inversion

and that such alleles arose as an adaptive response to sperm dysfunction caused by drive,

but further investigation is needed to establish this view.

We found morphological trait divergence in accessory gland size, which are small in SR

males, even after controlling for body size (fig. 1D). Previous work in *T. dalmanni* shows that

accessory gland size is linked with the mating rate (Baker et al. 2003; Rogers et al. 2005*a*).

This might explain why the mating frequency of SR males was low, being about 75% of the

rate for ST males over a 30-minute period (fig. 1E). In addition, SR males have small body

size and small eyespan for their body size (fig. 1), traits likely to reduce male mating success,

both in male-male agonistic interactions (Panhuis and Wilkinson 1999; Small et al. 2009) and

in attracting and mating with females (Wilkinson and Reillo 1994; Hingle et al. 2001; Cotton

et al. 2010). The increased allocation of resources to testes in SR males potentially causes a

reduction in the resources available for investment in accessory glands, as both traits

develop over several weeks post-eclosion (Baker et al. 2003; Rogers et al. 2008). Resource

competition with testes is not an obvious reason for reduced body size and eyespan which

are determined during larval development. However, expression of these traits might be

connected via juvenile hormone which has been shown to mediate a trade-off between

eyespan and testes in stalk-eyed flies (Fry 2006).

Small body size and eyespan are also likely to arise from the low genetic condition of drive

males. The *T. dalmanni* SR inversion(s) covers nearly all of the X chromosome, capturing one

third of the stalk-eyed fly genome (Johns et al. 2005; Paczolt et al. 2017). X^{SR} alleles will be

subject to weak natural selection due to reduced recombination and liable to accumulate

deleterious mutational effects (Kirkpatrick 2010). Consistent with a lack of recombination,

there are 955 fixed sequence differences between transcripts linked to X^{SR} and X^{ST}

(Reinhardt et al. 2014). Such mutations are expected to have a negative effect on costly,

condition-dependent traits, such as body size and eyespan, whose expression is affected by

multiple loci distributed throughout the genome (David et al. 2000; Cotton et al. 2004;

Bellamy et al. 2013). Given SR males have small eyespan, they will be unattractive and gain

fewer mating opportunities. Consequently, investment in accessory glands which enable

higher mating rates will give lower returns than the diversion of resources into larger testes

which allow SR males to produce ejaculates of equivalent size to those of ST males, and be

able to compete under the conditions of high sperm competition seen in stalk-eyed flies.

These ideas about linking resource allocation, condition and mating rates need further

investigation, in particular under the mating conditions that occur in the wild.

Here we demonstrate for the first time that through investment in testis, drive males can

maintain fertility, despite sperm destruction. Other responses to drive, such as genetic

suppression, polyandry and female choice, reduce the transmission advantage gained by

drive, and lead to reductions in the equilibrium frequency of drive (Hartl 1975; Taylor and

Jaenike 2002; Holman et al. 2015). In sharp contrast, increased investment in sperm

production intensifies the transmission of drive, because the fertility gain to the individual

male is also beneficial to the drive element itself. Such an association with meiotic drive has

neither been theoretically modelled nor empirically studied previously, but has implications

for the spread and equilibrium frequency of drive in natural populations.

Authors' contributions

All authors contributed to conceiving the project and methodology; LM and RK collected

data on fertility and morphology; SF collected data on mating frequency; LM analysed the

data; LM, KF and AP led the writing of the manuscript. All authors contributed critically to

the drafts and gave final approval for publication.

Acknowledgements

Funding was provided by a NERC Studentship held by LM, and by EPSRC (EP/F500351/1,

EP/I017909/1) awards to AP and NERC grants (NE/G00563X/1, NE/R010579/1) to KF and AP.

Data accessibility

Data will be uploaded to the Dryad Digital Repository

Ethical statement

No ethical approval was required for this research

Literature cited

Angelard, C., C. Montchamp-Moreau, and D. Joly. 2008. Female-driven mechanisms,

ejaculate size and quality contribute to the lower fertility of sex-ratio distorter males in

Drosophila simulans. BMC Evolutionary Biology 8:326.

Atlan, A., C. Capillon, N. Derome, D. Couvet, and C. Montchamp-Moreau. 2003. The

evolution of autosomal suppressors of sex-ratio drive in *Drosophila simulans*. Genetica

117:47–58.

Atlan, A., D. Joly, C. Capillon, and C. Montchamp-Moreau. 2004. *Sex-ratio* distorter of

Drosophila simulans reduces male productivity and sperm competition ability. Journal

of Evolutionary Biology 17:744–751.

Baker, R. H., R. Ashwell, T. Richards, K. Fowler, T. Chapman, and A. Pomiankowski. 2001.

Effects of multiple mating and male eye span on female reproductive output in the

stalk-eyed fly, *Cyrtodiopsis dalmanni*. Behavioral Ecology 12:732–739.

Baker, R. H., M. Denniff, P. Futerman, K. Fowler, A. Pomiankowski, and T. Chapman. 2003.

Accessory gland size influences time to sexual maturity and mating frequency in the

stalk-eyed fly, *Cyrtodiopsis dalmanni*. Behavioral Ecology 14:607–611.

Bellamy, L., N. Chapman, K. Fowler, and A. Pomiankowski. 2013. Sexual traits are sensitive

to genetic stress and predict extinction risk in the stalk-eyed fly, *Diasemopsis meigenii*.

Evolution 67:2662–2673.

Branco, A. T., Y. Tao, D. L. Hartl, and B. Lemos. 2013. Natural variation of the Y chromosome

suppresses sex ratio distortion and modulates testis-specific gene expression in

Drosophila simulans. Heredity 111:8–15.

Burt, A., and R. Trivers. 2006. Genes in Conflict: The biology of selfish genetic elements.

Cambridge, MA: BelknapHarvard. Harvard University Press, Cambridge, CA, CA.

Carroll, L. S., S. Meagher, L. Morrison, D. J. Penn, and W. K. Potts. 2004. Fitness effects of a

selfish gene (the *Mus t* complex) are revealed in an ecological context. Evolution

58:1318–1328.

Carvalho, A. B., and L. B. Klaczko. 1993. Autosomal suppressors of sex-ratio in *Drosophila*

mediopunctata. Heredity 71:546–551.

Carvalho, A. B., S. C. Vaz, and L. B. Klaczko. 1997. Polymorphism for Y-linked suppressors of

sex-ratio in two natural populations of *Drosophila mediopunctata*. Genetics 146:891–

902.

Cazemajor, M., C. Landré, and C. Montchamp-Moreau. 1997. The Sex-Ratio trait in

Drosophila simulans: Genetic analysis of distortion and suppression. Genetics 147:635–

642.

Corley, L. S., S. Cotton, E. McConnell, T. Chapman, K. Fowler, and A. Pomiankowski. 2006.

Highly variable sperm precedence in the stalk-eyed fly, *Teleopsis dalmanni*. BMC

Evolutionary Biology 6:53.

Cotton, A. J., S. Cotton, J. Small, and A. Pomiankowski. 2015. Male mate preference for

female eyespan and fecundity in the stalk-eyed fly, *Teleopsis dalmanni*. Behavioral

Ecology 26:376–385.

Cotton, A. J., M. Földvári, S. Cotton, and A. Pomiankowski. 2014. Male eyespan size is

associated with meiotic drive in wild stalk-eyed flies (*Teleopsis dalmanni*). Heredity

112:363–9.

Cotton, S., K. Fowler, and A. Pomiankowski. 2004. Condition dependence of sexual

ornament size and variation in the stalk-eyed fly *Cyrtodiopsis dalmanni* (Diptera:

Diopsidae). Evolution 58:1038–46.

Cotton, S., J. Small, R. Hashim, and A. Pomiankowski. 2010. Eyespan reflects reproductive

quality in wild stalk-eyed flies. Evolutionary Ecology 24:83–95.

David, P., T. Bjorksten, K. Fowler, and A. Pomiankowski. 2000. Condition-dependent

signalling of genetic variation in stalk-eyed flies. Nature 406:186–188.

Dormann, C. F., J. Elith, S. Bacher, C. Buchmann, G. Carl, G. Carré, J. R. G. Marquéz, et al.

2013. Collinearity: A review of methods to deal with it and a simulation study

evaluating their performance. Ecography 36:027–046.

Dyer, K. A., B. Charlesworth, and J. Jaenike. 2007. Chromosome-wide linkage disequilibrium

as a consequence of meiotic drive. Proceedings of the National Academy of Sciences

104:1587–1592.

Finnegan, S. R., N. J. White, D. Koh, M. F. Camus, K. Fowler, and A. Pomiankowski. 2019.

Meiotic drive reduces egg-to-adult viability in stalk-eyed flies. bioRxiv 690321.

Fry, C. L. 2006. Juvenile hormone mediates a trade-off between primary and secondary

sexual traits in stalk-eyed flies. Evolution and Development 8:191–201.

Gage, M. J. G. 1994. Associations between body size, mating pattern, testis size and sperm

lengths across butterflies. Proceedings of the Royal Society B: Biological Sciences

258:247–254.

Gay, L., D. J. Hosken, R. Vasudev, T. Tregenza, and P. E. Eady. 2009. Sperm competition and

maternal effects differentially influence testis and sperm size in *Callosobruchus*

maculatus. Journal of Evolutionary Biology 22:1143–1150.

Haig, D., and C. T. Bergstrom. 1995. Multiple mating, sperm competition and meiotic drive.

Journal of Evolutionary Biology 8:265–282.

Hartl, D. L. 1975. Modifier theory and meiotic drive. Theoretical Population Biology 7:168–

174.

Hartl, D. L., Y. Hiraizumi, and J. Crow. 1967. Evidence for sperm dysfunction as the

mechanism of segregation distortion in *Drosophila melanogaster*. Genetics 58:2240–

2245.

Hettyey, A., and J. D. Roberts. 2006. Sperm traits of the quacking frog, *Crinia georgiana*:

Intra- and interpopulation variation in a species with a high risk of sperm competition.

Behavioral Ecology and Sociobiology 59:389–396.

Hingle, A., K. Fowler, and A. Pomiankowski. 2001. The effect of transient food stress on

female mate preference in the stalk-eyed fly *Cyrtodiopsis dalmanni*. Proceedings of the

Royal Society B: Biological Sciences 268:1239–1244.

Holman, L., T. A. R. Price, N. Wedell, and H. Kokko. 2015. Coevolutionary dynamics of

polyandry and sex-linked meiotic drive. Evolution 69:709–720.

Hosken, D. J., and P. I. Ward. 2001. Experimental evidence for testis size evolution via sperm

competition. Ecology Letters 4:10–13.

Hurst, G. D. D., and J. H. Werren. 2001. The role of selfish genetic elements in eukaryotic

evolution. Nature Reviews Genetics 2:597–606.

Hurst, L. D., and A. Pomiankowski. 1991. Causes of sex ratio bias may account for unisexual

sterility in hybrids: A new explanation of Haldane's rule and related phenomena.

Genetics 128:841–858.

Jaenike, J. 1996. Sex-Ratio meiotic drive in the *Drosophila quinaria* group. The American

Naturalist 148:237–254.

———. 2001. Sex chromosome meiotic drive. Annual Review of Ecology and Systematics

32:25–49.

Johns, P. M., L. L. Wolfenbarger, and G. S. Wilkinson. 2005. Genetic linkage between a

sexually selected trait and X chromosome meiotic drive. Proceedings of the Royal

Society B: Biological Sciences 272:2097–2103.

Kirkpatrick, M. 2010. How and why chromosome inversions evolve. PLoS Biology

8:e1000501.

Larracuente, A. M., and D. C. Presgraves. 2012. The selfish *Segregation Distorter* gene

complex of *Drosophila melanogaster*. Genetics 192:33–53.

Lindholm, A. K., K. A. Dyer, R. C. Firman, L. Fishman, W. Forstmeier, L. Holman, H.

Johannesson, et al. 2016. The ecology and evolutionary dynamics of meiotic drive.

Trends in Ecology & Evolution 31:315–326.

Lindholm, A. K., K. Musolf, A. Weidt, and B. König. 2013. Mate choice for genetic

compatibility in the house mouse. Ecology and Evolution 3:1231–1247.

Lindholm, A. K., and T. A. R. Price. 2016. The evolutionary consequences of selfish genetic

elements. Current Zoology 62:655–658.

Manser, A., A. K. Lindholm, L. W. Simmons, and R. C. Firman. 2017. Sperm competition

suppresses gene drive among experimentally evolving populations of house mice.

Molecular Ecology 5784–5792.

Meade, L. C., D. Dinneen, R. Kad, D. M. Lynch, K. Fowler, and A. Pomiankowski. 2018.

Ejaculate sperm number compensation in stalk-eyed flies carrying a selfish meiotic

drive element. Heredity.

Paczolt, K. A., J. A. Reinhardt, and G. S. Wilkinson. 2017. Contrasting patterns of X-

chromosome divergence underlie multiple sex-ratio polymorphisms in stalk-eyed flies.

Journal of Evolutionary Biology 30:1772–1784.

Palopoli, M. F., and C. I. Wu. 1996. Rapid evolution of a coadapted gene complex: evidence

from the *Segregation Distorter* (SD) system of meiotic drive in *Drosophila*

melanogaster. Genetics 143:1675–88.

Panhuis, T. M., and G. S. Wilkinson. 1999. Exaggerated male eye span influences contest

outcome in stalk-eyed flies (Diopsidae). Behavioral Ecology and Sociobiology 46:221–

227.

Pinzone, C. A., and K. A. Dyer. 2013. Association of polyandry and sex-ratio drive prevalence

in natural populations of *Drosophila neotestacea*. Proceedings of the Royal Society B:

Biological Sciences 280:20131397.

Pitnick, S., G. T. Miller, J. Reagan, and B. Holland. 2001. Males' evolutionary responses to

experimental removal of sexual selection. Proceedings of the Royal Society B: Biological

Sciences 268:1071–1080.

Presgraves, D. C., E. Severance, and G. S. Wilkinson. 1997. Sex chromosome meiotic drive in

stalk-eyed flies. Genetics 147:1169–80.

Price, T. A. R., A. J. Bretman, T. D. Avent, R. R. Snook, G. D. D. Hurst, and N. Wedell. 2008*a*.

Sex ratio distorter reduces sperm competitive ability in an insect. Evolution 62:1644–

1652.

Price, T. A. R., A. J. Bretman, A. C. Gradilla, J. Reger, M. L. Taylor, P. Giraldo-Perez, A.

Campbell, et al. 2014. Does polyandry control population sex ratio via regulation of a

selfish gene? Proceedings of the Royal Society B: Biological Sciences 281:20133259.

Price, T. A. R., D. J. Hodgson, Z. Lewis, G. D. D. Hurst, and N. Wedell. 2008*b*. Selfish genetic

elements promote polyandry in a fly. Science 322:1241–1243.

Price, T. A. R., R. C. Hoskyns, H. Rapley, J. C. Evans, and N. Wedell. 2012. No evidence that

temperature-related fertility differences influence the distribution of a selfish genetic

element. Functional Ecology 26:657–665.

Price, T. A. R., and N. Wedell. 2008. Selfish genetic elements and sexual selection: Their

impact on male fertility. Genetica 134:99–111.

Reinhold, K., L. Engqvist, B. Misof, and J. Kurtz. 1999. Meiotic drive and evolution of female

choice. Proceedings of the Royal Society B: Biological Sciences 266:1341–1345.

Rogers, D. W., R. H. Baker, T. Chapman, M. Denniff, A. Pomiankowski, and K. Fowler. 2005*a*.

Direct and correlated responses to artificial selection on male mating frequency in the

stalk-eyed fly *Cyrtodiopsis dalmanni*. Journal of Evolutionary Biology 18:642–650.

Rogers, D. W., T. Chapman, K. Fowler, and A. Pomiankowski. 2005*b*. Mating-induced

reduction in accessory reproductive organ size in the stalk-eyed fly *Cyrtodiopsis*

dalmanni. BMC Evolutionary Biology 5:37.

Rogers, D. W., M. Denniff, T. Chapman, K. Fowler, and A. Pomiankowski. 2008. Male sexual

ornament size is positively associated with reproductive morphology and enhanced

fertility in the stalk-eyed fly Teleopsis dalmanni. BMC Evolutionary Biology 8.

Rydzewski, W. T., S. A. Carioscia, G. Liévano, V. D. Lynch, and M. M. Patten. 2016. Sexual

antagonism and meiotic drive cause stable linkage disequilibrium and favour reduced

recombination on the X chromosome. Journal of Evolutionary Biology 29:1247–1256.

Silver, L. M. 1985. Mouse *t* haplotypes. Annual Review of Genetics 19:179–208.

Simmons, L. W., and F. García-González. 2008. Evolutionary reduction in testes size and

competitive fertilization success in response to the experimental removal of sexual

selection in dung beetles. Evolution 62:2580–2591.

Small, J., S. Cotton, K. Fowler, and A. Pomiankowski. 2009. Male eyespan and resource

ownership affect contest outcome in the stalk-eyed fly, *Teleopsis dalmanni*. Animal

Behaviour 78:1213–1220.

Swallow, J. G., L. E. Wallace, S. J. Christianson, P. M. Johns, and G. S. Wilkinson. 2005.

Genetic divergence does not predict change in ornament expression among

populations of stalk-eyed flies. Molecular Ecology 14:3787–3800.

Tao, Y., J. P. Masly, L. Araripe, Y. Ke, and D. L. Hartl. 2007. A *sex-ratio* meiotic drive system in

Drosophila simulans. I: An autosomal suppressor. PLoS Biology 5:e292.

Taylor, D. R., and P. K. Ingvarsson. 2003. Common features of segregation distortion in

plants and animals. Genetica 117:27–35.

Taylor, D. R., M. J. Saur, and E. Adams. 1999. Pollen performance and sex-ratio evolution in a

dioecious plant. Evolution 53:1028–1036.

Taylor, J. E., and J. Jaenike. 2002. Sperm competition and the dynamics of X chromosome

drive: Stability and extinction. Genetics 160:1721–1731.

Wedell, N. 2013. The dynamic relationship between polyandry and selfish genetic elements.

Philosophical Transactions of the Royal Society B: Biological Sciences 368:20120049.

Werren, J. H. 2011. Selfish genetic elements, genetic conflict, and evolutionary innovation.

Proceedings of the National Academy of Sciences of the United States of America

108:10863–10870.

Wilkinson, G. S., P. M. Johns, E. S. Kelleher, M. L. Muscedere, and A. Lorsong. 2006. Fitness

effects of X chromosome drive in the stalk-eyed fly, *Cyrtodiopsis dalmanni*. Journal of

Evolutionary Biology 19:1851–1860.

Wilkinson, G. S., H. Kahler, and R. H. Baker. 1998*a*. Evolution of female mating preferences

in stalk-eyed flies. Behavioral Ecology 9:525–533.

Wilkinson, G. S., D. C. Presgraves, and L. Crymes. 1998*b*. Male eye span in stalk-eyed flies

indicates genetic quality by meiotic drive suppression. Nature 391:276–279.

Wilkinson, G. S., and P. R. Reillo. 1994. Female choice response to artificial selection on an

exaggerated male trait in a stalk-eyed fly. Proceedings of the Royal Society B: Biological

Sciences 255:1–6.

Wilkinson, G. S., and M. I. Sanchez. 2001. Sperm development, age and sex chromosome

485  meiotic drive in the stalk-eyed fly, *Cyrtodiopsis whitei*. Heredity 87:17–24.

486 Wilkinson, G. S., J. G. Swallow, S. J. Christianson, and K. Madden. 2003. Phylogeography of

487  sex ratio and multiple mating in stalk-eyed flies from southeast Asia. Genetica 117:37–

488  46.

489 Wolfenbarger, L. L., and G. S. Wilkinson. 2001. Sex-linked expression of a sexually selected

490  trait in the stalk-eyed fly, *Cyrtodiopsis dalmanni*. Evolution 55:103–110.

491 Zeh, J. A., and D. W. Zeh. 1997. The evolution of polyandry II: Post-copulatory defenses

492  against genetic incompatibility. Proceedings of the Royal Society B: Biological Sciences

493  264:69–75.

Figure 1

A, testes (T) and accessory glands (Ag) after dissection. *B–D*, male morphological and

reproductive trait size for SR (red) and ST (blue) males, plotted against male body size. *B,*

male eyespan. *C*, male testis area. *D*, male accessory gland area. SR males have smaller body

size, eyespan and accessory gland size, but larger testis size. Grey shading shows ± s.e. *E*,

mating frequency, measured as total number of copulations (mean ± s.e.) observed over 30

minutes.

Figure 2

A–B, upper: box plots (median and interquartile range) and *lower:* Kernel probability density

of measures of total fertility of SR (red) and ST (blue) males. *A,* mated to a single female. *B,*

mated to five females. Across both mating regimes, SR and ST males did not differ in the

number of eggs fertilised. *C–D,* absolute testis area plotted against total fertility. *C,* mated to

a single female. *D,* mated to five females. Across both mating regimes, total fertility

increased with testis area in SR and ST males. Grey shading shows ± s.e.