1 ***C. elegans* transgenerationally adapts to infection by increasing the expression of**
2 **cysteine synthases in progeny**
3
4

5 Nicholas O. Burton[1,*], Cristian Riccio[2,3], Alexandra Dallaire[2,4], Jon Price[2,4], Benjamin
6 Jenkins[5], Albert Koulman[5], Eric A. Miska[2,3,4]
7
8 [1] Centre for Trophoblast Research, Department of Physiology, Development and
9 Neuroscience, University of Cambridge, Cambridge, CB2 3EG, UK
10
11 [2]Gurdon Institute, University of Cambridge, Cambridge, CB2 1QN, UK
12
13 [3]Wellcome Sanger Institute, Wellcome Genome Campus, Cambridge CB10 1SA, UK
14
15 [4]Department of Genetics, University of Cambridge, Downing Street, Cambridge CB2 3EH,
16 UK
17
18 [5] Metabolic Research Laboratories, University of Cambridge, Cambridge, CB2 0QQ, UK
19
20
21 *Correspondence: nob20@cam.ac.uk

22
23
24
25
26
27
28
29
30
31

32 Keywords
33 *C. elegans*, transgenerational inheritance, infection, pathogen, heritable, *rhy-1, cysl-1, cysl-2,*
34 *wdr-23*

35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50

Abstract

Parental exposure to pathogens can prime offspring immunity in diverse organisms. The

mechanisms by which parental exposure to pathogens can prime offspring immunity remain

largely unknown. Here we report that the soil bacteria *Pseudomonas vranovensis* is a natural

pathogen of the nematode *Caenorhabditis elegans* and that parental exposure of animals to *P.*

vranovensis promotes offspring resistance to infection. This adaptation can be transmitted

transgenerationally such that infection of adults can enhance the immunity of their

descendants four generations later. We find that parental infection by *P. vranovensis* results

in increased expression of the cysteine synthases CYSL-1 and CYSL-2 and the regulator of

hypoxia inducible factor RHY-1 in progeny, that the expression of these three genes in

offspring is required for adaptation to *P. vranovensis,* and that the expression of these genes

is regulated by the WD40 repeat protein WDR-23.

Introduction

Intergenerational and transgenerational responses to environmental stress have been reported in evolutionarily diverse organisms[1–7]. These studies investigated a variety of types of stresses ranging from osmotic stress[1], to mitochondrial stress[5], to pathogen infection[8]. In each case, parental exposure to stress appeared to prime offspring to respond to a similar stress. For example, parental exposure of the nematode *Caenorhabditis elegans* to the opportunistic pathogen *Pseudomonas aeruginosa* was reported to alter offspring behaviour in a way that promotes offspring avoidance of *P. aeruginosa* and enhances offspring survival[8]. Despite this apparent adaptation, parental exposure of *C. elegans* to *P. aeruginosa* also has been reported to have deleterious effects; parental exposure to *P. aeruginosa* results in offspring that are more susceptible to osmotic stress[1]. Taken together, these observations suggest that parents might prime their offspring to adapt to stresses they have faced at the expense of offspring's ability to respond to other stresses. However, whether these observations represent rare examples and potential laboratory artifacts or if such programmed predictive adaptive responses might be ubiquitous throughout evolution remains unclear.

Among the diverse stresses an organism might encounter, pathogens such as viruses, some bacteria and eukaryotes are among the most ubiquitous stresses found in nature[9]. To adapt to the constant threat of pathogens, many organisms have evolved mechanisms by which parental exposure to pathogens can prime offspring immunity[6,10–15]. For example, in mammals, a mother can transfer specific antibodies to her offspring via milk to prime offspring immunity[15]. Similar observations of parents priming offspring immunity in response to pathogens have been reported in both plants[14] and invertebrates[10], even though

these organisms lack antibodies. These findings suggest that multiple independent

mechanisms have evolved for parents to prime offspring immunity.

The mechanisms by which parental exposure to pathogens might program predictive adaptive

responses in offspring in organisms that lack antibodies remain largely unknown. Here we

report that *C. elegans* can transgenerationaly adapt to a natural pathogen, *Pseudomonas*

vranovensis. In this case, we find that parental exposure to *P. vranovensis* results in an

increase in the expression of cysteine synthases in offspring which promotes resistance to *P.*

vranovensis. This mechanism is distinct from any previously proposed pathway[8].

Collectively, these observations describe a new mechanism by which parents can prime

offspring immunity in response to a bacterial infection.

<u>Results</u>

One of the major obstacles in determining the molecular mechanisms underlying heritable

adaptations to pathogen infection is the lack of robust models that can be easily analyzed in

the laboratory. To address this obstacle, we sought to develop a model of a heritable

adaptation to bacterial infection in the nematode *C. elegans* by testing whether *C. elegans*

could heritably adapt to potential bacterial pathogens from its natural environment. Previous

sampling of bacterial species from the natural environments of *C. elegans* identified 49 as yet

undescribed bacterial isolates that induce the expression of the immune response genes *irg-1*

or *irg-5*[16]. We tested whether parental exposure to these potentially pathogenic bacteria could

enhance offspring survival in response to future exposure to the same bacteria. We found that

parental exposure to two of the isolates, BIGb446 and BIGb468, significantly enhanced

offspring survival in response to future exposure to the same bacteria (Fig. 1a). Specifically,

approximately 95% of newly hatched larvae from parents fed a standard laboratory diet of *E.*

coli HB101 died within 24 hours of hatching on BIGb446 or BIGb468 (Fig. 1a). By contrast,

45% of embryos from adults briefly exposed to BIGb446 or BIGb468 were still alive at 24

hours (Fig. 1a) and a majority of these larvae survived through adulthood (Fig. 1b-c). In

addition, we found that adults were more resistant to BIGb446 than larval animals

(Supplementary Fig. 1) and that UV-killed bacteria did not cause lethality (Fig. 1d). We

conclude that parental exposure to bacterial isolates BIGb446 and BIGb468 enhances

offspring survival in response to future exposure to these bacteria.

To determine the species identity of bacterial isolates BIGb446 and BIGb468 we performed

long read whole genome sequencing and assembled the genomes of these bacteria. We found

that the genomes of both BIGb446 and BIGb468 were approximately 5.9 Mb (Supplementary

File 1 and 2) and were 99.38% identical across the entire genome (Supplementary Table 1

and Supplementary Fig. 2). We concluded that BIGb446 and BIGb468 are isolates of the

same species. We compared the 16S rRNA sequence of BIGb446 to known bacterial

genomes using BLAST. We found that the 16s rRNA sequence from BIGb446 is 99.93%

identical to *Pseudomonas vranovensis* strain 15D11. Previous studies of *Pseudomonas*

phylogeny have used the DNA sequences of *gyrB*, *rpoB*, and *recA* to differentiate species of

Pseudomonas, with similarity above 97% set as the accepted species threshold[17,18]. We found

that the sequence of *gyrB* was 98.05% identical, the sequence of *rpoB* was 99.44% identical,

and the sequence of *recA* was 98.66% identical to the sequences of homologous genes in *P.*

vranovensis strain 15D11. No other species of *Pseudomonas* was greater than 97% identical

to *Pseudomonas sp.* BIGb446 at any of these three genes. Furthermore, we compared the

average nucleotide identity (ANI) of our assembly of *Pseudomonas sp.* BIGb446 with the

genome of *P. vranovensis* strain 15D11 using OrthoANIu[19]. We found these two genomes

had an average nucleotide identity of 97.33%. We conclude that *Pseudomonas sp.* BIGb446

and *Pseudomonas sp.* BIGb468 are isolates of *Pseudomonas vranovensis*, a Gram-negative

soil bacteria[20].

In some cases, the effects of parental stress on offspring have been reported to be

intergenerational and only last a single generation[1]. In other cases, the effects of parental

stress on offspring have been reported to persist transgenerationally and thus affect

descendants several generations later[5,8,21]. To test whether exposure to BIGb446 had any

transgenerational effects on immunity we fed adult animals BIGb446 for 24 hours and

assayed the response of progeny two generations later. We found that a single exposure of

adult animals to BIGb446 enhanced their offspring's survival (one generation later) but did

not enhance the survival of their descendants two generations later (Fig. 2a), indicating that

the effects of parental exposure to BIGb446 only last a single generation under these

conditions. Some studies of the effects of parental environment on offspring have found that

consecutive exposure of multiple generations of animals to the same stress can enhance

adaptation to stress in descendants[5]. We therefore tested whether the exposure of five

consecutive generations (P0, F1, F2, F3, and F4) of animals to BIGb446 could enhance

progeny (F5) survival in response to BIGb446 but found that this was not the case (Fig. 2a).

Finally, we tested whether the exposure of four consecutive generations of animals to

BIGb446 could enhance survival in progeny for more than a single generation. Unlike a

single generation of exposure to BIGb446, we found that the enhanced survival of

descendants of animals exposed to BIGb446 for four consecutive generations persisted for

two generations (Fig. 2a). Furthermore, we found that three consecutive generations of

exposure to BIGb446 was required for resistance to BIGb446 to persist transgenerationally

(Fig. 2b). We conclude that under conditions of multiple consecutive generations of animals

being exposed to BIGb446 the effects of exposure to BIGb446 can be transmitted

transgenerationally.

Multiple signaling pathways and epigenetic modifiers have been proposed to mediate

heritable and transgenerational responses to environmental stress. We tested whether

mutations in any factors previously reported to be involved heritable or transgenerational

responses to stress were required for *C. elegans* heritable adaptation to BIGb446 including

small RNA mediated pathways (*hrde-1*[22], *prg-1*[8]), H3K9 methylation (*set-32*[23] and *met-2*[4]),

H3K4 methylations (*set-2, wdr-5.1, spr-5*)[5,21], DNA adenosine methylation (*damt-1*)[5], and the

RAS/ERK signaling pathway (*lin-45*)[1]. We found that none of these factors were required for

C. elegans adaptation to BIGb446 (Fig. 2c) These data suggest that adaptation to *P.*

vranovensis involves an as-yet-unknown mechanism of inheritance.

Bacterial isolates BIGb446 and BIGb468 were originally reported to promote the expression

of immune response gene *irg-5*[16]. The expression of *irg-5* in response to pathogen infection is

controlled by the p38-like MAP kinase PMK-1[24]. We tested whether PMK-1 is required for

resistance to BIGb446 by placing wild-type and *pmk-1* mutant adults on plates seeded with

BIGb446. We found that 100% of *pmk-1* mutants were dead after 48 hours while more than

40% of wild-type animals remained alive (Supplementary Fig. 1). These results indicate that

PMK-1 promotes resistance to BIGb446. We tested whether PMK-1 was required for

heritable adaptation to BIGb446 by exposing wild-type and *pmk-1* mutants to BIGb446 for

24 hours and assaying the survival of their offspring in response to repeated exposure to

BIGb446. We found that PMK-1 is not required for the heritable adaptation to BIGb446 (Fig.

2d).

C. elegans' heritable adaptation to *P. vranovensis* might be a general response to pathogen

infection or specific to *P. vranovensis.* To test whether parental infection by different

bacterial pathogens could protect offspring from BIGb446 we exposed adult animals to two

additional bacterial pathogens of *C. elegans*, *Pseudomonas aeruginosa* PA14[25] and

Photorhabdus luminescens Hb[26], and assayed the response of their offspring to BIGb446.

We found that parental infection by these two pathogens did not enhance offspring survival

in response to BIGb446 (Fig. 2e). These results indicate that adaptation to *P. vranovensis* is

not a generic response to pathogenic bacteria.

Previous studies found that *C. elegans* can heritably adapt to osmotic stress and starvation by

regulating gene expression and metabolism in offspring[1,3]. To identify how parental infection

by BIGb446 affects gene expression and metabolism in offspring we profiled gene

expression by RNA-seq and metabolite levels by LC/MS. Using RNA-seq we identified

1,153 genes that exhibit increased expression greater than 2-fold and 491 genes that

decreased expression greater than 2-fold in embryos from parents fed BIGb446 when

compared to embryos from parents fed HB101 (Fig. 3a and Supplementary Table 2). We also

quantified gene expression in young adults exposed to BIGb446. We found that 1,275 genes

exhibited a greater than 2-fold change in expression when compared to young adults fed

HB101 (Supplementary Table 2). Of these 1,275 genes, 398 were also observed to change in

embryos (Fig. 3b). We conclude that infection by *P. vranovensis* BIGb446 results in similar,

but distinct, effects on adult and embryonic gene expression. Using LC/MS we profiled 92

lipid metabolites, including those previously observed to heritably change in abundance in

response to osmotic stress[27], and found that 6 metabolites exhibit modest changes in

abundance in embryos from parents fed BIGb446 when compared to embryos from parents

fed HB101 (Fig. 3c and Supplementary Table 3). We conclude that parental exposure to the

pathogenic diet of *P. vranovensis* BIGb446 alters gene expression and metabolite abundance

in embryos.

We previously found that parental infection by *P. aeruginosa* PA14 and *P. luminescens* Hb

did not protect offspring from BIGb446 (Fig. 2e). We hypothesized that different parental

infections might result in different gene expression and metabolite changes in embryos and

that the specific changes caused by parental infection by BIGb446 are required for offspring

adaptation to BIGb446. To compare how parental infection by different bacterial pathogens

affects offspring gene expression and metabolism we exposed young adults to *P. aeruginosa*

PA14 or *P. luminescens* Hb for 24 hours and collected embryos from these animals to profile

gene expression by RNA-seq and metabolites by LC/MS. We found that only 41 genes

exhibited a greater than 2-fold change in expression in embryos from parents exposed to

PA14 when compared to embryos from parents exposed to HB101 and of these genes 29 also

exhibited altered gene expression in embryos from parents exposed to BIGb446 (Fig. 3d-e,

Supplementary Table 4). Separately, we found that only 26 genes exhibited a greater than 2-

fold change in expression in embryos from parents exposed to *P. luminescens Hb* when

compared to parents fed HB101 (Fig. 3d-e and Supplementary Table 4). Of these genes, only

5 also exhibited altered expression in embryos from parents exposed to BIGb446 (Fig. 3d).

Collectively, these results indicate that parental infection by *P. aeruginosa* and *P.*

luminescens had only a limited effect on offspring gene expression when compared to

parental infection by *P. vranovensis*. Only a single lipid metabolite measured exhibited a

statistically significant change in abundance in embryos from parents fed *P. aeruginosa* and

P. luminescens (Supplementary Table 3). We conclude that parental exposure to different

pathogenic diets has distinct effects on offspring gene expression.

We hypothesized that the specific changes in gene expression in embryos from parents

exposed to *P. vranovensis* BIGb446 mediate the observed adaptation to this pathogen.

Among the genes that we found to specifically increase in expression in response to BIGb446

was the glutathione-s-transferase *gst-31* (among the top 10 most upregulated genes (Table

S2)) and we confirmed that *C. elegans'* exposure to BIGb446 resulted in increased

expression of *gst-31* in both adults and embryos using a GFP reporter (Fig. 4a-b). To identify

factors that regulate animals' gene expression in response to BIGb446 we performed a

chemical mutagenesis screen for mutants that exhibit increased expression of *gst-31* even

when fed a normal laboratory diet of *E. coli* HB101. We identified four mutants from this

screen and all four contain mutations in the conserved WD40 repeat protein *wdr-23* (Fig. 4c).

Consistent with our findings, multiple previous studies have identified that *wdr-23* mutants

exhibit increased expression of glutathione-s-transferases, including *gst-31*[28–30]. We conclude

that WDR-23 negatively regulates the expression of *gst-31*.

We tested whether the loss of WDR-23 affected *C. elegans* adaptation to *P. vranovensis* and

found that multiple independent *wdr-23* mutants were unable to heritably adapt to BIGb446

(Fig. 4d). These results indicate that loss of WDR-23 prevents animals from being able to

adapt to BIGb446. WDR-23 is a negative regulator of the Nrf2-like transcription factor SKN-

1 and the loss of WDR-23 results in SKN-1 activation[31]. Activation of SKN-1 in turn

promotes the expression of oxidative stress response genes[30] and mutations in *skn-1* suppress

many of the defects observed in *wdr-23* mutants[31]. We exposed wild-type, *skn-1* partial loss-

of-function*, and *wdr-23;skn-1* double mutant adults to BIGb446 and assayed the response of

their offspring to BIGb446. We found that partial loss-of-function mutations in *skn-1* (null

mutants are embryonic lethal) also reduced animals' ability to adapt to BIGb446 and that this

mutation in *skn-1* did not suppress the adaptation defect of *wdr-23* mutants (Fig. 4d).

Collectively, these results indicate that mutations that both increase and decrease the activity

of SKN-1 disrupt animals' ability to adapt to *P. vranovensis*. We propose that proper

signaling via WDR-23 and SKN-1 is critical for *C. elegans* to adapt to *P. vranovensis*.

We previously observed that parental exposure to BIGb446 modifies gene expression in

offspring. We hypothesized that parental infection by BIGb446 might regulate the expression

of pathogen or stress response genes in offspring via WDR-23 and SKN-1. To test this

hypothesis, we first performed RNA-seq of wild-type and *wdr-23* mutant embryos. We

identified that 1,634 genes exhibited a greater than 2-fold change in expression in *wdr-23*

mutants (Fig. 4e and Supplementary Table 5). Of these genes, we found that 634 also exhibit

a greater than 2-fold change in expression in embryos from parents exposed to BIGb446 (Fig.

4e and Supplementary Table 5). Despite regulating an overlapping set of genes, we found

that parental exposure to BIGb446 and the loss of WDR-23 resulted in distinct gene

expression patterns (Fig. 4f). These data indicate that exposure to *P. vranovensis* and

signaling via WDR-23 regulate the expression of a significantly overlapping set of target

genes and are consistent with our hypothesis that parental exposure to BIGb446 modifies

gene expression in offspring in part via WDR-23.

94% of genes that exhibit a greater than 2-fold change in expression in response to both *P.*

vranovensis and loss of WDR-23 exhibit an increase in mRNA expression (Supplementary

Tables 2 and 5). Among these genes we identified the cysteine synthases CYSL-1 and CYSL-

2 (Fig. 5a and Supplementary Tables 2 and 5), which were previously reported to be involved

in breaking down bacterial toxins produced by *P. aeruginosa*[32]. We confirmed that exposure

of parents to BIGb446 increases CYSL-2 expression in embryos using a GFP reporter (Fig.

5b)[33]. We hypothesized that increased expression of *cysl-1* and *cysl-2* in offspring promotes

adaptation to *P. vranovensis*. We assayed seven independent alleles of *cysl-1* and two

independent mutants lacking *cysl-2* and found that these mutants were unable to adapt to

BIGb446 (Fig. 5c-d and Supplementary Fig. 3). In addition, we found that these defects could

be rescued by expressing a wild-type copy of *cysl-2* (Fig. 5d). We conclude that CYSL-1 and

CYSL-2 are required for adaptation to BIGb446.

CYSL-1 and CYSL-2 were previously found to function in a signaling pathway with the

regulator of hypoxia factor, RHY-1, the EGLN1 homolog EGL-9, and the hypoxia inducible

factor HIF-1 to regulate animals' response to hypoxia [33]. We found that one of these genes,

rhy-1, also exhibited increased expression in both animals exposed to BIGb446 and in *wdr-*

23 mutants (Fig. 6a). We tested whether mutants lacking RHY-1, EGL-9, and HIF-1 also

exhibited altered adaptation to BIGb446. We found that three independent mutations in *rhy-1*

resulted in animals that did not adapt to BIGb446 (Fig. 6b), similar to *cysl-1* and *cysl-2*

mutants. This defect was rescued by expressing a wild-type copy of *rhy-1* (Fig. 6b). By

contrast, we found that the loss of *hif-1* and *egl-9* did not affect adaptation to BIGb446 (Fig.

6c). These results indicate that RHY-1, CYSL-1, and CYSL-2 function separately from EGL-

9 and HIF-1 to promote animals' adaptation to *P. vranovensis*.

<u>Discussion</u>

Collectively, our results demonstrate that *P. vranovensis* is a pathogen of *C. elegans* and that

parental exposure of *C. elegans* to *P. vranovensis* can protect offspring from future infection

by increasing the expression of the cysteine synthases *cysl-1* and *cysl-2* and the regulator of

hypoxia inducible factor *rhy-1* in progeny. The ability of *P. vranovensis* to kill *C. elegans*

larvae within 24 hours and the observation that *C. elegans* activates genes associated with

oxidative stress in response to *P. vranovensis* suggests that *P. vranovensis* produces a toxic

molecule(s) that is lethal to *C. elegans*. The cysteine synthases CYSL-1 and CYSL-2 have

been reported to break down toxic molecules such as hydrogen cyanide and hydrogen sulfide

[32]. Based on these observations, we propose that *C. elegans* can transgenerationally adapt to

infection *by P. vranovensis* by increasing the expression of CYSL-1 and CYSL-2 in offspring

which protects animals by breaking down bacterial toxins. Thus, we demonstrate a new

transgenerationally adaptive mechanism for *C. elegans* to fight a natural pathogen.

The expression of *cysl-1*, *cysl-2* and *rhy-1* is regulated by signaling via WDR-23, however,

we found that mutations that both increase and decrease SKN-1 activity disrupt adaptation to

P. vranovensis. One possible explanation for these results is that specific signaling via WDR-

23 and SKN-1, likely in combination with signaling via other pathogen response pathways, is

required to elicit a distinct response to *P. vranovensis* and that completely abolishing

signalling via this pathway using genetic mutants results in an inappropriate response to *P.*

vranovensis and animals that are unable to adapt. Such a model is similar to other core

signaling pathways that regulate animals' response to multiple different stresses, such as the

insulin-like signaling pathway in *C. elegans* that functions in combination with signaling via

SSU-1 and NHR-1 to specify *C. elegans* distinct responses to both starvation and osmotic

stress[27]. Alternatively, it is possible that WDR-23 and SKN-1 regulate the expression of

CYSL-1, CYSL-2, and RHY-1 in parallel to infection by *P. vraonvensis.* In this case,

disrupted signaling via WDR-23 and SKN-1 might result in animals that are simply more

susceptible to infection and thus unable to adapt. Interestingly, loss of RHY-1 alone also

modifies CYSL-1 activity and results in significantly increased expression of CYSL-2[33]. It is

possible that RHY-1, WDR-23, and SKN-1 all function by regulating the activity of CYSL-1

and CYSL-2 and that the failure of *rhy-1, wdr-23,* and *skn-1* mutants to adapt to *P.*

vraonvensis is due to abnormal regulation of these two cysteine synthases.

A single generation of exposure to *P. vranovensis* only resulted in an intergenerational

adaptation to infection (Figure 2A). By contrast, we found that multiple successive

generations of exposure to *P. vranovensis* could result in transgenerational adaptation to

infection that persisted for two generations, however, the second generation exhibited a

significantly less robust adaptation than the offspring of infected animals (Figure 2A). This

transgenerational effect required 3 consecutive generations of exposure to *P. vranovensis*

(Figure 2B). These observations are consistent with an as yet unknown factor potentially

accumulating in germ cells and embryos and that is diluted out after two generations of

removal from the pathogen as a food source. Such a factor could be an epigenetic mark, a

signalling molecule, the altered abundance of a metabolite, or the presence of a bacterial

toxin. The identification of the molecular identity of this factor will be crucial to determining

the nature of this transgenerational response.

Recent studies of *C. elegans* response to the human opportunistic pathogen *P. aeruginosa*

have also described a transgenerational response to this pathogenic species of *Pseudomonas*[8].

In this paradigm researchers found that exposure of a single generation of *C. elegans* to *P.*

aeruginosa resulted in a heritable change in avoidance behaviour that persisted for four

generations and was dependent on the PIWI Argonaute homolog PRG-1[8]. By contrast, we

found that exposure of a single generation of *C. elegans* to *P. vranovensis* did not result in a

transgenerational adaptation to infection, but rather the heritable increase in immunity only

lasted one generation (Fig. 2a-b). Transgenerational effects of *P. vranovensis* infection were

only observed after several consecutive generations of exposure to *P. vranovensis* (Fig. 2b).

In addition, we found that *C. elegans* heritable adaptation to *P. vranovensis* did not require

PRG-1 (Fig. 2c). The explanation for why *C. elegans* elicits different responses to *P.*

aeruginosa when compared to *P. vranovensis* remains unclear. It is possible that *C. elegans*

evolved distinct responses to *P. vranovensis* when compared to *P. aeruginosa*. Alternatively,

studies of *C. elegans'* heritable responses to *P. aeruginosa* focused on changes in avoidance

behaviour while our studies of *C. elegans* heritable responses to *P. vranovensis* focused on

changes in survival under conditions where animals were unable to avoid the pathogen. It is

possible that heritable changes in behaviour in response to bacterial infection in *C. elegans*

are mediated by a small RNA pathway and PRG-1, while changes in immunity and the

expression of cysteine synthases are controlled by a mechanism that is PRG-1 independent.

Future studies will be important in differentiating between these possibilities and determining

how parental infection can heritably prime offspring immunity, if these mechanisms are

evolutionarily conserved, and how much they contribute to organisms' responses to

pathogens.

Finally, studies of *C. elegans* interaction with the opportunistic pathogen *P. aeruginosa*

indicate that parental exposure to *P. aeruginosa* can have both adaptive[8] and deleterious[1]

consequences for offspring. We suspect that similar trade-offs, where adaptation to one stress

comes at the cost of fitness in a different environment, are also likely to be the case for *C.*

elegans adaptation to *P. vranovensis.* Future studies will likely be critical in determining

what the costs of intergenerational and transgenerational responses to stress are, and we

propose that such studies might be able to identify links between different types of stress

responses. For example, previous studies have observed that *C. elegans* transcriptional

response to osmotic stress and pathogen infection appear to be related[34]. These findings, in

combination with observations that parental infection by *P. aeruginosa* results in offspring

that are more susceptible to osmotic stress[1], suggest that adapting to pathogen infection might

generally disrupt animals' ability to respond to osmotic stress and explain why animals are

not normally resistant to this pathogen.

Acknowledgments: We thank Buck Samuel, Gary Ruvkun, and Jonathan Ewbank for

bacterial isolates; Long Ma, Keith Choe, Derek Sieburth, Bob Horvitz, Na An, and the

Caenorhabditis Genetic Center, which is funded by the NIH National Center for Research

Resources (NCRR), for strains; Martin Welch and George Salmond for use of bacterial

culture equipment and space, and Martin Hemberg for feedback on RNA-seq analysis. N.O.B

is funded by a Next Generation Fellowship from the Centre for Trophoblast Research. C.R.

was supported by a Wellcome PhD programme. A.K. and B.J are funded by BBSRC grant

BB/M027252/1. This work was also supported by Cancer Research UK (C13474/A18583,

C6946/A14492) and the Wellcome Trust (104640/Z/14/Z, 092096/Z/10/Z) grants to E.A.M.

Author Contributions: N.O.B. conceived the project and designed the experiments. N.O.B

and E.A.M analysed the data. N.O.B., C. R., A.D., J.P., A.K., and B.J. performed the

experiments. N.O.B. wrote the manuscript.

Declarations of Interest: The authors declare that they have no competing interests.

Data and materials availability: RNA-seq data that support the findings of this study have

been deposited at the European Nucleotide Archive (ENA) under the accession code

PRJEB32993. The raw data for assembling the genomes of *P. vranovensis* isolate BIGb446 is

available at the ENA under the accession code ERS3670403 and BIGb468 is available under

the accession code ERS3670404. The raw data related to metabolomics is available on Dryad

using doi:10.5061/dryad.8t6q5f5. Raw data supporting all figures is provided in

Supplementary Table 6.

Figure 1. *C. elegans* heritably adapts to infection by *Pseudomonas vranovensis.* (a) Percent of

wild-type animals surviving on plates seeded with either *E. coli* HB101 or bacterial isolates

BIGb446 and BIGb468 after 24 hrs. Error bars, s.d. n = 3 experiments of >100 animals. (b)

Percent of wild-type animals surviving on plates seeded with bacterial isolate BIGb446. Error

bars, s.d. n = 3 experiments of 500 animals. (c) Images of wild-type animals surviving after

120 hrs of feeding on bacterial isolate BIGb446. 1000 animals were used at t = 0 in each

condition and surviving animals were resuspended in 20 µl M9 and imaged. Scale bars 1 mm.

(d) Percent of wild-type animals surviving on *E. coli* HB101 or bacterial isolate BIGb446

after 24 hrs. Error bars, s.d. n = 3 experiments of >100 animals. * = $p < 0.05$, ** = $p < 0.01$,

*** = $p < 0.001$, **** $p < 0.0001$.

Figure 2. *C. elegans* adaptation to *P. vranovensis* can be inherited transgenerationally. (a)

Percent of wild-type animals surviving on plates seeded with bacterial isolate BIGb446 after

24 hrs. Error bars s.d. $n = 3$ experiments of > 100 animals. (b) Percent of wild-type animals

surviving on plates seeded with bacterial isolate BIGb446 after 24 hrs. Error bars s.d. $n = 3$

experiments of 500 animals. Wild-type animals carried the integrated transgene *nIs470*. (c)

Percent of wild-type, *set-32(ok1457), prg-1(n4357), hrde-1(tm1200), damt-1(gk961032),*

wdr-5.1(ok1417), set-2(ok952), spr-5(by134), lin-45(n2018), and *met-2(n4256)* mutants

surviving on plates seeded with bacterial isolates BIGb446 after 24 hrs. Error bars, s.d. $n = 3$

experiments of >100 animals. (d) Percent of wild-type and *pmk-1(km25)* mutants surviving

on plates seeded with bacterial isolates BIGb446 after 24 hrs. Error bars, s.d. $n = 3$

experiments of >100 animals. (e) Percent of wild-type animals surviving on plates seeded

with bacterial isolate BIGb446 after 24 hrs. Error bars s.d. $n = 3$ experiments of 500 animals.

*** $= p < 0.001$, **** $p < 0.0001$.

Figure 3. Parental infection by *P. vranovensis* alters gene expression and metabolism in

offspring. (a) Gene expression changes in embryos from parents fed *P. vranovensis* BIGb446

when compared to embryos from parents fed *E. coli* HB101. Values represent averages from

three replicates. (b) Venn diagram of genes exhibiting a greater than 2-fold change in RNA

expression in adults fed *P. vranovensis* BIGb446 and embryos from adults fed *P. vranovensis*

BIGb446. *p*-value represents normal approximation to the hypergeometric probability (See

Statistics and Reproducibility) (c) μM abundance of lipid metabolites exhibiting a greater

than 2-fold change in abundance in embryos from parents fed *P. vranovensis* BIGb446 when

compared to embryos from parents fed *E. coli* HB101. Error bars, s.d. n = 3 replicates. (d)

Venn diagram of genes exhibiting a greater than 2-fold change in RNA expression in

embryos from parents fed *P. vranovensis* BIGb446, *P. aeruginosa* PA14, and *P. luminescens*

Hb. *p*-value represents normal approximation to the hypergeometric probability (See

Statistics and Reproducibility) (e) Principal component analysis (PCA) plot of mRNA

expression data from RNA-seq of embryos from parents fed *E. coli* HB101, *P. vranovensis*

BIGb446 and BIGb468, *P. aeruginosa* PA14, or *P. luminescens* Hb.

Figure 4. Signaling via WDR-23 regulates *C. elegans* response to *P. vranovensis.* (a)

Representative images of *gst-31::GFP* expression in wild-type embryos from parents fed

either *E. coli* HB101 or *P. vranovensis* BIGb446. Scale bars 10 μm. (b) Representative

images of *gst-31::GFP* expression in wild-type adults fed either *E. coli* HB101 or *P.*

vranovensis BIGb446 for 24 hrs. Scale bars 100 μm. (c) Schematic of *wdr-23* mutations that

cause increased *gst-31::GFP* expression when animals are fed *E. coli* HB101. Scale bar, 100

bp. (d) Percent of wild-type, *wdr-23(mac51, tm1817),* and *skn-1(zj15)* mutants surviving on

plates seeded with bacterial isolates BIGb446 after 24 hrs. Error bars, s.d. n = 3 experiments

of >100 animals. **** $p < 0.0001$. (e) Venn diagram of genes exhibiting a greater than 2-fold

change in RNA expression in embryos from parents fed *P. vranovensis* BIGb446 and *wdr-*

23(bur9) mutant embryos from parents fed *E. coli* HB101. *p*-value represents normal

approximation to the hypergeometric probability (See Statistics and Reproducibility). (f)

Principal component analysis (PCA) plot of mRNA expression data from RNA-seq of wild-

type and *wdr-23(bur9)* mutant embryos from parents fed *E. coli* HB101, *P. vranovensis*

BIGb446, *P. aeruginosa* PA14, or *P. luminescens* Hb.

A

Parental diet

B

gst-31::GFP

C

D

E

Genes exhibiting greater than 2-fold change in expression

$p < 1.0 \, e^{-100}$

F

Figure 5. CYSL-1 and CYSL-2 are required for *C. elegans* to adapt to *P. vranovensis.* (a)

Transcripts per million (TPM) of *cysl-2* in wild-type and *wdr-23(bur9)* mutant embryos from

parents fed *E. coli* HB101 or *P. vraonvensis* BIGb446. Error bars, s.d. n = 3 replicates. Data

is the same as found in Table S2 and S5. (b) Representative images of *cysl-2::GFP* in

embryos from parents fed *E. coli* HB101 or *P. vraonvensis* BIGb446. Scale bars 100 μm. (c)

Percent of wild-type and *cysl-1(ok762)* mutants surviving on plates seeded with bacterial

isolates BIGb446 after 24 hrs. Error bars, s.d. n = 3 experiments of >100 animals. (d) Percent

of wild-type and *cysl-2(ok3516)* mutants surviving on plates seeded with bacterial isolates

BIGb446 after 24 hrs. Error bars, s.d. n = 3 experiments of >100 animals. *** = $p < 0.001$,

**** $p < 0.0001$.

Figure 6. RHY-1 is required for *C. elegans* adaptation to *P. vranovensis*. (a) Transcripts per

million (TPM) of *rhy-1* in wild-type and *wdr-23(bur9)* mutant embryos from parents fed *E.*

coli HB101 or *P. vraonvensis* BIGb446. Error bars, s.d. n = 3 replicates. Data is the same as

found in Table S2 and S5. (b) Percent of wild-type and *rhy-1* mutants surviving on plates

seeded with bacterial isolates BIGb446 after 24 hrs. Error bars, s.d. n = 3 experiments of

>100 animals. (c) Percent of wild-type, *egl-9(n586), hif-1(ia4),* and *rhy-1(n5500)* mutants

surviving on plates seeded with bacterial isolates BIGb446 after 24 hrs. Error bars, s.d. n = 3

experiments of >100 animals. The wild-type data in panels B and C and Figure 5C are

identical as these experiments were run together. *** = p < 0.001, **** p < 0.0001.

Methods

Strains. *C. elegans* strains were cultured as described[35] and maintained at 20 °C unless noted

otherwise. The Bristol strain N2 was the wild-type strain.

LGI: *set-32(ok1457), prg-1(n4357), spr-5(by134), wdr-23(bur7, bur8, bur9, bur10, tm1817,*

   mac51)

LGII: *cysl-2(ok3516, syb1431), damt-1(gk961032), rhy-1(n5500, ok1402, tm12398))*

LGIII: *wdr-5.1(ok1417), hrde-1(tm1200), set-2(ok952), met-2(n4256)*

LGIV: *pmk-1(km25), skn-1(zj15), pgl-1(bn102), lin-45(n2018), nIs470[cysl-2::Venus; myo-*

   2::RFP]

LGV: *egl-9(n586), hif-1(ia4)*

LGX: *cysl-1(ok762, mr23, mr25, mr26, mr29, mr39, mr40)*

Unknown linkage: *burIs2[gst-31::gfp]*

Extrachromosomal arrays: *nEx1763[rhy-1(+); myo-2::RFP], burEx1 [cysl-2(+); myo-*

   3::RFP]

Sequencing of Pseudomonas vranovensis *BIGb446 and BIGb468*

Genomic DNA was prepped using a Gentra Puregene kit (QIAGEN). DNA was sheared to 10

kb using gTUBE (Covaris). Sheared DNA was barcoded and multiplexed for PacBio

sequencing using Template Prep Kit 1.0-SPv3 and Barcoded Adapter Kit 8A (PacBio).

Genomic DNA was sequenced using the PacBio Sequel system using version 3.0 sequencing

reagents (PacBio) and 1M v3 SMRT cell.

Genome assembly of BIGb446 and BIGb468

The genomes of BIGb446 and BIGb46 were assembled using HGAP4 from SMRT Link

version 5.1.0.26412 with estimated genome sizes of 5 MB.

Assays of adaptation to Pseudomonas vranovensis *BIGb446 and BIGb468*

P. vranovensis BIGb446 and BIGb468 was cultured in LB at 37 °C overnight. 1 mL of

overnight culture was seeded onto 50 mm NGM agar plates and dried in a laminar flow hood

(bacterial lawns completely covered the plate such that animals could not avoid the

pathogen). All plates seeded with BIGb446 or BIGb468 were used the same day they were

seeded. Young adult animals were placed onto 50 mm NGM agar plates seeded with 1 mL

either *E. coli* HB101 or *P. vranovensis* BIGb446 or BIGb468 for 24 hours at room

temperature. Embryos from these animals were collected and placed onto fresh NGM agar

plates seeded with BIGb446 or BIGb468. Percent surviving were counted after 24 hours at

room temperature unless otherwise noted.

Transgenerational adaptation to P. vranovensis

P. vranovensis BIGb446 was cultured in LB at 37 °C overnight. 1 mL of overnight culture

was seeded onto 50 mm NGM agar plates and dried in a laminar flow hood (bacterial lawns

completely covered the plate). All plates seeded with BIGb446 were used the same day they

were seeded. All animals in each generation were grown from embryos to young adults on

NGM agar plates seeded with *E. coli* HB101. Young adults were then moved to fresh plates

seeded with BIGb446 at room temperature for 24 hours. 24 hours of feeding on BIGb446 was

counted as one generation of exposure to a BIGb446 diet. Embryos from parents fed

BIGb446 were collected and placed onto plates seeded with *E. coli* HB101 until they were

young adults and then moved to fresh plates seeded with BIGb446 for additional generations.

Imaging of wild-type animals surviving after P. vranovensis *exposure*

1,000 embryos were placed onto NGM agar plates seeded with *P. vranovensis* BIGb446 at t =

0 in each condition and surviving animals at 120 hours were washed off of plates and

resuspended in 20 μl M9 and imaged.

Assays of C. elegans response to P. aeruginosa *and* P. luminescens

Young adult animals were placed onto slow-killing assay (*P. aeruginosa*) plates or NGM

agar (*P. luminescens*) plates seeded with either *P. aeruginosa* PA14 or *P. luminescens* Hb for

24 hours at room temperature. Embryos from these animals were collected and snap frozen in

liquid nitrogen for RNA sequencing and metabolomics analysis or placed onto fresh NGM

agar plates seeded with BIGb446. Percent of animals placed on BIGb446 surviving were

counted after 24 hours at room temperature.

Assay of adult survival

Greater than 100 young adult animals were placed onto placed onto NGM agar plates seeded

with *P. vranovensis* BIGb446 from overnight cultures in LB. Percent of animals surviving

was counted at 24 hour intervals. Animals were scored as alive if they were mobile and dead

if they were immobile and did not respond to touch.

RNA-seq

Young adult animals were placed onto NGM agar plates seeded with *E. coli* HB101, *P.*

vranovensis BIGb446 or BIGb468, or *P. luminescens* Hb or slow-killing assay plates seeded

with *P. aeruginosa* PA14 for 24 hours at room temperature. Adult animals or embryos

collected from adult animals after 24 hours were snap frozen in liquid nitrogen. Samples were

lysed using a BeadBug microtube homogenizer (Sigma) and 0.5 mm Zirconium beads

(Sigma). RNA was extracted using a RNeasy Plus Mini kit (Qiagen). mRNA was enriched

using an NEBNext rRNA Depletion kit (NEB). Libraries for sequencing were prepped using

an NEBNext Ultra II Library prep kit for Illumina (NEB) and loaded for paired-end

sequencing using the Illumina HiSeq 1500.

Mutagenesis screening

Approximately 20,000 L4 stage animals expressing *gst-31::GFP* (*burIs2*) were incubated

with 20 µL of EMS in 4 mL of M9 for 4 h at 20 °C, and approximately 200,000 F3 generation

embryos were placed onto NGM agar Petri plates seeded with *E. coli* HB101 and screened

for mutants that expressed *gst-31::GFP* even in the absence of pathogen infection.

gst-31::GFP *imaging*

To image adults and embryos, young adult animals expressing *burIs2* were placed onto NGM

agar plates seeded with BIGb446 at room temperature for 24 hours. Adults and embryos were

collected and immediately imaged using a Zeiss AXIO imager A1 microscope and a

Hamamatsu ORCA-ER camera.

cysl-2::GFP *imaging*

To image adults and embryos, young adult animals expressing *nIs470* were placed onto

NGM agar plates seeded with BIGb446 at room temperature for 24 hours. Embryos were

collected and immediately imaged using a Zeiss AXIO imager A1 microscope and a

Hamamatsu ORCA-ER camera.

UV-killing bacteria

1 mL of overnight culture of BIGb446 was seeded onto 50 mm NGM agar plates and dried in

the laminar flow hood. Seeded plates were then exposed to 20 μW/cm^2 for 1 hour. Complete

killing of bacteria was confirmed by testing inoculations of bacteria in LB overnight.

cysl-2 *cloning and rescue*

Genomic *cysl-2* DNA was amplified from wild-type animals using the primers

ACGATTGGGTTGGCTGTAAG and GGTCGTACGTGTTCGTTGTG. Extrachromosomal

arrays were generated by injecting the corresponding PCR fragment and co-injection marker

into the gonad of one-day old adults at the specified concentrations. *nobEx1* was generated by

injecting *cysl-2* genomic DNA at 20 ng/μl and *myo-3::RFP* was injected at 10 ng/μl. Final

injection DNA concentration was brought up to 150 ng/μl using DNA ladder (1kb

HyperLadder – Bioline).

Generation of cysl-2 *CRISPR alleles*

syb1431 was generated by SunyBiotech. *syb1431* contains a 50 bp frameshift deletion in the

first exon with the following flanking sequences

ACCGGTGGTGAGCTCATCGGAAACACCCCA and

GGTAGAGTACATGAACCCTGCCTGCTC.

LC/MS lipid profiling

C. elegans was prepared for LC-MS lipidomics and acyl-carnitine analysis as previously

described [36] with minor modifications. Briefly, ~40 μL of concentrated embryos were re-

suspended in 100 μL of water, then 0.4 mL of chloroform was added to each sample followed

by 0.2 mL of methanol containing the stable isotope labelled acyl-carnitine internal standards

(Butyryl-L-carnitine$_{-d7}$ at 5 μM and Hexadecanoyl-L-carnitine$_{-d3}$ at 5 μM). The samples were

then homogenised by vortexing then transferred into a 2 mL Eppendorf screw-cap tube. The

original container was washed out with 0.5 mL of chloroform: methanol (2: 1, respectively)

and added to the appropriate 2 mL Eppendorf screw-cap tube. This was followed by the

addition of 150 µL of the following stable isotope labelled internal standards (approximately

10 to 50 µM in methanol): Ceramide_$C16_{d31}$, LPC_$(C14:0_{d42})$, PC_$(C16:0_{d31}$ / C18:1),

PE_$(C16:0_{d31}$ / C18:1), PG_$(C16:0_{d31}$ / C18:1), PI_$(C16:0_{d31}$ / C18:1), PS_$(C16:0_{d62})$,

SM_$(C16:0_{d31})$, TG_$(45:0_{d29})$ and TG_$(48:0_{d31})$. Then, 400 µL of sterile water was added. The

samples were vortexed for 1 min, and then centrifuged at ~20,000 rpm for 5 minutes.

For the intact lipid sample preparation, 0.3 mL of the organic layer (the lower chloroform layer)

was collected into a 2 mL amber glass vial (Agilent Technologies, Santa Clara California,

USA) and dried down to dryness in an Eppendorf Concentrator Plus system (Eppendorf,

Stevenage, UK) run for 60 minutes at 45 °C. The dried lipid samples were then reconstituted

with 100 µL of 2:1:1 solution of propan-2-ol, acetonitrile and water, respectively, and then

vortexed thoroughly. The lipid samples were then transferred into a 300 µL low-volume vial

insert inside a 2 mL amber glass auto-sample vial ready for liquid chromatography separation

with mass spectrometry detection (LC-MS) of intact lipid species.

For the acyl-carnitine sample preparation, 0.2 mL of the organic layer (the lower chloroform

layer) and 0.2 mL of the aqueous layer (the top water layer) were mixed into a 2 mL amber

glass vial and dried down to dryness. The dried acyl-carnitine samples were then reconstituted

with 100 µL of water and acetonitrile (4: 1, respectively) and thoroughly vortexed. The acyl-

carnitine samples were then transferred into a 300 µL low-volume vial insert inside a 2 mL

amber glass auto-sample vial ready for liquid chromatography separation with mass

spectrometry detection (LC-MS) of the acyl-carnitine species.

Full chromatographic separation of intact lipids [37] was achieved using Shimadzu HPLC System

(Shimadzu UK Limited, Milton Keynes, United Kingdom) with the injection of 10 µL onto a

Waters Acquity UPLC® CSH C18 column; 1.7 µm, I.D. 2.1 mm X 50 mm, maintained at 55

°C. Mobile phase A was 6:4, acetonitrile and water with 10 mM ammonium formate. Mobile

phase B was 9:1, propan-2-ol and acetonitrile with 10 mM ammonium formate. The flow was

maintained at 500 µL per minute through the following gradient: 0.00 minutes_40% mobile

phase B; 0.40 minutes_43% mobile phase B; 0.45 minutes_50% mobile phase B; 2.40

minutes_54% mobile phase B; 2.45 minutes_70% mobile phase B; 7.00 minutes_99% mobile

phase B; 8.00 minutes_99% mobile phase B; 8.3 minutes_40% mobile phase B; 10

minutes_40% mobile phase B; 10.00 minutes_40% mobile phase B. The sample injection

needle was washed using 9:1, 2-propan-2-ol and acetonitrile with 0.1 % formic acid. The mass

spectrometer used was the Thermo Scientific Exactive Orbitrap with a heated electrospray

ionisation source (Thermo Fisher Scientific, Hemel Hempstead, UK). The mass spectrometer

was calibrated immediately before sample analysis using positive and negative ionisation

calibration solution (recommended by Thermo Scientific). Additionally, the heated

electrospray ionisation source was optimised at 50:50 mobile phase A to mobile phase B for

spray stability (capillary temperature; 380 °C, source heater temperature; 420 °C, sheath gas

flow; 60 (arbitrary), auxiliary gas flow; 20 (arbitrary), sweep gas; 5 (arbitrary), source voltage;

3.5 kV. The mass spectrometer resolution was set to 25,000 with a full-scan range of m/z 100

to 1,800 Da, with continuous switching between positive and negative mode. Lipid

quantification was achieved using the area under the curve (AUC) of the corresponding high

resolution extracted ion chromatogram (with a window of ± 8 ppm) at the indicative retention

time. The lipid analyte AUC relative to the associated internal standard AUC for that lipid class

was used to semi-quantify and correct for any extraction/instrument variation.

Acyl-carnitine chromatographic separation was achieved using an ACE Excel 2 C18-PFP (150

mm, I.D. 2.1 mm, 2 μm) LC-column with a Shimadzu UPLC system. The column was

maintained at 55 °C with a flow rate of 0.5 mL/min. A binary mobile phase system was used

with mobile phase A; water (with 0.1% formic acid), and mobile phase B; acetonitrile (with

0.1% formic acid). The gradient profile was as follows; at 0 minutes_0% mobile phase B, at

0.5 minutes_100% mobile phase B, at 5.5 minutes_100% mobile phase B, at 5.51 minutes_0%

mobiles phase B, at 7 minutes_0% mobile phase B. Mass spectrometry detection was

performed on a Thermo Exactive orbitrap mass spectrometer operating in positive ion mode.

Heated electrospray source was used; the sheath gas was set to 40 (arbitrary units), the aux gas

set to 15 (arbitrary units) and the capillary temperature set to 250°C. The instrument was

operated in full scan mode from m/z 75–1000 Da. Acyl-carnitine quantification was achieved

using the area under the curve (AUC) of the corresponding high resolution extracted ion

chromatogram (with a window of ± 8 ppm) at the indicative retention time. The acyl-carnitine

analyte AUC relative to the associated internal standard AUC was used to semi-quantify and

correct for any extraction/instrument variation. All lipid values were normalized to total lipid.

RNA-seq and principal component analysis

Cutadapt version 1.18 was used to remove adapter sequences

(AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGACGCTCTTCCG

ATC for HS678 and AGATCGGAAGAGCACACGTCTGAACTCCAGTCA (forward) and

AGATCGGAAGAGCGTCGTGTAGGGAAAGAGTGT (reverse) for HS755. Cutadapt was

also used to trim the 3' ends of reads when the phred quality value was below 20. Reads

shorter than 40 bp were discarded. The genome sequence in FASTA format and annotation

file in GFF3 format of *C. elegans* were downloaded from Ensembl release 96. The genome

sequence was index with the annotation with hisat2 version 2.1.0. hisat2 was used for

aligning reads to the reference genome and the maximum number of alignments to be

reported per read was set to 5000.Featurecounts from the conda subread package version

1.6.3 was used to count the number of reads per gene. The Ensembl release 96 annotation file

in GTF format for *C. elegans* was used. Fragments were counted when they overlapped an

exon by at least 1 nucleotide and fragments are reported at the gene level. The option for the

stranded protocol was turned off. Only read pairs that had both ends aligned were counted.

Given our average fragment length of 300 bp, a distance of between 50 and 600 nucleotides

was tolerated for read pairs. Read pairs that had their two ends mapping to different

chromosomes or mapping to the same chromosome but on different strands were not counted.

Multi-mapping reads were not counted.The raw counts table was imported into R 3.5.1 for

differential expression analysis with DESeq2 version 1.22.1 Normalisation was carried out

within each contrast. The PCA plots were produced with DESeq2 function plotPCA() after

variance stabilising transformation of the data. A snakemake workflow (CITE snakemake [38])

was created for the RNA-seq analysis and can it be found at

https://github.com/cristianriccio/celegans-pathogen-adaptation.

Genomic DNA alignment

MUMmer (*Version 3.0, default parameters*) and MUMmer-plot[39] were used to visualize

global alignments of BIGb446 and BIGb468 whole genome assemblies. Further to this

NUCmer [39] and the associated dnadiff script was used to produce statistics on the alignment

of the two bacterial genomes.

Statistics and reproducibility

ANOVA analysis with post hoc p-value calculations was used for Fig. 1a, 1d, 2a, 2b, 2c, 2d,

2e, 4d, 5c, 5d, 6b, 6c, and Supplementary Fig. 3. Two-tail t-tests were used for Fig. 1b and

Supplementary Fig. 1a. For Fig. 3b, 3d, and 4e the p-value was calculated using a normal

approximation to the hypergeometric probability, as in

http://nemates.org/MA/progs/representation.stats.html. For lipidomics data (Fig. 3c and

Supplementary Table 3) p-values were calculated using two-tail t-tests and statistical

significance was determined using Bonferonni correction for multiple hypotheses. $*$ = p <

0.05, $**$ = p < 0.01, $***$ = p < 0.001, $****$ p < 0.0001. No statistical method was used to

predetermine sample size. Sample sizes were chosen based on similar studies in the relevant

literature. The experiments were not randomized. The investigators were not blinded to

allocation during experiments and outcome assessment.

909
910

918   Supplementary Figure 1. Genes associated with transgenerational inheritance and pathogen

919   infection in *C. elegans* are not required for heritable adaptation to *P. vranovensis* BIGb446.

920   Percent of wild-type and *pmk-1(km25)* mutant adults surviving on NGM plates seeded with

921   P. vranovenis BIGb446. Error bars, s.d. n = 3 replicates of 100 animals. ** = p < 0.01

958   Supplementary Figure 2. BIGb446 and BIGb468 are isolates of a single species. MUMmer

959   plot of the alignments of contigs of BIGb446 and BIGb468 assembled genomes.

1005   Supplementary Figure 3. CYSL-1 and CYSL-2 are required for adaptation to *P. vranovensis*

1006   BIGb446. Percent of wild-type and *cysl-1(mr23, mr25, mr26, mr29, mr39, mr40)* and *cysl-*

1007   *2(syb1431)* mutants surviving on plates seeded with bacterial isolates BIGb446 after 24 hrs.

1008   Error bars, s.d. n = 3 experiments of >100 animals. **** $p < 0.0001$.

1009

1010

1011

1012

1013

1014

1015

1016

1017

1018

1019

1020

1021

1022

1023

1024

1044   Supplementary File 1. Genome assembly of *P. vranovensis* BIGb446. FASTA format file of

1045   *P. vranovensis* BIGb446 genomic DNA.

1046

1047   Supplementary File 2. Genome assembly of *P. vranovensis* BIGb468. FASTA format file of

1048   *P. vranovensis* BIGb468 genomic DNA.

1049

1050   Supplementary Table 1. Comparison of genomic DNA sequences of BIGb446 and BIGb468

1051   to *P. vranovensis.*

1052

1053   Supplementary Table 2. DEseq2 and TPM analysis of mRNA expression in wild-type

1054   embryos from parents fed *E. coli* HB101 or *P. vranovensis* BIGb446.

1055

1056   Supplementary Table 3. Profile of lipid metabolite abundances in wild-type embryos from

1057   parents fed *E. coli* HB101 or *P. vranovensis* BIGb446.

1058

1059   Supplementary Table 4. DEseq2 analysis of mRNA expression in wild-type embryos from

1060   parents fed *E. coli* HB101, *P. aeruginosa* PA14, or *P. luminescens* Hb.

1061

1062   Supplementary Table 5. DEseq2 analysis of mRNA expression in wild-type and *wdr-*

1063   *23(bur9)* mutant embryos from parents fed *E. coli* HB101. Input for wild-type embryos fed *E.*

1064   *coli* HB101 are the same as Supplementary Table 4 and these samples were analysed at the

1065   same time.

1066

1067   Supplementary Table 6. Statistics Source Data

1068

References and Notes:

1. Burton, N. O. *et al.* Insulin-like signalling to the maternal germline controls progeny response to osmotic stress. *Nat. Cell Biol.* **19**, 252–257 (2017).

2. Dantzer, B. *et al.* Density triggers maternal hormones that increase adaptive offspring growth in a wild mammal. *Science* **340**, 1215–1217 (2013).

3. Jobson, M. A. *et al.* Transgenerational Effects of Early Life Starvation on Growth, Reproduction, and Stress Resistance in Caenorhabditis elegans. *Genetics* **201**, 201–212 (2015).

4. Lev, I. *et al.* MET-2-Dependent H3K9 Methylation Suppresses Transgenerational Small RNA Inheritance. *Curr. Biol. CB* **27**, 1138–1147 (2017).

5. Ma, C. *et al.* N6-methyldeoxyadenine is a transgenerational epigenetic signal for mitochondrial stress adaptation. *Nat. Cell Biol.* **21**, 319–327 (2019).

6. Roth, O. *et al.* Paternally derived immune priming for offspring in the red flour beetle, Tribolium castaneum. *J. Anim. Ecol.* **79**, 403–413 (2010).

7. Vellichirammal, N. N., Gupta, P., Hall, T. A. & Brisson, J. A. Ecdysone signaling underlies the pea aphid transgenerational wing polyphenism. *Proc. Natl. Acad. Sci. U. S. A.* **114**, 1419–1423 (2017).

8. Moore, R. S., Kaletsky, R. & Murphy, C. T. Piwi/PRG-1 Argonaute and TGF-β Mediate Transgenerational Learned Pathogenic Avoidance. *Cell* (2019). doi:10.1016/j.cell.2019.05.024

9. Hamilton, W. D., Axelrod, R. & Tanese, R. Sexual reproduction as an adaptation to resist parasites (a review). *Proc. Natl. Acad. Sci. U. S. A.* **87**, 3566–3573 (1990).

10. Little, T. J. & Kraaijeveld, A. R. Ecological and evolutionary implications of immunological priming in invertebrates. *Trends Ecol. Evol.* **19**, 58–60 (2004).

11. Dhinaut, J., Chogne, M. & Moret, Y. Immune priming specificity within and across

generations reveals the range of pathogens affecting evolution of immunity in an insect.

J. Anim. Ecol. **87**, 448–463 (2018).

12. Yue, F. *et al.* Maternal transfer of immunity in scallop Chlamys farreri and its trans-

generational immune protection to offspring against bacterial challenge. *Dev. Comp.*

Immunol. **41**, 569–577 (2013).

13. Huang, C. C. & Song, Y. L. Maternal transmission of immunity to white spot syndrome

associated virus (WSSV) in shrimp (Penaeus monodon). *Dev. Comp. Immunol.* **23**, 545–

552 (1999).

14. Luna, E., Bruce, T. J. A., Roberts, M. R., Flors, V. & Ton, J. Next-generation systemic

acquired resistance. *Plant Physiol.* **158**, 844–853 (2012).

15. Hanson, L. A. *et al.* The transfer of immunity from mother to child. *Ann. N. Y. Acad. Sci.*

987, 199–206 (2003).

16. Samuel, B. S., Rowedder, H., Braendle, C., Felix, M.-A. & Ruvkun, G. Caenorhabditis

elegans responses to bacteria from its natural habitats. *Proc. Natl. Acad. Sci. U. S. A.* **113**,

E3941-3949 (2016).

17. Gomila, M., Pena, A., Mulet, M., Lalucat, J. & Garcia-Valdes, E. Phylogenomics and

systematics in Pseudomonas. *Front. Microbiol.* **6**, 214 (2015).

18. Tran, P. N., Savka, M. A. & Gan, H. M. In-silico Taxonomic Classification of 373

Genomes Reveals Species Misidentification and New Genospecies within the Genus

Pseudomonas. *Front. Microbiol.* **8**, 1296 (2017).

19. Yoon, S.-H., Ha, S.-M., Lim, J., Kwon, S. & Chun, J. A large-scale evaluation of

algorithms to calculate average nucleotide identity. *Antonie Van Leeuwenhoek* **110**,

1281–1286 (2017).

20. Tvrzova, L. *et al.* Pseudomonas moraviensis sp. nov. and Pseudomonas vranovensis sp.

nov., soil bacteria isolated on nitroaromatic compounds, and emended description of

Pseudomonas asplenii. *Int. J. Syst. Evol. Microbiol.* **56**, 2657–2663 (2006).

21. Greer, E. L. *et al.* Transgenerational epigenetic inheritance of longevity in Caenorhabditis

elegans. *Nature* **479**, 365–371 (2011).

22. Buckley, B. A. *et al.* A nuclear Argonaute promotes multigenerational epigenetic

inheritance and germline immortality. *Nature* **489**, 447–451 (2012).

23. Ashe, A. *et al.* piRNAs can trigger a multigenerational epigenetic memory in the

germline of C. elegans. *Cell* **150**, 88–99 (2012).

24. Bolz, D. D., Tenor, J. L. & Aballay, A. A conserved PMK-1/p38 MAPK is required in

caenorhabditis elegans tissue-specific immune response to Yersinia pestis infection. *J.*

Biol. Chem. **285**, 10832–10840 (2010).

25. Kim, D. H. *et al.* A conserved p38 MAP kinase pathway in Caenorhabditis elegans innate

immunity. *Science* **297**, 623–626 (2002).

26. Engelmann, I. *et al.* A comprehensive analysis of gene expression changes provoked by

bacterial and fungal infection in C. elegans. *PloS One* **6**, e19055 (2011).

27. Burton, N. O. *et al.* Neurohormonal signaling via a sulfotransferase antagonizes insulin-

like signaling to regulate a Caenorhabditis elegans stress response. *Nat. Commun.* **9**, 5152

(2018).

28. Tang, L., Dodd, W. & Choe, K. Isolation of a Hypomorphic skn-1 Allele That Does Not

Require a Balancer for Maintenance. *G3 Bethesda Md* **6**, 551–558 (2015).

29. Staab, T. A. *et al.* The conserved SKN-1/Nrf2 stress response pathway regulates synaptic

function in Caenorhabditis elegans. *PLoS Genet.* **9**, e1003354 (2013).

30. Xu, Z. *et al.* WDR-23 and SKN-1/Nrf2 Coordinate with the BLI-3 Dual Oxidase in

Response to Iodide-Triggered Oxidative Stress. *G3 Bethesda Md* **8**, 3515–3527 (2018).

31. Choe, K. P., Przybysz, A. J. & Strange, K. The WD40 repeat protein WDR-23 functions

with the CUL4/DDB1 ubiquitin ligase to regulate nuclear abundance and activity of

SKN-1 in Caenorhabditis elegans. *Mol. Cell. Biol.* **29**, 2704–2715 (2009).

32. Budde, M. W. & Roth, M. B. The response of Caenorhabditis elegans to hydrogen sulfide

and hydrogen cyanide. *Genetics* **189**, 521–532 (2011).

33. Ma, D. K., Vozdek, R., Bhatla, N. & Horvitz, H. R. CYSL-1 interacts with the O2-

sensing hydroxylase EGL-9 to promote H2S-modulated hypoxia-induced behavioral

plasticity in C. elegans. *Neuron* **73**, 925–940 (2012).

34. Rohlfing, A.-K., Miteva, Y., Hannenhalli, S. & Lamitina, T. Genetic and physiological

activation of osmosensitive gene expression mimics transcriptional signatures of

pathogen infection in C. elegans. *PloS One* **5**, e9010 (2010).

35. Brenner, S. The genetics of Caenorhabditis elegans. *Genetics* **77**, 71–94 (1974).

36. FOLCH, J., LEES, M. & SLOANE STANLEY, G. H. A simple method for the isolation

and purification of total lipides from animal tissues. *J. Biol. Chem.* **226**, 497–509 (1957).

37. Virtue, S. *et al.* Peroxisome Proliferator-Activated Receptor gamma2 Controls the Rate

of Adipose Tissue Lipid Storage and Determines Metabolic Flexibility. *Cell Rep.* **24**,

2005-2012.e7 (2018).

38. Köster, J. & Rahmann, S. Snakemake—a scalable bioinformatics workflow engine.

Bioinformatics **34**, 3600–3600 (2018).

39. Kurtz, S. *et al.* Versatile and open software for comparing large genomes. *Genome Biol.*

5, R12 (2004).

1163