Kin structure and roost fidelity in greater noctule bats

João D. Santos[1,2,*], Christoph F.J. Meyer[1,3], Carlos Ibáñez[4], Ana G. Popa-Lisseanu[4] & Javier Juste[4,5]

[1] Centre for Ecology, Evolution and Environmental Changes (cE3c), Faculty of Sciences, University of Lisbon, 1749-016 Lisbon, Portugal

[2] UMR AGAP, CIRAD, F-34398 Montpellier, France

[3] School of Science, Engineering and Environment, University of Salford, Salford, M5 4WT, United Kingdom

[4] Department of Evolutionary Ecology, Estación Biológica de Doñana (CSIC), Avenida Américo Vespucio 26, 41092 Seville, Spain

[5] Centro de Investigación Biomédica en Red de Epidemiología y Salud Pública, CIBERESP, Spain

*Correspondent: João Santos, present address: Cirad, AGAP, TA A-108 / 03, Avenue Agropolis, 34398 Montpellier Cedex 5, France; E-mail: dourado.jns@gmail.com, Tel: +33623720645

Running Header: Kin structure of the greater noctule

ABSTRACT

Roost fidelity is an important aspect of mammalian biology. Studying the mechanisms

underlying philopatry can help us understand a species' energetic requirements, ecological

constraints and social organisation. Temperate bat species notably exhibit a comparatively high

degree of female philopatry, resulting in maternity colonies segregated at the mitochondrial

level. We focus on the greater noctule, *Nyctalus lasiopterus*, to study this behaviour in more

depth. We make use of microsatellite data for 11 markers across 84 individuals residing in Maria

Luisa Park in Seville, Spain. At the time of sampling, this urban park boasted the highest number

of registered bats of this species, among which three social groups were observed to segregate

spatially. We studied the distribution of pairs of individuals across filial relationship categories

and relatedness estimates relative to the social group of each individual. This analysis was

complemented by information on roost-use frequency among a subset of genotyped bats. We

found no significant relationship between roost use and genetic distance, but there was evidence

of higher group-sharing with increased levels of relatedness. Mother-daughter pairs shared the

same group more often than expected, as did pairs of individuals of relatedness above 0.36. We

discuss the implications of these results in terms of the behavioural ecology of temperate bats

and for conservation efforts aimed at preserving them.

Keywords: philopatry, genetic structure, microsatellites, relatedness, *Nyctalus lasiopterus*,

Chiroptera

INTRODUCTION

Female philopatry, or fidelity to a birth site or region, is a common trait among mammals

(Greenwood, 1980; Clutton-Brock & Lukas, 2012), including bats (Burland & Wilmer, 2001).

Bats display an exceptionally high degree of female philopatry and colony fidelity (Kerth et al.,

2008). While this can be partly explained by the temporal stability and sparse distribution of the

roosts used by cave-dwelling species (Lewis, 1995; Kerth, 2008), it is still common in tree- and

foliage-roosting bats, for which roost availability may not constitute a limiting factor (Kerth,

2008). Philopatry in female temperate tree-roosting bats has thus been attributed to familiarity

with the natal area, roost and foraging locations, or the maintenance of social relationships

(Kerth, 2008; Olivera-Hyde et al., 2019).

Sociality theory posits that philopatry and traits linked to colony fidelity were selected for

in order to take advantage of the benefits of group living. Indeed, reported interactions among

females of bat colonies include communal nursing (Wilkinson, 1992), social grooming

(Wilkinson, 1986; Kozhurina, 1993; Kerth 2008), food sharing (Carter & Wilkinson, 2013) and

information transfer about roost sites and foraging locations (Wilkinson & Boughman, 1998;

Kerth & Reckardt, 2003; Chaverri et al., 2010). Strong female-biased philopatry together with

high longevity in bats (Barclay & Harder, 2003) should lead to the formation of matrilineal

societies and open the door for the fixation of cooperative behaviours through kin selection

(Hamilton 1964; Emlen & Oring, 1977; Clutton-Brock, 2009). However, the extent to which

these interactions are maintained by kin selection is unknown. For example, food sharing in

Desmodus rotundus was reported to be more dependent on reciprocity than relatedness (Carter &

Wilkinson, 2013). Likewise, information transfer in both the evening and Bechstein's bats

showed no evidence of kin selection (Wilkinson, 1992; Kerth & Reckardt, 2003). For other

behaviours, such as association and grooming, empirical support for different species is

incongruent (Metheny et al., 2008; Patriquin et al., 2013).

Studies monitoring roost associations have shown females will roost preferentially with

some females rather than others, resulting in a non-random association pattern within the colony

(Willis & Brigham, 2004; Metheny et al., 2008; Patriquin et al., 2013; Naďo et al., 2017).

Because tree-roosting bats occupy hollows not large enough for more than a small number of

individuals, regular fission and fusion of subgroups could help maintain larger social groups

spread over a given area (O'Donnell, 2000; Patriquin et al., 2016). A recent review of studies

reporting pairwise association and genetic relatedness data across nine species of bats spanning

four families found no correlation between the strength of associations between kin and the

occurrence of complex cooperative behaviours (Wilkinson et al., 2019). Rather, the authors

observed that species that switched roosts often tended to present stronger associations between

relatives. Understanding the causes underlying philopatry in this context is rendered particularly

difficult. The kind of philopatry at play – to the natal area (which may not necessarily relate to

social behaviour) or to the natal colony, becomes elusive.

The greater noctule, *Nyctalus lasiopterus* (Schreber, 1780), is one of the largest and rarest

European vespertilionid bats (Ibáñez et al., 2004). Greater noctules typically roost in large, old

trees (Alcalde et al., 2016). The sparse number of existing records suggests that the species'

distribution is circum-Mediterranean and patchy from Morocco to Uzbekistan (Ibáñez et al.,

2004). Western Andalusia, Spain, is unique for harboring several known maternity colonies of

greater noctules in three main breeding areas (Santos et al., 2016). Females consolidate colonies

in spring to breed, and remain until late August, when they disaggregate and disperse,

presumably to mating and hibernation sites (Ibáñez et al., 2009), although some cases of

permanent residency have been observed (C. Ibáñez and J. Juste, unpublished data). Few males

have been found in these colonies, contrasting with other sites in northern Spain and Europe

(Ibáñez et al., 2009).

In 2008, a study on the social structure of the greater noctule population of Maria Luisa

Park (MLP) in Seville (Spain) revealed it to follow a fission-fusion structure, with bats switching

roosts on average every three days and segregated in three spatially distinct social groups (9-17%

overlap), whereby the same tree holes were rarely used by bats of different groups (Popa-

Lisseanu et al., 2008; Fortuna et al., 2009). Moreover, the foraging areas of the three groups

(extending over 40 km away from the park) overlapped almost completely. Popa-Lisseanu et al.

(2008) further reported that females were loyal to their groups for at least five years, and that

juvenile females returned to their natal roosting areas during the study period. The authors

concluded that the three groups observed within the park could be considered independent

maternity colonies, defined as groups of reproductive females roosting together. However, more

recently, a genetic analysis encompassing the region of Andalusia, Spain, and including three

maternity colonies apart from Seville (which was considered as a single population), found

significant mitochondrial structure at the regional scale, but not within the park (Santos et al.,

2016).

In this study, we narrowed the geographic focus of our previous work on regional genetic

structure (Santos et al., 2016) to the urban park in Seville. Using bi-parentally inherited markers

(microsatellites), we asked whether the status of colony attributed to the different social groups

within the park could be verified genetically. If the groups represent socially and

demographically separated colonies, then we expect this behaviour to lead to genetic isolation,

detectable as: i) higher within- than between-group relatedness, conditional on the age of each

group, since higher within-group relatedness was not observed during colonization (Santos et al.,

2016), ii) a similar deviation in the distribution of related pairs of females across groups to that

reported by Santos et al. (2016) across the region (throughout the article, we refer to the groups

within the park as groups or social groups, and to the Seville population as population or colony).

Furthermore, within our genetic dataset, we include samples from individuals used by Popa-

Lisseanu et al. (2008) to identify temporal roosting associations. Because the three groups are

both physically and socially defined, we used this to refine our initial hypothesis: H0) group

identity, physical or social, does not impact individual decisions to return – genetic structure

should be absent, genetic and roosting data disassociated; H1) Group identity alone has inherent

impact – higher within-group relatedness should be observed but no roosting/genetic structure;

H2) Group identity impacts dyad structure because *N. lasiopterus* females chose to roost with

relatives – both dyad structure and roosting/genetic association should be observed. H3, the

absence of genetic isolation in the presence of roosting/genetic association, was not seriously

considered as it would imply a more complicated scenario of familial interactions. Regarding our

understanding of the philopatric behaviour of female giant noctules, H0 corresponds to a coarse

site fidelity, and could be explained by the limited number of alternative roosting opportunities

in the region. H1 implies a more specific preference for certain roosts or individuals, and H2

would indicate the maintenance of kin-relationships as the basis for the return.

Materials and Methods

Focal Species and Study Population

Very few breeding colonies of *N. lasiopterus* are known outside of the Iberian Peninsula:

France, Hungary, Slovakia, Belarus, Ukraine and Russia (Uhrin et al., 2006; Estók & Gombkötő,

2007; Dubourg-Savage et al., 2013; Dombrovski et al., 2016; Vlaschenko et al., 2016; Kovalov

et al., 2018).

The study population is located in MLP, Seville (37°22′29″N, 5°59′19″W). The 122-

year-old park extends over 38 ha and contains many large old trees that bear enough cavities to

harbour the three distinct colonies. Most roost trees belong to *Platanus* spp., *Gleditsia*

triacanthos, *Sophora japonica* and *Washingtonia filifera*. With a population previously estimated

at ca. 500 individuals, but now in dramatic decline due to an invasive parakeet species

(Hernández-Brito et al., 2018) that is taking over the roosting holes, it was once the largest

known breeding site of greater noctules (Popa-Lisseanu et al., 2008).

Sampling

We analysed biopsy samples of 82 adult female bats from MLP collected from 2004 to

2007. Each bat was assigned to one of three areas of the park identified by Popa-Lisseanu et al.

(2008) as characteristic of different social groups. Assignment was based on the tree in which

individuals roosted in when captured. Group designations (I, II and III) were maintained from

that earlier study. Twenty-four individuals were sampled from each group. Samples included 25

individuals that were radio-tracked to study the roost-switching behaviour of the population that

was used to identify and define the social group dynamics in the previous study (Popa-Lisseanu

et al., 2008).

Capture and experiments were performed under permit of the Ethical Committee of EBD-

CSIC and the techniques used meet the guidelines published by the American Society of

Mammalogists (Gannon and Sikes, 2007) on the use of wild mammals in research.

138   *Molecular Methods*

Total DNA was extracted from 3 mm diameter wing punches using a modified version of

the salt-based protocol developed by Aljanabi & Martinez (1997).

All individuals were genotyped at 11 nuclear microsatellite loci: Nle 2, 3 and 6-11

(developed for *Nyctalus leisleri*, see Boston et al*.,* 2008); EF4 (developed for *Eptesicus fuscus*,

Vonhof et al*.,* 2002); P20 and P217 (developed for *Pipistrellus sp.* Kaňuch et al., 2007). All

were tested in muscle tissue prior to genotyping (see Supplementary Methods for a detailed

description of DNA extraction, amplification, sequencing, and microsatellite genotyping) and

successfully employed in our previous study (Santos et al., 2016). Labelling followed Schuelke's

procedure (2000).

149   *Data Analysis*

All nuclear markers were in linkage equilibrium as assessed with FSTAT v. 2.9.3.2

(Goudet et al., 1995). Using CERVUS, we estimated observed and expected heterozygosities, as

well as deviations from Hardy-Weinberg equilibrium (HWE). Of the 11 microsatellites, three

(Nle6, P20 and P217) deviated significantly from HWE (see Supplementary Table S1).

Pairwise and mean relatedness values (r) were estimated using ML-Relate (Kalinowski et

al., 2006), which implements a corrected maximum-likelihood method for estimating relatedness

that allows loci with null alleles to be incorporated into the analysis (Wagner et al., 2006). Three

of the 11 loci used in this study showed the highest frequencies of null alleles (min=0.097,

mean=0.187). Incidentally, this might have been the cause of the departure from HWE, as it

would simulate a heterozygote deficiency. Because we did not want to abdicate the inferential

power of three loci, we resorted to the method by Wagner et al. (2006), implemented in ML-

Relate, and kept all 11 loci. Using this software, pairwise relations were classed as Parent-

Offspring (PO), Full-Sibling (FS), Half-Sibling (HS), or Unrelated (U), and each attributed a

value *r* of pairwise relatedness. Principal component analysis (PCA) was performed on the

squared matrix of pairwise relatedness values.

With reference to the three groups within the park, pairwise relationships were classed

according to whether or not both individuals had been captured within the same group. We tested

the null hypothesis that each group shared the same population mean as the whole population.

Firstly, analysis of molecular variance (AMOVA) was conducted between global and within-

group data sets of relatedness estimates, using the function 'f_oneway' of the python package

'scipy.stats' (Oliphant, 2007). We then compared within-group relatedness to relatedness across

the park independently for each group using a randomization test. We relied on two-tailed t-tests

as implemented in the function 'ttest_ind' of the python package 'scipy.stats' (Oliphant, 2007).

We approximated the distribution of the test statistic by comparing relatedness estimates across

the park to group-sized samples of randomly selected relatedness values. The procedure was

repeated 1000 times for each group. The final *p*-value was calculated as the proportion of

sampled statistics below that observed for each group.

Regarding group choice, we assumed as H0 that the probability of choosing the same

group as another individual at random follows a binomial distribution. We further considered this

probability to be conditional on the relative size of the group, not the relative sample size. Then,

the probability $P(O)$ of pairs of individuals falling across groups was estimated as $P(O) =$

$P_I(1 - P_I) + P_{II}(1 - P_{II}) + P_{III}(1 - P_{III})$. The values P_I, P_{II} and P_{III} were estimated as the

proportions of individuals registered to roost in groups I, II and III at the time of sampling (81,

61 and 114 bats respectively, Popa-Lisseanu et al. 2008), so that $P(O) = 0.644$. The probability

$P(I)$ of two individuals choosing the same group was estimated as $1 - P(O) = 0.356$. We

compared the expected and observed proportions across the data set and within the classes of

relation PO, FS, HS and U, and then studied the deviation of observed within-group pairs as a

function of relatedness values. We estimated the upper-tail *p*-value of observed proportions of

same-group pairs within each class assuming a binomial probability $P(I)$. We further estimated

upper-tail *p*-values of the proportion of dyads falling within the same group along windows of

relatedness. Proportions were calculated among relationships presenting values within a 0.15

window of relatedness around a central value ranging between 0.075 and 0.5 in increments of

0.01. Window size was selected based on the minimum, median and standard deviation of the

number of relationships across windows of a given size (see Supplementary Figure S1).

To explore the relationship between genetic relatedness and roost use we resorted to the

matrix of pairwise roost use similarity among 15 bats radio-tracked in 2004 used by Popa-

Lisseanu et al. (2008) to infer the existence of social groups within the park (Supplementary

Figure S2). Similarity was estimated using the Freeman-Tukey statistic (Krebs, 1989). All

individuals considered in that study were genotyped. A Mantel test of the two distance matrices

was performed using the mantel.rtest function of R library ade4 (Dray and Dufour 2007) with the

default number of repeats.

We constructed network graphs of roosting associations and parent-offspring dyads.

Roosting association edges were created as dyads presenting a similarity value above 0.05

(Freeman-Tukey statistic). Networks were constructed using the Fruchterman-Reingold

algorithm as implemented in the Networkx python package (Aric et al., 2008).

Results

We first evaluated genetic structure across the park considering its known subdivisions.

No genetic structure was observed through PCA (Fig. 1A), and one-way AMOVA of within-

group relatedness estimates was non-significant ($p = 0.16$, d.f. $= 1133$). We then considered the

interplay of group assignment, genetic relatedness and relationships. Of the total number of

possible associations within the park and groups (every female paired with every other female),

the percentage of those exceeding a relatedness degree of 0.25 accounted for 6% of the total

(Table 1). However, around 81% of females shared their group with at least one close relative (*r*

> 0.25, Table 1), while 95% shared the park (considering close relatives found in both the same

and different groups). We focused on each group to ascertain whether mean relatedness exceeded

expectations under randomness. The result of the t-test of pairwise relatedness means between

total and group III dyads was significant ($p < 0.01$, Fig. 1B), as was that between total and group

I ($p < 0.05$, Figure 1B).

We also looked at the impact of group-sharing on relationship categories and relatedness

estimates. The proportion of parent-offspring pairs that shared the same group differed

significantly ($p < 0.001$) from random expectations (Fig. 3). The same proportions for FS, HS

and U pairs did not deviate significantly from the expected proportions (Fig. 2). However, a

decreasing trend is observed when these groups are considered in order of decreasing

relatedness.

When the same test is performed on pairs of individuals of increasing relatedness, the

proportion of these pairs falling within the same group rapidly becomes significant (r > 0.18, Fig.

3A). When only pairs within a moving window of relatedness are considered (size= 0.15, step=

0.01), the same proportion becomes significant above a relatedness level of 0.36 (Fig. 3B).

Finally, we assessed the effect of roosting association on genetic relatedness. A Mantel

test revealed no significant association between the two variables (r = 0.06, simulated *p*-value =

0.27). This subset of individuals represented 37 half-sibling relationships, 2 full-sibs, 1 parent-

offspring pair and 65 unrelated dyads. Roost association between the mother-daughter pairs was

above average, but not significantly so (P = 0.09).

Discussion

Female fidelity to breeding colonies has been reported in a number of temperate bat

species (Rossiter et al., 2002; Castella et al., 2008; Juste et al., 2009), leading, in the extreme, to

the formation of closed societies as in *Myotis bechsteinii* (Kerth et al., 2000). The existence of

among-colony mitochondrial structure at the regional level supports a certain degree of

philopatry among female *N. lasiopterus* (Santos et al., 2016). Our results confirm that this

behaviour extends to social groups within the population of Seville. This behaviour leads to a

proportion of female pairs sharing the same group that increases with relatedness, becoming

significant at higher levels (r > 0.36) and resulting in a substantial proportion (62.5%, 25 / 40) of

predicted mother-daughter pairs sharing the same social group. While this value is below that

observed for the strictly philopatric Bechstein's bat (Kerth et al., 2002), it is considerably higher

than reported in big brown bats (9%; Vonhof et al., 2008). We further find that two of the groups

present significantly higher relatedness levels than the park as a whole, indicating that this

behaviour is potentially stable. The fact that group II females do not exhibit higher average

genetic relatedness could be explained if this group was recent, since group formation in this

species is not dependent on kin (Santos et al., 2016).

The reasons for philopatric behaviour have been widely discussed (Moussy et al., 2013),

and generally attributed to the females' acquaintance with their natal area, to facilitate access to

foraging and roosting sites, and to the benefits they might derive from social relationships and

cooperative behaviours with conspecifics (Kerth, 2008; Kerth et al., 2011). We tested for the

effect of roosting associations, previously shown to follow the same park sub-sections, and found

a lack of correlation between roost use and genetic relatedness. Despite the limited power of this

analysis due to small sample sizes, our results thus seem to refute our initial hypotheses H0 and

H2 in favour of H1: females tend to return to natal roosting areas, but this choice is not driven by

a desire to roost with kin. We can also discard explanations based on facilitated access; we are

not aware of differences in roost characteristics between the three areas of the park that could

account for the patterns observed, and the three groups have completely overlapping foraging

areas (Popa-Lisseanu et al., 2009). Philopatric behaviour in Seville can thus be explained either

by familiarity with the natal area, or the maintenance of social relationships that are not kin-

based. Associative learning is the imprinting of roosts by non-volant young when being carried

by their mothers between day roosts (O'Donnell, 2000). Based on reports of observational

learning, O'Donnell (2000) reasoned that if young *Chalinolobus tuberculatus* could have their

mother's social group pool of favourite roost trees imprinted through repetitive roost switching,

group structure would be maintained. The same mechanism could be at play in *N. lasiopterus.*

The alternative hypothesis is that roost fidelity functions to maintain social bonds,

helping returning females reconnect with acquaintances made at specific roosts. Studying the

social organization of a maternity colony of noctule bats *Nyctalus noctula* in captivity,

Kozhurina (1993) reported the best predictors of social organization to be age and sex (92.6 %

classification accuracy versus 29.6 % for family membership) and that the disparity between age

classes remained over time. It is important to add that occasions of social grooming occurred

strictly among members of the same social groups. A more recent study of the temporal and

genetic components of associative dynamics in *Nyctalus leisleri* also found a stronger impact of

cohort year on the composition of social groups inferred through network analysis (Naďo et al.,

2017). As pointed out by the authors, although genetic relatedness was also observed to

significantly impact association between dyads, this could be the result of different genetic

profiles between cohorts due to male-biased gene flow. Cooperative behaviours and long-term

associations among colony members, sometimes spanning several years, have been described in

numerous bat species (Wilkinson & Boughman, 1998; Carter & Wilkinson, 2013). Kozhurina

(1993) further observed that noctule bats start forming associations in their first year. If this is

true for greater noctules, maintaining these bonds, either for the comfort derived from familiarity

and predictability (e.g. reduced aggression, reduced demands on limited attention capacity

already strained by pup-rearing), or the increased likelihood of reciprocal cooperative behaviour,

could explain why individuals tend to return to the area of the park they were born in. This

theory has the advantage of providing an explanation for the formation of social groups, which

are a pre-requisite for the associative learning hypothesis.

In summary, we found that female *N. lasiopterus* will seek to return to the specific areas

or social groups they were born in, not just the region or breeding area, more often than expected

by chance. We further found that roost associations are not kin-based. While these findings

support the hypothesis that group structure is maintained through philopatry (Popa-Lisseanu et

al., 2008), it is likely that neither associative learning nor sociality account for this behaviour

entirely, but that instead both play a role. Both hypotheses explain the specificity of natal

philopatry in greater noctules. Associative learning has the obvious adaptive advantage of

allowing young adult females to benefit from their mothers' successful choice of colony. Strong

social bonds leading to cooperative behaviours, such as allo-nursing or information sharing,

bring fitness benefits to nursing mothers with already high energy expenditures, an explanation

which seems particularly plausible for females of this species in light of their challenging

carnivorous habits during part of the year (Dondini and Vergari, 2000; Ibáñez et al., 2001; Popa-

Lisseanu et al., 2007). While both hypotheses seem plausible, each rests on certain assumptions

regarding roosting behaviour or social interactions that lack confirmation in this species. There is

one case of reported co-operative nursing in captivity (Kozhurina, 1993), but no study to date has

captured roosting associations or behaviours with enough precision to infer the correlates of

social organization in maternity colonies of *N. lasiopterus*. Likewise, insufficient information is

available on roost fidelity to assess how it correlates with breeding status or to verify roost reuse.

Popa-Lisseanu et al. (2008) did report a decrease in roost switching during lactation, an

observation that perhaps failed to reach significance due to lack of statistical power. Repeating

this experiment with a larger sample size could be informative since a significant reduction in

roost switching would render the associative learning hypothesis less likely. In the meantime,

regardless of the mechanism involved, we argue that in light of our results, and those of previous

studies in this park, it is safe to re-classify the social groups in MLP as separate colonies.

Determining the relative importance of associative learning and social bonds in this

species is crucial, as it could direct future conservation efforts regarding the few known

maternity colonies. In order to discriminate between these two hypotheses, future studies should

focus on roost fidelity and roosting patterns as well as on long-term associations. Evidence of

roost reuse by individual females, as well as varying roost fidelity according to breeding status

would point to associative learning. These will have to be compared with the frequency of long-

term associations within the same colonies in order to determine the relative importance of social

bonds. More generally, discovering the mechanism behind philopatry represents an important

step in our understanding of the thought processes behind the individual decisions that structure

mammalian societies, and has implications for conservation policies aimed at preserving healthy

populations.

Acknowledgements

J. Nogueras and C. Ruiz helped collecting bat samples. We particularly thank J.L. García-

Mudarra and J.M. Arroyos-Salas for valuable advice and their technical insight. We also

acknowledge the staff of the Servicio de Parques y Jardines de Sevilla for their help and

continuous support of our research. Logistical support was provided by the Laboratorio de

Ecología Molecular, Estación Biológica de Doñana, CSIC (LEM-EBD). The regional

government of Andalusia provided permits for collecting samples and handling of bats. This

study was partially funded by the Spanish MICINN CGL 2009- 12393, the PPNN 021/2002 and

1981/2010. JDS would like to acknowledge support from the Erasmus Student Mobility

program.

337   **Supplementary Data**

Supplementary Methods – Detailed description of DNA extraction, purification, sequencing and

genotyping.

Supplementary Table S1 – Summary statistics and PCR specifications for each locus over all

colonies.

Supplementary Figure S1 - Exploratory analysis of the impact of window size on sampling

across relatedness steps.

Supplementary Figure S2 - Intra-colony network of roost-sharing between greater noctules in

Maria Luisa Park, Seville, Spain.

Supplementary Figure S3 - Fig. S3 Network of parent-offspring dyads among giant noctule bats

roosting in Maria Luisa Park, Seville, Spain.

349   **Data accessibility**

350   Microsatellite genotypes, sample ID and location were deposited in the Dryad Digital Repository

351   (doi: 10.5061/dryad.rc504). All data analysis scripts are accessible in the GitHub repository (doi:

352   10.5281/zenodo. 2653473).

353   **Conflicts of interest**

354   None declared

References

Aljanabi, S.M. & Martinez, I. (1997). Universal and rapid salt-extraction of high quality genomic
DNA for PCR-based techniques. *Nucleic Acids Res.* **25**, 4692–3.

Alcalde, J., Juste, J. & Paunović, M. (2016). *Nyctalus lasiopterus*. The IUCN Red List of
Threatened Species 2016: e.T14918A22015318.
http://dx.doi.org/10.2305/IUCN.UK.2016-2.RLTS.T14918A22015318.en. Downloaded
on 24 November 2018.

Aric, A.H., Daniel, A.S., Pieter J. S. (2008). Exploring network structure, dynamics, and function
using NetworkX. in Proceedings of the 7th Python in Science Conference (SciPy2008).
Gäel, V., Travis, V., and Jarrod, M. (Eds). Pasadena, CA USA.

Barclay, R.M.R. & Harder, L.D. (2003). Life histories of bats: life in the slow lane. In *Bat
Ecology*: 209–253. Kunz, T.H. & Fenton, M.B. (Eds.). University of Chicago Press.

Boston, E., Montgomery, I. & Prodöhl, P.A. (2008). Development and characterization of 11
polymorphic compound tetranucleotide microsatellite loci for the Leisler's bat, *Nyctalus
leisleri* (Vespertilionidae, Chiroptera). *Conserv. Genet.* **10**, 1501–1504.

Burland, T.M. & Wilmer, J.W. (2001). Seeing in the dark: molecular approaches to the study of
bat populations. *Biol. Rev.* **76**, 389–409.

Carter, G.G. & Wilkinson, G.S. (2013). Food sharing in vampire bats: reciprocal help predicts
donations more than relatedness or harassment. *Proc. R. Soc. B Biol. Sci.* **280**, 2012-2573.

Castella, V., Ruedi, M. & Excoffier, L. (2008). Contrasted patterns of mitochondrial and nuclear
structure among nursery colonies of the bat *Myotis myotis*. *J. Evol. Biol.* **14**, 708–720.

Chaverri, G., Gillam, E.H. & Vonhof, M.J. (2010). Social calls used by a leaf-roosting bat to
signal location. *Biol. Lett.* **6**, 441–444.

Clutton-Brock, T. (2009). Cooperation between non-kin in animal societies. *Nature* **462**, 51–57.

Clutton-Brock, T. & Lukas, D. (2012). The evolution of social philopatry and dispersal in female
mammals. *Mol. Ecol.* **21**, 472–492.

Dombrovski, V., Fenchuk, V. & Zhurauliou, D. (2016). New occurrence and the first breeding
record of *Nyctalus lasiopterus* in Belarus. *Vespertilio*, **18**: 55-59.

Dondini, G. & Vergari, S. (2000). Carnivory in the greater noctule bat (*Nyctalus lasiopterus*) in
Italy. *J. Zool.* **251**, 233–236.

Dubourg-Savage, M.J., Bec, J. & Gaches, L. (2013). First roosts of *Nyctalus lasiopterus* breeding
females in France. *Barbastella* **6**, 44–50.

Dray, S. & Dufour A. (2007). The ade4 Package: Implementing the Duality Diagram for
Ecologists. *J. Stat. Softw.* **22**, 1-20. doi: 10.18637/jss.v022.i04.

Emlen, S.T. & Oring, L.W. (1977). Ecology, sexual selection, and the evolution of mating
systems. *Science* **197**, 215–223.

Estók, P. & Gombkötő, P. (2007). Review of the Hungarian data of *Nyctalus lasiopterus*
(Schreber, 1780). *Fol. Hist. Nat. Mus. Matraensis* **31,** 167-172.

Fortuna, M.A., Popa-Lisseanu, A.G., Ibáñez, C. & Bascompte, J. (2009). The roosting spatial
network of a bird☐predator bat. *Ecology* **90**, 934-944.

Gannon WL, Sikes RS. Guidelines of the American Society of Mammalogists for the Use of
Wild Mammals in Research. *J. Mamm.* (2007). **88,** 809–823. doi:10.1644/06-MAMM-F-
185R1.1

Goudet, J. (1995). FSTAT (version 1.2): a computer program to calculate F-statistics. *J. Hered.*
86, 485–486.

Greenwood, P.J. (1980). Mating systems, philopatry and dispersal in birds and mammals. *Anim.*
Behav. **28**, 1140–1162.

Hamilton, W.D. (1964). The genetical theory of kin selection. *J. Theor. Biol.* **7**, 1-52.

Hernández-Brito, D., Carrete, M., Ibáñez, C., Juste, J. & Tella, J.L. (2018). Nest-site competition
and killing by invasive parakeets cause the decline of a threatened bat population. *R. Soc.*
Open Sci. **5**, 172477.

Ibáñez, C., Juste, J., Garcia-Mudarra, J.L. & Agirre-Mendi, P.T. (2001). Bat predation on
nocturnally migrating birds. *Proc. Natl. Acad. Sci.* **98**, 9700–9702.

Ibáñez, C., Guillén, A. & Bogdanowicz, W. (2004). *Nyctalus lasiopterus* — Riesenabendsegler.
In *Handbuch der Säugetiere Europas*: 695–715. Krapp, F. (Ed.). Wiebelsheim: Aula-
Verlag.

Ibáñez, C., Guillén, A., Agirre-Mendi, P.T., Juste, J., Schreur, G., Cordero, A.I., Popa-Lisseanu,
A.G. (2009). Sexual segregation in Iberian noctule bats. *J. Mammal.* **90**, 235–243.

Juste, J., Bilgin, R., Muñoz, J. & Ibáñez, C. (2009). Mitochondrial DNA signatures at different
spatial scales: from the effects of the Straits of Gibraltar to population structure in the
meridional serotine bat (*Eptesicus isabellinus*). *Heredity* **103**, 178–187.

Kalinowski, S.T., Wagner, A.P. & Taper, M.L. (2006). Ml-Relate: a computer program for
maximum likelihood estimation of relatedness and relationship. *Mol. Ecol. Notes* **6**, 576–
579.

Kaňuch, P., Fornůsková, A., Bartonička, T. & Bryja, J. (2007). Multiplex panels of polymorphic
microsatellite loci for two cryptic bat species of the genus *Pipistrellus*, developed by
cross-species amplification within the family Vespertilionidae. *Mol. Ecol. Notes* **7**, 871-873.

Kerth, G. (2008). Causes and consequences of sociality in bats. *Bioscience* **58**, 737.

Kerth, G., Mayer, F. & König, B. (2000). Mitochondrial DNA (mtDNA) reveals that female
Bechstein's bats live in closed societies. *Mol. Ecol.* **9**, 793–800.

Kerth G, Mayer F, Petit E. 2002. Extreme sex-biased dispersal in the communally breeding,
nonmigratory Bechstein's bat (Myotis bechsteinii). Mol. Ecol. 11:1491–1498. doi:
10.1046/j.1365-294X.2002.01528.x.

Kerth, G., Perony, N. & Schweitzer, F. (2011). Bats are able to maintain long-term social
relationships despite the high fission-fusion dynamics of their groups. *Proc. R. Soc. B Biol.*
Sci. **278**, 2761–2767.

Kerth, G. & Reckardt, K. (2003). Information transfer about roosts in female Bechstein's bats: an
experimental field study. *Proc. R. Soc. B Biol. Sci.* **270**, 511–515.

Kovalov, V., Hukov, V. & Rodenko, O. (2019). New record of *Nyctalus lasiopterus* (Schreber,
1780) in Ukraine with a new confirmation of carnivory. North-West. J. Zool. **15**, 91-95.

Kozhurina, E.I. (1993). Social organization of a maternity group in the noctule bat, *Nyctalus*
noctula (Chiroptera□: Vespertilionidae). *Ethology* **93**, 89–104.

Krebs, C.J. (1989). *Ecological methodology*. Harper & Row New York.

Lewis, S.E. (1995). Roost fidelity of bats: a review. *J. Mammal.* **76**, 481–496.

Metheny, J.D., Kalcounis-Rueppell, M.C., Willis, C.K.R., Kolar, K.A. & Brigham, R.M. (2007).
Genetic relationships between roost-mates in a fission–fusion society of tree-roosting big
brown bats (*Eptesicus fuscus*). *Behav. Ecol. Sociobiol.* **62**, 1043–1051.

Moussy, C., Hosken, D.J., Mathews, F., Smith, G.C., Aegerter, J.N. & Bearhop, S. (2013).
Migration and dispersal patterns of bats and their influence on genetic structure. *Mamm.*
Rev. **43**, 183–195.

Naďo, L., Chromá, R. & Kaňuch, P. (2017). Structural, temporal and genetic properties of social
groups in the short-lived migratory bat *Nyctalus leisleri*. *Behaviour* **154**, 785-807.

O'Donnell, C.F.J. (2000). Cryptic local populations in a temperate rainforest bat *Chalinolobus*
tuberculatus in New Zealand. *Anim. Conserv.* **3**, 287–297.

Oliphant, T.E. (2007). Python for scientific computing. *Comput. Sci. Eng.* **9**, 10–20.

Olivera-Hyde, M.O., Silvis, A., Hallerman, E.M., Ford, M. & Britzke, E. (2019). Relatedness
within and among *Myotis septentrionalis* colonies at a local scale. *Can. J. Zool.* **97**, 724–
735.

Patriquin, K.J., Palstra, F., Leonard, M.L. & Broders, H.G. (2013). Female northern myotis
(*Myotis septentrionalis*) that roost together are related. *Behav. Ecol.* **24**, 949–954.

Patriquin, K.J., Leonard, M.L., Broders, H.G., Ford, W.M., Britzke, E.R. & Silvis, A. (2016).
Weather as a proximate explanation for fission–fusion dynamics in female northern long-
eared bats. *Anim. Behav.* **122**, 47–57.

Popa-Lisseanu, A.G., Bontadina, F., Mora, O. & Ibáñez, C. (2008). Highly structured fission–
fusion societies in an aerial-hawking, carnivorous bat. *Anim. Behav.* **75**, 471–482.

Popa-Lisseanu, A. G., Bontadina, F. & Ibáñez, C. (2009). Giant noctule bats face conflicting
constraints between roosting and foraging in a fragmented and heterogeneous landscape. *J.*
Zool. **278**, 126-133.

Popa-Lisseanu, A. G., Delgado-Huertas, A., Forero, M. G., Rodríguez, A., Arlettaz, R. & Ibáñez,
C. (2007). Bats' conquest of a formidable foraging niche: the myriads of nocturnally
migrating songbirds. *PLoS One* **2**, e205.

Rossiter, S., Jones, G., Ransome, R. & Barratt, E. (2002). Relatedness structure and kin-biased
foraging in the greater horseshoe bat (*Rhinolophus ferrumequinum*). *Behav. Ecol. Sociobiol.*
51, 510–518.

Santos, J.D., Meyer, C.F.J., Ibáñez, C., Popa-Lisseanu, A.G. & Juste, J. (2016). Dispersal and
group formation dynamics in a rare and endangered temperate forest bat (*Nyctalus*
lasiopterus, Chiroptera: Vespertilionidae). *Ecol. Evol.* **6**, 8193–8204.

Schuelke, M. (2000). An economic method for the fluorescent labeling of PCR fragments. *Nat.*
Biotechnol. **18**, 233.

Uhrin, M., Kaňuch, P., Benda, P., Hapl, E., Verbeek, H. J., Krištín, A., Krištofík, J., Mašán, P. &
Andreas, M. (2006). On the greater noctule (*Nyctalus lasiopterus*) in central Slovakia.
Vespertilio **9,** 183-192.

Vlaschenko, A., Kravchenko, K., Prylutska, A., Ivancheva, E., Sitnikova, E. & Mishin, A.
(2016). Structure of summer bat assemblages in forests in European Russia. Turkish J. Zool.
40, 876-893.

Vonhof, M.J., Davis, C.S., Fenton, M.B. & Strobeck, C. (2002). Characterization of dinucleotide
microsatellite loci in big brown bats (*Eptesicus fuscus*), and their use in other North
American vespertilionid bats. *Mol. Ecol. Notes* **2**, 167-169.

Vonhof MJ, Strobeck C, Fenton MB. 2008. Genetic Variation and Population Structure in Big
Brown Bats (Eptesicus fuscus): Is Female Dispersal Important? *J. Mammal.* 89:1411–1420.
doi: 10.1644/08-MAMM-S-062.1.

Wagner, A.P., Creel, S. & Kalinowski, S.T. (2006). Estimating relatedness and relationships
using microsatellite loci with null alleles. *Heredity* **97**, 336–45.

Wilkinson, G.S. (1986). Social grooming in the common vampire bat, *Desmodus rotundus*.
Anim. Behav. **34**, 1880–1889.

Wilkinson, G.S. (1992). Communal nursing in the evening bat, *Nycticeius humeralis*. *Behav.*
Ecol. Sociobiol. **31**, 225–235.

Wilkinson, G.S. & Boughman, J.W. (1998). Social calls coordinate foraging in greater spear-
nosed bats. *Anim. Behav.* **55**, 337–350.

Wilkinson, G.S., Carter, G., Bohn, K.M., Caspers, B., Chaverri, G., Farine, D., Günther, L.,
Kerth, G., Knörnschild, M., Mayer, F., Nagy, M., Ortega, J. & Patriquin, K. (2019).
Kinship, association, and social complexity in bats. *Behav. Ecol. Sociobiol.* **73**, 7.

Willis, C.K.R. & Brigham, R.M. (2004). Roost switching, roost sharing and social cohesion:
forest-dwelling big brown bats, *Eptesicus fuscus*, conform to the fission–fusion model.
Anim. Behav. **68**, 495–505.

506   # Tables

507   Table 1: Mean relatedness estimates within groups and across the whole colony in Maria Luisa Park,

508   Seville, percentage of associations with $r > 0.25$ (close relatives) among total possible associations, and

509   percentage of females with close relatives within their groups as well as the whole colony. The total

510   sample consists of 84 individuals, 24 from each group, resulting in a total of 3486 possible dyads.

Group		*% associations*		
	Mean r	*SD*	*r>0.25*	*%*
I.	0.056	±0.106	6.3	89.2
II.	0.050	±0.099	5.6	75.0
III.	0.051	±0.099	5.6	78.6
Colony	**0.059**	**±0.098**	**5.7**	**95.2**

Figures

Fig. 1 Pairwise relatedness between adult female *Nyctalus lasiopterus* within and among social groups in

Maria Luisa Park, Sevilla, Spain. **A**) Biplot of a principal component analysis based on a relatedness matrix. **B**)

Boxplot of pairwise relatedness across the pooled data set and within each social group. t-tests were calculated for

group and colony sample means for each of the three groups (*p*-values shown above).

Fig. 2 Percentage of pairwise relations within and among social groups across relatedness categories. Dyads

were classed into relation groups using ML-Relate. Individual dyads were further classed according to whether they

were caught within the same areas of Maria Luisa Park. Grey: proportions of dyads between individuals from

different groups; black: proportion of dyads between individuals from the same group. *P*-value of observed

proportion of pairs from the same group for each class shown in bold. Categories PO, FS, HS and U consisted of 40,

47, 441 and 2958 dyads respectively. *P*-values were derived assuming a binomial probability of two individuals

sharing the same group $P(I) = 0.356$.

Fig. 3 Proportion of group sharing as a function of pairwise relatedness. Pairs of individuals were classified

according to whether both were captured in the same area of Maria Luisa Park. This proportion was then studied in

relation to the estimated relatedness of each pair. **A**) Proportion of dyads within the same group with relatedness

values above abscissae. **B**) Proportion of dyads within relatedness windows that fall within the same group (window

size= 0.15 *r*, step= 0.03 *r*). The upper-bound 95% confidence interval (UCI, lozenge-line) was estimated from a

binomial distribution of $p = (I)$. The dashed line indicates the expected proportion $P(I)$ of within-group pairs.