Differential Allele-Specific Expression Uncovers Breast Cancer Genes Dysregulated By *Cis* Noncoding Mutations

Pawel F Przytycki[1,2,3] and Mona Singh[1,2,*]

1 Department of Computer Science, Princeton University, Princeton, NJ 08544, USA

2 Lewis-Sigler Institute for Integrative Genomics, Princeton University, Princeton, NJ 08544, USA

3 Present address: Gladstone Institutes, San Francisco, CA 94158, USA

* Corresponding author: mona@cs.princeton.edu

Abstract

Identifying cancer-relevant mutations in noncoding regions is extremely challenging due to the large numbers of such mutations, their low levels of recurrence, and the general difficulty in interpreting their impact. To uncover genes that are dysregulated due to somatic mutations in *cis*, we introduce the concept of *differential allele-specific expression* (ASE) and develop methods to identify genes within an individual's cancer whose ASE differs from what is found in matched normal tissue. When applied to breast cancer tumor samples, our methods readily detect the known allele-specific effects of copy number variation and nonsense mediated decay. Further, genes that are found to recurrently exhibit differential ASE across samples are cancer relevant. Genes with *cis* mutations are enriched for differential ASE, and we find 147 potentially functional noncoding mutations *cis* to genes that exhibit significant differential ASE. Overall, our results suggest that differential ASE is a promising means for discovering gene dysregulation within an individual due to *cis* noncoding mutations.

Introduction

Cancer is a heterogeneous disease in which every tumor exhibits a unique combination of genotypic and phenotypic features.[1] A major aim of cancer genomics is to understand how acquired somatic mutations within cancer cells lead to the dysregulation that allows cells to grow and proliferate uncontrollably.[2] Towards this aim, consortia such as TCGA[3] and ICGC[4] have profiled the tumors of thousands of individuals across tens of cancer types to determine DNA sequences, gene expression levels, copy number variation, and other functional genomics data. Initial analyses have revealed that each individual's tumor typically has a unique combination of mutations, most of which play no functional role in cancer.[1,2]

While many computational methods have been developed to identify which of the numerous somatic alterations observed within a tumor are linked to cancer phenotypes, these have mostly focused on mutations affecting protein-coding regions.[5] However, the vast majority of cancer somatic mutations are found in noncoding regions,[6] as ~98% of the human genome is noncoding.[7,8] Further, since widespread gene expression dysregulation is common in cancer and the regulatory elements controlling gene expression are largely found within the noncoding portion of the genome,[9] methods to link noncoding somatic alterations within an individual's tumor to changes in gene expression would not only vastly advance our knowledge of cancer biology but also be a great aid in interpreting specific cancer genomes.

Perhaps the most commonly applied approaches to identify functional somatic mutations in cancer search for mutations that recur across tumor samples more than expected by chance. In the context of noncoding mutations, such approaches may additionally integrate functional impact scores or diverse functional genomics annotations relevant for gene regulation.[10–12] Overall, while recurrence-based approaches have proven to be highly successful in identifying cancer-relevant mutations within coding regions,[13,14] they have had limited success in uncovering such mutations within noncoding regions.[6] The best understood of these are mutations within the *TERT* promoter, which occur in about 30% of melanomas and are relatively common in other cancers and are associated with *TERT* overexpression.[15,16] However, further recurrence analysis across hundreds of deeply sequenced samples has only uncovered a handful of other candidate causal mutations in cancer.[17–20]

A parallel approach is to link somatic noncoding mutations to changes in gene expression. Samples with and without mutations within a proximal upstream region of a specific gene can be tested for association with different expression levels for that gene.[21,22] Mutation and gene pairs can also be evaluated in the context of known functional enhancers and promoters allowing the inference of more distally mutated loci associated with changes in expression in cancers.[23–25] However, in order to infer statistically significant differences in gene expression, these types of approaches also require that noncoding regulatory regions for a gene are recurrently mutated across individuals.

Our work begins with the observation that functional noncoding somatic mutations within regulatory regions should change the expression of one copy of the target gene without affecting the expression of the allele on the other chromosome. Thus, we aim to identify genes where only the expression of one allele is changed in cancer cells in order to hone in on *cis* noncoding mutations that lead to regulatory changes that are functional in cancers. However, unequal expression levels of alleles of a gene, or allele-specific expression (ASE), is a relatively common phenomenon in both healthy tissues and cancer.[26–29] While ASE has been used to measure the impact of cis-eQTLs across healthy tissues,[30] in the context of cancer, ASE is often detected in more than 25% of all genes,[27,28,31] making it difficult to apply for detecting either genes dysregulated in cancers due to alterations in *cis* or underlying functional noncoding somatic mutations.

To help narrow the search for genes whose regulation is affected by changes in *cis*, we introduce the concept of *differential ASE*, where the ASE of a gene within cancer cells is different from its ASE in paired

normal cells from the same individual and tissue type. Differential ASE aims to find dysregulated genes within an individual by directly comparing the level of ASE in a tumor sample to the level of ASE at the same heterozygous site in a matched normal sample. In this way the only genomic sequence differences between the paired samples are cancer related, and thus changes in ASE are as well. However, detecting ASE in tumor cells is complicated by intra-tumor heterogeneity. We thus introduce a series of increasingly complex models to detect differential ASE from gene expression data, including two models to deconvolve expression measurements derived from tumor samples in order to estimate the contributions from cancer cells.

Previous studies of ASE in cancer have not directly compared ASE levels for genes in tumors with their matched normal counterparts. For example, while genes with different levels of ASE have been observed when comparing cell lines derived from the germlines of those with familial pancreatic cancer and control samples, ASE in tumor cells was not considered.[32] An alternate approach which considered somatic coding mutations and uncovered those mutations that showed preferential ASE in tumor samples did not consider whether genes already exhibited ASE in that individual's normal tissue.[33] Another approach has been to test whether genes that have significant ASE in cancer tissues also have significant ASE in matched normal tissues in order to classify the effect as tumor specific or shared.[34] However, this merely determines if a gene as a whole exhibits significant ASE in one context but not the other. Our differential ASE approach is, to the best of our knowledge, the first that directly compares tumor ASE levels to normal ASE levels in matched samples to compute the amount of differential ASE and does so while also accounting for tumor heterogeneity.

We apply our approach to detect genes that exhibit differential ASE to paired breast cancer and normal samples profiled by TCGA.[3] First, as validation, we demonstrate that the dysregulation caused by copy number variation and nonsense mediated decay results in differential ASE. Next, we observe that genes that recurrently exhibit differential ASE include known cancer drivers and fall in cancer-related pathways. Furthermore, estrogen receptor positive tumors tend to show differential ASE of estrogen receptor genes. We then focus in on noncoding mutations derived from whole-exome sequencing data, as very few of the individuals with paired expression profiles have whole-genome sequencing data. We find that genes with *cis* noncoding mutations are enriched for differential ASE. Of the observed *cis* noncoding mutations, 96.2% do not result in differential ASE, and thus can be filtered out as they are highly unlikely to be functional. Of the remaining mutations, we find 147 mutations that occur in

annotated regulatory sites. We extend this analysis by looking at whole-genome sequencing data in four individuals and identify an additional five mutations within annotated regulatory sites that are upstream of genes that exhibit differential ASE. We conclude that differential ASE is a powerful approach to hone in on functional *cis* noncoding mutations.

Results

Overview of Differential Allele-Specific Expression

We define the differential ASE of a gene as the difference between its estimated ASE in cancer and paired normal tissue samples. Briefly, rather than computing ASE at each heterozygous site by comparing the fraction of RNA-seq reads in tumor samples that contain a specific nucleotide to an expected fraction of one half (as done in standard models to estimate ASE), we consider the observed fraction at the same site in a matched normal sample. Because tumor samples are heterogeneous and contain both cancer and normal cells, estimating ASE in cancer cells is challenging and we introduce three models for this task (see Figure 1). In the first model, diffASE-baseline, we directly compute the difference between the fraction of transcripts with the specific nucleotide in tumor and normal. In the second model, diffASE-purity, we use estimated tumor purity to deconvolve expression data from the tumor sample into proportions arising from cancer cells and normal cells. In the third model, diffASE-exp, we adjust our differential ASE estimates to account for differential expression between cancer and normal cells. For each model, we compute a per-site *p*-value to determine if the computed fraction of reads in cancer cells is significantly different from the fraction in normal cells, given the number of reads estimated to arise from cancer cells. Simulations (see Supplement, Section A) show that the models can successfully detect differential ASE in realistic settings for sample purity and read depth and that more complex models tend to better estimate differential ASE. Additionally, simulations predict that differential ASE is easier to detect in cases where cancer genes are overexpressed. Finally, because TCGA data is unphased, to obtain gene-level estimates of differential ASE using any of the three models, we take a read-depth weighted median value over all its heterozygous sites. Per-gene significance values are obtained by combining *p*-values across all sites within a gene using Fisher's method and computing false discovery rates (FDRs) via the Benjamini-Yekutieli procedure.[35] See Methods for more details.

The Landscape Of Differential Allele-Specific Expression in Breast Cancer

In the TCGA breast cancer data set, 91 individuals have matched RNA-seq data for both tumor and normal samples, along with copy number variation (CNV) and whole-exome sequencing data. Since

immune cells can infiltrate tumors and may confound our analyses, we used Cibersort[36] to remove tumor samples with significant expression correlation to immune cell types (*p*-value < 0.05). We ran the models on all genes for the remaining 46 individuals as described in Methods. While there are many gene-sample pairs that are considered to exhibit significant differential ASE by all three of our models, each model selected a slightly different set as significant across the individuals (Figure 2a).

To incorporate all three models, we consider a gene to exhibit significant differential ASE in an individual if in at least one model it has an FDR < 0.1 and an estimated differential ASE > 0.15. We found that across individuals, the median fraction of genes where ASE could be determined (i.e., with at least one heterozygous site) that exhibit significant differential ASE is 2.18%. In contrast, across individuals, a median of 47.0% of genes are significant when using standard ASE with the same FDR and ASE effect size thresholds (Figure 2b). The discrepancy between the number of significant genes when using standard ASE versus differential ASE indicates that the majority of ASE observed in a tumor sample by the conventional method is likely also present in the matched normal. In contrast to differential expression analysis, which considers a collection of samples and typically uncovers numerous differentially expressed genes between tumor and normal samples[37], differential ASE has specificity in detecting regulatory events in cancer and is able to detect per-individual regulatory events.

While a gene may exhibit differential ASE due to somatic alterations in its regulatory regions, it may also arise for other reasons, including nonsense mediated decay, aberrant methylation, or CNVs where amplifications or deletions of the gene on one copy lead to changes in expression for that copy. These situations provide a useful validation for our approach to uncover differential ASE. CNVs in particular have a large impact on ASE.[28] We observe that genes affected by CNVs have higher differential ASE than other genes (Supplement, Section B) and are enriched in the set of gene-sample pairs with significant differential ASE (one-sided hypergeometric *p*-value = 6.12e-37), confirming that our models can detect expected allele-specific changes in expression resulting from amplifications and deletions. Further, we find that, across genes with at least one heterozygous site, a per-sample median of 4.31% of genes with CNVs are significant across samples while only 1.83% without CNVs are significant. Our approach appears to be equally effective at detecting differential ASE arising from either amplifications or deletions: across samples, of genes with CNVs, a median of 4.46% of genes within regions of amplification and 4.22% of genes in regions containing deletions are significant (Figure 3a).

Differential Allele-Specific Expression Captures Effects of Coding Mutations

To examine the impact of mutations on differential ASE, we looked at coding mutations occurring in gene-sample pairs unaffected by CNVs. Nonsense mutations are expected to reduce the number of transcripts from the copy of DNA they appear on due to nonsense mediated decay.[38,39] We find that genes-sample pairs that exhibit significant differential ASE are 1.81-fold enriched for nonsense mutations as compared to the baseline fraction of all gene-sample pairs containing nonsense mutations (Figure 3b, dark purple), which represents a statistically significant level of enrichment (one-sided hypergeometric p-value = 6.67e-5). As a comparison, significant gene-sample pairs are 0.98-fold under-enriched for those pairs that do not have mutations in exome sequencing data (Figure 3b, dashed line). Furthermore, it has been shown that nonsense mediated decay is less likely to occur if a nonsense mutation occurs within 50 bases of the 3'UTR.[38,39] In agreement with this, of the 28 gene-sample pairs where such mutations are observed, we find only one with significant differential ASE; this is a 1.22-fold enrichment for such mutations (Figure 3b, light purple) and is not significantly different from random (one-sided hypergeometric p-value = 0.17).

Next we considered the effect of missense and silent mutations on differential ASE. The effect of these two mutation types in cancer on ASE has previously been observed.[5,39] In order to examine the effect of each given mutation type exclusively, we only consider gene-sample pairs that contain that type of mutation in an evolutionarily conserved site and no other mutations in conserved sites, except for intron mutations, of which there are too many of to exclude. We find that there is a modest 1.23-fold enrichment for gene-sample pairs with missense mutations (Figure 3c, red) and a lower 1.17-fold enrichment for gene-sample pairs with silent mutations (Figure 3c, blue) among all gene-sample pairs exhibiting significant differential ASE. The enrichment for gene-sample pairs with missense mutations is significant (one-sided hypergeometric p-value = 3.29e-4) while the enrichment for gene-sample pairs with silent mutations is not (one-sided hypergeometric p-values = 0.064). Interestingly, among all gene-sample pairs with significant differential ASE, the fraction with missense mutations does not differ significantly from the fraction with silent mutations (one-sided Binomial p-value = 0.39), suggesting that these mutations may affect ASE by impacting transcript stability.

Known Cancer Genes Recurrently Exhibit Differential Allele-Specific Expression

One of the most powerful techniques to identify cancer-relevant events is to consider recurrence across samples so we next examined genes that recurrently exhibit significant differential ASE across samples.

A gene recurrently exhibiting differential ASE at least six times across samples is statistically significant (Benjamini-Hochberg corrected one-sided Poisson binomial FDR < 0.1). Due to the strong effect of CNVs, we examined recurrence separately across genes without CNVs and those with, in both cases finding many cancer-related genes (Figure 4a, genes ordered by non CNV recurrence left and by CNV recurrence right). Among those without CNVs, among the top 10 most recurrent genes, two are known cancer genes from the Cancer Gene Census,[40] a significant enrichment (hypergeometric *p*-value = 0.02) and 7 show significant differential expression between cancer and normal when considering only the set of individuals that exhibit significant differential ASE in that gene, despite the fact that significance is difficult to obtain with such small numbers of samples. We find that the oncogene *MDM2* is significantly overexpressed in samples showing significant differential ASE (edgeR[41] *p*-value = 1.5e-3) indicating that the observed dysregulation may be contributing to cancer development. Among the other genes, *RABEP1* promotes breast cancer growth,[42] *GSR* is involved in oxidative stress response, *RPA1* is a component of DNA repair,[43] *MREG* is a functional partner of the *RAB27B* oncogene,[44] and *NUP88* has been linked to cancer progression.[45] Among the recurrently significant genes with CNVs, *NAT1* is known to activate carcinogens, and all 6 recurrences are amplifications,[44] while *ASAH1* is thought to play a role in cancer progression, and all 6 recurrences are deletions.[43] *ASAH1* additionally exhibits significant differential ASE in four samples that do not have CNVs in that gene. *KMO* is thought to play a role in cancer by preventing apoptosis, and all 6 recurrences are amplifications.[46] *NSL1* is an important component of cell division and interacts with *CASC5* (Cancer Susceptibility Candidate 5).[44] Finally, *PRUNE* is known to play a role in cell proliferation, and all 6 recurrences are amplifications.[43] It also exhibits significant differential ASE in two additional samples without CNVs.

To evaluate the list of all genes recurrently exhibiting differential ASE (for the full list, see Supplement, Section C) we looked for enrichment in cancer-related gene sets. We used gene set enrichment analysis on the list of genes ranked by recurrence for all pathways from Reactome as well as all diseases from the Disease Ontology.[47] The most significant pathway is "stabilization of p53" an important pathway for cancer development. Other relevant significant pathways include "apoptosis" as well as specific cancer related pathways such as "Autodegradation of Cdh1" and "APC-Cdc20 mediated degradation of Nek2A" (for a full list of significant pathways see Supplement, Section D). The top nine Disease Ontology gene sets are all cancers and breast cancer appears in the full list of significant terms (for a full list of significant disease ontologies see Supplement, Section D). These results indicate that genes recurrently exhibiting differential ASE are broadly involved in cancer processes.

Since breast cancer is subtyped partly based on the overexpression of estrogen receptors, we examined differential ASE in subtyped samples to determine the cause of dysregulation in those samples, as noncoding mutations may play a role.[48] Among the 22 individuals who were clinically subtyped as estrogen receptor positive (ER+) and had higher levels of alpha estrogen receptor gene (*ESR1*) in their cancer cells than in their normal cells, 5 individuals exhibited significant differential ASE for ESR1, while another 2 had amplifications due to a copy number alteration; this suggests that for a notable fraction of ER+ individuals, the overexpression of their *ESR1* gene is due to *cis*-regulatory alterations. Further, all samples with significant differential ASE in *ESR1* overexpressed it. We note that amongst the ER- patients, none exhibited differential ASE for ESR1. Finally, to determine if dysregulation was broadly similar among ER+ patients, we computed the distances between all individuals across genes by measuring the Euclidean distance between a binary vector for each individual of whether each gene was significant with respect to differential ASE or not in that individual. We find that the differential ASEs across genes between ER+ individuals are significantly more similar to each other than to ER- individuals (Wilcoxon *p*-value = 6.56e-6) suggesting that similar sets of genes are altered in *cis* across individuals within subtypes.

Functional Noncoding Mutations Lead to Differential Allele-Specific Expression

To measure the impact of noncoding mutations on differential ASE, we looked for enrichment in the presence of promoter, 5'UTR, 3'UTR, 5'Flank, and intronic mutations among gene-sample pairs exhibiting significant differential ASE. As with coding mutations, in order to examine the effect of each given mutation type exclusively, we only consider gene-sample pairs that contain that type of mutation in an evolutionarily conserved site exclusively, with the exception of mutations in introns. We find that gene-sample pairs with promoter mutations are 1.23-fold enriched in gene-sample pairs exhibiting significant differential ASE (Figure 4b, brown), which is not a statistically significant enrichment given the number of such mutations (one-sided hypergeometric *p*-value = 0.079). However, if we filter the set of promoter mutations to only those falling in sites annotated as regulatory by the curated site ORegAnno,[49] we observe a much larger 1.44-fold enrichment (Figure 4b, orange) that is statistically significant (one-sided hypergeometric *p*-value= 0.034). Moreover, when we restrict to promoter mutations that fall into sites that are not annotated as regulatory, we see only 1.06 fold enrichment (hypergeometric *p*-value= 0.25). When we consider gene-sample pairs with a mutation in their 5'UTR, we observe the strongest enrichments within gene-sample pairs with significant differential ASE (2.16-

fold enrichment, one-sided hypergeometric *p*-value = 4.59e-4, Figure 4c, green). We also observe a large enrichment for differential ASE in gene-sample pairs with mutations in the 3'UTR (1.73-fold enrichment, one-sided hypergeometric *p*-value = 1.32e-5, Figure 4b, blue), a small but significant enrichment for gene-sample pairs with mutations in their introns (1.16-fold enrichment, one-sided hypergeometric *p*-value = 2.56e-4, Figure 4c, purple), and no enrichment for those with mutations in the 3'Flank (1.03-fold enrichment, one-sided hypergeometric *p*-value = 0.20, Figure 4c, red). In contrast, when using standard ASE, of the noncoding mutations, only 3'UTR mutations had a very small but significant enrichment among gene-sample pairs that are significant (Supplement, Section E), indicating that differential ASE is crucial to measuring the effects of noncoding mutations.

One of the most powerful features of differential ASE is that it serves as a means to distinguish which *cis* noncoding mutations have the potential to cause dysregulation. Mutations *cis* to genes that have no observed differential ASE are not leading to dysregulation of that gene and thus can be filtered out as potential functional noncoding mutations. This is important because there are numerous noncoding mutations in any given sample, most of which have no functional impact. Among the samples we studied, there are 23,400 noncoding mutations observed in whole-exome sequencing data in conserved sites. However, only 769 of these are *cis* to genes with observed differential ASE (for a full table of counts by mutation type see Supplement, Section F). Filtering out 96.2% of mutations represents an enormous decrease in the search space for functional noncoding mutations. In comparison, standard ASE is only able to filter 46.3% of conserved noncoding mutations.

To generate a list candidate functional noncoding mutations, we consider only those that occur *cis* to genes with significant differential ASE at FDR < 0.1 and occur within a regulatory site as annotated by ORegAnno.[49] We note that multiple mutations can be implicated by the same gene, and thus to minimize alternative causes of dysregulation, we exclude any mutation for which the sample had a CNV or nonsense mutation in the same gene. These stringent requirements give us a list of nine promoter mutations, 15 5'UTR mutations, 15 3'UTR mutations, three 3'Flank mutation, and 105 intronic mutations (for a full list of mutations see Supplement, section G). Among these noncoding mutations, 5.6% are associated with known cancer genes from the CGC,[40] which is significantly enriched relative to the background rate of CGC genes (hypergeometric *p*-value = .011). This list also includes many other cancer relevant genes such as *PGRMC1*,[50] *RHOB*,[51] *HYOU1*,[52] *INPPL1*,[43] *GADD45B*,[43] and *CDKN1B*.[53]

We next sought to determine if there were potential functional mechanisms linking the effect of the observed mutations to changes in ASE. We tested all our mutations for their predicted regulatory effect using deepSEA.[54] We find that 22% of promoter mutations, 47% of 5'UTR mutations, 27% of 3'UTR mutations, two of the three 3'Flank mutation, and 8.6% of intronic mutations have a significant predicted functional impact. This indicates that a large proportion of our candidate mutations have the potential to affect gene regulation. We additionally used FIMO[55] with a set of transcription factors from HOCOMOCO[56] to test if our set of mutations could disrupt or create new transcription factor binding sites. We found that eight of the nine promoter mutations cause at least one change in transcription factor binding sites. These results indicate that a significant fraction of the mutations we observe *cis* to genes showing significant differential ASE have the potential to cause the observed dysregulation.

Finally, to assess the power of differential ASE as a method for whole-genome patient-specific analysis, we examined mutations in the promoters of four samples that have whole-genome sequencing data available. While these individuals are also included in the whole-exome sequencing data, using whole-gene sequencing allows us to look further upstream from genes to detect additional mutations. Among these four samples, we find 451 promoter mutations in conserved sites within 2000 bases upstream of transcription start sites. Only 10 of these mutations are upstream of genes with differential ASE. When we limit these mutations to know regulatory regions as annotated by ORegAnno, we observed five potentially functional mutations (for a full list of mutations see Supplement, Section H). These appear upstream of KIF1B, TNFAIP8, PNN, FCF1, and PUDP, the first three of which have previously been associated with cancer.[57–59] Furthermore, four out of these five mutations are predicted by deepSEA to have a significant functional impact. Thus, the powerful filtering provided by differential ASE makes it possible to focus on only the subset of mutations that are potentially functional in whole genome sequencing.

Discussion

We have shown that differential ASE is a powerful tool for honing in on genes that are dysregulated due to somatic *cis* alterations. We have demonstrated that genes that recurrently exhibit differential ASE across samples are cancer related, and that we can detect the effects of CNVs, nonsense mediated decay, coding mutations, and noncoding mutations. While ASE has previously been analyzed in cancer genomes, our work suggests that many genes exhibit ASE in cancer cells but that most of these do so

due to their expression patterns in normal cells. Standard ASE by itself is thus not an effective means to analyze cancer genomes.

Differential ASE analysis identifies 147 potentially functional noncoding mutations without relying on recurrence across individuals. While these mutations are strong candidates in terms of functional impact, they do not overlap any previously identified functional noncoding mutations in breast cancer. However, none of the individuals included in our dataset had a *TERT* promoter mutation or the previously described functional recurrent promoter mutations in breast cancer, such as the ones in the FOXA1 promoter.[15,17,20,60] This is not surprising due to the low levels of recurrence of even the most well studied noncoding somatic mutations. As a result, the overlap in predicted functional mutations by different methods is expected to be small, and further development of methods such as ours that are not based on recurrence is critical.

We introduced three models to detect differential ASE, each of which takes a different approach to handle the heterogeneity observed in tumors. These three models uncover complementary sets of gene-sample pairs exhibiting differential ASE, and in the future, it may be possible to develop even more sensitive methods to deconvolve mRNA expression. For instance, not all cancer cells carry any given mutation. Models could account for the Variant Allele Frequency (VAF) of any given mutation to adjust the fraction of mRNA that is expected to derive from cells with that mutation. Further, models could consider tumor mutation phylogenetic trees, as earlier occurring mutations are more likely to be those that drive the progression of cancer.[61]

One important caveat to our work is that there are many events that can cause expression dysregulation and we were only able to consider a small fraction of them. For example, genes that recurrently exhibit differential ASE may be subject to distal *cis* regulatory events; the use of known enhancers or Hi-C contact maps may be helpful in constraining the sets of mutations that need to be considered.[62] Alternatively, it is known that there are epigenetic factors that can lead to ASE.[34] Furthermore, even if we find that a gene is dysregulated and suggest observed *cis* mutations as potentially causative, there may be multiple detected *cis* mutations and additional analysis is necessary to detect which of those are causing the change in ASE.[60] Nevertheless, because of the large number of noncoding mutations in cancer genomes, it is vital to have techniques for filtering out those mutations that are unlikely to be

playing a regulatory role; indeed, we have found that the vast majority of noncoding mutations near genes do not appear to alter their expression.

Differential ASE analysis in cancer necessitates the use of a matched normal sample, and thus this analysis may be better suited to some cancer types than others. We chose to analyze TCGA breast cancer samples because they comprise the only large data set where, across the same individuals, there is matched tumor-normal RNA-seq data and copy number variant calling, along with either whole-exome or whole-genome sequencing. Just as sequencing the germline is essential for uncovering somatic alterations in cancers, further studies that obtain expression information for both tumor and matched normal samples would be especially helpful in order to facilitate methods such as ours that control for variation in the normal sample. Still ongoing large-scale cancer sequencing efforts such as ICGC have the potential to generate such data.[4] Looking further ahead, widespread analysis of differential ASE without matched tumor-normal samples may be enabled by single-cell sequencing, as it should be possible to identify normal cells within a tumor sample.

Overall, our results suggest that differential ASE is a powerful way to detect recurrent dysregulation and to hone in on function *cis* noncoding mutations. Our observation that the majority of ASE detected in tumor samples is also present in the matched normal indicates that differential analysis is fundamental to examining dysregulation in cancer. Finally, the detection of differential ASE does not rely on observing recurring mutations across samples, and thus we believe that it could be used as a powerful tool for personalized medicine by identifying patient-specific functional noncoding mutations.

Methods

Estimating per-site differential allele-specific expression

In each individual, for each gene, we wish to compute its differential ASE, or the difference between the gene's ASE in cancer cells and its ASE in normal cells. Since TCGA data is unphased, we first estimate differential ASE separately for each heterozygous site within the gene, and then combine these estimates. We propose three models with increasing complexity for the task of estimating differential ASE at a particular heterozygous site i (for a fixed gene within an individual) using paired RNA-seq data from tumor and normal samples.

At each heterozygous site, we refer to the variant that appears more frequently in the normal sample transcripts as the primary nucleotide. Let $R_{c,i}$ be the fraction of all RNA-seq reads mapped to site i that contain the primary nucleotide in cancer cells in the tumor sample. We want to compare this to $R_{n,i}$ the fraction of all RNA-seq reads from normal cells that contain the same nucleotide at site i. However, because the tumor sample is a heterogeneous mixture of normal and cancer cells, we can only directly compute $R_{s,i}$, the fraction of all RNA-seq reads in the tumor sample contain that the primary nucleotide at site i. The three models we propose estimate $R_{c,i}$, from $R_{s,I}$ in different ways and then use this value to compute differential ASE as $|R_{c,i} - R_{n,i}|$.

For our baseline model, **diffASE-baseline**, we compute $\Delta ASE_{baseline,i}$ by assuming that $R_{c,i} = R_{s,i}$.

Our next model, **diffASE-purity**, considers ρ, the purity of the tumor sample (i.e., the fraction of cells in a tumor sample that are cancer cells), and computes $\Delta ASE_{\rho,i}$. This model assumes that the fraction of RNA-seq reads at site i in the tumor sample that arise from cancer cells is proportional to ρ, and that the ASE of genes within normal cells within the tumor is the same as their ASE in the paired-normal cells. That is,

$$(1 - \rho)R_{n,i} + \rho R_{c,i} = R_{s,i}$$

and we can solve for $R_{c,i}$ and then use it to compute $\Delta ASE_{\rho,i}$.

Our final model, **diffASE-exp**, additionally considers changes in expression for a gene between paired normal and tumor samples, and computes $\Delta ASE_{exp,i}$. This model uses tumor sample and normal expression counts for a gene (e_s and e_n, respectively, measured in counts per million) to estimate f_t the fraction of all transcripts for that gene in the tumor sample that arise from cancer cells. We assume that normal cells in the tumor sample have the same expression level for the gene on average as in the paired normal. With this assumption, the transcripts expressed within a tumor sample are a combination of normal transcripts and cancer transcripts adjusted for purity. That is,

$$(1 - \rho)e_n + \rho e_c = e_s$$

We can compute the fraction of transcripts for the gene in a tumor sample that arise from cancer cells as

$$f_t = \frac{\rho e_c}{e_s}$$

or equivalently that

$$f_t = \frac{e_s - (1 - \rho)e_n}{e_s}$$

Then the RNA-seq reads in the tumor sample at site i within the gene arise from a mixture of normal and cancer transcripts weighted using f_t

$$(1 - f_t)R_{n,i} + f_t R_{c,i} = R_{s,i}$$

We can again solve for $R_{c,i}$ and use it to compute $\Delta ASE_{exp,i}$.

Estimating the per-site significance of differential ASE

For a heterozygous site i within an individual, we determine whether its estimated $R_{c,i}$ is significantly different from $R_{n,i}$ as follows. First, for each model, we estimate the total number of RNA-seq reads at site i arising from cancer cells, $T_{c,i}$, according to the assumptions of each model. For the diffASE-baseline model, we assume that $T_{c,i}$ is equal to the number of mapped reads at site i in the tumor sample. For the diffASE-purity model, we use ρ times the number of reads in the tumor sample. For the diffASE-exp model, we use f_t times the number of reads in the tumor sample. Next, for each model, using the estimated $R_{c,i}$ and $T_{c,i}$, we compute the number of reads with the primary nucleotide arising from cancer cells, and determine the probability of observing this number or a more extreme value. Our null model is $R_{c,i} = R_{n,i}$, and we use the normal approximation of the binomial distribution with a mean of $R_{n,i}T_c$ and a variance of $R_{n,i}(1 - R_{n,i})T_{c,i}$.

Per-gene estimates of differential ASE

For a given gene in an individual, for each model, we compute two gene level measures, one for statistical significance and one for effect size. For each individual, we only consider genes with at least one heterozygous site. To compute statistical significance for each gene, we combine the p-values at each heterozygous site using Fisher's method. Because sites with very high read depth would dominate

this measure, if a site has more than 100 reads mapped to it in the tumor sample, we downsample to a maximum of 100 reads. Similarly, if a gene has more than 10 heterozygous sites, we randomly sample ten of them ten times and take the median *p*-value of those randomizations. To estimate a gene-level differential ASE value, we take a depth weighted median of per site differential ASE, also considering a weight of at most 100 if there were more than 100 reads at a site.

For a particular method to estimate differential ASE, we say a gene has significant differential ASE if it is statistically significant at an FDR < 0.1 (Benjamini–Yekutieli method[35]) and has a gene-level estimated differential ASE > 0.15. The median number of significant genes changes by fewer than ten genes per sample for FDRs between 0.05 and 0.2 and weighted differential ASEs between 0.1 and 0.2 (Supplement, Section I). Finally, to combine models, we consider a gene to be significant overall if is significant in at least one model. We note that we found no statistically significant difference between gene-sample pairs with significant differential ASE and those without with respect to of the number of heterozygous sites across the length of the gene (one-sided Wilcoxon *p*-value = 0.86).

TCGA data acquisition and processing

In the Cancer Genome Atlas (TCGA)[3] breast invasive cancer (BRCA) cohort, 91 individuals have matched RNA-seq data for both tumor and normal samples, along with copy number variation and exome sequencing data. We downloaded somatic mutation calls identified by Mutect2[63] using the exome data. Across the 91 individuals we analyzed, Mutect2 identified 103,748 coding mutations and 303,415 noncoding mutations (for a full table of mutation counts by type, see Supplement, Section F). Of the samples we examined, four additionally have whole genome sequencing (WGS) data. We called mutations in the WGS data using Mutect2 with default parameters. For the WGS data, we considered promoter mutations to be any occurring within 2000bp upstream of a gene.

We used the *TxDb.Hsapiens.UCSC.hg38.knownGene* package in R[64] to download the locations of all annotated human genes excluding those on the Y chromosome. For each of these 22,732 genes, for each individual we considered all RNA-seq reads from the solid tissue normal BAM files that mapped uniquely to that gene (mapq quality code 255). For RNA-seq data, in cases where two vials were processed for a sample, we always used vial A. For each individual and gene pair, we used the *AllelicImbalance* package in R[65] to find every site in the gene that had at least 20 mapped reads of which the more frequent nucleotide is at most 90 percent, filtering out those sites that had a somatic mutation

observed in that individual. These sites comprise our collection of heterozygous sites for the gene in the individual. Next, for each individual and gene pair, for each heterozygous site identified in the gene in that individual, we counted both the total number of reads at each heterozygous site in both tumor and normal, as well as a count for the number of times the primary nucleotide appeared in normal and that same nucleotide appeared in the tumor.

For each individual and gene pair, we used its tumor and matched normal expression count data as measured in count files to compute the corresponding expression values in Counts Per Million (CPM) using *edgeR*[66] with library sizes normalized by the Trimmed Mean of M-values (TMM). Masked copy number segment files from TCGA for both tumor and matched normal samples were obtained. For each gene-individual pair, we calculated the average segmented copy number score across the gene for both tumor and normal samples. For an individual, a gene is identified as falling into an amplified (respectively, deleted) region in the tumor if the $\log_2$-fold change in average segmented copy number score is > 0.5 (respectively, < -0.5). For each tumor sample, purity estimates were obtained from Aran et al.[67] In particular, we used the combined purity estimate (CPE) when available, and for the samples where the authors did not provide a combined estimate, we used the median of the available purity measures that comprise their combined estimate.

For each tumor sample, we used the Cibersort webserver to determine the correlation of gene expression CPM values with the LM22 set of immune expression signatures.[36] We filtered out individuals whose tumors had a significant correlation with immune expression profiles at *p*-values < 0.05 leaving us with 46 individuals. In the Supplement, we consider results when filtering out larger numbers of individuals by considering tumor expression correlations with immune expression profiles at less stringent *p*-value thresholds (Supplement, Section I). We note that increasing the threshold to 0.1 only removes one additional individual.

Functional and genomic analysis

The Cancer Gene Census (CGC) from COSMIC[40] consists of 595 genes and comprises our list of known cancer genes. We compute Gene Set Enrichment using the package *ReactomePA* in R.[47] We compute differential expression using *edgeR*.[66] For our breast cancer subtype analysis, clinical data is obtained from the Broad GDAC Firehose,[68] and we classified any individuals with "estrogen receptor status" that was "positive" as ER+ and any with a status of "negative" as ER-.

We queried UCSCs genome browser[69] using *ucscTableQuery* via the *rtracklayer* package in R[70] to determine 20-way phyloP conservation score at the location of each somatic mutation.[71] Regulatory regions as annotated by ORegAnno 3.0 were obtained from UCSC's genome browser.[49] The functional impact of each noncoding mutation was assessed by the deepSEA webserver[54] using the specific location and the base change of the mutation. We considered a mutations to have a functional impact if it had a functional significance score < 0.01. To determine if noncoding somatic mutations disrupt transcription factor binding sites, we ran FIMO[55] with default parameters using a 40 base region around each mutation against consensus binding models from the HOmo sapiens COmprehensive MOdel COllection (HOCOMOCO) version 11.[56] We considered a mutation to change transcription factor binding if only one of either the germline sequence or the sequence resulting from the somatic mutation had a *p*-value < 1e-4 for a given motif.

We consider enrichment with respect to eight different sets of mutations (nonsense, missense, silent, promoter, 3'UTR, 5'UTR, 5'Flank, and intron as annotated by TCGA). For mutations other than nonsense mutations, we only consider mutations occurring within a location with a phyloP score of at least 0.1 and exclude mutations from the set if there is any other conserved non-intronic mutation in the same sample *cis* to or in the same gene because it would be unclear which mutation is contributing to the observed effect. Throughout, enrichment for a specific mutation type is computed as the fraction of gene-sample pairs exhibiting differential ASE that have the mutation as compared to the baseline fraction of gene-sample pairs with that type of mutation.

Author Contributions

PFP and MS designed the study. PFP performed the analysis. PFP and MS wrote manuscript. All authors read and approved the final manuscript.

Data availability

The data that support the findings of this study are publicly available on the TCGA data portal https://portal.gdc.cancer.gov/.

Code availability

All code used to generate results is available upon request.

Competing Financial Interests

The authors declare that they have no competing financial interests.

References

1.   Garraway LA, Lander ES. Lessons from the cancer genome. *Cell*. 2013;153(1):17-37. doi:10.1016/j.cell.2013.03.002

2.   Vogelstein B, Papadopoulos N, Velculescu VE, Zhou S, Diaz Jr. LA, Kinzler KW. Cancer Genome Landscapes. *Science (80-)*. 2013;339(6127):1546-1558. doi:10.1126/science.1235122

3.   TCGA Research Network. The Cancer Genome Atlas. http://cancergenome.nih.gov/.

4.   International Cancer Genome Consortium. International network of cancer genome projects. *Nature*. 2010;464(7291):993-998. doi:10.1038/nature08987

5.   Diederichs S, Bartsch L, Berkmann JC, et al. The dark matter of the cancer genome: aberrations in regulatory elements, untranslated regions, splice sites, non-coding RNA and synonymous mutations. *EMBO Mol Med*. 2016. doi:10.15252/emmm.201506055

6.   Khurana E, Fu Y, Chakravarty D, Demichelis F, Rubin MA, Gerstein M. Role of non-coding sequence variants in cancer. *Nat Rev Genet*. 2016;17(2):93-108. doi:10.1038/nrg.2015.17

7.   Lander ES, Linton LM, Birren B, et al. Initial sequencing and analysis of the human genome. *Nature*. 2001. doi:10.1038/35057062

8.   Craig Venter J, Adams MD, Myers EW, et al. The sequence of the human genome. *Science (80-)*. 2001. doi:10.1126/science.1058040

9.   The ENCODE Project Consortium. An integrated encyclopedia of DNA elements in the human genome. *Nature*. 2012;489(7414):57-74. doi:10.1038/nature11247

10.   Lochovsky L, Zhang J, Fu Y, Khurana E, Gerstein M. LARVA: An integrative framework for large-scale analysis of recurrent variants in noncoding annotations. *Nucleic Acids Res*. 2015. doi:10.1093/nar/gkv803

11.   Fu Y, Liu Z, Lou S, et al. FunSeq2: a framework for prioritizing noncoding regulatory variants in cancer. *Genome Biol*. 2014. doi:10.1186/s13059-014-0480-5

12.   Mularoni L, Sabarinathan R, Deu-Pons J, Gonzalez-Perez A, López-Bigas N. OncodriveFML: a general framework to identify coding and non-coding regions with cancer driver mutations. *Genome Biol*. 2016;17(1):128. doi:10.1186/s13059-016-0994-0

13.   Przytycki PF, Singh M. Differential analysis between somatic mutation and germline variation profiles reveals cancer-related genes. *Genome Med*. 2017;9(1):79. doi:10.1186/s13073-017-0465-6

14.   Lawrence MS, Stojanov P, Polak P, et al. Mutational heterogeneity in cancer and the search for new cancer-associated genes. *Nature*. 2013;499(7457):214-218. doi:10.1038/nature12213

15.   Vinagre J, Almeida A, Pópulo H, et al. Frequency of TERT promoter mutations in human cancers. *Nat Commun*. 2013;4. doi:10.1038/ncomms3185

16.   Huang FW, Hodis E, Xu MJ, Kryukov G V, Chin L, Garraway LA. Highly recurrent TERT promoter mutations in human melanoma. *Science*. 2013;339(6122):957-959. doi:10.1126/science.1229259

17.   Rheinbay E, Parasuraman P, Grimsby J, et al. Recurrent and functional regulatory mutations in breast cancer. *Nature*. 2017;547(7661):55-60. doi:10.1038/nature22992

18.   Weinhold N, Jacobsen A, Schultz N, Sander C, Lee W. Genome-wide analysis of noncoding regulatory mutations in cancer. *Nat Genet*. 2014;46(11):1160-1165. doi:10.1038/ng.3101

19.   Melton C, Reuter JA, Spacek D V., Snyder M. Recurrent somatic mutations in regulatory regions of human cancer genomes. *Nat Genet*. 2015;47(7):710-716. doi:10.1038/ng.3332

20.   Nik-Zainal S, Davies H, Staaf J, et al. Landscape of somatic mutations in 560 breast cancer whole-genome sequences. *Nature*. 2016. doi:10.1038/nature17676

21.   Fredriksson NJ, Ny L, Nilsson JA, Larsson E. Systematic analysis of noncoding somatic mutations and gene expression alterations across 14 tumor types. *Nat Genet*. 2014;46(12):1258-1263. doi:10.1038/ng.3141

22.   Gyorffy B, Pongor L, Bottai G, et al. An integrative bioinformatics approach reveals coding and non-coding gene variants associated with gene expression profiles and outcome in breast cancer molecular subtypes. *Br J Cancer*. 2018. doi:10.1038/s41416-018-0030-0

23.   Zhang W, Bojorquez-Gomez A, Velez DO, et al. A global transcriptional network connecting noncoding mutations to changes in tumor gene expression. *Nat Genet*. 2018;50(4):613-620. doi:10.1038/s41588-018-0091-2

24.   Hornshøj H, Nielsen MM, Sinnott-Armstrong NA, et al. Pan-cancer screen for mutations in non-coding elements with conservation and cancer specificity reveals correlations with expression and survival /631/67/69 /631/114 article. *npj Genomic Med*. 2018. doi:10.1038/s41525-017-0040-5

25.   Feigin ME, Garvin T, Bailey P, et al. Recurrent noncoding regulatory mutations in pancreatic ductal adenocarcinoma. *Nat Genet*. 2017. doi:10.1038/ng.3861

26.   Aguet F, Ardlie KG, Cummings BB, et al. Genetic effects on gene expression across human tissues. *Nature*. 2017;550(7675):204-213. doi:10.1038/nature24277

27.   Romanel A, Lago S, Prandi D, Sboner A, Demichelis F. ASEQ: Fast allele-specific studies from next-generation sequencing data. *BMC Med Genomics*. 2015;8(1). doi:10.1186/s12920-015-0084-2

28.   Mayba O, Gilbert H, Liu J, et al. MBASED: allele-specific expression detection in cancer tissues and cell lines. *Genome Biol*. 2014;15(8):405. http://genomebiology.com/2014/15/8/405.

29.   Spurr L, Li M, Alomran N, et al. Systematic pan-cancer analysis of somatic allele frequency. *Sci Rep*. 2018. doi:10.1038/s41598-018-25462-0

30.   Mohammadi P, Castel SE, Brown AA, Lappalainen T. Quantifying the regulatory effect size of cis-acting genetic variation using allelic fold change. *Genome Res*. 2017. doi:10.1101/gr.216747.116

31.   Ma X, Liu Y, Liu Y, et al. Pan-cancer genome and transcriptome analyses of 1,699 paediatric

leukaemias and solid tumours. *Nature*. 2018. doi:10.1038/nature25795

32. Aik CT, Fan JB, Karikari C, et al. Allele-specific expression in the germline of patients with familial pancreatic cancer: An unbiased approach to cancer gene discovery. *Cancer Biol Ther*. 2008;7(1):137-146. doi:10.4161/cbt.7.1.5199

33. Halabi NM, Martinez A, Al-Farsi H, et al. Preferential Allele Expression Analysis Identifies Shared Germline and Somatic Driver Genes in Advanced Ovarian Cancer. *PLoS Genet*. 2016;12(1). doi:10.1371/journal.pgen.1005755

34. Ongen H, Andersen CL, Bramsen JB, et al. Putative cis-regulatory drivers in colorectal cancer. *Nature*. 2014;512(1):87-90. doi:10.1038/nature13602

35. Benjamini Y, Yekutieli D. The control of the false discovery rate in multiple testing under dependency. *Ann Stat*. 2001. doi:10.1214/aos/1013699998

36. Newman AM, Liu CL, Green MR, et al. Robust enumeration of cell subsets from tissue expression profiles. *Nat Methods*. 2015;12(5):453-457. doi:10.1038/nmeth.3337

37. Koboldt DC, Fulton RS, McLellan MD, et al. Comprehensive molecular portraits of human breast tumours. *Nature*. 2012. doi:10.1038/nature11412

38. Lindeboom RGH, Supek F, Lehner B. The rules and impact of nonsense-mediated mRNA decay in human cancers. *Nat Genet*. 2016;48(10):1112-1118. doi:10.1038/ng.3664

39. Rhee J-K, Lee S, Park W-Y, Kim Y-H, Kim T-M. Allelic imbalance of somatic mutations in cancer genomes and transcriptomes. *Sci Rep*. 2017;7(1):1653. doi:10.1038/s41598-017-01966-z

40. Futreal PA, Coin L, Marshall M, et al. A census of human cancer genes. *Nat Rev Cancer*. 2004;4(3):177-183. doi:10.1038/nrc1299

41. Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics*. 2009;26(1):139-140. http://bioinformatics.oxfordjournals.org/cgi/doi/10.1093/bioinformatics/btp616%5Cnpapers2://publication/doi/10.1093/bioinformatics/btp616.

42. Yan GR, Xu SH, Tan ZL, Liu L, He QY. Global identification of miR-373-regulated genes in breast cancer by quantitative proteomics. *Proteomics*. 2011;11(5):912-920. doi:10.1002/pmic.201000539

43. O'Leary NA, Wright MW, Brister JR, et al. Reference sequence (RefSeq) database at NCBI: Current status, taxonomic expansion, and functional annotation. *Nucleic Acids Res*. 2016;44(D1):D733-D745. doi:10.1093/nar/gkv1189

44. Szklarczyk D, Morris JH, Cook H, et al. The STRING database in 2017: Quality-controlled protein-protein association networks, made broadly accessible. *Nucleic Acids Res*. 2017;45(D1):D362-D368. doi:10.1093/nar/gkw937

45. Agudo D, Gómez-Esquer F, Martínez-Arribas F, Núñez-Villar MJ, Pollán M, Schneider J. Nup88 mRNA overexpression is associated with high aggressiveness of breast cancer. *Int J Cancer*. 2004;109(5):717-720. doi:10.1002/ijc.20034

46. Wilson K, Auer M, Binnie M, et al. Overexpression of human kynurenine-3-monooxygenase protects against 3-hydroxykynurenine-mediated apoptosis through bidirectional nonlinear

feedback. *Cell Death Dis*. 2016;7:e2197. doi:10.1038/cddis.2016.87

47.   Yu G, He QY. ReactomePA: An R/Bioconductor package for reactome pathway analysis and visualization. *Mol Biosyst*. 2016;12(2):477-479. doi:10.1039/c5mb00663e

48.   Bailey SD, Desai K, Kron KJ, et al. Noncoding somatic and inherited single-nucleotide variants converge to promote ESR1 expression in breast cancer. *Nat Genet*. 2016;48(10):1260-1266. doi:10.1038/ng.3650

49.   Lesurf R, Cotto KC, Wang G, et al. ORegAnno 3.0: A community-driven resource for curated regulatory annotation. *Nucleic Acids Res*. 2016;44(D1):D126-D132. doi:10.1093/nar/gkv1203

50.   Neubauer H, Ma Q, Zhou J, et al. Possible role of PGRMC1 in breast cancer development. *Climacteric*. 2013;16(5):509-513. doi:10.3109/13697137.2013.800038

51.   Chen W, Niu S, Ma X, et al. RhoB acts as a tumor suppressor that inhibits malignancy of clear cell renal cell carcinoma. *PLoS One*. 2016;11(7). doi:10.1371/journal.pone.0157599

52.   Krętowski R, Stypułkowska A, Cechowska-Pasko M. Low-glucose medium induces ORP150 expression and exerts inhibitory effect on apoptosis and senescence of human breast MCF7 cells. *Acta Biochim Pol*. 2013;60(2):167-173.

53.   Chu I, Sun J, Arnaout A, et al. p27 Phosphorylation by Src Regulates Inhibition of Cyclin E-Cdk2. *Cell*. 2007;128(2):281-294. doi:10.1016/j.cell.2006.11.049

54.   Zhou J, Troyanskaya OG. Predicting effects of noncoding variants with deep learning-based sequence model. *Nat Methods*. 2015;12(10):931-934. doi:10.1038/nmeth.3547

55.   Grant CE, Bailey TL, Noble WS. FIMO: Scanning for occurrences of a given motif. *Bioinformatics*. 2011;27(7):1017-1018. doi:10.1093/bioinformatics/btr064

56.   Kulakovskiy I V., Vorontsov IE, Yevshin IS, et al. HOCOMOCO: Towards a complete collection of transcription factor binding models for human and mouse via large-scale ChIP-Seq analysis. *Nucleic Acids Res*. 2018;46(D1):D252-D259. doi:10.1093/nar/gkx1106

57.   Vitiello E, Ferreira JG, Maiato H, Balda MS, Matter K. The tumour suppressor DLC2 ensures mitotic fidelity by coordinating spindle positioning and cell-cell adhesion. *Nat Commun*. 2014. doi:10.1038/ncomms6826

58.   Day TF, Mewani RR, Starr J, et al. Transcriptome and Proteome Analyses of TNFAIP8 Knockdown Cancer Cells Reveal New Insights into Molecular Determinants of Cell Survival and Tumor Progression. In: *Methods in Molecular Biology*. ; 2017. doi:10.1007/978-1-4939-6539-7_7

59.   Yang X, Sun D, Dong C, Tian Y, Gao Z, Wang L. Pinin associates with prognosis of hepatocellular carcinoma through promoting cell proliferation and suppressing glucose deprivation-induced apoptosis. *Oncotarget*. 2016. doi:10.18632/oncotarget.9233

60.   Zhang Y, Manjunath M, Zhang S, Chasman D, Roy S, Song JS. Integrative genomic analysis predicts causative cis-regulatory mechanisms of the breast cancer–associated genetic variant rs4415084. *Cancer Res*. 2018;78(7):1579-1591. doi:10.1158/0008-5472.CAN-17-3486

61.   El-Kebir M, Satas G, Oesper L, Raphael BJ. Inferring the Mutational History of a Tumor Using Multi-state Perfect Phylogeny Mixtures. *Cell Syst*. 2016;3(1):43-53. doi:10.1016/j.cels.2016.07.004

62. Orlando G, Law PJ, Cornish AJ, et al. Promoter capture Hi-C-based identification of recurrent noncoding mutations in colorectal cancer. *Nat Genet*. 2018. doi:10.1038/s41588-018-0211-z

63. Cibulskis K, Lawrence MS, Carter SL, et al. Sensitive detection of somatic point mutations in impure and heterogeneous cancer samples. *Nat Biotechnol*. 2013;31(3):213-219. doi:10.1038/nbt.2514

64. Team BC, Maintainer BP. TxDb.Hsapiens.UCSC.hg38.knownGene: Annotation package for TxDb object(s). *R Packag version 340*. 2016.

65. Gådin JR, van't Hooft FM, Eriksson P, Folkersen L. AllelicImbalance: An R/ bioconductor package for detecting, managing, and visualizing allele expression imbalance data from RNA sequencing. *BMC Bioinformatics*. 2015;16(1). doi:10.1186/s12859-015-0620-2

66. Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics*. 2010;26(1):139-140. doi:10.1093/bioinformatics/btp616

67. Aran D, Sirota M, Butte AJ. Systematic pan-cancer analysis of tumour purity. *Nat Commun*. 2015;6. doi:10.1038/ncomms9971

68. Broad Institute TCGA Genome Data Analysis Center. Aggregate Analysis Features. *Broad Inst MIT Harvard*. 2016. doi:10.7908/C15Q4VFJ

69. Karolchik D. The UCSC Table Browser data retrieval tool. *Nucleic Acids Res*. 2004;32(90001):493D-496. doi:10.1093/nar/gkh103

70. Lawrence M, Gentleman R, Carey V. rtracklayer: An R package for interfacing with genome browsers. *Bioinformatics*. 2009;25(14):1841-1842. doi:10.1093/bioinformatics/btp328

71. Pollard KS, Hubisz MJ, Rosenbloom KR, Siepel A. Detection of nonneutral substitution rates on mammalian phylogenies. *Genome Res*. 2010;20(1):110-121. doi:10.1101/gr.097857.109

Figure 1: **Overview of Differential Allele-Specific Expression**. **a.** The differential ASE of a transcript in an individual's tumor considers both R_s, the fraction of gene transcripts in the tumor sample that contain a particular allele, along with R_n, the fraction in the matched normal sample. In this schematic, both the tumor and matched normal samples exhibit allelic imbalance. **b.** We develop three models to compute differential ASE at a heterozygous site i. Let $R_{s,i}$ and $R_{n,i}$ denote the fraction of all RNA-seq reads mapped to site i that have a given nucleotide at site i in the tumor sample and normal sample, respectively. The diffASE-baseline model (top) assumes that the tumor sample is entirely composed of cancer cells, thus $\Delta ASE_{baseline,i}$ is directly computed as the absolute difference between $R_{s,i}$ and $R_{n,i}$. The diffASE-purity model (middle) accounts for tumor samples being a mix of cancer and normal cells by using a tumor purity estimate ρ to deconvolve RNA reads in the tumor sample into reads arising from cancer cells and normal cells. Assuming that ASE estimated from the normal sample is a good approximation for ASE in the normal cells within the tumor sample, we can solve for $R_{c,i}$, the fraction of reads from cancer cells that contain the given nucleotide. $\Delta ASE_{\rho,i}$ is then the absolute difference between R_c and R_n. The diffASE-exp model (bottom) extends this concept by accounting for differential expression between cancer and normal cells. We assume that the normal cells in the tumor sample are on average expressed at the same levels as in the matched normal. This allows us to compute f_t, the fraction of RNA reads that are derived from cancer cells by subtracting away the ones that are derived from the normal cells in the sample. We then proceed as in the prior model using that fraction in the place of purity. For more details, see Methods.

Figure 2: **Landscape of Differential Allele Specific Expression in Breast Cancer. a.** Numbers of gene-sample pairs with significant differential ASE using each of the three methods. While there are many gene-sample pairs that are called significant by all three models, each model selected a slightly different set of genes as significant across the individuals. **b.** The distributions of genes exhibiting significant differential ASE per individual when estimating differential ASE between cancer and normal cells using any of the three models (blue) versus standard ASE computed using tumor sample RNA-seq data only (red). The median fraction of genes with at least one heterozygous site that exhibit significant ASE is 47.0%, whereas the median fraction of these genes exhibiting differential ASE is only 2.18%, indicating that the majority of ASE observed in a tumor sample by the conventional method is likely also present in the matched normal. For both panels, in order to be called significant, a minimum of differential ASE or ASE of > 0.15 is required along with an FDR < 0.1.

Figure 3: **Differential ASE Detects Known Causes of ASE. a.** The fractions of significant genes per individual when the genes contain amplifications, deletions, or no copy number variants. Across individuals, a median of 4.46% and 4.22% of genes with CNV amplifications and deletions, respectively, are significant while only 1.83% without CNVs are significant, confirming that our model can detect the expected allele-specific change in expression resulting from CNVs. **b.** Enrichment of gene-sample pairs that have nonsense mutations (dark purple) and the subset of nonsense mutations within 50bp of the 3'UTR (light purple) amongst all gene-sample pairs that exhibit significant differential ASE. For each mutation type, to the right of the bar, the number of gene-sample pairs that exhibit differential ASE and have that type of mutation is given. Enrichments with hypergeometric *p*-values < 0.05 are starred. Whereas there is a significant 1.81-fold enrichment for all nonsense mutations, nonsense mutations within 50bp of the 3'UTR exhibit a non-statistically significant 1.22-fold enrichment. Gene-sample pairs with no mutations have no enrichment amongst all genes with significant differential ASE (dashed vertical lines at 0.98). **c.** Gene-sample pairs with significant differential ASE are also enriched for missense mutations (red) and silent mutations (blue), but this enrichment is only significant for missense mutations.

Figure 4: **Regulatory Changes are Detected by Differential ASE. a.** Genes that most frequently exhibit differential ASE when ordered by recurrence across samples without CNVs (left) and ordered by recurrence across samples with CNVs (right). In both cases, the count for samples without CNVs is in blue and the count for samples with CNVs is in red. **b.** The enrichment (see Figure 3 caption) among gene-sample pairs exhibiting significant differential ASE for promoter mutations (brown) and the subset of promoter mutations that are annotated to regulatory regions (orange) as compared to no mutation (dashed line). **c.** The enrichment among gene-sample pairs exhibiting significant differential ASE for mutations in an intron (purple), 5'UTR (green), 3' flank (red), or 3' UTR (blue) as compared to no mutation (dashed line).