1

Light responses of melanopsin-expressing ganglion cells in the foetal mammalian retina

Jan Verweij[2, *], Shawnta Y. Chaney[2, *], Derek Bredl[2], Shruti Vemaraju[3], Gabriele M. König[6], Evi

Kostenis[6], Richard A. Lang[3, 4,5], David R. Copenhagen[1*]

[1]Departments of Ophthalmology and Physiology, University of California, San Francisco, San

Francisco, California Box 0444, 94158, USA

[2]Department of Ophthalmology, University of California, San Francisco, San Francisco,

California 94143, USA

[3]The Visual Systems Group, Abrahamson Pediatric Eye Institute, Division of Pediatric

Ophthalmology, Cincinnati Children's Hospital Medical Center, Cincinnati, OH 45229, USA

[4]Division of Developmental Biology, University of Cincinnati College of Medicine, Cincinnati, OH

45229, USA

[5]Department of Ophthalmology, University of Cincinnati College of Medicine, Cincinnati, OH

45229, USA

[6]Molecular, Cellular and Pharmacobiology Section, Institute of Pharmaceutical Biology,

University of Bonn, 53115 Bonn, Germany

*Co-first authors

Author contributions

DRC and RAL provided project leadership. JV, SYC, DRC and DB designed and performed

experiments and analysed data. DRC wrote the manuscript. JV, SYC, DB, SV and RAL

critiqued the study and edited the manuscript. GMK and EK provided crucial reagents and

advice.

Corresponding author:

David R. Copenhagen, Departments of Ophthalmology and Physiology, Box 0444, University

of California, San Francisco, San Francisco, California 94158, USA. Tel: 415-502-7351 (Office),

Fax: 415-476-6889. Email: cope@phy.ucsf.edu

Key points

- Melanopsin is a light-excitable photopigment expressed in a subset of ganglion cell neurons (mRGCs) in the retinas of many different species of vertebrates. In mature animals, light activation of mRGCs modulates many visual adaptive functions including pupil constriction, entrainment of circadian rhythms, mood and learning. In neonatal pups at ages prior to the developmental onset of visual signalling from rods and cones, melanopsin cells mediate photoaversive behaviour. In foetal pups, light activation of melanopsin cells accelerates maturation of the ocular vasculature. Here, we describe and physiologically characterize the light responses of melanopsin ganglion cells in the retinas of foetal pups.

- MRGCs in embryonic retinas respond to light at least four days prior to birth and exhibit responses to light of intensity as low as 3×10^{12} photons/cm^2/s.

- Phototransduction mechanisms include melanopsin activation of $G_{q/11}$ – G proteins, voltage-activated sodium currents, and voltage-gated L-type calcium currents.

- MRGCs are electrotonically coupled to other mRGCs in foetal retinas.

- We propose that melanopsin-expressing ganglion cells are excited by light while *in utero* and that this excitation relies, for the most part, on phototransduction pathways that have been described in postnatal retinas. Furthermore, we propose that foetal mRGCs have the requisite properties to modulate light-regulated maturation of the ocular vasculature and, perhaps, the development of visual pathways.

Abstract

Sensory stimulation plays a critical role in the maturation of sensory organs and systems. For example, when deprived of light before birth, foetal mouse pups *in utero* exhibit altered ocular vascular development. Normal vascular development depends on light excitation of melanopsin, a non-rod, non-cone photopigment that is expressed in a subset of ganglion cells (mRGCs) in the retina. However, there is no direct evidence that mRGCs in foetal eyes are light-responsive.

Very little is known about how light absorption leads to excitation in these foetal neurons. Using

mRGC-specific expression of the calcium indicators GCaMP3 and GCaMP6, we report that

foetal mouse mRGCs respond to light as early as 4 days before birth. Further, two distinct $G_{q/11}$-

G protein family antagonists, FR9000359 and YM-254890, abolish these light responses. TTX,

a blocker of voltage-activated sodium channels, reversibly represses light responses, and

FPL6417 and L-cis-diltiazem, which modify L-type calcium channels, respectively increase and

reduce light responses. Electrophysiological patch pipette recordings show that embryonic

mRGCs respond to light of intensity as low as 2.9 x 10^{12} photons/cm^2/s. The present findings

demonstrate a heretofore unproven but postulated light sensitivity in the retinas of foetal mice

and identify the transduction pathways involved. Surprisingly, mRGCs do not function as

completely independent photoreceptors but are electrotonically coupled with other mRGCs.

Given that melanopsin is expressed in foetal human retinas, these findings support the idea that

the eyes of foetal and early preterm infants are likely to exhibit functional photosensitivity. (241

words)

Introduction

Genetic programmes instruct the initial development of all organisms. Innate genetic

programming also regulates the maturation of the eye and the visual system. However, light

stimulation of maturing visual circuits also plays an important role in refining patterns of neural

connectivity and establishing various types of functionality (Breedlove, 2017). Most previous

studies have concentrated on the visually mediated refinement of visual centres at ages after

rods and cones in the retina begin responding to light to generate visual signals. However, a

unique class of intrinsically photosensitive melanopsin-expressing retinal ganglion cells

(mRGCs) that can generate visual signals in the absence of input from rods and cones has

been found in adult mammalian eyes (Berson et al., 2002; Do and Yau, 2010). In the eyes of

foetal mice, markers of melanopsin expression can be observed 9 days before birth and 19

days before the emergence of visual signalling from rods and cones. In human eyes, markers of

melanopsin expression appear at 8.5 weeks post-conception (31 weeks before birth) (Tarttelin

et al., 2003). The presence of genetic and immunohistological markers for melanopsin raises

the possibility that these mRGCs are intrinsically photosensitive and that light-mediated

signalling may occur in embryonic eyes. Thus, ambient light from the pregnant mother's

environment that penetrates to the eyes of foetuses might have the ability to control some

aspects of foetal development. Indeed, mouse foetuses that are light-deprived during the final

four days of gestation exhibit abnormal postnatal ocular vascular development (Rao et al.,

2013). Moreover, genetic deletion of melanopsin in foetal pups mimics the effects of light

deprivation (Rao et al., 2013). This finding suggests that ambient light passing through the

abdominal wall of the pregnant dam can activate melanopsin-expressing cells in foetal eyes and

trigger normal development of the ocular vasculature. A strong supporting argument in favour of

this hypothesis would be physiological evidence that mRGCs in the embryonic retina are light-

responsive. We report here that foetal mRGCs do respond to light.

Rods and cones in vertebrate retinas utilize opsin-based photopigments to capture light.

Rod and cone opsins couple to G-proteins belonging to the G_t (transducin) family to generate

photoresponses (Arshavsky et al., 2002). The photopigments in invertebrate photoreceptors,

unlike those in vertebrates, couple to non-transducin G-proteins. Rhodopsin in *Drosophila*

photoreceptors couples to G-proteins of the Gq family (Katz and Minke, 2009). Transduction of

light stimuli by melanopsin in postnatal mammalian retinas appears not to be coupled via G_t, but

the identity of the G-protein that mediates phototransduction remains controversial. Several

studies have demonstrated that mRGCs in postnatal mammals express one or more of four

different alpha subunits, *Gnaq*, *Gna11*, *Gna1* and *Gna15*, that are members of the $G_{q/11}$ gene

family (Chew et al., 2014; Graham et al., 2008; Hughes et al., 2015). Some studies have

suggested that melanopsin phototransduction is predominantly mediated by *Gnaq/11* coupling

(Hughes et al., 2015; Panda et al., 2005; Qiu et al., 2005). However, other reports concluded

that melanopsin functions independently of *Gnaq* and Gna*11* (Chew et al., 2014). In addition, in

expression systems, melanopsin can be coupled to $G_{i/o}$ (Bailes and Lucas, 2013) and to G_t

(Newman et al. 2003). Although the identification of melanopsin-G-protein coupling in postnatal

mice is somewhat controversial, a very recent study demonstrated that melanopsin-*Gnaq*

coupling is necessary and sufficient to trigger normal vascular development in the eyes of foetal

mice (Vemaraju et al. 2019). Direct measurement of light activity in mRGCs and

pharmacological intervention testing whether $G_{q/11}$ mediates the light responses of mRGCs in

foetal retinas and not merely ocular vascular development remains to be done.

We report here, using genetically encoded calcium indicator proteins expressed in

melanopsin cells, that embryonic mRGCs are light-responsive. We further show that

administration of the $G_{q/11}$ antagonists FR900359 and YM-254890 strongly attenuates these light

responses, supporting a role for melanopsin-mediated $G_{q/11}$ -coupled signalling in

phototransduction. These findings convincingly establish that the melanopsin-expressing

ganglion cells in retinas of foetal mice as young as embryonic day 15.5 are light-responsive.

Given that the *Opn4* gene is expressed in human eyes at gestational week 9 (Tartellin et al.

2003), these results also raise the very interesting possibility that light sensitivity may be

conserved in the human foetus and that it may endow the foetal eye with previously

unrecognized photosensitivity.

139   **Methods**

Animals

The experiments were performed in accordance with the policy of The Journal of

Physiology and the rules and regulations set forth in the NIH Guidelines for

the Care and Use of Laboratory Animals. The Institutional Animal Care and Use Committee of

the University of California, San Francisco (Animal Welfare Assurance Number: A3400-01)

approved the experiments. The experiments also conformed to the Society for Neuroscience

Policy on the Use of Animals in Neuroscience Research. The mice were housed in an AALAC-

accredited pathogen-free animal facility at UCSF with ad libitum access to food and water.

The experiments were performed on mice of either sex with an age ranging from embryonic day

15.5 (E15.5) to postnatal day 2 (P2). The mice were housed on a 12-h light-dark cycle with

lights on at 7 AM and off at 7 PM. The following mouse strains were used to breed the

experimental animals in this study: $Opn4^{cre}$ knock-in mice with Cre recombinase under the

control of the endogenous melanopsin promoter (Ecker et al., 2010); $Opn4^{GFP}$ transgenic mice

with the green fluorescent protein gene driven by the melanopsin promoter (Schmidt et al.,

2008); Ai14 strain mice with the floxed tdTomato gene in the rosa26 locus (Madisen et al.,

2010); C57Bl/6J wild-type mice (Jackson Laboratory, Sacramento, CA); melanopsin knockout

mice (Panda et al., 2003); and finally Ai38(B6.Cg-$Gt(ROSA)26Sor^{tm38(CAG-GCaMP3)Hze}$/J) and AI96

(B6;129S6-Gt(ROSA)26Sor$^{tm96(CAG-GCaMP6s)Hze}$/J) strains with the floxed GCaMP3 and GCaMP6

genes, respectively, which encode two different calcium sensors, in the rosa26 locus (Jackson

Laboratory, Sacramento, CA; Zariwala et al., 2012). Genotyping of $Opn4^{Cre}$ (Ecker et al., 2010),

$Opn4^{-}$ (Panda et al., 2002), $Ai14$ (Madisen et al., 2010) and $Ai38$ (Zariwala et al., 2012) mice

was performed as described in the referenced articles.

Tissue preparation

Pregnant females used in the calcium measurements were sacrificed by CO_2 inhalation followed

by cervical dislocation. For the electrophysiological experiments, pregnant females were

sacrificed by cervical dislocation. Following laparotomy, the uterine horns were removed and

transferred to cold ACSF (119 mM NaCl, 24 mM NaHCO$_3$, 1.25 mM NaH$_2$PO$_4$, 2.5 mM KCl, 2.5

mM CaCl$_2$, 1.5 mM MgSO$_4$, 10 mM glucose, 2 mM Na pyruvate, and 2 mM Na lactate) that had

been previously bubbled with carbogen (95%O$_2$/5% CO$_2$). The foetuses were dissected free of

their extra-embryonic membranes and decapitated, and the eyes were removed and placed in

carbogen-bubbled ACSF. For calcium imaging, the retina was freed from each eye under a

dissecting microscope, cut radially and teased, photoreceptor side down, onto a small piece of

plastic film; it was then flattened with a brush consisting of a single cat whisker while the plastic

film was dragged out of the solution on a microscope slide "boat ramp". The plastic film was

lowered upside down onto a polylysine-coated (0.1 mg/mL) glass coverslip (Number 1, round,

12-mm diameter), and the plastic film was floated off the coverslip using a drop of ACSF. These

steps were performed under ambient room light at room temperature (RT). The coverslips were

kept in carbogen-bubbled ACSF (RT) in darkness until they were mounted in the recording

chamber. For each experiment, a coverslip was placed in the recording chamber (Warner

Instruments, Hamden, CT) and perfused with carbogen-bubbled ACSF at ~35°C flowing at 0.4

to 0.8 mL per minute. For patch-clamping, retinas were dissected and kept in carbogen-bubbled

ACSF. Immediately before each experiment, a radially cut retina was teased ganglion cell side

down onto a plastic film using a single cat whisker brush, dragged up the "boat" ramp, and

lowered photoreceptor side down onto a coated coverslip. After floating off the plastic film, the

coverslip was mounted in the preparation chamber.

Patch clamping

The perfused recording chamber was mounted on the stage of an upright Zeiss Axioskop FS

microscope equipped with infrared DIC optics and epifluorescence (470/40 excitation filter and

525/50 emission filter). An additional optical channel transmitted excitation from a high-wattage

480 nm LED (Sutter Instruments, Novato, CA). A factory-calibrated radiometer (Model S370,

UDT Instruments, San Diego, CA) was used to measure the incident irradiance at the retina.

Recordings were performed in ACSF heated to ~34°C using 3 stages of heating (a lab-designed

warming blanket near the supply reservoirs followed by a feedback-controlled inline heater and

a coverslip preparation chamber with a heated bottom (Cell MicroControls, Norfolk, VA) with

flow at approximately 1 mL/s. Patch clamp recordings were performed using an EPC-10

amplifier controlled by "PatchMaster" software (Heka Instruments, Holliston, MA). Glass pipettes

were formed from Sutter glass (BF150-110) on a Sutter P-97 electrode puller (Sutter

Instruments, Novato, CA). The pipettes were filled with 1 of 3 different patch pipette solutions.

The standard solution contained (in mM) 120 K gluconate, 5 NaCl, 4 KCl, 2 EGTA, 10 HEPES,

0.3 NaGTP, 4 MgATP, 7 phosphocreatine, and Tris-KOH (pH 7.3) and had an impedance of 4-6

MOhm. The Alexa pipette solution contained (in mM) 116 K gluconate, 2 EGTA, 10 HEPES, 0.3

NaGTP, 4 MgATP, 7 phosphocreatine tris, neurobiotin 12.4 (Vector Laboratories, Burlingame,

CA), Alexa 633 0.07% (Thermo Fisher Scientific, Waltham, MA) and was adjusted to pH 7.3

with KOH. The standard Lucifer yellow pipette solution contained 10% H_2O, 12.4 neurobiotin

and 0.2% Lucifer yellow CH (Sigma-Aldrich, St. Louis, MO). For amphotericin recordings, cell-

attached recording pipettes were filled with standard patch solution supplemented with freshly

mixed amphotericin B (0.1 mM, Sigma). Liquid junction potentials were measured (9 to 11 mV),

and the measured membrane potentials were corrected accordingly.

In the electrophysiological studies designed to record from identified light- responding mRGCs,

we used retinas harvested from *Opn4^{cre}::Ai96* embryos. To locate light-responsive ganglion

cells, we first recorded the fluorescent response of the entire field of ganglion cells to ~5s

epifluorescent stimulation (470/40 nm, 3 x 10^{17} - 3 x 10^{19} photons/cm^2/s excitation, 525/50 nm

emission). The locations of light-responsive fluorescent cells were marked on the screen

monitor. The same area of the retina was then viewed with infrared DIC optics, and the patch

pipettes were aligned with the cells showing light responses. Fluorescent and DIC images were

collected using a CCD camera (Hamamatsu Digital Camera C4742) and Micro-Manager

software (Open-image.com, San Francisco, CA).

Intracellular calcium measurements

GCaMP calcium measurements:

Epifluorescence from isolated retinas was imaged in an inverted Nikon (TE2000S) microscope

(20x S-Fluor objective, NA 0.75). A Lambda LS 175-watt stand-alone xenon arc lamp and power

supply (Sutter Instruments, Novato, CA) provided the light for stimulation. The incident light

passed through an FITC/TRITC filter set (51004v2 - peak excitation wavelength ~480 nm;

Chroma Technology Corp., Bellows Falls, VT). Stimulus-evoked epifluorescence was focussed

onto a CCD camera (either a Hamamatsu C4742; 1024*1280 pixels or a Hamamatsu Orca

Flash 4.0 LT; 2048x2048 pixels, Hamamatsu Corp., San Jose, CA). The field of view was ~ 430

μm x 340 μm (C4742) or ~ 650 μm x 650 μm (4.0 LT). Images were acquired using 4x4 binning

and had a resulting resolution of 512 x 512 pixels. Metafluor software (Molecular Devices, Inc.,

San Jose, CA) controlled stimulation and data acquisition. The excitation light simultaneously

activated GCaMP3 or GCaMP6 and stimulated melanopsin photopigment. Light stimuli were

presented at 2 Hz and 3 Hz. Stimulation/acquisition periods were 530 ms (2 Hz) and 330 ms (3

Hz). The photon flux at the retina was routinely 4 x 10^{17} photons/cm^2/s. Regions of interest

encircling the somas of GCaMP3-fluorescent ganglion cells were set manually on images

recorded from the best-focussed position. Fluorescence images from complete fields of view

were recorded continuously. The summed responses from all pixels within each separate ROI

were also viewed in real time and recorded to disc. Photobleaching of GCaMP3 was shown to

be minimal based on the very small decrease in fluorescence in GCaMP-expressing cells

lacking photopigment. Photoresponses could be repeatedly and reproducibly measured for

periods of up to 60 min.

Calbryte-590 indicator dye

Exogenous calcium dye loading. We used Calbryte-590™ AM dye (573 nm Ex/ 588 nm Em;

AAT Bioquest, Sunnyvale CA) to monitor intracellular calcium. This dye localizes to the

cytoplasm and is 10-fold brighter than Rhod-2 AM (Zhao et al. 2019). A 5 mM stock solution of

the dye in anhydrous DMSO was prepared just prior to use and diluted to a working solution of

20 µM by adding stock Calbryte-590™ AM solution to carbogen-bubbled ACSF containing

Pluronic F-127 (0.04% f.c.). Isolated retinas were incubated in the ACSF/Calbryte-590™

solution for 2 hours in the dark with continuous carbogen circulation; they were then rinsed with

fresh ACSF, mounted ganglion cell side down on a polylysine-coated (0.1 mg/mL) glass

coverslip and placed in a heated (37°C) perfusion chamber. Calcium imaging was performed on

the same optical rig used for the GCaMP experiments.

Immunohistochemical labelling

For GCaMP3/melanopsin immunohistochemistry, the eye was removed, the cornea was

perforated with a razor blade, and the entire eye was immediately immersed in 4% (w/v)

paraformaldehyde (PFA) in PBS. In some experiments, the anterior part of the eye, including

the cornea and lens, was first carefully removed before immersing the entire posterior eyecup in

fixative. The eye was fixed in PFA in 0.1 M PBS for 30–90 min at RT. The eye was then stored

overnight in 25% sucrose, 0.01% sodium azide in 0.1 M PBS at 4°C. The following day, the

eyes were embedded in OCT and frozen on dry ice. Twenty-micron sections of the retina were

cut perpendicular to the vitreal surface with a cryostat, mounted on Super-Frost Plus slides

(Fisher Scientific, Pittsburgh, PA), and stored frozen at −20°C until immunostaining. ToPro3

(Life Technologies, Waltham, MA, Cat. # T3605) staining was used to mark nuclei. UF008

(1:500, Advanced Targeting Systems, San Diego, CA) and goat anti-GFP (1:500, Novus

Biologicals, Littleton, CO) were used to label melanopsin and GCaMP3/GCaMP6, respectively.

Drugs

FR900359 was isolated from the dried leaves of *Ardisia crenata* according to a

previously published procedure (Schrage et al., 2015). YM-254890, tetrodotoxin, FPL65176 and

L-cis-diltiazem were purchased from Wako Chemicals (Osaka, Japan), Calbiochem

[MilliporeSigma] (Temecula, CA), Tocris Bio-Techne Corporation (Minneapolis, MN) and

MilliporeSigma (Temecula, CA), respectively.

274   **Results**

Cells in foetal mouse retinas are light-responsive as detected by the fluorogenic calcium

indicator dye Calbryte-590^{TM}.

Melanopsin ganglion cells in foetal mouse pups differentiate before birth (Tartellin et al.

2003). In postnatal mouse retinas, mRGCs and perhaps other types of cells coupled to mRGCs

are light-responsive, as demonstrated by stimulus-evoked increases in intracellular calcium

levels (Sekaran et al. 2005). Photosensitive cells have not been reported in the retinas of

embryonic mice or any other embryonic mammal. To determine whether such photosensitive

cells exist, we searched for light responses by monitoring the activity of cells in embryonic

mouse retinas after exogenously bulk-loading the retinas with the calcium-sensitive dye

Calbryte-590™ AM (Zhao et al. 2019). After harvesting, the flat-mounted retinas were

fluorescently imaged, immersed in Calbryte-590^{TM} AM, mounted in a perfusion chamber, bathed

with ACSF at 35°C, and stimulated with light of wavelength 480 nm (see Methods). Figure 1A

shows an example of a Calbright-590™-loaded 340 µm x 430-µm region cropped from the full

image (650 nm x 650 nm). This retina was harvested from an E17.5 mouse pup. The image was

captured from the video recording at the onset of the light stimulus. Figure 1B shows an image

of the same retina captured 2.5 s after stimulus onset. Figure 1C, which is a difference image

(Fig. 1B minus Fig. 1A), accentuates the light-evoked increase in fluorescence that reflects the

increase in intracellular calcium in the cells. The time courses of calcium increases in four of the

cells, marked with circular regions of interest, are plotted in Fig. 1D. Peak changes in calcium

centred approximately 2 s after the onset of the light. In total, we recorded Calbryte-590™

images in 2 to 3 animals taken from 3 different litters at age E17.5 and 2 litters at E18.5. We

commonly detected more than 100 light-dependent calcium responses in each 430 mm x 340

mm region of the retina. Analysis of 4 of the brighter cells is shown in Figure 1.

Non-OPN4 opsins such as encephalopsin (OPN3) and neuropsin (OPN5) are expressed

in the retina. Cells that express these opsins might also exhibit light-evoked Calbyte-590[TM]

signals (Buhr et al., 2015, Kumbalasiri and Provencio, 2005). In mouse eyes, markers for

neuropsin and encephalopsin appear after 10.5 days of gestation (Tarttellin et al. 2003). We

tested foetal retinas from melanopsin knockout mice (*Opn4*[cre] homozygotes) and never

observed a light-evoked increase in Calbright-590[TM] fluorescence in these retinas. These

findings support the interpretation that the light-evoked calcium increases we observed in foetal

retinas were evoked in OPN4 cells and that no detectable calcium elevations were observed in

OPN3- or OPN5-expressing cells. It is noteworthy that light stimulation of OPN5 cells has been

shown to modulate ocular dopamine levels in early postnatal mice (Nyugen et al. 2019).

In summary, the data presented here provide strong evidence that a subpopulation of

cells within the foetal retina exhibits light-dependent calcium responses. However, we cannot

rule out the possibility that the observed light responses reflect light-evoked activity from non-

mRGCs as well as from mRGCs. For example, Calbryte-590™ AM dyes (and likely other AM

indicator dyes) can indiscriminately load into cells of various types, including astrocytes

(Tirschbirek et al. 2015). Thus, neurons or glial cells coupled to mRGCs could also exhibit light-

evoked activation (Sekaran 2003; Reifler 2017). To address these ambiguities, we designed an

experimental approach to selectively restrict the expression of calcium indicators to mRGCs.

319   Figure 1 near here

321
Melanopsin ganglion cells in foetal retinas are light-responsive as shown by the

responses of the calcium indicator GCaMP3 expressed exclusively in mRGCs.

We first asked whether the genetically encoded calcium indicator GCaMP3 is selectively

expressed in mRGCs of mouse retina. We utilized the Ai38 mouse strain in which GCaMP3 is

conditionally expressed. In the progeny of *Opn4*[cre/+]::*Ai38* mice, an allele that encodes the

EGFP-based Ca^{2+} reporter GCaMP3 is transcribed in cells that express Cre recombinase

(Zariwala et al. 2014). Because the *Opn4* gene is expressed relatively early in gestation, we

reasoned that we could expect to observe conditional GCaMP3 expression well before birth and

that it would be possible to detect light-evoked calcium signals that were restricted to

melanopsin-expressing cells. Figure 2 A-C shows flat-mounted retinas in which melanopsin

immunofluorescence co-localizes with EGFP immunoreactivity. This confirms that GCaMP3 is

expressed in mRGCs. Of note, we also observed a population of GCaMP3-positive cells that

showed little, if any, melanopsin immunoreactivity. The melanopsin-immunoreactive cells we did

observe are likely to be predominantly mRGCs of the M1 subtype, which express high levels of

melanopsin (Ecker et al. 2010). The other GCaMP3-EGFP-labelled cells may correspond to

other mRGC subtypes that express melanopsin Cre but are known to express lower levels of

melanopsin. In radial cryosections, an EGFP signal was evident in the inner plexiform layer and

was strong in neurons with RGC morphology (Fig. 2D). These anatomical findings demonstrate

that GCaMP3 is indeed localized to mRGCs. It should be noted that the images shown in Figure

1A-D were obtained from postnatal mice. Although CGaMP3 was well expressed in embryonic

retinas, melanopsin antibody staining of prenatal retinas was not consistently detected.

Therefore, based on the co-localization of antibody staining, it cannot be persuasively argued

that the GCaMP3 cells in the foetal retinas co-expressed functional melanopsin protein.

However, given that we were able to record light-evoked calcium increases in most of the

embryonic GCaMP3-positive cells of the inner retina, there is little doubt that melanopsin protein

was expressed in these foetal RGCs. The light sensitivity of mRGC-GCaMP3 embryonic

neurons in retinae was next characterized.

To search for light responses in foetal eyes, we mechanically isolated retinas from foetal

pups, placed them in a perfusion chamber and stimulated them with light at 480 nm, the

wavelength of light to which the melanopsin photopigment is most sensitive. Light stimulation

resulted in $[Ca^{2+}]_i$ increases in GCaMP3-expressing neurons localized across the ganglion cell

layer of the isolated retinas. Single-frame images of GCaMP3-filled RGCs show fluorescence in

the retina of an E16.5 mouse before the onset of the stimulus (Fig. 2E) and during a light

response (Fig. 2F). The complete time courses of changes in $[Ca^{2+}]_i$ in the 4 encircled cells (Fig.

2E) are shown in Fig. 2G. The black line in Fig. 2G shows the averaged response of all 79 cells

recorded in this experiment. These records reveal that the time-to-peak following light onset

ranged from 1.5 s to 4 s. The response kinetics were comparable to those reported in previous

studies of postnatal mRGCs (Ecker et al. 2010, Hartwick et al. 2008, Sekaran et al. 2005). We

also measured and plotted a cumulative histogram of peak response amplitudes (Fig. 2H). The

responses of the individual cells (Fig. 2G) were taken from the 20[th], 40[th], 60[th] and 80[th] percentile

groups (Fig. 2H). We discerned no correlation between peak amplitude and time course and no

obvious partitioning of responses, suggesting that the temporal differences noted in different

mRGC subtypes in postnatal mice are not evident at this age (Tu et al., 2005).

In total, we examined 1 to 4 littermates from 12 different embryonic litters. In

Opn4$^{cre/+}$::Ai38 foetal pups, we found light-responsive cells in 6 of 6 litters. In *Opn4$^{cre/+}$::Ai96*

foetal pups, we observed light responses in 16 of 16 litters. Using our standard irradiance

stimulus (3.8×10^{16} photons/cm^2/s), we recorded up to 90 responsive cells in 430 μm x 340 μm

regions of the retinas. The fact that as many as 90% of the GCaMP3-positive cells exhibited

light responses argues against the possibility that artefactual Cre expression may have occurred

in non-melanopsin-producing cells (Delwig et al., 2016; Ecker et al., 2010). We were able to find

detectable $[Ca^{2+}]_i$ elevations with stimulus irradiances as low as 3×10^{16} photons/cm^2/s. At

stimulus irradiances below that level, the signal-to-noise ratios were too low for reliable

detection.

No light responses were observed in control experiments performed in melanopsin

knockout retinas. *Opn4$^{cre/+}$::Ai38* mice crossed with *Opn4$^{cre/cre}$* mice produce some

Opn4$^{cre/cre}$::Ai38 pups. The mRGCs in these pups express GCaMP3 but not melanopsin. We

found no light-responsive cells in these foetal pups (Fig. 2G, inset). The experiments with

knockout retinas also serve to rule out the possibility that the light-evoked increase in GCaMP3

fluorescence in melanopsin-expressing cells is an artefact unrelated to melanopsin-mediated

calcium influx. Additionally, the recordings in knockout cells reveal that bleaching of GCaMP3 is

minimal under our experimental conditions. Specifically, the fluorescence amplitude 10 seconds

after light onset was 94 ± 3% of the initial amplitude (Fig. 2G, inset).

In summary, these results convincingly demonstrate that mRGCs in foetal mouse retinas are

responsive to light. In addition, the Calbryte590^{TM} measurements in OPN4 knockout mice

suggest that any photo-responses generated by other opsins, such as OPN3 and OPN5, were

below the limit of detection.

390   Figure 2 near here

Antagonists of the G$_{q/11}$ G-protein family attenuate mRGC light responses in foetal retina.

Light-activated opsins in invertebrate eyes signal via coupling to members of the G$_{q/11}$

family of G-proteins (Katz B & Minke B., 2009). Similarly, light-activated melanopsin in mRGCs

has been postulated to signal via a comparable G$_{q/11}$ pathway (Panda S et al., 2005; Hughes et

al., 2015; Graham DM et al., 2008; Jiang et al., 2018). However, a recent study reported that

deletion of G$_{q/11}$ G-proteins does not abolish light responses in postnatal mRGCs (Chew et al.,

2014). We tested whether G$_{q/11}$ signalling plays an essential role in embryonic mRGC light

responses. To do this, we examined the effects of two different cell- permeable antagonists,

FR900359 (Schrage et al., 2015) and YM-254890 (Kawasaki et al., 2003; Nishimura et al.,

2010), on light responses. These two compounds inhibit activation of the Gnaq, Gna11 and

Gna14 alpha subunits of $G_{q/11}$-G proteins. For these experiments, we used *Ai96* mice that

express the calcium indicator GCaMP6. GCaMP6 has superior sensitivity and a better

signal/noise ratio than GCaMP3 (Chen et al., 2013). The methods of breeding and fluorescence

recording were otherwise identical to those used with the *Ai38* animals in the previous

experiments. Recordings were made in *Opn4$^{cre/+}$::Ai96* retinas harvested from E15.5 and E16.5

foetal pups. The retinas were placed in control medium (ACSF), and light responses were

recorded from mRGCs before and after the addition of YM-254890 at 200 nM or FR900359 at 1

μM. Figure 3A shows an example of a recording from one retina; the light responses, quantified

as the increase in light-induced fluorescence, dF/F, were stable over a 40-min period in ACSF

but were progressively attenuated during the 30-min period following the addition of YM-

254890. Four representative light responses on an expanded time scale are shown (Fig. 3B).

We found no evidence of reversibility following washout. Poor reversibility has been reported

previously (Yasui et al., 2008). The time courses of the mean light responses from multiple

mRGCs in each retina are shown in Fig. 3C and D. Both YM-254890 (E15.5 retina, 32 mRGCs)

and FR900359 (E16.5 retina, 43 mRGCs) severely attenuated light responses after 30 min. To

rule out the possibility that the inhibitors merely killed the mRGCs or quenched the

fluorescence, we applied high potassium (50 mM) at the end of the inhibitor infusion. Figure 3D

shows an example in which potassium substitution was able to induce a dF/F larger than that

originally induced by light.

Prolonged incubation in 200 nM YM-254890 reduced the average dF/F of the light

responses to 7% ± 11% of that of the control responses (n=93). Prolonged incubation in 1 μM

FR900359 reduced the average dF/F of the light responses to 4% ± 18% that of the control

responses (n=219). Comparing light responses from individual cells prior to and during drug

application, we found highly significant drug effects across the ensemble of responsive cells.

According to the Wilcoxon Signed Rank Sum Test for paired data, the two-tailed P = $2.01948e^{-28}$

for YM and was even smaller for FR.

The use of two independent Gnaq/11/14 inhibitors diminishes the possibility that off-

target effects of either FR900359 or YM-254890 are responsible for the observed suppression

of light responsiveness. These results implicate $G_{q/11}$ signalling in the generation of light

responses in embryonic mRGCs. One caveat is that the averaged data shown in Figure 3 may

not reflect the presence of subpopulations of mRGCs that might utilize non-$G_{q/11}$ pathways.

436   Figure 3 near here

437
438

No subgroups of G$_{q/11}$-independent mRGCs in embryonic retinas are detected in

population data.

Jiang et al. (2018) reported that in adult mice most classes of mRGCs employ a G$_{q/11}$

cascade for transduction. However, they also reported that M4- type mRGCs utilize a second

alternative transduction cascade involving a cyclic nucleotide as the second messenger and

HCN channels. Jiang et al. (2018) reported that this pathway very likely does not use G$_{q/11}$

family members or TRP6/7 ion channels. In support of this conclusion, they showed that M4

mRGCs responded to light normally in cells devoid of *Gnaq*, *Gna11* and Gna14 as generated in

Gnaqfl/fl;Gna11$^{-/-}$;Gna14$^{-/-}$ mouse eyes transfected with Cre-GFP via adeno-associated virus

serotype 2 (AAV2-CMV-Cre-GFP).

We hypothesized that if a group of mRGCs in foetal retinas also does not use G$_{q/11}$

signalling, then a subpopulation of GCaMP6-expressing mRGCs should be resistant to the

action of YM and FR. Figure 4, below, shows cumulative data from 93 mRGCS tested in YM-

254890 and 219 mRGCs tested in FR900359. Our data indicate that only approximately 1% of

the cells might be resistant to these drugs. Thus, our experiments provide no supportive

evidence for the existence of a detectable subpopulation of foetal mRGCs that utilize a non-G

$_{q/11}$ transduction pathway.

458   Figure 4 near here

TTX-sensitive sodium currents amplify mRGC light responses in embryonic retinas.

Based on electrophysiological recording and calcium measurements from neonatal

mRGCs harvested from rat retinas, Hartwick et al. (2007) proposed a transduction pathway in

which melanopsin-driven G-protein activation leads to the opening of cationic TRPC6/7

channels. The resulting inward cation flux subsequently depolarizes the membrane, leading to

activation of TTX-sensitive sodium action potentials and then to activation of L-type calcium

channels, which, in turn, increases calcium ion influx. We sought to determine whether TTX-

sensitive sodium and L-type calcium currents support light responses in embryonic mRGCs.

The likelihood of finding a role for TTX-sensitive currents in embryonic mRGC responses is

enhanced by previous work demonstrating that a majority (78%) of randomly selected RGCs in

E15 mouse retinas expressed TTX-sensitive sodium currents (Rorig and Grantyn, 1994).

We found that TTX (1 µM) reversibly diminished the average light response amplitudes

of virtually all the foetal GCaMP6-expressing mRGCs we tested. Figure 5 shows the responses

recorded from 177 cells in 4 E16.5 retinas. Each solid data point (blue) denotes the light

response amplitude in TTX (ordinate) as a function of the amplitude in control ACSF prior to

TTX (abscissa). The open circles show the responses of the same cells after washout of TTX.

We found that TTX effectively reduced the light responses to below the detectable threshold in

21% of the cells (dF/F < 0.05). The aggregate differences between responses in TTX versus

washout are highly significant (Wilcoxon Signed Rank Test for paired data, two-tailed P = 1.01e^{-46}

). Of note, these data show that while TTX reversibly diminished the light responses in virtually

all of the tested mRGCs, the effects of TTX varied across the sample of cells. This variability is

at odds with Hartwick et al. (2007), who reported that TTX abolished responses nearly

completely in all mRGCs (n = 8). The observed difference in the efficacy of TTX could reflect

our sampling of a larger population of mRGCs. Alternatively, these findings might reflect cell-

specific differences in the maturation of INa(V) channels during embryonic and postnatal

development. Nonetheless, we conclude that these TTX-sensitive channels do serve to amplify

the light-evoked responses in a plurality of embryonic mRGCs.

494   Figure 5 near here

Modifiers of L-type voltage-gated calcium channels modulate the light responses of

mRGCs in E16.5 mouse retinas.

Previously, Hartwick et al. (2007) showed that L-type voltage-gated calcium channels

account for most of the calcium influx that occurs following light stimulation of enzymatically

isolated postnatal rat mRGCs. While there are no reports of L-type calcium channels specifically

in mRGCs of embryonic retinas, earlier studies suggested that L-type calcium channels are

expressed in unspecified, randomly selected RGCs in E15 mouse retinas (Rorig and Grantyn,

1994). Consistent with this finding, expression of two L-type channel subunits, Cacna1a and

Cacna1b, in the eyes of E15.5 mouse pups has been reported (Allen Brain Atlas, Cacna1a – RP

090930_03_A09 – sagittal; Cacna1b - RP_090707_03_G11 - sagittal). We tested

pharmacologically for evidence that L-type calcium channels contribute to the light responses of

GCaMP3-expressing mRGCs. In one set of experiments, we found that L-*cis*-diltiazem, a

blocker of L-type calcium channels (Catterall et al., 2005), significantly diminished light

responses in embryonic retinas (n=3). Figure 6 shows the results obtained from 31 cells that

were recorded simultaneously in one retina. Panel A shows examples of reversible L-cis-

diltiazem-induced suppression of light responses in 5 mRGCs. Panel B plots the responses

from all mRGCs in which the light response peak amplitude returned to a value between 80 and

120% of that observed under control conditions upon washout. The blue dots plot the responses

in L-cis-diltiazem, and the black dots plot the amplitude of the responses following washout. The

five circled data points correspond to the light responses shown in Fig 6A. The effect of L-cis-

diltiazem is highly significant according to the Wilcoxon Signed Rank Test for paired data (two-

tailed P = $9.31e^{-10}$).

519   Figure 6 near here

In a complementary set of experiments, we found that FPL64176, an activator of L-type

calcium channels (Zheng et al., 1990), increased light-evoked dF/Fs in embryonic mRGCs. In

previous studies, FPL64176 was shown to enhance the activity of retinal L-type calcium

channels. Specifically, FPL64176 enhanced L-type calcium-mediated evoked release of the

neurotransmitter glycine from amacrine cells (Bieda & Copenhagen, 2004). In addition,

FPL64176 potentiated spontaneous waves of activity that occur in neonatal retinas (Singer et

al., 2001). Figure 7 plots the peak light-evoked fluorescent responses observed before and

during exposure of explanted retinas to FPL64176 (2 μM). The data include the responses from

147 GCaMP3-expressing mRGCs from 2 different E16.5 retinas. The responses observed

during treatment with FPL64167 are plotted against the responses observed under control

conditions. On average, the df/F values in FPL64176 were 0.22 ± 0.02 (SEM) larger than dF/F

in control. The blue line shows the best linear regression fit to the FPL64176 data (y = 1.51x).

The black line shows the values that would be expected if FPL64176 had no effect on the light

responses. The Wilcoxon Rank Sum Test for paired data yielded a two-tailed P value of 1.172e^{-}

^{32}. Together, the L-cis-diltiazem and FPL-64176 findings strongly support a role for L-type

calcium channels in the generation of the light responses, similar to the transduction role

proposed by Hartwick et al. (2007) for postnatal mRGCs.

542   Figure 7 near here

Voltage-activated sodium currents exist ubiquitously in mRGCs from at least E16.5

Having found that the INa (V) antagonist TTX reduces and sometimes completely

abolishes light responses in embryonic mRGCs, we sought to characterize these sodium

currents electrophysiologically. Using patch pipettes, we recorded from mRGCs genetically

marked with either GCaMP6, tdTomato or EGFP derived from $Opn4^{cre}$::Ai96, $Opn4^{cre}$::Ai14, or

$Opn4^{EGFP}$ mouse strains, respectively. Patch pipettes were directed towards the fluorescently

marked cells in isolated flat-mounted retinas. In the GCaMP-labelled retinas ($Opn4^{cre}::Ai96$), we

recorded from only those RGCs that displayed light-evoked calcium responses as reported by

GCaMP6. An example of a voltage clamp recording from an mRGC in E18.5 retina is shown in

Fig. 8. Panels A-C show current responses to a series of sequentially larger 10 mV voltage

steps initiated at –70 mV (see Panel E). Brief, transient inward currents began to occur at – 40

mV and became faster with successive depolarizing pulses (Panel A). Perfusion with TTX (1

µM) reversibly blocked the inward currents (Panels B and C). The evoked TTX-sensitive

currents (Panel D) were isolated by subtracting the response in Panel B from Panel A. The

voltage dependence of the TTX-sensitive currents is plotted in Panel F. The peak current

amplitude (black dots) exhibited a sharp onset at -40 mV, reached a maximum at -30 mV and

then declined with further depolarizing steps. The kinetics and voltage dependency of the TTX

currents shown here resemble those of the canonical INa(V) currents observed in excitable

muscle and nerve cells across multiple phyla (Johnston & Wu, 1995).

The mean amplitudes of INa (V) currents increase substantially from E16.5 to P1-3.

INa (V) currents were detected in virtually all of the mRGCs recorded in 10 E16.5, 35

E18.5 and 7 P1-3 retinas. The peak INa (V) currents (not all data were analysed and included)

are plotted in Fig. 8 (Panel G). The mean peak currents in the aforementioned groups of retinas

were 320, 674 and 1986 pA, respectively. Comparable developmentally linked increases in the

amplitude of INa(V) currents have been reported in random samplings of RGCs in mouse and

rat retinas (Rorig and Grantyn, 1994; Schmid and Guenther, 1996). Our findings confirm that

functional voltage-dependent sodium channels are expressed in mRGCs as early as E16.5 and

further reveal an increase in the size of these currents increase during embryonic maturation. It

is evident that the peak INa (V) varied across the sampled population at each age. We were

unable to reliably and systematically determine whether the variability reflected the presence of

cells of different sizes or whether the expression of INa (V) channels per unit of cell membrane

varied. Of note, Schmid & Guenther (1996) divided whole-cell sodium current amplitude by cell

capacitance and found that INa (V) current density increased sizably from E18 to P6.

579   Fig. 8 near here

581   **Electrophysiological recordings demonstrate that irradiance with light as dim as 3 x 10^{12}**

582   **photons/cm^2/s evoke responses in individual foetal mRGCs.**

583   Our previous work demonstrated that the ambient light in the mouse housing room was

bright enough to trigger subsequent vascular development via stimulation of melanopsin in the

eyes of foetal pups (Rao et al. 2013). Specifically, normal regression of the hyaloid blood

vessels was retarded in foetal pups lacking expression of melanopsin (*OPN4$^{-/-}$*) and in WT pups

whose mothers were dark-reared from 16.5 days post-conception to the time of the pups' birth.

The effects of ambient light were not transmitted from the pregnant female to the pups but were

a direct effect of ambient light on the foetal pups themselves. This was shown by the fact that

environmental light still contributed to normal ocular vessel development when the pregnant

female was enucleated or when melanopsin was selectively deleted in the pregnant mothers but

not in the foetal pups. Thus, the light that struck the foetal eyes resulted from ambient light

transmission through the abdominal walls of the dams. Direct measurements of the attenuation

of the intensity of blue light across the dams' skin and muscle layers support the notion that if

mRGCs in embryonic retinas are as sensitive to light as mRGCs in postnatal retinas, the

embryonic mRGCs could easily be activated by low-intensity environmental light at floor level in

the mouse housing room (5.6 x 10^{13} photons/cm^2/s, Rao et al. 2013). However, the actual

photosensitivity of embryonic mRGCs is unknown. In the GCaMP experiments described above

(Figs 2-7), we were not able to measure mRGC sensitivity because the minimum stimulus

intensity needed to interrogate GCaMPs strongly activated melanopsin as well. To circumvent

this experimental obstacle, we turned to a non-optical, electrophysiological approach using

patch pipette recordings from individual mRGCs.

603   We recorded light-evoked responses from fluorescently marked mRGCs in foetal retina

preparations. We found a total of 15 mRGCs that exhibited detectable light responses. Figure 9

shows light responses recorded from an mRGC in the retina of an E18.5 pup. Action potentials

were observed at an irradiance of 2.9×10^{12} photons/cm^2/s (Fig. 9D4). The onset of spiking

displayed a latency of 21 s. The responses to steps of light at higher irradiances (5.8×10^{14}, 1.5

$\times 10^{14}$ and 1.35×10^{13} photons/cm^2/s) are plotted in Fig. 9D1, D2, and D3, respectively. Similar

to the mRGC responses recorded in neonatal and adult retinas, the response latencies

decreased as the irradiance increased (Tu et al. 2005, Wong KW, 2012). In our sample of cells,

one mRGC had a threshold of 1.6×10^{12} photons/cm^2/s. Seven of the 15 mRGCs had

thresholds between 6.2×10^{12} and 3.5×10^{13} photons/cm^2/s, and six of the 15 light responders

had thresholds between 1.5×10^{14} and 2.4×10^{14} photons/cm^2/s. It should be noted that all of

the GCaMP6-expressing mRGCs recorded electrophysiologically were initially identified by their

responsiveness to light at 3.10^{17} – 3.10^{19} photons/cm^2/s (Fig. 9A,B).

A population of low-amplitude spikes during light onset and occasional depolarizing

"bumps" with a duration of approximately 1-2 s are evident in these recordings. We interpret the

smaller spikes as spikelets conducted from neighbouring mRGCs as described by Arroyo et al.

(2016). The bumps, which occur with a frequency of approximately 1/min, probably reflect the

occurrence of spontaneous waves (Meister et al. (1990); Maffei L& Galli-Resta L (1990)).

622   Figure 9 near here

mRGCs in embryonic retinas are coupled via gap junctions.

Cells in the adult mammalian retina are extensively coupled through gap junctions both

to cells of the same type and to cells of different types. In the retinas of rats and mice, in

particular, mRGCs are coupled to amacrine cells (Muller et al. 2010; Reifler et al. 2015).

However, dye coupling between mRGCs in adult rodent retina has not been found, although

Müller et al. (2010) and Ecker et al. (2010) specifically searched for homologous coupling

between these cells. Precedent suggests that gap junction coupling could be age-dependent

and that gap junction coupling may therefore exist in the retinas of younger mice. In spinal cord

(Walton and Navarrete, 1991) and in the cerebral cortex (Kandler and Katz, 1991), gap junction

coupling between neurons can exist early in development but is absent or much reduced in

older animals. Consistent with a greater prominence of gap junction coupling in the retinas of

very young animals compared to those of adults, Sekaran et al. (2003) reported that MFA, a

putative blocker of gap junction coupling, reduced the number of light-responsive cells in early

postnatal retinas. This was interpreted as evidence for gap junction coupling between mRGCs

and other cells. Subsequently, Arroyo et al. (2016) found MFA-inhibitable functional and tracer

dye coupling between genetically marked mRGCs in young neonatal mice (P4 to P8). We

extended the work of Arroyo et al. (2016) to specifically test whether prenatal mRGCs are

coupled to one another via gap junctions, pursuing the idea that younger mice may be more

likely to express gap junction communication. We tested whether MFA, a compound that is

commonly used to block gap junction coupling, reduced or eliminated light responses (Sekaran

et al. 2003; Veruki et al. 2010). We examined the effects of MFA on light-evoked calcium

elevations across a population of GCaMP-expressing mRGCs in fields of cells in flat-mounted

retinas. Figure 10A shows the combined results obtained from 81 cells in two separate E16.5

retinas. MFA reversibly attenuated the amplitudes of the cells' responses to light. The black dots

in the top panel plot the response amplitudes after washout as a function of the response

amplitudes observed prior to MFA application. The black dots are best fitted by a linear

regression corresponding to y = 1.04 x. The data show good recovery from MFA. The blue dots

plot the response amplitudes of each mRGC in MFA as a function of the response amplitudes

observed prior to MFA application. The best-fitted linear regression line is y = 0.21 x. Non-

parametric Wilcoxon Rank Sum Tests indicate that the reduction observed in MFA is highly

significant (P = 8.3 x 10^{-25}). These results are consistent with the spreading of light-evoked

excitation between mRGCs via gap junctions. We performed the MFA experiments in 7 fields

from 5 embryonic retinas. In 4 fields, MFA attenuated the responses reversibly. In the other 3

fields, the light responses recovered to more than 80% of their pre-drug levels in only a minority

of the mRGCs. This apparent lack of reversibility is likely due to not having waited long enough

after initiating washout (> 20 min). It should also be noted that in *OPN4cre::Ai96* mice we could

only interrogate responses in mRGCs. Therefore, we were unable to use this method to rule out

gap junction coupling to other non-melanopsin-expressing neurons. Additionally, a caveat

associated with relying on MFA to identify gap junction coupling is that carbenoxolone, a

previously used MFA-related compound, has been reported to attenuate L-type calcium currents

in mRGCs (Bramley et al. 2011). Finding reduced numbers of light-responsive cells in MFA

could also be attributed to the reduction of calcium currents.

To explore whether our MFA findings did indeed indicate coupling between mRGCs, we

used patch pipettes to inject 17 embryonic mRGCs with neurobiotin as a means of tracing gap

junction-mediated dye coupling to adjacent cells. Figure 10B shows an example of an mRGC

that was injected with neurobiotin. Three somas were neurobiotin- and EGFP-positive, indicative

of migration of the tracer from one GCaMP6-expressing mRGC to at least two other mRGCs

(stars). At least five other somas were neurobiotin-positive (red) but lacked EGFP

immunoreactivity. In summary, in 12 of 17 injections, at least one nearby mRGC cell was filled

with neurobiotin. The maximum number of dye-coupled mRGCs observed to be connected an

injected mRGC was 4. In 10 of the 17 injected mRGCs, the axons were filled with neurobiotin.

Neurobiotin also filled from 0 to 7 non-mRGCs that were identified by the absence of OPN4cre-

driven EGFP or tdTomato from their cell bodies. We were often able to follow neurobiotin-filled

dendritic processes from the mRGC somas to the other dye-filled cells. These results provide

strong evidence that mRGCs are coupled to other mRGCs as well as to other classes of inner

retinal neurons in foetal mouse retinas.

682   **Figure 10** Near here

683
684

685   **Discussion**

The photopigment melanopsin is expressed in a subset of retinal ganglion cells termed

mRGCs. Markers for melanopsin first appear in embryonic mouse and human retinas well

before birth (embryonic day 11.5 and gestational week 8.6, respectively). mRGCs in postnatal

retinas can be excited autonomously by light independently of visual inputs from rods and

cones, the canonical photoreceptors in the eye. We recently demonstrated that exposure of

pregnant females to light regulates the normal maturation of ocular vasculature in subsequently

born pups. Based on our finding that the deletion of melanopsin in the foetal pups phenocopied

darkness, we inferred that light stimulation of mRGC in the eyes of the foetuses modulates

vascular maturation. There have been no reports directly demonstrating that mRGCs in foetal

retinas are light-sensitive, raising some doubts about whether the ambient light in the mouse

colony in which the pregnant females were housed was sufficient to excite mRGCs in the eyes

of the foetuses. Here, using both intracellular calcium indicators and patch pipette recordings,

we show that foetal mouse mRGCs respond to light as early as E15.5, four days before birth.

We report further that foetal mRGCs share many of the phototransduction pathways that have

been previously identified in mRGCs in the retinas of postnatal animals. Pharmacological

experiments reveal that light responses are initiated by the coupling of light-activated

melanopsin to a $G_{q/11}$-G-protein cascade. We also show that calcium influx through L-type

calcium channels and cation influx through TTX-sensitive sodium channels boost the amplitude

of photoresponses. Electrophysiological recordings from individual embryonic mRGCs

demonstrated that these cells can respond to light as dim as 2.9 x 10^{12} photons/cm^2/s, providing

strong supportive evidence that the ambient light in rooms where the pregnant females were

housed is sufficient to excite mRGCs in foetuses. Additionally, we found that when the dye

neurobiotin was injected into individual single mRGCs, it flowed into other nearby mRGCs via

gap junction coupling. This finding contrasts with reports performed in adult rodent retinas in

which no mRGC/mRGC coupling was discovered. Finding photosensitivity in the embryonic

retinas of mice raises the question of whether human foetuses and early preterm infants could

be sensitive to light and whether light exposure might modulate ocular development. Given that

light stimulation of mRGCs plays a role in entraining circadian rhythms and mood in postnatal

animals, some consideration should be given to the consequences of light exposure

experienced by human foetal or preterm infants. Light exposure sources might include lighting

regimens in neonatal care units and light exposure during intrauterine surgery. Such exposure

might influence ocular development and/or behavioural states.

No evidence was found that other photopigment-expressing cells in the foetal retinas of

mice are light-sensitive.

721   Could cells other than mRGCs in the embryonic eye confer light sensitivity that might

affect ocular development? The present study focussed on the light sensitivity of mRGCs in

retina. The results of the Calbryte 590^{TM} experiments (Fig. 1) argue against the idea that OPN3-

and OPN5-expressing cells generate light responses in the foetal mouse retina. Strictly

speaking, we did not address the question of whether other non-retinal cell types that might

express melanopsin prenatally could regulate ocular development. However, the preponderance

of evidence presented here and in a previous study (Rao et al. 2013) strongly supports the idea

that light stimulation of mRGCs is sufficient to account for light modulation of ocular vasculature

maturation. In adult rodents, melanopsin expression has been found in the cornea, the iris and

the retinal pigmented epithelium (RPE). We recently showed that melanopsin-containing fibres

and cells in the corneas of postnatal animals were not light-responsive, suggesting that

melanopsin might play a role in other sensory modalities such as thermal detection (Delwig et

al. 2018). While light-sensitive melanopsin cells have been identified in the iris, there are no

reports that these cells are light-sensitive in embryonic retinas or even that melanopsin markers

are found in the embryonic irides. Melanopsin gene expression has been reported in cultured

lines derived from RPE. There is no evidence that native RPE or cultured RPE cells lines are

light-responsive. Further study focussed on the iris and RPE would be required to ascertain

whether light stimulation of these cell types alters the vascular maturation of the eye.

mRGCs are coupled electrotonically to one another in the foetal retina.

Syncytial networks of electrotonically coupled neurons facilitate the spread of excitation
across regions of neural tissue and coordinate the timing of excitability and rhythmic oscillations
(Nagy et al., 2018). Electrotonic coupling between heterologous and homologous classes of
neurons is common in both mammalian and non-mammalian neural tissues. Moreover, it is
commonly found that electrotonic coupling is more prominent in the tissues of younger animals
(Walton and Navarrete, 1991, Kandler and Katz, 1991). In the retinas of adult mice, electrotonic
coupling of mRGCs to other mRGCs seems to be absent [Müller et al. (2010), Ecker et al.
(2010)], but it is observed in younger postnatal mice (Sekaran et al. 2003; Arroyo et al. 2016). In
the present study, we found electrotonic coupling between mRGCs. Additionally, we have
evidence for electrotonic coupling from mRGCs to heretofore-unidentified non-mRGC cell types
in the inner retinas of foetal mice. Electrotonic coupling between mRGCs and other
heterologous neurons raises the possibility that light exposure during foetal and early postnatal
maturation could excite neighbouring ganglion cells that project axons to the higher visual
centres. In postnatal animals, light deprivation produced by suturing shut the eyelid of one eye
disturbs the normal segregation of eye-specific targets in higher visual centres. This suggests
that light excitation of mRGCs in neonatal animals before the normal age of eyelid opening may
play a previously unrecognized role in retinotopic mapping in the higher visual centres.
Another notion drawn from the results presented here is that foetal light exposure might
play a role in the formation of ocular dominance columns in primates. Horton and his colleague
(Horton & Hocking, 1996) found ocular dominance columns in newborn macaques that were
delivered by Caesarean section and kept in total darkness for a week after birth. Horton and
Hocking argued that the formation of columns was genetically programmed based on the
assumption that there had been no light-induced activity in the retinas of foetal monkeys. Given
our findings in mice, it is worth considering the idea that ambient light might stimulate mRGCs in
embryonic monkeys and thereby play a previously unrecognized role in the formation of ocular
dominance columns.
767
Light excitation of mRGCs in foetal retinas could modify patterns of retinogeniculate
connections.
Retinal ganglion cells extend axons to the lateral geniculate nucleus several days before
birth. The postsynaptic contact regions in the LGN exhibit eye-specific contact regions. A
consensus opinion is that spontaneous activity, often observed as propagated waves of activity,
plays a critical role in segregation (Shatz CJ & Stryker MP (1988); Shatz CJ (1996); Meister et
al. (1991); Maffei & Galli-Resta, (1990). Renna et al. (2011) and Arroyo et al. (2016) reported

that light modulated the temporal structure of retinal waves via activation of mRGCs in the

retinas of young neonatal mouse pups. Renna et al. (2011) also reported that mRGC excitation

perturbed the eye-specific targeting of RGC axons in the LGN. Given our demonstration that

light can activate mRGCs in foetal retina, we suggest that light exposure of pregnant females

might play a role in establishing retinogeniculate connectivity in the maturing pups. Light could

induce activity in mRGC axons targeting the LGN, or it could induce activity in neighbouring

RGCs that might be electrotonically coupled to mRGCs. Another alternative is that mRGC

excitation could modulate the structure of spontaneous waves that propagate across the foetal

retina. Further studies will be needed to establish whether the light-induced activity of mRGCs

as young as E15.5 is conducted to the LGN.

Are human foetuses sensitive to light and might ambient lighting conditions at early

gestational ages affect eye development?

Third-trimester human foetuses respond acutely to light directed through the mother's

abdominal wall. Examples of photic induced responses include modulation of heart rate and

body movement (Kiuchi et al. 2000) and visually evoked brain activity (Eswaran et al. 2004,

McCubbin et al. 2007). Since rods and cones begin functioning at approximately gestational

week 35, the acute actions of light stimulation could be activating rods and cones as well as

mRGCs. In any case, the occurrence of light-evoked responses in human foetuses

demonstrates that light can excite photoreceptors in the foetal eye after passing through the

abdominal wall of the pregnant mother.

Human eye development might also be sensitive to ambient light at gestational ages

near the end of the first trimester. Human foetuses at gestational day 60 are at the same

developmental age as mouse foetuses at gestational day 16.5 (Clancy et al. 2001), the age at

which ambient light modulates subsequent vascular maturation in mice. Relevant to the

possibility that reduced daily photoperiod might influence eye development in humans, an

analysis of 276,911 conscripts of the Israeli Defense Forces between 2000 and 2004 found an

association between an increased incidence of moderate and severe myopia and season of

birth. Conscripts for whom the end of the first *in utero* trimester occurred during the winter

months exhibited an increased risk of myopia (Mandel et al. 2008). One intriguing explanation is

that light exposure of foetuses during the first trimester influence eye growth. While Mandel et

al. (2008) ruled out family planning as a variable many other confounds need to be considered

including diseases experienced during winter months, Vitamin D deficiencies, birth weights and

other factors. In another analysis, Yang et al. (2013) reported that infants whose early gestation

occurred during the winter months were at increased risk of severe retinopathy of prematurity

(ROP). These findings raise the possibility that ambient light passing across the abdominal wall

of pregnant mothers early in gestation might be considered as a factor that could affect eye

development.

813
814 **References**

815
816 Arroyo DA, Kirkby LA & Feller MB (2016). Retinal Waves Modulate an Intraretinal Circuit of
817   Intrinsically Photosensitive Retinal Ganglion Cells. *J Neurosci.* 36:6892-905.

818
819 Arshavsky V.Y., Lamb T.D., Pugh E.N. Jr. (2002) G proteins and phototransduction. *Annu Rev*
820   *Physiol.*; 64(): 153-87.

821
822 Bailes H.J. & Lucas R.J. (2013) Human melanopsin forms a pigment maximally sensitive to blue
823   light (lambdamax approximately 479 nm) supporting activation of G(q/11) and G(i/o)
824   signalling cascades. *Proc Biol Sci* 280(1759):20122987.

825
826 Berson D.M., Dunn F.A., Takao M. (2002) Phototransduction by retinal ganglion cells that set
827   circadian clock. *Science.* 95:1070-3.

828
829 Bieda MC & Copenhagen DR (2004). N-type and L-type calcium channels mediate glycinergic
830   synaptic inputs to retinal ganglion cells of tiger salamander. *Vis Neurosci.* 21:545-50.

831
832 Bramley JR, Wiles EM, Sollars PJ & Pickard GE. (2011) Carbenoxolone blocks the light-evoked
833   rise in intracellular calcium in isolated melanopsin ganglion cell photoreceptors. *PLoS*
834   *One.* 6(7):e22721. doi: 10.1371/journal.pone.0022721. Epub 2011 Jul 29.

835
836 Breedlove MA (2017) Foundations of Neural Development, Sinauer Associates, Sunderland,
837   MA

838
839 Buhr ED, Yue WW, Ren X, Jiang Z, Liao HW, Mei X, Vemaraju S, Nguyen MT, Reed RR, Lang
840   RA, Yau KW, & Van Gelder RN (2015) Neuropsin (OPN5)-mediated photoentrainment of
841   local circadian oscillators in mammalian retina and cornea. *Proc Natl Acad Sci U S A*
842   112:13093-8.

843
844 Catterall WA, Perez-Reyes E, Snutch TP, Striessnig J. (2005) International Union of
845   Pharmacology. XLVIII. Nomenclature and structure-function relationships of voltage-gated
846   calcium channels. *Pharmacol Rev* 57:411-25.

847
848 Chen T.W.*,* Chen TW, Wardill TJ, Sun Y, Pulver SR, Renninger SL, Baohan A, Schreiter ER,
849   Kerr RA, Orger MB, Jayaraman V, Looger LL, Svoboda K. & Kim DS. (2013) Ultrasensitive
850   fluorescent proteins for imaging neuronal activity. *Nature* 499(7458):295-300.

851
852 Chew K.S., Schmidt T.M., Rupp A.C., Kofuji P. & Trimarchi J.M. (2014) Loss of Gq/11 genes
853   does not abolish melanopsin phototransduction. *PLoS One* 9(5):e98356.
854

855   Clancy B, Darlington RB & Finlay BL (2001) Translating developmental time across mammalian
856   species. *Neuroscience* 105:7-17.
857
858   Delwig A, Larsen DD, Yasumura D, Yang CF, Shah NM & Copenhagen DR (2016) Retinofugal
859   projections from melanopsin-expressing retinal ganglion cells revealed by intraocular
860   injections of cre-dependent virus. *PLoS One.* 11(2):e0149501. doi:
861   10.1371/journal.pone.0149501.
862
863   Delwig A, Chaney SY, Bertke AS, Verweij J, Quirce S, Larsen DD, Yang C, Buhr E, Van Gelder
864   R, Gallar J, Margolis T, Copenhagen DR. (2018) Melanopsin expression in the cornea. *Vis
865   Neurosci.* 35:E004. doi: 10.1017/S0952523817000359.
866
867   Do M.T.H. & Yau K.W. (2010) Intrinsically photosensitive retinal ganglion cells. *Physiol Rev.*
868   90:1547-81. doi: 10.1152/physrev.00013.2010.
869
870   Ecker JL, Dumitrescu ON, Wong KY, Alam NM, Chen SK, LeGates T, Renna JM, Prusky GT,
871   Berson DM & Hattar S (2010) Melanopsin-expressing retinal ganglion-cell photoreceptors:
872   cellular diversity and role in pattern vision. *Neuron* 67(1):49-60.
873
874   Eswaran H, Lowery CL, Wilson JD, Murphy P, & Preissl H (2004). Functional development of
875   the visual system in human fetus using magnetoencephalography. *Exp Neurol.* 190 Suppl
876   1:S52-8.
877
878   Graham DM, Wong KY, Shapiro P, Frederick C, Pattabiraman K & Berson DM. (2008)
879   Melanopsin ganglion cells use a membrane-associated rhabdomeric phototransduction
880   cascade. *J Neurophysiol* 99(5):2522-2532.
881
882
883   Hartwick AT, Bramley JR, Yu J, Stevens KT, Allen CN, Baldridge WH, Sollars PJ, Pickard GE
884   (2007) Light-evoked calcium responses of isolated melanopsin-expressing retinal ganglion
885   cells. *J Neurosci* 27(49):13468-13480.
886
887   Horton JC & Hocking DR (1996) An adult-like pattern of ocular dominance columns in striate
888   cortex of newborn monkeys prior to visual experience. *J Neurosci.* 16:1791-807.
889
890   Hughes S, Jagannath A, Hickey D, Gatti S, Wood M, Peirson SN, Foster RG & Hankins MW..
891   (2015) Using siRNA to define functional interactions between melanopsin and multiple G
892   Protein partners. *Cell Mol Life Sci* 72(1):165-179.
893
894   Johnston D & Wu S M-S. Foundations of Cellular Neurophysiology 1995, The MIT Press,
895   Cambridge Massachusetts
896
897   Johnson J, Wu V, Donovan M, Majumdar S, Rentería RC, Porco T, Van Gelder RN &
898   Copenhagen DR. (2010) Melanopsin-dependent light avoidance in neonatal mice. *Proc
899   Natl Acad Sci U S A* 107(40):17374-17378.
900

901   Kandler K, Katz, LC (1995). Neuronal coupling and uncoupling in the developing nervous
902   system, *Curr. Opin. Neurobiol* 5, 98–105.
903
904
905   Katz B & Minke B. (2009) Drosophila photoreceptors and signaling mechanisms*. Front. Cell*
906   *Neurosci.* 10.3389/neuro.03.002.2009.
907
908   Kawasaki T, Taniguchi M, Moritani Y, Hayashi K, Saito T, Takasaki J, Nagai K, Inagaki O &
909   Shikama H. (2003) Antithrombotic and thrombolytic efficacy of YM-254890, a Gq/11
910   inhibitor, in a rat model of arterial thrombosis. *Thromb Haemost* 90(3):406-413.
911
912   Kiuchi M, Nagata N, Ikeno S & Terakawa N (2000) The relationship between the response to
913   external light stimulation and behavioral states in the human fetus: how it differs from
914   vibroacoustic stimulation. Early Hum Dev. 2000 May;58(2):153-65.
915
916   Kumbalasiri T. & Provencio I. (2005) Melanopsin and other novel mammalian opsins. *Exp Eye*
917   *Res* 81(4):368-375.
918
919   Madisen L, Zwingman TA, Sunkin SM, Oh SW, Zariwala HA, Gu H, Ng LL, Palmiter RD,
920   Hawrylycz MJ, Jones AR, Lein E & Zeng H. (2010) A robust and high-throughput Cre
921   reporting and characterization system for the whole mouse brain. *Nat Neurosci* 13(1):133-
922   140.
923
924   Maffei L & Galli-Resta (1990) Correlation in the discharges of neighboring rat retinal ganglion
925   cells during prenatal life *Proc Natl Acad Sci USA* 87:2861-4.
926
927   Mandel Y, Grotto I, El-Yaniv R, Belkin M, Isreal E, Polat U & Bartov E. (2008) Season of birth,
928   natural light, and myopia. *Ophthalmology*. 115:686-92. Epub 2007 Aug 15.
929
930   McNeill DS, Sheely CJ, Ecker JL, Badea TC, Morhardt D, Guido W & Hattar S. (2011)
931   Development of melanopsin-based irradiance detecting circuitry. *Neural Dev*. 6:8. doi:
932   10.1186/1749-8104-6-8.
933
934   Melyan Z, Tarttelin EE, Bellingham J, Lucas RJ & Hankins MW (2005) Addition of human
935   melanopsin renders mammalian cells photoresponsive. *Nature* 433(7027):741-745.
936
937   Meister M, Wong RO, Baylor DA & Shatz CJ.(1991) Synchronous bursts of action potentials in
938   ganglion cells of the developing mammalian retina. *Science* 252:939-43.
939
940
941   McCubbin J, Murphy P, Eswaran H, Preissl H, Yee T, Robinson SE & Vrba J.
942   (2007) Validation of the flash-evoked response from fetal MEG. *Phys Med Biol.* 52:5803-
943   13. Epub 2007 Sep 10.
944
945

946   Morin L.P., Blanchard J.H. & Provencio I. (2003) Retinal ganglion cell projections to the hamster
947   suprachiasmatic nucleus, intergeniculate leaflet, and visual midbrain: bifurcation and
948   melanopsin immunoreactivity. *J Comp Neurol* 465(3):401-416.

950   Müller LP, Do MT, Yau KW, He S and Baldridge WH. (2010) Tracer coupling of intrinsically
951   photosensitive retinal ganglion cells to amacrine cells in the mouse retina. *J Comp Neurol*.
952   518:4813-24.

953   Nagy JI, Pereda AE & Rash JE (2018). Electrical synapses in mammalian CNS: Past eras,
954   present focus and future directions*. Biochim Biophys Acta Biomembr.* 1860:102-123. doi:
955   10.1016/j.bbamem.2017.05.019. Epub 2017 Jun 1.

956   Newman LA, Walker MT, Brown RL, Cronin TW & Robinson PR (2003) Melanopsin forms a
957   functional short-wavelength photopigment *Biochemistry* 42:12734-8.

959   Nishimura A, Kitano K, Takasaki J, Taniguchi M, Mizuno N, Tago K, Hakoshima T, Itoh H.
960   (2010) Structural basis for the specific inhibition of heterotrimeric Gq protein by a small
961   molecule, *Proc Natl Acad Sci U S A* 107: 13666-13671.

963   Nguyen MT, Vemaraju S, Nayak G, Odaka Y, Buhr ED, Alonzo N, Tran U, Batie M, Upton BA,
964   Darvas M, Kozmik Z, Rao S, Hegde RS, Iuvone PM, Van Gelder RN & Lang RA. (2019)
965   An opsin-5 dopamine pathway mediates light-dependent vascular development in the eye.
966   *Nat Cell Biol.* 21:420-429. doi: 10.1038/s41556-019-0301-x. Epub 2019 Apr 1.

968   Panda S, Provencio I, Tu DC, Pires SS, Rollag MD, Castrucci AM, Pletcher MT, Sato TK,
969   Wiltshire T, Andahazy M, Kay SA, Van Gelder RN & Hogenesch JB. (2003) Melanopsin is
970   required for non-image-forming photic responses in blind mice. *Science* 301(5632):525-
971   527.

973   Panda S, Sato TK, Castrucci AM, Rollag MD, DeGrip WJ, Hogenesch JB, Provencio I & Kay
974   SA. (2002) Melanopsin (Opn4) requirement for normal light-induced circadian phase
975   shifting. *Science* 298(5601):2213-2216.

977   Panda S, Nayak SK, Campo B, Walker JR, Hogenesch JB & Jegla T. (2005) Illumination of the
978   melanopsin signaling pathway. *Science* 307(5709):600-604.

980   Provencio I., Jiang G., De Grip W.J., Hayes W.P. & Rollag M.D. (1998) Melanopsin: An opsin in
981   melanophores, brain, and eye. *Proc Natl Acad Sci U S A* 95(1):340-345

983   Qiu X, Kumbalasiri T, Carlson SM, Wong KY, Krishna V, Provencio I & Berson DM. (2005)
984   Induction of photosensitivity by heterologous expression of melanopsin. *Nature*
985   433(7027):745-749.
986

Rao S, Chun C, Fan J, Kofron JM, Yang MB, Hegde RS, Ferrara N, Copenhagen DR & Lang
RA. (2013) A direct and melanopsin-dependent fetal light response regulates mouse eye
development. *Nature* 494(7436):243-246.

Reifler AN, Chervenak AP, Dolikian ME, Benenati BA, Li BY, Wachter RD, Lynch AM,
Demertzis ZD, Meyers BS, Abufarha FS, Jaeckel ER, Flannery MP, Wong KY. (2015) All
Spiking, Sustained ON Displaced Amacrine cells receive Gap-Junction Input from
Melanopsin Ganglion Cells. *Curr Biol* 25:2878.

Renna JM, Wang S & Berson DM (2011) Light acts through melanopsin to alter retinal waves
and segregation of retinogeniculate afferents. *Nat Neurosci.* *14*:827-9.

Rörig B & Grantyn R (19914) Ligand- and voltage-gated ion channels are expressed by
embryonic mouse retinal neurones Neuroreport. 5:1197-1200.

Schrage R, Schmitz AL, Gaffal E, Annala S, Kehraus S, Wenzel D, Büllesbach KM, Bald T,
Inoue A, Shinjo Y, Galandrin S, Shridhar N, Hesse M, Grundmann M, Merten N,
Charpentier TH, Martz M, Butcher AJ, Slodczyk T, Armando S, Effern M, Namkung Y,
Jenkins L, Horn V, Stößel A, Dargatz H, Tietze D, Imhof D, Galés C, Drewke C, Müller CE,
Hölzel M, Milligan G, Tobin AB, Gomeza J, Dohlman HG, Sondek J, Harden TK, Bouvier
M, Laporte SA, Aoki J, Fleischmann BK, Mohr K, König GM, Tüting T, Kostenis E. (2015)
The experimental power of FR900359 to study Gq-regulated biological processes. *Nature*
communications 6:10156.

Sekaran S, Lupi D, Jones SL, Sheely CJ, Hattar S, Yau KW, Lucas RJ, Foster RG & Hankins
MW. (2005) Melanopsin-dependent photoreception provides earliest light detection in the
mammalian retina. *Curr Biol* 15(12):1099-1107.

Sekaran S, Foster RG, Lucas RJ, & Hankins MW (2003) Calcium imaging reveals a network of
intrinsically light-sensitive inner-retinal neurons. *Curr Biol* 13(15):1290-1298.

Schmid S, Guenther E. (1996) Developmental regulation of voltage-activated Na+ and Ca2+
currents in rat retinal ganglion cells. *Neuroreport.* 7:677-81.

Schmidt TM, Taniguchi K, Kofuji P. (2008) Intrinsic and extrinsic light responses in melanopsin-
expressing cells during mouse development *J Neurophysiol.* 100:371-84. doi:
10.1152/jn.00062.2008. Epub 2008 May 14

Shatz CJ & Stryker MP (1988) Prenatal tetrodotoxin infusion blocks segregation of
retinogeniculate afferents. *Science* 242:87-9;

Shatz CJ (1996) Emergence of order in visual system development. *Proc Natl Acad Sci USA*
23:602-8.

Singer JH, Mirotznik RR, Feller MB. (2001) Potentiation of L-type calcium channels reveals
nonsynaptic mechanisms that correlate spontaneous activity in the developing mammalian
retina. *J Neurosci.* 21:8514-22.
1034
1035
1036   Taniguchi M, Suzumura K, Nagai K, Kawasaki T, Takasaki J, Sekiguchi M, Moritani Y, Saito T,
1037   Hayashi K, Fujita S, Tsukamoto S & Suzuki K. (2004) YM-254890 analogues, novel cyclic
1038   depsipeptides with Galpha(q/11) inhibitory activity from Chromobacterium sp. QS3666.
1039   *Bioorg Med Chem* 12(12):3125-3133.
1040
1041   Tarttelin EE, Bellingham J, Bibb LC, Foster RG, Hankins MW, Gregory-Evans K, Gregory-Evans
1042   CY, Wells DJ & Lucas RJ. (2003) Expression of opsin genes early in ocular development
1043   of humans and mice. *Exp Eye Res* 76(3):393-396.
1044
1045   Tischbirek C, Birkner A, Jia H, Sakmann B, & Konnerth A. (2015) Deep two-photon brain
1046   imaging with a red-shifted fluorometric Ca2+ indicator. *Proc Natl Acad Sci U S A.*
1047   112:11377-82. doi: 10.1073/pnas.1514209112. Epub 2015 Aug 24.
1048
1049   Tu D*C,* Zhang D, Demas J, Slutsky EB, Provencio I, Holy TE & Van Gelder RN. (2005)
1050   Physiologic diversity and development of intrinsically photosensitive retinal ganglion cells.
1051   *Neuron* 48(6):987-999.
1052

1053   Vemaraju S, Nayak G, , Miller WE, Copenhagen DR & Lang RA (2019) Fetal stage melanopsin
1054   (OPN4) and GNAQ (Gαq) signaling regulates vascular development of the eye *BioRxiv*
1055   doi: https://doi.org/10.1101/537225

1056
1057   Veruki ML, Oltedal L & Hartveit E (2010). Electrical coupling and passive membrane properties
1058   of AII amacrine cells. *J. Neurophysiol* 103, 1456-66. doi: 10.1152/jn.01105.2009. Epub
1059   2010 Jan 20.
1060
1061
1062   Walton KD, Navarrete R (1991). Postnatal changes in motoneurone electrotonic coupling
1063   studied in the *in vitro* rat lumbar spinal cord *J Physiol.* 433, 283-305.
1064
1065   Wong KY (2012) A retinal ganglion cell that can signal irradiance continuously for 10 hours. *J*
1066   *Neurosci* 32(33):11478-11485.
1067
1068   Wettschureck N*,* Rütten H, Zywietz A, Gehring D, Wilkie TM, Chen J, Chien KR & Offermanns
1069   S. (2001) Absence of pressure overload induced myocardial hypertrophy after conditional
1070   inactivation of Galphaq/Galpha11 in cardiomyocytes. *Nat Med* 7(11):1236-1240.
1071
1072   Yang MB, Rao S, Copenhagen DR & Lang RA (2013) Length of day during early gestation as a
1073   predictor of risk for severe retinopathy of prematurity. *Ophthalmology* 120:2706-2713.

1074
1075 Yasui F, Miyazu M, Yoshida A, Naruse & Takai A. (2008) Examination of signalling pathways
1076   involved in muscarinic responses in bovine ciliary muscle using YM-254890, an inhibitor of
1077   the Gq/11 protein. *Br J Pharmacol.* 154(4):890-900. doi: 10.1038/bjp.2008.140. Epub 2008
1078   Apr 21.
1079
1080 Zariwala HA, Borghuis BG, Hoogland TM, Madisen L, Tian L, De Zeeuw CI, Zeng H, Looger LL,
1081   Svoboda K & Chen TW. (2012) A Cre-dependent GCaMP3 reporter mouse for neuronal
1082   imaging in vivo. *J Neurosci* 32(9):3131-3141.
1083
1084 Zhao Q, Guo H, Ruggu P, Jixiang, L, Liao J & Diwu, Z (2019) New Red Fluorescent Calcium
1085   Indicators for Functional Analysis of GPCRs and Ca2+ Channel Targets. *Biophysical*
1086   *Journal* (2019): 240a
1087
1088 Zheng W, Rampe D & Triggle DJ (1990) Pharmacological, radio ligand binding, and
1089   electrophysiological characteristics of FPL 64176, a novel nondihydropyridine Ca2+
1090   channel activator, in cardiac and vascular preparations. *Mol Pharmacol* 40:734-41.

1091
1092
1093
1094
1095   Biographies

1096
Jan Verweij acquired his doctorate in neuroscience at the University of Amsterdam under
the mentorship of Martin Kammermans, where he studied synaptic and electrotonic
interaction between rods and cones in goldfish retina]. He joined the laboratory of Denis
Dacey at the University of Washington, where he studied synaptic interactions in monkey
retina. Then, he joined the laboratory of Julie Schnapf at UCSF to study functional
interactions between rods and cones in monkey retina.

Shawnta Y. Chaney received her B.S. in Biology and her Ph.D. in Biochemistry at the
University of Houston, Houston TX. There, she studied the effects of gestational lead
(Pb2+) exposure on mouse retinas. She further studied retinal development as a
postdoctoral scholar in David Copenhagen's lab at UCSF, where she focussed on
embryonic and early postnatal light activation of melanopsin ganglion cells. Shawnta is now
an Ocular Scientist at UNITY Biotechnology, where she tests therapies for age-related
ocular diseases.

1110

1111 **Additional information**

1112 **Competing interests**

1113 The authors declare that they have no competing interests.

1114
1115 **Author contributions**

1116

1117 DRC and RAL provided project leadership. JV, SYC, DRC and DB and designed and performed

1118 experiments and analysed data. DRC wrote the manuscript. JV, SYC, DB, SV and RAL

1119 critiqued the study and edited the manuscript. GMK, EK provided crucial reagents and advice.

1120

1121 **Funding**
1122
Funding was provided by the following sources: National Eye Institute (R01 EY02636 to RAL

and DRC; R01 EY 01869 to DRC, R21 EY025435 to DRC, P30 EY002162 to UCSF)

Ophthalmology, Additional support was provided by That Man May See (San Francisco), an

unrestricted grant from Research to Prevent Blindness (UCSF Ophthalmology) and the German

research foundation- (DFG) funded research unit FOR2372, projects no. 1582/10-1, KO

1582/10-2 to E.K. and G.M.K.

1129

1130

1131

1132 **Acknowledgements**

1133 We thank Anton Delwig and Michael Stryker for valuable discussion and advice during the

1134 course of this research. We thank Andrew Ishida for his critique of the manuscript. For technical

1135 advice, we thank Franklin Caval-Holme, James Bothelo III, Travis Porco and Jessica Wong.

1136

1137

1138

Figure 1 Cells in foetal mouse retinas are light-responsive.

Cells in embryonic mouse retinas exhibit light-dependent increases in intracellular calcium. (A-C) Representative images of light-evoked fluorescence recorded from cells in an isolated, flat-mounted retina of an E17.5 mouse after loading of the retina with Calbryte-590 calcium-sensing dye. Images of a 340 x 430 mm region of the retina captured at T=0 s and at T = 2.5 s (A, B) are displayed. The colour-coded regions of interest (ROIs) show 4 cells in which light evoked a very discernible increase in fluorescence, indicating an increase in intracellular calcium. (C) Difference image of the two frames ($T_{2.5} - T_0$) highlighting the regions in which increased dye fluorescence was observed after light onset. (D) Plots of the temporal response profiles of the cells marked by ROIs. The ordinate plots dF/F, where F is the baseline fluorescence immediately after the onset of the stimulus and prior to the onset of the response. Stimulus light of 480 nm wavelength was applied at 4 x 10^{17} photons/cm^2/s. The scale bars in A, B and C indicate 50 μm.

Figure 2. The genetically encoded intracellular calcium indicator GCaMP3 reveals light responses of melanopsin RGCs in embryonic retinas. Retinas from melanopsin knockout mice were unresponsive to light.

GCaMP3-EGFP co-localizes with melanopsin and indicates light responsiveness of foetal mRGCs in *Opn4^{cre/+}*::Ai38 retinas. (A) A flat-mount retina from an *Opn4^{cre/+}*::*Ai38* postnatal day 7 (P7) pup showing its melanopsin immunoreactivity (red). (B) GCaMP3-EGFP-immunoreactive cells (green). (C) Merged image of (A) and (B). The white circles show co-labelled cells; the purple circles mark cells in which GCaMP3+ immunoreactivity is strong and melanopsin immunoreactivity is absent or very dim. White circles with an asterisk mark the occasional melanopsin+, GCaMP3-negative cells. These particular cells probably reflect failure of the GCaMP3 allele to recombine. (D) GCaMP3-EGFP immunoreactivity in a perpendicular cryosection of P30 retina. Immunoreactivity is observed in the ganglion cell layer (GC), the inner plexiform layer (IPL) and within a few structures in the outer plexiform layer (OPL). (E, F) Retinas isolated from embryonic pups were placed in a perfusion chamber and stimulated with 480 nm light. (E) GCaMP3 fluorescence at the onset of the light response in one representative E16.5 retina. (F)

Fluorescence recorded at the peak of the light response in the same retina. Visual comparison of the images in (E) and (F) shows qualitatively brighter images in (F). (G) Time course of light-induced increases in df/F for 4 light-responsive cells (numbered 1 to 4 and colour-coded). The thick black trace plots the average response of 79 light-responsive cells in this experiment. These plots allow a more quantitative comparison of light responses in individual mRGCs. The scale bars indicate 50 μm. (G inset) Deletion of melanopsin (genotype $Opn4^{cre/cre}::Ai38$) reveals, first, that melanopsin is required for light responses and, second, that the observed light-evoked increases in fluorescence do not result from unrecognized effects of light on GCaMP3-EGFP.

Figure 3. Antagonists that block the functions of the Gnaq, Gna11 and Gna14 alpha subunits of $G_{q/11}$-G proteins **of the $G_{q/11}$ family severely attenuate the light responses of mRGCs in foetal retinas.**

The effects of YM-254890 (200 nM) and FR900359 (1 μM), two different antagonists of Gnaq, Gna11 and Gna14, on light-evoked calcium elevations were tested in 4 explanted E15.5 and 3 explanted E16.5 OPN4cre::Ai96 retinas. (A) Time course of the experiment in an E16.5 retina showing 12-s light stimuli (red dots, 1-7) over the course of 6000 s. Responses to light stimuli 1-3 were recorded during 3300 s in control ACSF, and responses to light stimuli 4-7 were recorded after application of YM-254890. (B) Selected light responses from 4 mRGCs before and during perfusion with YM-254890. The scale bar shows df/F of 0.1. (C) Mean response amplitudes observed in 32 simultaneously recorded mRGCs in the same E15.5 retina analysed in (B). (D) Mean response amplitudes observed in 43 simultaneously recorded mRGCs in an E16.5 retina

perfused with medium containing 1 mM FR900359 (from ~400 to 1800 s). When dF/F responses were close to zero at 1400 seconds, the retina was perfused with 50 mM K-ACSF in which 47.5 mM NaCl was substituted for 47.5 mM KCl. The inclusion of KCl in the ACSF increased the calcium fluorescent signals to a level brighter than that induced by light before perfusion with FR9000359. All of the cells plotted in (C) and (D) had an initial criterion light response dF/F > 0.05. The error bars indicate the SEM.

Figure 4 Ensemble data confirming that YM254890 and FR900359 uniformly attenuate light responses in virtually all embryonic mRGCs.

 (A) Combined light-evoked dF/F's of 93 cells from E15.5 and E16.5 *OPN4cre::Ai96* retinas. All cells had a light-evoked dF/F > 0.05 in control conditions. The black dots indicate the lowest dF/F after 1500-2500 s incubation in 200 nM YM-254890 versus control dF/F for each cell. Prolonged incubation in 200 nM YM-254890 reduced the average dF/F of the light responses to 7% ± 11% that of the control responses (n=93). The data are best fitted by the linear regression y = 0.09x (solid red line). The dashed line indicates the predicted location of the dots if YM-254890 had no effect. (B) Combined light-evoked dF/F's of 219 cells from 3 embryonic day 16.5 retinas. All cells exhibited dF/F > 0.05 under control conditions. The black dots indicate the lowest dF/F after a 1300-3100 s incubation in 1 mM FR900359 versus control dF/F for each cell. Prolonged incubation in 1 mM FR900359 reduced the average dF/F of the light responses to 4% ± 18% that of the control responses (n=219). The data are best fitted by the linear regression y = 0.03x (solid red line). These results are highly significant (Wilcoxon Signed Rank Test for paired data [two-tailed P = $2.01948e^{-028}$ for YM and even smaller for FR]).

Figure 5 TTX variably but reversibly reduces the amplitude of light responses across the population of embryonic mRGCs.

(A) We measured the action of TTX on the fluorescent light responses of GCaMP6-expressing mRGCs in *OPN4cre::Ai96* embryonic retinas. Combined data from 4 E16.5 retinas are shown. The criteria for selecting each cell were that dF/F in control ACSF was > 0.05 and that the response amplitude after TTX washout recovered to between 80% and 120% of the control value. The open circles plot the recovery dF/F versus control dF/F for each of the 177 light-responsive mRGCs. The solid blue dots plot dF/F in 1 μM TTX versus control dF/F for each cell. The black line (y = 1.00 x) indicates the best-fitting linear regression line through the recovery data and the origin 0,0. The blue line (y = 0.36 x) is the best-fitting linear regression line for the TTX data. The differences between the results obtained during TTX treatment and under control conditions are highly significant (Wilcoxon Signed Rank Test for paired data, two-tailed P = $1.01e^{-46}$). (B) Histogram of the data shown in (A). The blue bars plot dF/F in TTX divided by the dF/F under control conditions, and the grey bars plot dF/F in recovery divided by the dF/F under control conditions.

Figure 6 The L-type calcium channel antagonist L-cis-diltiazem (LCD) reversibly attenuates light-dependent increases in calcium in E16.5 mRGCs

(A) dF/F versus time in control ACSF, in LCD and after return to ACSF. Plots show representative recordings for 5 of 48 light-responsive cells from a single E16.5 *OPN4cre::Ai96* retina. The bars at the bottom indicate the timing of the light stimuli. (B) Plots of dF/F in LCD and recovery versus dF/F in control for 31 of 48 light-responsive cells in one retina in which the post-washout values recovered to between 80 and 120% of the control values. The dotted lines show the best linear regression that fits the data (y = 1.06x and y = 0.05x for recovery and LCD, respectively). The data from the 5 cells shown in (A) are shown as red circles. The Wilcoxon Rank Test for paired data yields a two-tailed P-value of 9.31323e^{-10}, indicating that L-cis-diltiazem at 25 μM highly significantly diminishes light responses in mRGCs.

Figure 7 The L-type calcium channel agonist FPL64176 increases light- evoked dF/Fs recorded from embryonic mRGCs.

A) dF/F in 2 mM FPL64176 versus dF/F in control for 147 mRGCs combined from 2 E16.5 *OPN4cre::Ai96* retinas. The blue line represents the best linear regression fit: y = 1.51x. The black line represents the parity of response in control and FPL64176. The Wilcoxon Rank test for paired data yielded a two-tailed P-value of $1.17193e^{-32}$, indicating that dF/F is very significantly larger during exposure of the mRGCs to FPL64176 than under control conditions. No recovery data are shown, as FPL washout is known to be very poor (Bieda and Copenhagen, 2004).

Figure 8 Patch pipette recordings reveal that TTX-sensitive sodium currents in mRGCs increase in amplitude during maturation from E16.5 to P1-3.

A-F show examples of TTX-sensitive currents recorded from an mRGC in an E18.5 retina (*OPN4cre::Ai96*) under voltage clamp conditions. A) Currents evoked under control conditions with the voltage clamp protocol shown in E. B) Currents evoked after several minutes in 1 μm TTX. C) Currents evoked after washout of TTX. D) Difference current, Control - TTX. E) Voltage clamp step protocol. F) The black dots indicate the maximum inward current for each trace from D versus the step potential (measurement of INa(V). To survey the amplitudes of peak Ina(V) across multiple cells in retinas of 3 different ages without applying TTX, we used an analytical method in which the final current sample of the voltage step in A was subtracted from the lowest current amplitude across the entire voltage step. These values, plotted as red dots, yield a good estimate of peak INa(V), as shown by comparison with the results obtained in TTX (black dots). G) Peak sodium currents in mRGCs during embryonic maturation. The dots indicate peak INa(V) estimates for 10, 35 and 7 mRGCs recorded from E16.5, E18.5 and P1-3 retinas, respectively. The vertical bars indicate the average ± standard error for each age (320 ± 73, 674 ± 72 and 1986 ± 153 pA, respectively). The data include whole-cell and perforated recordings and were not corrected for series resistance. In summary, these findings demonstrate that voltage-activated sodium currents are evident in

mRGCs at least 4 days before birth and that the magnitude of these currents increases with age.

Light responses in an embryonic day 18.5 mRGC

Figure 9

Light responses recorded from an mRGC in the retina of a foetal (E18.5) mouse pup.

To select mRGCs for electrophysiological patch pipette recordings, we first surveyed the retina for light-induced calcium responses, which should be restricted to melanopsin cells (OPN4cre::Ai96). Then, using infrared Nomarski imaging, we searched for light-responsive mRGCs whose cell bodies were exposed on the inner surface of the retina. A and B show light-induced GCaMP6 increases in fluorescence in three cells. The images in A and B were obtained immediately after and 1.35 s after the onset of light stimulation, respectively. Although cells 1, 2 and 3 were light-responsive, the better-focussed cell 1 was targeted for electrophysiological recordings. C) Current clamp traces showing the responses of cell 1 to current injections to -15 and -5 pA from a holding current of -35 pA. The steps to -5 pA evoked repetitive spiking. D) Current-clamped light responses versus intensity from cell 1. The intensity of the 480-nm light stimulus, expressed as photons/cm^2/s, is shown on the right. The bottom bar shows the timing of the light. The asterisks denote presumed spontaneous waves.

Figure 10

MFA, a gap junction-blocking drug, and intracellular neurobiotin injection reveal gap junction coupling between mRGCs in embryonic retinas.

A) Meclofenamic acid (50 mM) strongly and reversibly reduces light responses in E16.5 mRGCs. (Top panel) Light-evoked dF/F of 81 fluorescent cells in 2 retinas containing mRGCs expressing GCaMP6 (OPN4cre::Ai96). dF/F in 50 mM MFA (blue dots) and following washout (black dots) was plotted versus control dF/F. The data include 81 of 142 light-responsive cells in which the response after recovery was at least 80% and no more than 120% of the control value (note that 142 cells responded to light onset under control conditions with dF/F's > 0.05 but that in many of these cells the response failed to recover to at least 80% of the control value). The data were fitted with lines passing through the origin. The best fitting linear regression lines were y = 1.04 x and y= 0.21 x for recovery and MFA, respectively. The two-tailed P-value, obtained using the Wilcoxon Rank Test for paired data, was 8.3 x 10^{-25}, indicating that MFA highly significantly attenuates light-evoked dF/Fs. MFA reversibly attenuated light responses in essentially all the mRGCs.

B) Example of a neurobiotin injection (red) into an E18.5 mRGC. Three somas were neurobiotin- and EGFP-positive (green), indicative of tracer migration from one GCaMP6-expressing mRGC to at least two other mRGCs (stars). Five other somas (dots) were neurobiotin-positive (red) but lacked EGFP immunoreactivity, implying that mRGCs are coupled to non-melanopsin cells. The axon of the filled cell continued well beyond the picture to the left. The white scale bar indicates 20 μm.