Title: Hippocampal Neuron Functional Phenotype Influences the Stability of Its Neural Code

Authors

Kinsky, Nathaniel R.[1*]

Mau, William[1]

Sullivan, David W.[1]

Levy, Samuel J. [1,2]

Ruesch, Evan A.[1]

Hasselmo, Michael E.[1]

[1]Center for Memory and Brain, Boston University, Commonwealth Ave., Boston, MA 02215,

USA.

[2]Graduate Program for Neuroscience

[3]Corresponding author: kinsky@bu.edu

ABSTRACT

Trajectory-dependent splitter neurons in the hippocampus encode information about a

rodent's prior trajectory during performance of a continuous alternation task. As such, they

provide valuable information for supporting memory-guided behavior. Here, we employed

single-photon calcium imaging in freely moving mice to investigate the emergence and fate of

trajectory-dependent activity through learning and mastery of a continuous spatial alternation

task. We found that the quality of trajectory-dependent information in hippocampal neurons

correlated with task performance. We thus hypothesized that, due to their utility, splitter neurons

would exhibit greater stability than would traditional place cells. We found that splitter neurons

were more likely to remain active and retained more consistent spatial information across

multiple days than did place cells. Furthermore, we found that both splitter neurons and place

cells emerged rapidly and maintained stable trajectory-dependent/spatial activity thereafter. Our

results suggest that a neuron's functional coding properties influence its long-term stability.

INTRODUCTION

Place cells in the hippocampus encode the current position of many different animals and humans (Ekstrom et al., 2003; Geva-Sagiv, Romani, Las, & Ulanovsky, 2016; Miller et al., 2013; Muller & Kubie, 1987; Muller, Kubie, & Ranck, 1987; Niediek & Bain, 2014; O'Keefe, 1976; O'Keefe & Dostrovsky, 1971) supporting the known role of the hippocampus in spatial memory and navigation across species (Morris, Garrud, Rawlins, & O'Keefe, 1982; Vorhees & Williams, 2014). However, the hippocampus is also widely known for its role in supporting the encoding, retrieval, and consolidation of non-spatial long-term memories (Corkin, 1984; Eichenbaum, 2004; Milner, Corkin, & Teuber, 1968), suggesting that it must represent variables beyond an animal's current location. Indeed, recent studies have demonstrated that the hippocampus encodes the dimensions of a given task, from odors (Muzzio et al., 2009; Wood, Dudchenko, & Eichenbaum, 1999) to time (Howard et al., 2014; Kraus, Robinson II, White, Eichenbaum, & Hasselmo, 2013; MacDonald, Lepage, Eden, & Eichenbaum, 2011; Manns, Howard, & Eichenbaum, 2007; Pastalkova, Itskov, Amarasingham, & Buzsáki, 2008; Robinson et al., 2017; Salz et al., 2016) to tones (Aronov, Nevers, & Tank, 2017). One early demonstration that the hippocampus encodes dimensions beyond an animal's current location was the discovery of trajectory-dependent neurons or splitter neurons (Frank, Brown, & Wilson, 2000; Wood, Dudchenko, Robitsek, & Eichenbaum, 2000), cells whose firing rate within a particular position was modulated based on the animal's past or future trajectory in a spatial alternation task. The generation of this neural correlate suggests a potential mechanism allowing the hippocampal code to support both memory and decision based planning.

Several studies have demonstrated place cell firing fields move, or remap, their locations in response to new learning during a spatial learning task (Dupret, O'Neill, Pleydell-Bouverie, &

Csicsvari, 2010; McKenzie, Robinson, Herrera, Churchill, & Eichenbaum, 2013). These studies

highlight that the flexible adjustment of place field locations is important for learning new

information. Conversely, the ability of hippocampal neurons to maintain the same firing location

in the absence of learning might support long-term memory retrieval. In support of this idea, a

recent study illustrated that neurons with place fields located near a hidden goal were more stable

over time than cells with fields in other locations (Zaremba et al., 2017). Two other experiments

found that increasing rodents' attention to a task selectively heightened stability in neurons that

encoded task-relevant features (Kentros, Agnihotri, Streater, Hawkins, & Kandel, 2004; Muzzio

et al., 2009). These studies, along with the finding that place cells with fields in close proximity

to a goal location exhibit heightened activity in post-learning sleep (Dupret et al., 2010), suggest

that the utility of a neuron's information to task performance influences its long-term stability.

Thus, since splitter neurons provide immediately relevant information for performing a

spatial alternation task, we hypothesized that these neurons are important for successful task

performance. Furthermore, we hypothesized that due to their utility, splitters may exhibit

different long-term dynamics when compared to place cells. Specifically, we addressed three

lines of inquiry. First, does the level of trajectory-dependent information within the hippocampus

correlate with behavioral performance? Second, given the steady evolution of hippocampal

activity patterns across days (Cai et al., 2016; Mau et al., 2018; Rubin, Geva, Sheintuch, & Ziv,

2015; Ziv et al., 2013), do splitter neurons remain part of the active population longer than other

cells, thus providing a longer lasting memory or planning signal to guide behavior? Third, once a

neuron establishes trajectory-dependent activity, is it less prone to remapping than other

neurons? These questions are particularly relevant since trajectory-dependent activity has been

observed in other tasks (Ferbinteanu & Shapiro, 2003; Smith & Mizumori, 2006b) and could be

employed more generally by the hippocampus to guide the appropriate behavior based on

environmental cues (Smith & Mizumori, 2006a).

To track neurons across long timescales, we paired a continuous spatial alternation task

with *in vivo* miniscope recordings of GCaMP6f activity in dorsal CA1 of freely-moving mice.

This technology allowed us to not only track the long-term activity of neurons, but also to

adequately characterize the heterogeneity of trajectory-dependent activity in the hippocampus,

since we can simultaneously record from a large number of neurons in each session. We first

found that trajectory-dependent coding correlates with task performance, suggesting it is

important for supporting memory-guided behavior. Second, we established that a neuron's

functional coding properties, also referred to as functional phenotype below, are important for

predicting its long-term activity: splitter neurons were more likely to be persistently active in the

days following their onset than were return arm place cells indicating that neurons which provide

more adaptive information might provide longer lasting input to downstream structures. Third,

we found that trajectory-dependent neurons display more consistent long-term information about

an animal's location than pure place cells. Fourth, we found that the population as a whole

displayed a rapid onset of trajectory-dependent activity followed by stable coding of trajectory

thereafter. Last, we discovered that recruitment of context-dependent splitter cells peaked several

days into training, whereas place cell recruitment peaked on the first day. These results combined

suggest that neurons which develop the most behaviorally important coding properties are

preferentially stabilized in both their short and long-term dynamics enabling them to more

consistently and effectively support memory-guided behavior. Our research paves the way for

future studies investigating how heterogeneity in the neural code might support acquisition and

retention of more complex behavioral tasks.

RESULTS

Behavior and Imaging

Food deprived mice (n=4) with neurons expressing GCaMP6f in region CA1 of the

dorsal hippocampus were trained to perform a continuous spatial alternation task on a figure-8

maze (Figure 1A) while we simultaneously recorded calcium activity using a miniaturized

microscope. Mice exhibited a range of learning rates, taking from 5 to 21 sessions to acquire the

task, which was defined as the third consecutive session of performance at or above our criteria

of 70% (Figure 1B). Mice performed continuous alternation at or greater than criteria on average

throughout the course of the experiment (Figure 1C). We utilized custom-written software

(Kinsky, Sullivan, Mau, Hasselmo, & Eichenbaum, 2018; Mau et al., 2018) to extract neuron

ROIs (Figure 1D), construct their corresponding calcium traces, and identify each ROI's putative

spiking activity (Figure 1E). Using this technique, we recorded from large numbers of neurons

(243-1205 neurons per ~30 minute-session) and successfully tracked them across days by

comparing the distance between neuron ROI centroids (Supplemental Figure 1A) and verifying

that ROIs did not change orientation of their major elliptical axis between sessions

(Supplemental Figure 1B).

Figure 1: Experimental Setup and Imaging

A) Alternation Maze. Blue = Left turn trajectories, Red = Right turn trajectories, *= location of food
reward.
B) Example learning curve for one mouse. Red dashed = acquisition criterion (70%), red asterisk =
task acquisition day.
C) Performance summary for all four mice, all sessions included. Red dashed = criterion, black
dashed = chance.
D) Example maximum projection from one imaging session with 10 neuron ROIs overlaid.
E) Example calcium traces for ROIs depicted in D. Red lines on the ascending phase of each
calcium event indicate inferred spiking activity.
126

Trajectory-Dependent Activity is Maintained Across Days

The initial studies establishing the existence of trajectory-dependent splitter cells in the

hippocampus were performed using electrophysiology in rats (Frank et al., 2000; Wood et al.,

2000). Thus, we first wondered if we could detect trajectory-dependent activity in a different

species while using a technique with much lower temporal resolution. To do so, we constructed

tuning curves representing the probability a given neuron had calcium activity at each spatial bin

(1cm) along the stem in correct trials only, and classified neurons as trajectory-dependent

splitters if at least 3 bins displayed a significant difference between their tuning curves ($p < 0.05$,

permutation test). We found that we were capable of not only identifying trajectory-dependent

cells on a given day (60 ± 23, mean $\pm$ s.e.m. across all four mice), but that in many cases these

neurons maintained the same functional phenotype across multiple days (Figure 2A-B).

Significant trajectory-dependent activity was exhibited by 10% of neurons active on the maze

stem across all sessions (12%, 5%, 12%, and 9% for individual mice); note that this method for

identifying trajectory-dependent activity is more conservative than that used in previous studies

(Frank et al., 2000; Ito, Zhang, Witter, Moser, & Moser, 2015; Wood et al., 2000). Apparent

trajectory-dependent activity could also potentially result from factors such as systematic

variations in the mouse's lateral position along the stem. We addressed this in two ways. First,

we limited the portion of the maze we considered the stem to exclude any areas where the mouse

exhibited stereotypical turning behavior by eye (Figure 2A-B, bottom). Second, we performed an

ANOVA for each splitter neuron which included the animal's upcoming trajectory, position

along the stem, speed, and lateral position along the stem as covariates (Wood et al., 2000). We

found that a high proportion of our splitter neurons were significantly modulated by upcoming

turn direction after accounting for speed and lateral stem position (89%, 72%, 76%, and 83% for

individual mice). Together, these results indicate that trajectory-dependent coding exists in

mouse CA1 and in many cases maintains the same activity profile across both short and long

timescales. To the best of our knowledge, this is the first demonstration of hippocampal

trajectory-dependent activity using calcium imaging in mice.

Additionally, we observed the mean location of spatial firing along the stem progressed

backward during the task such that calcium activity occurred at earlier and earlier portions of the

stem with time (Supplemental Figure 2A). This is consistent with a study reporting the

backwards-migration of spatial firing with experience (Mehta, Quirk, & Wilson, 2000).

Interestingly, we did not find any evidence of consistent migration of spatial firing locations

between sessions (Supplemental Figure 2B).

Figure 2: Trajectory-Dependent Activity Persists Across Days

A) *Top:* Calcium event rasters along the stem for correct trials for two sessions recorded one day
163   apart, sorted by turn direction at the end of the stem. Blue = left, Red = right. *Middle:* Calcium
164   event probability curves for each turn direction.*p<0.05, shuffle-test. *Bottom:* occupancy
165   normalized calcium event map with reliability (REL) and discriminability (DIS) scores shown.
166   Red dashed = extent of stem considered in above plots.
B) Same as A, but for a different mouse and for sessions 7 days apart.

Trajectory-Dependent Activity Correlates with Performance

Trajectory-dependent neurons provide information vital to task performance that might

be utilized by downstream structures to inform proper motor actions (Albouy et al., 2008; Kahn

et al., 2017; Wise & Murray, 1999). This idea is supported by a study which found that

trajectory-dependent activity markedly diminished during error trials (Ferbinteanu & Shapiro,

2003). Thus, we predicted that successful task performance would be associated with prominent

trajectory-dependent information in the neural code of neurons active on the stem. We utilized

two metrics to measure different attributes of trajectory-dependent activity: 1) reliability, which

measured the consistency of a cell to fire on its preferred trial type along the entire stem, and 2)

discriminability, which measured the magnitude of difference between left and right turn tuning

curves along the entire stem (Methods). While most splitter neurons generally had high

reliability and discriminability values, neurons with sparser calcium activity for one turn

direction could exhibit low reliability and high discriminability (Figure 2A). Conversely, splitter

neurons that reliably increased their event rate for one turn direction but still exhibited activity

for the other turn direction could have high reliability but low discriminability (Figure 2B).

We obtained significant correlations between each metric and the animal's performance

in a given session (Figure 3A, B top) suggesting that trajectory-information carried in

hippocampal neurons might support working memory. To bolster this argument, we also trained

a decoder to classify future turn direction using a linear discriminant analysis (LDA) at each

spatial bin along the stem based on the neural activity of the population. We found that the

accuracy of the LDA decoder positively correlated with the animal's performance on a given

day, which indicated that better separation between upcoming left and right trajectories by the

neural code was related to increased memory (Figure 3C, top). These results held for

discriminability and LDA accuracy, but not reliability, when we averaged performance across

mice (Figure 3A-C, bottom). Last, for each cell, we correlated the left turn and right turn tuning

curves and subtracted those values from 1 $(1 - \rho)$ as another metric for trajectory-dependent

information. This metric is very conservative because it produces low values (indicating high-

trajectory dependent information) for splitter neurons that shift their location along the stem

between trial types but not for splitter neurons that modulate event rates in the same location.

Despite this we found a significant relationship between performance and $1 - \rho$ when averaging

within mice (Figure 3D, bottom). We obtained similar results when we focused on local metrics

of trajectory-dependent activity rather than their average along the entire stem (Supplemental

Figure 3). Together, these results indicate that trajectory-dependent information might facilitate

accurate task performance.

Figure 3: The Quality of Trajectory-Dependent Activity Correlates with Performance

A) *Top:* Performance for each session versus the mean reliability value for all cells from that session.
Circles = all sessions, all mice.*ρ=0.47, p=1.2x10^{-4} Pearson correlation. *Bottom:* Same as *Top* but
for each mouse, triangles = average for each mouse.
B) Same as A, but for the mean discriminability value. *ρ=-0.45, p=0.0043, **ρ=-0.98, p=0.012
(bottom), Pearson correlation.
C) Same as A, but for the mean LDA decoder accuracy. *ρ=0.25, p=0.047, **ρ=0.94, p=0.003
Pearson correlation.
D) Same as A, but for the mean value of $1 -$ Spearman's ρ between left and right tuning curves.
*ρ=0.91, p=0.046.

Trajectory-Dependent Neurons are more likely to Remain Active over Long Timescales than

Arm Place Cells

Multiple studies (Cai et al., 2016; Kinsky et al., 2018; Mau et al., 2018; Rubin et al.,

2015; Ziv et al., 2013) have shown that hippocampal neurons exhibit significant turnover across

days with fewer staying active within the same environment as time progresses. However, these

studies all treated the CA1 population as one homogeneous group. Thus, we wondered if splitter

neurons, which exhibit highly task relevant information, would be preferentially stabilized within

the CA1 network when compared to traditional place cells. As such the hippocampus would

maintain a more consistent population of neurons which could be utilized for guiding this

behavior. To address this question, we calculated the probability that each pool of neurons

remained active in subsequent sessions. We found that splitter cells were more likely to remain

active in a later session than arm place cells for both short (Figure 4A) and long (Figure 4B) time

lags between sessions. This heightened likelihood that splitter neurons remained active persisted

up to 15 days later when considering all mice together (Figure 4C-D) and also held when we

compared splitters to place cells with activity on the stem (stem place cells, Figure 4E-F). To

mitigate any sampling biases due to the higher event rate of splitter neurons (Supplemental

Figure 4A), we performed an additional analysis where we included only the most active place

cells such that their mean event rate matched that of splitters. We obtained similar results when

comparing splitter neurons to arm place cells but not stem place cells (Supplemental Figure 4B-

F). Since many stem place cells exhibited trajectory-dependent activity that fell short of meeting

our splitter neuron criteria, this suggests that a neuron's activity level, along with its functional

phenotype, also influences whether it stays active on following days. These findings combined

support the idea that the strength of task relevant information carried by a neuron influences its

 likelihood to maintain activity at later time points, which could be exploited for successful

 memory-guided behavior across days.

Figure 4: Splitter Neurons are more likely to Remain Active between Sessions than Arm Place Cells

A) Probability splitters and arm place cells (PCs) stay active day later for all mouse. *p=6.4x10^{-11},
one-sided signed-rank test.
B) Same as B but for all mice and for sessions 7 days apart. *p = 1.1x10^{-10}, one-sided signed-rank
test.
C) Probability splitters and arm place cells stay active versus lag between sessions. Dots:
probabilities from individual session-pairs, lines: mean probability at each time lag. Green =
splitters, Red = arm PCs.
D) Difference between the probability that splitters stay active versus the probability that arm PCs
stay active between sessions. Dots: probability differences for individual session-pairs. Black
solid/dashed lines: Mean and 25%/75% quantiles of data at each time point. Red bars =
significant differences after Holm-Bonferroni correction of one-sided signed-rank test. See Table
1 for signed-rank test p-values at all lags, α = 0.05.
E) and F) Same as C-D for splitter neurons vs. stem PCs.

Table 1: One-sided Signed-Rank Significance Values for Probability Splitter vs. Place Cells (PCs)

Remain Active between Sessions. PCs subsampled to match mean event rate of splitter neurons.

Lag (days)	1	2	3	4	5	6	7
vs. Arm PCs	1.3e-10	9.1e-7	9.4e-5	4.3e-5	2.2e-4	4.5e-7	2.7e-8
vs. Stem PCs	2.e-9	5.3e-6	0.016	1.5e-3	0.017	5.4e-6	1.7e-4
Lag (days)	8	9	10	11	12	13	14
vs. Arm PCs	1.2e-5	1.0e-4	4.9e-3	0.014	1.2e-4	2.5e-5	8.9e-4
vs. Stem PCs	1.1e-4	1.7e-3	0.082	0.097	5.2e-3	3.3e-4	0.012
Lag (days)	15	16					
vs. Arm PCs	1.2e-4	9.8e-3					
vs. Stem PCs	6.7e-3	2.0e-3					

We next wondered how the information provided by splitter cells differs from that of

other neuron functional phenotypes. To investigate, we decided to compare the long-term spatial

coding properties of trajectory-dependent splitter neurons on the stem to return arm place cells

(Figure 5A). When examining spatial calcium activity over the entire map across sessions, we

found that splitter neurons had a significantly higher 2D event map correlation values than arm

place cells (Figure 5B) and that this effect persisted up to 15 days later (Figure 5C) indicating

that they were more stable overall. We observed similar results when we compared splitter

neurons to stem place cells (Table 2, Supplemental Figure 5). This indicates that, in addition to

staying active over longer time-scales, trajectory-dependent splitter neurons might also better

guide memory task performance by providing a more consistent representation of space than

place cells.

Figure 5: Splitters Maintain More Consistent Spatial Information than Place Cells

A) Example 2D occupancy normalized calcium event maps from the same splitter neuron (top) and arm place cell (bottom) between sessions on adjacent days. The higher Spearman correlation between splitter neuron event maps indicates more consistent spatial activity.

B) Mean spatial correlations for splitter neurons versus return arm PCs for all sessions seven days apart
from all. *p=4.5x10^{-6}, one-sided signed-rank test.
C) Mean spatial correlations for splitter neurons and arm PCs versus lag between sessions for all
mice/sessions. Red = Arm PCs, green = splitters. Red bars = significant differences after Holm-
Bonferroni correction of one-sided signed-rank test, α = 0.05. See Table 2 for raw p-values at all lags.

Table 2: One-sided Signed-Rank Significance Values for Mean Spatial Correlation Values of Splitter vs. Arm PCs or

Splitters vs. Stem PCs.

Lag (days)	1	2	3	4	5	6	7
vs. Arm PCs	8.6e-8	5.0e-4	1.7e-3	4.6e-3	4.5e-3	2.6e-4	2.6e-3
vs. Stem PCs	4.7e-6	4.5e-5	7.5e-4	5.7e-5	5.7e-3	3.2e-3	1.2e-3
Lag (days)	8	9	10	11	12	13	14
vs. Arm PCs	0.059	0.032	0.013	0.064	0.015	3.2e-4	0.021
vs. Stem PCs	0.23	6.2e-3	1.2e-4	0.024	0.59	1.4e-3	0.44
Lag (days)	15	16					
vs. Arm PCs	0.15	0.78					
vs. Stem PCs	0.047	0.25					

Trajectory-Dependent Neurons Display a Rapid Onset Followed by Stable Activity

Next, we examined the ontogeny of trajectory-dependent behavior. We hypothesized two

different scenarios could support the emergence of splitters. In line with a study showing that

unstable neurons can support well-learned behavior (Driscoll, Pettit, Minderer, Chettih, &

Harvey, 2017), splitters could slowly ramp up/down their splitting behavior or they could come

online suddenly and turn off just as suddenly. On the other hand, previous research presented the

idea that neurons pre-disposed to become place cells can come online suddenly after a head-

scanning/attention event (Monaco, Rao, Roth, & Knierim, 2014), which is potentially supported

by the presence of reliable sub-threshold depolarizations of those neurons caused by calcium

activity in its dendritic arbor (Bittner, Milstein, Grienberger, Romani, & Magee, 2017;

Diamantaki et al., 2018; Lee, Lin, & Lee, 2012; Sheffield & Dombeck, 2014, 2019). In line with

this idea, splitter neurons could rapidly develop trajectory-dependent activity and then maintain

that activity thereafter. To address this question, we identified the day when each neuron we

recorded first exhibited significant trajectory-dependent activity, and then tracked whether that

neuron retained trajectory-dependent activity along a similar proportion of the maze stem

(splitting extent) in subsequent sessions. We found evidence for heterogeneity in the ontogeny of

splitting, with some neurons exhibiting a rapid onset of trajectory-dependent activity (Figure 6A)

while others ramped up their trajectory-dependent activity in the days prior to becoming a splitter

(Figure 6B). Each onset type appeared to maintain stable trajectory-dependent activity afterward

since, for individual mice, splitting extent remained higher in the day following splitter onset

when compared to the day preceding splitter onset (Figure 6C). The rapid onset of trajectory-

dependent activity and stable maintenance thereafter was readily apparent when examining

group data over longer time scales (+/- 10 days, Figure 6D). We obtained similar results for peak

reliability and peak discriminability along the stem (Figure 6E-F). In contrast, we observed only

a weak trend for reliability and discriminability averaged along the whole stem (Supplemental

Figure 6), supporting the observation that splitter neurons maintained consistent trajectory-

dependent activity along a local portion (~10-20%) of the stem after their onset. We observed a

similar trend for place cells using mutual information as a metric of place coding strength

(Figure 6G), suggesting that similar rules govern the onset and fate of trajectory-dependent and

spatial coding in hippocampal neurons.

Figure 6: Splitters Come Online Abruptly and Maintain Stable Fields

A) and B) Example splitter across days for two different mice illustrating sudden onset of trajectory-
dependent activity followed by stable trajectory-dependent activity thereafter.
C) Extent of significant splitting (trajectory-dependent activity) along the stem +/- 1 days from
splitter onset for one representative mouse. p = 7.6x10^{-19} Kruskal-Wallis ANOVA, *p < 1.0e-5,
**p=0.00025 post-hoc Tukey test. Circles = splitting extent for each neuron.
D) Extent of splitting +/- 10 days from splitter onset for all mice. p = 3.1x10^{-14} Kruskal-Wallis
ANOVA, *p < 6.0x10^{-5} post-hoc Tukey test. Circles = mean splitting extent of all neurons active
on the stem for each session.
E) Max reliability score +/- 10 days from splitter onset for all mice. p = 2.8x10^{-12} Kruskal-Wallis
ANOVA, *p < 2x10^{-4} post-hoc Tukey test.
F) Mean discriminability score +/- 10 days from splitter onset for all mice. p = 4.7x10^{-6} Kruskal-
Wallis ANOVA, *p < 0.003 post-hoc Tukey test.
G) Mean mutual information +/- 10 days from return place cell onset for all mice. We obtained
similar results for stem place cell onset (data not shown). p = 4.7x10^{-17} Kruskal-Wallis ANOVA,
*p < 6x10^{-4} post-hoc Tukey test.

Place Cell Onset Coincides With or Precedes Splitter Onset

We next wondered if hippocampal neurons displayed significant spatial tuning before,

during, or after they exhibited trajectory-dependent firing. As shown above, splitter cells produce

accurate spatially-modulated activity (Figure 5) and have a similar onset/offset trajectory to place

cells (Figure 6); thus, we hypothesized that the onset of trajectory-dependent firing in

hippocampal neurons would either coincide with or follow their onset as place cells. To test this

idea, we first tallied the onset day of each cell phenotype. We found, while both functional

phenotypes were present from day 1 and continued to come online throughout the experiment

(Figure 7A,C), the bulk of place cells were recruited on day 1. In contrast, and in agreement with

a previous study (Bower, Euston, & McNaughton, 2005), the recruitment of splitter cells did not

peak until several days later (Figure 7A,C), suggesting that trajectory-dependent activity tended

to emerge more slowly than spatial activity. This could occur independently in two different

groups of neurons, or it could occur serially with each neuron first becoming a splitter cell only

after becoming a place cell. Thus, to test if this delay in splitter cell ontogeny occurred in the

same cells, we directly compared the day a cell became a place cell to the day it began to exhibit

trajectory-dependent activity. We found that in the majority of neurons, trajectory-dependent

activity onset occurred simultaneously with place field onset, while a different population of

neurons exhibited trajectory-dependent activity only after first becoming place cells (Figure

7B,D). Thus, place cells and splitter cells occupy an overlapping population of neurons with

spatial responsivity coinciding with or preceding trajectory-dependent coding.

Figure 7: Place Cell Onset Coincides With or Precedes Splitter Onset

    A) Histogram of the first (onset) day a neuron exhibits a splitter cell or place cell phenotype for one mouse. *p = 3.7×10^{-17}, one-sided Kolmogorov-Smirnov test.

    B) Difference between splitter cell onset day and place cell onset day for one mouse. *p=2.5×10^{-24} χ^2 goodness-of-fit test, mean = 2.3, median = 2.

    C) Same as A) but for all mice. *p = 2.2×10^{-17}, one-sided Kolmogorov-Smirnov test.

    D) Same as B) but for all mice. *p=5.3×10^{-107} χ^2 goodness-of-fit test, mean = 1.6, median = 1.

DISCUSSION

From an evolutionary perspective, one adaptive function of memory is the ability to provide information vital to survival. Thus, maintaining activity and consistency in neurons encoding information pertinent to survival might provide a mechanism for preferentially strengthening connections with downstream structures via consistent replay of the same sequences (Buzsáki, 2015; Diba & Buzsáki, 2007; Louie & Wilson, 2001; Maboudi et al., 2018; Pfeiffer & Foster, 2013). Conversely, if the pool of neurons available to encode a given memory remains fixed, then forgetting of incidental information through the turnover/silencing of neurons

not required for survival is adaptive (Hardt, Nader, & Nadel, 2013) because it could increase the

numbers of neurons available to encode other relevant information (Richards & Frankland,

2017). Here, we utilized *in vivo* calcium imaging with miniaturized microscopes to explore this

idea (Figure 1) by investigating the development and fate of trajectory-dependent splitter neurons

(Frank et al., 2000; Wood et al., 2000) (Figure 2). To the best of our knowledge, this is the first

demonstration that trajectory-dependent hippocampal activity exists in mice and that it can be

detected with calcium imaging. Since trajectory-dependent splitter neurons contain information

relevant to proper task performance (Figure 3, see also Ferbinteanu & Shapiro, 2003), we

hypothesized that they would exhibit relatively high stability when compared to other neuron

phenotypes.

Several lines of evidence support this hypothesis. First, splitter neurons are more likely to

remain active across long time scales than neurons that only provide information about the

animal's current location on the return arm (Figure 4). Second, splitters come online abruptly and

then maintain a stable readout of trajectory up to 10 days after becoming a splitter (Figure 6).

Splitters also provide a more consistent signal of the animal's current location than do other

neurons (Figure 5), further supporting their long-term stability. Last, we found that splitter cells

are a dynamic subpopulation of place cells with the onset of place coding generally preceding the

onset of trajectory-dependent activity (Figure 7). This finding concurs with the slow increase of

trajectory-dependent activity with experience found in a previous study (Bower et al., 2005).

These data combined support the idea that neuron phenotype influences its subsequent stability

(Zaremba et al., 2017) and the consistency of the information it provides to downstream

structures. More broadly, this study supports the idea that adaptive memories are encoded in a

stable subpopulation of neurons, freeing the remaining pool of neurons to undergo plasticity

during new learning (Grosmark & Buzsáki, 2016; van de Ven, Trouche, McNamara, Allen, &

Dupret, 2016).

Our study utilizes single-photon imaging to perform longitudinal tracking of hippocampal

neuron activity and confirms existing studies that show increasing turnover of coactive neurons

with time (Cai et al., 2016; Rubin et al., 2015; Ziv et al., 2013). However, a recent study

performed in songbirds demonstrated that imaging artifacts, specifically small shifts in the z-

plane of single-photon imaging, could entirely account for putative cell turnover (Katlowitz,

Picardo, & Long, 2018). Thus, the turnover we and others observe in hippocampal neurons could

likewise be artefactual. While relevant, this concern is mitigated in our study for a number of

reasons. First, the Katlowitz et al. (2018) study was performed in the basal ganglia of songbirds

while they performed a stereotyped behavior supported by highly stable firing responses of

neurons over short and long timescales (Guitchounts, Markowitz, Liberti, & Gardner, 2013;

Hahnloser, Kozhevnikov, & Fee, 2002; Margoliash & Yu, 2009). In contrast, our study was

performed in CA1 of the mouse hippocampus, a highly plastic brain region exhibiting complete,

monthly turnover of afferent connections (Attardo, Fitzgerald, & Schnitzer, 2015; Pfeiffer et al.,

2018) that also exhibits a high degree of drift in neuron firing responses over relatively short

time-scales (Mankin et al., 2012; Manns et al., 2007). Second, studies utilizing activity-

dependent tagging of neurons also find that the overlap between active cells in the mouse

hippocampus declines with time between sessions (Cai et al., 2016; Kitamura et al., 2017),

supporting long-term hippocampal cell turnover as a real phenomenon. Most importantly, our

study compares the *relative* turnover rates of two different cell phenotypes: splitter cells and

place cells. Thus, even if day-to-day misalignments in the z-plane forced neurons out of focus,

this would occur equally for both splitters and place cells. Therefore, concerns about imaging

artifacts cannot explain our finding that splitter cells are more persistently active across long time

scales than place cells.

One notable study found that lesions or optogenetic silencing of nucleus reuniens, an

important communication hub between the medial prefrontal cortex and dorsal CA1 of the

hippocampus, significantly reduced trajectory-dependent activity in rat CA1 neurons while

having no impact on a rat's performance of a spatial alternation task (Ito et al., 2015). Those

results directly challenge our finding that the quality of trajectory-dependent information

contained in CA1 activity patterns correlates with a mouse's performance (Figure 3). One

potential reason for this discrepancy is that their intervention only partially reduced trajectory-

dependent information without eliminating it, allowing the splitter cells remaining to provide

adequate information for proper task performance. In fact, optogenetic silencing of nucleus

reuniens produced a smaller deficit in trajectory-dependent activity than did lesions; even lesions

eliminated trajectory-dependent activity predicting future trajectories only. Information related to

past trajectories, which could be utilized by downstream structures to help make the correct

upcoming turn, was maintained. Second, relatively easy tasks might be less resistant to a partial

disruption and rats performed at close to ceiling levels in the Ito et al. (2015) study. Our mice

performed at lower levels, though still well above chance, indicating that the spatial alternation

task might place higher attentional and cognitive demands on mice than on rats. Last, Ito et al.

(2015) also utilized the difference in peak firing rate on left versus right trials as a metric for

trajectory-dependent activity. This calculation does not account for trajectory-dependent

information provided by neurons that maintain similar firing rates, but shift their firing location

along the stem between left and right trials (see Figure 2B). Thus, trajectory-dependent neural

activity could still be important for proper task performance.

Rodents with hippocampal lesions are capable of performing a continuous alternation task

(Ainge, van der Meer, Langston, & Wood, 2007). This raises the question: how important is

trajectory-dependent activity if mice can perform the task without the hippocampus at all? We

have two responses to this question. First, long-term lesions test necessity, not sufficiency, since

these lesions can induce compensatory plasticity that could allow non-hippocampal regions to

support the task (Packard & McGaugh, 1996). Second, under normal conditions the

hippocampus might still be the default brain region for task performance in spatial alternation.

This is emphasized by Goshen et al. (2011), who demonstrated that mice cannot perform long-

term recall of a putatively hippocampal-independent contextual fear memory (Bontempi,

Laurent-Demir, Destrade, & Jaffard, 1999; Debiec, LeDoux, & Nader, 2002; Frankland,

Bontempi, Talton, Kaczmarek, & Silva, 2004; Kim & Fanselow, 1992; Kitamura et al., 2017,

2009; Winocur, Frankland, Sekeres, Fogel, & Moscovitch, 2009) when hippocampal inactivation

is limited to a short time period before the task; however, mice became capable of successful

long-term memory recall when this inactivation was extended over a long time period prior to

performing the task. This study and others (Meira et al., 2018; Sparks, Lehmann, Hernandez, &

Sutherland, 2011; Sutherland, O'Brien, & Lehmann, 2008; Wang, Teixeira, Wheeler, &

Frankland, 2009; Wiltgen et al., 2010) support the idea that the hippocampus is vital for long-

term recall under normal conditions and that redundant pathways are recruited for episodic

memory retrieval only if chronic aberrant activity is detected in the hippocampus.

Through what mechanism do trajectory-dependent neurons maintain a greater stability

across long time-scales? After the initial onset of trajectory-dependent behavior, these neurons

could receive feedback from dopaminergic neurons originating in the ventral tegmental area

(VTA) during learning (Gomperts, Kloosterman, & Wilson, 2015) or from locus coeruleus (LC)

neurons during post-learning sleep (Takeuchi et al., 2016) that could strengthen afferent

connections to splitter neurons. This could also occur during sharp-wave ripple related replay of

prior trajectories (Diba & Buzsáki, 2007; Pfeiffer & Foster, 2013) in conjunction with

simultaneous dopaminergic inputs from VTA (Gomperts et al., 2015). However, this mechanism

would also strengthen all cells active en route to the goal location whether they carried

information about trajectory or not. One recent study found that trajectories leading to larger

rewards were preferentially replayed over trajectories leading to smaller rewards (Michon, Sun,

Kim, Ciliberti, & Kloosterman, 2019). Thus, one possibility is that since trajectory-dependent

neurons are more useful for predicting how to obtain reward than pure place cells, they might be

preferentially reactivated during sharp-wave ripple events, an idea that warrants future testing.

Taken together, our results highlight the influence of cell phenotype on its subsequent

stability, and suggest that the emergence of task-related trajectory-dependent coding coincides

with or follows the emergence of spatial coding in neurons. Future work should investigate

mechanisms supporting the stability and emergence of trajectory-dependent neurons.

REFERENCES

Ainge, J. A., van der Meer, M. a a, Langston, R. F., & Wood, E. R. (2007). Exploring the role of

context-dependent hippocampal activity in spatial alternation behavior. *Hippocampus*,

17(10), 988–1002. https://doi.org/10.1002/hipo.20301

Albouy, G., Sterpenich, V., Balteau, E., Vandewalle, G., Desseilles, M., Dang-Vu, T., …

Maquet, P. (2008). Both the Hippocampus and Striatum Are Involved in Consolidation of

Motor Sequence Memory. *Neuron*, *58*(2), 261–272.

https://doi.org/10.1016/j.neuron.2008.02.008

Aronov, D., Nevers, R., & Tank, D. W. (2017). Mapping of a non-spatial dimension by the

hippocampal–entorhinal circuit. *Nature*, *543*(7647), 719–722.

https://doi.org/10.1038/nature21692

Attardo, A., Fitzgerald, J. E., & Schnitzer, M. J. (2015). Impermanence of dendritic spines in live

adult CA1 hippocampus. *Nature*, *523*(7562), 592–596. https://doi.org/10.1038/nature14467

Bittner, K. C., Milstein, A. D., Grienberger, C., Romani, S., & Magee, J. C. (2017). Behavioral

time scale synaptic plasticity underlies CA1 place fields. *Science*, *357*(6355), 1033–1036.

https://doi.org/10.1126/science.aan3846

Bontempi, B., Laurent-Demir, C., Destrade, C., & Jaffard, R. (1999). Time-dependent

reorganization of brain circuitry underlying long-term memory storage. *Nature*, *400*(6745),

671–675. https://doi.org/10.1038/23270

Bower, M. R., Euston, D. R., & McNaughton, B. L. (2005). Sequential-Context-Dependent

Hippocampal Activity Is Not Necessary to Learn Sequences with Repeated Elements. *The*

Journal of Neuroscience, *25*(6), 1313–1323. https://doi.org/10.1523/JNEUROSCI.2901-

04.2005

Buzsáki, G. (2015). Hippocampal sharp wave-ripple: A cognitive biomarker for episodic

memory and planning. *Hippocampus*, *25*(10), 1073–1188.

https://doi.org/10.1002/hipo.22488

Cai, D. J., Aharoni, D., Shuman, T., Shobe, J., Biane, J., Lou, J., … Silva, A. J. (2016). A shared

neural ensemble links distinct contextual memories encoded close in time. *Nature*, *534*,

115–118. https://doi.org/10.1038/nature17955

Corkin, S. (1984). Lasting Consequences of Bilateral Medial Temporal Lobectomy: Clinical

Course and Experimental Findings in H.M. *Seminars in Neurology*.

https://doi.org/10.1055/s-2008-1041556

Debiec, J., LeDoux, J. E., & Nader, K. (2002). Cellular and systems reconsolidation in the

hippocampus. *Neuron*, *36*(3), 527–538. Retrieved from

http://www.ncbi.nlm.nih.gov/pubmed/12408854

Diamantaki, M., Coletta, S., Nasr, K., Zeraati, R., Laturnus, S., Berens, P., … Burgalossi, A.

(2018). Manipulating Hippocampal Place Cell Activity by Single-Cell Stimulation in Freely

Moving Mice. *Cell Reports*, *23*(1), 32–38. https://doi.org/10.1016/j.celrep.2018.03.031

Diba, K., & Buzsáki, G. (2007). Forward and reverse hippocampal place-cell sequences during

ripples. *Nature Neuroscience*, *10*(10), 1241–1242. https://doi.org/10.1038/nn1961

Driscoll, L. N., Pettit, N. L., Minderer, M., Chettih, S. N., & Harvey, C. D. (2017). Dynamic

Reorganization of Neuronal Activity Patterns in Parietal Cortex. *Cell*, *170*(5), 986-999.e16.

https://doi.org/10.1016/j.cell.2017.07.021

Dupret, D., O'Neill, J., Pleydell-Bouverie, B., & Csicsvari, J. (2010). The reorganization and

reactivation of hippocampal maps predict spatial memory performance. *Nature*

Neuroscience, *13*(8), 995–1002. https://doi.org/10.1038/nn.2599

Eichenbaum, H. B. (2004). Hippocampus: Cognitive processes and neural representations that

underlie declarative memory. *Neuron*, *44*(1), 109–120.

https://doi.org/10.1016/j.neuron.2004.08.028

Ekstrom, A. D., Kahana, M. J., Caplan, J. B., Fields, T. A., Isham, E. A., Newman, E. L., &

Fried, I. (2003). Cellular networks underlying human spatial navigation. *Nature*, *425*(6954),

184–188. https://doi.org/10.1038/nature01964

Ferbinteanu, J. D., & Shapiro, M. L. (2003). Prospective and retrospective memory coding in the

hippocampus. *Neuron*, *40*(6), 1227–1239. https://doi.org/10.1016/S0896-6273(03)00752-9

Frank, L. M., Brown, E. N., & Wilson, M. A. (2000). Trajectory encoding in the hippocampus

and entorhinal cortex. *Neuron*, *27*(1), 169–178. Retrieved from

http://www.ncbi.nlm.nih.gov/pubmed/10939340

Frankland, P. W., Bontempi, B., Talton, L. E., Kaczmarek, L., & Silva, A. J. (2004). The

involvement of the anterior cingulate cortex in remote contextual fear memory. *Science*,

304(5672), 881–883. https://doi.org/10.1126/science.1094804

Geva-Sagiv, M., Romani, S., Las, L., & Ulanovsky, N. (2016). Hippocampal global remapping

for different sensory modalities in flying bats. *Nature Neuroscience*, *19*(May), 1–11.

https://doi.org/10.1038/nn.4310

Gomperts, S. N., Kloosterman, F., & Wilson, M. A. (2015). VTA neurons coordinate with the

hippocampal reactivation of spatial experience. *ELife*, *4*(OCTOBER2015), 1–22.

https://doi.org/10.7554/eLife.05360.001

Goshen, I., Brodsky, M., Prakash, R., Wallace, J., Gradinaru, V., Ramakrishnan, C., &

Deisseroth, K. (2011). Dynamics of retrieval strategies for remote memories. *Cell*, *147*(3),

678–689. https://doi.org/10.1016/j.cell.2011.09.033

Grosmark, A. D., & Buzsáki, G. (2016). Diversity in neural firing dynamics supports both rigid

and learned hippocampal sequences. *Science*, *351*(6280), 1440–1443.

https://doi.org/10.1126/science.aad1935

Guitchounts, G., Markowitz, J. E., Liberti, W. A., & Gardner, T. J. (2013). A carbon-fiber

electrode array for long-term neural recording. *Journal of Neural Engineering*, *10*(4),

046016. https://doi.org/10.1088/1741-2560/10/4/046016

Hahnloser, R. H. R., Kozhevnikov, A. A., & Fee, M. S. (2002). An ultra-sparse code underliesthe

generation of neural sequences in a songbird. *Nature*, *419*(6902), 65–70.

https://doi.org/10.1038/nature00974

Hardt, O., Nader, K., & Nadel, L. (2013). Decay happens: The role of active forgetting in

memory. *Trends in Cognitive Sciences*, *17*(3), 111–120.

https://doi.org/10.1016/j.tics.2013.01.001

Howard, M. W., MacDonald, C. J., Tiganj, Z., Shankar, K. H., Du, Q., Hasselmo, M. E., &

Eichenbaum, H. (2014). A Unified Mathematical Framework for Coding Time, Space, and

Sequences in the Hippocampal Region. *Journal of Neuroscience*, *34*(13), 4692–4707.

https://doi.org/10.1523/jneurosci.5808-12.2014

Ito, H. T., Zhang, S. J., Witter, M. P., Moser, E. I., & Moser, M.-B. (2015). A prefrontal-

thalamo-hippocampal circuit for goal-directed spatial navigation. *Nature*, *522*(7554), 50–55.

https://doi.org/10.1038/nature14396

Kahn, K., Johnson, M., González-Martínez, J., Park, H.-J., Bulacio, J., Thompson, S., … Kerr,

M. S. D. (2017). The Role of Associative Cortices and Hippocampus during Movement

Perturbations. *Frontiers in Neural Circuits*, *11*(April), 1–11.

https://doi.org/10.3389/fncir.2017.00026

Katlowitz, K. A., Picardo, M. A., & Long, M. A. (2018). Stable Sequential Activity Underlying

the Maintenance of a Precisely Executed Skilled Behavior. *Neuron*, *98*(6), 1133-1140.e3.

https://doi.org/10.1016/j.neuron.2018.05.017

Kentros, C. G., Agnihotri, N. T., Streater, S., Hawkins, R. D., & Kandel, E. R. (2004). Increased

attention to spatial context increases both place field stability and spatial memory. *Neuron*,

42(2), 283–295. https://doi.org/10.1016/S0896-6273(04)00192-8

Kim, J. J., & Fanselow, M. S. (1992). Modality-specific retrograde amnesia of fear. *Science*,

256(5057), 675–677. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/1585183

Kinsky, N. R., Sullivan, D. W., Mau, W., Hasselmo, M. E., & Eichenbaum, H. B. (2018).

Hippocampal Place Fields Maintain a Coherent and Flexible Map across Long Timescales.

Current Biology, *28*(22), 1–11. https://doi.org/10.1016/J.CUB.2018.09.037

Kitamura, T., Ogawa, S. K., Roy, D. S., Okuyama, T., Morrissey, M. D., Smith, L. M., …

Tonegawa, S. (2017). Engrams and circuits crucial for systems consolidation of a memory.

Science, *78*(April), 73–78. https://doi.org/10.1126/science.aam6808

Kitamura, T., Saitoh, Y., Takashima, N., Murayama, A., Niibori, Y., Agayby, B., … Inokuchi,

K. (2009). Adult neurogenesis modulates the hippocampus-dependent period of associative

fear memory. *Cell*, *139*(4), 814–827. https://doi.org/10.1016/j.cell.2009.10.020

Kraus, B. J., Robinson II, R. J., White, J. A., Eichenbaum, H. B., & Hasselmo, M. E. (2013).

Hippocampal "Time Cells": Time versus Path Integration. *Neuron*, *78*(6), 1090–1101.

https://doi.org/10.1016/j.neuron.2013.04.015

Lee, D., Lin, B.-J., & Lee, A. K. (2012). Hippocampal Place Fields Emerge upon Single-Cell

Manipulation of Excitability During Behavior. *Science*, *337*(6096), 849–853.

https://doi.org/10.1126/science.1221489

Louie, K., & Wilson, M. A. (2001). Temporally structured replay of awake hippocampal

ensemble activity during rapid eye movement sleep. *Neuron*, *29*(1), 145–156.

https://doi.org/10.1016/S0896-6273(01)00186-6

Maboudi, K., Acsády, L., de Jong, L. W., Pfeiffer, B. E., Foster, D. J., Diba, K., & Kemere, C.

(2018). Uncovering temporal structure in hippocampal output patterns. *ELife*, *7*, e34467.

https://doi.org/10.7554/eLife.34467

MacDonald, C. J., Lepage, K. Q., Eden, U. T., & Eichenbaum, H. B. (2011). Hippocampal "time

cells" bridge the gap in memory for discontiguous events. *Neuron*, *71*(4), 737–749.

https://doi.org/10.1016/j.neuron.2011.07.012

Mankin, E. A., Sparks, F. T., Slayyeh, B., Sutherland, R. J., Leutgeb, S., & Leutgeb, J. K.

(2012). Neuronal code for extended time in the hippocampus. *Proceedings of the National*

Academy of Sciences, *109*(47), 19462–19467. https://doi.org/10.1073/pnas.1214107109/-

/DCSupplemental.www.pnas.org/cgi/doi/10.1073/pnas.1214107109

Manns, J. R., Howard, M. W., & Eichenbaum, H. B. (2007). Gradual Changes in Hippocampal

Activity Support Remembering the Order of Events. *Neuron*, *56*, 530–540.

https://doi.org/10.1016/j.neuron.2007.08.017

Margoliash, D., & Yu, A. C. (2009). Temporal Hierarchical Control of Singing in Birds. *Science*,

273(5283), 1871–1875.

Mau, W., Sullivan, D. W., Kinsky, N. R., Hasselmo, M. E., Howard, M. W., & Eichenbaum, H.

B. (2018). The Same Hippocampal CA1 Population Simultaneously Codes Temporal

Information over Multiple Timescales. *Current Biology*, *28*(10), 1499-1508.e4.

https://doi.org/10.1016/j.cub.2018.03.051

McKenzie, S., Robinson, N. T. M., Herrera, L., Churchill, J. C., & Eichenbaum, H. B. (2013).

Learning Causes Reorganization of Neuronal Firing Patterns to Represent Related

Experiences within a Hippocampal Schema. *The Journal of Neuroscience*, *33*(25), 10243–

10256. https://doi.org/10.1523/JNEUROSCI.0879-13.2013

Mehta, M. R., Quirk, M. C., & Wilson, M. A. (2000). Experience-dependent asymmetric shape

of hippocampal receptive fields. *Neuron*, *25*(3), 707–715. https://doi.org/10.1016/S0896-

6273(00)81072-7

Meira, T., Leroy, F., Buss, E. W., Oliva, A., Park, J., & Siegelbaum, S. A. (2018). A

hippocampal circuit linking dorsal CA2 to ventral CA1 critical for social memory

dynamics. *Nature Communications*, (2018), 1–14. https://doi.org/10.1038/s41467-018-06501-w

Michon, F., Sun, J. J., Kim, C. Y., Ciliberti, D., & Kloosterman, F. (2019). Post-learning Hippocampal Replay Selectively Reinforces Spatial Memory for Highly Rewarded Locations. *Current Biology*, *29*(9), 1436-1444.e5. https://doi.org/10.1016/j.cub.2019.03.048

Miller, J. F., Neufang, M., Solway, A., Brandt, A., Trippel, M., Mader, I., … Schulze-Bonhage, A. (2013). Neural Activity in Human Spatial Context of Retrieved Memories. *Science*, *342*(1111–1114).

Milner, B., Corkin, S., & Teuber, H. L. (1968). Further analysis of the hippocampal amnesic syndrome: 14-year follow-up study of H.M. *Neuropsychologia*, *6*(3), 215–234. https://doi.org/10.1016/0028-3932(68)90021-3

Monaco, J. D., Rao, G., Roth, E. D., & Knierim, J. J. (2014). Attentive scanning behavior drives one-trial potentiation of hippocampal place fields. *Nature Neuroscience*, *17*(5), 725–731. https://doi.org/10.1038/nn.3687

Morris, R. G. M., Garrud, P., Rawlins, J. N., & O'Keefe, J. (1982). Place navigation impaired in rats with hippocampal lesions. *Nature*, *297*(5868), 681–683. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/7088155

Muller, R. U., & Kubie, J. L. (1987). The effects of changes in the environment on the spatial firing of hippocampal complex-spike cells. *The Journal of Neuroscience*, *7*(7), 1951–1968. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/3612226

Muller, R. U., Kubie, J. L., & Ranck, J. B. (1987). Spatial firing patterns of hippocampal complex-spike cells in a fixed environment. *The Journal of Neuroscience*, *7*(7), 1935–1950. https://doi.org/10.1016/S0301-0082(96)00019-6

Muzzio, I. A., Levita, L., Kulkarni, J., Monaco, J., Kentros, C. G., Stead, M., … Kandel, E. R.

(2009). Attention enhances the retrieval and stability of visuospatial and olfactory

representations in the dorsal hippocampus. *PLoS Biology*, *7*(6).

https://doi.org/10.1371/journal.pbio.1000140

Niediek, J., & Bain, J. (2014). Human single-unit recordings reveal a link between place-cells

and episodic memory. *Frontiers in Systems Neuroscience*, *8*(6954), 184–187.

https://doi.org/10.3389/fnsys.2014.00158

O'Keefe, J. (1976). Place Units in the Hippocampus of the Freely Moving Rat. *Experimental*

Neurology, *51*(1), 78–109.

O'Keefe, J., & Dostrovsky, J. O. (1971). The hippocampus as a spatial map . Preliminary

evidence from unit activity in the freely-moving rat. *Brain Research*, *34*(1), 171–175.

Packard, M. G., & McGaugh, J. L. (1996). Inactivation of hippocampus or caudate nucleus with

lidocaine differentially affects expression of place and response learning. *Neurobiology of*

Learning and Memory, *65*(1), 65–72. https://doi.org/10.1006/nlme.1996.0007

Pastalkova, E., Itskov, V., Amarasingham, A., & Buzsáki, G. (2008). Internally Generated Cell

Assembly Sequences in the Rat Hippocampus. *Science*, *321*(September), 1322–1328.

Pfeiffer, B. E., & Foster, D. J. (2013). Hippocampal place-cell sequences depict future paths to

remembered goals. *Nature*, *497*(7447), 74–79. https://doi.org/10.1038/nature12112

Pfeiffer, T., Poll, S., Bancelin, S., Angibaud, J., Inavalli, V. K., Keppler, K., … Nägerl, U. V.

(2018). Chronic 2P-STED imaging reveals high turnover of dendritic spines in the

hippocampus in vivo. *ELife*, *7*, 1–17. https://doi.org/10.7554/elife.34700

Resendez, S. L., Jennings, J. H., Ung, R. L., Namboodiri, V. M. K., Zhou, Z. C., Otis, J. M., …

Stuber, G. D. (2016). Visualization of cortical, subcortical and deep brain neural circuit

dynamics during naturalistic mammalian behavior with head-mounted microscopes and

chronically implanted lenses. *Nature Protocols*, *11*(3), 566–597.

https://doi.org/10.1038/nprot.2016.021

Richards, B. A., & Frankland, P. W. (2017). The Persistence and Transience of Memory.

Neuron, *94*(6), 1071–1084. https://doi.org/10.1016/j.neuron.2017.04.037

Robinson, N. T. M., Priestley, J. B., Rueckemann, J. W., Garcia, A. D., Smeglin, V. A., Marino,

F. A., & Eichenbaum, H. B. (2017). Medial Entorhinal Cortex Selectively Supports

Temporal Coding by Hippocampal Neurons. *Neuron*, *94*(3), 677-688.e6.

https://doi.org/10.1016/j.neuron.2017.04.003

Rubin, A., Geva, N., Sheintuch, L., & Ziv, Y. (2015). Hippocampal ensemble dynamics

timestamp events in long-term memory. *ELife*, *4*(December), 1–16.

https://doi.org/10.7554/eLife.12247

Salz, D. M., Tiganj, Z., Khasnabish, S., Kohley, A., Sheehan, D., Howard, M. W., &

Eichenbaum, H. B. (2016). Time Cells in Hippocampal Area CA3. *The Journal of*

Neuroscience, *36*(28), 7476–7484. https://doi.org/10.1523/JNEUROSCI.0087-16.2016

Sheffield, M. E. J., & Dombeck, D. A. (2014). Calcium transient prevalence across the dendritic

arbour predicts place field properties. *Nature*, *517*(7533), 200–204.

https://doi.org/10.1038/nature13871

Sheffield, M. E. J., & Dombeck, D. A. (2019). Dendritic mechanisms of hippocampal place field

formation. *Current Opinion in Neurobiology*, *54*, 1–11.

https://doi.org/10.1016/j.conb.2018.07.004

Smith, D. M., & Mizumori, S. J. Y. (2006a). Hippocampal place cells, context, and episodic

memory. *Hippocampus*, *16*(9), 716–729. https://doi.org/10.1002/hipo.20208

Smith, D. M., & Mizumori, S. J. Y. (2006b). Learning-Related Development of Context-Specific

Neuronal Responses to Places and Events: The Hippocampal Role in Context Processing.

The Journal of Neuroscience, *26*(12), 3154–3163.

https://doi.org/10.1523/JNEUROSCI.3234-05.2006

Sparks, F. T., Lehmann, H., Hernandez, K., & Sutherland, R. J. (2011). Suppression of

neurotoxic lesion-induced seizure activity: Evidence for a permanent role for the

hippocampus in contextual memory. *PLoS ONE*, *6*(11).

https://doi.org/10.1371/journal.pone.0027426

Sutherland, R. J., O'Brien, J., & Lehmann, H. (2008). Absence of systems consolidation of fear

memories after dorsal, ventral, or complete hippocampal damage. *Hippocampus*, *18*(7),

710–718. https://doi.org/10.1002/hipo.20431

Takeuchi, T., Duszkiewicz, A. J., Sonneborn, A., Spooner, P. A., Yamasaki, M., Watanabe, M.,

… Morris, R. G. M. (2016). Locus coeruleus and dopaminergic consolidation of everyday

memory. *Nature*, 1–18. https://doi.org/10.1038/nature19325

van de Ven, G. M., Trouche, S., McNamara, C. G., Allen, K., & Dupret, D. (2016). Hippocampal

Offline Reactivation Consolidates Recently Formed Cell Assembly Patterns during Sharp

Wave-Ripples. *Neuron*, 1–7. https://doi.org/10.1016/j.neuron.2016.10.020

Vorhees, C. V., & Williams, M. T. (2014). Assessing spatial learning and memory in rodents.

ILAR Journal, *55*(2), 310–332. https://doi.org/10.1093/ilar/ilu013

Wang, S.-H., Teixeira, C. M., Wheeler, A. L., & Frankland, P. W. (2009). The precision of

remote context memories does not require the hippocampus. *Nature Neuroscience*, *12*, 253–

255. https://doi.org/10.1038/nn.2263

Wiltgen, B. J., Zhou, M., Cai, Y., Balaji, J., Karlsson, M. G., Parivash, S. N., … Silva, A. J.

(2010). The hippocampus plays a selective role in the retrieval of detailed contextual

memories. *Current Biology : CB*, *20*(15), 1336–1344.

https://doi.org/10.1016/j.cub.2010.06.068

Winocur, G., Frankland, P. W., Sekeres, M., Fogel, S., & Moscovitch, M. (2009). Changes in

context-specificity during memory reconsolidation: selective effects of hippocampal

lesions. *Learning & Memory (Cold Spring Harbor, N.Y.)*, *16*(11), 722–729.

https://doi.org/10.1101/lm.1447209

Wise, S. P., & Murray, E. A. (1999). Role of the hippocampal system in conditional motor

learning: Mapping antecedents to action. *Hippocampus*, *9*(2), 101–117.

https://doi.org/10.1002/(SICI)1098-1063(1999)9:2<101::AID-HIPO3>3.0.CO;2-L

Wood, E. R., Dudchenko, P. A., & Eichenbaum, H. B. (1999). The global record of memory in

hippocampal neuronal activity. *Nature*, *397*(6720), 613–616. https://doi.org/10.1038/17605

Wood, E. R., Dudchenko, P. A., Robitsek, R. J., & Eichenbaum, H. B. (2000). Hippocampal

neurons encode information about different types of memory episodes occurring in the same

location. *Neuron*, *27*(3), 623–633. Retrieved from

http://www.ncbi.nlm.nih.gov/pubmed/11055443

Zaremba, J. D., Diamantopoulou, A., Danielson, N. B., Grosmark, A. D., Kaifosh, P. W.,

Bowler, J. C., … Losonczy, A. (2017). Impaired hippocampal place cell dynamics in a

mouse model of the 22q11.2 deletion. *Nature Neuroscience*, *20*(11), 1612–1623.

https://doi.org/10.1038/nn.4634

Ziv, Y., Burns, L. D., Cocker, E. D., Hamel, E. O., Ghosh, K. K., Kitch, L. J., … Schnitzer, M. J.

(2013). Long-term dynamics of CA1 hippocampal place codes. *Nature Neuroscience*, *16*(3),

264–266. https://doi.org/10.1038/nn.3329

SUPPLEMENTAL FIGURES

Supplemental Figure 1: Neuron Extraction and Across-Session Registration (Related to Figure 1)

    A) Example neuron registration between two sessions. Blue/Green = pixels corresponding to putative ROIs extracted in the 1st/2nd session only. Yellow = pixels corresponding to portions of ROIs active in both sessions. Red = outline of ROIs matched as the same neuron between sessions.

    B) The small size of changes in ROI orientation between sessions indicate proper neuron registration between sessions.

Supplemental Figure 2: Backward Migration of Spatial Firing Across but not Between Sessions

(Related to Figure 2)

A) The centroid of spatial firing on the stem relative to its mean location across the entire session
drifts backwards throughout the session. Circles = centroid shifts for each neuron active on the
stem. Example session from one mouse for right turns only. *r = -0.28, p=5.1e-5, t = -4.1 for null
hypothesis that slope = 0.
B) Average change in centroid location between adjacent sessions for all mice between two sessions
indicates that place field location does not drift between sessions. p = 0.67 t-test.

Supplemental Figure 3: The Quality of Local Trajectory-Dependent Information Correlates with

Performance

A) *Top:* Performance for each session versus the peak discriminability value for all cells from that
session. Circles = all sessions, all mice. *Bottom:* Same as *Top* but for each mouse, triangles =
average for each mouse. *ρ=0.35, p=0.0056, ** ρ=0.99, p=0.0066 Pearson correlation.

B) Same as A, but for the peak reliability value. *ρ=-0.27, p=0.031 Pearson correlation.

Supplemental Figure 4: Decreased Turnover Rates for Non Event-Rate Matched Splitters versus

Place Cells (Related to Figure 4)

A) Histogram from one session showing that Ca^{2+} event rates are higher for splitters for than for arm
PCs. Blue/red dashed: mean number Ca^{2+} events for splitter/arm PCs.
B) Probability splitters and PCs stay active one day later for all mice. *p=3.2x10^{-6}, one-sided signed-
rank test.
C) Probability splitters and PCs stay active seven days later for all mice. *p=3.7x10^{-6}, one-sided
signed-rank test.
D) Difference between the probability that splitters stay active versus the probability stem PC stay
active. Dots: probability differences for individual session-pairs. Black solid/dashed lines: Mean
and 25%/75% quantiles of data at each time point. See Table 1 for one-sided signed-rank test p-
values at all lags.
E) Probability splitters and arm place cells stay active versus lag between sessions. Dots:
probabilities from individual session-pairs, lines: mean probability at each time lag. Green/red:
splitters/arm PCs.
F) Same as D for splitters vs. arm PCs. Red bars = significant differences after Holm-Bonferroni
correction of one-sided sign test, α = 0.05.

Table 3: One-sided Signed-Rank Significance Values for Probability Splitter vs. Return Arm Place Cells (APCs) are Present.

Lag (days)	1	2	3	4	5	6	7
vs. Arm PCs	7.0e-7	7.3e-5	0.003	0.018	0.013	0.005	1.4e-5
Lag (days)	8	9	10	11	12	13	14
vs. Arm PCs	7.8e-5	0.05	0.11	0.22	0.01	0.13	0.17
Lag (days)	15						
vs. Arm PCs	9.8e-4						

Supplemental Figure 5: Spatial Consistency of Splitters vs. Stem Place Cells (PCs) (Related to Figure 5)

A) Mean spatial correlations for splitter neurons versus stem PCs and return arm PCs for all sessions seven days apart from all mice. $*p=8.6 \times 10^{-6}$, $**p=4.5 \times 10^{-6}$, $**p=6.5 \times 10^{-5}$ one-sided signed-rank test.

B) Mean spatial correlations for splitter neurons and stem PCs versus lag between sessions for all mice/sessions. Red = Arm PCs, green = splitters, red bars = significant differences after Holm-Bonferroni correction of one-sided sign-test, $\alpha = 0.05$. See Table 2 for p-values at all lags.

Supplemental Figure 6: Tracking the Onset of Trajectory-Dependent Activity along the Entire

Stem

A) Mean reliability along the stem +/- 1 days from splitter onset for one representative mouse. p = 0.008 Kruskal-Wallis ANOVA, *p < 0.03 post-hoc Tukey test. Circles = mean reliability score for each neuron.

B) Mean reliability +/- 10 days from splitter onset for all mice. p = 0.0024 Kruskal-Wallis ANOVA, *p < 0.02 post-hoc Tukey test. Circles = mean of mean reliability of all neurons active on the stem for each session.

C) Mean discriminability along the stem +/- 1 days from splitter onset for one representative mouse. p = 1.7×10^{-6} Kruskal-Wallis ANOVA, *p < 0.0004 post-hoc Tukey test. Circles = mean discriminability score for each neuron.

D) Mean discriminability +/- 10 days from splitter onset for all mice. p = 0.0014 Kruskal-Wallis ANOVA, *p < 0.006 post-hoc Tukey test. Circles = mean of mean discriminability of all neurons active on the stem for each session.

METHODS

Animals

Five male C57/BL6 mice (Jackson Laboratories), age 3-14 months and weighing 25-30g were used. One mouse was excluded from analysis after performing the experiment due to the inability to correct motion artifacts in his imaging videos. Mice were housed socially with 1-3 other mice in a vivarium on a 12hr light/dark cycle with lights on at 7am and given free access to food and water. All mice were singly housed after surgery. All procedures were performed in compliance with the guidelines of the Boston University Animal Care and Use Committee.

Viral Constructs

We used an AAV9.*Syn*.GCaMP6f.WPRE.SV40 virus from the University of Pennsylvania Vector Core at an initial titer of ~4×10^{13} GC/mL and diluted it to ~$5\text{-}6 \times 10^{12}$ GC/mL with sterilized 0.05 phosphate buffered saline (KPBS) prior to infusion into CA1.

Stereotactic Surgeries

All surgeries were performed in accordance with previously published procedures (Kinsky et al., 2018; Resendez et al., 2016) in accordance with the Boston University Animal Care and Use Committee. Briefly, we performed two stereotactic surgeries and one base-plate implant on naïve mice, aged 3-8 months. Surgeries were performed under 1-2% isoflurane mixed with oxygen. Mice were given 0.05mL/kg buprenorphine (Buprenex) for analgesia, 5.0mL/kg of the anti-inflammatory drug Rimadyl (Pfizer), and 400mL/kg of the antibiotic Cefazolin (Pfizer) immediately after induction. They received the same dosage of Buprenex, Cefazolin, and Rimadyl twice daily for three days following surgery and were carefully monitored to ensure they never dropped below 80% of their pre-operative weight during convalescence. In the first

surgery, a small craniotomy was performed at AP -2.0, ML +1.5 (right) and 250nL of GCaMP6f

virus was injected 1.5mm below the brain surface at 40nL/min. The needle remained in place a

minimum of 10 minutes after the infusion finished at which point it was slowly removed, the

mouse's scalp was sutured, and the mouse was removed from anesthesia and allowed to recover.

3-4 weeks after viral infusion, mice received a second surgery to attach a gradient index

(GRIN) lens (GRINtech, 1mm x 4mm). After performing an ~2mm craniotomy around the

implant area, we carefully aspirated cortex using a blunted 25ga and 27ga needle under constant

irrigation with cold, sterile saline until we visually identified the medial-lateral striations of the

corpus callosum. We carefully removed these striations using a blunted 31ga needle while

leaving the underlying anterior-posterior striations intact, after which we applied gelfoam to stop

any bleeding. We then lowered the GRIN lens until it touched the brain surface and then

proceeded to lower it another 50µm to counteract brain swelling during surgery (note that in two

mice we first implanted a sleeve cannula with a round glass window on the bottom without

depressing an additional 50µm and then cemented in the GRIN lens during base plate

attachment). We then applied Kwik-Sil (World Precision Instruments) to provide a seal between

skull and GRIN lens and then cemented the GRIN lens in place with Metabond (Parkell),

covered it in a layer of Kwik-Cast (World Precision Instruments), and then removed the animal

from anesthesia and allowed him to recover after removing any sharp edges remaining from

dried Metabond and providing any necessary sutures.

Finally, after ~2 weeks we performed a procedure in which the mouse was put under

anesthesia but no tissue was cut in order to attach a base plate for easy future attachment of the

microscope. To do so, we attached the base plate to the camera via a set screw, carefully lowered

the camera objective and aligned it to the GRIN lens by eye, and visualized fluorescence via

857 nVistaHD v2.0/v3.0 until we observed clear vasculature and putative cell bodies expressing

858 GCaMP6f (Resendez et al., 2016), then raised the camera up ~50µm before applying Flow-It

859 ALC Flowable Composite (Pentron) between the underside of the baseplate and the cured

860 Metabond on the mouse's skull. After light curing we applied opaque Metabond over the Flow-It

861 ALC epoxy to the sides of the baseplate to provide additional strength and to block ambient light

862 infiltration.

863 *Imagine Acquisition and Processing*

864  Brain imaging data was obtained using nVista HD (Inscopix) v2/v3 at 1440 x 1280 pixels

865 and a 20 Hz sample rate. Two mice were lightly anesthetized (~60 seconds) to facilitate camera

866 attachment and then given ~15 minutes to recover prior to any recordings; the camera was

867 attached to the other two mice while they were awake. Prior to neuron/calcium event

868 identification we first pre-processed each movie using Mosaic (Inscopix) software which

869 entailed a) spatially downsampling by a factor of 2 (1.18 µm/pixel), b) performing motion

870 corrections, and c) cropping the motion-corrected movie to eliminate any dead pixels or areas

871 with no calcium activity. We then extracted a minimum projection of the pre-processed movie

872 for later neuron registration. We replaced isolated dropped frames (maximum 2 consecutive

873 frames) with the previous good frame, and in the rare case where more than 2 frames dropped in

874 a row these frames were excluded from all analyses.

875 *Neuron and Calcium Event Identification*

876  We utilized custom-written, open-source MATLAB software (available at

877 https://github.com/SharpWave/Tenaspis) to identify putative neuron ROIs and their calcium

878 events in accordance with previously published results (Kinsky et al., 2018; Mau et al., 2018). A

neuron had to have at least four calcium events in order to be considered active on a given

session.

Across-Session Neuron Registration

We utilized custom-written, open-source MATLAB software (available at

https://github.com/nkinsky/ImageCamp) to perform neuron registration across sessions in

accordance with previously published results (see Supplemental Figure 1). We checked the

quality of neuron registration between each session-pair in two ways: 1) by plotting the

distribution of changes in ROI orientations between session and comparing it to chance,

calculated by shuffling neuron identity between session 1000 times, and 2) plotting ROIs of all

neurons between two sessions and looking for systematic shifts I neuron ROIs that could lead to

false negatives/positives in the registration. During the course of these checks, we noticed the

quality of registration between sessions dropped significantly approximately halfway through the

experiment for two mice. Thus, we excluded any registrations occurring between the first and

second halves of the experiment for these two mice. Furthermore, the second half of the

experiment was excluded for these two mice when calculating the absolute onset session of place

cells and splitter cells but was included when calculating the relative onset day for each cell type.

Several other session-pairs exhibiting poor registrations based on the criteria above were also

excluded, though these were rare.

Behavioral Tracking and Parsing

Behavioral data was recorded via an overhead camera with Cineplex v2/v3 software

(Plexon) at a 30Hz sample rate. Cineplex produced automated tracking of the animal's position

by comparing each frame to a baseline image without the animal in the arena. Imaging and

behavioral data were synchronized by TTL pulse at the beginning of the recording. Each video

was inspected by eye for errors in automated tracking and fixed manually via custom-written

MATLAB software. After fixing all erroneous data points, the animal's position was interpolated

to determine his location at each imaging movie time point.

Histology

Mice were killed and transcardially perfused with 10% KPBS followed by formalin.

Brains of perfused mice were then extracted and post-fixed in formalin for 2-4 more days after

which they were placed in a 30% sucrose solution in KPBS for 1-2 additional days. The brains

were then frozen and sliced on a cryostat (Leica CM 3050S) in 40 μm sections after which they

were mounted and coverslipped with Vectashield Hardset mounting medium with DAPI (Vector

Laboratories). We then imaged slides at 4x, 10x, and 20x on a Nikon Eclipse Ni-E

epifluorescence microscope to verify proper placement of the GRIN lens above the CA1

pyramidal cell layer.

Experimental Outline

After recovery from surgery, mice were food deprived to maintain no less than 85% of

their pre-surgery weight. Mice were subsequently exposed to a variety of arenas in order to

habituate them to navigating with the camera attached. Prior to training on the alternation task,

all mice were given 1-4 habituation sessions on the alternation maze. The maze floor (inner

dimension = 64 x 29 cm) and walls (height = 18cm) were constructed from 3/8 inch (0.95cm)

thick plywood and the barriers between arms were constructed from two 53cm long 1.5 x 5.5

inch (3.8 x 14 cm) pine framing studs. The finished maze consisted of a central stem and two

return arms, each 7.5cm wide with 5.7cm wide openings at each of the central stem through

which mice could exit or enter the return arms. Two food wells ~ 0.25 cm deep were created

toward the end of each return arm to hold chocolate sprinkles: they were centered 12.5 cm from

the end of the maze where mice exited the return arm/entered the center stem. Food was placed

in these wells through a small opening in the side of the maze. The arena was sealed with

urethane prior to exploration.

Three of the mice were first trained to loop on each side of the maze independently in 3-

10 minutes blocks by blocking off access to the other side with Plexiglas dividers in order to

familiarize mice with the general task demands, arena, and location of food reward (chocolate

sprinkles); the other mouse received one habituation session where he was allowed to freely

traverse the maze. Following habituation, mice were placed in the center stem and rewarded at

the well on the reward arm regardless of the first turn direction. On subsequent trials, mice were

only rewarded if they turned the opposite direction of the previous trial. Mice were allowed to

run freely and were only blocked when they a) attempted to reverse course on the central stem,

b) attempted to exit the return arm after they had committed to it, or c) attempted to run straight

across to the other arm without turning down the central stem after obtaining reward. A mouse

was considered committed to an arm after his tail entirely crossed from the edge of the central

arm into the stem. Mice generally ran ballistically down the center stem and were allowed to

pause once they entered the return arm and after they obtained reward. Food reward was only

delivered once the mouse had committed to a return arm in order to avoid providing an auditory

cue of reward location. Two mice were forced to alternate in a subset of sessions/trials. One

mouse encountered a lapse in performance mid-way through the experiment and began

perseverating on one turn direction in blocks: he was subsequently given a number of trials at the

beginning of each session where he was forced to turn each direction by blocking off one turn

direction with a Plexiglas divider, after which he was then allowed to freely choose turn

directions. The other mouse was initially forced to alternate at the end of his habituation looping

sessions. All forced trials were not considered during later data analysis. Mice received 1-2

sessions per day. Sessions were terminated each day after 30 minutes or when the mouse stopped

consistently running ballistically down the center arm, whichever came first. The experiment

lasted 27, 16, 29, and 36 days for the four mice involved.

Place Cell Identification

Place cell identification was performed as described in Kinsky et al.(2018).

Trajectory-Dependent/Splitter Cell Identification

Prior to performing any analysis, each mouse's trajectory data was aligned to that from

the first habituation session. This was done by 1) manually rotating the data to correct for any

day-to-day changes in maze angle relative to the recording camera, 2) calculating the edges of

the mouse's trajectory as the data points located at the 2.5% and 97.5% points in the cumulative

density function of his x/y position data, and 3) adjusting the data by applying the necessary

translation and scaling (minimal) to overlay each session's trajectory on the first session. After

aligning data across sessions, the mouse's trajectory on each trajectory was parsed into his

progression through the different sections of the maze, starting at the a) **base**, then moving down

the b) **center stem** into the c) **choice** point, then turning into the d) left/right **entry** to the e)

return arm, and finally entered the f) **approach** to the center stem just after the reward port.

The center stem portion was manually identified for each mouse as the point where the mouse's

trajectory into/out of each return arm stopped diverging. This was done in order to mitigate the

possibility that trajectory-depending activity was controlled entirely by stereotyped sensory

inputs, e.g. the mouse hugging/whisking the left side of the center stem after right turn trial.

After parsing the animal's behavior into these sections, the center stem was broken up

into ~1cm bins and the event rate for each neuron was calculated for each trial. Tuning curves for

each trial type (left or right turn) were then constructed, which consisted of each neuron's mean

event rate for all correct trials at each spatial bin. The difference between these curves was then

calculated. To assess significance, we again constructed tuning curves for left/right trials and

calculated their difference, but after randomly shuffling trial turn identity 1000 times to establish

the likelihood the observed difference between tuning curves could emerge by chance. We then

defined splitters/trajectory-depending cells as neurons which had at least three bins whose real

tuning curve difference exceeded 950 of shuffled values. In order to exclude spurious

identification of splitters we only included neurons that produced a calcium event on the stem of

the maze on at least 5 trials.

We calculated several different metrics to quantify the level of trajectory-dependent

activity in each neuron. First, we calculated discriminability by summing the absolute value of

the difference between tuning curves along all stem bins and then dividing by the sum of tuning

curves along all stem bins. Second, we calculated reliability in the following manner: a) we

shuffled trial identity 1000 times and calculated the difference between shuffled tuning curves,

then b) calculated the proportion of shuffles in which the real difference between tuning curves

exceeded that of shuffled, then c) calculated reliability as the mean of this proportion along all

the stem bins. Note that splitter neurons by definition must have at least three bins with a

reliability value above 0.95 (see above). Last, we calculated the correlation between left and

right tuning curves. Note that this metric is very conservative since it produces low correlations

for splitters who shift the location of their peak activity between left and right trials along the

length of the stem (Figure 2B) but not for splitters who modulate their event rate in the same

place along the stem (Figure 2A).

In order to check the robustness of our results and control for any trajectory-dependent

information resulting from stereotyped deviations in speed or lateral position along the stem, we

also performed an additional analysis in line with previous studies (Ito et al., 2015; Wood et al.,

2000). To do so, we first divided the stem into five bins and calculated the average transient

probability in each bin for all trials. We then performed an ANOVA analysis using the anovan

function in MATLAB for each trajectory-dependent splitter neuron we detected. We used trial

type (left/right), stem bin, stem bin x trial type, speed, and the animal's lateral position as

covariates and mean transient probability as our dependent variable. Finally, we considered any

neuron to be a trajectory-dependent splitter neuron if had had a significant effect of trial type or

trial type x stem bin after accounting for speed and lateral position.

Linear Discriminant Decoding Analysis

A linear discriminant decoder was trained on data from 50% of trials on a given session

using the fitdiscr function in MATLAB. Calcium event activity for each neuron at each time

point when the mouse was on the center stem were used as the input variables and the mouse's

upcoming turn direction was used as the response variable. Only correct trials were considered

for training. The decoder was then used to predict the turn direction of the other 50% of trials,

after which the process was repeated 999 times using a different random 50% of trials for

training/decoding. The decoding accuracy was then calculated in ~3.3cm bins along the stem,

and the mean accuracy across all bins was taken as the decoding accuracy for that session.

Functional Phenotype Designation and Analysis of Neurons that Remain Active between

Sessions

We first performed neuron registration between sessions and classified neurons as staying

active if they were identified by our cell extraction algorithm on both sessions and produced at

least five calcium events (while the mouse was running) through the course of the first recording

session being considered in the registration. We then categorized cells into three different

functional phenotypes, 1) context-dependent splitter cells, 2) arm place cells, and 3) stem place

cells. Splitter cells were designated based on the criteria listed above. Neurons that produced no

calcium activity on the stem of the maze and met our place cell criteria were defined as return

arm place cells. Neurons that produced calcium activity on the stem and met our place cell

criteria but not our splitter neuron criteria were designated as stem place cells. In order to ensure

sufficient precision in calculating the probability a functional phenotype stayed active, a session-

pair was excluded from analysis if there were fewer than ten cells in either category in the first

session being registered. This analysis was performed in two ways: 1) including all cells found

for each phenotype, and 2) matching mean event rate between neuron phenotype by excluding

the lowest firing rate place cells. In the rare event that place cells had a higher mean firing rate

than splitter cells, no place cells were excluded.

Phenotype Ontogeny Analysis

We tracked the ontogeny splitter cells in three steps. First, we registered all the neurons we

recorded across the entire experiment. Second, we identified the first day/session that a neuron

passed our statistical criteria to be considered a splitter and defined that session as its onset.

Finally, we calculated multiple metrics for the quantity of trajectory-dependent activity produced

by each of these neurons (see 0 above) in all the sessions preceding and following onset,

excluding any sessions that occurred on the same day. The methodology for tracking place cell

onset was identical, except mutual information was used as a metric of spatial information

provided by each cell.

1038

AUTHOR CONTRIBUTIONS

Conceptualization: NRK. Methodology: NRK. Software: NRK, WM, DJS, SJL. Validation:

NRK. Formal Analysis: NRK, WM. Investigation: NRK, WM. Resources: NRK, MEH. Data

Curation: NRK, WM, EAR. Writing – original draft preparation: NRK. Writing – review and

editing: NRK, WM, DJS, EAR, SJL, MEH. Visualization: NRK, WM. Supervision: MEH, NRK.

Project administration: NRK, MEH. Funding Acquisition: MEH.

1045

ACKNOWLEDGEMENTS

First and foremost we would like to thank Dr. Howard Eichenbaum for his help in all stages of

this project. We would like to also thank Dr. Jon Rueckemann for input on data analysis and

interpretation and Dr. Nick Robinson for feedback during manuscript preparation. We would like

to thank Dr. Steve Gomperts, Dr.Annabelle Singer, Dr.Amar Sahay, Dr.Kamran Diba and the

members of their lab, as well as Dr.Ian Davison, Dr.Steve Ramirez, Dr.Ehren Newman, and Dr.

Mark Kramer, as well as Dr. Andrew Alexander, Dr. Jake Hinman, Dr. John Bladon, Dr. Ryan

Place, Dan Sheehan, Winny Ning, Dan Orlin, Jiawen Chen, and Catherine Mikkelsen for

feedback and critiques during early stages of data analysis and writing. We would like to thank

Annalyse Kohley, Hanish Polavarapu, and Scott Bovino for help with animal training and

behavioral tracking quality control. We would like to thank Denise Parisi, Shelley Russek, and

Sandra Grasso for administrative support. We would like to acknowledge the GENIE Program,

specifically Vivek Jayaraman, Ph.D., Douglas S. Kim, Ph.D., Loren L. Looger, Ph.D., Karel

Svoboda, Ph.D. from the GENIE Project, Janelia Research Campus, Howard Hughes Medical

Institute, for providing the GCaMP6f virus. Finally, we would like to acknowledge Inscopix, Inc.

for making single-photon calcium imaging miniscopes widely available, and specifically Lara

Cardy and Vardhan Dani for all their technical support throughout and after the experiment.

1063

1064   **GRANT SUPPORT**

1065   This work was supported by ONR MURI N00014-16-1-2832, NIH R01 MH060013,

1066   NIH RO1 MH052090, NIH RO1 MH051570, and NSF NRT UtB: Neurophotonics DGE-

1067   1633516.

1068

1069   The authors declare no competing interests.