How does individual variation in sociality influence fitness

in prairie voles?

Anne C. Sabol[a,*], Connor T. Lambert[b], Brian Keane[b], Nancy G. Solomon[b], Ben Dantzer[a,c]

[a]Department of Psychology, University of Michigan, Ann Arbor, MI, USA

[b]Department of Biology, Miami University, Oxford, Ohio, USA

[c]Department of Ecology and Evolutionary Biology, University of Michigan, Ann Arbor, MI, USA

* Correspondence:

Anne Sabol

11200 SW 8[th] St.

Miami, FL 33199

acsabol72@gmail.com

(937) 243-8767

Abstract

Comparative studies aid in our understanding of specific conditions favoring the initial evolution

of different types of social behaviors, yet there is much unexplained intraspecific variation in the

expression of social behavior that comparative studies have not yet addressed. The proximate

causes of this individual variation in social behavior within a species have been examined in

some species but its fitness consequences have been less frequently investigated. In this study,

we quantified the fitness consequences of variation in the sociality of prairie voles (*Microtus*

ochrogaster). We characterized sociality of voles in semi-natural enclosures using an automated

behavioral tracking system paired with social network analyses to quantify the degree of spatial

and temporal co-occurrence of different voles. We then assessed the relationship between

sociality with mating success (number of different conspecifics with which an individual

produced offspring) and reproductive success (total number of offspring surviving to first

capture). We measured the number of social connections each individual had with all voles and

only with opposite-sex voles by calculating unweighted degree through social network analyses.

Both female and male voles varied in the number of social connections they had with all

conspecifics and with opposite-sex conspecifics. Voles with an intermediate number of social

connections with voles of both sexes had higher mating success overall. In our analyses that

considered all social connections with voles of both sexes, voles with an intermediate number of

social connections produced more offspring. Males with a very high or low number of social

connections also had the lowest average body mass. Overall, our results suggest some limit on

the fitness benefits of sociality. Although there was substantial individual-variation in our

measure of vole social behavior, intermediate levels of social connections may be most

favorable.

Keywords

Prairie vole, Social behavior, Social network analyses

Introduction

Sociality comes in many forms across a diversity of taxa, ranging from loose, temporary associations during breeding to long-term group associations whose members exhibit complex social behaviors such as cooperative hunting or alloparental care. Different forms of sociality are thought to provide a variety of fitness benefits, including shared resources, reduced predation, communication, or better access to mates (Eisenberg et al., 1972; Alexander, 1974; van Schaik 1983; Emlen, 1984; Krause and Ruxton, 2002, Silk, 2007). Sociality may also come with costs associated with group-living, including increased disease transmission (Ewald, 1994; Langwig et al., 2012; Kappeler et al., 2015), parasite load (Côté, and Poulinb, 1995; Whiteman and Parker, 2004), physiological stress (Creel et al., 2013), reproductive interference by other group members (Clutton-Brock et al., 1998; Lukas and Huchard, 2014), as well as the time and energy costs devoted to developing and maintaining social connections that could otherwise be devoted towards activities directly related to individual survival or reproduction such as foraging or parental care. Given that these fitness benefits and costs of sociality may vary in direction or magnitude due to ecological circumstances such as food abundance, weather patterns, or temporal variability in these factors (Emlen, 1994; Hatchwell and Komdeur, 2000; Shuster and Wade, 2003; Schradin and Pillay, 2005; Rubenstein and Lovette, 2007; Schradin et al., 2010; Shen et al., 2017), it is no surprise that there is considerable variation in the presence or expression of different types of social behaviors among species.

Comparative studies have been useful in revealing some of the factors favoring the evolution of different types of social behaviors. For example, phylogenetic comparative meta-analyses show that social monogamy is more commonly found in mammalian species that experience low population densities (Lukas and Clutton-Brock, 2013) and genetic monogamy is also more common in mammalian species with low population densities as well as those that exhibit parental care (Lambert et al., 2018). Other comparative studies have shown that cooperative breeding, where subordinate group members care for offspring produced by

dominant breeders, is more likely to be found in mammalian species that live in arid areas

(Faulkes et al., 1997; Lukas and Clutton-Brock, 2017) or in avian species that inhabit areas

where rainfall is low and unpredictable (Jetz and Rubenstein, 2011). While these studies help

illuminate some factors affecting the evolution of social behaviors like social monogamy or

cooperative breeding, they typically cannot address the causes of intraspecific variation in

sociality.

Sociality is not always expressed to the same level or in the same way within a species

and there are often a variety of social strategies within the same species (Lott, 1991; Clutton-

Brock, 1989; Getz et al., 1993). The magnitude of variation in the expression of social behavior

within a species may not be as large as that among species but it requires explanation

nonetheless. Although some studies have investigated some of the proximate causes of

individual variation in specific social behaviors such as social monogamy (Young and Wang,

2004; Donaldson and Young, 2008; Ophir et al., 2008; Okhovat et al., 2015; Walum and Young,

2018) or alloparental care (Dantzer et al., 2017), fewer studies document its fitness

consequences. This is not surprising as it requires detailed studies that document the social

behavior of individuals and then relate their degree of sociality to their survival and reproduction.

Most previous studies on this topic have been conducted in primates where investigators

documented how the strength of social relationships ("social bonds") impacted offspring

survival, mating success, or longevity (e.g., Silk et al., 2003, 2009, 2010; Schülke, et al., 2010).

By investigating the association between social behavior and fitness within a species, we may

be able to understand how individuals balance the tradeoffs between the costs and benefits of

sociality and therefore obtain an even greater understanding of the evolution of sociality.

We characterized the variability of social behavior in individual female and male prairie

voles (*Microtus ochrogaster*) and its association with their mating success, reproductive

success, and body condition. Prairie voles are an ideal study system to investigate sociality

because both sexes can exhibit natural variation in sociality by displaying different reproductive

strategies (Solomon and Jacquot, 2002). Both females and males can exhibit strategies ranging

from pair bonded "residents", with an established territory to non-territorial, unpaired

"wanderers" and both are known to switch their strategy over the course of their lifetime (Getz

and Hofmann, 1986; Getz et al., 1993; Solomon and Jacquot, 2002; McGuire and Getz, 2010;

Shuster et al., 2019). Further, genetic monogamy and social monogamy are distinct with some

female-male pairs exhibiting high levels of social and genetic monogamy, some pairs being

socially monogamous but not genetically monogamous, and some individuals exhibiting no

socially or genetically monogamous behavior or mating patterns (Solomon et al., 2004). Thus,

prairie voles may vary in the number and strength of social associations they have with other

individuals. For example, a genetically monogamous female and male likely have a very strong

social association with each other whereas a wandering individual may have many weak

connections with multiple opposite-sex voles. Additionally, voles may vary in the number of

social connections they have with other individuals than their partner because under certain

environmental conditions, such as high population density (Getz et al., 1993; Cochran and

Solomon 2000; Lucia et al. 2008), voles can also form extended family groups when offspring

delay dispersal from the natal nest.

We characterized the social behavior of individual prairie voles in semi-natural

enclosures using an automated monitoring system. Prairie voles were all marked uniquely with

passive integrated technology (PIT) tags and their movements were continuously monitored by

an array of radio-frequency identification (RFID) antennas. We used patterns of spatial and

temporal co-occurrence generated from this system with social network analyses to estimate

patterns of social association (unweighted degree). We have previously shown that opposite-

sex voles that exhibit close social associations, as generated by this automated method of data

collection, are also more likely to be caught in the same trap together, exhibit overlapping home

ranges, and show a strong social preference for one another in a choice experiment (Sabol et

al., 2018), suggesting that these measures reflect the strength of social associations. Therefore,

we classified individuals with a greater number of social network connections with opposite-sex

or same-sex voles as more social, although we note that the valence of these connections

(agonistic or affiliative) is not known. We characterized the sociality of all

voles throughout the breeding season using social network analyses, recorded their survival

through this period and used parentage analyses to quantify their mating success (number of

different individuals with which they produced offspring) and reproductive success (number of

offspring produced that survived to emergence from the natal nest).

We predicted that voles with a greater number of social network connections (i.e., more

social) would have higher mating and reproductive success but lower body condition due to the

trade-offs associated with high levels of this type of sociality. We only investigated body

condition in males because any changes in body mass in females is likely directly related to

pregnancy, We predicted that male voles that were more social (had more social network

connections) would have lower body condition (mass) because polygyny typically involves

energetic costs and risky travel associated with finding female mating partners and interacting

with male competitors, which may therefore reduce survival (Blanckenhorn et al 1995; Armitage

1998).

Methods

Study site and study animals

All fieldwork was conducted at Miami University's Ecology Research Center in Oxford,

Ohio from May to August 2017. Voles were released into two separate 0.1 ha enclosures (33 m

x 33 m). The 20-gauge sheet metal walls of the enclosures were 75 cm tall and embedded 45

cm into the ground to prevent vole movements among enclosures. Enclosure walls were topped

with an electrified wire to prevent other small to medium sized mammals (e.g. raccoons and

weasels) from entering the enclosures and disturbing traps. Although this system likely

prevented many mammalian predators from entering the enclosures, the enclosures were open

and accessible to avian predators and snakes. We had multiple animals that were confirmed to

be depredated by owls and also occasionally saw large snakes within the enclosures. Prior to

releasing prairie voles into the enclosures, we live-trapped within the enclosures for 3

continuous days to capture any small mammals (e.g. *Microtus pennsylvanicus*, *Peromyscus*

maniculatus, or *Blarina brevicauda*) and released them outside of the enclosures.

We released laboratory-bred 7th and 8th generation prairie voles (descended from voles

originally captured in Illinois) into two enclosures. The pedigree of the laboratory population was

known and to avoid inbreeding, we did not place opposite-sex siblings or parents and their

offspring into the same enclosure. All founding voles were sexually mature (> 31 d, Solomon

1993) but had never bred. Each enclosure was founded with a different density: Enclosure 1

was established by releasing 48 voles (24 females, 24 males) and Enclosure 2 initially

contained 24 voles (12 females, 12 males). These represented densities of 480 voles/ha and

240 voles/ha respectively, which were within the range of vole densities observed in natural

populations (10 to 600 voles/ha: Getz et al., 1994; Getz et al., 2001). The different starting

density was employed to assess the role of density on vole social behavior but, as shown below

(Fig. 1), vole density in Enclosure 1 (high density) decreased over the course of the field

season. Additionally, we do not find or report an effect of density on vole social behavior in this

dataset, only whether there is a difference between the enclosures in general for the whole

season. The vegetation within enclosures consisted primarily of perennial grasses and forbs,

which provided food and cover. Voles were not provided with supplemental food besides the

cracked corn, a low-quality food, used to bait the live-traps.

Recording social behavior of voles

Prior to release into the enclosures, all founding voles were implanted with a PIT tag

(Biomark: Boise, Idaho) with a unique identification number. Once released, we monitored our

populations through a proprietary system of 12 RFID antennas connected to a central recording

station (BioMark RM310/SM303, Boise, ID) that we have used previously to create social

networks of prairie voles and infer their social associations (Sabol et al., 2018). These antennas

were placed within the enclosures in two different 3 x 4 arrays (Figure A1) that were rotated

from array 1 to array 2 every 3 days and moved from one enclosure to the other every 6 days

during the 14-week study. The antennas recorded the individual PIT tag numbers of each vole

that passed within 15 cm of the antenna once a second for the entire time the animal was within

this radius of the antenna. When multiple tags were within the 15 cm radius, the system

alternated in recording the tag numbers so that both could be detected. This allowed us to

record the natural movements and social associations of individuals in both populations, which

we have previously shown to be comparable and more detailed than traditional methods of

recording social associations in these populations (Sabol et al., 2018).

Monitoring vole reproduction

We live-trapped both enclosures by placing two Ugglan live- traps (Granhab, Hillerstorp,

Sweden), baited with cracked corn, per grid stake on a 5 x 5 m trapping grid. These live-traps

had a metal cover over the top to shield voles from the elements while in the traps. At the

beginning of the experiment, we live-trapped nearly every day, setting traps in the evening

(2230-2300 h) and checking them approximately two hours later and then leaving them set to be

checked the next morning (0700 h). After the first two weeks of the experiment, we set traps

approximately 3 times per week per enclosure (Monday, Wednesday, and Friday) alternating

between the two enclosures so that voles in each enclosure were trapped three days over the

course of two weeks. Traps were set in the evening (2230-2300 h) and checked the following

morning at 0700 h. If we needed to put radiocollars on voles (see below), we also set traps from

1800-1900 h and then checked them the same evening from 2000-2100 h. Prairie vole gestation

and time to weaning are each approximately 21 days (Richmond & Conaway, 1969) and,

therefore, the offspring produced within the enclosures usually first emerge from the nest

approximately 5-6 weeks after the adults are released (N. Solomon, B. Keane, personal

observations). We therefore stopped setting traps overnight from week 6-14 of the experiment

to ensure that offspring would not spend an entire night without their parents. During this time,

we set traps in the evening (2230-2300 h) and checked them approximately two hours later but

did not leave traps set overnight. Finally, we did not trap when there were severe thunderstorms

and made up this trapping session on a different day during the week, if possible. We

supplemented this regular grid trapping by placing additional traps at nest entrances after we

located them using VHF telemetry and/or UV powder tracking (Lemen and Freemen, 1985).

During each capture, we identified each individual using a handheld PIT tag reader,

recorded the location where the vole was live-trapped, the other individuals caught in the same

trap, body mass (g, with Pesola spring scale), and assessed reproductive condition. Males were

recorded as being either scrotal or non-scrotal depending on the development of the testes.

Females were recorded as non-reproductive, pregnant, or lactating depending on whether

developing fetuses could be felt or if nipples were pronounced. Once offspring were live trapped

for the first time, we gave them a unique toe clip until they were large enough (>25 g) to be PIT

tagged. Reproductive success of adult voles was estimated as the total number of offspring they

produced that we were able to trap.

Parentage analyses

Before adult voles were released into the enclosures, we collected a small piece of ear

tissue and stored it in 70% ethanol in a -80^0 C freezer. When offspring were trapped for the first

time, the tissue from the identifying toe clip was saved and temporarily stored in a -20^0 C freezer

until samples could be moved to a -80^0 F freezer. We extracted DNA using DNeasy Blood and

Tissue kits (Qiagen). We followed the manufacturer's protocol except that tissue samples were

incubated at 70^0 C, DNA was eluted in 100 µl of molecular grade water instead of 200 µl of buffer

AE, and DNA samples were incubated at room temperature for 5 minutes instead of 1 minute.

Once DNA was extracted, we genotyped the samples for all adults and offspring at seven

microsatellite loci (Keane et al., 2007). We then ran an allele frequency analysis on the population

of voles in each enclosure separately using Cervus 3.0.7 (see Mabry et al., 2011 for full details).

To determine parentage, we used Cervus 3.0.7 parentage analyses with known sexes, which

calculates a likelihood ratio for each potential mother and father in order to determine the most

likely biological parents in the population for each offspring (Mabry et al., 2011). We were able to

determine both parents (trio confidence level) with a 95% confidence level for 33/41 (80.5%)

offspring, so only these 33 offspring were included in the analyses of mating and reproductive

success.

Ethical Note

All procedures involving live animals were in accordance with the guidelines provided by

the American Society of Mammalogists for the use of wild mammals in research (Sikes et al.,

2016) and were approved by Miami University's Institutional Animal Care and Use Committee

(protocol number 979) as this was where all work with live animals occurred.

Statistical Analyses

All statistical analyses were done in R version 3.4.1 (R Core Team, 2017). Figure 1 was

made in R while all other figures were made in ggplot2 version 2.2.1 (Wickham, 2009). All linear

models and generalized linear models were run in R. For all models listed below, we assessed

collinearity among the predictor variables using variance inflation factors (VIFS) in the package

car, version 3.0-0 (Fox and Weisberg, 2011).

Density and Body Mass

Population density was calculated based on the number of unique individuals caught

within each two-week period (over each two-week period both enclosures were trapped with

equal effort except for occasional cancellations due to weather). To investigate population

density over the course of the field season, we used a linear model with density (log-

transformed with base 10 to improve normality of residuals) with the fixed effects of enclosure

and weeks in the study and the interaction of these terms. Sex ratio was calculated by dividing

the number of adult males by the number of total adults for each two-week period. We used a

binomial generalized linear model to investigate sex ratio with the fixed effects of enclosure and

weeks in the study and the interaction of these terms. VIFs for all non-interaction terms were all

< 3.57.

Social Network Analyses

We measured the number of social connections (unweighted degree, hereafter degree)

between same-sex or opposite-sex voles based on co-occurrence data from the RFID

antennas. Individuals with a high degree would have had instances of spatial and temporal co-

occurrence with many other voles whereas those with a low degree had few. We conducted all

social network analyses using the R package asnipe version 1.1.4 (Farine, 2017b). In order to

generate our social networks, we took the PIT tag readings from the RFID antennas and ran

them through a Gaussian Mixture Model with each day labeled separately (Psorakis et al.,

2012). This model goes through the raw data of the PIT tag readings and creates groups based

on when tag readings at the same antenna are clumped throughout time. Therefore, there is not

a uniform time period used to create these groups, they are based on how our data were

distributed over time. This model uses clusters of tag readings as "centres of mass" where data

are concentrated and then determines the groups based on the amount and distribution in time

of tag readings in each cluster to determine where to split groups (Psorakis et al., 2012). The

duration of these group events ranged from 0 seconds (so voles were both at the antenna at the

same time) to 66,161 seconds with an average of 655.2 ± 3,352.8 seconds. This then creates a

group by individual matrix where being in the same spatial and temporal "group" counts as an

association between individuals. As we were only interested in the number of connections each

individual had (not the strength of these connections), we used a binary, unweighted

measurement of degree where any non-zero association was counted as a "1". Thus, anytime

we refer to the number of social connections in this paper, we calculated this using the

unweighted degree. For more details about the construction of the social networks see Sabol et

al. (2018).

Reproductive Success Models

For all models including the number of mates (mating success) or the number of

offspring produced that survived to emergence from the natal nest (reproductive success) as the

response variable, we used Poisson generalized linear models. For each response variable we

ran two models using social network data, one including all social connections in order to

investigate sociality overall and one including only the opposite-sex social connections. These

models all had fixed effects of the number of social connections, the interaction of this with sex,

the number of social connections squared to assess non-linear effects of social connections on

mating success or reproductive success (one model including all connections and another for

each response variable including only opposite-sex connections), the interaction of this with sex,

enclosure, and survival (calculated as the proportion of the field season the individual survived

based on last detection). To test if mating success and reproductive success were related, we

ran a separate model with the number of offspring produced as the response variable and fixed

effects of the number of mates with which individuals produced offspring, the interaction of this

with sex, and enclosure. None of the GLMs were over-dispersed as all the dispersion

parameters were <1, which we tested using R package AER version 1.2.5 (Kleiber and Zeil,

2008). VIFs were all < 3.5 except interaction and squared terms, which were predictably high.

Body Mass

To investigate body mass, we calculated the average body mass for each male vole for

the entire field season (range 1-19 measurements, average 7.75 measurements). Females

were not included because we were using body mass as a proxy for body quality, and female

mass would be affected by both pregnancy status and body condition. We then used a general

linear model for average body mass with the number of social connections (one model including

all connections another including only opposite-sex connections), the number of social

connections squared (one model including all connections another including only opposite-sex

connections), enclosure, survival (calculated as the proportion of the field season the individual

survived based on last detection), and the number of mass measurements we had for each

individual. We visually assessed the distribution of the data and residuals for normality. VIFs

were all < 2.2 except interaction or squared terms and survival. However, when survival was

excluded from the model, VIFs for all of the other terms were < 3.5 except interactions and

squared terms. Including survival did not alter the statistical significance of any of the results

shown below so we left it in.

Randomized Models

For every model that included unweighted degree (the number of social connections),

we used the network permutation method in asnipe (Farine, 2017b). This method is useful

because it helps control for the fact that social network data are not independent. This method

also allows us to investigate our hypotheses more specifically by allowing us to test if the

observed relationships are significantly different from random networks with the same structure

as our social networks (see Farine et al., 2015; Spiegel et al. 2017 for other similar uses of this

method). The network permutation method takes a piece of data from the group by individual

matrix and swaps it for a different individual (Farine, 2013). Specifically, we ran 10,000

randomized models where each time another piece of data from the individual by group matrix

was swapped. We also restricted swaps to only voles in the same enclosure that were recorded

on the RFID antennas during the same day to control for voles that did not survive the entire

season. Further, for the opposite-sex networks we restricted swaps to include only voles of the

same sex so that we were only comparing our opposite-sex network to other opposite-sex

networks, not all possible combinations. We then compared the regression coefficients from the

model for each variable that includes a social network statistic to corresponding b-values from

randomized networks and calculated a new *P*-value based on the number of randomized

models that produced a b-value with a higher absolute value than the absolute value of the

observed model. Therefore, our *P*-value shows us whether the relationship we have observed is

stronger than the relationship from 10,000 randomizations of our dataset (Farine, 2013). We ran

each set of randomizations three times to ensure that the *P*-values were consistently significant

in each of the randomizations. We present all three *P*-values from these randomizations and,

conservatively, only consider that a relationship is statistically significant if all three

randomizations revealed *P*-values <0.05.

Results

Enclosure density & adult sex ratio

The number of adult voles in each enclosure declined over the course of the field

season due to mortality (effect of weeks in the study, b = -0.14, SE = 0.012, t_{12} = -11.45, *P* <

0.0001, Fig. 1a, note estimates on $\log_{10}$ scale) and the significant interaction between weeks in

the study and enclosure indicated that vole density decreased more strongly in the high density

enclosure than in the low density enclosure (enclosure x weeks: b = 0.083, SE = 0.017, t_{12} =

4.79, *P* = 0.00045). For example, the starting density of voles in the high-density enclosure was

480 voles/ha (week 0 in Fig. 1a) but was reduced to 200 voles/ha in the middle of the

experiment (week 8 in Fig. 1a) and to 60 voles/ha at the end of the experiment (week 14 in Fig.

1a). The starting density in the low-density enclosure was 240 voles/ha (week 0 in Fig. 1a) but

was reduced to 180 voles/ha in the middle of the experiment (week 8 in Fig. 1a) and 100

voles/ha at the end of the experiment (week 14 in Fig. 1a). In total, only 12.5% of voles in the

high-density enclosure were still alive at the end of the experiment whereas 41.7% of the voles

in the low-density enclosure were still alive at the end of the experiment.

Unlike density, adult sex ratio did not differ significantly during the course of the

experiment (effect of time, b = -0.04, SE = 0.16, z = -0.26, df = 12, *P* = 0.80, Fig. 1b) or between

the two enclosures (effect of enclosure, b = 0.15, SE = 1.84, z = 0.082, df = 12, *P* = 0.94, Fig.

1b).

Effects of sociality on mating success

Overall, both female and male voles that had an intermediate (i.e., the middle of the

range of observed values) number of social connections produced offspring with a greater

number of different mates (i.e., had higher mating success). In the model considering all social

interactions with same- and opposite-sex individuals, voles that had an intermediate number of

social connections (degree) with all possible individuals had higher mating success (effect of

social connections^2: b = -0.012, z = -0.88, P-values from randomized networks = 0.012, 0.0093,

<0.0001, Table 1, Fig. 2a) but this relationship was slightly different between the sexes (sex x

social connections^2: b = 0.014, z = 0.94, P = 0.010, 0.0018, <0.0001, Table 1, Fig. 2a). In both

females and males, those with an intermediate number of social connections had the highest

mating success, therefore the interaction with sex and the number of social connections^2 on

mating success seemed to be largely due to males having slightly more overall social

connections than females while female mating success peaked at a lower number of social

connections (Fig. 2a). There is also a qualitative difference in the shape of the curve, with

female mating success peaking at a lower number of social connections but then dropping off

more steeply, while male mating success peaked at a higher number of social connections but

declined more gradually (Fig. 2a).

The same relationship was true when only opposite-sex connections were considered.

Mating success was highest for female and male voles with an intermediate number of

opposite-sex social connections (social connections^2: b = -0.028, z = -1.01, P = 0.0011, 0.0035,

0.001, Fig. 2b), although the magnitude of this effect slightly differed between the sexes (sex x

social connections^2: b = 0.057, z = 1.36, P = <0.0001, <0.0001, 0.0068, Table 1, Fig. 2b). This

latter difference between the sexes seems to be driven by the difference in the number of social

connections between the sexes with females tending to have slightly more social connections

than males.

Overall, both female and male voles in the low-density enclosure had higher mating

success than individuals in the higher density enclosure (from model for all social connections: b

= 1.89, z = 4.55, *P* < 0.0001; from model for all opposite-sex social connections: b = 2.03, z =

4.72, *P* < 0.0001, Table 1). Individuals that survived in the enclosures for longer had higher

mating success (all social connections: b = 2.07, z = 2.33, *P* = 0.020; opposite-sex social

connections: b = 1.75, z = 2.09, *P* = 0.037, Table 1).

Effects of sociality on reproductive success

Both female and male voles with an intermediate number of social connections with all

voles produced more offspring that survived to emergence from the natal nest (b = -0.0085, z =

-0.77, *P* = 0.002, 0.035, 0.015, Table 2, Fig. 3a). This result did not consistently differ by sex

(sex x social connections^2: b = 0.0082, z = 0.63, *P* = 0.021, 0.11, 0.022, Table 2, Fig. 3a).

However, when only considering opposite-sex social connections (Fig. 3b), these relationships

were not significant. The number of offspring that voles produced was not related to the number

of opposite-sex social connections for males or females (effect of opposite-sex social

connections: b = 0.15, z = 0.52, *P* = 0.20, 0.53, 0.54, Table 2, Fig. 3b) and voles with an

intermediate number of opposite-sex social connections did not produce significantly more

offspring (effect of opposite-sex social connections^2: b = -0.022, z = -0.93, *P* = 0.14, 0.099, 0.12,

Table 2, Fig. 3b). This relationship did not consistently vary with sex, as the difference between

the sexes for opposite-sex connections^2 where the inverted u-shaped relationship was slightly

lessened in males and was not significant in all three sets of randomizations (sex x opposite-sex

connections^2: b = 0.038, z-value = 0.92, *P* = 0.063, 0.010, 0.012, Table 2, Fig 3b) and the

difference between opposite sex connections between males and females where males tended

to have fewer connections than females was also not significant in all three sets of

randomizations (sex x number of opposite-sex connections: b = -0.39, z = -1.04, *P* = 0.047,

0.024, 0.11). Male and female voles that survived for longer produced more offspring (all social

connections: b = 2.49, z = 3.03, P = 0.0025; opposite-sex social connections: b = 2.19, z = 2.83,

P = 0.0046, Table 2).

 Effects of mating tactics on reproductive success

Overall, there was a fitness advantage to being polygynous (mating with a greater

number of individuals) in both females and males. Individuals with more mates produced more

offspring that survived to emergence from the natal nest (b = 0.64, z = 4.88, *P* <0.0001, Table 3,

Fig. 4) and this relationship did not differ by sex as the interaction between number of mates

and sex was not significant (b = -0.093, z = -0.56, *P* = 0.58, Table 3). Individuals in the lower

density enclosure had higher reproductive success than individuals in the higher density

enclosure (b = 1.53, z = 3.37, *P* = 0.00076, Table 3).

Effects of sociality on body mass

Males with an intermediate number of connections had a higher average body mass

(effect of social connections^2: b = -0.066, t = -2.36, *P* = <0.0001, <0.0001, <0.0001, Table 4,

Fig. 5a). This relationship was also significant when only opposite-sex connections were

considered (effect of opposite-sex social connections^2: b = -0.20, t = 1.54, *P* = 0.0024, 0.0031,

0.0005, Table 4, Fig. 5b). Average body mass was not significantly different between the two

enclosures (all social connections: b = -1.03, t = -0.40, *P* = 0.69; opposite-sex social

connections: b = -0.16, t = -0.058, *P* = 0.95, Table 4). Survival did not predict average body

mass in either model (all social connections: b = 13.37, t = 1.52, P = 0.14; opposite-sex social

connections: b = 10.80, t = 1.22, *P* = 0.23, Table 4). Although the number of measurements we

obtained to calculate each individual's average body mass varied (Fig. 5), it did not affect our

measure of average body mass in either model (all social connections: b = -0.12, t = -0.20, *P* =

0.84; opposite-sex social connections: b = -0.20, t = -0.31, *P* =0.76, Table 4).

428   Discussion

As expected, both female and male prairie voles varied in the frequency of social

interactions, as reflected in our social network analyses that quantified social network degree

using the frequency of temporal and spatial co-occurrence generated by our RFID system.

Female and male voles with an intermediate number of social interactions had the greatest

mating success and produced the greatest number of offspring, though the latter was only the

case when we included all interactions with other voles, and not when only opposite-sex

connections were considered.

We tested our predictions about the fitness benefits of sociality using two different sets

of social network data, one including social connections with all voles and one only including

connections with opposite-sex individuals. This allowed us to investigate the potential costs and

benefits of sociality overall (e.g., social interactions of a female vole with other females and

males) as well as specifically social interactions between opposite-sex individuals, which may

be more directly relevant to mating and reproductive success. We found that voles with an

intermediate number of social connections with all voles (i.e., more social) had significantly

higher mating and reproductive success whereas voles with an intermediate number of

opposite-sex connections had significantly higher mating success but not higher reproductive

success. The process of running the randomizations was somewhat different for the two sets of

networks (all social connections or just opposite-sex connections), which could contribute to the

observed differences in the results between the two sets of networks. We limited permutations

for the opposite-sex network to within the same sex (where female social network data were

being swapped for another female and male data were being swapped for another male) so

these swaps could only be done between approximately half as many individuals each time. By

keeping the sex of each individual in the association consistent, the structure of the

randomizations was more like the data it was being compared to instead of comparing all

possible connections (Farine, 2017a). However, since the permutations were done on the raw

data and then we pulled only the opposite-sex connections from these networks, it is possible

that some of the 10,000 permutations affected the same-sex connections in the raw data

(therefore changing a social interaction that is not included in data used for the linear model)

and therefore may not have changed the estimate for the relationship between opposite-sex

connections and the response variable every time, whereas in the models with all connections

included, a relevant social interaction would have changed every time, which would then change

the model estimate for the relationship between social connections and the response variable

some amount every time. Since we ran such a large number of these permutations, this may not

have affected the overall result of the tests, but it is a limitation of the method.

Using our measure of sociality, our results suggest the possibility of stabilizing selection

on sociality because voles that co-occurred spatially and temporally with very few or very many

conspecifics had the lowest mating success and, at least when considering all social

connections, had the lowest reproductive success. Both McGuire et al. (2002) and Solomon and

Keane (2018) showed that large social groups do not increase female reproductive success.

McGuire et al. (2002) also showed that female prairie voles that lived in large groups had fewer

offspring survive to 12 or 30 days of age. Similarly, Solomon and Keane (2018) showed that

females did not benefit from living in large social groups in two other natural populations. These

studies are consistent with our results that a very large number of social connections (which

should occur in large social groups) does not increase reproductive success of breeding

females, and our results show that this is also true for males.

We only recorded the association between one measure of sociality and mating and

reproductive success in one year and the effects of the number of social connections on fitness

could be altered when environmental conditions change. For example, the Female Dispersion

Hypothesis would predict that if our measure of sociality reflects the socially monogamous

behavior of male prairie voles, our measure of sociality should be positively correlated with male

mating success when females are spatially clumped as males that have more social

connections with clumped females should have higher mating success (Shuster and Wade,

2003; Dobson et al., 2010; Lukas and Clutton-Brock, 2013). In natural populations of prairie

voles, density is quite variable across years (Getz et al., 1993, 2001) and some previous

observational studies of prairie voles in field settings suggested that socially monogamous

behavior is more common at low densities (McGuire et al.,1990; Solomon et al., 2009; but see

Getz and McGuire,1993). There is also some evidence that resource distribution may impact the

mating strategy of prairie voles and this effect may be mediated through its influence on density

(Streatfeild et al., 2011). This suggests the possibility that selection on the social behavior of

prairie voles varies among years due to changes in population or female density but additional

multi-year studies measuring a broader array of social behaviors in free-living voles are needed

to test this prediction.

One possible explanation for an intermediate level of sociality being associated with the

highest mating success and potentially highest reproductive success is that this reflects a

tradeoff between devoting time to social interactions (although we do not know the type of social

interaction occurring) with other voles and time to other behaviors like foraging and parental

care. Although high levels of sociality can have beneficial effects on individual fitness, it may

also carry costs for an individual's health or physical condition (Nunn et al., 2015). Indeed, we

found that male voles with the most social connections had the lowest body mass, suggesting

that there may be a reduction in body condition associated with a very high level of sociality.

This could reflect the energetic costs associated with having many social connections or living in

a large group (e.g., Lutermann et al., 2013), or these could be agonistic interactions with males

on neighboring territories, resulting in males investing more time in territory defense than males

with fewer neighbors. Why males with very few social connections were also lighter in body

mass is not clear but these males may have been of lower phenotypic quality given that they

had few social connections, low body mass, and low mating and reproductive success.

Alternatively, having fewer social connections could result in a loss of body mass if these males

had no assistance in territory defense and thus, expended more energy than males with more

social connections (e.g., having a female social partner). Females likely face many of these

same tradeoffs, but as we did not test quality in females (due to changes in mass being linked to

pregnancy) and so further study is needed to investigate this relationship in females.

Our results suggest that it is not advantageous for voles to have social connections with

too many opposite-sex conspecifics. One possible explanation is that individuals with an

intermediate number of social connections may better balance the trade-off between the

number and quality or strength of social relationships. For example, individuals with the highest

social network degree may just have many weak social or agonistic connections, which may not

result in more matings or increased reproductive success. Individuals with an intermediate

social network degree may have more affiliative social connections that are strong enough to

result in matings than voles at either extreme. This also is reflected in the fact that voles have

many more social connections than actual mating partners (Fig. 2), where the range of the

number of mates varies from 0 to 4 while the range of the number of social connections with

opposite-sex individuals is from 0 to 14. This is supported by studies of the association between

the strength of social connections and fitness in cercopithecine primates (baboons) where

females with strong social bonds with other females in their group have higher offspring survival

(Silk et al., 2003, 2009) or longevity (Silk et al., 2010). Similar relationships between the

strength of social bonds and fitness have also been found in male primates; male Assamese

macaques (*Macaca assamensis*) with strong social bonds to other males (including unrelated

males) sired more offspring than those with fewer strong bonds with other males (Schülke, et

al., 2010). As the number of social connections increase, the strength of association of each of

these social connections may decline (Whitehead 2008), thus prairie voles may be constrained

by the number of social connections in which they can invest enough time to result in successful

mating or rearing of offspring given that prairie voles exhibit biparental care. Individuals that can

best balance this trade-off between the number and strength of social connections may have

the highest mating and reproductive success.

It is of course likely that the fitness benefits of the quantity versus quality of social

connections may vary according to whether the modal social structure of the species is group-

living (such as primate species mentioned above where strong social bonds increase fitness) or

its mating system. For example, Ryder et al. (2009) found a positive association between the

number of social connections (social network degree) and number of offspring sired in male

manakins. As this is a lekking species, coordinated male displays may make male-male

connections a more important factor for mating success than in prairie voles. Additionally, the

short-term coalitions at leks may make the strength of the relationship less important than in

species like prairie voles. Studies like these are rare and so future studies across a broader

array of species with different mating systems will be needed to fully characterize the

relationship between the number and strength of social connections and measures of fitness.

Doing so will help provide insight into how individuals within a species balance the fitness costs

and benefits of social behavior, thereby providing a complementary approach to comparative

studies regarding the evolution of social behavior.

Acknowledgements

We thank the Ecology Research Center (ERC) at Miami University for permitting us to conduct

this study and Jeremy Fruth, field manager at the ERC, for coordinating the project. We also

thank James Lichter, Sage Sparks, Caleb Collins, Tony Leon, Meghan Charles, and Staci

Burrow for assisting on the project. We also thank the American Society of Mammalogists

(Grant-in-Aid of Research to A. Sabol) and University of Michigan (B. Dantzer) for funding.

Data Statement

All raw data and R code are available under CC-BY from FigShare at the link

https://figshare.com/projects/How_does_individual_variation_in_sociality_influence_fitness_in_p

rairie_voles_/71969).

Literature Cited

Alexander, R. D. (1974). The evolution of social behaviour. *Annual Review of Ecology, Evolution,*

and Systematics, 5, 325-383.

Armitage, K. B. (1998). Reproductive strategies of yellow-bellied marmots: Energy conservation

and differences between the sexes. *Journal of Mammalogy, 79*, 385–393.

Blanckenhorn, W. U., Preziosi, R. F. & Fairbairn, D. J.(1995). Time and energy constraints and

the evolution of sexual size dimorphism — to eat or to mate? *Evolutionary Ecology, 9,* 369-381.

doi:10.1007/BF01237760

Clutton-Brock, T. H. (1989). Review lecture: mammalian mating systems. *Proceedings of the*

Royal Society B, 236, 339-372.

Clutton-Brock, T.H., Brotherton, P.N.M., Smith, R., McIllrath, G.M., Kansky, R., Gaynor, D., O'rian,

M.J., Skinner, J.D. (1998). Infanticide and expulsion of females in a cooperative mammal.

Proceedings of the Royal Society B, 265, 2291-2295.

Cochran, G. R., & Solomon, N. G. (2000). Effects of food supplementation on the social

organization of prairie voles (*Microtus ochrogaster*). *Journal of Mammalogy*, *81*, 746-757.

Côté, I.M., & Poulinb R. (1995). Parasitism and group size in social animals: a meta-analysis.

Behavioral Ecology, 6, 159-165.

Creel, S., Dantzer, B., Goymann, W. & Rubenstein, D.R. (2013). The ecology of stress: effects of

the social environment. *Functional Ecology, 27,* 66–80.

Dantzer, B., Goncalves, I.B., Spence-Jones, H.C., Bennett, N.C., Heistermann, M., Ganswindt, A.,

Dubuc, C., Gaynor, D., Manser, M.B., Clutton-Brock, T.H. (2017). The influence of stress

hormones and aggression on cooperative behaviour in subordinate meerkats. *Proceedings of*

the Royal Society B, 284, 20171248.

Dobson, F. S., Way, B. M., & Baudoin C. (2010). Spatial dynamics and the evolution of social

monogamy in mammals. *Behavioral Ecology, 21,* 747–752. doi:10.1093/beheco/arq048

584 Donaldson, Z. R., & Young, L. J. (2008). Oxytocin, vasopressin, and the neurogenetics of

585  sociality. *Science, 322,* 900-904. doi:10.1126/science.1158668

586 Eisenberg, J.F., Muckenhirn, N.A., Rudran, R. (1972). The relation between ecology and social

587  structure in primates. *Science, 176*, 863-874.

588 Emlen, S.T. (1984). *Cooperative breeding in birds and mammals. Behavioural Ecology: An*

589  *Evolutionary Approach*. J.R. Krebs and N.B. Davies (Eds.), pp. 305-339, Blackwell, Oxford.

590 Emlen, S.T. (1994). Benefits, constraints and the evolution of the family. *Trends in Ecology and*

591  *Evolution, 9,* 282-285.

592 Ewald, P. W. (1994) *Evolution of Infectious Disease*. Oxford: Oxford University Press.

593 Farine, D. R. (2013). Animal social network inference and permutations for ecologists in R using

594  *asnipe*. *Methods in Ecology and Evolution, 4,* 1187-1194. doi:10.1111/2041-210X.12121

595 Farine, D. R., Firth, J. A., Aplin, L. M., Crates, R. A., Culina, A., Garroway, C.J., Hinde, C. A.,

596  Kidd, L. R., Milligan, N. D., Psorakis, I., Radersma, R., Verhelst, B., Voelkl, B., & Sheldon, B. C.

597  (2015). The role of social and ecological processes in structuring animal populations: a case

598  study from automated tracking of wild birds. *Royal Society Open Science, 2.*

599  doi:10.1098/rsos.150057

600 Farine, D. R. (2017a). A guide to null models for animal social network analysis. *Methods in*

601  *Ecology and Evolution, 8,* 1309-1320. doi:10.1111/2041-210X.12772

602 Farine, D. R. (2017b). *asnipe: Animal Social Network Inference and Permutations for E*cologists.

603  R package version 1.1.4. Available online at: https://CRAN.R-project.org/package=asnipe

604 Faulkes, C. G., Bennett, N. C., Bruford, M. W., O'brien H. P., Aguilar G. H., & Jarvis, J. U. M.

605  (1997). Ecological constraints drive social evolution in the African mole–rats. *Proceedings of the*

606  *Royal Society of London B: Biological Sciences, 267*. doi:10.1098/rspb.1997.0226

607 Fox, J., & Weisberg, S. (2011). An {R} Companion to Applied Regression, Second Edition.

608  Thousand Oaks CA: Sage. URL: http://socserv.socsci.mcmaster.ca/jfox/Books/Companion

Getz, L. L., & Hofmann, J. E. (1986). Social organization in free-living prairie voles, *Microtus*

ochrogaster. *Behavioral Ecology and Sociobiology, 18*, 275-282.

Getz, L. L., Hofmann, J. E., McGuire, B., & Dolan, T. W. III. (2001). Twenty-five years of

population fluctuations of *Microtus ochrogaster* and *M. pennsylvanicus* in three habitats in east-

central Illinois. *Journal of Mammalogy, 82*, 22–34.

Getz, L. L., & McGuire, B. (1993). A comparison of living singly and in male-female pairs in the

prairie vole, *Microtus ochrogaster*. *Ethology, 94,* 265-278. doi:10.1111/j.1439-

0310.1993.tb00444.x

Getz, L. L., McGuire, B., Pizzuto, T., Hofmann, J. E., & Frase, B. (1993). Social organization of

the prairie vole (*Microtus ochrogaster*). *Journal of Mammalogy. 74,* 44–58. doi:

10.2307/1381904

Getz, L. L., McGuire, B., Hofmann, J., Pizzuto, T., & Frase, B. (1994). Natal dispersal and

philopatry in prairie voles (*Microtus ochrogaster*): settlement, survival, and potential reproductive

success. *Ethology Ecology & Evolution, 6,* 267-284.

Hatchwell, B. J., & Komdeur, J. (2000). Ecological constraints, life history traits and the evolution

of cooperative breeding. *Animal Behaviour, 59,* 1079–1086. doi:10.1006/anbe.2000.1394

Jetz, W., & Rubenstein, D. R. (2011). Environmental uncertainty and the global biogeography of

cooperative breeding in birds. *Current Biology, 21,* 72-78. doi:10.1016/j.cub.2010.11.075.

Kappeler, P. M., Cremer, S., & Nunn, C. L. (2015). Sociality and health: impacts of sociality on

disease susceptibility and transmission in animal and human societies. *Philosophical*

Transactions of the Royal Society B: Biological Sciences, 370. doi:10.1098/rstb.2014.0116

Keane, B., Bryant, L., Goyal, U., Williams, S., Kortering, S. L., Lucia, K. E., Richmond, A. R., &

Solomon, N. G. (2007). No effect of body condition at weaning on survival and reproduction in

prairie voles. *Canadian Journal of Zoology, 85,* 718-727.

Kleiber, C., & Zeileis, A. (2008). Applied econometrics with R. New York: Springer-Verlag. ISBN

978-0-387-77316-2. https://CRAN.R-project.org/package=AER" https://CRAN.R-

project.org/package=AER

Krause, J., & Ruxton, G. D. (2002). *Living in groups.* Oxford: Oxford University Press.

Lambert, C. T., Sabol, A. C., & Solomon, N. G. (2018). Genetic monogamy in socially

monogamous mammals is primarily predicted by multiple life history factors: a meta-analysis.

Frontiers in Ecology and Evolution, 6, 139. doi:10.3389/fevo.2018.00139

Langwig, K. E., Frick, W. F., Bried, J. T., Hicks, A. C., Kunz, T. H. & Marm Kilpatrick, A. (2012),

Sociality, density-dependence and microclimates determine the persistence of populations

suffering from a novel fungal disease, white-nose syndrome. *Ecology Letters, 15:* 1050-1057.

doi:10.1111/j.1461-0248.2012.01829.x

Lemen, C. A., & Freeman, P. W. (1985). Tracking mammals with fluorescent pigments: a new

technique. *Journal of Mammalogy, 66,* 134–136.

Lott, D. F. (1991). *Intraspecific variation in the social systems of wild vertebrates.* Cambridge:

Cambridge University Press.

Lucia, K. E., Keane, B., Hayes, L. D., Lin, Y. K., Schaefer, R. L., & Solomon, N. G. (2008).

Philopatry in prairie voles: an evaluation of the habitat saturation hypothesis. Behavioral.

Ecology, 19, 774–783. doi: 10.1093/beheco/arn028

Lukas, D., & Clutton-Brock, T. H. (2013). The evolution of social monogamy in mammals.

Science, 341, 526 -530. doi:10.1126/science.1238677

Lukas, D., & Clutton-Brock, T. H. (2017). Climate and the distribution of cooperative breeding in

mammals. *Royal Society Open Science, 4.* doi:10.1098/rsos.160897

Lukas, D., & Huchard, E. (2014). The evolution of infanticide by males in mammalian societies.

Science, 346, 841-844. doi:0.1126/science.1257226

Lutermann, H., Bennett, N. C., Speakman, J. R., & Scantlebury, M. (2013). Energetic benefits of

sociality offset the costs of parasitism in a cooperative mammal. *PLOS ONE*, *8*, e57969.

doi:10.1371/journal.pone.0057969

Mabry, K. E., Streatfeild, C. A., Keane, B., & Solomon, N. G. (2011). avpr1a length polymorphism

is not associated with either social or genetic monogamy in free-living prairie voles. *Animal*

Behaviour, 81, 11-18.

McGuire, B., & Getz, L.L. (2010). Alternative male reproductive tactics in a natural population of

prairie voles, *Microtus ochrogaster. Acta Theriol, 55,* 261-270. doi:10.4098/j.at.0001-

7051.077.2009

McGuire, B., Getz, L. L., & Oli, M. K. (2002). Fitness consequences of sociality in prairie voles,

Microtus ochrogaster: influence of group size and composition. *Animal Behaviour, 64,* 645-654.

doi:10.1006/anbe.2002.3094.

McGuire, B., Pizzuto, T., Getz, L.L. (1990). Potential for social interaction in a natural population

of prairie voles (*Microtus ochrogaster*). *Canadian Journal of Zoology, 68,* 391-398.

Nunn, C.L., Jordán, F., McCabe, C.M., Verdolin, J.L., Fewell, J.H. (2015). Infectious disease and

group size: more than just a numbers game. *Philosophical Transactions of the Royal Society B,*

370, 20140111.

Okhovat, M., Berrio, A., Wallace, G., Ophir, A. G., & Phelps, S. M. (2015). Sexual fidelity trade-

offs promote regulatory variation in the prairie vole brain. *Science, 350,* 1371-1374.

Ophir, A.G., Wolff, J.O., Phelps, S.M. (2008). Variation in neural V1aR predicts sexual fidelity and

space use among male prairie voles in semi-natural settings. *PNAS, 105*, 1249-1254.

Psorakis, I., Roberts, S. J., Rezek, I., & Sheldon, B. C. (2012). Inferring social network structure in

ecological systems from spatio-temporal data streams. *Journal of the Royal Society Interface, 9,*

3055–3066. doi: 10.1098/rsif.2012.0223

R Core Team (2017). *R: A Language and Environment for Statistical Computing*. Vienna: R

Foundation for Statistical Computing. Available online at: https://www.R-project.org/.

Richmond, M., & Conaway, C. H. (1969). Management, breeding, and reproductive performance

of the vole, *Microtus ochrogaster*, in a laboratory colony. *Laboratory Animal Care, 19,* 80-87.

Rubenstein, D. R., & Lovette, I. J. (2007) Temporal environmental variability drives the evolution

of cooperative breeding in birds. *Current Biology, 17,* 1414-1419.

doi:10.1016/j.cub.2007.07.032.

Ryder, T.B., Parker, P.G., Blake, J.G., & Loiselle, B.A. (2009). It takes two to tango: reproductive

skew and social correlates of male mating success in a lek-breeding bird.

Proceedings of the Royal Society B: Biological Sciences, 276, 2377-2384.

doi:10.1098/rspb.2009.0208

Sabol, A. C., Solomon, N. G., & Dantzer, B. (2018). How to study socially monogamous behavior

in secretive animals? Using social network analyses and automated tracking systems to study

the social behavior of prairie voles. *Frontiers in Ecology and Evolution, 6,* 178.

doi:10.3389/fevo.2018.00178

Schradin, C., & Pillay, N. (2005). Intraspecific variation in the spatial and social organization of the

African striped mouse, *Journal of Mammalogy, 86*, 99–107.

Schradin, C., König, B., Pillay, N. (2010). Reproductive competition favours solitary living while

ecological constraints impose group-living in African striped mice. *Journal of Animal Ecology,*

73, 515-521.

Schülke, O., Bhagavatula, J., Vigilant, L., & Ostner, J. (2010). Social bonds enhance reproductive

success in male macaques. *Current Biology, 20,* 2207-2210.

doi:10.1016/j.cub.2010.10.058.

Shen, S-F., Emlen, S.T., Koenig, W.D., Rubenstein, D.R. (2017). The ecology of cooperative

breeding behaviour. *Ecology Letters, 20,* 708-720.

Shuster, S. M., & Wade, M. J. (2003). *Mating systems and mating strategies*. Princeton: Princeton

University Press.

Shuster, S. M., Willen, R. M., Keane, B., & Solomon, N. G. (2019). Alternative mating tactics in

socially monogamous prairie voles, Microtus ochrogaster. *Frontiers in Ecology and Evolution*, 7.

doi:10.3389/fevo.2019.00007

Sikes, R. S., Bryan, J. A., Byman, D., Danielson, B. J., Eggleston, J., Gannon, M. R., Gannon, W.

L., Hale, D. W., Jesmer, B. R., Odell, D. K., Olson, L. E., Stevens, R. D., Thompson, T. A.,

Timm, R. M., Trewhitt, S. A., & Willoughby, J. R. (2016). 2016 Guidelines of the American

Society of Mammalogists for the use of wild mammals in research and education. *Journal of*

Mammalogy 97, 663–688.

Silk, J. B. (2007). The adaptive value of sociality in mammalian groups. *Philosophical*

Transactions of the Royal Society B: Biological Sciences, *362*, 539–559.

doi:10.1098/rstb.2006.1994

Silk, J. B., Alberts, S. C., & Altmann, J. (2003). Social bonds of female baboons enhance infant

survival. *Science*, *302*, 1231-1234. doi:10.1126/science.1088580

Silk, J.B., Beehner, J.C., Bergman, T.J., Crockford, C., Engh, A.L., Moscovice, L.R., Wittig, R.M.,

Seyfarth, R.M., Cheney, D.L. (2009). The benefits of social capital: close social bonds among

female baboons enhance offspring survival. *Proceedings of the Royal Society B, 276,* 3099-

3104.

Silk, J.B., Beehner, J.C., Bergman, T.J., Crockford, C., Engh, A.L., Moscovice, L.R., Wittig, R.M.,

Seyfarth, R.M., Cheney, D.L. (2010). Strong and consistent social bonds enhance the longevity

of female baboons. *Current Biology, 20,* 1359-1361.

Solomon, N. G. (1993). Comparison of parental behavior in male and female prairie voles

(*Microtus ochrogaster*). *Canadian Journal of Zoology, 71,* 1991–1994.

Solomon, N. G., & Keane, B. (2018). Dispatches from the field: sociality and reproductive success

in prairie voles. *Animal Behaviour*, *143*, 193–203. doi:10.1016/j.anbehav.2018.07.001

Solomon, N. G., & Jacquot, J. J. (2002). Characteristics of resident and wandering prairie voles,

Microtus ochrogaster. *Canadian Journal of Zoology, 80*, 951-955.

Solomon, N. G., Keane, B., Knoch, L. R., & Hogan, P. J. (2004). Multiple paternity in socially

monogamous prairie voles (Microtus ochrogaster). *Canadian Journal of Zoology, 82,* 1667–

1671.

Solomon, N.G., Richmond, A. R., Harding, P. A., Fries, A., Jacquemin, S., Schaefer, R. L., Lucia,

K. E., & Keane, B. (2009). Polymorphism at the avpr1a locus in male prairie voles correlated

with genetic but not social monogamy in field populations. *Molecular Ecology, 18,* 4680-4695.

Spiegel, O., Sih, A., Leu, S. T., & Bull, C. M. (2017). Where should we meet? Mapping social

network interactions of sleepy lizards shows sex-dependent social network structure. *Animal*

Behaviour, 136, 207-215. doi:10.1016/j.anbehav.2017.11.001.

Streatfeild, C. A., Mabry, K. E., Keane, B., Crist, T. O., & Solomon, N. G. (2011). Intraspecific

variability in the social and genetic mating systems of prairie voles, *Microtus ochrogaster*.

Animal Behaviour, 82, 1387–1398. doi: 10.1016/j.anbehav.2011.09.023

van Schaik, C. (1983). Why are diurnal primates living in groups? *Behaviour, 87*, 120-144.

Walum, H., & Young, L. J. (2018). The neural mechanisms and circuitry of the pair bond. *Nature*

Reviews Neuroscience, *19*, 643–654. doi:10.1038/s41583-018-0072-6

Whitehead, H. (2008). *Analyzing animal societies: quantitative methods for vertebrate social*

analysis. Chicago: University of Chicago Press.

Whiteman, N. K., & Parker, P. G. (2004). Effects of host sociality on ectoparasite population

biology. *Journal of Parasitology, 90*, 939-947. doi:10.1645/GE-310R

Wickham, H. (2009). Ggplot2: Elegant graphics for data analysis. New York: Springer-Verlag

Young, L. J., & Wang, Z. (2004). The neurobiology of pair bonding. *Nature Neuroscience*, *7*,

1048–1054. doi:10.1038/nn1327

760 **Table 1**. Effects of the number of social connections on vole mating success (number of

761 different individuals a vole produced offspring with).

Social Network	Variable	Estimate	SE	z-value	P-value from GLM	P-values from Randomization Tests
All individuals	**Intercept**	**-2.66**	**0.65**	**-4.06**	**<0.0001**	NA
	Sex (M)	0.68	0.92	0.73	0.46	NA
	Enclosure (LD)	**1.89**	**0.41**	**4.55**	**<0.0001**	NA
	Survival	**2.07**	**0.89**	**2.33**	**0.020**	NA
	Degree	0.083	0.21	0.40	0.69	0.12, 0.0098, 0.14
	Degree^2	**-0.012**	**0.013**	**-0.88**	**0.38**	**0.012, 0.0093, <0.0001**
	Sex (M) x Degree	**-0.21**	**0.23**	**-0.90**	**0.37**	**0.021, 0.0017, <0.0001**
	Sex (M) x Degree^2	**0.014**	**0.015**	**0.94**	**0.35**	**0.01, 0.0018, <0.0001**
Opposite-sex individuals	**Intercept**	**-2.81**	**0.70**	**-4.08**	**<0.0001**	NA
	Sex (M)	0.78	0.91	0.86	0.39	NA
	Enclosure (LD)	**2.03**	**0.43**	**4.72**	**<0.0001**	NA
	Survival	**1.75**	**0.84**	**2.09**	**0.037**	NA
	Degree	0.19	0.31	0.63	0.53	0.18, 0.19, <0.0001
	Degree^2	**-0.028**	**0.028**	**-1.01**	**0.31**	**0.0011, 0.0035, 0.001**
	Sex (M) x Degree	**-0.53**	**0.38**	**-1.42**	**0.15**	**<0.0001,<0.0001, 0.0046**
	Sex (M) x Degree^2	**0.057**	**0.042**	**1.36**	**0.17**	**<0.0001, <0.0001, 0.0068**

Degree refers to the number of social connections for each individual either with all voles or

opposite-sex voles. Note that relationships involving social network data have three *P*-values

because those regression coefficients were compared to those from randomized networks three

times to determine if they were consistently significant (see methods for details). Survival refers to

the proportion of days the vole was in the enclosure based on when it was last recorded. "LD" is

low-density enclosure.

Table 2. Effects of the number of social connections on vole reproductive success (number of

offspring produced that survived until emergence from the natal nest).

Social Network	Variable	Estimate	SE	z-value	P-value from GLM	P-values from Randomization Tests
All individuals	**Intercept**	**-2.90**	**0.66**	**-4.14**	**<0.0001**	NA
	Sex (M)	0.61	0.93	0.65	0.52	NA
	Enclosure (LD)	**2.10**	**0.40**	**5.24**	**<0.0001**	NA
	Survival	**2.49**	**0.82**	**3.03**	**0.0025**	NA
	Degree	0.053	0.19	0.28	0.78	0.33, 0.25, 0.19
	Degree^2	**-0.0085**	**0.011**	**-0.77**	**0.44**	**0.002, 0.035, 0.015**
	Sex (M) x Degree	-0.14	0.21	-0.64	0.52	0.12, 0.18, 0.12
	Sex (M) x Degree^2	0.0082	0.013	0.63	0.53	0.021, 0.11, 0.022
Opposite-sex individuals	**Intercept**	**-2.97**	**0.67**	**-4.41**	**<0.0001**	NA
	Sex (M)	0.58	0.91	0.64	0.53	NA
	Enclosure (LD)	**2.21**	**0.41**	**5.40**	**<0.0001**	NA
	Survival	**2.19**	**0.77**	**2.83**	**0.0046**	NA
	Degree	0.15	0.28	0.52	0.60	0.20, 0.53, 0.54
	Degree^2	-0.022	0.024	-0.93	0.35	0.14, 0.099, 0.12
	Sex (M) x Degree	-0.39	0.38	-1.04	0.30	0.047, 0.024, 0.11
	Sex (M) x Degree^2	0.038	0.041	0.92	0.36	0.063, 0.010, 0.02

Degree refers to the number of social connections for each individual either with all voles or

opposite-sex voles. Note that relationships involving social network data have three *P*-values

because those regression coefficients were compared to those from randomized networks three

times to determine if they were consistently significant (see methods for details). Survival refers

to the proportion of days the vole was in the enclosure based on when it was last recorded. "LD"

is low-density enclosure.

Table 3. Overall effects of vole mating success (number of different individuals with which a

vole produced offspring) on reproductive success (number of offspring produced that survived

until emergence from the natal nest).

Variable	Estimate	SE	z-value	P-value from GLM
Intercept	**-2.13**	**0.44**	**-4.83**	**<0.0001**
Number of Mates	**0.64**	**0.13**	**4.88**	**<0.0001**
Sex (M)	0.26	0.49	0.53	0.60
Enclosure (LD)	**1.53**	**0.45**	**3.37**	**0.00076**
# of Mates x Sex (M)	-0.093	0.17	-0.56	0.58

780 Since this model did not include social network data we did not perform the randomizations to

781 generate three *P*-values, we present the *P*-values from the GLM. "LD" is low-density enclosure.

783   **Table 4**. Effects of the number of social connections on average body mass for males.

Social Network	Variable	Estimate	SE	t-value	P-value from LM	P-values from Randomization Tests
All individuals	**Intercept**	**43.42**	**2.70**	**16.10**	**<0.0001**	NA
	Enclosure (LD)	-1.03	2.57	-0.40	0.69	NA
	Survival	13.37	8.78	1.52	0.14	NA
	# of measures	-0.12	0.60	-0.20	0.84	NA
	Degree	**0.95**	**0.56**	**1.70**	**0.099**	**<0.0001, 0.0001, <0.0001**
	Degree^2	**-0.066**	**0.028**	**-2.36**	**0.025**	**<0.0001, <0.0001, <0.0001**
Opposite-sex individuals	**Intercept**	**44.31**	**2.81**	**15.78**	**<0.0001**	NA
	Enclosure (LD)	-0.16	2.80	-0.058	0.95	NA
	Survival	10.80	8.89	1.22	0.23	NA
	# of measures	-0.20	0.63	-0.31	0.76	NA
	Degree	**1.54**	**1.22**	**1.26**	**0.22**	**0.0035, 0.0036, 0.0332**
	Degree^2	**-0.20**	**0.13**	**-1.54**	**0.13**	**0.0024, 0.0031, 0.0005**

Degree refers to the number of social connections for each individual either with all voles or

opposite-sex voles. Relationships involving social network data have three *P*-values because

those regression coefficients were compared to those from randomized networks three times to

determine if they were consistently significant (see methods). "# of measures" refers to number

of times we measured body mass, which were used to generate average mass for each

individual. Survival refers to proportion of days the vole was in the enclosure. "LD" is low-density

enclosure.

Figure 1. a) The number of females, males, and offspring in each of our two enclosures over

time during the study based on the number of unique individuals live-trapped during each two-

week period of the field season. Note that the area of enclosures is equal, so the number of

voles in each enclosure can be used to compare relative density between the two, b) The sex

ratio in each enclosure, calculated as the proportion of total adult voles trapped during each

two-week period of the field season that were males, over time of the study.

Figure 2. Both females and males with an intermediate number of social connections with a) all

voles in their enclosure (both same- or opposite-sex individuals) or b) only opposite-sex

individuals in their enclosure had higher mating success (defined as the number of different

individuals with which they produced offspring). Points are jittered with males shown in blue and

females in red. Full statistical results shown in Table 1.

Figure 3. a) Both females and males with an intermediate number of social connections with all

voles in their enclosure (both same- and opposite-sex individuals) produced more offspring that

survived to emergence from the natal nest. b) Female and male voles with an intermediate

number of social connections with only opposite-sex individuals in their enclosure also tended to

produce more offspring. Points are jittered with red points being females and blue points being

males. Full statistical results shown in Table 2.

Figure 4. Both female and male prairie voles that had higher mating success (produced

offspring with a greater number of different mates) produced a great number of offspring that

survived to emergence from the natal nest. Points for females and males are jittered. Full results

shown in Table 3.

Figure 5. Male voles with an intermediate number of social connections a) with both female and

male voles in their enclosure or b) with just female voles, were significantly heavier over the

course of this study. Body mass for males was averaged for the entire duration of this study.

The number of times we measured body mass ("N") varied among males so the size of each

point is scaled based on the number of recorded mass measurements we have for each

individual. Full results shown in Table 4.

Figure A1. Layout of the two RFID arrays in the enclosures. The RFID system was kept at each

array in each enclosure for three days in the order: array 1 enclosure 1, array 2 enclosure 1,

array 1 enclosure 2, and array 2 enclosure 2 and then repeated for the duration of the field

season.