How does individual variation in sociality influence fitness in prairie
voles?

Anne C. Sabol[a,*], Connor T. Lambert[b], Brian Keane[b], Nancy G. Solomon[b], Ben Dantzer[a,c]

[a]Department of Psychology, University of Michigan, Ann Arbor, MI, USA

[b]Department of Biology, Miami University, Oxford, Ohio, USA

[c]Department of Ecology and Evolutionary Biology, University of Michigan, Ann Arbor, MI, USA

* Corresponding author e-mail address: acsabol72@gmail.com

Abstract

Comparative studies aid in our understanding of specific conditions favoring the initial evolution

of different types of social behaviors, yet there is much unexplained intraspecific variation in the

expression of social behavior that comparative studies have not yet addressed. The proximate

causes of this individual variation in social behavior within a species have been examined in

some species but its fitness consequences have been less frequently investigated. In this study,

we quantified the fitness consequences of variation in the sociality of prairie voles (*Microtus*

ochrogaster). We characterized sociality of female and male voles in semi-natural enclosures

using an automated behavioral tracking system paired with social network analyses to quantify

the degree of spatial and temporal co-occurrence of different voles. We then assessed the

relationship between these metrics of sociality with individual mating success (number of

different conspecifics with which an individual produced offspring) and reproductive success

(total number of offspring surviving to first capture). We measured the number of social

connections each individual had with all voles and only with opposite-sex voles by calculating

unweighted degree through social network analyses. Both female and male voles varied in the

number of social connections they had with all voles (both sexes) and with opposite-sex

conspecifics. Both females and males with an intermediate number of social connections with

voles of both sexes had higher mating success overall. In our analyses that considered all social

connections with voles of both sexes, voles with an intermediate number of social connections

produced more offspring. Males with a very high or low number of social connections also had

the lowest average body mass. Overall, our results suggest some limit on the fitness benefits of

sociality. Although there was substantial individual-variation in our measure of vole social

behavior, intermediate levels of social connections may be most favorable.

Introduction

Sociality comes in many forms across a diversity of taxa, ranging from loose, temporary

associations during breeding to long-term group associations whose members exhibit complex

social behaviors such as cooperative hunting or alloparental care. Different forms of sociality are

thought to provide a variety of fitness benefits, including shared resources, reduced predation,

communication, or better access to mates (Eisenberg et al., 1972; Alexander, 1974; van Schaik

1983; Emlen, 1984; Krause and Ruxton, 2002, Silk, 2007). Sociality may also come with costs

associated with group-living, including increased disease transmission (Ewald, 1994; Langwig et

al., 2004; Kappeler et al., 2015), parasite load (Coote and Poulinb, 1995; Whiteman and Parker,

2004), physiological stress (Creel et al., 2013), reproductive interference by other group

members (Clutton-Brock et al., 1998; Lukas and Huchard, 2014), as well as the time and energy

costs devoted to developing and maintaining social connections that could otherwise be

devoted towards activities directly related to individual survival or reproduction such as foraging

or parental care. Given that these fitness benefits and costs of sociality may vary in direction or

magnitude due to ecological circumstances such as food abundance, weather patterns, or

temporal variability in these factors (Emlen, 1994; Hatchwell and Komdeur, 2000; Shuster and

Wade, 2003; Schradin and Pillay, 2005; Rubenstein and Lovette, 2007; Schradin et al., 2010;

Shen et al., 2017), it is no surprise that there is considerable variation in the presence or

expression of different types of social behaviors among species.

Comparative studies have been useful in revealing some of the factors favoring the

evolution of different types of social behaviors. For example, phylogenetic meta-analyses show

that social monogamy is more commonly found in mammalian species that experience low

population densities (Lukas and Clutton-Brock, 2013) and genetic monogamy is also more

common in mammalian species with low population densities as well as those that exhibit

parental care (Lambert et al., 2018). Other comparative studies have shown that cooperative

breeding, where subordinate group members care for offspring produced by dominant breeders,

is more likely to be found in mammalian species that live in arid areas (Faulkes et al., 1997;

Lukas and Clutton-Brock, 2017) or in avian species that inhabit areas where rainfall is low and

unpredictable (Jetz and Rubenstein, 2011). While these studies help illuminate some factors

affecting the evolution of social behaviors like social monogamy or cooperative breeding, they

typically cannot address the causes of intraspecific variation in sociality.

Sociality is not always expressed to the same level or in the same way within a species

and there are often a variety of social strategies within the same species (Lott, 1991; Clutton-

Brock, 1989; Getz et al., 1993). The magnitude of variation in the expression of social behavior

within a species may not be as large as that among species but it requires explanation

nonetheless. Although some studies have investigated some of the proximate causes of

individual variation in specific social behaviors such as social monogamy (Young and Wang,

2004; Donaldson and Young, 2008; Ophir et al., 2008; Okhovat et al., 2015; Walum and Young,

2018) or alloparental care (Dantzer et al., 2017), fewer studies document its fitness

consequences. This is not surprising as it requires detailed studies that document the social

behavior of individuals and then relate their degree of sociality to their survival and reproduction.

Most previous studies on this topic have been conducted in primates where investigators

documented how the strength of social relationships ("social bonds") impacted offspring

survival, mating success, or longevity (e.g., Silk et al., 2003, 2009, 2010; Schulke et al., 2010).

By investigating the association between social behavior and fitness within a species, we may

be able to understand how individuals balance the tradeoffs between the costs and benefits of

sociality and therefore obtain an even greater understanding of the evolution of sociality.

We characterized the variability of social behavior in individual female and male prairie

voles (*Microtus ochrogaster*) and its association with their mating success, reproductive

success, and body condition. Prairie voles are an ideal study system to investigate sociality

because both sexes can exhibit natural variation in sociality by displaying different reproductive

strategies (Solomon and Jacquot, 2002). Both females and males can exhibit strategies ranging

from pair bonded, with an established territory to non-territorial, unpaired "wanderers" and both

are known to switch their strategy over the course of their lifetime (Getz and Hofmann, 1986;

Getz et al., 1993; Solomon and Jacquot, 2002; McGuire and Getz, 2010; Shuster et al., 2019).

Further, genetic monogamy and social monogamy are distinct with some female-male pairs

exhibit high levels of social and genetic monogamy, some pairs being socially monogamous but

not genetically monogamous, and some individuals exhibiting no socially or genetically

monogamous behavior or mating patterns (Solomon et al., 2004). Thus, prairie voles may vary

in the number and strength of social associations they have with other individuals. For example,

a genetically monogamous female and male likely have a very strong social association with

each other whereas a wandering individual may have many weak connections with multiple

opposite-sex voles. Additionally, voles may vary in the number of social connections they have

with other individuals than their partner because under certain environmental conditions, such

as high population density (Getz et al., 1993; Cochran and Solomon 2000; Lucia et al. 2008),

voles can also form extended family groups when offspring delay dispersal from the natal nest.

We characterized the social behavior of individual prairie voles in semi-natural

enclosures using an automated monitoring system. Prairie voles were all marked uniquely with

passive integrated technology (PIT) tags and their movements were continuously monitored by

an array of radio-frequency identification (RFID) antennas recording the unique code

transmitted by the PIT tag when they were within range as well as the date and time this

information was transmitted. We used patterns of spatial and temporal co-occurrence generated

from this system with social network analyses to estimate patterns of social association. We did

this by measuring the number of social connections (unweighted degree, hereafter degree)

between same-sex or opposite-sex voles based on co-occurrence data from the RFID

antennas. Individuals with a high degree would have had instances of spatial and temporal co-

occurrence with many other voles whereas those with a low degree had few. We have

previously shown that opposite-sex voles that exhibit close social associations, as generated by

this automated method of data collection, are also more likely to be caught in the same trap

together, exhibit overlapping home ranges, and a show a strong social preference for one

another in a choice experiment (Sabol et al., 2018), suggesting that these measures of degree

reflect the strength of social associations. We characterized the sociality of all voles throughout

the breeding season using social network analyses that generated measures of unweighted

degree, recorded their survival through this period and used parentage analyses to quantify

their mating success (number of different individuals with which they produced offspring) and

reproductive success (number of offspring produced that survived to emergence from the natal

nest).

We classified individuals with a greater number of social network connections with

opposite-sex or same-sex voles as more social, as this corresponded to voles that co-occurred

with more individual voles during the field season. We predicted that voles with a greater

number of social network connections (i.e., more social) would have higher mating and

reproductive success but lower body condition due to the trade-offs associated with high levels

of this type of sociality. We only investigated body condition in males because any changes in

body mass in females is likely directly related to pregnancy, We predicted that male voles that

were more social (had more social network connections) would have lower body condition

(mass) because polygyny typically involves energetic costs and risky travel associated with

finding female mating partners and interacting with male competitors, which may therefore

reduce survival (Blanckenhorn et al 1995; Armitage 1998).

Methods

136   *Study site and study animals*

137   All fieldwork was conducted at Miami University's Ecology Research Center in Oxford,

Ohio from May to August 2017. Voles were released into two separate 0.1 ha enclosures (33 m

x 33 m). The 20-gauge sheet metal walls of the enclosures were 75 cm tall and embedded 45

cm into the ground to prevent vole movements among enclosures. Enclosure walls were topped

with an electrified wire to prevent other small to medium sized mammals (e.g. raccoons and

weasels) from entering the enclosures and disturbing traps. Although this system likely

prevented many mammalian predators from entering the enclosures, the enclosures were open

and accessible to avian predators and snakes. We had multiple animals that were confirmed to

be depredated by owls and also occasionally saw large snakes within the enclosures. Prior to

releasing prairie voles into the enclosures, we live-trapped within the enclosures for 3

continuous days to capture any small mammals (e.g. *Microtus pennsylvanicus*, *Peromyscus*

maniculatus, or *Blarina brevicauda*) and released them outside of the enclosures.

149   We released laboratory-bred 7th and 8th generation prairie voles (descended from voles

originally captured in Illinois) into two enclosures. The pedigree of the laboratory population was

known and to avoid inbreeding, we did not place opposite-sex siblings or parents and their

offspring into the same enclosure. All founding voles were sexually mature (> 31 d, Solomon

1991) but had never bred. Each enclosure was founded with a different density: Enclosure 1

was established by releasing 48 voles (24 females, 24 males) and Enclosure 2 initially

contained 24 voles (12 females, 12 males). These represented densities of 480 voles/ha and

240 voles/ha respectively, which were within the range of vole densities observed in natural

populations (10 to 600 voles/ha: Getz et al., 1994; Getz et al., 2001). The different starting

density was employed to assess the role of density on vole social behavior but, as shown below

(Fig. 1), vole density in Enclosure 1 (high density) decreased over the course of the field

season. Additionally, we do not find or report an effect of density on vole social behavior in this

datset. The vegetation within enclosures consisted primarily of perennial grasses and forbs,

which provided food and cover. Voles were not provided with supplemental food besides the

cracked corn, a low-quality food, used to bait the live-traps.

164   *Recording social behavior of voles*

Prior to release into the enclosures, all founding voles were implanted with a PIT tag

(Biomark: Boise, Idaho) with a unique identification number. Once released, we monitored our

populations through a proprietary system of 12 RFID antennas connected to a central recording

station (BioMark RM310/SM303, Boise, ID) that we have used previously to create social

networks of prairie voles and infer their social associations (Sabol et al., 2018). These antennas

were placed within the enclosures in two different 3 x 4 arrays (Figure S1) that were rotated

from array 1 to array 2 every 3 days and moved from one enclosure to the other every 6 days

during the 14-week study. The antennas recorded the individual PIT tag numbers of each vole

that passed within 15 cm of the antenna once a second for the entire time the animal was within

this radius of the antenna. When multiple tags were within the 15 cm radius, the system

alternated in recording the tag numbers so that both could be detected. This allowed us to

record the natural movements and social associations of individuals in both populations, which

we have previously shown to be comparable and more detailed than traditional methods of

recording social associations in these populations (Sabol et al., 2018).

179   *Monitoring vole reproduction*

We live-trapped both enclosures by placing two Ugglan live- traps (Granhab, Hillerstorp,

Sweden), baited with cracked corn, per grid stake on a 5 x 5 m trapping grid. These live-traps

had a metal cover over the top to shield voles from the elements while in the traps. At the

beginning of the experiment, we live-trapped nearly every day, setting traps in the evening

(2230-2300 h) and checking them approximately two hours later and then leaving them set to be

checked the next morning (0700 h). After the first two weeks of the experiment, we set traps

approximately 3 times per week per enclosure (Monday, Wednesday, and Friday) alternating

between the two enclosures so that voles in each enclosure were trapped three days over the

course of two weeks. Traps were set in the evening (2230-2300 h) and checked the following

morning at 0700 h. If we needed to put radiocollars on voles (see below), we also set traps from

1800-1900 h and then checked them the same evening from 2000-2100 h. Prairie vole gestation

and time to weaning are each approximately 21 days (Richmond & Conaway, 1969) and,

therefore, the offspring produced within the enclosures usually first emerge from the nest

approximately 5-6 weeks after the adults are released (N. Solomon, B. Keane, unpublished

data). We therefore stopped setting traps overnight from week 6-14 of the experiment to ensure

that offspring would not spend an entire night without their parents. During this time, we set

traps in the evening (2230-2300 h) and checked them approximately two hours later but did not

leave traps set overnight. Finally, we did not trap when there were severe thunderstorms and

made up this trapping session on a different day during the week, if possible. We supplemented

this regular grid trapping by placing additional traps at nest entrances after we located them

using VHF telemetry and/or UV powder tracking (Lemen and Freemen, 1985).

During each capture, we identified each individual using a handheld PIT tag reader,

recorded the location where the vole was live-trapped, the other individuals caught in the same

trap, body mass (g, with Pesola spring scale), and assessed reproductive condition. Males were

recorded as being either scrotal or non-scrotal depending on the development of the testes.

Females were recorded as non-reproductive, pregnant, or lactating depending on whether

developing fetuses could be felt or if nipples were pronounced. Once offspring were live trapped

for the first time, we gave them a unique toe clip until they were large enough (>25 g) to be PIT

tagged. Reproductive success of adult voles was estimated as the total number of offspring they

produced that we were able to trap.

Parentage analyses

Before adult voles were released into the enclosures, we collected a small piece of ear

tissue and stored it in 70% ethanol in a -80^0 C freezer. When offspring were trapped for the first

time, the tissue from the identifying toe clip was saved and temporarily stored in a -20^0 C freezer

until samples could be moved to a -80^0 F freezer. We extracted DNA using DNeasy Blood and

Tissue kits (Qiagen). We followed the manufacturer's protocol except that tissue samples were

incubated at 70^0 C, DNA was eluted in 100 μl of molecular grade water instead of 200 μl of buffer

AE, and DNA samples were incubated at room temperature for 5 minutes instead of 1 minute.

Once DNA was extracted, we genotyped the samples for all adults and offspring at seven

microsatellite loci (Keane et al., 2007). We then ran an allele frequency analysis on the population

of voles in each enclosure separately using Cervus 3.0.7 (see Mabry et al., 2011 for full details).

To determine parentage, we used Cervus 3.0.7 parentage analyses with known sexes, which

calculates a likelihood ratio for each potential mother and father in order to determine the most

likely biological parents in the population for each offspring (Mabry et al., 2011). We were able to

determine both parents (trio confidence level) with a 95% confidence level for 33/41 (80.5%)

offspring, so only these 33 offspring were included in the analyses of mating and reproductive

success.

Statistical Analyses

All statistical analyses were done in R version 3.4.1 (R Core Team, 2017).

Density and Body Mass

Population density was calculated based on the number of unique individuals caught

within each two-week period (over each two-week period both enclosures were trapped with

equal effort except for occasional cancellations due to weather). To investigate population

density over the course of the field season, we used a linear model with density (log-

transformed with base 10 to improve normality of residuals) with the fixed effects of enclosure

and weeks in the study and the interaction of these terms. Sex ratio was calculated by dividing

the number of adult males by the number of total adults for each two-week period. We used a

binomial generalized linear model to investigate sex ratio with the fixed effects of enclosure and

weeks in the study and the interaction of these terms.

Social Network Analyses

We conducted all social network analyses using the R package asnipe version 1.1.4

(Farine, 2017). In order to generate our social networks, we took the PIT tag readings from the

RFID antennas and ran them through a Gaussian Mixture Model with each day labeled

separately (Psorakis et al., 2012). This model goes through the raw data of the PIT tag readings

and creates groups based on when tag readings at the same antenna are clumped throughout

time. Therefore, there is not a uniform time period used to create these groups, they are based

on how our data were distributed over time. The duration of these group events ranged from 0

seconds (so voles were both at the antenna at the same time) to 66,161 seconds with an

average of 655.2 ± 3,352.8 seconds. This then creates a group by individual matrix where being

in the same spatial and temporal "group" counts as an association between individuals. As we

were only interested in the number of connections each individual had (not the strength of these

connections), we used a binary, unweighted measurement of degree where any non-zero

association was counted as a "1". Thus, anytime we refer to the number of social connections in

this paper, we calculated this using the unweighted degree. For more details about the

construction of the social networks see Sabol et al., 2018.

Reproductive Success Models

For all models including the number of mates (mating success) or the number of

offspring produced that survived to emergence from the natal nest (reproductive success) as the

response variable, we used Poisson generalized linear models. For each response variable we

ran two models using social network data, one including all social connections in order to

investigate sociality overall and one including only the opposite-sex social connections. These

models all had fixed effects of the number of social connections, the interaction of this with sex,

the number of social connections squared to assess non-linear effects of social connections on

mating success or reproductive success (one model including all connections and another for

each response variable including only opposite-sex connections), the interaction of this with sex,

enclosure, and survival (calculated as the proportion of the field season the individual survived

based on last detection). To test if mating success and reproductive success were related, we

ran a separate model with the number of offspring produced as the response variable and fixed

effects of the number of mates with which individuals produced offspring, the interaction of this

with sex, and enclosure. None of the GLMs were over-dispersed as all the dispersion

parameters were <1.

Body Mass

To investigate body mass, we calculated the average body mass for each male vole for

the entire field season (range 1-19 measurements, average 7.75 measurements). Females

were not included because we were using body mass as a proxy for body quality, and female

mass would be affected by both pregnancy status and body condition. We then used a general

linear model for average body mass with the number of social connections (one model including

all connections another including only opposite-sex connections), the number of social

connections squared (one model including all connections another including only opposite-sex

connections), enclosure, survival (calculated as the proportion of the field season the individual

survived based on last detection), and the number of mass measurements we had for each

individual. We visually assessed the distribution of the data and residuals for normality.

Randomized Models

For every model that included unweighted degree (the number of social connections),

we used the network permutation method in asnipe (Farine, 2017). This method is useful

because it helps control for the fact that social network data are not independent. This method

also allows us to investigate our hypotheses more specifically by allowing us to test if the

observed relationships are significantly different than random networks with the same structure

as our social networks (see Farine et al., 2015; Spiegel et al. 2017 for other similar uses of this

method). The network permutation method takes a piece of data from the group by individual

matrix and swaps it for a different individual (Farine, 2013). Specifically, we ran 10,000

randomized models where each time another piece of data from the individual by group matrix

was swapped. We also restricted swaps to only voles in the same enclosure that were recorded

on the RFID antennas during the same day to control for voles that did not survive the entire

season. Further, for the opposite-sex networks we restricted swaps to include only voles of the

same sex so that we were only comparing our opposite-sex network to other opposite-sex

networks, not all possible combinations. We then compared the b-value from the model for each

variable that includes a social network statistic to corresponding b-values from randomized

networks and calculated a new P-value based on the number of randomized models that

produced a b-value with a higher absolute value than the absolute value of the observed model.

Therefore, our P-value shows us whether the relationship we have observed is stronger than

the relationship from 10,000 randomizations of our dataset (Farine, 2013). We ran each set of

randomizations three times to ensure that the P-values were consistently significant. We

present all three P-values from these randomizations and, conservatively, only consider that a

relationship is statistically significant if all three randomizations revealed P-values <0.05.

Results

310   *Enclosure density & adult sex ratio*

The number of adult voles in each enclosure declined over the course of the field

season due to mortality (effect of weeks in the study, b = -0.14, t_{12} = -6.07, *P* < 0.0001, Fig. 1A,

note estimates on $\log_{10}$ scale) and the significant interaction between weeks in the study and

enclosure indicated that vole density decreased more strongly in the high density enclosure

than in the low density enclosure (enclosure x weeks: b = 0.083, t_{12} = 4.79, *P* = 0.00045). For

example, the starting density of voles in the high-density enclosure was 480 voles/ha (week 0 in

Fig. 1A) but was reduced to 200 voles/ha in the middle of the experiment (week 8 in Fig. 1A)

and to 60 voles/ha at the end of the experiment (week 14 in Fig. 1A). The starting density in the

low-density enclosure was 240 voles/ha (week 0 in Fig. 1A) but was reduced to 180 voles/ha in

the middle of the experiment (week 8 in Fig. 1A) and 100 voles/ha at the end of the experiment

(week 14 in Fig. 1A). In total, only 12.5% of voles in the high-density enclosure were still alive at

the end of the experiment whereas 41.7% of the voles in the low-density enclosure were still

alive at the end of the experiment.

Unlike density, adult sex ratio did not differ significantly during the course of the

experiment (effect of time, b = -0.04, z = -0.26, df = 12, *P* = 0.80, Fig. 1B) or between the two

enclosures (effect of enclosure, b = 0.15, z = 0.082, df = 12, *P* = 0.94, Fig. 1B).

Effects of sociality on mating success

Overall, both female and male voles that had an intermediate (i.e., the middle of the

range of observed values) number of social connections produced offspring with a greater

number of different mates (i.e., had higher mating success). In the model considering all social

interactions with same- and opposite-sex individuals, voles that had an intermediate number of

social connections with all possible individuals had higher mating success (effect of social

connections^2: b = -0.012, z = -0.88, *P*-values from randomized networks = 0.012, 0.0093,

<0.0001, Table 1, Fig. 2A) but this relationship was slightly different between the sexes (sex x

social connections^2: b = 0.014, z = 0.94, *P* = 0.010, 0.0018, <0.0001, Table 1, Fig. 2A).

However, in both females and males, those with an intermediate number of social connections

had the highest mating success, and the difference in the interaction with sex and the number of

social connections on mating success seemed to be largely due to males having slightly more

overall social connections than females (Fig. 2A).

The same relationship was true when only opposite-sex connections were considered.

Mating success was highest for female and male voles with an intermediate number of

opposite-sex social connections (social connections^2: b = -0.028, z = -1.01, P = 0.0011, 0.0035,

0.001, Fig. 2B), although the magnitude of this effect slightly differed between the sexes, though

this seems to be driven by the difference in the number of social connections between the sexes

with females tending to have more social connections than males (sex x social connections^2: b

= 0.057, z = 1.36, P = <0.0001, <0.0001, 0.0068, Table 1, Fig. 2B).

Overall, both female and male voles in the low-density enclosure had higher mating

success than individuals in the higher density enclosure (all social connections: b = 1.89, z =

4.55, P < 0.0001; opposite-sex social connections: b = 2.03, z = 4.72, P < 0.0001, Table 1).

Individuals that survived in the enclosures for longer had higher mating success (all social

connections: b = 2.07, z = 2.33, P = 0.020; opposite-sex social connections: b = 1.75, z = 2.09,

P = 0.037, Table 1).

353   *Effects of sociality on reproductive success*

Both female and male voles with an intermediate number of social connections produced

more offspring that survived to emergence from the natal nest (b = -0.0085, z = -0.77, P =

0.002, 0.035, 0.015, Table 2, Fig. 3A). This result did not consistently differ by sex (sex x social

connections^2: b = 0.0082, z = 0.63, P = 0.021, 0.11, 0.022, Table 2, Fig. 3A). However, when

only considering opposite-sex social connections, these relationships were not significant. The

number of offspring that voles produced was not related to the number of opposite-sex social

connections for males or females (b = 0.15, z = 0.52, P = 0.20, 0.53, 0.54, Table 2, Fig. 3B) and

voles with an intermediate number of opposite-sex social connections did not produce

significantly more offspring (b = -0.022, z = -0.93, *P* = 0.14, 0.099, 0.12, Table 2, Fig. 3B). This

relationship did not consistently vary with sex, although there is a tendency for a difference

between the sexes for opposite-sex connections[2] where the inverted u-shaped relationship was

slightly lessened in males and males tended to have fewer connections than females (sex x

number of opposite-sex connections: b = -0.39, z = -1.04, *P* = 0.047, 0.024, 0.11; sex x

opposite-sex connections[2]: b = 0.038, z-value = 0.92, *P* = 0.063, 0.010, 0.012, Table 2, Fig 3B).

Male and female voles that survived for longer produced more offspring (all social connections:

b = 2.49, z = 3.03, P = 0.0025; opposite-sex social connections: b = 2.19, z = 2.83, P = 0.0046,

Table 2).

371   Overall, there was a fitness advantage to being polygynous (mating with a greater

number of individuals) in both females and males. Individuals with more mates produced more

offspring that survived to emergence from the natal nest (b = 0.64, z = 4.88, *P* <0.0001, Table 3,

Fig. 4) and this relationship did not differ by sex as the interaction between number of mates

and sex was not significant (b = -0.093, z = -0.56, *P* = 0.58, Table 3). Individuals in the lower

density enclosure had higher reproductive success than individuals in the higher density

enclosure (b = 1.53, z = 3.37, *P* = 0.00076, Table 3).

378   *Effects of sociality on body mass*

379   Males with an intermediate number of connections had a higher average body mass

(effect of degree[2] squared: b = -0.066, t = -2.36, *P* = <0.0001, <0.0001, <0.0001, Table 4, Fig.

5A). This relationship was also significant when only opposite-sex connections were considered

(b = -0.20, t = -1.54, *P* = 0.0024, 0.0031, 0.0005, Table 4, Fig. 5B). Average body mass was not

significantly different between the two enclosures (all social connections: b = -1.03, t = -0.40, *P*

= 0.69; opposite-sex social connections: b = -0.16, t = -0.058, *P* = 0.95, Table 4). Survival did

not predict average body mass in either model (all social connections: b = 13.37, t = 1.52, P =

0.14; opposite-sex social connections: b = 10.80, t = 1.22, *P* = 0.23, Table 4). Although the

number of measurements we obtained to calculate each individual's average body mass varied

(Fig. 5), it did not affect our measure of average body mass in either model (all social

connections: b = -0.12, t = -0.20, *P* = 0.84; opposite-sex social connections: b = -0.20, t = -0.31,

P =0.76, Table 4).

Discussion

We examined the fitness consequences of variation in the frequency of social

interactions, as estimated using unweighted degree from social network analyses. As expected,

both female and male prairie voles varied in the frequency of social interactions, as reflected in

our social network analyses that quantified social network degree using the frequency of

temporal and spatial co-occurrence generated by our RFID system. Female and male voles with

an intermediate number of social interactions had the greatest mating success and produced

the greatest number of offspring, though the latter was only the case when we included all

interactions with other voles, and not when only opposite-sex connections were considered.

We investigated the number of connections each vole had with other conspecifics as our

measure of sociality (based on co-occurrence data). In social network analyses, this is

effectively the unweighted degree of an individual. We have previously shown that spatial and

temporal co-occurrence is correlated with vole behavior in a standardized partner preference

test and the degree of home range overlap between two voles (Sabol et al., 2018). Thus, while

this is a rough characterization of vole sociality, it seems to reflect other aspects of vole social or

mating behavior, such as home range use and overlap or behavioral interactions between two

voles, although it is important to note that our measure of sociality does not characterize the

strength, quality, or nature of these social associations. For example, voles that are socially

monogamous may have one strong social connection with their social partner regardless of

whether they are genetically monogamous or not, but non-genetically monogamous voles may

have more connections with other opposite-sex individuals. Furthermore, being connected does

not necessarily equate to an affiliative relationship, as some voles may have frequent temporal

and spatial co-occurrence that are agonistic in nature due to the necessity of defending

territories, mates, or offspring.

We tested our predictions about the fitness benefits of sociality using two different sets

of social network data, one including social connections with all voles and one only including

connections with opposite-sex individuals. This allowed us to investigate the potential costs and

benefits of sociality overall (e.g., social interactions of a female vole with other females and

males) as well as specifically social interactions between opposite-sex individuals, which may

be more directly relevant to mating and reproductive success. We found that voles with an

intermediate number of social connections with all voles (i.e., more social) had significantly

higher mating and reproductive success whereas voles with an intermediate number of

opposite-sex connections had significantly higher mating success but not higher reproductive

success. The process of running the randomizations was somewhat different for the two sets of

networks (all social connections or just opposite-sex connections), which could contribute to the

observed differences in the results between the two sets of networks. We limited permutations

for the opposite-sex network to within the same sex (where female social network data were

being swapped for another female and male data were being swapped for another male) so

these swaps could only be done between approximately half as many individuals each time. By

keeping the sex of each individual in the association consistent, the structure of the

randomizations was more like the data it was being compared to instead of comparing all

possible connections (Farine, 2017). However, since the permutations were done on the raw

data and then we pulled only the opposite-sex connections from these networks, it is possible

that some of the 10,000 permutations affected the same-sex connections in the raw data

(therefore changing a social interaction that is not included in data used for the linear model)

and therefore may not have changed the estimate for the relationship between opposite-sex

connections and the response variable every time, whereas in the models with all connections

included, a relevant social interaction would have changed every time, which would then change

the model estimate for the relationship between social connections and the response variable

some amount every time. Since we ran such a large number of these permutations, this may not

have affected the overall result of the tests, but it is a limitation of the method.

Using our measure of sociality, our results suggest the possibility of stabilizing selection

on sociality because voles that co-occurred spatially and temporally with very few or very many

conspecifics had the lowest mating success and, at least when considering all social

connections, had the lowest reproductive success. The results of McGuire et al. (2002) and

Solomon and Keane (2018) show that large social groups do not increase female reproductive

success. McGuire et al. (2002) showed that female prairie voles that lived in large groups had

fewer offspring survive to 12 or 30 days of age. Similarly, Solomon and Keane (2018) showed

that females did not benefit from living in large social groups in two other natural populations.

These studies are consistent with our results that a very large number of social connections

(which should occur in large social groups) does not increase reproductive success of breeding

females, and our results show that this is also true for males.

We only recorded the association between one measure of sociality and mating and

reproductive success in one year and the effects of the number of social connections on fitness

could be altered when environmental conditions change. For example, the Female Dispersion

Hypothesis would predict that if our measure of sociality reflects the socially monogamous

behavior of male prairie voles, our measure of sociality should be positively correlated with male

mating success when females are spatially clumped as males that have more social

connections with clumped females should have higher mating success (Shuster and Wade,

2003; Dobson et al., 2010; Lukas and Clutton-Brock, 2013). In natural populations of prairie

voles, population density is quite variable across years (Getz et al., 1993, 2001) and some

previous observational studies of prairie voles in field settings suggested that socially

monogamous behavior is more common at low densities (McGuire et al.,1990; Solomon et al.,

2009; but see Getz and McGuire,1993). There is also some evidence that resource distribution

may impact the mating strategy of prairie voles and this effect may be mediated through its

influence on density (Streatfeild et al., 2011). This suggests the possibility that selection on the

social behavior of prairie voles varies among years due to changes in population or female

density but additional multi-year studies measuring a broader array of social behaviors in free-

living voles are needed to test this prediction.

One possible explanation for an intermediate level of sociality being associated with the

highest mating success and potentially highest reproductive success is that this reflects a

tradeoff between devoting time to social interactions (although we do not know the type of social

interaction occurring) with other voles and time to other behaviors like foraging and parental

care. Although high levels of sociality can have beneficial effects on individual fitness, it may

also carry costs for an individual's health or physical condition (Nunn et al., 2015). Indeed, we

found that male voles with the most social connections had the lowest body mass, suggesting

that there may be a reduction in body condition associated with a very high level of sociality.

This could reflect the energetic costs associated with having many social connections or living in

a large group (e.g., Lutermann et al., 2013), or these could be agonistic interactions with males

on neighboring territories, resulting in males investing more time in territory defense than males

with fewer neighbors. Why males with very few social connections were also lighter in body

mass is not clear but these males may have been of lower phenotypic quality given that they

had few social connections, low body mass, and low mating and reproductive success.

Alternatively, having fewer social connections could result in a loss of body mass if these males

had no assistance in territory defense and thus, expended more energy than males with more

social connections (e.g., having a female social partner). Females likely face many of these

same tradeoffs, but as we did not test quality in females (due to changes in mass being linked to

pregnancy) and so further study is needed to investigate this relationship in females.

Our results suggest that it is not advantageous for voles to have social connections with

too many opposite-sex conspecifics. One possible explanation is that individuals with an

intermediate number of social connections may better balance the trade-off between the

number and quality or strength of social relationships. For example, individuals with the highest

social network degree may just have many weak social or agonistic connections, which may not

result in more matings or increased reproductive success. Individuals with an intermediate

social network degree may have more affiliative social connections that are strong enough to

result in matings than voles at either extreme. This is supported by studies of the association

between the strength of social connections and fitness in cercopithecine primates (baboons)

where females with strong social bonds with other females in their group have higher offspring

survival (Silk et al., 2003, 2009) or longevity (Silk et al., 2010). Similar relationships between the

strength of social bonds and fitness have also been found in male primates; male Assamese

macaques (*Macaca assamensis*) with strong social bonds to other males (including unrelated

males) sired more offspring than those with fewer strong bonds with other males (Schulke et al.,

2010). As the number of social connections increase, the strength of association of each of

these social connections may decline (Whitehead 2008), thus prairie voles may be constrained

by the number of social connections in which they can invest enough time to result in successful

mating or rearing of offspring given that prairie voles exhibit biparental care. Individuals that can

best balance this trade-off between the number and strength of social connections may have

the highest mating and reproductive success.

It is of course likely that the fitness benefits of the quantity versus quality of social

connections may vary according to whether the modal social structure of the species is group-

living (such as primate species mentioned above where strong social bonds increase fitness) or

its mating system. For example, Ryder et al. (2009) found a positive association between the

number of social connections (social network degree) and number of offspring sired in male

manakins. As this is a lekking species, coordinated male displays may make male-male

connections a more important factor for mating success than in prairie voles. Additionally, the

short-term coalitions at leks may make the strength of the relationship less important than in

species like prairie voles. Studies like these are rare and so future studies across a broader

array of species with different mating systems will be needed to fully characterize the

relationship between the number and strength of social connections and measures of fitness.

Doing so will help provide insight into how individuals within a species balance the fitness costs

and benefits of social behavior, thereby providing a complementary approach to comparative

studies regarding the evolution of social behavior.

Acknowledgements

We thank the Ecology Research Center (ERC) at Miami University for permitting us to conduct

this study and Jeremy Fruth, field manager at the ERC, for coordinating the project. We also

thank James Lichter, Sage Sparks, Caleb Collins, Tony Leon, Meghan Charles, and Staci

Burrow for assisting on the project. We also thank the American Society of Mammalogists

(Graint-in-Aid of Research to A. Sabol) and University of Michigan (B. Dantzer) for funding.

Data Statement

All raw data and R code will be made available under CC-BY from FigShare upon acceptance.

Literature Cited

Alexander, R. D. (1974). The evolution of social behaviour. *Annual Review of Ecology, Evolution, and Systematics, 5,* 325-383.

Armitage, K. B. (1998). Reproductive strategies of yellow-bellied marmots: Energy conservation and differences between the sexes. *Journal of Mammalogy, 79*, 385–393.

Clutton-Brock, T. H. (1989). Review lecture: mammalian mating systems. *Proceedings of the Royal Society B, 236,* 339-372.

Clutton-Brock, T.H., Brotherton, P.N.M., Smith, R., McIllrath, G.M., Kansky, R., Gaynor, D., O'rian, M.J., Skinner, J.D. (1998). Infanticide and expulsion of females in a cooperative mammal. *Proceedings of the Royal Society B, 265,* 2291-2295.

Côté, I.M., & Poulinb R. (1995). Parasitism and group size in social animals: a meta-analysis. *Behavioral Ecology, 6*, 159-165.

Creel, S., Dantzer, B., Goymann, W. & Rubenstein, D.R. (2013). The ecology of stress: effects of the social environment. *Functional Ecology, 27,* 66–80.

Dantzer, B., Goncalves, I.B., Spence-Jones, H.C., Bennett, N.C., Heistermann, M., Ganswindt, A., Dubuc, C., Gaynor, D., Manser, M.B., Clutton-Brock, T.H. (2017). The influence of stress hormones and aggression on cooperative behaviour in subordinate meerkats. *Proceedings of the Royal Society B, 284,* 20171248.

Dobson, F. S., Way, B. M., & Baudoin C. (2010). Spatial dynamics and the evolution of social monogamy in mammals. *Behavioral Ecology, 21,* 747–752. doi:10.1093/beheco/arq048

Donaldson, Z. R., & Young, L. J. (2008). Oxytocin, vasopressin, and the neurogenetics of sociality. *Science, 322,* 900-904. doi:10.1126/science.1158668

Eisenberg, J.F., Muckenhirn, N.A., Rudran, R. (1972). The relation between ecology and social structure in primates. *Science, 176*, 863-874.

Emlen, S.T. (1984). *Cooperative breeding in birds and mammals. Behavioural Ecology: An Evolutionary Approach*. J.R. Krebs and N.B. Davies (Eds.), pp. 305-339, Blackwell, Oxford.

Emlen, S.T. (1994). Benefits, constraints and the evolution of the family. *Trends in Ecology and Evolution, 9,* 282-285.

Farine, D. R. (2013). Animal social network inference and permutations for ecologists in R using *asnipe. Methods in Ecology and Evolution, 4,* 1187-1194. doi:10.1111/2041-210X.12121

Farine, D. R., Firth, J. A., Aplin, L. M., Crates, R. A., Culina, A., Garroway, C.J., Hinde, C. A., Kidd, L. R., Milligan, N. D., Psorakis, I., Radersma, R., Verhelst, B., Voelkl, B., & Sheldon, B. C. (2015). The role of social and ecological processes in structuring animal populations: a case

584   study from automated tracking of wild birds. *Royal Society Open Science, 2.*
585   doi:10.1098/rsos.150057
586
587   Farine, D. R. (2017a). A guide to null models for animal social network analysis. *Methods in*
588   *Ecology and Evolution, 8,* 1309-1320. doi:10.1111/2041-210X.12772
589
590   Farine, D. R. (2017b). *asnipe: Animal Social Network Inference and Permutations for*
591   *E*cologists. R package version 1.1.4. Available online at: https://CRAN.R-
592   project.org/package=asnipe
593
594   Faulkes, C. G., Bennett, N. C., Bruford, M. W., O'brien H. P., Aguilar G. H., & Jarvis, J. U. M.
595   (1997). Ecological constraints drive social evolution in the African mole–rats. *Proceedings of the*
596   *Royal Society of London B: Biological Sciences, 267*. doi:10.1098/rspb.1997.0226
597
598   Getz, L. L., & Hofmann, J. E. (1986). Social organization in free-living prairie voles, *Microtus*
599   *ochrogaster. Behavioral Ecology and Sociobiology, 18*, 275-282.
600
601   Getz, L. L., Hofmann, J. E., McGuire, B., & Dolan, T. W. III. (2001). Twenty-five years of
602   population fluctuations of *Microtus ochrogaster* and *M. pennsylvanicus* in three habitats in east-
603   central Illinois. *Journal of Mammalogy, 82*, 22–34.
604
605   Getz, L. L., & McGuire, B. (1993). A comparison of living singly and in male-female pairs in the
606   prairie vole, *Microtus ochrogaster. Ethology, 94,* 265-278. doi:10.1111/j.1439-
607   0310.1993.tb00444.x
608
609   Getz, L. L., McGuire, B., Pizzuto, T., Hofmann, J. E., & Frase, B. (1993). Social organization of
610   the prairie vole (*Microtus ochrogaster*). *Journal of Mammalogy. 74,* 44–58. doi:
611   10.2307/1381904
612
613   Getz, L. L., McGuire, B., Hofmann, J., Pizzuto, T., & Frase, B. (1994). Natal dispersal and
614   philopatry in prairie voles (*Microtus ochrogaster*): settlement, survival, and potential
615   reproductive success. *Ethology Ecology & Evolution, 6,* 267-284.
616
617   Hatchwell, B. J., & Komdeur, J. (2000). Ecological constraints, life history traits and the
618   evolution of cooperative breeding. *Animal Behaviour, 59,* 1079–1086.
619   doi:10.1006/anbe.2000.1394
620
621   Jetz, W., & Rubenstein, D. R. (2011). Environmental uncertainty and the global biogeography of
622   cooperative breeding in birds. *Current Biology, 21,* 72-78. doi:10.1016/j.cub.2010.11.075.
623
Kappeler, P. M., Cremer, S., & Nunn, C. L. (2015). Sociality and health: impacts of sociality on
disease susceptibility and transmission in animal and human societies. *Philosophical*
Transactions of the Royal Society B: Biological Sciences, 370. doi:10.1098/rstb.2014.0116
Keane, B., Bryant, L., Goyal, U., Williams, S., Kortering, S. L., Lucia, K. E., Richmond, A. R.,
& Solomon, N. G. (2007). No effect of body condition at weaning on survival and reproduction
in prairie voles. *Canadian Journal of Zoology, 85,* 718-727.

Krause, J., & Ruxton, G. D. (2002). *Living in groups*. Oxford: Oxford University Press.
630
631   Lambert, C. T., Sabol, A. C., & Solomon, N. G. (2018). Genetic monogamy in socially
632   monogamous mammals is primarily predicted by multiple life history factors: a meta-analysis.
633   *Frontiers in Ecology and Evolution, 6,* 139. doi:10.3389/fevo.2018.00139
634
635   Lemen, C. A., & Freeman, P. W. (1985). Tracking mammals with fluorescent pigments: a new
636   technique. *Journal of Mammalogy, 66,* 134–136.
637
638   Lott, D. F. (1991). *Intraspecific variation in the social systems of wild vertebrates.* Cambridge:
639   Cambridge University Press.
640
641   Lukas, D., & Clutton-Brock, T. H. (2013). The evolution of social monogamy in mammals.
642   *Science, 341,* 526 -530. doi:10.1126/science.1238677
643
644   Lukas, D., & Clutton-Brock, T. H. (2017). Climate and the distribution of cooperative breeding
645   in mammals. *Royal Society Open Science, 4*. doi:10.1098/rsos.160897
646
647   Lukas, D., & Huchard, E. (2014). The evolution of infanticide by males in mammalian societies.
648   *Science, 346,* 841-844. doi:0.1126/science.1257226
649
650   Lutermann, H., Bennett, N. C., Speakman, J. R., & Scantlebury, M. (2013). Energetic benefits of
651   sociality offset the costs of parasitism in a cooperative mammal. *PLOS ONE*, *8*, e57969.
652   doi:10.1371/journal.pone.0057969
653
654   Mabry, K. E., Streatfeild, C. A., Keane, B., & Solomon, N. G. (2011). avpr1a length
655   polymorphism is not associated with either social or genetic monogamy in free-living prairie
656   voles. *Animal Behaviour, 81*, 11-18.
657
658   McGuire, B., & Getz, L.L. (2010). Alternative male reproductive tactics in a natural population
659   of prairie voles, *Microtus ochrogaster*. *Acta Theriol, 55,* 261-270. doi:10.4098/j.at.0001-
660   7051.077.2009
661
662   McGuire, B., Getz, L. L., & Oli, M. K. (2002). Fitness consequences of sociality in prairie voles,
663   Microtus ochrogaster: influence of group size and composition. *Animal Behaviour, 64,* 645-654.
664   doi:10.1006/anbe.2002.3094.
665
666   McGuire, B., Pizzuto, T., Getz, L.L. (1990). Potential for social interaction in a natural
667   population of prairie voles (*Microtus ochrogaster*). *Canadian Journal of Zoology, 68,* 391-398.
668
669   Nunn, C.L., Jordán, F., McCabe, C.M., Verdolin, J.L., Fewell, J.H. (2015). Infectious disease
670   and group size: more than just a numbers game. *Philosophical Transactions of the Royal Society
671   B, 370,* 20140111.
672
673   Okhovat, M., Berrio, A., Wallace, G., Ophir, A. G., & Phelps, S. M. (2015). Sexual fidelity
674   trade-offs promote regulatory variation in the prairie vole brain. *Science, 350,* 1371-1374.

676 Ophir, A.G., Wolff, J.O., Phelps, S.M. (2008). Variation in neural V1aR predicts sexual fidelity
677 and space use among male prairie voles in semi-natural settings. *PNAS, 105*, 1249-1254.
678

679 Psorakis, I., Roberts, S. J., Rezek, I., & Sheldon, B. C. (2012). Inferring social network structure
680 in ecological systems from spatio-temporal data streams. *Journal of the Royal Society Interface,*
681 *9,* 3055–3066. doi: 10.1098/rsif.2012.0223
682

683 R Core Team (2017). *R: A Language and Environment for Statistical Computing.* Vienna: R
684 Foundation for Statistical Computing. Available online at: https://www.R-project.org/.
685

686 Richmond, M., & Conaway, C. H. (1969). Management, breeding, and reproductive performance
687 of the vole, *Microtus ochrogaster*, in a laboratory colony. *Laboratory Animal Care, 19,* 80-87.
688

689 Rubenstein, D. R., & Lovette, I. J. (2007) Temporal environmental variability drives the
690 evolution of cooperative breeding in birds. *Current Biology, 17,* 1414-1419.
691 doi:10.1016/j.cub.2007.07.032.
692

693 Ryder, T.B., Parker, P.G., Blake, J.G., & Loiselle, B.A. (2009). It takes two to tango:
694 reproductive skew and social correlates of male mating success in a lek-breeding bird.
695 *Proceedings of the Royal Society B: Biological Sciences, 276,* 2377-2384.
696 doi:10.1098/rspb.2009.0208
697

698 Sabol, A. C., Solomon, N. G., & Dantzer, B. (2018). How to study socially monogamous
699 behavior in secretive animals? Using social network analyses and automated tracking systems to
700 study the social behavior of prairie voles. *Frontiers in Ecology and Evolution, 6,* 178.
701 doi:10.3389/fevo.2018.00178
702

703 Schradin, C., & Pillay, N. (2005). Intraspecific variation in the spatial and social organization of
704 the African striped mouse, *Journal of Mammalogy, 86*, 99–107.
705

706 Schradin, C., König, B., Pillay, N. (2010). Reproductive competition favours solitary living
707 while ecological constraints impose group-living in African striped mice. *Journal of Animal*
708 *Ecology, 73,* 515-521.
709

710 Schülke, O., Bhagavatula, J., Vigilant, L., & Ostner, J. (2010). Social bonds enhance
711 reproductive success in male macaques. *Current Biology, 20,* 2207-2210.
712 doi:10.1016/j.cub.2010.10.058.
713

714 Shen, S-F., Emlen, S.T., Koenig, W.D., Rubenstein, D.R. (2017). The ecology of cooperative
715 breeding behaviour. *Ecology Letters, 20,* 708-720.
716

717 Shuster, S. M., & Wade, M. J. (2003). *Mating systems and mating strategies*. Princeton:
718 Princeton University Press.
719

Shuster, S. M., Willen, R. M., Keane, B., & Solomon, N. G. (2019). Alternative mating tactics in
socially monogamous prairie voles, Microtus ochrogaster. *Frontiers in Ecology and Evolution*, 7.
doi:10.3389/fevo.2019.00007

Silk, J. B. (2007). The adaptive value of sociality in mammalian groups. *Philosophical*
Transactions of the Royal Society B: Biological Sciences, *362*, 539–559.
doi:10.1098/rstb.2006.1994

Silk, J. B., Alberts, S. C., & Altmann, J. (2003). Social bonds of female baboons enhance infant
survival. *Science*, *302*, 1231-1234. doi:10.1126/science.1088580

Silk, J.B., Beehner, J.C., Bergman, T.J., Crockford, C., Engh, A.L., Moscovice, L.R., Wittig,
R.M., Seyfarth, R.M., Cheney, D.L. (2009). The benefits of social capital: close social bonds
among female baboons enhance offspring survival. *Proceedings of the Royal Society B, 276,*
3099-3104.

Silk, J.B., Beehner, J.C., Bergman, T.J., Crockford, C., Engh, A.L., Moscovice, L.R., Wittig,
R.M., Seyfarth, R.M., Cheney, D.L. (2010). Strong and consistent social bonds enhance the
longevity of female baboons. *Current Biology, 20,* 1359-1361.

Solomon, N. G. (1993). Comparison of parental behavior in male and female prairie voles
(*Microtus ochrogaster*). *Canadian Journal of Zoology, 71,* 1991–1994.

Solomon, N. G., & Keane, B. (2018). Dispatches from the field: sociality and reproductive
success in prairie voles. *Animal Behaviour*, *143*, 193–203. doi:10.1016/j.anbehav.2018.07.001

Solomon, N. G., & Jacquot, J. J. (2002). Characteristics of resident and wandering prairie voles,
Microtus ochrogaster. *Canadian Journal of Zoology, 80*, 951-955.

Solomon, N. G., Keane, B., Knoch, L. R., & Hogan, P. J. (2004). Multiple paternity in socially
monogamous prairie voles (Microtus ochrogaster). *Canadian Journal of Zoology, 82,* 1667–
1671.

Solomon, N.G., Richmond, A. R., Harding, P. A., Fries, A., Jacquemin, S., Schaefer, R. L.,
Lucia, K. E., & Keane, B. (2009). Polymorphism at the avpr1a locus in male prairie voles
correlated with genetic but not social monogamy in field populations. *Molecular Ecology, 18,*
4680-4695.

Spiegel, O., Sih, A., Leu, S. T., & Bull, C. M. (2017). Where should we meet? Mapping social
network interactions of sleepy lizards shows sex-dependent social network structure. *Animal*
Behaviour, 136, 207-215. doi:10.1016/j.anbehav.2017.11.001.

Streatfeild, C. A., Mabry, K. E., Keane, B., Crist, T. O., & Solomon, N. G. (2011). Intraspecific
variability in the social and genetic mating systems of prairie voles, *Microtus ochrogaster*.
Animal Behaviour, 82, 1387–1398. doi: 10.1016/j.anbehav.2011.09.023
765

766   van Schaik, C. (1983). Why are diurnal primates living in groups? *Behaviour, 87*, 120-144.
767
768   Walum, H., & Young, L. J. (2018). The neural mechanisms and circuitry of the pair bond.
769   *Nature Reviews Neuroscience*, *19*, 643–654. doi:10.1038/s41583-018-0072-6
770
771   Whitehead, H. (2008). *Analyzing animal societies: quantitative methods for vertebrate social*
772   *analysis*. Chicago: University of Chicago Press.
773
774   Whiteman, N. K., & Parker, P. G. (2004). Effects of host sociality on ectoparasite population
775   biology. *Journal of Parasitology, 90*, 939-947. doi:10.1645/GE-310R
776
777   Young, L. J., & Wang, Z. (2004). The neurobiology of pair bonding. *Nature Neuroscience*, *7*,
778   1048–1054. doi:10.1038/nn1327
779

          Figures

Figure 1. A) The number of females, males, and offspring in each of our two enclosures over

time during the study based on the number of unique individuals live-trapped during each two-

week period of the field season. Note that the area of enclosures is equal, so the number of

voles in each enclosure can be used to compare relative density between the two, B) The sex

ratio in each enclosure, calculated as the proportion of total adult voles trapped during each

two-week period of the field season that were males, over time of the study.

791

792 **Figure 2**. Both females and males with an intermediate number of social connections with A) all

793 voles in their enclosure (both same- or opposite-sex individuals) or B) only opposite-sex

794 individuals in their enclosure had higher mating success (defined as the number of different

795 individuals with which they produced offspring). Points are jittered with males shown in blue and

796 females in red. Full statistical results shown in Table 1.

Figure 3. A) Both females and males with an intermediate number of social connections with all

voles in their enclosure (both same- and opposite-sex individuals) produced more offspring that

survived to emergence from the natal nest. B) Female and male voles with an intermediate

number of social connections with only opposite-sex individuals in their enclosure also tended to

produce more offspring. Points are jittered with red points being females and blue points being

males. Full statistical results shown in Table 2.

808   **Figure 4**. Both female and male prairie voles that had higher mating success (produced

809   offspring with a greater number of different mates) produced a great number of offspring that

810   survived to emergence from the natal nest. Points for females and males are jittered. Full results

811   shown in Table 3.

Figure 5. Male voles with an intermediate number of social connections with A) both female and male voles in their enclosure or B) just female voles were significantly heavier over the course of this study. Body mass for males was averaged for the entire duration of this study. The number of times we measured body mass ("N") varied among males so the size of each point is scaled based on the number of recorded mass measurements we have for each individual. Full results shown in Table 3.

Table 1. Effects of the number of social connections on vole mating success (number of

different individuals a vole produced offspring with). Degree refers to the number of social

connections for each individual either with all voles or opposite-sex voles. Note that

relationships involving social network data have three *P*-values because those b values were

compared to those from randomized networks three times to determine if they were consistently

significant (see methods for details). Survival refers to the proportion of days the vole was in the

enclosure based on when it was last recorded. "LD" is low-density enclosure.

Social Network	Variable	Estimate	SE	z-value	P-value from GLM	P-values from Randomization Tests
All individuals	**Intercept**	**-2.66**	**0.65**	**-4.06**	**<0.0001**	NA
	Sex (M)	0.68	0.92	0.73	0.46	NA
	Enclosure (LD)	**1.89**	**0.41**	**4.55**	**<0.0001**	NA
	Survival	**2.07**	**0.89**	**2.33**	**0.020**	NA
	Degree	0.083	0.21	0.40	0.69	0.12, 0.0098, 0.14
	Degree^2	**-0.012**	**0.013**	**-0.88**	**0.38**	**0.012, 0.0093, <0.0001**
	Sex (M) x Degree	**-0.21**	**0.23**	**-0.90**	**0.37**	**0.021, 0.0017, <0.0001**
	Sex (M) x Degree^2	**0.014**	**0.015**	**0.94**	**0.35**	**0.01, 0.0018, <0.0001**
Opposite-sex individuals	**Intercept**	**-2.81**	**0.70**	**-4.08**	**<0.0001**	NA
	Sex (M)	0.78	0.91	0.86	0.39	NA
	Enclosure (LD)	**2.03**	**0.43**	**4.72**	**<0.0001**	NA
	Survival	**1.75**	**0.84**	**2.09**	**0.037**	NA
	Degree	0.19	0.31	0.63	0.53	0.18, 0.19, <0.0001
	Degree^2	**-0.028**	**0.028**	**-1.01**	**0.31**	**0.0011, 0.0035, 0.001**
	Sex (M) x Degree	**-0.53**	**0.38**	**-1.42**	**0.15**	**<0.0001,<0.0001, 0.0046**
	Sex (M) x Degree^2	**0.057**	**0.042**	**1.36**	**0.17**	**<0.0001, <0.0001, 0.0068**

Table 2. Effects of the number of social connections on vole reproductive success (number of

offspring produced that survived until emergence from the natal nest). Degree refers to the

number of social connections for each individual either with all voles or opposite-sex voles. Note

that relationships involving social network data have three *P*-values because those b values

were compared to those from randomized networks three times to determine if they were

consistently significant (see methods for details). Survival refers to the proportion of days the

vole was in the enclosure based on when it was last recorded. "LD" is low-density enclosure.

Social Network	Variable	Estimate	SE	z-value	P-value from GLM	P-values from Randomization Tests
All individuals	**Intercept**	**-2.90**	**0.66**	**-4.14**	**<0.0001**	NA
	Sex (M)	0.61	0.93	0.65	0.52	NA
	Enclosure (LD)	**2.10**	**0.40**	**5.24**	**<0.0001**	NA
	Survival	**2.49**	**0.82**	**3.03**	**0.0025**	NA
	Degree	0.053	0.19	0.28	0.78	0.33, 0.25, 0.19
	Degree^2	**-0.0085**	**0.011**	**-0.77**	**0.44**	**0.002, 0.035, 0.015**
	Sex (M) x Degree	-0.14	0.21	-0.64	0.52	0.12, 0.18, 0.12
	Sex (M) x Degree^2	0.0082	0.013	0.63	0.53	0.021, 0.11, 0.022
Opposite-sex individuals	**Intercept**	**-2.97**	**0.67**	**-4.41**	**<0.0001**	NA
	Sex (M)	0.58	0.91	0.64	0.53	NA
	Enclosure (LD)	**2.21**	**0.41**	**5.40**	**<0.0001**	NA
	Survival	**2.19**	**0.77**	**2.83**	**0.0046**	NA
	Degree	0.15	0.28	0.52	0.60	0.20, 0.53, 0.54
	Degree^2	-0.022	0.024	-0.93	0.35	0.14, 0.099, 0.12
	Sex (M) x Degree	-0.39	0.38	-1.04	0.30	0.047, 0.024, 0.11
	Sex (M) x Degree^2	0.038	0.041	0.92	0.36	0.063, 0.010, 0.02

839 **Table 3.** Overall effects of vole mating success (number of different individuals with which a

840 vole produced offspring) on reproductive success (number of offspring produced that survived

841 until emergence from the natal nest). Since this model did not include social network data we

842 did not perform the randomizations to generate three *P*-values, we present the *P*-values from

843 the GLM. "LD" is low-density enclosure.

Variable	Estimate	SE	z-value	P-value from GLM
Intercept	**-2.13**	**0.44**	**-4.83**	**<0.0001**
Number of Mates	**0.64**	**0.13**	**4.88**	**<0.0001**
Sex (M)	0.26	0.49	0.53	0.60
Enclosure (LD)	**1.53**	**0.45**	**3.37**	**0.00076**
# of Mates x Sex (M)	-0.093	0.17	-0.56	0.58

Table 4. Effects of the number of social connections on average body mass for males. Degree

refers to the number of social connections for each individual either with all voles or opposite-

sex voles. Relationships involving social network data have three *P*-values because those b

values were compared to those from randomized networks three times to determine if they were

consistently significant (see methods). "# of measures" refers to number of times we measured

body mass, which were used to generate average mass for each individual. Survival refers to

proportion of days the vole was in the enclosure. "LD" is low-density enclosure.

Social Network	Variable	Estimate	SE	t-value	P-value from LM	P-values from Randomization Tests
All individuals	**Intercept**	**43.42**	**2.70**	**16.10**	**<0.0001**	NA
	Enclosure (LD)	-1.03	2.57	-0.40	0.69	NA
	Survival	13.37	8.78	1.52	0.14	NA
	# of measures	-0.12	0.60	-0.20	0.84	NA
	Degree	**0.95**	**0.56**	**1.70**	**0.099**	**<0.0001, 0.0001, <0.0001**
	Degree^2	**-0.066**	**0.028**	**-2.36**	**0.025**	**<0.0001, <0.0001, <0.0001**
Opposite-sex individuals	**Intercept**	**44.31**	**2.81**	**15.78**	**<0.0001**	NA
	Enclosure (LD)	-0.16	2.80	-0.058	0.95	NA
	Survival	10.80	8.89	1.22	0.23	NA
	# of measures	-0.20	0.63	-0.31	0.76	NA
	Degree	**1.54**	**1.22**	**1.26**	**0.22**	**0.0035, 0.0036, 0.0332**
	Degree^2	**-0.20**	**0.13**	**-1.54**	**0.13**	**0.0024, 0.0031, 0.0005**

853
854

Supplementary Material

870   **Figure S1.** Layout of the two RFID arrays in the enclosures. The RFID system was kept at each

871   array in each enclosure for three days in the order: array 1 enclosure 1, array 2 enclosure 1,

872   array 1 enclosure 2, and array 2 enclosure 2 and then repeated for the duration of the field

873   season.