Systematic perturbation of yeast essential genes using base editing

Philippe C Després[1,2,3], Alexandre K Dubé[1,2,3,4], Motoaki Seki[5], Nozomu Yachie*[5,6,7] and Christian R Landry*[1,2,3,4]

1. Département de Biochimie, Microbiologie et Bio-informatique, Faculté de sciences et génie, Université Laval, Québec, Québec, G1V 0A6, Canada

2. PROTEO, le regroupement québécois de recherche sur la fonction, l'ingénierie et les applications des protéines, Université Laval, Québec, Québec, G1V 0A6, Canada

3. Centre de Recherche en Données Massives (CRDM), Université Laval, Québec, Québec, G1V 0A6, Canada

4. Département de Biologie, Faculté de sciences et Génie, Université Laval, Québec, Québec, G1V 0A6, Canada

5. Research Center for Advanced Science and Technology, Synthetic Biology Division, University of Tokyo, Tokyo, 4-6-1 Komaba, Meguro-ku, 153-8904, Japan

6. Department of Biological Sciences, Graduate School of Science, the University of Tokyo, Tokyo, Japan

7. Institute for Advanced Biosciences, Keio University, Tsuruoka, Japan

*To whom correspondence should be addressed. CRL: Tel: 1-418-656-3954, Fax 1-418-656-7176, christian.landry@bio.ulaval.ca NY: Tel +81-3-5452-5242 (x55242), Fax +81-3-5452-5241 (x55241), yachie@synbiol.rcast.u-tokyo.ac.jp

Abstract

Base editors engineered from an inactivated Cas9 and a DNA editing enzyme offer an

unprecedented opportunity for the precise modification of genes, but have yet to be used at a

genome-scale throughput. Here, we test the ability of an editor based on a cytidine deaminase,

the Target-AID base editor, to systematically modify genes genome-wide using the set of yeast

essential genes. We tested the effect of mutating around 17,000 sites across more than 1,500

genes in a single experiment. We identified over 1,100 sites where mutations have a significant

impact on fitness. Using previously determined and preferred Target-AID mutational outcomes,

we predicted the protein variants caused by each of these gRNAs. We found that gRNAs with

significant effects on fitness are enriched in variants predicted to be deleterious by independent

methods based on site conservation and predicted protein destabilization. Finally, we identify key

features to design effective gRNAs in the context of base editing. Our results show that base

editing is a powerful tool to identify key amino acid residues at the scale of proteomes.

Introduction

Recent technical advances have allowed the investigation of the genotype-phenotype map at high resolution by experimentally measuring the effect of all possible nucleotide substitutions in a DNA sequence. While saturated mutagenesis informs us on the effect of many mutations, it usually covers a single locus or a fraction of it (Fowler and Fields 2014; Gray *et al.* 2018). Because such data is only available at sufficient coverage for a very small number of proteins, general rules on substitution effects must be extrapolated to other, often unrelated proteins. At a lower level of resolution, genome-scale mutations data has mostly been acquired through large-scale loss-of-function strain collections, where the same genetic change (for example, complete gene deletion) is applied to all genes (Winzeler *et al.* 1999; Giaever *et al.* 2002; C. elegans Deletion Mutant Consortium 2012). This approach is a powerful way to isolate each gene's contribution to a phenotype, including fitness, but limits our understanding of the role of specific positions within a locus.

CRISPR-Cas9 based approaches usually cause protein loss of function through indel formation (Shalem *et al.* 2014) or by modifying gene expression levels (Qi *et al.* 2013; Sander and Joung 2014; Smith *et al.* 2016) at many loci in parallel. Again, these approaches generally limit the information gain to one perturbation per locus. There is therefore a strong tradeoff between the resolution of the existing assays and the number of loci or genes investigated. Recent developments in the field now allow for the exploration of the effects of many mutations per gene across the genome. For instance, in yeast, methods for high throughput strain library construction have allowed the measurement of thousands of variant fitness effects in parallel across the genome (Sharon *et al.* 2018; Bao *et al.* 2018; Roy *et al.* 2018). These approaches rely on CRISPR-Cas9 based genome modifications requiring the formation of double-strand breaks followed by repair using donor DNA, which often relies on complex strain and plasmid

constructions. An alternative approach would be to use base editors, which allow the introduction

of the mutations of interest directly in the genome.

Base editors use DNA modifying enzymes fused to modified Cas9 or Cas12 proteins to create

specific point mutations in a target genome (Nishida *et al.* 2016; Gaudelli *et al.* 2017; reviewed in

Rees and Liu 2018). Such base editors have recently been used to perform site-specific forward

mutagenesis in human cell lines. The two main approaches, Targeted AID-mediated mutagenesis

(TAM) (Ma *et al.* 2016) and CRISPR-X (Hess *et al.* 2016), target specific regions of the genome

where they induce mutations randomly. This generates a library of mutant genotypes that can be

competed to find beneficial and deleterious variants under selective pressure. As the relative

fitness measurements depend on targeted sequencing of the locus of interest, these approaches

are difficult to adapt to high throughput multiplexed screens where tens of thousands of sites can

be targeted within the same gRNA libraries.

Here, we present a method that bridges the flexibility of Target-AID mutagenesis and the

multiplexing capacities of genome editing depletion screens. By using a base editor with a narrow

and well-defined activity window (Nishida *et al.* 2016), we selected gRNAs generating a limited

number of predictable edits in yeast essential genes. This allowed us to use gRNAs as a readout

for the effect of the mutations, similar to commonly used barcode-sequencing approaches to

measure fitness effects.

Results

Large-scale base editing screening

We used Target-AID mutagenesis to simultaneously assess mutational effects at over 17,000

putative sites in the yeast genome. We scanned yeast essential genes for sites amenable to

editing by the Target-AID base editor as well as targets with other specific properties, including

intronic sequences (Figure 1A, Figure S1). Because all essential genes have the same fitness

effects when deleted (Giaever *et al.* 2002), focusing on these genes allowed to limit the variation

in fitness that could be due to the relative importance of individual genes for growth rather than to

the importance of specific positions.

To ensure we could predict gRNA mutational outcomes with accuracy, we included in the library

only gRNAs with one to two nucleotides with a high probability of being edited based on the known

activity window of Target-AID in yeast (Nishida *et al.* 2016). We could then predict mutagenesis

outcomes for gRNAs computationally. We took into account that Target-AID is known to produce

both C-to-G and C-to-T mutations in yeast, with a 1.5 to 2 fold preference for C-to-G (Nishida *et*

al. 2016; Després *et al.* 2018). We also extended the analysis to include other point mutants at

possible secondary editing sites within the activity window (see methods). As such, we could

associate most gRNAs targeting protein-coding DNA to a primary C-to-G and C-to-T outcome (C-

to-G #1 and C-to-T #1), as well as to possible secondary outcomes if applicable (C-to-G #2 and

C-to-T #2). We also filtered out gRNAs that did not target between the 0.5th and 75th percentile

of the length of annotated genes to limit position biases that could influence the efficiency of stop-

codon generating guides (Doench *et al.* 2014; Michel *et al.* 2017).

The gRNA library was cloned into a high-throughput co-selection base editing vector (Després *et*

al. 2018). We performed pooled mutagenesis followed by bulk competition (Figure S2) to identify

mutations with significant fitness effects. As the relative abundance of each gRNA in the extracted

plasmid pool depends on the abundance of the subpopulation of cells bearing them, any fitness

effect caused by the mutation they induce will influence their relative abundance. Variation in

plasmid abundance was measured using targeted next-generation sequencing of the variable

gRNA locus on the base editing vector in a manner similar to GeCKO approaches (Sanjana *et al.*

2014; Shalem *et al.* 2014).

Figure 1. High-throughput forward mutagenesis by Target-AID base editing identifies sensitive sites across the yeast genome. A) Experimental design. Essential genes were scanned for sites appropriate for Target-AID mutagenesis. Mutational outcomes include silent (grey triangle) and missense (black triangle) mutations, as well as stop codons (*). DNA fragments bearing the gRNA sequences were synthesized as an oligonucleotides pool and cloned into a co-selection base editing vector. Using gRNAs as molecular barcodes, the abundance of cell subpopulations bearing mutations was measured during mutagenesis and bulk competition. Mutations with fitness effects were inferred from a reduction in the relative barcode read count. B) Cumulative distribution of z-scores of the $\log_2$ fold-change in gRNA abundance between mutagenesis and the end of the bulk competition experiment averaged between replicates (see Figure S2). A 5% false positive threshold was calculated by fitting a distribution of abundance variation z-score of the sequenced gRNAs with synthesis errors (SE gRNAs) and is represented by a dotted black line. The distribution of target types in the 1,118 gRNAs with Negative Effects (GNE) is shown in the inset. C) Positions of base editing target sites in the yeast genome. Telomeric regions are depleted in target sites because very few essential genes are located there. GNEs are shown in red, and other gRNAs are in black. The orientation of the line matches the targeted strand relative to the annotated coding sequence. D) Decline in barcode abundance (on a log scale) between timepoints after mutagenesis for gRNAs targeting *GLN4*, a tRNA synthetase. Median barcode abundance across the entire library through time is shown in green. The red lines represent the gRNAs categorized as having a significant effect (GNE) for this gene, while non-significant gRNAs (NSG) are shown in black. The gRNA with the most extreme z-score targets residue G267. E) Mutagenesis of *GLN4*-G267 confirms its essential role for protein function (See methods and Figure S3A). Tetrad dissection of a heterozygous deletion mutant bearing an empty vector results in only two viable spores, while the wild-type copy in the same vector restores growth.

Dissection of the two heterozygous mutants bearing a plasmid with the most probable single mutant based
on the known activity window of Target-AID shows both mutations are lethal.

After applying a stringent filtering threshold based on barcode read count at the mutagenesis step

(Figure S2), we identified a total of ~17,000 gRNAs for which we could evaluate fitness effects.

Replicate data for gRNAs passing the minimal read count selection criteria show high correlation

across experimental time points (Figure S3) and cluster by experimental step (Figure S4),

showing that the approach is reproducible. Using the distribution of abundance variation of non-

functional gRNAs with synthesis errors as a null distribution (see methods), we identified 1,118

gRNAs across 605 genes or loci with significant negative effects (GNE) on cell survival or

proliferation. GNEs are distributed evenly across the yeast genome (Figure 1B and 1C),

suggesting no inherent bias against specific regions.

An example of barcode abundance variation through time for all gRNAs targeting *GLN4* is shown

in Figure 1D. *GLN4* is an essential gene coding for a glutamine t-RNA synthetase. To confirm the

effect of the predicted mutations, we transformed a centromeric plasmid bearing a wild-type or

mutated copy of the gene under the control of its native promoter (Ho *et al.* 2009) in a

heterozygous deletion background (Giaever *et al.* 1999). Following dissection, spore survival was

compared between wild-type and mutated copy of *GLN4* (Figure S5). Using this approach, we

confirmed the strong fitness effect of the best scoring GNE for *GLN4*, as the most probable

mutations generated are in fact lethal (Figure 1D).

Comparison of GNE induced mutations with variant effect predictions
If GNEs indeed induce specific deleterious mutations, these mutations should be predicted to be

more deleterious than those of Non-Significant gRNAs (NSG). We tested two recently published

resources for variant effect prediction: Envision (Gray *et al.* 2018) and Mutfunc (Wagih *et al.*

2018). Envision is based on a machine learning approach that leverages large-scale saturated

mutagenesis data of multiple proteins to perform quantitative predictions of missense mutation

effects on protein function. The lower the Envision score, the higher the effect on protein function.

Mutfunc aggregates multiple types of information such as residue conservation through the use

of SIFT (Ng and Henikoff 2003) as well as structural constraints to provide a binary prediction of

variant effect. Mutations with a low SIFT score have a lower chance of being tolerated, while those

with a positive ΔΔG are predicted to destabilize protein structure or interactions. Both Envision

and the Mutfunc aggregated SIFT data cover the majority of the most probable mutations

generated by the gRNA library (Figure S6A). The structural modeling information had much lower

coverage, covering at best around 12% of the most probable mutations (Figure S6B).

Figure 2: GNE induced mutations are enriched in predicted deleterious effects A) SIFT score
distributions for the most likely induced mutations of both GNEs (blue) and NSGs (red). The thresholds for
the categories used in the enrichment calculations in B) are shown as black dotted lines. SIFT scores
represent the probability of a specific mutation being tolerated based on evolutionary information: the first
threshold of 0.05 was set by the authors in the original manuscript (Ng and Henikoff 2003) but might be
permissive considering the number of mutations tested in our experiment. All GNE vs NSG score
comparisons are significant (Welch's t-test p-values: 1.19×10^{-24}, 3.01×10^{-24}, 9.00×10^{-12}, 1.55×10^{-12}). The
box cutoff is due to the large fraction of mutations for which the SIFT score is 0. B) Enrichment folds of
GNEs over NSGs for different variant effect prediction measurements. Envision score (Env.), SIFT score
(SIFT), protein folding stability based on solved protein structures (Struct. ΔΔG), protein folding based on
homology models (Model ΔΔG) and protein-protein interaction interface stability based on structure data
(Inter. ΔΔG). The raw values used to calculate ratios are shown in Supplementary table 1 The prediction
based on conservation and experimental data are grouped under 'Predictors' and those based on the
computational analysis of protein structures and complexes under 'Structural'.

As expected, mutations generated by GNEs showed significantly lower SIFT scores (Figure 2A)

and showed enrichment for strong effects predicted by SIFT, and Envision. Indeed, all four most

probable substitutions created by GNEs are about twice more likely to be predicted to have a

large deleterious effect by Envision or a very low chance of being tolerated as predicted by SIFT

compared to NSG gRNAs. The high homogeneity of Envision scores across the proteome makes

it harder to interpret. As such, the shift in score values is more subtle but supports that GNE

mutations are generally more likely to be deleterious as well (Figure S6C, Figure S7A).

Mutation with destabilizing effects as predicted by structural data also appeared to be enriched,

but low residue coverage limits the strength of this association. This is supported by the raw $\Delta\Delta G$

value distributions, which show a significant tendency for GNE mutations to be more destabilizing

(Figure S7B,C,D). However, the shift in distribution only achieved significance for certain mutation

predictions based on solved structures and homology models. While low residue coverage limits

our statistical power, this weak apparent enrichment for mutations affecting protein stability may

reflect the marginal stability of the target proteins (DePristo *et al.* 2005), resulting in individual

destabilizing mutations having a limited effects on fitness. As expected from known experimental

data on mutagenesis outcomes (Nishida *et al.* 2016), signal was usually stronger for the most

probable C to G mutation.

Sensitive sites provide new biological insights

Because our screen specifically targeted essential genes, many gRNAs cause mutations in highly

conserved regions. To illustrate this, we focus on the highest scoring GNE targeting *GLN4*, a

tRNA synthetase, shown in Figure 1D. The gRNA 33725 mutates a glycine at position 267 into

either an arginine or a serine. Glycine 267 is part of the "HIGH" motif, characteristic of class I

tRNA synthetases, and is involved in ATP binding and catalysis and is highly conserved through

evolution (Eriani *et al.* 1990). As expected, the region around the "HIGH" motif shows both a low

evolutionary rate based on inter-species comparisons and a much lower variant density in yeast

populations compared to other domains of Gln4 (Figure S3B), showing conservation both on a

short and long timescales. Surprisingly, mutagenesis experiments in the bacterial homolog MetRS

concluded that mutating this residue from glycine to alanine did not alter significantly catalysis

while mutating it to proline had a strong disruptive effect (Schmitt *et al.* 1995). We found that

mutating Gly 267 either to Arg and Ser was enough to cause protein loss of function (Figure 1D).

Other sensitive sites identified in *GLN4* by our screen are also clustered in regions with slow

evolutionary rates. Interestingly, one of those mutations affects residue R568, which has been

hypothesized to play a conserved role from bacteria to yeast in the anti-codon and glutamine

recognition process (Grant *et al.* 2013).

Since Target-AID can only generate a limited range of amino acid substitutions from a specific

coding sequence, we investigated whether any of these mutational patterns were enriched in

GNEs (Figure 3A, source data in Supplementary tables S2, S3, and S4). We found several

deviations from random expectations in both C-to-G and C-to-T mutation ratios as well as in

mutation combination ratios. Three out of four of the mutation pair patterns involving glycine were

enriched in GNEs. For example, the Glycine to Arginine or Serine mutation pair (as exemplified

by guide 33725 targeting *GLN4*) is the second most enriched pattern, being almost four-fold

overrepresented in GNE outcomes. This pattern is consistent with the fact that Arginine has

properties highly dissimilar to those of Glycine (Sneath 1966), making these substitutions highly

deleterious. Furthermore, as Glycine residues are often important components of cofactor binding

motifs (eg.: Phosphates) (Copley and Barton 1994) this observation might reflect a tendency for

GNEs to alter these sites. Interestingly, genes for which more than one GNE were detected were

enriched for molecular function terms linked to cofactor binding (Supplementary table 5). This

suggests that the GNEs might indeed have had a tendency to affect protein function through

mechanisms other than protein or interaction interface destabilization. These protein properties

depend on many residues, making them more robust to single amino acid substitutions, whereas

cofactor binding may depend specifically on a handful of residues, making these sites critical for

function.

As expected, there is a strong enrichment for patterns that result in mutation to stop codons: both C-to-G patterns (Tyrosine to stop and Serine to stop) but only one C-to-T pattern (Tryptophan to stop) was overrepresented significantly. Stop codon generation in one outcome also drove enrichment in the other: for example, the link between Serine to Stop (C-to-G) appears to be the cause of the Serine to Leucine (C-to-T) overrepresentation. Both mutation pairs involving mutating a Tryptophan to a stop via a C-to-T mutation: this is not surprising, as the alternative mutations Tryptophan to Serine or Cysteine are also highly disruptive (Sneath 1966). Changes between similar amino acids were also generally depleted in GNEs, for example the Alanine to Glycine/Valine pair. Mutations in intronic sequences and putative non-functional peptides were also underrepresented, as were most patterns leading to silent mutations. These results show the power of this approach to identify important functional sites across the genome.

Figure 3 GNE mutations are enriched for specific amino acid substitution patterns and identify critical sites for protein function. A) Fold depletion and enrichment volcano plots for the most probable mutations induced by GNEs in the screen. Enrichment and depletion values were calculated by comparing the relative abundance of each mutation among GNEs and NSGs using Fisher's exact test. Mutation patterns significantly depleted are shown in blue, while those that are enriched are in red. The significance

threshold was set using the Holm-Bonferroni method at 5% FDR and is shown as a dotted grey line. B)
Protein variant frequency among 1000 yeast isolates (black dots) and residue evolutionary rate across
species (blue line) for *RAP1*. The target site for the GNEs targeting T486 is highlighted by a red line while
the other detected GNEs target sites are shown by a grey line. C) Tetrad dissections confirm most *RAP1*
GNE induced mutations indeed have strong fitness effects, as well as other substitutions targeting these
sites.

While the precise targeting of our method also allows us to investigate amino acid residues with

known functional annotations such as post-translational modifications. We found no significant

enrichment for gRNAs mutating directly annotated PTMs (ratio$^{GNE\ PTM}$ = 19/1118, ratio$^{NSG\ PTM}$

243/15536, Fisher's exact test p=0.71). This is consistent with the hypothesis that many PTM

sites may have little functional importance (Landry *et al.* 2009) and thus their mutations may have

no detectable effects for a large part. The same was also observed for gRNAs mutating residues

near known PTMs that could disturb recognition sites (ratio$^{GNE\ nearPTM}$ = 130/1118, ratio$^{NSG\ nearPTM}$

= 1698/15536, Fisher's exact test p=0.43). However, GNEs that do target annotated PTM sites

might provide additional evidence supporting the importance of these sites in particular. For

example, the best scoring GNE in the well-studied transcriptional regulator *RAP1* is predicted to

mutate residue T486. This threonine has been reported as phosphorylated in two previous studies

(Albuquerque *et al.* 2008; Holt *et al.* 2009), but the functional importance of this phosphorylation

has not been explored yet. Residue T486 is located in a disordered region in the DNA binding

domains (Konig *et al.* 1996), which part of the only *RAP1* fragment essential for cell growth

(Graham *et al.* 1999; Wu *et al.* 2018).

Because the available wild-type *RAP1* plasmid (see methods) does not complement gene

deletion growth phenotype, we used a different strategy for confirmation that relied on CRISPR-

mediated knock-in (see methods and Figure S8). While we could not confirm that the two most

likely mutations predicted to be caused by the GNE had a detectable fitness effect in these

conditions, we found that phosphomimetic mutations at this position were lethal (Figure 3C and

D) while most other amino acids were well tolerated. This suggests that the constitutive

phosphorylation of this residue would be highly deleterious. We could also confirm deleterious

effects for GNE induced mutations targeting residues R523 and A540, while mutations at residue

A510 had no detectable effect on fitness (Figure 3C and D). As we only tested progeny survival

on rich media and at a permissive temperature and the screen was performed in synthetic media

at 30°C, these mutants might still affect cell phenotype but in an environment-dependent manner.

gRNA properties influence mutagenesis efficiency

There are still very few high-throughput experimental datasets available that allow the

investigation of which gRNA properties affect editing efficiency in the context of base editing. Such

large-scale data was key in developing models to optimize Cas9 nuclease activity in other types

of genome editing experiments, which revealed that sequence specific motifs and thermodynamic

RNA properties can be key features (Doench *et al.* 2014, 2016; Wong *et al.* 2015). As gRNAs

showing high Cas9 nuclease activity might have poor base editing activity (Kim *et al.* 2017),

existing datasets are not easily transferable. We therefore examine what gRNA and target

sequence features could influence mutagenesis efficiency. To do so, we focused on the subset

of gRNAs with the potential to generate stop codons (stop codon generating gRNAs, SGG) in

essential genes (Figure 4A). Successful mutagenesis by SGGs should result in cell death or no

proliferation, and a sharp decrease in read abundance, also serving as a positive control for

fitness effects within the screen. As most gRNAs were designed to target the first 75% of the

coding sequences of essential genes, it is expected that stop codons in these genes would lead

to a loss of function.

Figure 4: gRNA and target properties affect mutagenesis efficiency. A) Since Target-AID can generate
both C to G and C to T mutations, many codons can be targeted to create premature stop codons. B)
Cumulative z-score density of SGGs grouped by the mutational outcome generating the stop codon. A
higher rate of GNE is observed for gRNAs for which a C-to-G mutation at the highest editing activity position
generates a stop codon mutation. The significance threshold is shown as a black dotted line C) Cumulative
z-score density of SGG with a different number of mutational outcomes that could result in a stop codon.
gRNAs for which more than one mutational outcome results in a stop codon show a higher mutagenesis
success rate. D) Cumulative z-score density of SGG targeting the coding or non-coding strand. Stop codon
generating gRNAs targeting the non-coding strand of essential genes show higher efficiencies compared
to those targeting the coding strand. E) Distributions of modeled RNA/DNA duplex melting temperature for
all stop codon generating gRNAs, those with no significant effects, and SGG GNE. P-values were
calculated using the two-sample Kolmogorov-Smirnov test. F) Stop codon generating gRNAs GNE
enrichment compared to the expected GNE ratio for different melting temperature ranges.

Data from the original Target-AID study (Nishida *et al.* 2016) suggests that the most prevalent

outcome for an edited site is a C-to-G transversion. Our data support this observation, as gRNAs

which would lead to a C-to-G mutation at the highest activity site of the editing window have the

highest GNE detection rate (Figure 4B). It was also suggested that Target-AID could modify

multiple nucleotides within the activity window that could be edited during mutagenesis. Our data

support this observation, as gRNAs for which two outcomes have the potential to generate a stop

codon are markedly more efficient than those with only one stop codon outcome (Figure 4C). This

finding also extends to gRNAs that do not generate stop codons (Figure S9A).

We observed that the targeted strand relative to transcription greatly influenced editing efficiency

(Figure 4D). This strand effect can be explained by multiple factors. First, there are multiple

outcomes leading to mutation to a stop codon starting from a TGG codon (shown in Figure 4A).

This codon is the only one that can be targeted on the non-coding strand to generate a stop

codon. Second, repair efficiency has been shown to be higher for the transcribed strand in yeast

(Reis *et al.* 2012). Finally, as the non-coding strand is the one which is transcribed, a deamination

event there might lead to consequences at the protein level more rapidly because it does not

need DNA replication to be present on both strands. gRNAs that do not generate stop codons

also have a higher chance of having a fitness effect if they target the non-coding strand (Figure

S9B), but we did not observe any effects of the chromosomal strand on efficiency (Figure S9C).

One other parameter with a high impact on mutagenesis rate is the predicted melting temperature

of the RNA-DNA duplex formed by the gRNA sequence and its target DNA sequence (Figure 4E).

The distribution of the melting temperature shows a clear shift between stop codon generating

gRNAs that have an effect on fitness and those that do not. gRNAs with low values have a lower

chance of being detected as having effects, while gRNAs with higher values are enriched for GNE

(Figure 4F). This observation also extends to gRNAs that do not generate stop codons (Figure

S9D, E). This enrichment cannot be attributed to technical biases in library preparation or high-

throughput sequencing that would tend to lower their abundance as melting temperature shows

practically no correlation with read count at every time point (Figure S10). Furthermore, this effect

is not caused by target position bias within target genes or a strong correlation between GC

content and the targeted position (Figure S11). As binding energy can differ drastically even within

groups of gRNAs with similar GC content (Figure S9F), this could provide a useful criterion to help

select efficient gRNAs.

Discussion

We tested whether the Target-AID base editor is amenable for genome-wide mutagenesis. . Using the yeast essential genes as test cases, we identified hundreds of gRNAs and this genome position with significant effects on cellular fitness when mutated. The precision and traceability of Target-AID genome editing allowed us to predict the mutational outcomes of GNE and to confirm their effects using orthogonal approaches. We used this data to investigate which factors influence base editing efficiency and found multiple gRNAs and target properties that affect mutagenesis and that could be optimized for future experiments for specific genomic space. By focusing on a few highly relevant variants, we highlighted the power of our approach to generate new biological insights.

In previously published methods such as TAM and CRISPR-X (Hess *et al.* 2016; Ma *et al.* 2016), the semi-random nature of the editing forces the use of mutant allele frequencies as a readout for mutational fitness effects, potentially limiting the scale of the experiments because only one genomic region can be targeted at a time. To complement these approaches, we use more predictable base editing to increase dramatically the number of target loci, albeit at the cost of a lower mutational density. Although our results demonstrate the feasibility of base editing screening at a large scale with applications beyond stop codon generation, our approach still requires further development. For instance, the use of a base editor with multiple possible mutagenesis outcomes complexifies the prediction of editing outcomes, which can, in turn, make GNE confirmation challenging. Using a base editor that channels mutational outcomes such as cytidine deaminase-uracil glycosylase inhibitor (UGI) fusion can address this problem but decreases the number of mutations explored during the experiment. However, recently published data on cytidine deaminase-UGI fusion off-target activity has shown they could lead to off-target editing in vivo at a much higher rate compared to adenine base editors or the Cas9 nuclease (Jin *et al.* 2019; Zuo *et al.* 2019). Although there is currently no high throughput data on the off-target

activity of Target-AID, data generated in yeast in the original publication suggests far lower rates

than those recently reported in mammalian cells (Nishida *et al.* 2016).

We provide key empirical data on parameters that can be used to optimize base editing efficiency,

based on gRNA dependent properties such as target strand and GC content. The results we

observed differ from what has been reported for Cas9-based genome editing, in which high gRNA

RNA/DNA duplex binding has been associated with lower mutagenesis efficiency (Wong *et al.*

2015). Our data thus confirms the observation that parameters associated with Cas9 editing

cannot readily be transferred to base editors (Kim *et al.* 2017). Furthermore, the temperature at

which experiments are performed might affect efficiency for certain gRNAs with low gRNA-DNA

duplex binding energy and should be considered when designing base editing experiments in

different organisms. However, it remains to be confirmed whether the enrichment for certain

gRNA properties we observed are specific to Target-AID or will also be transferable to other base

editors as this may depend on the enzymatic properties of these proteins.

The field of base editing is rapidly evolving, with new tools being developed constantly. One of

the most recent additions to this fast-growing toolkit is engineered Cas9 enzymes with broadened

PAM specificities (Nishimasu *et al.* 2018), which have already been shown to be compatible with

base editors. More flexible PAM requirements are especially useful for base editing applications,

as they increase the number of sites to be edited and also the number of potential gRNAs per

site, increasing the chances of choosing optimal properties and thus greater efficiency (Dandage

et al. 2019). Data experiments at a scale which bridges saturation mutagenesis methods and

genome-wide knock-out studies will expand our understanding of mutational effects and resilience

within cellular networks.

Methods

Generation of a gRNA library for Target-AID mutagenesis of essential genes in yeast

The Target-AID base editor has an activity window between base 15 to 20 in the gRNA sequence starting from the PAM, and the efficiency at these different positions was characterized in Nishida *et al.* 2016. This allowed us to predict the mutational outcomes for a specific gRNA provided the number of editable bases in the window is not too high. To select gRNAs, we parsed a database of gRNA targets for the *S. cerevisiae* reference genome sequences (strain S288c) (Dicarlo *et al.* 2013) and applied several selection criteria. Since the screen was to be performed in the BY4741 strain, all gRNAs (unique seed sequence, no NAG site) within the database were aligned to the reference genome of that strain using Bowtie (Langmead *et al.* 2009). Only gRNAs with a single perfect alignment were kept for subsequent steps. To select gRNAs amenable to Target-AID base editing, we selected gRNAs with cytosines within the highest activity window of the editor (positions -17 to -19 starting from the PAM). To limit the total number of possible mutational outcomes, gRNAs with three cytosines within the window were removed as well as those with two cytosines at the highest activity positions. Next, we filtered out any gRNA containing a BsaI restriction site to prevent errors during the library cloning step.

The list of essential genes (n=1156) (Winzeler *et al.* 1999; Giaever *et al.* 2002) was used to discriminate between gRNAs targeting essential or non-essential genes (retrieved from http://www-sequence.stanford.edu/group/yeast_deletion_project/Essential_ORFs.txt). Among non-essential genes, data from Qian et al. 2012 (Qian *et al.* 2012) was used to create categories of fitness effects. If the fitness score (averaged across media and replicates) of a gene was below 0.75, it was categorized as "high effect" on fitness. We excluded auxotrophic marker genes as well as *CAN1*, *LYP1*, and *FCY1* because those could be used as co-selection markers (Després *et al.* 2018). Gene deletions with an averaged fitness score between 0.999 and 1.001 were categorized as having "no detectable effect" on fitness. We selected gRNAs targeting essential

and high effect genes, as well as gRNAs targeting a set of 38 randomly chosen no effect genes.

To further limit the space of gRNAs examined, only gRNAs mapping from the 0.5[th] percent to the

75[th] percent of coding sequences were chosen. We also added gRNAs targeting all known yeast

introns (Ares lab Database 4.3) (Grate and Ares 2002) and putative non-functional peptides

(Smith *et al.* 2014) selected with the same strategy except for the constraints on gRNA position

within the sequence of interest. This resulted in a set of 39,989 gRNAs: library properties are

summarized in Figure S1.

Library construction

The plasmids, oligonucleotides, and media used in this study are presented as supplementary

tables S6, S7 and S8 respectively. The oligo pool was synthesized by Arbor Biosciences

(Michigan, USA) and was cloned into the pDYSCKO vector using Golden Gate Assembly (New

England Biolabs, Massachusetts, USA) with the following reaction parameters:

NEB GG buffer 10X	2 µl
pDYSCKO [75ng/ul]	1 µl
Oligo pool [2ng/ul]	1 µl
NEB GG mix	1 µl
Water	15 µl

The ligation mix was transformed in *E. coli* strain MC1061 (*[araD139]$_{B/r}$ Δ(araA-leu)7697 ΔlacX74*

galK16 galE15(GalS) λ- e14- mcrA0 relA1 rpsL150(strR) spoT1 mcrB1 hsdR2, Casadaban and

Cohen 1980) using a standard chemical transformation protocol and plated on ampicillin selective

media to select for transformants. Serial dilution of cells after outgrowth were plated and then

used to calculate the total number of clones produced by the cloning reaction. Quality control of

the assembly was performed by Sanger sequencing ~10 clones per assembly reaction. Cells

were scraped from plates by adding ~5 ml of sterile water, incubating a few minutes at room

temperature, and then using a glass rake to resuspend colonies. Resuspended plates were then

pooled together in a single flask per reaction, which was then used to make glycerol stocks of the

library and cell pellets for plasmid extraction. The Qiagen Midi-Prep kit (Qiagen, Germany) was

used to extract plasmid DNA from cell pellets by following the manufacturer's instructions. The

DNA concentration of each eluate was then measured using a NanoDrop (Thermofisher,

Massachusetts, USA), and a normalized master library for yeast transformation was assembled

by combining equal quantities of each assembly pool.

Library transformation in yeast

Competent BY4741 (*MATa his3Δ1 leu2Δ0 met15Δ0 ura3Δ0*) cells were first transformed with the

pKN1252 (p415-GalL-Target-AID) plasmid using a standard lithium acetate method (Gietz and

Schiestl 2007). Transformants were selected by plating cells on SC-L. After 48 h of growth,

multiples colonies were used to inoculate a starter liquid culture for competent cells preparation

using the standard lithium acetate protocol (Gietz and Schiestl 2007): a culture volume of 200 ml

was used to generate enough competent cells for mass transformation. The large-scale library

transformation was performed by combining 40 transformation reactions performed with 40 ul of

competent cells and 5 ul plasmid library (240 ng/ul) after the outgrowth stage and plating 100 ul

aliquots on SC-UL: cells were then allowed to grow at 30°C for 48 h. A 1/1000 serial dilution of

the cell recovery was plated in 5 replicates and used to calculate the number of transformants

obtained. The total number of transformants reached 3.48 x10^6 CFU, corresponding to about

100X coverage of the plasmid pool.

Target-AID mutagenesis and competition screening

The mutagenesis protocol is an upscaled version of our previously published method and is

shown in Figure S2. Transformants were scraped by spreading 5 ml sterile water on plates and

then resuspending cells using a glass rake. All plates were pooled together in the same flask, and

the OD of the yeast resuspension was measured using a Tecan Infinite F200 plate reader (Tecan,

Switzerland). Pellets corresponding to about 6 x 10^8 cells were washed twice with SC-UL without

a carbon source and then used to inoculate a 100 ml SC-UL +2% glucose culture at 0.6 OD two

times to generate replicates A and B. Cells were allowed to grow for 8 hours before 1×10^9 cells

were pelleted and used to inoculate a 100 ml SC-UL + 5% glycerol culture. After 24 hours, 5 x

10^8 cells were pelleted and either put in SC-UL + 5% galactose for mutagenesis or SC-UL + 5%

glucose for a mock induction control. Target-AID expression (from pKN1252) was induced for 12

hours before 1×10^8 cells were pelleted and used to inoculate a canavanine (50 µg/ml) co-

selection culture in SC-ULR. After 16 hours of incubation, 5×10^7 cells of each culture were used

to inoculate 100 ml SC-UR, which was grown for 12 hours before 5×10^7 cells were used to

inoculate a final 100 ml SC-UR culture which was grown for another 12 hours. Cell pellets were

washed with sterile water between each step, and all incubation occurred at 30°C with agitation.

~2×10^7 cells were taken for plasmid DNA extraction at the end of each mutagenesis and

competition screening step.

Yeast plasmid DNA extraction

Yeast plasmid DNA was extracted using the ChargeSwitch Plasmid Yeast Mini Kit (Invitrogen,

California, USA) by following the manufacturer's protocol with minor modifications: Zymolase

4000 U/ml (Zymo Research, California, USA) was used instead of lyticase, and cells were

incubated for 1 hour at room temperature, one min at -80°C, and then incubated for another 15

minutes at room temperature before the lysis step. Plasmid DNA was eluted in 70 µl of E5 buffer

(10 mM Tris-HCl, pH 8.5) and stored at -20°C for use in library preparation.

Next-generation library sequencing preparation

Libraries were prepared by using two PCR amplification steps, one to amplify the gRNA region of

the pDSYCKO plasmid pool and the second to add sample barcodes as well as the Illumina p5

and p7 sequences (Yachie *et al.* 2016). Oligonucleotides for library preparation are shown in the

first part of the oligonucleotide table. Reaction conditions for the first PCR were as follows:

Phusion HF buffer (NEB) 5X	5 µl
dNTPs 10 mM	0.5 µl
pDYSCKO_gRNA_for 10 µM	1.25 µl
pDYSCKO_gRNA_rev 10 µM	1.25 µl
Phusion polymerase	0.5 µl
Template DNA (<1 ng/µl)	5 µl
PCR grade water	11.7 µl

493   Thermocycler protocol:

Temperature (°C)	Time (s)	Cycles
98	30	1
98	10	
58	15	16
72	5	
72	5	1

495   The resulting product was verified on a 2% agarose gel colored with Midori Green Advance

496   (Nippon Genetics, Japan) and then gel-extracted and purified using the FastGene Gel/PCR

497   Extraction Kit (Nippon Genetics, Japan). The purified products were used as the template for the

498   second PCR reaction, with the following conditions:

Phusion Mastermix-HF (NEB)	10 µl
P5-barcode-X oligo 1.333 µM	3.75 µl
P7-barcode-Y oligo 1.333 µM	3.75 µl
Template DNA (~1 ng/µl)	2.5 µl

500   Thermocycler protocol:

Temperature (°C)	Time (s)	Cycles
98	30	1
98	10	15
60	10	
72	60	
72	300	1

PCR products were verified on a 2% agarose gel colored with Midori Green Advance (Nippon

Genetics, Japan) and then gel-extracted and purified using the FastGene Gel/PCR Extraction Kit

(Nippon Genetics, Japan). Library quality control and quantification were performed using the

KAPA Library Quantification Kit for Illumina platforms (Kapa Biosystems, Massachusetts, USA)

following the manufacturer's instructions. Libraries were then run on a single lane on HiSeq 2500

(Illumina, California, USA) with paired-end 150 bp in fast mode.

Large-scale screen sequencing data analysis

The custom Python scripts used to analyze the data are available as supplementary material,

packages and software used are presented in Supplementary table 9. Briefly, reads were

separated into three subsequences for alignment: the P5 barcode, the gRNA, and the P7 barcode.

Each of these was aligned using Bowtie (Langmead *et al.* 2009) to an artificial reference genome

containing either the barcodes or gRNA sequences flanked by the common amplicon sequences.

The gRNA sequences are aligned both with 0 or 1 mismatch allowed, and misalignment position

and type were stored. Information on barcode and gRNA alignment for each read was stored and

combined to generate a barcode count per library table, a list of mismatches in alignments for

each gRNA in each library, as well as mismatch types and counts for the same gRNA across all

libraries.

Synthesis error within oligonucleotide libraries is one of the major limits of current large-scale

genome editing screening methods. These errors can introduce gRNA sequences that cannot

perform mutagenesis because the gRNA sequence does not match a site in the genome. We

refer to those gRNAs as SE gRNAs. In our experiment, the stringent selection criteria used to

select gRNAs limited the risk of off-target effects even for gRNAs with one mismatch, minimizing

the risk that a synthesis error gRNA could lead to editing at another site in the genome. We

therefore decided to use highly abundant SE gRNAs as negative controls to obtain a null

distribution of abundance variation for gRNAs with no fitness effects. To differentiate synthesis

errors from sequencing errors, we used the mismatch type and count table to assess whether a

particular mismatched gRNA constitutes a too large fraction of the reads associated with a gRNA

to be simply a repeated sequencing error. For each error, we test if:

530
$$\frac{N_{readsformismatch}}{N_{perfectalignment}} > 0.075$$

and discarded the reads associated with the specific mismatch alignment. This threshold was

obtained by iteratively testing different threshold values in an effort to maximize the gain in gRNA

counts while minimizing the noise added by incorrect assignments. Read counts per library for

abundant ($N_{readsformismatch} > 1,000$) SE gRNAs were kept to serve as negative controls when

measuring fitness effects, resulting in a set of 1,032 abundant SE gRNAs. gRNAs absent from

more than half of the libraries (4446 out of 39,989) were removed from the analysis before gRNA

abundance calculations.

Detecting mutations with high fitness effects

Barcode sequencing competition experiments use DNA barcodes to measure the relative

abundance of many different subpopulations of cells grown in the same pool (Robinson *et al.*

2014). Since each gRNA is linked to its possible mutagenesis outcomes, we can use relative

gRNA abundance to detect mutations with significant fitness effects. To do so, the $\log_2$ of the

relative abundance of a barcode after mutagenesis is compared with its abundance at the end of

the screen:

$$\Delta\log_{2_{gRNA}} = \log_2\left(\frac{N_{readsgRNAt_1}}{N_{readst_1}}\right) - \log_2\left(\frac{N_{readsgRNAt_0}}{N_{readst_0}}\right)$$

For each gRNA, the measured fitness effect is the product of the effect of the mutational outcomes

on growth and of the mutation rate within the cell subpopulation bearing this particular gRNA.

Relative counts will also vary stochastically because of variation in sequencing coverage

depending on the time point and replicate. To reduce the impact of these effects, a minimal read

count at the end of the galactose induction step was used to filter out low abundance gRNAs. We

found a minimal read threshold of n=54 provided a good tradeoff between the number of gRNAs

eligible for analysis and inter-replicate correlation.

Using the distribution of $\Delta\log2$ values, we calculated a z-score for each gRNA in both replicates.

We then averaged z-scores between replicates and compared the score distributions between

SE and Non-SE gRNAs. This revealed the presence of a left-skewed tail in the z-score distribution

of valid gRNAs, which is absent in the SE. Because the number of SE gRNAs is smaller than the

one of functional gRNAs by almost two orders of magnitude, a type I error (false positives)

empirical threshold based solely on a weighted SE z-score distribution was not practical. To

resolve this, we fitted a Gumbell left skewed distribution to the SE gRNAs z-score distribution and

used it to approximate the type I error rate as a function of the z-score. We set a significance

threshold such as that all gRNAs at z-scores for which the estimated false positive rate is below

or equal to 5% are considered GNEs.

Complementation assays

Experiments were performed in heterozygous deletion mutants from the YKO project

heterozygous deletion strain set (Dharmacon, Colorado, USA). For each gene, a single colony

streaked from the glycerol stock was used to prepare competent cells using the previously

described lithium acetate protocol. To generate mutant alleles of the genes of interest, we

performed site-directed mutagenesis on the appropriate MoBY collection plasmid (Ho *et al.* 2009).

These centromeric plasmids encode the yeast gene of interest under the control of their native

promoters and terminators. Mutagenesis reactions were performed with the following reaction

setup:

Kapa HiFi buffer (Kapa biosciences) 5X	5 µl
dNTPs 10µM	0.75 µl
mutation_for 10µM (see table 7)	0.75 µl
mutation_rev 10µM (see table 7)	0.75 µl
Kapa Hot-start polymerase	0.5 µl
Template plasmid DNA (15ng/ul)	0.75 µl
PCR grade water	16.5 µl

574 Thermocycler protocol:

Temperature (°C)	Time (s)	Cycles
95	300	1
98	20	
60	15	20
72	720	
72	1080	1

After amplification, the mutagenesis product was digested with DpnI for 2 hours at 37°C and 5 ul

was transformed in *E. coli* strain BW23474 (F-, Δ*(argF-lac)169*, Δ*uidA4::pir-116*, *recA1*,

rpoS396(Am), *endA9(del-ins)::FRT*, *rph-1*, *hsdR514*, *rob-1*, *creC510,* Haldimann *et al.* 1996).

Transformants were plated on 2YT+Kan+Chlo and grown at 37°C overnight. Plasmid DNA was

then isolated from clones and sent for Sanger sequencing (CHUL sequencing platform, Université

Laval, Québec City, Canada) to confirm mutagenesis success.

Competent cells of target genes were transformed with the appropriate mutant plasmids as well

a the original plasmid bearing the wild-type gene and the empty vector (Zhao *et al.* 2016), and

transformants were selected by plating on SC-U (MSG). Multiple independent colonies per

transformation were then put on sporulation media until sporulation could be confirmed by

microscopy. For tetrad dissection, cells were resuspended in 100ul 20T zymolyase (200mg/ml

dilution in water) and incubated for 20 minutes at room temperature. Cells were then centrifuged

and resuspended in 50ul 1M sorbitol before being streaked on a level YPD plate. All dissections

were performed using a Singer SporePlay microscope (Singer Instruments, UK). Plate pictures

were taken after five days incubation at room temperature except for the RAP1 plasmid

complementation test for which the picture was taken after three days. Pictures are shown in

Supplementary image 1.

Strain construction for confirmations in *RAP1*

Because the MoBY collection plasmid for RAP1 cannot fully complement the gene deletion

(Supplementary image file 1), we instead performed confirmations by engineering mutations a

diploid strain to create heterozygous mutants. *RAP1* was first tagged with a modified version of

fragment DHFR F[1,2] (the first half) of the mDHFR enzyme (Tarassov *et al.* 2008). The

mDHFR[1,2]-FLAG cassette was amplified using gene-specific primers and previously described

reaction parameters (Tarassov *et al.* 2008). Cells were transformed with the cassette using the

previously described transformation protocol and were plated on YPD+Nourseothricine (YPD+Nat

in Media table). Positive clones were identified by colony PCR and successful fragment fusion

was confirmed by Sanger sequencing (CHUL sequencing platform). We then mated the confirmed

clones with strain Y8205 (*Matα can1::STE2pr-his5 lyp1::STE3prLEU2 Δura3 Δhis3 Δleu2*, Kindly

gifted by Charlie Boone) by inoculating a 4ml YPD culture with overnight starter cultures of both

strains and letting the culture grow overnight. Cells were then streaked on YPD+Nat and diploid

cells were identified by colony PCR using mating type diagnosis primers (Huxley *et al.* 1990).

To create heterozygous deletion mutants of the target gene, we amplified a modified version of

the *URA3* cassettes that could then be targeted with the CRISPR-Cas9 system to integrate our

mutations of interest using homologous recombination at the target locus. The oligonucleotides

we used differ from those commonly used in that they amplify the cassette without the two LoxP

sites present at both ends. We found it necessary to remove those sites as one common

mutational outcome after introducing a double-stranded break in the *URA3* cassette was inter-

LoxP site recombination without the integration of donor DNA at the target locus. These modified

cassettes recombine with DNA upstream the target gene on one end and the mDHFR F[1,2]

fusion on the other, ensuring that the heterozygous deletion is always performed at the locus that

is already tagged. Cassettes were transformed using the standard lithium acetate method, and

cells were plated on SC-U (MSG) selective media. Heterozygous deletion mutants were then

confirmed by colony PCR.

CRISPR-Cas9 mediated Knock-in of targeted mutations

Mutant alleles of target genes were amplified in two fragments using template DNA from the

haploid tagged strain (See Figure S8). The two fragments bearing mutations are then fused

together by a second PCR round to form the final donor DNA. This DNA was then co-transformed

with a plasmid bearing Cas9 and a gRNA targeting the URA3 cassette for HDR mediated editing

using a standard protocol (Ryan *et al.* 2016). Clones were then screened by PCR to verify donor

DNA and mutation integration at the target locus. The targeted region of *RAP1* was then Sanger

sequenced (CHUL sequencing platform, Univesité Laval, Québec City, Canada) to confirm the

presence of the mutation of interest. Heterozygous mutants were sporulated on solid media until

sporulation could be confirmed by microscopy using the same protocol previously described. The

plates were then replica plated on YPD+Nat media, and the pictures were taken after five days at

room temperature Supplementary image 2.

Evolutionary rate measurements and protein variant abundance

Evolutionary rates were calculated using the Rate4site software (Mayrose *et al.* 2004) using

multiple sequence alignments and phylogenies from PhylomeDB V4 (Huerta-Cepas *et al.* 2014)

as input and using the raw calculated rates as output. Variant data was compiled using data from

the   1002   Yeast   Genome   Project   (http://1002genomes.u-strasbg.fr/files/

allReferenceGenesWithSNPsAndIndelsInferred.tar.gz). Strain-specific protein coding sequence

were aligned to the S288c sequence using Fastx36 (Pearson *et al.* 1997) with the following

parameters: `fastx36 -p -s -VT10 -T 6 -m 10 -n -3 querymultifasta.fasta`

`ref_orf.db 12\> fasta_out`. Alignments were then parsed with a custom Python script to

identify variants. Variant abundance was measured as the number of strains in the dataset in

which a specific variant was found. If the coding sequence contained ambiguous nucleotides (ex.:

R or Y), separate coding sequences were generated for each possibility and each possible variant

was considered as a separate occurrence.

Analysis of the properties of stop codon generating gRNAs

To analyse the sequence and target properties of gRNA inducing the creation of stop codons,

data from multiple sources was compiled. For each target gene, length and chromosomal strand

was obtained from the Saccharomyces Genome Database using the Yeastmine query interface

(Cherry *et al.* 2012). Distance to centromere was obtained by calculating the minimal distance

between the start of the gene and one extremity of the centromere coordinates. RNA:DNA duplex

melting temperature of gRNA sequence with target genomic DNA was calculated using the

MeltingTemp module from Biopython (Cock *et al.* 2009), which uses values taken from Sugimoto

et al (Sugimoto *et al.* 1995). Correlation between gRNA/DNA duplex melting temperatures was

assessed using Spearman's rank correlation.

Variant effect prediction resources analysis and GO enrichment

All prediction data except the Envision scores were extracted from the aggregated data of the

Mutfunc database (Wagih *et al.* 2018). Precomputed values were downloaded directly from the

FTP server (http://ftp.ebi.ac.uk/pub/databases/mutfunc/mutfunc_v1/yeast/). This database

includes precomputed SIFT scores for 5498 yeast proteins, as well as predicted variant ddG

values based on protein structure (n=1057), homology models (n=1703) and protein-protein

interaction interfaces (n=1109). Mutations with $\Delta\Delta$G>1 considered destabilizing.

Precomputed values from Envision (Gray *et al.* 2018) were downloaded directly from the database

website (https://envision.gs.washington.edu/shiny/envision_new/, file yeast_predicted_2017-03-

12.csv). This file contained 34857830 mutation effect predictions spread across 4011 genes. The

distribution of Envision scores for the genes targeted in the experiment that are included in the

database are shown in (Figure S6).

Gene enrichments were performed using the PANTHER gene list analysis tool (Mi *et al.* 2019).

The list of genes for which 2 or more GNEs were detected was tested for enrichment against all

genes targeted by the library using Fisher's exact test and False Discovery Rate calculations. The

Gene Ontology datasets used were: GO molecular function complete, GO biological process

complete, and GO cellular component complete.

Acknowledgments

This work was supported by the Canadian Institutes of Health Research Foundation grant 387697

to CRL., as well as project grants 364920, 384483, a Frederick Banting and Charles Best graduate

scholarship and a Vanier graduate scholarship to P.C.D, by Université Laval via an André

Darveau Fellowship to P.C.D., the Fonds Québécois de Recherche en Santé via a Master's

training award to P.C.D. and the Japan Society for the Promotion of Science grant numbers

S15734 and S17161 to C.R.L. and N.Y. The authors thank Mathieu Hénault, Johan Hallin, and

Dan Yamamoto Evans for comments on the manuscript, as well as Maria Isabel Acosta Lopez

for assistance during the strain construction process.

Author contributions

682 PCD, AKD, NY and CRL designed research. PCD and AKD performed experiments. PCD and

683 MS generated NGS sequencing data. All data analysis was performed by PCD with input from

684 CRL. PCD and CRL wrote the manuscript with input from all authors.

Conflict of interest

686 None to declare

References

Albuquerque C. P., M. B. Smolka, S. H. Payne, V. Bafna, J. Eng, *et al.*, 2008 A multidimensional
chromatography technology for in-depth phosphoproteome analysis. Mol. Cell. Proteomics
7: 1389–96. https://doi.org/10.1074/mcp.M700468-MCP200

Bao Z., M. HamediRad, P. Xue, H. Xiao, I. Tasan, *et al.*, 2018 Genome-scale engineering of
Saccharomyces cerevisiae with single-nucleotide precision. Nat. Biotechnol.
https://doi.org/10.1038/nbt.4132

C. elegans Deletion Mutant Consortium T. C. elegans D. M., 2012 Large-Scale Screening for
Targeted Knockouts in the Caenorhabditis elegans Genome. G3;
Genes|Genomes|Genetics 2: 1415–1425. https://doi.org/10.1534/g3.112.003830

Casadaban M. J., and S. N. Cohen, 1980 Analysis of gene control signals by DNA fusion and
cloning in Escherichia coli. J. Mol. Biol. 138: 179–207. https://doi.org/10.1016/0022-
2836(80)90283-1

Cherry J. M., E. L. Hong, C. Amundsen, R. Balakrishnan, G. Binkley, *et al.*, 2012
Saccharomyces Genome Database: The genomics resource of budding yeast. Nucleic
Acids Res. https://doi.org/10.1093/nar/gkr1029

Cock P. J. A., T. Antao, J. T. Chang, B. A. Chapman, C. J. Cox, *et al.*, 2009 Biopython: Freely
available Python tools for computational molecular biology and bioinformatics.
Bioinformatics. https://doi.org/10.1093/bioinformatics/btp163

Copley R. R., and G. J. Barton, 1994 A Structural Analysis of Phosphate and Sulphate Binding
Sites in Proteins. J. Mol. Biol. 242: 321–329. https://doi.org/10.1006/jmbi.1994.1583

Dandage R., P. C. Després, N. Yachie, and C. R. Landry, 2019 beditor: A Computational
Workflow for Designing Libraries of Guide RNAs for CRISPR-Mediated Base Editing.
Genetics 212: 377–385. https://doi.org/10.1534/genetics.119.302089

DePristo M. A., D. M. Weinreich, and D. L. Hartl, 2005 Missense meanderings in sequence
space: a biophysical view of protein evolution. Nat. Rev. Genet. 6: 678–687.
https://doi.org/10.1038/nrg1672

Després P. C., A. K. Dubé, L. Nielly-Thibault, N. Yachie, and C. R. Landry, 2018 Double
Selection Enhances the Efficiency of Target-AID and Cas9-Based Genome Editing in
Yeast. G3 (Bethesda). g3.200461.2018. https://doi.org/10.1534/g3.118.200461

Dicarlo J. E., J. E. Norville, P. Mali, X. Rios, J. Aach, *et al.*, 2013 Genome engineering in
Saccharomyces cerevisiae using CRISPR-Cas systems. Nucleic Acids Res. 41: 4336–
4343. https://doi.org/10.1093/nar/gkt135

Doench J. G., E. Hartenian, D. B. Graham, Z. Tothova, M. Hegde, *et al.*, 2014 Rational design
of highly active sgRNAs for CRISPR-Cas9-mediated gene inactivation. Nat. Biotechnol. 32:
1262–1267. https://doi.org/10.1038/nbt.3026

Doench J. G., N. Fusi, M. Sullender, M. Hegde, E. W. Vaimberg, *et al.*, 2016 Optimized sgRNA
design to maximize activity and minimize off-target effects of CRISPR-Cas9. Nat.
Biotechnol. 34: 184–191. https://doi.org/10.1038/nbt.3437

Eriani G., M. Delarue, O. Poch, J. Gangloff, and D. Moras, 1990 Partition of tRNA synthetases
into two classes based on mutually exclusive sets of sequence motifs. Nature 347: 203–
206. https://doi.org/10.1038/347203a0

Fowler D. M., and S. Fields, 2014 Deep mutational scanning: a new style of protein science.
Nat. Methods 11: 801–7. https://doi.org/10.1038/nmeth.3027

Gaudelli N. M., A. C. Komor, H. A. Rees, M. S. Packer, A. H. Badran, *et al.*, 2017
Programmable base editing of A•T to G•C in genomic DNA without DNA cleavage. Nature
551: 464–471. https://doi.org/10.1038/nature24644

Giaever G., D. D. Shoemaker, T. W. Jones, H. Liang, E. A. Winzeler, *et al.*, 1999 Genomic
profiling of drug sensitivities via induced haploinsufficiency. Nat. Genet. 21: 278–83.
https://doi.org/10.1038/6791

Giaever G., A. M. Chu, L. Ni, C. Connelly, L. Riles, *et al.*, 2002 Functional profiling of the
Saccharomyces cerevisiae genome. Nature 418: 387–391.
https://doi.org/10.1038/nature00935

Gietz R. D., and R. H. Schiestl, 2007 High-efficiency yeast transformation using the LiAc/SS
carrier DNA/PEG method. Nat. Protoc. 2: 31–34. https://doi.org/10.1038/nprot.2007.13

Graham I. R., R. A. Haw, K. G. Spink, K. A. Halden, and A. Chambers, 1999 In vivo analysis of
functional regions within yeast Rap1p. Mol. Cell. Biol. 19: 7481–90.
https://doi.org/10.1128/mcb.19.11.7481

Grant T. D., J. R. Luft, J. R. Wolfley, M. E. Snell, H. Tsuruta, *et al.*, 2013 The structure of yeast
glutaminyl-tRNA synthetase and modeling of its interaction with tRNA. J. Mol. Biol. 425:
2480–2493. https://doi.org/10.1016/j.jmb.2013.03.043

Grate L., and M. Ares, 2002 Searching yeast intron data at Ares lab web site. Methods
Enzymol. https://doi.org/10.1016/S0076-6879(02)50975-7

Gray V. E., R. J. Hause, J. Luebeck, J. Shendure, and D. M. Fowler, 2018 Quantitative
Missense Variant Effect Prediction Using Large-Scale Mutagenesis Data. Cell Syst.
https://doi.org/10.1016/j.cels.2017.11.003

Haldimann A., M. K. Prahalad, S. L. Fisher, S. K. Kim, C. T. Walsh, *et al.*, 1996 Altered
recognition mutants of the response regulator PhoB: a new genetic strategy for studying
protein-protein interactions. Proc. Natl. Acad. Sci. U. S. A. 93: 14361–6.
https://doi.org/10.1073/pnas.93.25.14361

Hess G. T., L. Frésard, K. Han, C. H. Lee, A. Li, *et al.*, 2016 Directed evolution using dCas9-
targeted somatic hypermutation in mammalian cells. Nat. Methods.
https://doi.org/10.1038/nmeth.4038

Ho C. H., L. Magtanong, S. L. Barker, D. Gresham, S. Nishimura, *et al.*, 2009 A molecular
barcoded yeast ORF library enables mode-of-action analysis of bioactive compounds. Nat.
Biotechnol. 27: 369–377. https://doi.org/10.1038/nbt.1534

Holt L. J., B. B. Tuch, J. Villén, A. D. Johnson, S. P. Gygi, *et al.*, 2009 Global analysis of Cdk1
substrate phosphorylation sites provides insights into evolution. Science 325: 1682–6.
https://doi.org/10.1126/science.1172867

Huerta-Cepas J., S. Capella-Gutiérrez, L. P. Pryszcz, M. Marcet-Houben, and T. Gabaldón,
2014 PhylomeDB v4: Zooming into the plurality of evolutionary histories of a genome.
Nucleic Acids Res. https://doi.org/10.1093/nar/gkt1177

Huxley C., E. D. Green, and I. Dunham, 1990 Rapid assessment of S. cerevisiae mating type by
PCR. Trends Genet. 6: 236.

Jin S., Y. Zong, Q. Gao, Z. Zhu, Y. Wang, *et al.*, 2019 Cytosine, but not adenine, base editors
induce genome-wide off-target mutations in rice. Science eaaw7166.
https://doi.org/10.1126/science.aaw7166

Kim D., K. Lim, S. T. Kim, S. H. Yoon, K. Kim, *et al.*, 2017 Genome-wide target specificities of
CRISPR RNA-guided programmable deaminases. Nat. Biotechnol.
https://doi.org/10.1038/nbt.3852

Konig P., R. Giraldo, L. Chapman, and D. Rhodes, 1996 The crystal structure of the DNA-
binding domain of yeast RAP1 in complex with telomeric DNA. Cell 85: 125–36.
https://doi.org/10.1016/S0092-8674(00)81088-0

Landry C. R., E. D. Levy, and S. W. Michnick, 2009 Weak functional constraints on
phosphoproteomes. Trends Genet. 25: 193–7. https://doi.org/10.1016/j.tig.2009.03.003

Langmead B., C. Trapnell, M. Pop, and S. L. Salzberg, 2009 Ultrafast and memory-efficient
alignment of short DNA sequences to the human genome. Genome Biol. 10: R25.
https://doi.org/10.1186/gb-2009-10-3-r25

Ma Y., J. Zhang, W. Yin, Z. Zhang, Y. Song, *et al.*, 2016 Targeted AID-mediated mutagenesis
(TAM) enables efficient genomic diversification in mammalian cells. Nat. Methods 13:
1029–1035. https://doi.org/10.1038/nmeth.4027

Mayrose I., D. Graur, N. Ben-Tal, and T. Pupko, 2004 Comparison of site-specific rate-inference
methods for protein sequences: Empirical Bayesian methods are superior. Mol. Biol. Evol.

https://doi.org/10.1093/molbev/msh194

Mi H., A. Muruganujan, X. Huang, D. Ebert, C. Mills, *et al.*, 2019 Protocol Update for large-scale
genome and gene function analysis with the PANTHER classification system (v.14.0). Nat.
Protoc. 14: 703–721. https://doi.org/10.1038/s41596-019-0128-8

Michel A. H., R. Hatakeyama, P. Kimmig, M. Arter, M. Peter, *et al.*, 2017 Functional mapping of
yeast genomes by saturated transposition. Elife 6. https://doi.org/10.7554/eLife.23570

Ng P. C., and S. Henikoff, 2003 SIFT: Predicting amino acid changes that affect protein
function. Nucleic Acids Res.

Nishida K., T. Arazoe, N. Yachie, S. Banno, M. Kakimoto, *et al.*, 2016 Targeted nucleotide
editing using hybrid prokaryotic and vertebrate adaptive immune systems. Science (80-.).
353: 553–563. https://doi.org/10.1126/science.aaf8729

Nishimasu H., X. Shi, S. Ishiguro, L. Gao, S. Hirano, *et al.*, 2018 Engineered CRISPR-Cas9
nuclease with expanded targeting space. Science eaas9129.
https://doi.org/10.1126/science.aas9129

Pearson W. R., T. Wood, Z. Zhang, and W. Miller, 1997 Comparison of DNA Sequences with
Protein Sequences. Genomics 46: 24–36. https://doi.org/10.1006/geno.1997.4995

Qi L. S., M. H. Larson, L. A. Gilbert, J. A. Doudna, J. S. Weissman, *et al.*, 2013 Repurposing
CRISPR as an RNA-guided platform for sequence-specific control of gene expression. Cell
152: 1173–83. https://doi.org/10.1016/j.cell.2013.02.022

Qian W., D. Ma, C. Xiao, Z. Wang, and J. Zhang, 2012 The Genomic Landscape and
Evolutionary Resolution of Antagonistic Pleiotropy in Yeast. Cell Rep. 2: 1399–1410.
https://doi.org/10.1016/j.celrep.2012.09.017

Rees H. A., and D. R. Liu, 2018 Base editing: precision chemistry on the genome and
transcriptome of living cells. Nat. Rev. Genet. 19: 770–788. https://doi.org/10.1038/s41576-
018-0059-1

Reis A. M. C., W. K. Mills, I. Ramachandran, E. C. Friedberg, D. Thompson, *et al.*, 2012
Targeted detection of in vivo endogenous DNA base damage reveals preferential base
excision repair in the transcribed strand. Nucleic Acids Res. 40: 206–219.
https://doi.org/10.1093/nar/gkr704

Roy K. R., J. D. Smith, S. C. Vonesch, G. Lin, C. S. Tu, *et al.*, 2018 Multiplexed precision
genome editing with trackable genomic barcodes in yeast. Nat. Biotechnol.
https://doi.org/10.1038/nbt.4137

Ryan O. W., S. Poddar, and J. H. D. Cate, 2016 Crispr–cas9 genome engineering in
Saccharomyces cerevisiae cells. Cold Spring Harb. Protoc. 2016: 525–533.
https://doi.org/10.1101/pdb.prot086827

Sander J. D., and J. K. Joung, 2014 CRISPR-Cas systems for editing, regulating and targeting
genomes. Nat. Biotechnol. 32: 347–55. https://doi.org/10.1038/nbt.2842

Sanjana N. E., O. Shalem, and F. Zhang, 2014 Improved vectors and genome-wide libraries for
CRISPR screening. Nat. Methods 11: 783–784. https://doi.org/10.1038/nmeth.3047

Schmitt E., M. Panvert, S. Blanquet, and Y. Mechulam, 1995 Transition state stabilization by the
"high" motif of class I aminoacyl-tRNA synthetases: The case of Escherichia coli methionyl-

tRNA synthetase. Nucleic Acids Res. https://doi.org/10.1093/nar/23.23.4793

Schymkowitz J., J. Borg, F. Stricher, R. Nys, F. Rousseau, *et al.*, 2005 The FoldX web server:
an online force field. Nucleic Acids Res. 33: W382–W388.
https://doi.org/10.1093/nar/gki387

Shalem O., N. E. Sanjana, E. Hartenian, X. Shi, D. A. Scott, *et al.*, 2014 Genome-scale
CRISPR-Cas9 knockout screening in human cells. Science (80-.). 343: 84–87.
https://doi.org/10.1126/science.1247005

Sharon E., S. A. A. Chen, N. M. Khosla, J. D. Smith, J. K. Pritchard, *et al.*, 2018 Functional
Genetic Variants Revealed by Massively Parallel Precise Genome Editing. Cell.
https://doi.org/10.1016/j.cell.2018.08.057

Smith J. E., J. R. Alvarez-Dominguez, N. Kline, N. J. Huynh, S. Geisler, *et al.*, 2014 Translation
of Small Open Reading Frames within Unannotated RNA Transcripts in Saccharomyces
cerevisiae. Cell Rep. 7: 1858–1866. https://doi.org/10.1016/j.celrep.2014.05.023

Smith J. D., S. Suresh, U. Schlecht, M. Wu, O. Wagih, *et al.*, 2016 Quantitative CRISPR
interference screens in yeast identify chemical-genetic interactions and new rules for guide
RNA design. Genome Biol. 17: 45. https://doi.org/10.1186/s13059-016-0900-9

Sneath P. H., 1966 Relations between chemical structure and biological activity in peptides. J.
Theor. Biol. 12: 157–95.

Sugimoto N., S. Nakano, M. Katoh, A. Matsumura, H. Nakamuta, *et al.*, 1995 Thermodynamic
parameters to predict stability of RNA/DNA hybrid duplexes. Biochemistry 34: 11211–6.

Tarassov K., V. Messier, C. R. Landry, S. Radinovic, M. M. Serna Molina, *et al.*, 2008 An in vivo
map of the yeast protein interactome. Science 320: 1465–70.
https://doi.org/10.1126/science.1153878

Wagih O., M. Galardini, B. P. Busby, D. Memon, A. Typas, *et al.*, 2018 A resource of variant
effect predictions of single nucleotide variants in model organisms. Mol. Syst. Biol. 14:
e8430. https://doi.org/10.15252/MSB.20188430

Winzeler E. A., D. D. Shoemaker, A. Astromoff, H. Liang, K. Anderson, *et al.*, 1999 Functional
characterization of the S. cerevisiae genome by gene deletion and parallel analysis.
Science (80-.). 285: 901–906. https://doi.org/10.1126/science.285.5429.901

Wong N., W. Liu, and X. Wang, 2015 WU-CRISPR: characteristics of functional guide RNAs for
the CRISPR/Cas9 system. Genome Biol. 16: 218. https://doi.org/10.1186/S13059-015-
0784-0

Wu A. C. K., H. Patel, M. Chia, F. Moretto, D. Frith, *et al.*, 2018 Repression of Divergent
Noncoding Transcription by a Sequence-Specific Transcription Factor. Mol. Cell 72: 942-
954.e7. https://doi.org/10.1016/J.MOLCEL.2018.10.018

Yachie N., E. Petsalaki, J. C. Mellor, J. Weile, Y. Jacob, *et al.*, 2016 Pooled-matrix protein
interaction screens using Barcode Fusion Genetics. Mol. Syst. Biol. 12: 863.

Zhao L., Q. Yang, J. Zheng, X. Zhu, X. Hao, *et al.*, 2016 A genome-wide imaging-based
screening to identify genes involved in synphilin-1 inclusion formation in Saccharomyces
cerevisiae. Sci. Rep. 6: 30134. https://doi.org/10.1038/srep30134

Zuo E., Y. Sun, W. Wei, T. Yuan, W. Ying, *et al.*, 2019 Cytosine base editor generates

substantial off-target single-nucleotide variants in mouse embryos. Science (80-.).

eaav9973. https://doi.org/10.1126/SCIENCE.AAV9973

SUPPLEMENTARY MATERIAL: Supplementary Figures 1-11

Figure S1: A gRNA library for the systematic mutagenesis of yeast essential genes and other targets of interest. A) Number of genes targeted by the gRNA library for the different target classes. B) Total number of gRNAs targeting genes in the different target classes. C) Distribution of number gRNAs for each gene targeted in the different classes. D) Distribution of minimal (light grey) and median (dark grey) pairwise sequence distance between all gRNA sequences in the library.

Figure S2: Experimental workflow for Target-AID mutagenesis and co-selection. The mutagenesis method closely follows the base editing protocol previously described (Després *et al.* 2018). After a pooled transformation step, cells were scraped and splitted into two replicates for pre-cultures. After each step of the protocol, plasmid DNA was extracted from a cell sample and used to amplify and sequence the gRNA pool. The red stars indicate time points used for fitness effects analysis: read counts after galactose induction were used as T0 and were compared with read counts after two rounds of competition. The mock induction steps mimics the induction conditions but galactose in the media is replaced by glucose. This prevents the editing enzyme from being expressed because glucose represses the GAL pathway. After canavanine co-selection, cells go through two competition rounds in synthetic media were selective pressure for the Target-AID bearing plasmid is lost. The entire experiment was completed within less than 25 generations after galactose induction, limiting the impact of compensatory and spontaneous mutations.

Figure S3: Read abundance rank order is strongly correlated between replicates. For
each time point, Spearman rank correlation of gRNA $\log_{10}$ read abundance after basic filtering is
shown. The minimal read count after galactose induction, which served as the principal filtering
criteria, is shown on the galactose subpanel.

Figure S4: Barcode abundance correlation clusters different experimental steps of the screen. Pairwise Spearman rank correlation of barcode counts was used to cluster the libraries obtained at the different time points described in Figure S2. The lower level of correlation between the galactose induction and mock induction timepoints compared to other associated steps could reflect higher stochasticity in growth caused by cell to cell variation in the metabolic switch from glycerol to sugars as the main carbon source as well as editing in the case of the galactose timepoint.

**Figure S5 Plasmid-based confirmation workflow by complementation test and
evolutionary information on *GLN4*.** A) Detailed protocols for the different steps are presented
in the methods. First, directed mutagenesis is used to introduce the mutation of interest (shown
in red) in the MoBY collection plasmid of the targeted gene (YFG). This vector is then
transformed into the heterozygous collection deletion strain (BY4743, *MATa/α his3Δ1/his3Δ1
leu2Δ0/leu2Δ0 LYS2/lys2Δ0 met15Δ0/MET15 ura3Δ0/ura3Δ0*) of the gene of interest. The
transformants are sporulated and their tetrads are dissected. If the mutated allele carried by the
plasmid cannot complement the gene deletion, then only the two progenies bearing the wild-
type copies will be viable. B) Protein variant frequency among 1000 yeast isolates (black dots)
and residue evolutionary rate across species (blue line) for *GLN4*. The target site for the most
deleterious GNE is highlighted by a red line and other GNE target sites are shown as grey lines.

Figure S6 gRNA predicted mutation coverage for Mutfunc and Envision data. Mutfunc integrates both the SIFT prediction scores and FoldX (Schymkowitz *et al.* 2005), $\Delta\Delta$G predictions for solved protein structures, homology models, and protein-protein interaction interfaces. gRNAs which do not generate missense mutations were included in the calculations. A) Coverage for the SIFT and Envision variant effect predictors for the four most probable single mutants created by gRNAs detected in the experiment. B) Coverage for $\Delta\Delta$G predictions for solved protein structures, homology models, and protein-protein interaction interfaces for the four most probable single mutants created by gRNAs detected in the experiment. C) Distribution of Envision scores across all sites in the database for all proteins targeted by the set of gRNAs detected in the screen (n=7,556,573). The median score is shown as a dotted black line.

Figure S7 GNE and non-significant gRNA effect prediction distributions. A) Envision score
distributions for the four most probable mutations induced by GNEs (blue) and NSGs (red). Welch's
t-test p-values for comparisons: 5.00×10^{-6}, 0.002, 0.007, 7.75×10^{-5}. B) Predicted folding energy
variation ($\Delta\Delta G$) of GNE and NSG induced protein mutants compared to the wild-type structure
based on resolved protein structure. Welch's t-test p-values for comparisons: 0.016, 0.441, 0.195,
0.689. C) Predicted folding energy variation ($\Delta\Delta G$) of GNE and NSG induced protein mutants
compared to the wild-type structure based on homology models of protein structure. Welch's t-
test p-values for comparisons: 0.0001, 0.006, 0.148, 0.007. D) Binding energy variation ($\Delta\Delta G$) of
GNE and NSG induced mutant protein-protein interfaces compared to the wild-type based on a
resolved structure on the interface. Welch's t-test p-values for comparisons: 0.285, 0.303, 0.033,
0.95.

Figure S8 Fitness affecting variant by CRISPR knock in confirmation workflow. Detailed
protocols for the different steps are presented in the methods. Starting from the wild-type
laboratory strain BY4741, the gene of interest (*YFG*, blue) is first tagged with a modified version
of the DHFR F[1,2] cassette (dark gray and green). The tagged strain is then crossed with a MAT*α*
strain (Y8205) to create a heterozygous diploid. A *URA3* deletion cassette (black) that recombines
with the *YFG* upstream sequence and the start of the mDHFR fragment is then used to generate
a heterozygous KO strain. In parallel, genomic DNA is extracted from the tagged haploid strain.
This DNA is then used as a template to amplify two fragments of *YFG* bearing the mutation of
interest (shown in red) using a set of overhanging primers. The two fragments are then combined
by fusion PCR to obtain the donor DNA used in the next step. Using a modified Cas9 vector (Ryan
et al. 2016) that expresses a gRNA targeting the *URA3* cassette, the mutated allele is introduced
at the KO locus to create a heterozygous mutant strain. The diploid cells can then be sporulated,
and tetrad dissection allows observation of any phenotype linked with the mutation of interest.

Figure S9 Properties influencing stop codon GNEs are generalizable to non-stop codon generating GNEs. A) Cumulative z-score density for gRNAs that do not generate stop codons depending on the number of mutable sites. A higher rate of GNE is observed for gRNAs which can lead to the editing of multiple nucleotides (Two-sample Kolmogorov-Smirnov test, p=3.91x10^{-29}). The significance threshold is shown as a black dotted line. B) Cumulative z-score density for NSGs on orf target strand. gRNAs targeting the non-coding strand of the ORF have a higher likelihood of being GNEs (Two-sample Kolmogorov-Smirnov test, p=1.87x10^{-16}). C) gRNA z-score cumulative density for both SGGs and non-SGGs grouped by the chromosomal strand they target. In SGGs, the target strand does not impact z-score distributions (Two-sample Kolmogorov-Smirnov test, p=0.753) and GNE proportions (Fisher's exact test, p=0.149). For non-SGGs, the chromosomal strand has a small influence on z-score distributions (Two-sample Kolmogorov-Smirnov test, p=0.035) and GNE proportions (Fisher's exact test, p=0.002) D) Distributions of modeled RNA/DNA duplex melting temperature for all non-SGGs generating gRNAs, the NSG subset, and the GNEs subset. P-values were calculated using the two-sample Kolmogorov-Smirnov test. F) Non-SGGs GNE enrichment compared to the expected GNE ratio for different melting temperature ranges. F) gRNA/DNA duplex melting temperature as a function of gRNA GC content for all gRNAs for which fitness effects were measured. The higher and lower efficiency thresholds are based on the enrichments shown in panel E and Figure 4F.

Figure S10 gRNA/DNA duplex melting temperature is not linked to systematic sequencing biases. Spearman rank correlation between replicate averaged read count and predicted gRNA/DNA duplex melting temperature is shown across timepoints. The minimal read count after galactose induction, which served as a filtering criterion, is shown on the galactose subpanels. gRNAs for which no reads were detected in one of the time points were included when computing the correlation but are not shown on the graphs because of log scaling.

Figure S11 GNE density is independent of target nucleotide position bias. A) In SGGs, GNE and NSG target sites that are evenly distributed across the target genes, and GNEs do not show any bias (Two-sample Kolmogorov-Smirnov). B) Non-SGG GNEs do not show any positional bias. C) A significant but small negative correlation is observed between gRNA target relative position and GC content of SGGs (Spearman's rank correlation). The very small observed effect coupled with the absence of position bias suggests that relative target position bias does not drive the link between GC content and gRNA efficiency. D) Similarly, a small but significant but small negative correlation is also observed between gRNA relative position and GC content for non-SGGs (Spearman's rank correlation).

1006