1
2
3
4
5 **Single EGF mutants unravel the mechanism for stabilization of Epidermal Growth Factor**
6 **Receptor (EGFR) system**
7
8
9

10 Stefano Pascarelli[1], Dalmira Merzhakupova[1], Gen-Ichiro Uechi[1], Paola Laurino[1*]
11
12
13
14 [1] Okinawa Institute of Science and Technology Graduate University, 1919-1 Tancha, Onna-son, Okinawa
15 904-0495 Japan.
16
17
18 * Corresponding author
19 E-mail: paola.laurino@oist.jp
20 **Running title**: EGF mutants elucidate receptor transduction

21
22 **Keywords**: epidermal growth factor (EGF), epidermal growth factor receptor (EGFR), protein
23 phosphorylation, receptor tyrosine kinase, mutant, molecular dynamics, dimerization, protein-protein
24 interaction
25

Abstract

The Epidermal Growth Factor Receptor (EGFR) is a membrane-anchored tyrosine kinase that is able to respond to multiple extra-cellular stimuli in a selective way. Previous studies have pointed out that the modularity of this system could be carried out by ligand-induced differences in the stability of the dimerized receptor. Nevertheless, nobody has ever explored this hypothesis by observing the effects of single-mutant ligands so far. Herein, we identified residues responsible for inducing functional divergence among paralog ligands using a newly developed approach. Then, we mutated these positions and assessed the mutants' effects on the receptor by employing a combination of molecular dynamics and experimental techniques. Although having comparable binding affinities for EGFR to the wild type, the EGF mutants induced different responses at both the receptor and the cellular level. This study shows for the first time that a single mutation in EGFR ligand is enough for changing the outcome of the activation pathway at the cellular level. These results also support the theory of biased signaling in the tyrosine kinase receptor system and show a promising new way to study ligand-receptor interactions.

Introduction

The Epidermal Growth Factor (EGF)-like domain ligand-receptor signaling system is involved in many biological events in multicellular organisms (1). This system is constituted by one receptor (EGFR) and seven distinct peptide ligands (EGF, Epidermal Growth Factor; HBEGF, Heparin-Binding Epidermal Growth Factor; EPGN, Epigen; BTC, Betacellulin; EREG, Epiregulin; AREG, Amphiregulin). Upon binding to the receptor, these ligands can activate multiple intracellular downstream pathways through a network of intramolecular interactions with several feedback loops (2). For all ligands, binding induces a transition in the EGFR from monomer or inactive dimer to an active dimer state (3,4). However, different ligands are able to promote divergent outcomes, even at saturating concentrations; thus, the mechanism responsible of the modular downstream pathway activation is independent of ligand affinity or potency, and likely encompasses intrinsic effects (5). It is well known that the EGFR system plays a key role in cancer development. In particular, some studies have shown that overexpression of EGFR or its ligands may induce different types of cancer (6). A better understanding of the interaction between EGFR and its ligands could lead to the development of targeted therapies (7).

Protein-protein interactions (PPIs), such as that between EGFR and its ligands, are well-studied examples of molecular co-evolution in biological systems. These interactions are sometimes defined by one part (receptor) that binds several counterparts (ligands). In these instances, receptor and ligands experience different selective constraints, where receptors tend to evolve more slowly due to the necessity of binding multiple ligands (8). Furthermore, paralogs, proteins related by a duplication event, are less likely to retain the same function as orthologs, proteins related by a speciation event (9). Then, the paralogous ligands rather than the receptor are a good candidate to test the functional divergence in the biased signaling system. However, this approach is susceptible to indirect factors, such as different ligand binding selectivity, thus resulting in the activation of unrelated pathways. In this work, we decided to test single mutants of one ligand, EGF, by identifying and modifying sites responsible for a divergent function among paralog ligands. Recent studies have shown that some EGF residues like Arg-41 and Leu-47 are highly conserved and important for high binding affinity to EGFR (10). Another study highlighted Tyr-13, Leu-15 and His-16 in EGF as essential for downstream activity of ErbB1 (11). These outcomes were based on structural analyses of ligands and experimental validation. While bioinformatic tools such as contact prediction (12) or molecular dynamics (13) can give a good overall picture of ligand-receptor interactions, the contribution of single ligand residues to the modularity of the system still remains unclear.

Ligand-induced selective activation of downstream pathway has been observed in G-protein coupled receptors, a phenomenon known as "biased signaling" (14,15). A recent report hints that this mechanism might also take place in the EGFR tyrosine kinase (16). Initially, the main contributor to the modularity of the system was thought to be the affinity of the ligands to the receptor. However, the discovery of the ligand EPGN, which induces a potent mitogen effect despite low binding affinity (17), and multiple cell-line studies (5) have changed this perspective. A plausible explanation was initially formulated by Wilson *et al.*, where different EGFR ligands induce different receptor dimer stabilities, altering the phosphorylation dynamics(5); Later, new experimental evidence supported this theory by observing a transient activation in stable receptor dimers induced by EGF; whereas the activation of the weak receptor dimer induced by EPGN or EREG persisted for a longer time (18). A difference in dimer stabilities could result in alterations of the receptor oligomerization state, previously shown as a determinant of the intracellular kinase phosphorylation pattern (19). Now, the open question is how these ligands induce different dimer stabilities. Interactions between the interface of the ligand and EGFR have been shown to influence the stability of the dimerized receptor (20), a factor that is related to ligand-specific signaling bias (18,21). It is known that the EGFR dimer can be observed in a symmetrical, "flush" conformation or an asymmetrical, "staggered" conformation (22) depending on the presence/absence of the membrane, glycosylation, and the number of bound ligands (23). Whether and how this plays a role in biased signaling is still unknown.

Here we show how single amino acid substitutions on the ligands effect the biased signaling in the EGF-EGFR ligand-receptor system. We developed a new bioinformatic tool to analyze the co-evolution of the ligand-receptor pair and identify candidate function-altering mutations. The identified mutants induced an altered phosphorylation dynamics and different cellular phenotype. This is the first study to explain differences in biased signaling of EGFR using single-residue EGF mutants. Furthermore, our co-evolutionary analysis can be applied readily to other ligand-receptor interactions.

Results

DIRpred
Firstly, we developed a method for predicting residues that are likely to be responsible for functional divergence among paralogs sharing a common interactor (referred as ligands and receptor from now on). We called the method DIRpred (Divergence Inducing Residues prediction). Our approach combines residue-specific conservation measures to identify positions that are conserved among orthologs while diverging among paralogs. The DIRpred analysis is based on the assumption that conservation of a residue in orthologs of a specific ligand shows whether a residue is important for either structural or functional reasons, while conservation of a residue among paralogous ligands denotes the importance of a residue for interaction with the common interactor (the main shared property of all ligands). Thus, residues that are highly conserved in orthologs but not in paralogs of a ligand are likely related to the ligand's specific function. Unlike other existing methods for prediction of specificity determining residues (a review can be found in (24)), we included inter-protein coevolution measures in order to narrow down those residues that are responsible for a specific interaction. The DIRpred score is calculated as the sum of the four components (I: orthologs conservation, II: complement of the paralog's conservation, III: ligand-receptor co-evolution, IV: complement of ligand internal co-evolution). Optionally, the analysis can be conducted using a structural alignment of the paralogs (MSTA) instead of the sequence alignment (MSA). An implementation of the DIRpred analysis was done using Python. The pipeline accepts multiple sequence or structure alignments and a reference protein to conduct the analysis. The output consists of a single tabular file containing the four individual scores and the combined prediction score for each protein site, and a plotted recap of the results (Figure S9).

DIRpred analysis of EGF

We applied the DIRpred algorithm to predict the residues in EGF that induce functional divergence among
paralogs (Figure 2). Since the analysis requires prior knowledge of paralogs and orthologs of the target
gene, we first performed a phylogenetic analysis to confirm that the reported paralogous ligands of EGF
are monophyletic (Figure S1). The same tree was used in the statistical validation of the individual DIRpred
scores. A random 53 amino acid long sequence was evolved 100 times on the EGFR ligands phylogenetic
tree using Pyvolve(25). Assuming that the distribution of DIRpred scores will be normal, we estimated the
probability that a site without functional constraints would have the same or higher than observed score (P-
value) (26). We included the results of the statistical analysis to the output of DIRpred for EGF (Figure S10,
Table S1). The analysis highlighted residue Asn-32, Asp-46, Lys-48 and Trp-50 as potential candidates for
paralog functional divergence. Asn-32 and Lys-48 show a very small conservation in both sequence and
structure alignments resulting in a high partial score (II). Asp-46 has a relatively high coevolution with the
receptor (III), while Trp-50 has high scores overall (Figure S2).
138
Three out of the four positions were mentioned in previous reports, however none of them was individually
mutated. The tryptophan in position 50 is a strong outlier in our bioinformatics analysis, along with Trp-49
(Figure 2). Their score is high even when using conservation measures that do not take amino acid type
change into account (data not shown). However, Trp-50 was a better candidate for testing because of its
outward facing position, while Trp-49 is involved in buried protein contacts (27). Trp-49 and Trp-50 could
be responsible in facilitating the interaction of EGF with the membrane phospholipids, as it happens for
Pro-7 and Leu-8 (28); Trp-49 and Trp-50 are not burying inside the bilayer when EGF and a membrane are
in solution alone (29), though this might be different when in complex with the receptor. For example, this
could be achieved by Trp-50 through the formation of a helix, clustering together with Val-34 and Arg-45
around the conserved Leu-47 (30). Mutation N32R is on the interface between ligand and receptor. The
slightly higher affinity is probably due to the presence of the guanidinium group of R which is positively
charged and could interact with Gln-16 of EGFR ECD (Figure S8). Interestingly, mutations of the
corresponding position in chicken TGFA were able to alter the mitogenicity without strongly affecting the
binding affinity (31). While no previous literature reported about Asp-46 before, Lys-48 was found in two
mutants that showed higher affinity (32).
154
Next, we manually chose which amino acid to introduce on the four sites depending on several factors. The
main contribution to the decision was given by the paralogs alignment, while also considering the amino
acid type and the ligand functional divergence. The paralogs were divided into two groups based on their
kinetics parameters of interaction with the receptor. After that, we selected an amino acid that infers a
significant change in biochemical properties and that is found in the paralogs group without EGF. When
multiple choices were possible, priority was given to EPGN or EREG, since these two ligands are the ones
observed to induce a biased signaling in Freed *et al* (18). The four designed EGF mutants with a single
amino-acid substitution that were selected for functional characterization are N32R, D46T, K48T and
W50Y.
164
Biochemical properties of the EGF mutants
To determine the integrity of the secondary structure of the mutated ligands and the functional effects of
these amino acid substitutions, we first performed *in-vitro* analysis. Initially, Circular Dichroism (CD)
spectroscopy was used to confirm that the secondary structure of the mutants was maintained (Figure S3).
The β-sheet content of the EGF mutants (ranging from 0.41 to 0.54 %) was not substantially different than
the WT EGF (0.44 %) while the other secondary structure varies. It is not surprising that only the beta
sheet content can be detected by CD considering that EGF is a peptide (53 amino acids) constituted by
two β-sheets connected by a loop and the regions at the C- and N- terminus are flanking. Then, the K_d of
each mutant for the soluble extracellular domain of EGFR was determined by Isothermal Titration
Calorimetry (ITC). The binding affinity of the mutants D46T, K48T, and W50Y, respectively 1.12 µM,
1.62 µM, and 1.86 µM, were not significantly different than the calculated K_d value of WT EGF, 1.28 µM

(Figure 3). Only N32R has 2-fold higher affinity for EGFR compared to WT EGF (Figure 3C). Thus, the
mutations do not appear to affect the ligand secondary structure and the binding affinity to the receptor.
178
Biological effects of the EGF mutants
Next, we sampled the effect of the EGF single mutants on EGFR expression and phosphorylation in A431
cells. A431 is a human epidermoid carcinoma cell line that overexpresses EGFR. For this reason, we
postulated that any changes at the receptor level would be amplified. We measured the amount of total and
phosphorylated EGFR protein at multiple timesteps by Western Blot (WB) after treating with saturating
concentrations (100 nM) of WT or mutant ligands (Figure 4). Remarkably, D46T mutant showed a reduced
level of EGFR phosphorylation (pEGFR) up to 30 minutes after treatment (Figure 4A). Meanwhile, the
pEGFR bands for K48T and W50Y samples appear slightly stronger than the WT in the first three timesteps,
however the difference is not significant. Treatment with the ligands also caused a reduction in the total
amount of EGFR 1 h and 6 h after the dosage compared to the control (Figure 4B). However, the reduction
of D46T sample did not appear as marked as the other samples. This result shows an inverse relationship
between receptor activation and receptor quantity, a fact that could be explained by the biased signaling
theory. At later time points, the effects converge back to the same range of the No Ligand (NL) control
sample (Figure S4). Thus, while the EGF mutants showed comparable binding affinities to the receptor, at
least one of the mutants showed a different phosphorylation pattern compared to the WT ligand.
Furthermore, the observed differences in the total amount of EGFR protein indicate a possible alteration in
the membrane expression or recycling of the receptor.
196
In order to observe the effects of the mutants on the cellular phenotype, we performed a cell growth assay
using an IncuCyte® live-cell analysis system on two fibroblast cell lines (Bj5-tα and Albino Swiss mouse
3T3). With the same platform, we also performed an apoptotic assay on A431 cells by measuring the
reduction in cell population, using the Annexin V green reagent to confirm the induction of apoptosis. After
one day of initial incubation, we subjected the cells to 1, 10 or 100 nM concentration of growth factors.
Treating Bj5-tα fibroblasts with 100 nM WT EGF resulted in a reduction in the proliferation rate compared
to the control. The reduction in proliferation was accompanied by a change in cellular morphology that
could be a signal of differentiation (Supplementary data 1). However, treatment with any of the mutants
decreased the ability of the ligand to suppress growth (Figure 5A). The D46T and K48T mutants differed
most from the WT (Figure S5A). We observed a similar trend in the Albino Swiss mouse 3T3 cell line
(Figure S5B). In both cell lines, the effects are concentration dependent (Figure S6). Next, we further tested
the mutants in an apoptosis assay using the A431 cell lines. This cell line is known to exhibit apoptosis
when treated with high (>10 nM) concentrations of the EGF peptide (34). The WT showed the highest
decrease in cell population 48h after treating with 100 nM WT, while the four mutants had an intermediate
response between WT and the control. Cells treated with the W50Y mutant have the closest level to the
WT (Figure 5B). In WT EGF cells, we observed evidence of apoptosis in cell's globular processes and
increased ratio of Annexin to confluence signal (Figure S11). Meanwhile, D46T and K48T treated cells
displayed signs of cellular differentiation (Figure 5C). Significantly, these results show that single amino
acid changes in the EGF ligand display differential effects on the EGFR transduction mechanism.
216
The observation that the four mutations in EGF alter signal transduction without disrupting any contacts
between the ligand and the receptor can be explained by the "loss of symmetry" model of EGFR signaling
(18). Freed *et al.* observed how stable symmetrical dimers show a short-lived nature of signaling, while
asymmetric dimers conduct a sustained activation, lasting longer than 24 h. Such a model provides an
explanation on how a low level of phosphorylation after EGF treatment causes modifications in the
apoptotic behavior observed in the cellular growth experiments. Compared with WT EGF, the D46T variant
show a constantly lower phosphorylation signal, possibly through the formation of a less stable, asymmetric
dimer. Thus, the observed effects of the EGF mutations on EGFR phosphorylation and dimer stability are
consistent with the "loss of symmetry" theory. However, the fact that some mutants have a similar

phosphorylation pattern as the WT EGF but show different cellular phenotypes suggests that there might
be other factors at play.
228
229
Molecular dynamics
To understand how EGF single mutants affect the receptor signaling transduction, we performed full
atomistic molecular dynamic (MD) simulations of the extracellular EGFR in complex with WT or mutant
ligands (100 ns each). We modeled the receptor starting from the asymmetric, "unstable" conformation of
5WB7 (22). In this way, we expected to observe a fast rearrangement for those simulations where a ligand
more favorably induce a stable dimer. In previous literature, an unstable conformation was observed when
removing one of the EGF ligands (23). We also performed one simulation of the EGFR dimer in complex
with only one WT ligand as a comparison (1ligEGF). The MD simulations quickly converged to a stable
RMSD (Figure 6A). For each simulation, we calculated the number of H-bonds between the two receptor
dimers, and between ligand and receptor. In the WT EGF simulation, the EGFR dimer had an average of
15 H-bonds, higher than any other simulation (Figure 6D). However, the number of bonds between receptor
and ligand was not altered (Figure 6E). In addition, during the course of our simulations we also noted
differences in the RMSF specifically located at the dimerization arm domain (Figure 6B). The
conformational space sampled by the dimerization arm of K48T simulation was much wider than the WT
simulation (Figure 6C). To analyze the temporal distribution of these motion, we measured the distance
between Pro-272 and Gly-288 of different EGFR monomers. This distance was chosen because it was able
to discriminate between the EGFR dimer in complex with EREG (PDB ID 5WB7) and EGF (PDB ID
1IVO). In 5WB7, one of the two distances is much bigger (~ 1.10 nm vs 0.4 nm) compared to 1IVO (Figure
6F). In our simulations, we observe WT showing a sharp peak at 1 nm, while the mutants and 1ligEGF have
a secondary peak at higher distances (Figure 6G). Thus, the mutations in EGF appear to affect the stability
of the EGFR dimer without affecting the stability of the EGF-EGFR interaction.
251
252
253 **Discussion**
254
The prediction of functional residues is a well-developed field (35), where conservation measures are
considered a key factor to rely on. Tools such as ConSurf (36) and the ET-like methods (37) are able to
identify slowly evolving positions that are involved in folding, interaction, or catalytic activity (35).
However, the specific reason why a residue is conserved often remains unclear. In this work, we show a
new method to identify residues that affects specific functions in a system of interest. Our approach
combines a conservation score calculated from the structural alignment of paralogs and among orthologs,
with an intra- and inter-molecular co-evolution score with previously known interactors. Conservation and
coevolution give a complementary signal, thus improving the overall predictive performance (38). The
coevolution score was introduced to DIRpred to highlight residues that are directly responsible of an
interaction with the receptor, at the expenses of those that interact intramolecularly within the ligand. While
the first part efficiently identifies Asn-32 and Lys-48 as putative interactors, the second part does not
properly give a penalty to residue Tyr-29. The interaction of this position with His-10 is also conserved in
EGF but with reversed positioning (Figure S7), resulting in a low paralog conservation (then high
contribution to DIRpred score). While it is still possible to optimize the co-evolution scoring function,
integrating conservation and coevolution measures is a promising way to recall information of specific
functions involving protein-protein interactions.
271
With the exception of N32R, all mutant ligands do not change in binding affinity to EGFR of compared to
the wild type ligand. However, the mutants showed a different cellular effect on Bj-5ta cells (Figure 5). A
delayed proliferation response compared to control might seem counterintuitive for fibroblast (39). Though,
Bj-5ta hTERT immortalized fibroblasts show features that distinguish them from *in vivo* fibroblast, like
differential gene expression and epiregulin-dependent proliferation (40). A decrease in cell proliferation

after EGF treatment could be the result of competition between the endogenously synthesized epiregulin
and EGF, thus altering the balance between proliferation and differentiation.
A431 cells constitutively express EGFR at high levels. Treatment with EGF has been observed to promote
STAT1 dependent apoptosis (41). This pathway is dependent on the internalization of the ligand-receptor
complex by the endocytosis process, a key factor in the ligand-induced biased signaling (16). In our
experiments, A431 cells treated with mutant EGF ligands show a higher growth rate, therefore a decreased
rate of apoptosis compared to WT. Given the setting of the experiment, modularity in the endocytosis
pathway is a straightforward explanation to the observed differences in growth rate and in protein levels
and phosphorylation. Interestingly, D46T and K48T treated cells showed an increase in long and thin
cellular processes in all replicates (Figure 5). This mutant was also observed with a differences in the EGFR
phosphorylation compared to WT (Figure 4). A possible explanation could be the EGF-induced tubular
formation, an alternative EGF-induced pathway reported in intestinal epithelial cells (42). In both the
cellular growth experiments, the growth rate of the mutants is intermediate between the control and EGF,
and concentration dependent. This phenomenon, and the cell-line specific response of growth hormones is
consistent with previous literature reports (43).
292
The analysis of the fluctuations in the dimerization arm reveals an underlying bias in ligand-induced
receptor dimerization, originally not visible from the static images of structure comparison. K48T is the
mutant that induce the biggest deviation from the WT. Although, all mutants show a transition. While giving
an initial outlook on the effect of the mutants, our MD simulations do not take into account other factors
that could be important in the mechanism of action of EGFR. Differential multimerization (44),
oligomerization (19), receptor glycosylation, and the interaction with the membrane (23) are factors where
the underlying bias induced by the mutant ligands could also have an effect in.
300
In this work, we showed how a single mutation of EGF is able to alter the specific functional relationship
with the receptor. For functional divergence to arise, it could take as little as mutating 15% of the sequence
(45). However, the EGFR ligands divergence date back to the vertebrate ancestor of R1/R2 whole genome
duplication, up to 500 million years ago (46), as hinted by the low (~25%) sequence identity. From our
results, the sequence distance does not reflect the distance in function. In fact, the functional divergence of
EGF was altered with just a single targeted mutation.
307
In conclusion our data suggests that a single mutant ligand induces a conformational change of the receptor
that then affects receptor dimer stability, with plausible effects on phosphorylation level and downstream
pathway activation (47). This shows how the persistence of biased signaling in EGFR is in an unstable
equilibrium, where the observed conservation of diverging sites among paralogs is naturally reinforced to
maintain the functional divergence. To determine whether a short distance in function space among paralogs
is a consequence or a necessity for living systems, further studies will be required.
314
315
316 **Experimental procedures**
317
318 **Sequence and structure analysis**
319 The sequences of EGFR ligands and the multiple sequence alignment of EGF orthologs were obtained from
320 the Ensembl database (48). Multiple sequence alignment (MSA) of all ligands was performed with MAFFT
321 software (49). X-ray structures were obtained from the PDB database (50). Structural alignments were
322 created using Chimera (51).
323
Phylogenetic analysis
From the multiple sequence alignment of EGF from different species, nearly identical sequences were
removed. The *Drosophila melanogaster* EGF sequence was added as an outgroup in the EGF phylogeny,
while *Caenorhabditis elegans* EGF was used as outgroup for the ligand phylogeny. MSA and phylogenetic

tree images were created using Unipro UGENE software (52). Three phylogenetic trees were made using
the Neighbor Joining (NJ), Maximum Likelihood (ML), and Bayesian (MrB) methods in IQTREE (53),
using ModelFinder to scan for the best-fit evolutionary model and parameters (54).
331
Divergence Inducing Residue PREDiction
We identified sites in EGF responsible for functional divergence using a method that combines evolutionary
and co-evolutionary data, called DIRpred (Divergence Inducing Residue PREDiction). The DIRpred
scoring function combines four components to evaluate each residue: I) The combined conservation scores
in the ortholog alignments. This score is calculated by averaging the conservation of a reference ligand site
with the conservation in the respective positions of the other ligands' orthologs alignments; II) The
complementary value $(1 - x)$ of the conservation score in the paralogs alignment. This score can be obtained
either from sequence alignment (paralogs MSA) or structural alignment (paralogs MSTA). III) The highest
co-evolution score between a reference ligand site and all of the receptor sites. IV) The complementary
value $(1 - x)$ of the highest co-evolution score between a reference ligand site and all of the other ligand
sites in the joint orthologs alignment of all ligands (Figure 7).
343
Conservation scores were calculated using three different formulations: 1) IDENTITY, 2) BLOSUM, and
3) JSDw. The IDENTITY score measures the frequency of appearance of a reference residue in the MSA.
The BLOSUM score takes the amino acid substitution frequency into account using the BLOSUM62 matrix,
normalized by the maximum and minimum score for the BLOSUM matrix. The JSDw score is based on
Jensen Shannon divergence with a window of residues (55). The fasta sequences were imported using
BioPython package, while analysis and plots were performed with Python3 package SEABORN. EGF
index positions from the paralogs MSA and MSTA were respectively used to align the MSA and MSTA
based scores. A schematic representation of how the four scores were obtained is showed in Figure S9.
The code used in the analysis of the DIRpred score and plots and the data used in this paper are shared on
Github: https://github.com/oist/DIRpred.
354
Statistical validation of EGF DIRpred
The statistical analysis of EGF DIRpred scores was done as in Mirny and Gelfand (56) with some
modifications. The key point is that a null-hypothesis dataset should take into account the higher similarity
found between orthologs, rather than between paralogs. Such a dataset was obtained through a simulated
evolution performed using the Python package Pyvolve (25). The sequences in the output dataset were
required to have a relationship akin to those of the EGF homologs. Using the WAG model (57), a random
53 amino acid-long protein sequence was evolved from the root of the EGF homologs tree until each leaf
node, simulating a scenario of neutral evolution. Then, the sequences were divided in orthologs and
paralogs using the original tree classification. The output was used as input for the DIRpred pipeline to
obtain four partial scores for each site. After 100 times repetition, the partial scores were gathered together
to form four background distributions. Through the assumption of normality, it was possible to estimate the
probability of a site to have a higher or equal score as a randomly evolved site, i.e., without functional
constrains (26). The scores were considered significant only on the side where they contribute positively to
the total DIRpred score, for example, when there is a lower level of paralogs conservation than expected.
For this reason, only one side of the normal distribution is used for the calculation of the P-value. (FIGS10).
In this way, it was possible to isolate the highly conserved cysteine positions, that usually get a misleading
high overall DIRpred score. About the relative contributions of individual scores, an equal weighting
system was preferred over the use of arbitrary values, which might not be always optimal. However, the
pipeline allows the user to provide his own weighting system that might be more optimized for the user's
study subject.
375
Selection of the mutations
Along with the DIRpred score, the choice of positions for mutation was influenced by two manually curated
factors: the distance from the receptor and the amino acid variation among ligand types. We designed

mutations with the aim of inferring a transition to the amino acid properties in sites where the ligands show
a different pattern. Overlapping residues at a given position were divided into two groups, based on an
EGF-like and non EGF-like stabilization of the receptor dimer. This property was previously shown to
follow binding affinity (18). Residues that introduced a noticeable shift in amino acid properties between
the two groups were selected. For example, position N32 is hydrophobic in the high affinity ligands group,
positively charged in the low affinity group and negatively charged in EGF. Finally, we carefully analyzed
exceptional cases in the DIRpred scoring. Some of the residues that show high score have intramolecular
interactions with another amino acid in the ligand. These residues, if mutated, will lose EGF structural
stability (namely "residue swapping" behavior showed in Figure S7). The decision of which mutation to
introduce was made using the paralogs alignment, with a preferential choice over the residue found in
EREG or EPGN. Positions 32, 48 and 50 have high DIRpred score. Position 46 was included although
having a lower score, because the substitution pattern matches the two ligands groupings. Furthermore
position 46, 48 and 50 were preferred because, given previous experiments and the overall scores, the EGF
C-terminus tail seems to play a critical role in the ligand function; see for example (58).

Synthetic Peptides
395 The wild-type, N32R, D46T, K48T, and W50Y variants of EGF were ordered from Scrum Net Co with
396 purity >90% and quantity 5 mg/mL. These peptides were used for ITC measurements, Circular Dichroism
397 (CD) measurements, proliferation studies, and Western Blot (WB) analyses.

Cell Lines
The *Bj5-tα human normal fibroblast cell line* was purchased from ATCC. Cells were grown in DMEM with
10% fetal bovine serum (FBS), and 5 µg/mL hygromycin B. The *Swiss Albino 3T3* mouse *normal fibroblast*
cell line was obtained from the RIKEN Cell Bank. Cells were grown in DMEM, 10% FBS, 50 ug/mL
gentamycin at 37°C in a 5% CO_2 atmosphere with 95% humidity. The *A431 human epithelial carcinoma*
adherent cell line (RIKEN Cell Bank) is a model skin cancer cell line with overexpressed EGFR used for
oncogenic pathway studies (59).Cells were cultured in DMEM supplemented with 10% FBS (Sigma-
Aldrich), 50 ug/mL gentamycin antibiotic or a combination of 100 unit/ml Penicillin G (Nacalai Tesque)
and 100 µg/ml streptomycin sulfate (Nacalai Tesque). Experiments were conducted at 37°C in a 5% CO_2-
enriched air atmosphere with 95% humidity. Cell lines were grown and used for Western Blot and cell
proliferation studies.

Cell Proliferation Assay
We measured cell proliferation using a label-free, non-invasive, cellular confluence assay with IncuCyte
Live-Cell Imaging Systems (Essen Bioscience). Human Bj5-tα (2,500 cells / well) and Mouse Swiss Albino
3T3 (1,000 cells/well) were seeded overnight on a 96-well plate (Corning) at 37°C in an incubator. The
next day, cells were treated with WT EGF and mutants at 1 nM, 10 nM and 100 nM concentrations and
placed in an XL-3 incubation chamber maintained at 37°C. The plate was scanned using a 4x objective at
2-hr intervals over 3 days. Cell confluence was measured using IncuCyte Analysis Software. The IncuCyte
Analyzer gives real-time confluence data based on segmentation of high-definition phase-contrast images.
Cell proliferation is shown as an increase in confluence rate relative to control.

Apoptosis Assay
Experiments were performed with the A431 human cancer cell line. 5,000 cells/well were seeded on a 96-
well plate (Corning) and incubated at 37°C for 24 hr. Media were replaced with fresh DMEM containing
WT EGF, or EGF mutants at 1, 10, and 100 nM concentrations and fluorescent annexin V green reagent.
Plates were pre-warmed prior to data acquisition to avoid condensation and expansion of the plate, which
affect autofocus. Images were captured every 2 hrs (4x) for 3 days in the IncuCyte system. Cell proliferation
is reported as in the previous assay.

Statistics

430 Proliferation and apoptosis experiments were performed in duplicates. All results are shown as the
431 mean±s.d. Raw data was analyzed by two-way ANOVA with 95% confidence level. The multiple test
432 was corrected using Bonferroni post hoc test. Prism 8 software was used for statistical analysis.
433 Asterisks in the pictures show P-values using GraphPad convention: $0.1234 >$ (ns), $0.0332 >$ (*), 0.0021
434 $>$ (**), $0.0002 >$ (***), $0.0001 >$ (****).
435
Isothermal Titration Calorimetry (ITC)
All ITC studies employed a MicroCal PEAQ-ITC System (Malvern). For titration, both EGFR ECD
(Sigma-Aldrich) and EGF variants were dialyzed into the same reaction buffer Milli-Q H_2O (22 μm) at
25°C. Each titration involved serial injections of 13×3 μL aliquots of EGF variants (200 μM) into a
solution of EGFR ECD (20 μM) in the sample cell. In each case, the reference cell was filled with the same
reaction buffer as the control to determine the heat upon binding of the two components. The measured heat
constant value was subtracted from the heat per injection prior to analysis of the data. The experiment was
replicated twice. Results were analyzed by MicroCal PEAQ-ITC Analysis Software.
444
Circular Dichroism (CD)
446 Far UV measurements were taken at a protein concentration of 0.1 μM, using a cuvette with a path length
447 of 0.1 cm. Secondary structure content was calculated from far UV spectra using CAPITO software (60).
448 Five scans in the 190-240-nm wavelength range were taken.
449
Western Blot Analysis
A431 epidermoid carcinoma cells were harvested using Lysis Buffer (0.4% SDS, 1 mM DTT, 1%). Samples
were incubated at 37°C for 10 min with Benzonase and centrifuged at 15,000 rpm at 22°C for 10 min.
Supernatants were used for further analysis. Sample concentrations were measured with a BCA protein
assay kit (ThermoFisher Scientific). Lysate were mixed with 4x Sample Loading Laemmli Buffer and
incubated at 90°C for 5 min. Equal amounts of protein were loaded in 12% Mini PROTEAN® TGX™
SDS-PAGE gel (Bio-Rad) and transferred to PVDF membranes (Trans-Blot Turbo RTA Mini 0.2 μm
PVDF Transfer Kit). Membranes were blocked for 10 min with Bullet Blocking One (Nacalai) and reacted
with monoclonal rabbit anti-EGFR antibody (Cell Signaling Technology, Inc.), Phospho-EGF Receptor
(Tyr1173) (Cell Signaling Technology, Inc.), and rabbit anti-α-tubulin pAb (MBL) at dilution of 1:1000.
Samples were incubated with Goat Anti-Rabbit IgG HRP at a 1:5000 dilution and chemiluminescent signals
were detected by CDP Plus (ThermoFisher Scientific) and ChemiDoc touch MP (Bio-rad).
462
Molecular Dynamics
All the simulations were performed with Gromacs version 2020 (61) using charmm36-mar2019 force field
(62) and SPC216 for water. PDB ID 5wb7 was used as model and template. EGF structure model was
extracted from PDB ID 1ivo and superimposed to the ligand using UCSF Chimera Match-Maker algorithm
(63). Mutants were generated using Swissmodel webserver (64), while missing atoms were compensated
using Scwrl4 (65). A system was composed of EGFR dimer in complex with one or two EGF wild type
(WT), N32R, D46T, K48T or W50Y were solvated and neutralized using NaCl ions in a dodecahedral box.
Then, energy temperature and pressure equilibrations were performed to the system following the
guidelines in Lemkul 2019 (66).
472
A 100 ns production simulation was run using the Verlet cut-off scheme (67) for non-bonded interactions
and LINCS as constraint algorithm (68). All the simulations reached convergence of RMSD. Long range
electrostatic interactions were computed using the Particle Mesh Ewald method (69) using a dedicated GPU.
The parameter files for all runs are included in the supplementary data [#ADD#]. We checked for
differences in relative motions between the three simulations by extracting and concatenating the backbone
trajectories using catDCD plugin of VMD (70), then performing a PCA using Bio3d R package (71).
479
480

Data availability
The source code and alignments data to reproduce the analysis of this paper are available at
https://zenodo.org/badge/latestdoi/287415954.

Acknowledgments
We thank Tadashi Yamamoto, Ichiro Maruyama for providing the antibodies, Keiko Kono for the technical
help with the WB experiments. We are grateful for the help and support provided by the Scientific
Computing section of Research Support Division and in particular Jan Moren at OIST. We thank Madhuri
Gade, Mirco Dindo, Harry Wilson and Benjamin Clifton for critical reading. We thank Pierre Matricon for
the discussion on the molecular dynamics data, Russel Alenton on the cell proliferation assay and Christine
Guzman on the western blot analyses.

Author contributions
PL, SP, DM conceived and designed the project; SP developed and performed the computational models
and analyses; DM and GU performed the experiments; all authors analyzed the data and wrote the
manuscript.

Funding and additional information
Financial support by the Okinawa Institute of Science and Technology to P.L. is gratefully acknowledged.

Conflict of interest
The authors declare that they have no conflict of interest with the contents of this article.

References

1.   Gazdar, A. F. (2009) Activating and resistance mutations of EGFR in non-small-cell lung
      cancer: role in clinical response to EGFR tyrosine kinase inhibitors. *Oncogene* **28**, S24-
      S31
2.   Oda, K., Matsuoka, Y., Funahashi, A., and Kitano, H. (2005) A comprehensive pathway
      map of epidermal growth factor receptor signaling. *Mol Syst Biol* **1**, 2005.0010
3.   Landau, M., Fleishman, S. J., and Ben-Tal, N. (2004) A putative mechanism for
      downregulation of the catalytic activity of the EGF receptor via direct contact between its
      kinase and C-terminal domains. *Structure* **12**, 2265-2275
4.   Purba, E. R., Saita, E. I., and Maruyama, I. N. (2017) Activation of the EGF Receptor by
      Ligand Binding and Oncogenic Mutations: The "Rotation Model". *Cells* **6**
5.   Wilson, K. J., Gilmore, J. L., Foley, J., Lemmon, M. A., and Riese, D. J., 2nd. (2009)
      Functional selectivity of EGF family peptide growth factors: implications for cancer.
      Pharmacol Ther **122**, 1-8
6.   Nicholson, R. I., Gee, J. M., and Harper, M. E. (2001) EGFR and cancer prognosis.
      European journal of cancer (Oxford, England : 1990) **37 Suppl 4**, S9-15
7.   Xu, M. J., Johnson, D. E., and Grandis, J. R. (2017) EGFR-targeted therapies in the post-
      genomic era. *Cancer Metastasis Rev* **36**, 463-473

8.   Alvarez-Ponce, D., Feyertag, F., and Chakraborty, S. (2017) Position Matters: Network
Centrality Considerably Impacts Rates of Protein Evolution in the Human Protein-Protein
Interaction Network. *Genome Biol Evol* **9**, 1742-1756
9.   Jensen, R. A. (2001) Orthologs and paralogs - we need to get it right. *Genome Biol* **2**,
Interactions1002
10. Groenen, L. C., Nice, E. C., and Burgess, A. W. (1994) STRUCTURE-FUNCTION-
RELATIONSHIPS FOR THE EGF/TGF-ALPHA FAMILY OF MITOGENS. *Growth*
Factors **11**, 235-257
11. Souriau, C., Gracy, J., Chiche, L., and Weill, M. (1999) Direct selection of EGF mutants
displayed on filamentous phage using cells overexpressing EGF receptor. *Biol. Chem.* **380**,
451-458
12. Skwark, M. J., Raimondi, D., Michel, M., and Elofsson, A. (2014) Improved Contact
Predictions Using the Recognition of Protein Like Contact Patterns. *PLOS Computational*
Biology **10**, e1003889
13. Gocheva, G., and Ivanova, A. (2019) A Look at Receptor-Ligand Pairs for Active-
Targeting Drug Delivery from Crystallographic and Molecular Dynamics Perspectives.
Mol Pharm **16**, 3293-3321
14. Evans, B. A., Sato, M., Sarwar, M., Hutchinson, D. S., and Summers, R. J. (2010) Ligand-
directed signalling at beta-adrenoceptors. *Br J Pharmacol* **159**, 1022-1038
15. Kenakin, T. (2011) Functional selectivity and biased receptor signaling. *J Pharmacol Exp*
Ther **336**, 296-302
16. Ali, R., Brown, W., Purdy, S. C., Davisson, V. J., and Wendt, M. K. (2018) Biased
signaling downstream of epidermal growth factor receptor regulates proliferative versus
apoptotic response to ligand. *Cell Death Dis* **9**, 976
17. Strachan, L., Murison, J. G., Prestidge, R. L., Sleeman, M. A., Watson, J. D., and Kumble,
K. D. (2001) Cloning and biological activity of epigen, a novel member of the epidermal
growth factor superfamily. *The Journal of biological chemistry* **276**, 18265-18271
18. Freed, D. M., Bessman, N. J., Kiyatkin, A., Salazar-Cavazos, E., Byrne, P. O., Moore, J.
O., Valley, C. C., Ferguson, K. M., Leahy, D. J., Lidke, D. S., and Lemmon, M. A. (2017)
EGFR Ligands Differentially Stabilize Receptor Dimers to Specify Signaling Kinetics.
Cell **171**, 683-695.e618
19. Huang, Y., Bharill, S., Karandur, D., Peterson, S. M., Marita, M., Shi, X., Kaliszewski, M.
J., Smith, A. W., Isacoff, E. Y., and Kuriyan, J. (2016) Molecular basis for multimerization
in the activation of the epidermal growth factor receptor. *Elife* **5**
20. Mehrabi, M., Mahdiuni, H., Rasouli, H., Mansouri, K., Shahlaei, M., and Khodarahmi, R.
(2018) Comparative experimental/theoretical studies on the EGFR dimerization under the
effect of EGF/EGF analogues binding: Highlighting the importance of EGF/EGFR
interactions at site III interface. *Int J Biol Macromol* **115**, 401-417
21. Knudsen, S. L., Mac, A. S., Henriksen, L., van Deurs, B., and Grovdal, L. M. (2014) EGFR
signaling patterns are regulated by its different ligands. *Growth Factors* **32**, 155-163
22. Liu, P., Cleveland, T. E. t., Bouyain, S., Byrne, P. O., Longo, P. A., and Leahy, D. J. (2012)
A single ligand is sufficient to activate EGFR dimers. *Proc Natl Acad Sci U S A* **109**,
10861-10866
23. Arkhipov, A., Shan, Y., Kim, E. T., and Shaw, D. E. (2014) Membrane interaction of bound
ligands contributes to the negative binding cooperativity of the EGF receptor. *PLoS*
Comput Biol **10**, e1003742

24. Chagoyen, M., García-Martín, J. A., and Pazos, F. (2015) Practical analysis of specificity-
determining residues in protein families. *Briefings in Bioinformatics* **17**, 255-261
25. Spielman, S. J., and Wilke, C. O. (2015) Pyvolve: A Flexible Python Module for
Simulating Sequences along Phylogenies. *PloS one* **10**, e0139047
26. Chakrabarti, S., Bryant, S. H., and Panchenko, A. R. (2007) Functional specificity lies
within the properties and evolutionary changes of amino acids. *J Mol Biol* **373**, 801-810
27. Gallay, J., Vincent, M., Li de la Sierra, I. M., Alvarez, J., Ubieta, R., Madrazo, J., and
Padron, G. (1993) Protein flexibility and aggregation state of human epidermal growth
factor. A time-resolved fluorescence study of the native protein and engineered single-
tryptophan mutants. *Eur J Biochem* **211**, 213-219
28. Tynan, C. J., Roberts, S. K., Rolfe, D. J., Clarke, D. T., Loeffler, H. H., Kastner, J., Winn,
M. D., Parker, P. J., and Martin-Fernandez, M. L. (2011) Human epidermal growth factor
receptor (EGFR) aligned on the plasma membrane adopts key features of Drosophila
EGFR asymmetry. *Mol Cell Biol* **31**, 2241-2252
29. Li De La Sierra, I. M., Vincent, M., Padron, G., and Gallay, J. (1992) Interaction of
recombinant human epidermal growth factor with phospholipid vesicles. A steady-state
and time-resolved fluorescence study of the bis-tryptophan sequence (TRP49-TRP50).
European Biophysics Journal **21**, 337-344
30. Hommel, U., Harvey, T. S., Driscoll, P. C., and Campbell, I. D. (1992) Human epidermal
growth factor. High resolution solution structure and comparison with human transforming
growth factor alpha. *J Mol Biol* **227**, 271-282
31. Puddicombe, S. M., Chamberlin, S. G., MacGarvie, J., Richter, A., Drummond, D. R.,
Collins, J., Wood, L., and Davies, D. E. (1996) The significance of valine 33 as a ligand-
specific epitope of transforming growth factor alpha. *The Journal of biological chemistry*
271, 15367-15372
32. Lahti, J. L., Lui, B. H., Beck, S. E., Lee, S. S., Ly, D. P., Longaker, M. T., Yang, G. P., and
Cochran, J. R. (2011) Engineered epidermal growth factor mutants with faster binding on-
rates correlate with enhanced receptor activation. *FEBS letters* **585**, 1135-1139
33. Gill, G. N., and Lazar, C. S. (1981) Increased phosphotyrosine content and inhibition of
proliferation in EGF-treated A431 cells. *Nature* **293**, 305-307
34. Gulli, L. F., Palmer, K. C., Chen, Y. Q., and Reddy, K. B. (1996) Epidermal growth factor-
induced apoptosis in A431 cells can be reversed by reducing the tyrosine kinase activity.
Cell Growth Differ **7**, 173-178
35. Nemoto, W., Saito, A., and Oikawa, H. (2013) Recent advances in functional region
prediction by using structural and evolutionary information - Remaining problems and
future extensions. *Computational and structural biotechnology journal* **8**, e201308007
36. Ashkenazy, H., Abadi, S., Martz, E., Chay, O., Mayrose, I., Pupko, T., and Ben-Tal, N.
(2016) ConSurf 2016: an improved methodology to estimate and visualize evolutionary
conservation in macromolecules. *Nucleic Acids Res.* **44**, W344-W350
37. Lichtarge, O., Bourne, H. R., and Cohen, F. E. (1996) An evolutionary trace method
defines binding surfaces common to protein families. *J. Mol. Biol.* **257**, 342-358
38. Teppa, E., Wilkins, A. D., Nielsen, M., and Buslje, C. M. (2012) Disentangling
evolutionary signals: conservation, specificity determining positions and coevolution.
Implication for catalytic residue prediction. *BMC Bioinformatics* **13**, 235
39. Carpenter, G., and Cohen, S. (1979) Epidermal growth factor. *Annual review of*
biochemistry **48**, 193-216

40. Lindvall, C., Hou, M., Komurasaki, T., Zheng, C., Henriksson, M., Sedivy, J. M.,
Bjorkholm, M., Teh, B. T., Nordenskjold, M., and Xu, D. (2003) Molecular
characterization of human telomerase reverse transcriptase-immortalized human
fibroblasts by gene expression profiling: activation of the epiregulin gene. *Cancer Res* **63**,
1743-1747

41. Grudinkin, P. S., Zenin, V. V., Kropotov, A. V., Dorosh, V. N., and Nikolsky, N. N. (2007)
EGF-induced apoptosis in A431 cells is dependent on STAT1, but not on STAT3. *Eur J*
Cell Biol **86**, 591-603

42. Ibuka, S., Matsumoto, S., Fujii, S., and Kikuchi, A. (2015) The P2Y(2) receptor promotes
Wnt3a- and EGF-induced epithelial tubular formation by IEC6 cells by binding to integrins.
J Cell Sci **128**, 2156-2168

43. Bjorkelund, H., Gedda, L., and Andersson, K. (2011) Comparing the epidermal growth
factor interaction with four different cell lines: intriguing effects imply strong dependency
of cellular context. *PloS one* **6**, e16536

44. Macdonald-Obermann, J. L., and Pike, L. J. (2014) Different epidermal growth factor
(EGF) receptor ligands show distinct kinetics and biased or partial agonism for homodimer
and heterodimer formation. *The Journal of biological chemistry* **289**, 26178-26188

45. Zhou, J., Liu, D., Sa, Z., Huang, W., Zou, Y., and Gu, X. (2017) Effective estimation of
the minimum number of amino acid residues required for functional divergence between
duplicate genes. *Mol Phylogenet Evol* **113**, 126-138

46. Laisney, J., Braasch, I., Walter, R. B., Meierjohann, S., and Schartl, M. (2010) Lineage-
specific co-evolution of the Egf receptor/ligand signaling system. *BMC Evol. Biol.* **10**, 16

47. Bakker, J., Spits, M., Neefjes, J., and Berlin, I. (2017) The EGFR odyssey – from activation
to destruction in space and time. *Journal of Cell Science* **130**, 4087-4096

48. Zerbino, D. R., Achuthan, P., Akanni, W., Amode, M. R., Barrell, D., Bhai, J., Billis, K.,
Cummins, C., Gall, A., Giron, C. G., Gil, L., Gordon, L., Haggerty, L., Haskell, E., Hourlier,
T., Izuogu, O. G., Janacek, S. H., Juettemann, T., To, J. K., Laird, M. R., Lavidas, I., Liu,
Z. C., Loveland, J. E., Maurel, T., McLaren, W., Moore, B., Mudge, J., Murphy, D. N.,
Newman, V., Nuhn, M., Ogeh, D., Ong, C. K., Parker, A., Patricio, M., Riat, H. S.,
Schuilenburg, H., Sheppard, D., Sparrow, H., Taylor, K., Thormann, A., Vullo, A., Walts,
B., Zadissa, A., Frankish, A., Hunt, S. E., Kostadima, M., Langridge, N., Martin, F. J.,
Muffato, M., Perry, E., Ruffier, M., Staines, D. M., Trevanion, S. J., Aken, B. L.,
Cunningham, F., Yates, A., and Flicek, P. (2018) Ensembl 2018. *Nucleic Acids Res.* **46**,
D754-D761

49. Katoh, K., and Standley, D. M. (2013) MAFFT Multiple Sequence Alignment Software
Version 7: Improvements in Performance and Usability. *Mol. Biol. Evol.* **30**, 772-780

50. Berman, H. M., Westbrook, J., Feng, Z., Gilliland, G., Bhat, T. N., Weissig, H., Shindyalov,
I. N., and Bourne, P. E. (2000) The Protein Data Bank. *Nucleic Acids Res* **28**, 235-242

51. Goddard, T. D., Huang, C. C., Meng, E. C., Pettersen, E. F., Couch, G. S., Morris, J. H.,
and Ferrin, T. E. (2018) UCSF ChimeraX: Meeting modern challenges in visualization and
analysis. *Protein Sci.* **27**, 14-25

52. Okonechnikov, K., Golosova, O., Fursov, M., and Team, U. (2012) Unipro UGENE: a
unified bioinformatics toolkit. *Bioinformatics* **28**, 1166-1167

53. Nguyen, L. T., Schmidt, H. A., von Haeseler, A., and Minh, B. Q. (2015) IQ-TREE: A Fast
and Effective Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies.
Mol. Biol. Evol. **32**, 268-274

54. Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K. F., von Haeseler, A., and Jermiin, L. S.
(2017) ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat.*
Methods **14**, 587-+

55. Capra, J. A., and Singh, M. (2007) Predicting functionally important residues from
sequence conservation. *Bioinformatics* **23**, 1875-1882

56. Mirny, L. A., and Gelfand, M. S. (2002) Using Orthologous and Paralogous Proteins to
Identify Specificity-determining Residues in Bacterial Transcription Factors. *J. Mol. Biol.*
321, 7-20

57. Whelan, S., and Goldman, N. (2001) A general empirical model of protein evolution
derived from multiple protein families using a maximum-likelihood approach. *Mol Biol*
Evol **18**, 691-699

58. Puddicombe, S. M., Wood, L., Chamberlin, S. G., and Davies, D. E. (1996) The interaction
of an epidermal growth factor/transforming growth factor alpha tail chimera with the
human epidermal growth factor receptor reveals unexpected complexities. *The Journal of*
biological chemistry **271**, 30392-30397

59. Giard, D. J., Aaronson, S. A., Todaro, G. J., Arnstein, P., Kersey, J. H., Dosik, H., and
Parks, W. P. (1973) In vitro cultivation of human tumors: establishment of cell lines
derived from a series of solid tumors. *J Natl Cancer Inst* **51**, 1417-1423

60. Wiedemann, C., Bellstedt, P., and Gorlach, M. (2013) CAPITO--a web server-based
analysis and plotting tool for circular dichroism data. *Bioinformatics* **29**, 1750-1757

61. Abraham, M. J., Murtola, T., Schulz, R., Páll, S., Smith, J. C., Hess, B., and Lindahl, E.
(2015) GROMACS: High performance molecular simulations through multi-level
parallelism from laptops to supercomputers. *SoftwareX* **1-2**, 19-25

62. Huang, J., and MacKerell, A. D., Jr. (2013) CHARMM36 all-atom additive protein force
field: validation based on comparison to NMR data. *J Comput Chem* **34**, 2135-2145

63. Meng, E. C., Pettersen, E. F., Couch, G. S., Huang, C. C., and Ferrin, T. E. (2006) Tools
for integrated sequence-structure analysis with UCSF Chimera. *BMC Bioinformatics* **7**,
339

64. Waterhouse, A., Bertoni, M., Bienert, S., Studer, G., Tauriello, G., Gumienny, R., Heer, F.
T., de Beer, T. A P., Rempfer, C., Bordoli, L., Lepore, R., and Schwede, T. (2018) SWISS-
MODEL: homology modelling of protein structures and complexes. *Nucleic Acids Res.* **46**,
W296-W303

65. Krivov, G. G., Shapovalov, M. V., and Dunbrack, R. L., Jr. (2009) Improved prediction of
protein side-chain conformations with SCWRL4. *Proteins* **77**, 778-795

66. Lemkul, J. A. (2019) From Proteins to Perturbed Hamiltonians: A Suite of Tutorials for
the GROMACS-2018 Molecular Simulation Package [Article v1.0]. *Living J. Comp. Mol.*
Sci. **1(1)**, 5068

67. Páll, S., and Hess, B. (2013) A flexible algorithm for calculating pair interactions on SIMD
architectures. *Computer Physics Communications* **184**, 2641-2650

68. Hess, B., Bekker, H., Berendsen, H. J. C., and Fraaije, J. G. E. M. (1997) LINCS: A linear
constraint solver for molecular simulations. *Journal of Computational Chemistry* **18**, 1463-
1472

69. Essmann, U., Perera, L., Berkowitz, M. L., Darden, T., Lee, H., and Pedersen, L. G. (1995)
A smooth particle mesh Ewald method. *The Journal of Chemical Physics* **103**, 8577-8593

70. Humphrey, W., Dalke, A., and Schulten, K. (1996) VMD: visual molecular dynamics. *J*
Mol Graph **14**, 33-38, 27-38

71. Grant, B. J., Rodrigues, A. P., ElSawy, K. M., McCammon, J. A., and Caves, L. S. (2006)
Bio3d: an R package for the comparative analysis of protein structures. *Bioinformatics* **22**,
2695-2696
72. Okonechnikov, K., Golosova, O., Fursov, M., and team, t. U. (2012) Unipro UGENE: a
unified bioinformatics toolkit. *Bioinformatics* **28**, 1166-1167
73. Colell, E. A., Iserte, J. A., Simonetti, F. L., and Marino-Buslje, C. (2018) MISTIC2:
comprehensive server to study coevolution in protein families. *Nucleic Acids Res* **46**,
W323-w328

720
721

722 **Abbreviations**
723

724 **EGF**, Epidermal Growth Factor; **EGFR**, Epidermal Growth Factor Receptor; **HBEGF**, Heparin-Binding
725 Epidermal Growth Factor; **EPGN**, Epigen; **BTC**, Betacellulin; **EREG**, Epiregulin; **AREG**,
726 Amphiregulin; **PPI**, Protein-Protein Interaction; **MSA**, Multiple Sequence Alignment; **MSTA**, Multiple
727 STructural Alignment; **ITC**, Isothermal Titration Calorimetry; **DMEM**, Dulbecco's Modified Eagle
728 Medium; **ECD**, Extra Cellular Domain; **WT**, Wild Type; **CD**, circular dichroism.
729
730

731 **Figures**
732

733
734 **Figure 1. Study rationale.** EGF mutants were identified using an *ad hoc* methodology that combines
735 conservation and coevolution measures, DIRpred. Unlike WT EGF, the mutant ligands induce an unstable
736 conformation of the receptor dimer, similarly to a dimer with a single WT EGF. Treating A431 and
737 fibroblast cells with either WT EGF or mutant EGF resulted in a different response both in the
738 phosphorylation dynamics and the cell proliferation phenotype.
739

740
Figure 2. DIRpred analysis of EGF. (A) The relative ranking score of the four mutants. The outer circle
represents paralogs multiple sequence analysis (MSA)-based scores, while the inner circle represents
paralogs multiple structural alignment (MSTA)-based scores. The darker color indicates a higher ranking
in the EGF sites-based scoring. The ranking values are reported in Table S1. (B) Cross-conservation plot.
The plot is obtained by crossing the two conservation scores. Interestingly, no point lies in the bottom right
half of the plot (high paralogs conservation and low orthologs conservation), suggesting that paralogs and
orthologs conservation are not independent. This observation points out that there is no organism-specific
adaptation shared by all ligands at the protein sequence level. (C) Cross-co-evolution score. The plot is
obtained by crossing the two co-evolution scores, the ligand-receptor (L-R) co-evolution score (III) on the
y axis and the ligand-ligand (L-L) co-evolution score (IV) on the x axis.
751

752
Figure 3. ITC measurements of EGF mutants and the EGFR receptor. (A) ITC analysis of WT EGF
ligand and mutants N32R, D46T, K48T, and W50Y binding to the ECD of the EGFR receptor at 25° C.
Measurements were taken by adding WT EGF or mutants at 200 μM to the ECD of EGFR at 20 μM. **(B)**
Four zoom-in of X-Ray structure of the extracellular domain (ECD) of EGFR bound to EGF (PDB 1IVO).
In cyan cartoon the ECD of EGFR. Each zoom-in focuses on the mutated residue. Highlighted in yellow
stick side chain of the mutated residues and in cyan stick side chain of the residue of ECD in proximity (<
5 Å with the mutated residue. (C) K_d calculated from the ITC measurements using the program Affinimeter
KinITC Kintecs Software.
761

Figure 4. Effects of WT and single-mutant EGF on the receptor in A431 cells at different time points. (A) Short-term effects on the phosphorylation level of EGFR Tyr-1173 after treating A431 cells with saturating concentrations of different ligands. The bar plot reports the duplicate results. The membrane containing EGFR and tubulin was separated after the transfer. D46T treated cells show a statistically significant reduction in the level of phosphorylation compared to the other samples. (B) Western blot at 1 and 6 h for the long-term effects. The samples were tested two times each biological duplicate. The bars represent mean ± s.d. Band intensity estimates were calculated using BIO-RAD ImageLab software (BioRad). Plots and statistics were performed using PRISM software (GraphPad).

Figure 5. Growth assays of cells treated with EGF variants. (A) Effect of different concentrations of EGF variants on proliferation of the human normal fibroblast Bj5-tα cell line. Data represent the growth rate percentage calculated from the confluence of cells relative to the control (mean ± standard deviation) for each concentration of EGF. Percent confluence was estimated 24 h after the treatment (two replicates per treatment). (B) Apoptosis effect of WT or mutant EGF on A431 cells. EGF-induced apoptosis is measured as the reduction in total population compared to the control. (C) Comparison of A431 cell growth after treatment with 100 nM WT EGF and EGF variants D46T and K48T. Cells were labeled with fluorescent Annexin V Green Reagent. Plates were pre-warmed prior to data acquisition to avoid condensation and expansion of the plate, which affect autofocus. Images were captured every 2 h (4x magnification) for 3 days in the IncuCyte system.

784
Figure 6. Molecular Dynamics of the EGFR extracellular domain. (A) Root Mean Square Deviation
from the initial structure. (B) Top, Root Mean Square Fluctuations of the 100 ns simulations. The WT level
is marked in bold and dashed red line. Bottom, focus on the RMSF of the dimerization arm domain. The
two plots represent the two EGFR in the receptor dimer. (C) Dynamics of the dimerization arm domain in
WT (red) and K48T (green) simulations. A structure every 20 ns of simulation was taken and aligned to
1IVO reference structure (blue surface and black ribbon). The structure at 20 ns is represented in solid
colors. K48T simulation shows a more dynamic dimerization arm. (D) Number of H-bonds formed between
the two receptors during the 100 ns simulation time. (E) Number of H-bonds formed between the receptor
that shows a dynamic dimerization arm and the corresponding ligand during the 100 ns simulation time.
(F) Distance used to investigate the fluctuations of the dimerization domain highlighted on the structure.
Pro-272 (purple) and Gly-288 (red) are highlighted both on 5WB7 (green and teal) and in 1IVO (grey). The
distance between the alpha carbons of the 5WB7 couple is shown in brown and is notably longer than 1IVO.
(G) Distribution of the distance between Pro-272 and Gly-288 in all the simulations. WT simulation shows
a single peak at ~1nm, while in all other ligands the distance has two peaks.
799

I) Orthologs conservation

$$I(p) = \frac{\sum_l^{para} cons\_score(l_p)}{N_l}$$

EGF

HB-EGF

TGFA

II) Paralogs MSA/MSTA conservation

$$II(p) = 1 - cons\_score(msa_p)$$

MSA

MSTA

conservation scores

Identity

$$cons_{ID}(C_p) = \sum_p^C 1 \; if \; p = ref \Big/ N_C$$

Blosum

$$cons_{BLOS}(C_p) = \frac{\sum_p^C blos62(ref,p) - \min(blos62)}{\max(blos62) - \min(blos62)} \Big/ N_C$$

JSDw (Capra et al. 2007)

$$cons_{JSDw}(C_p) = \lambda RE_{p_c,r} + (1-\lambda)RE_{q,r}$$

Ligands

ECD of
EGFR

III) Ligand-receptor Co-evolution

$$III(p) = \max(coevo\_score(ref_p, EGFR_*))$$

$$DIRpred(p) = \alpha I + \beta II + \gamma III + \delta IV$$

Where

$$\alpha + \beta + \gamma + \delta = 1$$

IV) Ligands internal Co-evolution

$$IV(p) = 1 - \max(coevo\_score(para_p, para_*))$$

800
Figure 7. DIRpred rationale. The prediction of Divergence Inducing Residues consists of a linear
combination of four site-specific scores, in roman numerals. In I, the orthologs MSA of each paralog is
used to compute a conservation score (as exemplified from the histograms on the alignments). Alignment
images were produced using Unipro software (72). In II, the human paralog sequences or structures were
aligned and the same conservation score function was used. In the purple box, the conservation score
functions are represented. In III, the co-evolution score between EGF and each receptor site was computed
using Mutual Information (MI). The yellow lines on the structure connect the highest scoring co-evolving
residues between ligand and receptor. In IV, MI co-evolution score was computed from the combined
ligands orthologs alignment. The circular plot was performed using MISTIC2 (73).
810