1

Epidermal Growth Factor single mutants highlighted by homologs cross-conservation

approach differentially affect the EGF Receptor downstream pathway

Dalmira Merzhakupova[1¶], Stefano Pascarelli[1 ¶], Paola Laurino[1*]

[1] Okinawa Institute of Science and Technology Graduate University, 1919-1 Tancha, Onna-son, Okinawa 904-

0412 Japan.

* Corresponding author

E-mail: paola.laurino@oist.jp

¶ These authors contributed equally to this work.

Abstract

Epidermal Growth Factor Receptor is a tyrosine kinase receptor that plays a key role in cells.

Its ligands contribute to the system modularity by influencing the stability of the dimerized

receptor. A synthetically modified ligand could be a potential anti-cancer drug if able to

compete with the endogenous ligands while modulating a different pathway activation.

Herein, we developed a cross-conservation approach to identify functionally important EGF

residues. Four EGF mutants have been selected and studied. These mutants show binding

affinities for EGFR similar to EGF, while the effect at the cellular level changed, increasing

growth rate in fibroblasts and reducing apoptosis in skin cancer cells. These results support

the theory of biased signaling in the tyrosine kinase receptor system.

Abbreviations

EGF, Epidermal Growth Factor

EGFR, Epidermal Growth Factor Receptor

PPI, Protein-Protein Interaction

MSA, Multiple Sequence Alignment

MSTA, Multiple STructural Alignment

ITC, Isothermal Titration Calorimetry

DMEM, Dulbecco's Modified Eagle Medium

ECD, Extra Cellular Domain

WT, Wild Type

Introduction

Protein-protein interactions (PPIs) regulate many biological processes (1). PPIs are one of the

most interesting and well-studied examples of molecular co-evolution in biological systems.

These interactions are sometimes defined by one part (receptor) that binds several

counterparts (ligands). Receptors and ligands experience different selective constraints, and

receptors tend to evolve more slowly due to the necessity of binding multiple ligands.

Identification of key residues in a ligand that may affect binding and the resulting cellular

phenotype should provide new understanding of PPI co-evolution (2).

In recent years, different experimental techniques have been developed to define the effects

of single mutant proteins at the cellular level (3). Often these approaches generate false-

positive and false-negative that can be misinterpreted and leading to unclear conclusions (3).

Bioinformatic tools can be exploited for a more detailed analysis of PPIs co-evolution. The

combination of sequence and structural alignment methods have made possible to identify

amino acids essential for ligand-receptor interactions, but the ligand effect at the cellular

level still remains unclear (4).

The epidermal growth factor (EGF)-like domain ligand – ErbB1(EGFR) receptor signaling

system is involved in many biological events in multicellular organisms (5) and is considered

to be ancient (6). Few studies have shown that overexpression of oncogenic receptors and

ligands may induce different types of cancers (7). EGF ligands are also involved in ion

transport, glycolysis, and synthesis of proteins and nucleic acids (8). They also induce

stimulation of fibroblasts in early phases of wound healing (9). As anti-EGFR antibodies

often lead to inconsistent outcomes, design of EGF analogues remained an attractive target

for biomedical applications (10).

Ligand-induced selective activation of downstream pathway has been observed in G-coupled

protein receptor as "biased signaling" (11, 12). Recent reports hints that this mechanism

might take place also in EGFR tyrosine kinase (13). Comprehensive analysis of specific

residues in EGF ligands from different species and among different EGFR ligands (EGF,

HBEGF, EPGN, BTC, EREG, AREG, TGFA) might allow the design of mutants with

different or improved functions. Recent studies showed that some EGF residues like R41 and

L47 are highly conserved and important for high binding affinity to EGFR in the A431

cancer cell line (14). Another study highlighted Y13, L15 and H16 residues as essential for

downstream activity of ErbB1 (15). These outcomes were based on structural analyses of

ligands and experimental validation. Although, to the best of our knowledge a study to test a

difference in the biased signaling of a single-residue mutated EGF ligand was not performed

yet.

In this paper we show a novel approach to study PPIs, through cross-conservation analysis.

We combined bioinformatics and experimental tools to study co-evolution of the EGF-EGFR

ligand-receptor system at the residue level (Figure 1A). The identified residues have

phenotypic implications at the cellular level, influencing protein activation in the EGFR

downstream pathway. Overall, this approach has been critical to identify residues that play

important role in biased signaling.

Materials and Methods

Sequence and structure analysis

The sequences of EGFR ligands and the multiple sequence alignment of EGF orthologs were

obtained from Ensembl database (16). Multiple sequence alignment of all ligands was

performed with MAFFT software (17). X-ray structures were obtained from the PDB

database (18). Structure alignments were created using Chimera (19).

Phylogenetic analysis

From the multiple sequence alignment of EGF from different species, nearly identical

sequences were removed. The fruit fly EGF sequence was added as an outgroup in the EGF

phylogeny tree, while *Caenorhabditis elegans* EGF was used as outgroup in the tree of all

ligands. The image of MSA and phylogenetic trees was obtained using unipro UGENE

software (20). Three phylogenetic trees were made using Neighbor Joining (NJ), Maximum

Likelihood (ML), and Bayesian (MrB) methods.

As additional method, trees were also made with IQTREE (21), using ModelFinder to scan

for the most fit evolutionary model and parameters (22).

Calculation of Cross conservation score

From the evolutionary MSA of EGF orthologs and ligands MSA (or MSTA) of EGF paralogs,

two conservation measures were obtained. The conservation score was calculated in three

different ways: 1) IDENTITY, 2) BLOSUM, and 3) JSDw. The IDENTITY score measures

the frequency of appearance of an EGF residue in other ligands. In the BLOSUM score,

reference position substitutions are weighted using the BLOSUM62 matrix. The JSDw score

is based on Jensen Shannon divergence with a window of residues, and is the method used by

Pupko *et al.* in ConSurf (23). The cross-conservation plot and analysis were performed with

the Python package SEABORN. The two levels of conservation were plotted, and a cross-

conservation score was obtained by computing the distance from the diagonal.

The code used in the analysis of the cross-conservation score and plots and the data used in

this paper are shared on Github: https://github.com/oist/CrossConservation**.**

Mutation rationale

Along with the cross-conservation score, the choice of positions for mutation was influenced

by two other factors: the distance from the receptor, and the amino acid variation among

ligand types. Each ligand has different binding affinity and activate different pathways;

therefore, we designed mutations with the aim of changing pathway activation, by taking into

consideration the residues types in the ligands that show a different cellular effect. Finally,

we carefully analyzed exceptional cases in the cross conservation scoring. Some of the

residues that show high score have intramolecular interactions with another amino acid.

These residues, if mutated, will not only change interaction with the receptor but also lose

EGF structural stability (namely "residue swapping" behavior showed in Figure S2). The

decision of which mutation to introduce was made using EGFR ligands alignment.

Overlapping residues at a given position were divided into two groups based on EGF-like and

non EGF-like stabilization of the receptor dimer. This separation was previously shown to

follow binding affinity (24). Residues that introduced a noticeable shift in amino acid

properties in the two groups were selected. The chosen positions have higher-than-median

cross conservation scores. Furthermore position 46, 48 and 50 were preferred because, given

previous experiments and the overall scores, the EGF C-terminus tail seems to play a critical

role in the ligand function, for example (25). N32R was also chosen since it is highly

conserved in other ligands in the corresponding evolutionary alignment (e.g. CVC in TGFA,

or CRC in EREG), therefore hinting for a high cross-conservation score when computed for

other ligands.

Synthetic Peptides

Wild-type EGF (protein sequence:

N'NSDSECPLSHDGYCLHDGVCMYIEALDKYACNCVVGYIGERCQYRDLKWWELR-

C')

and EGF variants (See below the list of peptides) with purity >90% and quantity 5 mg/mL

were ordered from Scrum Net Co. These peptides were used for ITC measurements, Circular

Dichroism (CD) measurements, proliferation studies, and Western Blot (WB) analyses.

The list of Mutations:

Name	Mut1	Mut2	Mut3	Mut4
Position	46	48	50	32
Amino acid substitutions	D46T	K48T	W50Y	N32R

Cell Lines

The *Bj5-tα human normal fibroblast cell line* was purchased from ATCC. Cells were grown

in DMEM with 10% fetal bovine serum (FBS), and 5 µg/mL hygromycin B.

The *Swiss Albino 3T3* mouse *normal fibroblast cell line* was obtained from the RIKEN Cell

Bank. Cells were grown in DMEM, 10% FBS, 50 ug/mL gentamycin at $37^{o}C$ in a 5% CO_2

atmosphere with 95% humidity.

The *A431 human epithelial carcinoma adherent cell line* (RIKEN Cell Bank) is a model skin

cancer cell line with overexpressed EGFR used for oncogenic pathway studies (G. Carpenter

et.al.,1983). Cells were cultured in DMEM supplemented with 10% FBS (Sigma-Aldrich), 50

ug/mL gentamycin antibiotic.

Experiments were conducted at 37°C in a 5% CO_2-enriched air atmosphere with 95%

humidity. Cell lines were grown and used for cell ELISA and cell proliferation studies.

Cell Proliferation Assay

We measured cell proliferation using a label-free, non-invasive, cellular confluence assay with IncuCyte Live-Cell Imaging Systems (Essen Bioscience). Human Bj5-tα (2,500 cells / well) and Mouse Swiss Albino 3T3 (1,000 cells/well) were seeded overnight on a 96-well plate (Corning) at 37°C in an incubator. The next day, cells were treated with WT EGF and mutants at 1 nM, 10 nM and 100 nM concentrations and placed in an XL-3 incubation chamber maintained at 37°C. The plate was scanned using a 4x objective at 2-hr intervals over 3 days. Cell confluence was measured using IncuCyte Analysis Software. The IncuCyte Analyzer gives real-time confluence data based on segmentation of high-definition phase-contrast images. Cell proliferation is shown as relative rate increase in confluence to control.

Apoptosis Assay

Experiments were performed with the A431 human cancer cell line. 5,000 cells/well were seeded on a 96-well plate (Corning) and incubated at 37°C for 24 hr. Media were replaced with fresh DMEM containing WT EGF, or EGF mutants at 1, 10, and 100 nM concentrations and fluorescent annexin V green reagent. Plates were pre-warmed prior to data acquisition to avoid condensation and expansion of the plate, which affect autofocus. Images were captured every 2 hrs (4x) for 3 days in the IncuCyte system. Cell proliferation is reported as in the previous assay.

Statistics

Proliferation and apoptosis experiments were performed in duplicates. All results are shown as the mean±s.d. Raw data was analyzed by two-way ANOVA with 95% confidence level. The multiple test was corrected using Bonferroni post hoc test. Prism 8 software was used

for statistical analysis. Asterisks in the pictures show P-values using GraphPad

convention: 0.1234 > (ns), 0.0332 > (*), 0.0021 > (**), 0.0002 > (***), 0.0001 > (****).

Isothermal Titration Calorimetry (ITC)

All ITC studies employed a MicroCal PEAQ-ITC System (Malvern). For titration, both

EGFR ECD (Sigma-Aldrich) and EGF variants were dialyzed into the same reaction buffer

Milli-Q H_2O (22 µm) at 25°C. Each titration involved serial injections of 13×3 µL aliquots

of EGF variants (200 µM) into a solution of EGFR ECD (20 µM) in the sample cell. In each

case, the reference cell was filled with the same reaction buffer as the control to determine

the heat upon binding of the two components. The measured heat constant value was

subtracted from the heat per injection prior to analysis of the data. The experiment was

replicated twice. Results were analyzed by MicroCal PEAQ-ITC Analysis Software.

Circular Dichroism (CD)

Far UV measurements were taken at a protein concentration of 0.1 µM, using a cuvette with a

path length of 0.1 cm. Secondary structure content was calculated from far UV spectra using

CAPITO software (26). Five scans in the 190-240-nm wavelength range were taken.

Western Blot Analysis

A431 epidermoid carcinoma cells were harvested using Lysis Buffer (0.4% SDS, 0.2%

BETA-ME, 1% Bromophenol Blue, 10% glycerol. Samples were incubated at 65$^°$C for 10

min, sonicated, and centrifuged at 15,000 rpm at 22$^°$C for 10 min. Supernatants were used for

further analysis. Sample concentrations were measured with a Pierce^{TM} BCA protein assay

kit (ThermoFisher Scientific). Proteins were mixed with 2x Sample Loading Laemmli Buffer

and incubated at 65$^°$C for 10 min. Equal amounts of protein were loaded in 4-15% Mini-

PROTEAN® TGX™ SDS-PAGE gel (Bio-Rad) and transferred to PolyVinylidene

DiFluoride (PVDF) membranes. Membranes were blocked for 10 min with Turbo Transfer

Buffer and probed with monoclonal rabbit anti-EGFR antibody (Santa Cruz Biotechnology,

INC), monoclonal rabbit anti-PLCy, and anti-phosphorylated PLCy antibodies (Santa Cruz

Biotechnology, INC), monoclonal mouse anti-scr and rabbit anti-phosphorylated src

antibodies (Santa Cruz Biotechnology, INC), at dilution 1:1000. Samples were incubated

with Goat Anti-Rabbit IgG StarBright Blue 700 at a 1:2000 dilution and Anti-Tubulin

hFAB™ Rhodamine Antibody as a loading control at a 1:3,000 dilution for 3 hours and

washed with Blocking Buffer and Milli-Q H_2O (22-µm filtration). Immunoreactive

fluorescently labeled samples were visualized and analyzed with ImageLab Software.

Results

Initially, a phylogeny of all EGFR ligands was built (Figure S3), presenting a high degree of

monophyly among the seven paralogs. This monophyly justifies the comparison of different

ligands in our further analysis. Then, we developed a method for identifying amino acid

residues that are possibly involved in specific functions among paralogs, using what we

named cross-conservation approach. We applied the cross-conservation approach on the

seven ligands of EGFR to find out which mutations would have been key to switch function

in EGF. Cross-conservation analysis highlights functional positions in proteins, based upon

previous knowledge of their interacting partners, as explained in the methods section. The

combination of two residue conservation measures is used to generate a cross-conservation

plot (Figure 1B), by combining the alignment of all the ligands (paralogs that still share

binding to same receptor), using either the Multiple Sequence Alignment (MSA, Figure A in

Figure S1) or the Multiple Structural Alignment (MSTA, Figure B in Figure S1), and the

alignment of orthologs sequences of EGFs (herein so called evolutionary alignment, Figure C

in Figure S1).

Figure 1. Cross-conservation analysis.

(A)   Structure of extracellular EGFR-EGF complex (from PDB: 1ivo). EGFR extracellular

domain in cyan cartoon. EGF peptide is represented as cartoon in a gradient of blue to red

from the N-term to the C-term. I, II and III indicate the three ECD domains. C-tail of EGF is

disordered and in close proximity of domain III of EGFR. (B) Cross-conservation plot. The

plot is obtained by crossing the conservation measures of ligand alignments (rhombus for

MSA conservation, squares for MSTA conservation) and evolutionary alignments (Figure

S1). Positions highlighted in purple have been chosen for experimental verification. The

color gradient shows the N-/C-end displacement of the amino acid consistently with panel A.

Distance from the diagonal (e.g., 'd' in the plot) is used to calculate the cross-conservation

score. Interestingly, no point lies in the bottom right half of the plot suggesting that ligand

and evolutionary conservation are not independent and differently influenced by evolution

pressure. The C-terminus amino acids have higher cross-conservation score on average,

highlighting that this part has a functional role. (C) Extract of the paralog ligands alignments

focusing on the chosen positions, showing a low degree of conservation. (D) Extract of the

logo generated from the orthologs evolutionary alignment (Figure S1 panel C). Positions

chosen for mutation are highly conserved, therefore resulting in a high cross-conservation

score.

The cross-conservation score is calculated as the distance (d, Figure 1B) from the diagonal.

The list of residues sorted by score is shown in S1 Table. According to our analysis, residues

with high scores are concentrated on the C-terminal tail. This result is in line with previous

reports on the C-tail of EGF (27-29). Interactions between the interface of the ligand and

EGFR have been shown to influence the stability of the dimerized receptor (30), a factor that

is related to ligand-specific signaling bias (24, 31).

Based on the factors described in the methods section, we designed EGF mutants with single

amino-acid substitutions (N32R, D46T, K48T and W50Y). We characterized the mutants

biochemically and at the cellular level. Using circular dichroism (CD) experiments, we

confirmed that the secondary structure of these mutants was maintained (Figure S4). Then the

K_d for EGFR was determined *in vitro* by Isothermal Titration Calorimetry (ITC). ITC

measurements of the binding of all mutants to the ECD of EGFR exhibited similar K_d values

to the WT EGF. Only N32R has 2-fold higher affinity for EGFR compared to WT EGF

(Figure 2B).

Figure 2. ITC measurements of EGF mutants and the EGFR receptor.

(A) ITC analysis of WT EGF ligand and mutants N32R, D46T, and K48T binding to the

ECD of the EGFR receptor at 25°C. Measurements were taken by adding WT EGF at 200

μM to the ECD of EGFR at 20 μM. (B) Four zoom-in of X-Ray structure of the extracellular

domain (ECD) of EGFR bound to EGF (PDB 1IVO). In cyan cartoon the ECD of EGFR.

Each zoom-in focuses on the mutated residue. Highlighted in yellow stick side chain of the

mutated residues and in cyan stick side chain of the residue of ECD in proximity (< 5 Å with

the mutated residue. (C) K_d calculated from the ITC measurements using the program

Affinimeter KinITC Kintecs Software.

Mutation N32R is on the interface between ligand and receptor (Figure S5). The slightly

higher affinity is probably due to the presence of the guanidinium group of R which is

positive charged and could interact with Q16 of EGFR ECD.

Surprisingly, although the biochemical parameters are not substantially different, EGF

variants differently affected cell growth in cell proliferation studies. Compared to EGF, all

the mutants exhibited an intermediate effect on the proliferation of human fibroblasts bj5-tα

(Figure 3A). D48T and K48T mutants differed most from the WT (Figure S7). A similar

response was observed in Albino Swiss mouse 3T3 cell line (Figure S6), and in both cell

lines, the effects are concentration dependent.

In order to validate the differential effect, we performed an apoptosis assay using A431 cell

lines. After 24h of growth, the cells were treated with varying concentration of EGF or

mutated ligand. 48h after 100 nM treatments, EGF showed the highest decrease in cell

proliferation, while the four mutants had an intermediate effect, with W50T being the closest

to the WT (Figure 3B). Evidence of apoptosis was found from the globular processes formed

in cells treated with WT EGF (Figure 3C).

Figure 3. Results of cell growth assays for cells treated with EGF variants.

(A) Effect of different concentrations of EGF variants on proliferation of the human normal

fibroblast bj5-tα cell line. Data represent the growth rate percentage calculated from the

confluence of cells relative to the control (mean+- standard deviation) for each concentration

of EGF. Percent confluence was estimated 24h after the treatment (two replicates/treatment).

(B) Effect of EGF mutants on the growth of A431 cells. EGF induced apoptosis is measured

as the reduction in cell proliferation compared to the control. (C) Comparison of A431 cell

growth after treatment with 100 nM WT EGF and EGF variants D46T and K48T. Cells were

labeled with fluorescent annexin V green reagent. Plates were pre-warmed prior to data

acquisition to avoid condensation and expansion of the plate, which affect autofocus. Images

were captured every 2 hrs (4x) for 3 days in the IncuCyte system.

Significantly herein we report the effects of one amino acid change on the C-terminus tail of

EGF ligand. To explain this behavior, we proceed analyzing the downstream proteins

involved in the EGFR network. Both D46T and K48T reduced the expression of

Phospholipase-Cγ1 (PLCγ1), which is a downstream signaling protein required for EGFR-

induced squamous cell carcinoma (Figure 4A). The low levels of PLCγ protein lead to a

decreased amount of PKCζ protein (Figure 4A). Our findings show that modularity in the

downstream kinase cascade could be induced by one single amino acid substitution.

Figure 4. EGF variants D46T and K48T affect the EGFR downstream signaling

pathway.

(A) Western blot analysis of EGFR-regulated downstream gene expression of EGF variant-

treated A431 cancer cells. Expression of Epidermal Growth Factor Receptor (EGFR),

Phospholipase-Cγ1 (PLCγ1) and PKCζ protein in A431 cancer cell line after treatment with

100 nM WT, EGF variant D46T or variant K48T. Samples were collected on Day 3 after

treatment (two duplicates). Samples were incubated with Goat Anti-Rabbit IgG StarBright

Blue 700 at a 1:2000 dilution and Anti-Tubulin hFAB™ Rhodamine Antibody as a loading

control at a 1:3,000 dilution for 3 hrs and washed with Blocking Buffer and Milli-Q H$_2$O (22

µm filtration). Immunoreactive fluorescent labeled samples were visualized and analyzed

with ImageLab Software. (B) A schematic representation of one of the EGF-EGFR-mediated

signaling pathways that may be initiated in the A431 epidermoid cancer cell line. Arrows

indicate the positive action of downstream gene expression, whereas arrows with flat tips

indicate inhibition of gene expression. The "P+" symbol represents phosphorylation of

downstream-regulated proteins. The dashed line represents the potential cellular effect

regulated by altering gene expression levels involved in the depicted pathway.

Discussion

The prediction of functional residues is a well-developed field (32), where conservation of

each residue in a protein is considered a key factor to rely on. Tools like ConSurf (23) and the

ET-like methods (33) are able to identify slowly evolving positions that are involved in

folding, interaction, or catalytic activity of protein (32). Though, the specific reason why a

residue is conserved remains often unclear. In this work, we show that the conservation score

in the structural alignment of paralog sequences, combined with the orthologs alignment

conservation score is a promising way to identify important residues that affects the

downstream pathway and cellular behavior.

Cross-conservation analysis is based on the following assumptions: 1) Orthologs

evolutionary alignment conservation shows whether a residue is important for either

structural or functional reasons. 2) Ligands alignment conservation scores denote the

importance of a residue for receptor binding (the main shared property of all ligands). In our

analysis we rely on these two assumptions to conclude that highly conserved residues in the

evolutionary alignment (Figure C in Figure S1) that are not conserved in the ligand alignment

(Figure A in Figure S1) have ligand-specific relevance related to their function.

It is interesting to note that conserved positions in the paralogs alignments are a subset of

those conserved in the orthologs alignment. By subtracting the first ones, positions with a

shared function across all ligands are filtered out. Then, Cross-conservation analysis

overcomes the limitations of previous methods and highlights ligand-specific functional

residues.

In particular, the two tryptophan in positions 49 and 50 are strong outliers on our

bioinformatics analysis (Figure 1B). Their score is high even when using conservation

measures that do not take amino acid type change into account (data not shown). We

specifically chose W50 for further testing because of its outward facing position, on the

hypothesis that it might mediate previously unknown interactions. Mutant W50Y did not bind

stronger than WT EGF but it showed a cellular effect. This result further highlights the

importance of the disordered C-tail in the signal transduction.

Biochemical characterization of ligand-receptor interaction was done in solution using ITC.

The experiment was performed using the soluble extra-cellular domain of the receptor

followed by the cellular experiments. Tested mutants have binding affinities similar to that of

WT EGF (Figure 2) except mutant N32R which showed slightly higher affinity. N32R was

the only position chosen which is not on the C-terminus tail. Interestingly, while retaining the

same binding affinity to the receptor, all mutants showed a different cellular effect on Bj-5ta

cells, compared to EGF. A delayed proliferation response compared to control might appear

as counterintuitive for fibroblast (8). Though, Bj-5ta hTERT immortalized fibroblasts show

features that distinguish them from in vivo fibroblast, like differential gene expression and

epiregulin-dependent proliferation (34). Competition between endogenously synthesized

epiregulin and EGF might generate a different balance in the receptor stabilization, therefore

leading to a decreased cell proliferation after EGF treatment, as shown from the cell culture

videos in the Supplementary Materials.

A431 constitutionally express EGFR in high levels. Treatment with EGF has been observed

to promote STAT1 dependent apoptosis (35). This pathway is sustained by the internalization

of the ligand-receptor complex by the endocytosis process, a key factor in the ligand induced

biased signaling (13). In cancer cell line A431, the cells treated with mutants show a higher

growth rate, therefore a decreased rate in apoptosis compared to WT. Given the setting of the

experiment, modularity in the endocytosis pathway is a probable explanation.

Interestingly, K48T treated cells showed an increase in long and thin processes during growth

in all replicates (Figure 3). A possible hypothesis could be EGF-induced tubular formation, a

further EGF induced pathway reported in intestinal epithelial cells (36).

In both experiments, the growth rate of the mutants is intermediate between the control and

EGF, and concentration dependent. This phenomenon, and the cell-line specific response of

growth hormones is consistent with previous literature reports (37).

Since the four mutations are located in the disordered C-terminus, we cannot infer whether

they disrupt an important contact for EGF high-affinity binding. In fact, we observe the same

binding affinity with the isolated ECD as WT EGF. However, we can assume that they might

induce a different balance of the receptor dimerization, which then affect the downstream

pathway. Previous reports have also identified the importance of the C-terminus for binding

specificity (38) and binding affinity (39) as we did. Some studies have examined the effect of

individual positions at the C-terminus of EGF (40), although nobody has investigated the

residues reported herein. Though, other EGF mutants with a high cross-conservation score

have been previously reported to alter the EGF effect (41-43).

We speculate that the mutations might induce a conformational change of the receptor that

might affect interactions in the highly modulated endocytosis (44). Cancer treatment has

focused on the EGF receptor and deactivation of its intra-cellular tyrosine-kinase (45). As the

design of EGFR-based drugs remains complex, our study may support the hypothesis that the

D46T and K48T EGF mutants can be used as templates to design anti-cancer drugs.

Author contributions

PL, SP, DM conceived and designed the project; SP developed and performed the

computational models and analyses; DM performed the experiments; all authors analyzed the

data and wrote the manuscript.

Acknowledgments

We thank Tadashi Yamamoto's Unit and Russel Alenton for technical help with the cell

proliferation assay, and Keiko Kono's Unit for the technical help with the WB experiments.

We thank Madhuri Gade, Hayato Yanagida, Mary Collins and Mirco Dindo for critical

reading.

Funding

Financial support by the Okinawa Institute of Science and Technology to P.L. is gratefully

acknowledged.

References

1.   Wu JM, Vallenius T, Ovaska K, Westermarck J, Makela TP, Hautaniemi S. Integrated
network analysis platform for protein-protein interactions. Nat Methods. 2009;6(1):75-7.
2.   Mann M, Jensen ON. Proteomic analysis of post-translational modifications. Nat
Biotechnol. 2003;21(3):255-61.
3.   Rao VS, Srinivas K, Sujini GN, Kumar GN. Protein-protein interaction detection:
methods and analysis. International journal of proteomics. 2014;2014:147648.
4.   Laisney J, Braasch I, Walter RB, Meierjohann S, Schartl M. Lineage-specific co-
evolution of the Egf receptor/ligand signaling system. BMC Evol Biol. 2010;10:16.
5.   Gazdar AF. Activating and resistance mutations of EGFR in non-small-cell lung
cancer: role in clinical response to EGFR tyrosine kinase inhibitors. Oncogene. 2009;28:S24-
S31.
6.   Cantor AJ, Shah NH, Kuriyan J. Deep mutational analysis reveals functional trade-
offs in the sequences of EGFR autophosphorylation sites. Proc Natl Acad Sci U S A.
2018;115(31):E7303-E12.
7.   Nicholson RI, Gee JM, Harper ME. EGFR and cancer prognosis. European journal of
cancer (Oxford, England : 1990). 2001;37 Suppl 4:S9-15.
8.   Carpenter G, Cohen S. Epidermal growth factor. Annual review of biochemistry.
1979;48:193-216.

9.   Hardwicke J, Schmaljohann D, Boyce D, Thomas D. Epidermal growth factor therapy
and wound healing--past, present and future perspectives. The surgeon : journal of the Royal
Colleges of Surgeons of Edinburgh and Ireland. 2008;6(3):172-7.

10.   Misale S, Yaeger R, Hobor S, Scala E, Janakiraman M, Liska D, et al. Emergence of
KRAS mutations and acquired resistance to anti-EGFR therapy in colorectal cancer. Nature.
2012;486(7404):532-6.

11.   Evans BA, Sato M, Sarwar M, Hutchinson DS, Summers RJ. Ligand-directed
signalling at beta-adrenoceptors. Br J Pharmacol. 2010;159(5):1022-38.

12.   Kenakin T. Functional selectivity and biased receptor signaling. J Pharmacol Exp
Ther. 2011;336(2):296-302.

13.   Ali R, Brown W, Purdy SC, Davisson VJ, Wendt MK. Biased signaling downstream
of epidermal growth factor receptor regulates proliferative versus apoptotic response to
ligand. Cell Death Dis. 2018;9(10):976.

14.   Groenen LC, Nice EC, Burgess AW. STRUCTURE-FUNCTION-RELATIONSHIPS
FOR THE EGF/TGF-ALPHA FAMILY OF MITOGENS. Growth Factors. 1994;11(4):235-
57.

15.   Souriau C, Gracy J, Chiche L, Weill M. Direct selection of EGF mutants displayed on
filamentous phage using cells overexpressing EGF receptor. Biol Chem. 1999;380(4):451-8.

16.   Zerbino DR, Achuthan P, Akanni W, Amode MR, Barrell D, Bhai J, et al. Ensembl
2018. Nucleic Acids Res. 2018;46(D1):D754-D61.

17.   Katoh K, Standley DM. MAFFT Multiple Sequence Alignment Software Version 7:
Improvements in Performance and Usability. Mol Biol Evol. 2013;30(4):772-80.

18.   Berman HM, Westbrook J, Feng Z, Gilliland G, Bhat TN, Weissig H, et al. The
Protein Data Bank. Nucleic Acids Res. 2000;28(1):235-42.

19.   Goddard TD, Huang CC, Meng EC, Pettersen EF, Couch GS, Morris JH, et al. UCSF
ChimeraX: Meeting modern challenges in visualization and analysis. Protein Sci.
2018;27(1):14-25.

20.   Okonechnikov K, Golosova O, Fursov M, Team U. Unipro UGENE: a unified
bioinformatics toolkit. Bioinformatics. 2012;28(8):1166-7.

21.   Nguyen LT, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: A Fast and Effective
Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. Mol Biol Evol.
2015;32(1):268-74.

22.   Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS.
ModelFinder: fast model selection for accurate phylogenetic estimates. Nat Methods.
2017;14(6):587-+.

23.   Ashkenazy H, Abadi S, Martz E, Chay O, Mayrose I, Pupko T, et al. ConSurf 2016:
an improved methodology to estimate and visualize evolutionary conservation in
macromolecules. Nucleic Acids Res. 2016;44(W1):W344-W50.

24.   Freed DM, Bessman NJ, Kiyatkin A, Salazar-Cavazos E, Byrne PO, Moore JO, et al.
EGFR Ligands Differentially Stabilize Receptor Dimers to Specify Signaling Kinetics. Cell.
2017;171(3):683-95.e18.

25.   Puddicombe SM, Wood L, Chamberlin SG, Davies DE. The interaction of an
epidermal growth factor/transforming growth factor alpha tail chimera with the human
epidermal growth factor receptor reveals unexpected complexities. The Journal of biological
chemistry. 1996;271(48):30392-7.

26.   Wiedemann C, Bellstedt P, Gorlach M. CAPITO--a web server-based analysis and
plotting tool for circular dichroism data. Bioinformatics. 2013;29(14):1750-7.

27.   Mehrabi M, Mansouri K, Soleymani B, Hoseinkhani Z, Shahlaie M, Khodarahmi R.
Development of a human epidermal growth factor derivative with EGFR-blocking and

depleted biological activities: A comparative in vitro study using EGFR-positive breast
cancer cells. Int J Biol Macromol. 2017;103:275-85.
28.   Panosa C, Tebar F, Ferrer-Batalle M, Fonge H, Seno M, Reilly RM, et al.
Development of an epidermal growth factor derivative with EGFR blocking activity. PloS
one. 2013;8(7):e69325.
29.   Engler DA, Montelione GT, Niyogi SK. Human epidermal growth factor. Distinct
roles of tyrosine 37 and arginine 41 in receptor binding as determined by site-directed
mutagenesis and nuclear magnetic resonance spectroscopy. FEBS letters. 1990;271(1-2):47-
50.
30.   Mehrabi M, Mahdiuni H, Rasouli H, Mansouri K, Shahlaei M, Khodarahmi R.
Comparative experimental/theoretical studies on the EGFR dimerization under the effect of
EGF/EGF analogues binding: Highlighting the importance of EGF/EGFR interactions at site
III interface. Int J Biol Macromol. 2018;115:401-17.
31.   Knudsen SL, Mac AS, Henriksen L, van Deurs B, Grovdal LM. EGFR signaling
patterns are regulated by its different ligands. Growth Factors. 2014;32(5):155-63.
32.   Nemoto W, Saito A, Oikawa H. Recent advances in functional region prediction by
using structural and evolutionary information - Remaining problems and future extensions.
Computational and structural biotechnology journal. 2013;8:e201308007.
33.   Lichtarge O, Bourne HR, Cohen FE. An evolutionary trace method defines binding
surfaces common to protein families. J Mol Biol. 1996;257(2):342-58.
34.   Lindvall C, Hou M, Komurasaki T, Zheng C, Henriksson M, Sedivy JM, et al.
Molecular characterization of human telomerase reverse transcriptase-immortalized human
fibroblasts by gene expression profiling: activation of the epiregulin gene. Cancer Res.
2003;63(8):1743-7.
35.   Grudinkin PS, Zenin VV, Kropotov AV, Dorosh VN, Nikolsky NN. EGF-induced
apoptosis in A431 cells is dependent on STAT1, but not on STAT3. Eur J Cell Biol.
2007;86(10):591-603.
36.   Ibuka S, Matsumoto S, Fujii S, Kikuchi A. The P2Y(2) receptor promotes Wnt3a- and
EGF-induced epithelial tubular formation by IEC6 cells by binding to integrins. J Cell Sci.
2015;128(11):2156-68.
37.   Bjorkelund H, Gedda L, Andersson K. Comparing the epidermal growth factor
interaction with four different cell lines: intriguing effects imply strong dependency of
cellular context. PloS one. 2011;6(1):e16536.
38.   Sato K, Nakamura T, Mizuguchi M, Miura K, Tada M, Aizawa T, et al. Solution
structure of epiregulin and the effect of its C-terminal domain for receptor binding affinity.
FEBS letters. 2003;553(3):232-8.
39.   Wingens M, Walma T, van Ingen H, Stortelers C, van Leeuwen JE, van Zoelen EJ, et
al. Structural analysis of an epidermal growth factor/transforming growth factor-alpha
chimera with unique ErbB binding specificity. The Journal of biological chemistry.
2003;278(40):39114-23.
40.   Matsunami RK, Yette ML, Stevens A, Niyogi SK. MUTATIONAL ANALYSIS OF
LEUCINE-47 IN HUMAN EPIDERMAL GROWTH-FACTOR. J Cell Biochem.
1991;46(3):242-9.
41.   van de Poll ML, Lenferink AE, van Vugt MJ, Jacobs JJ, Janssen JW, Joldersma M, et
al. A single amino acid exchange, Arg-45 to Ala, generates an epidermal growth factor (EGF)
mutant with high affinity for the chicken EGF receptor. The Journal of biological chemistry.
1995;270(38):22337-43.
42.   Lahti JL, Lui BH, Beck SE, Lee SS, Ly DP, Longaker MT, et al. Engineered
epidermal growth factor mutants with faster binding on-rates correlate with enhanced
receptor activation. FEBS letters. 2011;585(8):1135-9.

542 43. Lui BH, Cochran JR, Swartz JR. Discovery of improved EGF agonists using a novel
543 in vitro screening platform. J Mol Biol. 2011;413(2):406-15.
544 44. Bakker J, Spits M, Neefjes J, Berlin I. The EGFR odyssey – from activation to
545 destruction in space and time. Journal of Cell Science. 2017;130(24):4087-96.
546 45. Sotelo MJ, García-Paredes B, Aguado C, Sastre J, Díaz-Rubio E. Role of cetuximab
547 in first-line treatment of metastatic colorectal cancer. World journal of gastroenterology.
548 2014;20(15):4208-19.
549

Supplementary material

Figure S1. EGF Alignments.

(A) Multiple sequence alignment of the mature forms of EGF paralog. Protein were chosen

for retaining binding to the EGF receptor in human. Sequences were obtained from

ENSEMBL database and aligned with MAFFT software. On top, sequence logo representing

the most conserved residue positions, obtained with WebLogo, UC Berkeley. (B) Multiple

structure alignment of EGF paralogs. 1IVO pdb file containing crystallized EGFR (teal) and

EGF was used as a template to align the structures of the other EGFR ligands. The ligands

pdb files are: 2RNL – AREG; 1IP0 – BTC; 5WB8 – EPGN; 5WB7 – EREG; 1MOX –

HBEGF; 1XDT – TGFA. All ligands residues are colored by conservation level (red high,

blue low) in the MSTA. The superposition and coloring were performed using CHIMERA.

(C) EGF orthologs evolutionary alignment. Orthologous sequences were obtained from

ENSEMBL and filtered of duplicate sequences (eg. very closely related species). Fruitfly

(*Drosophila melanogaster*) was used as outgroup. On top, bar plot representing the

conservation value of each position. The conservation measures of these alignments have

been used to make the cross-conservation plot.

S1 Table. Cross-conservation score of all EGF positions.

The table reports the cross-conservation score calculated using either the ligands MSA or the

ligands MSTA. Highlighted in red, the position with the residues swap issue. In yellow,

positions that were initially considered for mutagenesis. In green, the positions used in the

further analysis. On the top right, conservation score plot by amino acid position. It is

interesting to note how the highest peaks are concentrated in the C-terminus part.

Figure S2. Highlight on the EGF-specific residue swapping problem.

EGF is colored in teal, while HB-EGF is colored in white. The tyrosine in EGF position 29

(29Y) is not found on the corresponding position of HB-EGF. Instead, a proline is found

(130P). Though, the interaction between the two residue is probably maintained because, in

place of the histidine 10 (10H) of EGF, HB-EGF shows a tyrosine (112Y). When this type of

residue swapping is ligand-specific, the lowering of the conservation measures over-estimates

the real change happening in that position.

Figure S3. EGFR ligands evolutionary tree.

The tree was obtained using Maximum Likelihood algorithm. *C elegans* EGF was used as

outgroup. The EGFR ligands cluster neatly. Interestingly, fish ligands are found to have

longer branches on average. The tree was made using iqtree ModelFinder. The best-fit model

used is JTT+R6.

Figure S4. The validation of ordered structure of WT EGF and variants.

0.2 mg/ml of the sample was added to 10 µm cuvette and analyzed by Circular Dichroism

(CD) spectrometer J-1000. All samples showed the presence of a comparable amount of β-

sheet. In contrast, K48T and W50Y had another peak at 215 nm. The CD of EGF was found

similar to previously published results. The plot and analysis was made using CAPITO web

tool (26).

Figure S5. Crystalized structure of WT EGF with the highlighted position N32, In the

interaction with the position Q16 and G18 of the EGFR receptor backbone.

Figure S6. Effect of 1nM, 10nM and 100nM concentration of EGF variants on the

proliferation of mouse normal fibroblast Albino Swiss 3T3 cell line.

The data represent the relative growth percentage to control based on the confluence of cells

(mean ± standard deviation) for each concentration of EGF.

Figure S7. Effect of 1nM, 10nM and 100nM concentration of EGF variants on the

proliferation of human normal fibroblast bj5-tα cell line.

The data represent the relative growth percentage to control based on the confluence of cells

(mean ± standard deviation) for each concentration of EGF. The percentage confluence was

estimated 48h after the treatment (two replicates/treatment).

Figure S8. Effect of 1nM and 10nM concentration of EGF variants on the proliferation

of A431 skin cancer cells.

The data represent the relative growth percentage to control based on the confluence of cells

(mean ± standard deviation) for each concentration of EGF variants. The percentage

confluence was estimated 48h after the treatment (two replicates/treatment).

A   **Bj5t-α Human Normal Fibroblast Cell Growth**

24h

Relative Growth Rate (%)

Concentration

1 nM, 10 nM, 100 nM

Legend: Control, WT EGF, K48T, N32R, D46T, W50Y

B   **A431 Human Cancer Cell Growth**

48h

Relative Growth Rate (%)

100 nM

Concentration

Legend: control, WT, K48T, N32R, D46T, W50Y

C   **A431 Human Cancer Cells**

100 nM

0h   48h

WT EGF

D46T

K48T